Dating the bacterial tree of life based on ancient symbiosis

Sishuo Wang[1,2], Haiwei Luo[1,3*]

[1]Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory
of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[2]Department of Microbiology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[3]Institute of Environment, Energy and Sustainability, The Chinese University of Hong Kong,
Shatin, Hong Kong SAR

[*]**Corresponding author:**

Haiwei Luo
School of Life Sciences, The Chinese University of Hong Kong
Hong Kong SAR
Phone: (+852) 39436121
E-mail: haiweiluo@cuhk.edu.hk
16

Author Contributions: SW and HL conceived the study and wrote the manuscript. SW
performed the experiments and analysed the data.

Competing Interest Statement: No conflict of interest declared.

Classification: Biological Sciences – Evolution; Physical Sciences - Earth, Atmospheric, and
Planetary Sciences

Key words: molecular clock, bacterial tree of life, timetree, symbiosis, relative time
constraint, early life evolution

This PDF file includes:

Main Text

Figures 1 to 4

Abstract

Obtaining a timescale for bacterial evolution is crucial to understand early life evolution but is difficult owing to the scarcity of bacterial fossils and absence of maximum age constraints of the available fossils. Here, we introduce multiple new time constraints to calibrate bacterial evolution based on ancient symbiosis. This idea is implemented using a bacterial tree constructed with mitochondria-originated genes where the mitochondrial lineage representing eukaryotes is embedded within Proteobacteria, such that the date constraints of eukaryotes established by their abundant fossils are propagated to ancient co-evolving bacterial symbionts and across the bacterial tree of life. Importantly, we formulate a new probabilistic framework that considers uncertainty in inference of the ancestral lifestyle of modern symbionts to apply 19 relative time constraints (RTC) each informed by host-symbiont association to constrain bacterial symbionts no older than their eukaryotic host. Moreover, we develop an approach to incorporating substitution mixture models that better accommodate substitutional saturation and compositional heterogeneity for dating deep phylogenies. Our analysis estimates that the last bacterial common ancestor (LBCA) occurred approximately 4.0-3.5 billion years ago (Ga), followed by rapid divergence of major bacterial clades. It is robust to alternative root ages, root positions, tree topologies, fossil ages, ancestral lifestyle reconstruction, gene sets, among other factors. The timetree obtained enables the validation of various hypotheses, such as the survival of life during the late heavy bombardment, the absence of a connection between ancient stromatolites and cyanobacteria, and the presence of aerobic enzymes before the oldest geochemical records of molecular oxygen.

Significance Statement

Bacteria, with their vast diversity and ancient history, play a crucial role in shaping Earth's biogeochemistry. However, the scarcity of fossils complicates the determination of their evolution timescale and its link to Earth's history. To address this issue, we have devised and implemented novel methods that utilize ancient symbiosis and eukaryotic fossils to calibrate bacterial evolution by molecular clock. We obtain a comprehensive genus-level evolutionary timeline of bacteria that sheds light on their profound influence on the development and diversity of life on our planet, as well as its environmental dynamics. This research greatly contributes to our understanding of microbial evolution and its implications for Earth's past and present.

Introduction

Bacteria are the dominating form of life in the first two thirds of the history of our planet, and their evolution and diversification have greatly shaped the geosphere of Earth (1). Many hypotheses have been proposed to interpret outstanding questions like how bacteria affected the evolution of biogeochemical cycles, and when major innovations of bacteria such as photosynthesis occurred and why they did not appear earlier (2–4). Interpreting these questions depends on temporal links between bacterial evolution and the corresponding geobiological events. However, efforts to date the early evolution of bacteria based on relaxed molecular clock (5–7) often fail to reach convergence. A major challenge is that bacteria have rarely left calibrations well suited for molecular dating, i.e., fossils attributable to or biomarkers diagnostic for certain clades (4). Further, because fossils or biomarkers by themselves provide only a lower minimum bound (8, 9), there are no reliable maximum time constraints to calibrate bacterial evolution. Accordingly, the time estimates heavily depend on the maximum age assigned to the root, i.e., the LBCA, whose origin time involves even more debate (10, 11). Moreover, previous studies in dating the early evolution of bacteria (3, 4, 11) are mostly based on site-homogeneous amino acid substitution models that do not well account for among-site compositional heterogeneity and substitutional saturation (12). In deep-time phylogenetics, this may lead to an underestimate of branch length (expected number of substitutions per site) (10), which potentially affects time estimation.

Recent studies have used eukaryotic fossils to date the evolution of Proteobacteria (11) and Cyanobacteria (13, 14) where mitochondrial and plastid-endosymbiosis are believed to have occurred, respectively. These successes imply the value of the abundant eukaryotic fossils in timing bacterial evolution by molecular clock, but more temporal information of eukaryote evolution is waiting to be harvested. One such source is bacteria-eukaryote symbiosis: as such symbioses occurred in multiple ancient bacterial lineages (15, 16), it is

tempting to fit it into a recently developed framework called relative time constraint (RTC)

dating (17, 18) which constrains the order of divergence by assuming the age of some nodes

of the tree to be younger than others. Despite some encouraging progress, the current

implementation of this method treats all RTCs as if they are known with full certainty.

However, this assumption might be oversimplified because of the uncertainty in identifying

the relative time order based on horizontal gene transfer (HGT) (19) or other evolutionary

events. Here, we addressed the above challenges by leveraging combined evidence of

sequences, fossils, and ancient symbiosis to calibrate an integrative timescale of bacterial

evolution. We developed and implemented a framework to propagate date constraints of

eukaryotes established by abundant fossils to ancient co-evolving bacterial symbionts and

other bacteria by requiring that the origin of symbionts postdates their hosts' while

accommodating the uncertainty in ancestral host reconstruction. We established timelines of

bacterial evolution at different taxonomy resolutions and discussed how they may affect our

views of the evolution of early life and habitat.

Results

A new framework for dating deep-time bacterial evolution informed by host-symbiont

association

We overcame the paucity of bacterial fossils in dating the bacterial tree by leveraging the

temporal information of eukaryotes recorded in their abundant fossils. Briefly, by

incorporating eukaryotes into bacterial phylogeny as mitochondrial lineages affiliated with α-

proteobacteria based on mitochondrial endosymbiosis (20, 21), it is tempting to co-estimate

the ages of both the eukaryotic host and bacterial symbiont (11) while applying the recently

developed RTC-based molecular dating (17, 18, 22). In the context of host-symbiont

association, this means constraining the origin time of the symbiont's crown group to be no

earlier than that of the host's total group, thereby providing maximum time constraints to

symbionts whose temporal information may further be propagated across bacteria.

It is well possible that the ancestor of modern symbionts adapted to a different host or

even a free-living lifestyle. We formulate a three-step approach accounting for the possibility

of host shift of symbionts through ancestral state reconstruction (ASR) and fit it into RTC-

based dating in a probabilistic framework, which we call probability-based RTC (pRTC; see

Methods for details). First, ASR is performed to estimate the probabilities that the ancestral

nodes of modern symbionts are associated with different hosts or are free-living. Second,

MCMC molecular dating is performed using mitochondria-originated genes with fossil

calibrations but without RTC to generate posterior samples of timetrees where the ages of

symbionts and eukaryotic hosts, which are placed as the mitochondrial clade sister to α-

proteobacteria (11, 20, 21), are co-estimated. Third, the posterior samples of timetrees

collected in Step 2 are examined to remove those where the host's age is younger than the

symbiont's, weighted by the probability of different states (hosts) taken by the ancestor of

modern symbionts estimated in Step 1, using a rejection sampling approach. The idea is

further illustrated in Fig. 1A. Assume that ASR infers the ancestral lifestyles of "symbiont1"

as $P(S_1 = mammal) = 0.9$ and $P(S_1 = freeliving) = 0.1$, and for "symbiont2" as

$P(S_1 = mammal) = 0.5$ and $P(S_1 = freeliving) = 0.5$. For the two timetrees in the

posterior samples shown in Fig. 1A, the probabilities of acceptance are $(0.9 \times 1 +$

$0.1 \times 1) \times (0.5 \times 1 + 0.5 \times 1) = 1$, and $(0.9 \times 1 + 0.1 \times 1) \times (0.5 \times 0 + 0.5 \times 1) = 0.5$,

respectively (see Eqs. 3-5 in Methods). If the above procedure is performed for all posterior

timetrees collected from Step 2, the posterior dates of all bacterial lineages, particularly the

symbiont, might shift toward the present due to the use of RTC informed by host-symbiont

association (Fig. 1A). As ancient bacterial symbionts are distributed in different deep-

branching lineages (15, 23), this method has a great potential to date the bacterial tree of life.

Integrating eukaryote temporal information by a Bayesian sequential approach

We reconstructed the phylogeny of bacteria using the 265 genomes and 60 genes carefully

selected in a comprehensive phylogenomics study (24) using the best-fit site-heterogeneous

profile mixture model LG+G+C60 (Table S1A; Note S1). We applied the time non-reversible

model (25) which infers the root of the bacterial tree as part of maximum-likelihood

phylogenetic reconstruction where different root positions may have different likelihoods.

This method does not depend on any outgroup, thereby avoiding long branch attraction

artefacts caused by using highly divergent sequences to root deep phylogenies. Our analyses

rooted the bacterial phylogeny in the neighbourhood of a clade consisting of Fusobacteriota

and DST (Deinococcota, Synergistota, and Thermotogota). The analyses also supported the

division of most bacterial lineages into Gracilicutes and Terrabacteria, and placed the CPR

(Patescibacteria) clade (Fig. S1), initially suggested as the earliest-split bacterial group (26,

27), as a late-split clade sister to Chloroflexi (Fig. S1; Table S2; Note S2). This is generally

consistent with a recent study which roots the bacterial tree using an independent method

based on gene-species tree reconciliation (24).

To apply pRTC based on host-symbiont co-evolution, we integrated eukaryotes and

bacteria in a unified framework to co-estimate their times using the mitochondrial

endosymbiosis-based dating strategy (11). Apparently, the ages of symbionts are most

strongly constrained if the host date is inferred with a high precision, thus less uncertainty.

We therefore took the advantages of the genome-scale data of eukaryotes by employing a

Bayesian sequential approach in which the posterior dates of a small number of organisms

with many genes are estimated to serve as the time priors of a second analysis with many

organisms but a few genes (28, 29). Note that sequential Bayesian molecular dating is

conceptually distinct from secondary calibrations (using time estimates from other studies as

calibrations) by accommodating the uncertainty in time estimation in the first-step analysis

within a Bayesian framework (Note S1.2). We estimated the divergence times of 29

representative eukaryotes with 11 fossil calibrations (Figs. S2-S3) and 320 orthologs

conserved across eukaryotes (30). The posterior dates of eukaryote nodes were in agreement

with most studies (11, 31–33). Next, we approximated the posterior times of each node in the

eukaryote tree by a best-fit probability distribution (Fig. S4; Note S1.2.3), which were further

used to calibrate the eukaryote nodes for dating the bacterial tree where eukaryotes were

placed as a subtree sister to α-proteobacteria (11, 20, 21).

A time-calibrated bacterial phylogeny based on ancient symbiosis

To apply the pRTC dating approach based on ancient symbiosis, we first performed

comprehensive MCMC-based ASR to infer the ancestral lifestyles of seven well-known

ancient clades where symbionts are widely distributed, four at a phylum level (Spirochaetota,

Tenericutes, Chlamydiae, and Elusimicrobiota) and three at an order level (Rickettsiales,

Holosporales, and Legionellales). The ASR was performed using the 16S rRNA gene, which

best captures bacterial lifestyle diversity by including environmental and uncultured samples,

based on the lifestyle information of modern members within each of the ancient bacterial

clades mainly consisting of symbionts (Fig. S5). We additionally included two genus-level

symbionts (*Buchnera* and *Blattabacterium*) which are classical host-symbiont co-evolution

examples (16). A total of 19 ancestral nodes consisting of 28 symbiont lineages from the

above nine symbiont clades were selected to represent the host-symbiont association-based

RTCs in subsequent analyses (Figs. 2, S6; Note S3.1).

Further, we identified 19 genes well conserved across bacteria (Figs. S7-S8) from 108

genes with a mitochondrial origin reported in a recent study (21). We performed molecular

clock analysis with these 19 genes on a 148-genome focal dataset consisting of 29

eukaryotes, 28 symbionts, and 91 non-symbionts. The phylogeny was based on the tree

reconstructed above using the non-reversible model and with eukaryotes placed as the sister

to α-Proteobacteria. The posterior dates of eukaryotic lineages inferred in the first-step

sequential analysis above, and five lineage-specific fossil- or biomarker-calibrations within

bacteria (see Methods and Note S3.2), were used as time calibrations. We employed

MCMCTree's approximate likelihood method, which obtains a 1000× speed-up without loss

of accuracy compared with the exact likelihood method (34), well-suited for our large-scale

molecular clock analysis. However, the amino acid substitution models (e.g. LG+G) used to

calculate branch lengths in MCMCTree (and many other molecular dating software such as

BEAST and MrBayes) do not consider across-site compositional heterogeneity and often

show poor model fit in reconstructing deep phylogenies (10, 12).

We therefore developed a bootstrap-based approach to incorperating branch length

estimates inferred under the site-heterogeneous profile mixture substitution model

LG+G+C60, which better accommodates issues in deep-time phylogenetics like substitution

saturation (10), into MCMCTree's approximate likelihood method (Fig. 1B; see also

Methods). Briefly, maximum-likelihood trees were built for 1000 bootstrap alignments, and

the inverse of the bootstrap covariance matrix of their branch lengths were calculated to

approximate the hessian matrix which is necessary for MCMCTree's approximate method.

Simulations showed considerable improvements in the estimates of both ages and absolute

rates by integrating site-heterogeneous models with this approach, partcularly in the absence

of internal calibrations or if the divergence times are older than 3.0 Ga (Figs. S9-S10; see

Note S4 for details).

Next, we followed Eq. (5) (Methods) to apply a rejection sampling approach to filter out

the posterior timetrees obtained without applying RTC from Step 2, based on the probabilities

of the ancestor of modern symbionts being associated with different eukaryotic hosts or free-

living inferred from Step 1 (Figs. 1A, S8C). Collectively, using a carefully selected focal

dating scheme (Data S3), we estimated that the LBCA occurred at around 3648 Ma (95%

HPD [highest posterior density] 4111-3162 Ma) (Fig. 2), followed by the split between

Terrabacteria and Gracilicutes estimated to be 3629 Ma (95% HPD 4103-3167 Ma). Most

major clades, many of which are beyond a phylum level, diverged during 3500-3000 Ma

(Fig. 2). All selected symbiotic bacteria lineages (purple nodes in Fig. 2) originated no earlier

than 1700 Ma, approximately the appearance time of the LCA of eukaryotes.

We gradually included eukaryotic temporal constraints, the LG+G+C60 substitution

model, and host-symbiont association-based RTCs to improve date estimates of bacteria

(lower triangle in Fig. 3A). The inclusion of eukaryote's timing (*Strategy2*) led to a ~13%

decrease in the posterior dates compared with *Strategy1* where the five bacterial calibrations

were the only constraints (*Strategy1*). When the improved branch length estimates based on

the LG+G+C60 model was additionally included (*Strategy3*), those nodes between 3000 and

2000 Ma were estimated to be older by 7% (*Strategy2* vs. *Strategy3*). Further, when pRTC

was applied on the basis of *Strategy3*, the posterior dates of almost all nodes systematically

decreased ~5% (*Strategy3* vs. *Focal* in Fig. 3A), suggesting that the constraint on the time

order imposed by host-symbiont co-evolution was propagated through the bacterial tree of

life. Collectively, the combination of the use of time constraints provided by eukaryote

temporal information, site-heterogeneous profile mixture models, and host-symbiont

association-based RTCs estimated the divergence times of analysed bacteria ~15% younger

than those obtained without any of these settings (*Strategy1* vs. *Focal* in Fig. 3A; this

represents a net reduction and some effects cancelled out as illustrated above). As to the

posterior rate for each branch, the use of C60 led to an almost doubling of rate estimate

(upper triangle in Fig. 3A; *Strategy2* vs. *Strategy3*), indicating that the absolute rates are

largely underestimated by the compositionally homogeneous LG+G model. Moreover, when

we successively increased the maximum time prior of the root from 4500 Ma to 6000 Ma, the

posterior dates of bacterial lineages increased by more than 20% with *Strategy1*, but

remained almost unchanged with the focal scheme (Figs. 3B, S11A), indicating that the

(maximum) time constraints imposed by eukaryotes effectively constrained the timing of

bacterial evolution, such that their time estimates were robust to the root calibration where

much debate has arisen (10, 35).

Accommodating the uncertainty in divergence time estimation

We evaluated the influence of multiple sources of uncertainties in molecular clock analysis

on the posterior dates. First, fossil-based calibrations are known to play a major role in

molecular dating (36). The absolute time calibrations consisted of 11 eukaryotic ones, five

from within bacteria, and one at the root (LBCA). Replacing the 1.047 Ga-old fossil of

Bangiomorpha, the most widely recognized oldest crown-group eukaryote (32, 33), by the

recently discovered 1.6 Ga-old *Rafatazmia* (37), pushed the LCA of eukaryotes from ~1600

Ma to ~2100 Ma (Fig. S3), but had little impact on the time estimates of bacteria (*Euk1* in

Fig. 3C). A similar pattern was observed when the recently discovered 0.89 Ga-old sponge

fossil instead of the 0.55 Ga-old one (Note S3.2) were alternatively used as the minimum age

of the total-group animals (*Euk2* in Fig. 3C). As to the five bacterial calibrations, the two

cyanobacteria fossils and the two biomarkers displayed considerable impacts on the posterior

dates as their removal led to a decrease of ~7% in time estimates (*noCyanoFossil* and

noBiomarker in Fig. 3C). Replacing the 1.6 Ga-old Nostocales fossil calibration (38) by a

more disputable estimate based on the 2.0 Ga-old fossil (39) led to an increase in posterior

dates by more than ~10% (Fig. S11B). Except the above, the use of all alternative fossil

calibrations returned similar time estimates of both eukaryotes and bacteria (Figs. S11B,

S12).

Second, another source of time constraint is the use of ASR-based RTC. In the above

analysis, the ancestral states (hosts) between nodes within the same tree were considered

independent with each other such that their joint probability can be approximated by the

multiplied marginal probability (Fig. S6B). To check if this approximation affected the final

time estimates, we directly estimated the joint probability of their ancestral states using a

stochastic character mapping (SCM) method (see Methods). This did not change the posterior

dates (*joint_prob* in Fig. 3C). Further, while we have conducted comprehensive work in

classifying the traits into different hosts or free-living, there exist different interpretations of

the phenotypic traits particularly free-living, which might affect ASR. Additionally, we tried

different cut-offs of 16S OTU and a different algorithm (maximum-likelihood) in ASR, all of

which resulted in highly consistent time estimates (Fig. 3C). Including 13 more symbiont

lineages returned similar time estimates (*more_RTCs* in Fig. S12), suggesting that adding

host association-based RTCs would not lead to further change in the posterior ages of

bacteria.

Third, depencence of the results on the phylogenomic tree topology was investigated.

When molecular clock analysis was performed with three alternative root positions, two near

the root (*root0*) used in the focal analysis (*root1* and *root2* in Fig. 3D; Fig. S1), and one

(*rootCPR*) at the CPR clade as suggested by earlier studies (26, 27), the posterior dates

remained similar for most clades, except those that were alternatively placed as an early-split

phylogenetic position, particularly the CPR. Alternative phylogenetic positions of

mitochondria or between eukaryotic groups (Fig. S8D-S8E) displayed minor impacts on the

posterior dates of bacteria (Fig. S11C).

Fourth, the use of different genes in dating may lead to different divergence time

estimates. Mitochondrial genes tend to evolve faster than their orthologs in bacterial genomes

(11, 20). In theory, the issue can be accommodated by relaxed molecular clock models, but

this is not guaranteed (6, 7). Also, MCMCTree assumes a single tree topology for all genes,

which might not always hold true due to HGT, unresolved paralogy, or limited phylogenetic

signal of individual genes. To minimize the bias caused by gene selection, we ranked the 19

mitochondria-originated genes by i) the relative difference in their estimated evolutionary

rates between eukaryotic and bacterial lineages, and ii) $\triangle$LL, an index to measure the

species- and gene-tree incongruence (10). We selected the 5, 10, and 15 top-ranking (best)

genes using each of the above two measures and repeated molecular clock analyses (Fig. 3D).

Additionally, we analysed the full set of 32 mitochondria-originated genes (*all_genes* in Fig.

3D). Most analyses showed consistent results compared with using the 19-gene dataset; the

schemes using only the 5 top-ranking genes (schemes *top5*) displayed larger uncertainty in

the date estimates of certain nodes (e.g., MBDD), but this is likely due to their small number

of genes.

Fifth, other factors potentially affecting the dating analysis were examined. The largest

change in posterior dates were observed when the autocorrelated rates (AR) model, which

was consistently supported as the preferred model (Fig. S13), was replaced by the poorly

supported independent rates (IR) model: the divergence times of most lineages shifted toward

the present by ~20% (*IR* in Fig. 3C). The priors on the birth-death model which provides time

priors for nodes without time calibration and on the lineage-specific rate showed limited

impact (Fig. S11B). In addition, analysis with an independently sampled and expanded

dataset with 285 bacterial genomes (24) showed similar time estimates (*Secondary* in Fig.

S12), suggesting that the time estimates presented in the main analysis were not dependent on

a specific taxon sampling. Note that the overall acceptance rate of the rejection sampling was

less than 5%, meaning that more than 95% posterior were removed in Step 3. Because

running time of MCMCTree is roughly proportional to the number of partitions, to save time,

all analyses were performed on a single partitioned alignment, but similar patterns were

found when two partitions were used (Fig. S14).

A taxonomically high-resolution timetree of bacteria

Based on the time estimates from the above analysis where most lineages were sampled at a

phylum level, we proceeded with a sequential approach to date a taxonomically higher-

resolution timetree of bacteria. Due to the large computational burden associated with

Bayesian-based molecular clock methods, we applied the fast dating program RelTime.

Speficially, we used lineages included in the 148-genome dataset as the backbone, and used

their posterior dates estimated in the above analysis as calibrations to further infer a full-scale

bacterial timetree comprising 8898 tips each representing a genus based on GDTB (r95)

classification (Figs. 4A, S15; Data S4). The three major clades of Proteobacteria, α-, β-, and

γ-Proteobacteria, were estimated to originate ~2.1, ~1.6, and ~1.9 Ga, respectively, greatly

postdating the origin of those previously known as δ-Proteobacteria whose origin time was

estimated no later than 3.0 Ga (Fig. 4A). This supports the recent proposal to exclude δ-

Proteobacteria from the phylum Proteobacteria (40), which was further renamed as the

MBDD group (Myxococcota, Bdellovibrionota, Desulfomonadota, and Desulfobacterota)

(24). Firmicutes as a whole was the oldest phylum concerning the crown group, originating at

~3.3 Ga. Family-, order-, class, and phylum-level taxonomy groups were estimated to diverge

at 0.90 ± 0.31, 1.4 ± 0.36, 2.1 ± 0.46, 2.5 ± 0.56 Ga (mean $\pm$ standard deviation; Fig. 4B).

Further, we found a significant correlation between substitution rate of 16S rRNA gene and

divergence time at the family level ($p = 1.98E\text{-}44$, F-test; Fig. 4C). A simple linear regression

analysis suggests an average substitution rate of ~0.015 nucleotides/site/100 Ma for 16S

rRNA gene, but there are great differences between lineages (Fig. 4C). In other words, the

divergence time between bacteria may be implied by comparing the sequence of 16S rRNA

gene where 1% nucleotide difference roughly corresponds to divergence time of $1/0.015=67$

Ma. This indicates a longer time span for 1% nucleotide difference than previously estimated,

likely because the bacteria analysed in prior studies are mainly fast-evolving endosymbionts

(41–43). The correlation was robust to linear mixed models where the rate differnece of 16S

rRNA gene evolution in different lineages was controlled for (Note S1.4; Data S5).

Discussion

A major challenge in dating bacterial evolution is the paucity of credible calibrations (10, 11),

particularly the maximum time constraints. Previously, this was usually done by assigning a

somewhat arbitrary maximum age to the root (LBCA), which greatly varies between studies

due to the uncertainty in palaeobiological evidence and phylogenetic relationship around the

origin of life (10, 44). We addressed this problem by leveraging the timing information of

eukaryotes based on ancient symbiosis. This provides two additional layers of timing

information besides bacterial calibrations. First, we linked the evolution of bacteria and

eukaryotes by mitochondrial endosymbiosis. Because several eukaryotic calibrations have

maximum time bounds achieved based on the time estimates of the most recent geological

formation that ought to haven fossils of the clade of interest but that does not, this introduces

many reliable (maximum) time bounds to indirectly constrain bacterial evolution by

informing the substitution rates. Second, based on RTCs informed by host-symbiont

association, the timing of eukaryote evolution directly constrains the origin time of symbionts

to be no earlier than their eukaryotic host, which is further propagated across the bacterial

phylogeny.

The idea of RTC in molecular dating has recently caught much attention (17, 18, 45).

However, to our knowledge, none of the available RTC applications has considered the

uncertainty in the inference of RTC. In other words, previously the relative time order

established by horizontal gene transfer (i.e., gene recipient lineage cannot be older than gene

donor lineage) or any other forms of RTC was assumed to be known with certainty, which

might be oversimplified due to the uncertainties in identifying these evolutionary events. Our

new approach removes this assumption by weighting the different ancestral hosts by their

probability established in ASR through a rejection sampling framework. This greatly extends

the RTCs based on host-symbiont co-evolution to dating the bacterial tree, which involves

many ancient symbioses and large uncertainty in assigning the ancestral host. In addition,

prior studies that have applied the idea of host-symbiont association in molecular dating

typically used time estimates of the host from other studies (i.e., secondary calibrations) to

calibrate symbiont evolution, which may cause wrong time estimates by propagating errors

associated with the original time estimates (46). For example, the estimated origin times of

animals range from 1355 to 650 Ma according to 11 published recorded in the TimeTree

database (47). Apparently, calibrating with different times of animals can lead to dramatically

different time estimates of the symbionts. Our strategy co-estimates the origin times of hosts

and symbionts based on mitochondrial endosymbiosis (11), avoiding methodological and

data differences in comparing times estimates from different studies.

Great care should be taken when interpreting the estimated timeline of the bacterial tree

of life and reconstructed evolutionary events taking place anciently. Nonetheless, it is

remarkable to note the congruence between time estimates obtained using different time

constraints, different root positions, independent sets of genomes, and different settings in

molecular clock analysis. Our time estimate of LBCA is ~0.5 Ga younger than estimated in

some studies (4, 48). As evident in Fig. 3A, this might be due to the lack of enough maximum

time constraints and the use of poorly fit substitution models for inferring deep phylogenies

in these early efforts. Our study also improves molecular clock inferences by using a most-

updated phylogeny of the bacterial tree of life. For example, some groups once thought basal

in the bacterial tree, e.g., Aquificae (4) and CPR (26), are suggested as later-split lineages in

recent (10, 24) and the present study based on improved phylogenetics methods, leading to

different time estimates of corresponding lineages.

The estimated timescale of bacterial evolution allows testing hypotheses involving early

life evolution. For example, it is generally believed that the late heavy bombardment likely

occurred on early Earth ~4.1-3.8 Ga, during which large asteroids intensely pummelled Earth,

melted most parts of its surface, and killed off any forms of life that might have begun to

emerge (49, 50). Because primordial form of life likely had existed before LBCA, our

estimates for the age of LBCA at 4.0-3.5 Ga hint that life had survived the cataclysm,

consistent with modelling studies showing that much of Earth's crust and the microbes living

therein were unlikely affected by the intense bombardment (49). The robustness of the

posterior age of the LBCA to the maximum time prior assigned to it (Fig. 3B) provides

support to reject the hypothesis that life originated extra-terrestrially (35), where the LBCA

should be much older than the age of Earth. Further, under the most parsimonious hypothesis

that oxygenic cyanobacteria arose in the cyanobacterial stem lineage (45, 51), our analysis

placed the appearance of the first oxygenic cyanobacteria between 3 Ga (total-group

Cyanobacteria) and 2.5 Ga (crown-group Cyanobacteria) (Fig. 2). This suggests that one

cannot link stromatolites found in ancient sediments dated ~3.5 Ga to oxygenic cyanobacteria

despite their dominance in stromatolites nowadays (52, 53). This is supported by recent

evidence showing that stromatolites preserved in Archaean rocks neither required oxygen-

producing cyanobacteria for formation (54) nor were necessarily biogenic (55).

The new bacterial timetree also makes it possible to infer the mechanisms that drive the

origin of enzymes that involve oxygen. Accumulating evidence from isotope analyses

suggests the presence of oxygen on Earth back to 3.1 Ga (56–58), which could be produced

by cyanobacteria as mentioned above, and/or abiotically from H_2O_2 (59), potentially leading

to the formation of "oxygen oasis" thus extracellular access to O_2. A recent study (60) tracked

the origin of most oxygen-utilizing and -producing enzymes families to the Gracilicutes-

Terrabacteria split, which according to our estimate is ~3.6 Ga, predating the above events

for ~0.5 Ga (Fig. 2). This hints that life might have access to oxygen earlier than the oldest

geochemical record of the presence of oxygen (60), either extra- or intracellularly [possibly

under anaerobic conditions using alternative electron acceptors as observed in Thaumarchaea

(61) and the NC10 group (62)] but more likely the latter.

Our work shows that there is much more information beyond bacterial fossils for

calibrating bacterial evolution. The estimated timeline of bacterial evolution may well serve

as a foundation for studying the evolution of bacteria, their diversification rate, and its

correlation with Earth's geochemistry. It is hopeful that some of these questions can be

answered, and competing hypothesis can be reconciled by future molecular clock studies in

combination with modern phylogenetics and comparative genomics methods.

Methods

Phylogenomic reconstruction

A total of 62 marker genes considered conserved across bacteria were retrieved from a recent

phylogenomics study of the bacterial tree (24). Their sequences were aligned using MAFFT

v7.22261 (63) and trimmed using TrimAl v1.4 (settings: "-resoverlap 0.55 -seqoverlap 60")

(64). Unrooted phylogenomic trees were constructed based on the concatenated alignment

with time-reversible Markov model and each of the three empirical site-heterogeneous profile

mixture models C20, C40, and C60 using a PMSF approximation (guide tree built by

LG4M+G+I) with 1000 ultrafast bootstraps (65). The model LG+G+C60 was selected as the

best-fit model according to AIC and BIC, and the tree constructed by this model was used in

the main analysis. Rooted bacterial trees were constructed using the recently developed time-

non-reversible model implemented in IQ-Tree (25) that took the unrooted tree as either a

fixed tree or a start tree (see Note S2 for details).

Identification of mitochondria-originated genes conserved across the bacterial tree

We retrieved the sequences of the 108 genes identified to have a mitochondrial origin from

the study (21). We identified their homologs in eukaryotes encoded by either the

mitochondrial or nuclear genome based on BLAST with an e-value cut-off of 1e-30. Next, we

followed (66) to keep 32 genes present in at least one of the following three gene sets namely

bac120 (120 genes) (67), Battistuzzi2009 (25 genes) (4), and Coleman2021 (62 genes) (24),

which include orthologs well conserved and likely vertically transmitted across bacteria. Note

that two of the 62 orthologs originally included in Coleman2021 correspond to the same id

(K04485), and were both discarded. For each of the remaining 32 genes, we built a gene tree

using IQ-Tree, based on which we further removed i) two genes that did not show monophyly

between mitochondria and α-proteobacteria, indicating unresolved paralogy, and ii) 11 genes

where mitochondria/α-proteobacteria were placed at a basal position to all other bacteria with

different rooting approaches, indicating fast-evolving genes. This returned 19 mitochondria-

originated genes conserved across the bacterial tree, which were used in the main molecular

clock analysis (Table S3; Fig. S7).

Ancestral lifestyle reconstruction

The 16S rRNA genes of each interested symbiont group were collected from the NCBI

Entrez Nucleotide database with the keywords "taxon_name[ORGN] AND 16S[TITL] AND

rrna AND 1000:2000[SLEN] NOT spacer NOT trna" (April 2022). We also retrieved 16S

rRNA genes predicted by RNAmmer v1.2 (68) from genomes deposited in Genbank (August

2021). Those without any information of host or isolation source in the metadata were

removed (see SI). All remaining sequences were clustered using CD-HIT v4.7 (69) with

98.7% as the identity cut-off (70) into operational taxonomic units (OTUs), and one

representative was chosen from each OTU (see SI for alternative strategies in sequence

selection). We applied the stochastic character mapping (SCM) (71) to directly sample from

the joint posterior distribution, using *make.simmap* from the R package "phytools" v0.2-40

(72) to estimate ancestral hosts (or lifestyles) of modern symbionts. To account for

phylogenetic uncertainty, we applied the so-called most-recent-common-ancestor approach

(73) by identifying the LCA of the group of species of interest, reconstructing the state of that

node, and then summarizing the information across 100 ultrafast bootstrap (UFB) trees

reconstructed by IQ-Tree (settings: -m GTR+G). For each of the 100 UFB trees, 100

simulated stochastic character maps were generated using the MCMC algorithm implemented

in phytools under the "all-rates different" (ARD) model which allows transition rates at all

directions to be different, totalling $100 \times 100 = 10000$ simulations. All ASR analyses were

performed on phylogram—a phylogeny where branch lengths indicate amount of

evolutionary change, built by the 16S rRNA gene, as done in previous studies (11, 74).

Specifically, let S_i be the inferred lifestyle (or state) of the i^{th} internal node of symbionts,

which can take any possible hosts observed in the group of modern symbionts (including

free-living), denoted as s_i. By SCM, the marginal probability that the i^{th} internal node of

symbionts takes the state s_i may be approximated by

$$P(S_i = s_i) \approx \frac{n_{\{S_i = s_i\}}}{n}, \tag{1}$$

where $n_{\{S_1 = s_1\}}$ is the number that node i takes the state s_i during *n* rounds of SCMs.

Likewise, the same approach can be extended to estimate the joint probability that internal

nodes 1 to *k* take states $s_1, \ldots, s_k$ respectively by

$$485 \qquad P(S_1 = s_1, \dots, S_k = s_k) \approx \frac{n_{\{S_1=s_1,\dots,S_k=s_k\}}}{n}, \qquad (2)$$

486 where $n_{\{S_1=s_1,\dots,S_k=s_k\}}$ is the number that the event $\{S_1 = s_1, \dots, S_k = s_k\}$ happens during n

487 rounds of SCMs.

489 pRTC informed by host-symbiont association

Denote the data (i.e., sequence alignments) for molecular dating as D, and the divergence

times as $t = (t_1, \dots, t_k)$ where $t_1, \dots, t_k$ indicate the divergence time of the k ancestral nodes

of modern symbionts. Let S_i be the inferred lifestyle (or state) of the i^{th} ancestral node of

symbionts, which can take any possible hosts observed in the group of modern symbionts or

free-living, denoted as s_i. Let A indicate the event that the origin time of all symbionts is no

earlier than that of their host, i.e., $A = \{t_1 \le t_{S_1}, \dots, t_k \le t_{S_k}\}$. It follows that

$$496 \qquad f(t|D, A) \propto f(t|D)P(A|t, D)$$

$$497 \qquad\qquad\qquad = f(t|D)P(A|t) \qquad (3)$$

where $f(t|D, A)$ is the posterior of divergence times t given sequence data D and the

assumption A, $f(t|D)$ is the posterior of divergence times t given only sequence data D, and

$P(A|t)$ indicates the probability A is satisfied given time t. We further assume that the

inference of ancestral lifestyles of modern symbionts depends on only the phylogenetic

comparative data estimated by SCM and is independent of time. The second term on the

right-hand side of Eq. (3) can be rewritten as

$$504 \qquad P(A|t) = P\left(t_1 \le t_{S_1}, \dots, t_k \le t_{S_k} \middle| t\right)$$

$$505 \qquad\qquad = \sum_{s_1,\dots,s_k} P\left(t_1 \le t_{S_1}, \dots, t_k \le t_{S_k} \middle| t, S_1 = s_1, \dots, S_k = s_k\right) P(S_1 = s_1, \dots, S_k = s_k | t)$$

$$506 \qquad\qquad = \sum_{s_1,\dots,s_k} P\left(t_1 \le t_{s_1}, \dots, t_k \le t_{s_k} \middle| t, s_1, \dots, s_k\right) P(S_1 = s_1, \dots, S_k = s_k)$$

507
$$= \sum_{s_1,..,s_k} \left(\prod_{i=1}^{k} I(t_i, t_{s_i}) \right) P(S_1 = s_1, \ldots, S_k = s_k),$$

508 where $I(t_i, t_{s_i})$ is defined as

509
$$I(t_i, t_{s_i}) = \begin{cases} 1, t_i \leq t_{s_i} \\ 0, t_i > t_{s_i} \end{cases},$$

Particularly, we define $I(t_i, t_{fl}) = 1$ (free-living ancestral state) since the RTC that

symbiont's age is no older than the host's is automatically satisfied for free-living bacteria.

Further, assuming that the states of the ancestral nodes of symbiotic bacteria are

independent of each other so that $P(S_1 = s_1, \ldots, S_k = s_k) = \prod_{i=1}^{k} P(S_i = s_i)$, which

generally holds if the ancestral nodes are not phylogenetically too closely related (see Fig.

S6B for details), the above can be rewritten as

516
$$P(A|t) = \prod_{i=1}^{k} \sum_{s_i} I(t_i, t_{s_i}) P(S_i = s_i). \tag{4}$$

517 Collectively, by introducing Eq. (4) into Eq. (3), we have

518
$$f(t|D, A) \propto f(t|D) \left[\prod_{i=1}^{k} \sum_{s_i} I(t_i, t_{s_i}) P(S_i = s_i) \right]. \tag{5}$$

The above indicates that estimating the posterior dates under the assumption that the

origin time of all symbionts is no earlier than their hosts' while considering the uncertainty in

the inference of ancestral hosts can be obtained by:

i) Perform ASR to obtain the probability of each ancestral state for selected nodes in the

group of modern symbionts of interest.

ii) Perform MCMC molecular clock analysis with fossil calibrations but without any RTC.

iii) For each MCMC posterior sample collected from Step ii), perform a rejection sampling

by accepting it with a probability according to Eq. (5) (based on the ASR results obtained

from Step i).

Obviously, Step 2 is independent of Step 1, and Step 3 is based on Step 1 and Step 2.

Sequential Bayesian approach to dating the bacterial tree of life

We applied a sequential Bayesian approach (28, 29) to establish a timeline of the bacterial

tree, which allows using the genome-scale data and abundant fossil-based calibrations of

eukaryotes to better constrain symbionts' ages through the pRTC framework (see

Supplementary Note 1.3.2 for details). Sequential Bayesian molecular dating contains two

steps: i) run a dating analysis of the first subset of data and collect the posterior samples of

times and rates, ii) approximate the posteriors of each node in the first-step analysis by a

parametric probability distribution and use it as the prior in a subsequent analysis on a second

subset of data (see SI for details). We first used MCMCTree from PAML v4.10.0 (75) (see

Note S1.2 for details) to date an eukaryote tree with 29 selected species based on our

previous study (11) and 320 orthologs (30) with 11 fossil-based calibrations (Note S3.2), then

approximated the posterior samples of the dates obtained in the first-step analysis using a

best-fit probability distribution available in MCMCTree. The approximated dates were used

as the priors to calibrate the bacterial tree using the abovementioned 19 mitochondria-

originated genes conserved across bacteria (no overlap in genes between the first- and the

second-step analysis).

The posterior dates obtained in the above first-step sequential analysis were used

together with the following bacterial calibrations to date the bacterial tree (Note S3.2). The

calibrations from within bacteria were the 1.6 Ga-old akinete fossils for the filamentous

cyanobacteria Nostocales (38), the 1.7 Ga-old microfossils for the coccoid cyanobacteria

order Pleurocapsales (76), the GOE ~2.32 Ga for photosynthetic cyanobacteria (77), and the

hydrocarbon biomarkers for green and purple sulphur bacteria both dated at 1.64 Ga (78).

The root minimum was given as 2.32 Ga, the oldest among the above five. A maximum was

set as the age of Earth (4.5 Ga) for all the above calibrations within bacteria. If not otherwise

specified, in all molecular clock analyses we followed Betts et al. 2018 to apply a uniform

distribution as the prior: a soft tail was applied as the upper calibration bound, which means a

small probability (by default 2.5%) where the age is beyond the bound, and a hard bound was

set as the minimum age meaning that the age cannot be lower than the bound. This makes

sense as minimum bounds are based on solid fossil or biomarker records while the maxima

are often based on indirect evidence (8, 9).

A bootstrap-based approach to improving MCMCTree branch length estimate by

incorporating the mixture substitution model

We applied MCMCTree's approximate likelihood method (34, 79) which estimated the

branch lengths (expected number of amino acid substitutions per site) in the first step, and in

the second step, the branch length estimates, the gradient and the Hessian matrix are used to

calculate the likelihood by the second-order Taylor approximation to the likelihood function.

Different from previous studies which often use the LG+G substitution model, we applied the

bootstrap-based method developed in the present study to integrate the site-heterogeneous

profile mixture model LG+G+C60 (80) in MCMCTree analysis. We based our method on the

assumption that the maximum likelihood estimates of branch length follow a multivariate

normal (MVN) distribution, according to the maximum likelihood (ML) estimation theory

stating that ML estimators are asymptotically multivariate normally distributed (81). The idea

was originally proposed by Thorne et al. 1998 and later implemented in MCMCTree as the

approximate likelihood method (34). The approximate likelihood method (34) first obtains

the maximum-likelihood estimates (MLEs) of the branch length $\hat{\boldsymbol{\theta}} = (\hat{\theta}_1, \dots, \hat{\theta}_n)^T$, plus the

gradient $\mathbf{g}$ and the Hessian matrix $\mathbf{H}$, which are the first and second derivatives of the log-

likelihood evaluated at $\widehat{\boldsymbol{\theta}}$ respectively. Next, according to Eq. (1) in the study (34), Taylor

expansion at the second order derivative was applied to approximate the log-likelihood

579
$$\ell(\boldsymbol{\theta}) \approx \ell(\widehat{\boldsymbol{\theta}}) + \mathbf{g}^T \Delta\boldsymbol{\theta} + \frac{1}{2}\Delta\boldsymbol{\theta}^T \mathbf{H}\Delta\boldsymbol{\theta} \tag{6}$$

where $\boldsymbol{\theta} = (\theta_1, \theta_2, \dots, \theta_n)^T$ denotes the parameters to estimate (branch length) and $\ell(\boldsymbol{\theta})$ is the

log-likelihood as a function of branch lengths.

While the gradient is zero at the maximum likelihood estimate ($\mathbf{g} = \mathbf{0}$), directly

calculating $\mathbf{H}$ under IQ-Tree's profile mixture and other complex substitution model is

difficult and has not been available. However, note that when the sample size is large enough,

the Hessian matrix evaluated at $\widehat{\boldsymbol{\theta}}$ is approximately the negative inverse matrix of the

covariance matrix of the MLEs $\mathbf{V}(\widehat{\boldsymbol{\theta}})$ (81):

587
$$\mathbf{H} \approx -\mathbf{V}(\widehat{\boldsymbol{\theta}})^{-1}. \tag{7}$$

588   Hence, we instead applied a bootstrap method to estimate $\mathbf{V}(\widehat{\boldsymbol{\theta}})$ by the bootstrap covariance

589   matrix $\widehat{\mathbf{V}}(\widehat{\boldsymbol{\theta}}^*)$ (82) by taking B bootstrap alignments and estimating the MLEs of branch

590   lengths $\widehat{\boldsymbol{\theta}}^*$:

591
$$\mathbf{V}(\widehat{\boldsymbol{\theta}}) \approx \widehat{\mathbf{V}}(\widehat{\boldsymbol{\theta}}^*) = \frac{1}{B-1}\sum_{b=1}^{B}(\widehat{\boldsymbol{\theta}}^{(b)} - \overline{\widehat{\boldsymbol{\theta}}^*})(\widehat{\boldsymbol{\theta}}^{(b)} - \overline{\widehat{\boldsymbol{\theta}}^*})^T \tag{8}$$

where $\widehat{\boldsymbol{\theta}}^{(b)}$ indicate the MLEs of the branch lengths in the b^{th} bootstrap sample. Specifically,

1000 bootstraps ($B = 1000$) were performed using the site-heterogeneous profile mixture

model LG+G+Cxx with a fixed tree topology by IQ-Tree, therefore generating 1000

bootstrap MLEs for each branch length and allowing calculating the bootstrap covariance

matrix $\widehat{\mathbf{V}}(\widehat{\boldsymbol{\theta}}^*)$. Accordingly, the Hessian matrix can be approximated by solving the inverse

matrix of $-\widehat{\mathbf{V}}(\widehat{\boldsymbol{\theta}}^*)$. We imported the information of branch length estimates, i.e. the MLEs of

branch length, the gradient as a zero matrix, and the Hessian matrix approximated by the

above procedure, into a single file *in.BV*, which was used for time estimation using

MCMCTree's approximate likelihood method (34). A series of simulations were performed

to show the feasibility of this approach to integrating profile mixture substitution model into

MCMCTree analysis (see Note S4).

A taxonomically high-resolution timeline of the bacterial tree

A representative genome was downloaded from each genus deposited in the Genome

Taxonomy Database (GTDB; version r95) (83). We selected 14 out of the 25 orthologs

conserved across prokaryotes identified by a prior study (4), the other 11 being removed to

avoid squaring of likelihood using the sequential approach [otherwise the same genes are

used twice; see Note S1.2.2 and the studies (28, 29)]. These genes were then used to construct

a phylogenomic tree based on the first two codon positions of coding sequences using

GTR+G model and the "-fast" mode with IQ-Tree to reduce computational burden. We

applied RelTime implemented in MEGA X (84), which is based on estimating the relative

rates (85), to establish an evolutionary timescale for bacteria at the genus-level resolution.

The posterior dates obtained in the 148-genome dataset were approximated by a normal

(shown in Fig. 4) or uniform (in alternative analysis) distribution. Because RelTime

calculates divergence time by measuring the relative rate between neighbouring branches

(85) which is insensitive to overlaps in time priors between parent and child nodes (also

known as "truncation effect"; see Note S1.2.2), the age estimates using RelTime may not

change much from the specified calibrations. However, to minimize the impact of the overlap

in the calibrations between parent and child nodes, we alternatively removed those

calibrations where more than 10% overlap in the specified distributions between the parent

and child nodes to keep a maximum number of calibrations without significant overlap with

each other (Fig. S15D), and re-performed RelTime analysis (Fig. S15A-S15C). Taxonomic

classification was according to GTDB version r95 (86). Genetic divergence

(substitutions/site) between 16S rRNA sequences was calculated using IQ-Tree under

GTR+G+I model.

Data availability

The scripts to do pRTC and bootstrap-based approach hessian for MCMCTree are available

at https://github.com/evolbeginner/rrtc and https://github.com/evolbeginner/bs_inBV/

respectively. Other scripts and data are available at

https://figshare.com/s/1b39783f1ddda9797914.

Acknowledgements

We are particularly grateful to Jianhao Lv for the critical comments on the mathematical

derivation of the formulae, and the professional insights in statistics. We also thank

Daniel Mills and Philip Donoghue for insights in the interpretation of eukaryotic fossils,

Mario dos Reis, José Barba-Montoya and Suha Naser for technical support in phylogenetics

analysis, and Andrew Meade, Stilianos Louca and Liam Revell for guidance in ancestral state

reconstruction. We thank László Nagy, Torda Varga and Tomáš Flouris for kindly sharing

with us software. We also thank Luo Lab members Tianhua Liao and Hao Zhang for

discussion, and PhyloPic (https://www.phylopic.org/) for the free images of organisms. The

work is supported by the Hong Kong Research Grants Council General Research Fund

(14116922), the Natural Science Foundation of China (42293294), and the Hong Kong

Research Grants Council Area of Excellence Scheme (AoE/M-403/16).

References

1.   P. G. Falkowski, T. Fenchel, E. F. Delong, The microbial engines that drive earth's

biogeochemical cycles. *Science (80-.).* **320**, 1034–1039 (2008).

2.   S. Louca, *et al.*, Bacterial diversification through geological time. *Nat. Ecol. Evol.* **2**,

1458–1467 (2018).

3.   F. U. Battistuzzi, A. Feijao, S. B. Hedges, A genomic timescale of prokaryote

evolution: Insights into the origin of methanogenesis, phototrophy, and the

colonization of land. *BMC Evol. Biol.* **4**, 44 (2004).

4.   F. U. Battistuzzi, S. B. Hedges, A major clade of prokaryotes with ancient adaptations

to life on land. *Mol. Biol. Evol.* **26**, 335–343 (2009).

5.   E. Zuckerkandl, L. Pauling, "Evolutionary Divergence and Convergence in Proteins"

in *Evolving Genes and Proteins*, (1965), pp. 97–166.

6.   A. J. Drummond, S. Y. W. Ho, M. J. Phillips, A. Rambaut, Relaxed phylogenetics and

dating with confidence. *PLoS Biol.* **4**, 699–710 (2006).

7.   Z. Yang, B. Rannala, Bayesian estimation of species divergence times under a

molecular clock using multiple fossil calibrations with soft bounds. *Mol. Biol. Evol.*

(2006) https:/doi.org/10.1093/molbev/msj024.

8.   C. R. Marshall, Using the Fossil Record to Evaluate Timetree Timescales. *Front.*

Genet. **10**, 1049 (2019).

9.   M. J. Benton, P. C. J. Donoghue, Paleontological evidence to date the tree of life. *Mol.*

Biol. Evol. **24**, 26–53 (2007).

10.   E. R. R. Moody, *et al.*, An estimate of the deepest branches of the tree of life from

ancient vertically evolving genes. *Elife* **11**, e66695 (2022).

11.   S. Wang, H. Luo, Dating Alphaproteobacteria evolution with eukaryotic fossils. *Nat.*

Commun. **12**, 3324 (2021).

12.   R. Gouy, D. Baurain, H. Philippe, Rooting the tree of life: The phylogenetic jury is

673   still out. *Philos. Trans. R. Soc. B Biol. Sci.* **370** (2015).

674   13.   P. Sánchez-Baracaldo, J. A. Raven, D. Pisani, A. H. Knoll, Early photosynthetic

675   eukaryotes inhabited low-salinity habitats. *Proc. Natl. Acad. Sci.* (2017)

676   https:/doi.org/10.1073/pnas.1620089114.

677   14.   P. M. Shih, J. Hemp, L. M. Ward, N. J. Matzke, W. W. Fischer, Crown group

678   Oxyphotobacteria postdate the rise of oxygen. *Geobiology* **15**, 19–29 (2017).

679   15.   C. Toft, S. G. E. Andersson, Evolutionary microbial genomics: Insights into bacterial

680   host adaptation. *Nat. Rev. Genet.* **11**, 465–475 (2010).

681   16.   J. Perreau, N. A. Moran, Genetic innovations in animal–microbe symbioses. *Nat. Rev.*

682   *Genet.* (2021) https:/doi.org/10.1038/s41576-021-00395-z.

683   17.   C. Magnabosco, K. R. Moore, J. M. Wolfe, G. P. Fournier, Dating phototrophic

684   microbial lineages with reticulate gene histories. *Geobiology* **16**, 179–189 (2018).

685   18.   A. A. Davín, *et al.*, Gene transfers can date the tree of life. *Nat. Ecol. Evol.* **2**, 904–909

686   (2018).

687   19.   M. Ravenhall, N. Škunca, F. Lassalle, C. Dessimoz, Inferring Horizontal Gene

688   Transfer. *PLoS Comput. Biol.* **11**, e1004095 (2015).

689   20.   J. Martijn, J. Vosseberg, L. Guy, P. Offre, T. J. G. Ettema, Deep mitochondrial origin

690   outside the sampled alphaproteobacteria. *Nature* **557**, 101–105 (2018).

691   21.   S. A. Muñoz-Gómez, *et al.*, Site-and-branch-heterogeneous analyses of an expanded

692   dataset favour mitochondria as sister to known Alphaproteobacteria. *Nat. Ecol. Evol.* **6**,

693   253–262 (2022).

694   22.   G. J. Szöllõsi, *et al.*, Relative Time Constraints Improve Molecular Dating. *Syst. Biol.*

695   **71**, 797–809 (2022).

23.   J. P. McCutcheon, N. A. Moran, Extreme genome reduction in symbiotic bacteria. *Nat.*

Rev. Microbiol. **10**, 13–26 (2012).

24. G. A. Coleman, *et al.*, A rooted phylogeny resolves early bacterial evolution. *Science*

(80-.). **372** (2021).

25. S. Naser-Khdour, B. Quang Minh, R. Lanfear, Assessing Confidence in Root

Placement on Phylogenies: An Empirical Study Using Nonreversible Models for

Mammals. *Syst. Biol.* **71**, 959–972 (2022).

26. L. A. Hug, *et al.*, A new view of the tree of life. *Nat. Microbiol.* **1** (2016).

27. Q. Zhu, *et al.*, Phylogenomics of 10,575 genomes reveals evolutionary proximity

between domains Bacteria and Archaea. *Nat. Commun.* **10** (2019).

28. S. Álvarez-Carretero, *et al.*, A species-level timeline of mammal evolution integrating

phylogenomic data. *Nature* **602**, 263–267 (2022).

29. M. dos Reis, *et al.*, Phylogenomic datasets provide both precision and accuracy in

estimating the timescale of placental mammal phylogeny. *Proc. R. Soc. B Biol. Sci.*

279, 3491–3500 (2012).

30. J. F. H. Strassert, I. Irisarri, T. A. Williams, F. Burki, A molecular timescale for

eukaryote evolution with implications for the origin of red algal-derived plastids. *Nat.*

Commun. **12**, 1879 (2021).

31. L. W. Parfrey, D. J. G. Lahr, A. H. Knoll, L. A. Katz, Estimating the timing of early

eukaryotic diversification with multigene molecular clocks. *Proc. Natl. Acad. Sci. U.*

S. A. **108**, 13624–13629 (2011).

32. H. C. Betts, *et al.*, Integrated genomic and fossil evidence illuminates life's early

evolution and eukaryote origin. *Nat. Ecol. Evol.* **2**, 1556–1562 (2018).

33. T. M. Gibson, *et al.*, Precise age of Bangiomorpha pubescens dates the origin of

eukaryotic photosynthesis. *Geology* **46**, 135–138 (2018).

34. M. dos Reis, Z. Yang, Approximate likelihood calculation on a phylogeny for

Bayesian Estimation of Divergence Times. *Mol. Biol. Evol.* **28**, 2161–2172 (2011).

35.   E. G. Nisbet, N. H. Sleep, The habitat and nature of early life. *Nature* **409**, 1083–1091

(2001).

36.   S. Y. W. Ho, S. Duchêne, Molecular-clock methods for estimating evolutionary rates

and timescales. *Mol. Ecol.* **23**, 5947–5965 (2014).

37.   S. Bengtson, T. Sallstedt, V. Belivanova, M. Whitehouse, Three-dimensional

preservation of cellular and subcellular structures suggests 1.6 billion-year-old crown-

group red algae. *PLoS Biol.* **15**, e2000735 (2017).

38.   A. Tomitani, A. H. Knoll, C. M. Cavanaugh, T. Ohno, The evolutionary diversification

of cyanobacteria: Molecular-phylogenetic and paleontological perspectives. *Proc.*

Natl. Acad. Sci. U. S. A. **103**, 5442–5447 (2006).

39.   B. Amard, J. Bertrand-Sarfati, Microfossils in 2000 Ma old cherty stromatolites of the

Franceville Group, Gabon. *Precambrian Res.* **81**, 197–221 (1997).

40.   D. W. Waite, *et al.*, Proposal to reclassify the proteobacterial classes

deltaproteobacteria and oligoflexia, and the phylum thermodesulfobacteria into four

phyla reflecting major functional capabilities. *Int. J. Syst. Evol. Microbiol.* **70**, 5972–

6016 (2020).

41.   N. A. Moran, M. A. Munson, P. Baumann, H. Ishikawa, A molecular clock in

endosymbiotic bacteria is calibrated using the insect hosts. *Proc. R. Soc. B Biol. Sci.*

253, 167–171 (1993).

42.   H. Ochman, S. Elwyn, N. A. Moran, Calibrating bacterial evolution. *Proc. Natl. Acad.*

Sci. U. S. A. **96**, 12638–12643 (1999).

43.   C. H. Kuo, H. Ochman, Inferring clocks when lacking rocks: The variable rates of

molecular evolution in bacteria. *Biol. Direct* **4**, 35 (2009).

44.   A. H. Knoll, Paleobiological perspectives on early microbial evolution. *Cold Spring*

Harb. Perspect. Biol. **7**, 1–17 (2015).

45.   G. P. Fournier, *et al.*, The Archean origin of oxygenic photosynthesis and extant

cyanobacterial lineages. *Proc. R. Soc. B Biol. Sci.* **288**, 20210675 (2021).

46.   J. J. Schenk, Consequences of secondary calibrations on divergence time estimates.

PLoS One **11**, e0148228 (2016).

47.   S. Kumar, G. Stecher, M. Suleski, S. B. Hedges, TimeTree: A Resource for Timelines,

Timetrees, and Divergence Times. *Mol. Biol. Evol.* **34**, 1812–1819 (2017).

48.   J. Marin, F. U. Battistuzzi, A. C. Brown, S. B. Hedges, The Timetree of Prokaryotes:

New Insights into Their Evolution and Speciation. *Mol. Biol. Evol.* **34**, 437–446

(2017).

49.   O. Abramov, S. J. Mojzsis, Microbial habitability of the Hadean Earth during the late

heavy bombardment. *Nature* **459**, 419–422 (2009).

50.   P. Boehnke, T. M. Harrison, Illusory late heavy bombardments. *Proc. Natl. Acad. Sci.*

U. S. A. **113**, 10802–10806 (2016).

51.   P. Sánchez-Baracaldo, T. Cardona, On the origin of oxygenic photosynthesis and

Cyanobacteria. *New Phytol.* **225**, 1440–1446 (2020).

52.   A. C. Allwood, M. R. Walter, B. S. Kamber, C. P. Marshall, I. W. Burch, Stromatolite

reef from the Early Archaean era of Australia. *Nature* **441**, 714–718 (2006).

53.   H. J. Hofmann, K. Grey, A. H. Hickman, R. I. Thorpe, Origin of 3.45 Ga coniform

stromatolites in Warrawoona Group, Western Australia. *Bull. Geol. Soc. Am.* **111**,

1256–1262 (1999).

54.   J. W. Schopf, The paleobiological record of photosynthesis. *Photosynth. Res.* **107**, 87–

101 (2011).

55.   N. McLoughlin, L. A. Wilson, M. D. Brasier, Growth of synthetic stromatolites and

wrinkle structures in the absence of microbes - Implications for the early fossil record.

Geobiology **6**, 95–105 (2008).

56.   S. A. Crowe, *et al.*, Atmospheric oxygenation three billion years ago. *Nature* **501**,

774   535–538 (2013).

775   57.   X. Wang, *et al.*, A Mesoarchean shift in uranium isotope systematics. *Geochim.*

776   *Cosmochim. Acta* **238**, 438–452 (2018).

777   58.   N. J. Planavsky, *et al.*, Evidence for oxygenic photosynthesis half a billion years

778   before the Great Oxidation Event. *Nat. Geosci.* **7**, 283–286 (2014).

779   59.   H. He, *et al.*, An abiotic source of Archean hydrogen peroxide and oxygen that pre-

780   dates oxygenic photosynthesis. *Nat. Commun.* (2021) https:/doi.org/10.1038/s41467-

781   021-26916-2.

782   60.   J. Jabłońska, D. S. Tawfik, The evolution of oxygen-utilizing enzymes suggests early

783   biosphere oxygenation. *Nat. Ecol. Evol.* **5**, 442–448 (2021).

784   61.   B. Kraft, *et al.*, Oxygen and nitrogen production by an ammonia-oxidizing archaeon.

785   *Science (80-.).* (2022) https:/doi.org/10.1126/science.abe6733.

786   62.   K. F. Ettwig, *et al.*, Nitrite-driven anaerobic methane oxidation by oxygenic bacteria.

787   *Nature* (2010) https:/doi.org/10.1038/nature08883.

788   63.   K. Katoh, D. M. Standley, MAFFT Multiple Sequence Alignment Software Version 7:

789   Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

790   64.   S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: A tool for automated

791   alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–

792   1973 (2009).

793   65.   B. Q. Minh, *et al.*, IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

794   Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

795   66.   T. Liao, S. Wang, E. E. Stüeken, H. Luo, Phylogenomic Evidence for the Origin of

796   Obligate Anaerobic Anammox Bacteria Around the Great Oxidation Event. *Mol. Biol.*

797   *Evol.* **39**, msac170 (2022).

798   67.   D. H. Parks, *et al.*, Recovery of nearly 8,000 metagenome-assembled genomes

799   substantially expands the tree of life. *Nat. Microbiol.* **2**, 1533–1542 (2017).

800   68.   K. Lagesen, *et al.*, RNAmmer: Consistent and rapid annotation of ribosomal RNA

801   genes. *Nucleic Acids Res.* **35**, 3100–3108 (2007).

802   69.   W. Z. Li, A. Godzik, Cd-hit: a fast program for clustering and comparing large sets of

803   protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

804   70.   P. Yarza, *et al.*, Uniting the classification of cultured and uncultured bacteria and

805   archaea using 16S rRNA gene sequences. *Nat. Rev. Microbiol.* **12**, 635–645 (2014).

806   71.   J. P. Huelsenbeck, R. Nielsen, J. P. Bollback, Stochastic Mapping of Morphological

807   Characters. *Syst. Biol.* **52**, 131–158 (2003).

808   72.   L. J. Revell, phytools: An R package for phylogenetic comparative biology (and other

809   things). *Methods Ecol. Evol.* **3**, 217–223 (2012).

810   73.   M. Pagel, A. Meade, D. Barker, Bayesian estimation of ancestral character states on

811   phylogenies. *Syst. Biol.* **53**, 673–684 (2004).

812   74.   S. Wang, A. Meade, H.-M. Lam, H. Luo, Evolutionary Timeline and Genomic

813   Plasticity Underlying the Lifestyle Diversity in Rhizobiales. *mSystems* **5**, e00438-20

814   (2020).

815   75.   Z. Yang, PAML 4: Phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**,

816   1586–1591 (2007).

817   76.   Y. Zhang, S. Golubic, Endolithic microfossils (cyanophyta) from early Proterozoic

818   stromatolites, Hebei, China. *Acta Micropalaeontologica Sin.* **4**, 1–3 (1987).

819   77.   G. Luo, *et al.*, Rapid oxygenation of Earth's atmosphere 2.33 billion years ago. *Sci.*

820   *Adv.* **2** (2016).

78.   J. J. Brocks, *et al.*, Biomarker evidence for green and purple sulphur bacteria in a

stratified Palaeoproterozoic sea. *Nature* **437**, 866–870 (2005).

79.   J. L. Thorne, H. Kishino, I. S. Painter, Estimating the rate of evolution of the rate of

molecular evolution. *Mol. Biol. Evol.* **15**, 1647–1657 (1998).

80.   L. S. Quang, O. Gascuel, N. Lartillot, Empirical profile mixture models for

phylogenetic reconstruction. *Bioinformatics* **24**, 2317–2323 (2008).

81.   Y. Pawitan, *In all likelihood: statistical modelling and inference using likelihood*

(Oxford University Press, 2001).

82.   B. Efron, "The Bootstrap" in *The Jackknife, the Bootstrap and Other Resampling*

Plans, (1982), p. 36.

83.   D. H. Parks, *et al.*, GTDB: An ongoing census of bacterial and archaeal diversity

through a phylogenetically consistent, rank normalized and complete genome-based

taxonomy. *Nucleic Acids Res.* **50**, D785–D794 (2022).

84.   S. Kumar, G. Stecher, M. Li, C. Knyaz, K. Tamura, MEGA X: Molecular evolutionary

genetics analysis across computing platforms. *Mol. Biol. Evol.* **35**, 1547–1549 (2018).

85.   K. Tamura, *et al.*, Estimating divergence times in large molecular phylogenies. *Proc.*

Natl. Acad. Sci. U. S. A. (2012) https:/doi.org/10.1073/pnas.1213199109.

86.   D. H. Parks, *et al.*, A complete domain-to-species taxonomy for Bacteria and Archaea.

Nat. Biotechnol. (2020) https:/doi.org/10.1038/s41587-020-0501-8.

 Figure legends

Figure 1. Graphical illustration of the two new methods in molecular dating developed in the

present study. (A) Procedure of the pRTC dating informed by host-symbiont association.

Suppose that there are two symbiotic lineages, Symbiont1 from non-Proteobacteria (blue) and

Symbiont2 from α-Proteobacteria (purple), and most of their members are mammal

symbionts (orange) but some are free-living (yellow). First, ASR is performed on the

phylogram of 16S rRNA tree to obtain the probability of each ancestral state taken by the

LCA of the two "symbiont" clades respectively, i.e., $P(S_1)$ and $P(S_2)$. Second, a phylogeny

consisting of symbionts and mammals is constructed by mitochondria-originated genes, and

MCMCTree is run with fossil calibrations but without RTCs. Third, rejection sampling is

performed to remove posterior timetrees obtained in the previous step based on the

requirement that symbiont's origin must postdate the host. Specifically, we require that the

age of the crown group of the symbionts (node1 and node2) no older than that of the total-

group mammals (node3), weighted by $P(S_i = mammal)$. The use of mammals' total group

is because there could be stem-group mammals that are unsampled or go extinct but that have

Symbiont1 or Symbiont2 as symbiont. For the free-living state, we define $I(t_i, t_{fl}) = 1$ as

the RTC is automatically satisfied for free-living bacteria (see Methods). In the final timetree,

the divergence time of symbiont lineages becomes younger than that inferred before host-

association is considered, suggesting that incorporating RTC informed by host-association

improves the molecular clock inferences. (B) Overflow of the bootstrap-based approach to

incorperating branch length estimates inferred under site-heterogeneous profile mixture

substitution models (LG+G+Cxx). $\boldsymbol{\theta} = (\theta_1, \theta_2, \dots, \theta_n)^T$ denotes the parameters to estimate

(branch length, i.e., expected number of amino acid substitutions per site) and $\ell(\boldsymbol{\theta})$ is the log-

likelihood as a function of branch lengths. Briefly, the maximum-likelihood estimate (MLE)

of branch length $\hat{\boldsymbol{\theta}}$ is obtained by LG+G+Cxx, and the bootstrap covariance matrix of branch

lengths $\mathbf{V}(\hat{\boldsymbol{\theta}}^*)$ is calculated to approximate the hessian matrix (second-order derivative of the

log-likelihood of branch length), which are further used for time estimation by MCMCTree.

Figure 2. An evolutionary timeline of the bacterial tree of life established by the three-step

pRTC procedure. The root position is determined using IQ-Tree's non-reversible substitution

model in maximum-likelihood phylogenomic tree reconstruction using the 19 mitochondria-

originated genes. Node bars indicate the 95% HPD interval of age estimates. Nodes with

orange circles indicate the bacterial calibration points (including the root). Eukaryotic

calibrations obtained from sequential Bayesian dating are given in Fig. S4. RTCs of the 28

bacterial symbionts and five eukaryotic hosts (LCA of eukaryotes, total-group animals, total-

group ciliates, total-group insects, crown-group fungi) are denoted by pink and yellow

circles, respectively. Because the 28 symbionts are selected from "symbiont" clades, a total

of 28-9=19 internal nodes serve as the RTCs for bacteria. The organism images are placed at

the origin times of eukaryotes of interest co-estimated with bacterial lineages. Detailed

posterior ages for clades of interest are summarised in Table S4. DST: Deinococcota,

Synergistota, and Thermotogota; Armat-Eremi: Armatimonadetes and Eremiobacteraeota.

CPR: Candidate phyla radiation; FCB: Fibrobacterota, Chlorobiota, Bacteroidota; PVC:

Planctomycetota, Verrucomicrobiota, Chlamydiota, and related lineages; ACD: Aquificota,

Campylobacterota, and Deferribacterota. MBDD (δ-Proteobacteria): Myxococcota,

Bdellovibrionota, Desulfomonadota, and Desulfobacterota. The images of the choanozoan,

and the cyanobacteria and mitochondria are credited to Urutseg and Kevin Song respectively

under CC BY-SA 3.0 (https://creativecommons.org/licenses/by-sa/3.0/) with slight

modifications. All other organism images are distributed under CC0.

Figure 3. Assessing the uncertainty in Bayesian molecular clock analysis. (A) Comparison of the posterior mean ages (lower triangle; unit: Ma) and rates (upper triangle; unit: number of substitutions per site per Ga) of bacteria estimated with different strategies. *Strategy 1* (traditional strategy used in prior studies): five bacterial calibrations with hard minimum bounds and a soft maximum bound <4.5 Ga at LBCA, no eukaryote timing information, substitution model LG+G, no RTCs. *Strategy 2*: bacterial calibrations, eukaryote times by sequential molecular dating, substitution model LG+G, no RTCs. *Strategy 3*: bacterial calibrations, eukaryote times by sequential molecular dating, substitution model LG+G+C60, no RTCs. Focal: bacterial calibrations, eukaryote times by sequential molecular dating, substitution model LG+G+C60, with RTCs. (B) Posterior mean ages of the selected clades calculated under root maximum prior ages from 4500 to 6000 Ma with the focal strategy (dashed line) and Strategy 1 (solid line). (C) Divergence times of bacteria estimated by alternative schemes (*y*-axis; Data S3) versus the one in the focal strategy (*x*-axis). The blue bars denote the 95% HPD. *Euk1*: the minimum time bound of crown-group red algae alternatively set as 1.6 Ga based on *Rafatazmia*; *Euk2*: the minimum time bound of animal crown group alternatively set as 0.89 Ga (Note S3.2); *noCyanoFossil*: the two cyanobacteria minimum time bounds (total-group Nostocales and Pleurocapsales) removed; *noBiomarker*: the two bacterial biomarker minimum time bounds (total-group Chromatiaceae and Chlorobi) removed; *noGOE*: the minimum time bound based on GOE removed. *Joint_prob*: rejection sampling based on calculating the joint probability (see Fig. S6B) in Eq. (2). *Alt_lifestyle*: ASR inferred with alternative classification of lifestyles of modern symbionts (Note S3.1). *OTU97*: ASR performed with OTU at a cut-off of 97% sequence identity. *ASR_ML*: maximum likelihood algorithm instead of MCMC used in ASR. *IR*: the independent rate (IR) model is used instead of the auto-correlated rate (AR) model. (D) Changes in the posterior dates shown as 95% HPD interval resulting from different root positions of the

phylogenomics tree (*root1*, *root2*, and *rootCPR*), and the use of different genes. *rate_top* and

deltaLL_top: the 5, 10, and 15 (best) top-ranking genes according to relative rate difference

between mitochondrial and bacterial lineages or $\triangle$LL which measures the the degree of

species- and gene-tree incongruence (Note S1.2.4). *all_genes*: molecular dating performed on

all 32 mitochondria-originated genes conserved across bacteria. The centre is the posterior

mean age. FA: Firmicutes and Actinomycetes. MBDD (δ-Proteobacteria): Myxococcota,

Bdellovibrionota, Desulfomonadota, and Desulfobacterota.

Figure 4. A high-taxonomy resolution timetree of the bacterial tree of life. (A) Divergence

times of the 8898-tip bacterial tree, consisting of a randomly selected representative from all

genera according to GTDB r95, estimated using RelTime and 118 calibrations (black dots),

i.e., posterior dates calculated from the 148-genome molecular clock analysis. Ma: million

years ago. (B) The divergence times at a family-, order-, class-, and phylum level inferred

from the current study. Only families, orders, classes, and phyla with at least five members

are considered. (C) A simple linear regression analysis of the relationship between 16S rRNA

divergence (number of substitutions per site; *y*-axis) and absolute divergence time (*x*-axis) at

a family level. The intercept is fixed to zero. Shaded areas indicate the 95% confidence

interval of the estimated slope. For each family with at least five members, the largest

difference in 16S sequence between any of its members is considered the 16S divergence of

the family. The analysis is not performed at a higher taxonomic level as 16S sequence likely

reaches saturation due to the large divergence time. Detailed results of this simple linear

regression and alternative linear models are provided in Data S5 and Note S1.4.

A Figure 4

Terrabacteria
- Fusobacteriota/DST
- Actinomycetes
- Cyano-Margulis
- Chloroflexota
- Armat-Eremi
- CPR

Gracilicutes
- Elusimicrobiota
- Spirochaetota
- PVC
- FCB
- Acidobacteriota
- ACD
- Nitrospirae
- MMBD (δ-Proteobacteria)
- Proteobacteria

Ma

γ
β
α

B

time unit (Ma)

family   order   class   phylum
taxonomy level

C

16S rRNA divergence (substitutions/site)

$y=0.000156x$

phylum
- Actinobacteriota
- Cyanobacteria
- Firmicutes
- CPR
- Proteobacteria
- Others

Divergence time (Ma)