DNA conserved in diverse animals since the Precambrian controls genes for embryonic development

Martin C. Frith[1,2,3] and Shengliang Ni[2]

[1]Artificial Intelligence Research Center, AIST
[2]Graduate School of Frontier Sciences, University of Tokyo
[3]Computational Bio Big Data Open Innovation Laboratory, AIST

Summary

DNA that controls gene expression (e.g. enhancers, promoters) has seemed almost never to be conserved between distantly-related animals, like vertebrates and arthropods. This is mysterious, because development of such animals is partly organized by homologous genes with similar complex expression patterns, termed "deep homology".

Here we report twenty-five regulatory DNA segments conserved across bilaterian animals, of which seven are also conserved in cnidaria (coral and sea anemone). They control developmental genes (e.g. *Nr2f*, *Ptch*, *Rfx1/3*, *Sall*, *Smad6*, *Sp5*, *Tbx2/3*), including six homeobox genes: *Gsx*, *Hmx*, *Meis*, *Msx*, *Six1/2*, and *Zfhx3/4*. The segments contain perfectly or near-perfectly conserved CCAAT boxes, E-boxes, and other sequences recognized by regulatory proteins. More such DNA conservation will surely be found soon, as more genomes are published and sequence comparison is optimized. This reveals a control system for animal development conserved since the Precambrian.

Introduction

Genes often remain similar across vast gulfs of evolution. For example, the genes that encode ribosomal RNA (rRNA) in humans and bacteria have recognizably-similar DNA sequences. Gene expression is controlled by DNA segments near the genes, which curiously lack such long-term conservation. Some are conserved across vertebrate animals, but few have been found conserved between vertebrates and invertebrates [1–3].

Early animal evolution produced bilaterian animals, which then split into two superphyla: protostomes and deuterostomes (fig. 1). A pioneering study by Royo et al. found several regulatory DNA segments conserved across deuterostomes, but found none of them in any non-deuterostome, except, amazingly, two in sea anemone, which is not even bilaterian [4]. These two are enhancers of the *Sox21* and *Hmx* genes. Soon after, Clarke et al. found two regulatory elements conserved between deuterostomes and protostomes [5]: an enhancer of the *Id* gene in gastropods (but no other protostomes), and an enhancer of *Znf503* in tick (but no other protostomes). Beyond this handful of exceptions, it seems that gene-regulating DNA is not conserved across bilaterians, suggesting that their developmental programs are not conserved either.

Phylum	Gsx enhancer	Hmx enhancer	Meis promoter	Msx enhancer	Six1 enhancer	Six2 promoter	Zfhx3/4 promoter	Coe intron element	Nr2f promoter	Rfx1/3 promoter	Sall promoter	Smad6 promoter	Sox21 enhancer	Sp5 promoter	Tbx2/3 promoter	Tbx3 enhancer	Znf207 3'-UTR	Cpeb2/3/4 3'-UTR	Id enhancer	Lin28b promoter	Mex3 3'-UTR	Ptch promoter	Ssbp2 promoter	Znf503 enhancer	Znf503/703 enhancer	Animal
chordate	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	human
chordate	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	chimaera
chordate	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	lancelet
hemichordate	○	○	○	○		○	○		○	○	○	○	○	○	○	○	○		○		○	○	○	○	○	acorn worm
echinoderm				○			○		○	○	○	○	○	○		○			○				○	○		sea urchin
arthropod																		○								lobster
arthropod									○	○		○											○			centipede
arthropod									○	○		○											○	○		millipede
arthropod									○		○	○										○	○	○		scorpion
arthropod										○		○				○							○	○		tick
arthropod									○	○	○	○		○								○	○	○		horseshoe crab
arthropod									○																	sea spider
onychophora	○				○	○		○	○	○		○		○	○				○	○	○		○	○	○	velvet worm
priapulid	○								○	○		○				○		○	○		○			○		cactus worm
mollusc	○							○	○	○		○							○		○	○	○			chiton
mollusc			○							○		○							○		○	○				sea hare
mollusc	○	○	○		○	○	○		○	○		○		○	○				○		○	○	○	○		abalone
mollusc					○				○	○									○		○		○			owl limpet
mollusc		○			○				○	○		○							○			○	○			oyster
mollusc	○								○	○		○	○						○		○		○			nautilus
mollusc	○									○		○							○		○		○			octopus
mollusc										○																solenogaster
annelid	○					○			○	○		○	○						○				○			peanut worm
annelid						○			○	○		○							○				○			sandworm
annelid				○		○			○	○		○							○		○		○			Satsuma tubeworm
annelid									○	○				○					○				○			shingle tubeworm
brachiopod	○	○	○	○	○	○			○	○		○				○		○	○		○	○	○	○	○	shamisen shell
phoronid	○									○		○							○				○			horseshoe worm
nemertea	○								○	○		○	○	○					○				○			bootlace worm
cnidaria			○																							carnation coral
cnidaria		○	○						○			○	○	○												stony coral
cnidaria		○	○						○			○	○													lace coral
cnidaria		○							○																	starlet sea anemone
cnidaria		○	○						○					○							○					pale anemone

Figure 1: Animals in which each regulatory DNA element was detected. Regulatory segments are colored according to their nearby gene: homeobox genes red, other transcription factors blue. Each animal's phylum is written on the left, deuterostomes in purple and protostomes in green.

Regulatory DNA conserved across vertebrates or beyond tends to regulate genes that control embryonic development [2]. These genes often encode transcription factors: proteins that control other genes by interacting with their regulatory DNA. One family of transcription factor genes contain homeobox sequences, which encode a DNA-binding structure called the homeodomain: these are especially used in development. Development involves the appearance of different territories in the embryo that express different combinations of transcription factors, and then develop into different body parts. These territories form in response to varying concentrations of signalling molecules, such as BMP, Wnt, and hedgehog proteins, that are secreted from cells. Often, one gene contributes to many parts of development, presumably because development evolved by redeploying these genes.

It is curious that developmental genes are widely conserved across animals, but their regulatory DNA rarely is. One theory is that conserved regulatory segments correspond to conserved body plans such as vertebrate [1]. This does not explain, however, why the expression patterns of developmental genes are often deeply conserved across animals [2]. One possible answer is that regulatory function can be conserved even as the DNA evolves so much that sequence similarity is lost [6].

Here, we find many more DNA segments that control developmental gene expression and are conserved across bilaterian animals, and even in non-bilaterian corals and sea anemone. They were found by exploiting the wealth of recently-published animal genomes, and by optimizing DNA homology detection [7]. This reveals a system of DNA sequences conserved since the Precambrian that control animal development.

Results

The first step was to find conserved DNA segments between the genomes of human and chimaera, a cartilaginous fish. These genomes are related closely enough that many conserved segments can be found, but distantly enough that most of their DNA lacks similarity. Another reason for using chimaera is that its genome has evolved slowly [8]. Next, conserved regions were removed if they encode protein, rRNA, tRNA, snRNA, snoRNA, or miRNA. Pseudogenes were also removed, i.e. DNA that used to encode protein, rRNA, tRNA, etc. (Such DNA might also regulate gene expression, but its conservation can be explained by what it does or did encode.) The remaining conserved regions cover 5.7 million base-pairs in chimaera: 1/174-th of the genome.

The next step was to seek conserved segments between these chimaera regions and various invertebrate genomes. By not using the whole chimaera genome, we get 174-fold fewer false matches, so we can find 174-fold weaker similarities. Each similarity was given an E-value, which means the number of times such a similarity would be expected between random sequences of shuffled bases. More specifically, between random sequences with 40% g+c and the same lengths as: the one invertebrate genome, and all the chimaera regions. Similarities were accepted with either E-value ≤ 0.0001, or E-value ≤ 10 and near homologous genes. For example, a chimaera segment near the *smad6* gene matched octopus DNA with E-value 0.46, and the nearest gene in octopus is the homolog of *smad6*.

As a negative control, each search was repeated after reversing (but not complementing) the invertebrate genome. DNA doesn't evolve by reversal, because that would require flipping the $3'$-to-$5'$ bonds between all the nucleotides. In most cases a reversed genome had about 10 matches with E-value ≤ 10, and the lowest E-value ever seen was 0.0009.

Further sensitivity was gained by using conserved parts of invertebrate genomes. For example, by using oyster DNA segments that are conserved in scallop (1/59-th of the oyster genome), sensitivity

was boosted a further 59-fold, revealing matches near the *Ptch* and *Meis* genes.

These methods found 25 DNA segments with similarity between vertebrates and non-deuterostome animals (fig. 1). They can be explained by either common ancestry or convergent evolution. It seems implausible that most are due to convergent evolution, because they are in a wide range of animals (fig. 1, 2) and they preserve position and orientation relative to the genes (fig. 3). The segments were not found in some invertebrates (fig. 1): either they were lost during evolution of those animals, or they are present but undetected. The segments are labelled "promoter" if they occur at a gene's transcription start site in human (the best-annotated genome), else "enhancer". One segment lies in an intron, and three lie in 3′-UTRs (UnTranslated Regions of exons, downstream of the translated protein-coding region).

Interestingly, all of these genes, except *Znf207*, have multiple (2–4) copies in genomes of jawed vertebrates. For example, jawed vertebrates have three similar *Meis* genes (*Meis1*, *Meis2*, and *Meis3*), two *SoxB2* genes (called *Sox14* and *Sox21*), two *sine oculis* genes (*Six1* and *Six2*), and two *NET* genes (*Znf503* and *Znf703*). *Sp5* is a semi-exception: vertebrates have two copies, named *Sp5* and *Sp5-like*, but *Sp5-like* was lost in mammals and birds [9]. In early vertebrate evolution, there were two rounds of whole-genome duplication. Most gene duplicates were rapidly lost, so that vertebrates do not have many more genes than invertebrates, but some genes remain in 2–4 copies, including nearly all the ones here.

The conserved DNA segments were detected at all, some, or just one of these multi-copy genes. For example, the conserved *Gsx* enhancer was found near both *Gsx* genes in human and chimaera (fig. 2, 3), but the *SoxB2* enhancer was only found at *Sox21*. The *sine oculis* enhancer was only found at *Six1*, whereas the *sine oculis* promoter was only found at *Six2*. One *NET* enhancer was found at both *Znf503* and *Znf703*, the other only at *Znf503*. While "not found" does not necessarily mean "absent", it seems that duplication of these genes enabled divergence of their regulation and developmental roles.

The functions of these genes and conserved DNA segments are partially known. *Gsx* and *Msx* encode transcription factors that define territories in early development of the CNS (central nervous system), in a manner conserved across bilaterian animals [10–12]. They produce an ancient sensory/inter/motor neuron system. *Msx* is expressed at the lateral edges of the CNS, and produces Rohon–Beard mechanosensory neurons. *Gsx* is expressed more medially and produces interneurons, and the most medial territory produces motor neurons. The *Gsx* enhancer seems to contribute by preventing lateral expression of *Gsx*, thus defining the brain's pallium-subpallium boundary [13, 14]. *Msx* plays many other roles in development: the enhancer is the previously-identified *Msx1* "proximal element" that produces expression in mouse embryo nasal epithelium, myotome, limb mesenchyme, eye, ear, roof plate, second arch, genital ridge and epiphysis [15].

The *Six1/2* genes control development of eyes and other cranial sensory organs (e.g. olfactory epithelium, inner ear, taste papillae), and also contribute to development of kidney, thymus, parathyroid gland, and more [16]. The *Six1* enhancer is the previously-described Six1-13 element, which produced variable expression patterns of unclear importance [16]: it would be interesting to re-examine this element.

Tbx2/3 control formation of the brain's neurohypophysis [17], and contribute to development of vertebrate limbs, heart, liver, and more [18]. The *Tbx3* enhancer was shown to affect expression of *Tbx3* and limb length in mice and horses [19].

The *Coe* genes encode transcription factors affecting subpallium development, expressed in *Gsx*+ brain regions of mice and annelids [10]. They also define subtypes of body-wall motor neuron in mice and roundworms [20], and promote development of fat cells [21]. The conserved DNA segment is in

```
tagcaacagttgccccggtgagggaaacggaggcgccatagccacggtagtc  human RFX1
tagcaaccgttgccttgtgaat------caagcgccatagtcaccgtagtc   human RFX3
tagcaactgttgccttgaggac--------gagcgccatagccaccgtagtc  chimaera rfx1
tagcaactgttgccttgtgact--------cagcgccatagtcaccgtagtc  chimaera rfx3
tggcaacagttgccttggttgt-------gtgcgccatagccaccgtagtc   lancelet
tggcaactgttgccttgagaat--------aaacgccatagccaccgtagtc  acorn worm
tagcaacagttgccttgagaat--------aaacgccatagtcaccgtagtc  sea urchin
tggcaactgttgccttaggaat-------ccgacgccatagacacggtagct  centipede
tggcaactgttgccctaggaat-------aagcgccatagacacggtaact   millipede
tggcaactgttgcccttggagt--------caacgccataactacggaagct  tick
tggcaactgttgcccttggaat-------caacgccataccgacggtaacc   horseshoe crab
tggcaacagttgccttaggaat--------aaacgccatagccaccgtagct  velvet worm
tggcaactgttgcccaagg----------aaacgccatagacacggtagct   cactus worm
tggcaacagttgccgtaggaat-------aaacgccatagccacggtagct   chiton
tggcaactgttgccgtaggaat--------aaacgccatagacacggtagct  sea hare
tggcaacagttgccgtaggaat-------aaacgccatagccaccgtagct   abalone
tggcaacagttgccgtaggaat-------aaacgccatagccaccgtagct   owl limpet
tggcaactgttgccgtaggaat-------aaacgccatagccaccgtagct   oyster
tggcaactgttgccgtaggaat-------aaacgccatagccacggtagtt   nautilus
tggcaactgttgccctaggaat-------aaacgccatagccacggtatct   octopus
tggcaactgttgccctaggaat-------aaacgccatagacacggtagct   solenogaster
tggcaacagttgccgtaggaat-------aaacgccatagccaccgtagct   peanut worm
tggcaacagttgccgtaggaat-------aaacgccatagccacggtagct   sandworm
tggcaacagttgccgtaggaat-------aaacgccatagccacggtagct   Satsuma tubeworm
tggcaacagttgccgtaggaat-------aaacgccatagccaccgtagct   shingle tubeworm
tggcaacagttgccgtaggaat-------aaacgccatagccaccgtagct   shamisen shell
tggcaacagttgccgtaggaat-------aaacgccatagccaccgtagct   horseshoe worm
tggcaacagttgccgtaggaat--------aaacgccatagacaccgtagct  bootlace worm
```

```
aaccaatggcg...tgacacg----agctccatataaggca-gcgatctccatag  human NR2F1
-----------...caacccggctgctgccttataaggcgcgccgtcgccatgg   human NR2F2
aaccaatggtg...tgacatg----agctccatataaggca-gcgatctccatag  chimaera nr2f1
atccaatgcgg...tcgggtg---tagcgccatataaggaa-ggcatctccatag  chimaera nr2f2
agccaatggga...tgacatg---tcactccatatatggta-gtaattgtcatag  lancelet
agccaatcagc...tgacatt---tctttccatataaggca-ataattgtcatag  acorn worm
-----------...tgacacc---gcctgccatataaggca-gtgggtgtcatgg  sea urchin
agccaatgaga...tgacacg-ctcgcgccatatatggaa-gcagttgccatgg   centipede
ggccaatggca...tgacacg--------------------------------    millipede
agccaatgaga...tgacacg--ttttgccatatatggaa-gtggttaccatgc   scorpion
agccaatgaga...tgacacg--ttttaccatatatggaa-tacgttgccatgc   horseshoe crab
gcccactgtg-...tgacatg---tctatccttataaggaa---tatggtcggtg  sea spider
agccaatgaga...tgacatg---ttttaccatatatggag-ttagtatccatgg  velvet worm
gaccaatagga...tgacaca---gtactccatataggaa-gtagtcgccatag   chiton
gaccaatagga...tgacacc---gctctccatataagga--gaggtaaccatag  abalone
ggccaatcaga...tgactcc---cagcgccatataagga--ggtgtaaccatag  owl limpet
--ccaatagga...tgacatt---agactccttatatggaa---aactgccatag  oyster
gaccaatggga...tggcagt---gtcatccatatatggg--cttgtatccatag  nautilus
agccaatgagc...tgacatc---acctgccttatatggca-gtggttgctatag  peanut worm
gaccaatagga...tgacatg---ccctgccatataaggaa-gtggtttccatgg  sandworm
-----------...tgacacg---gcgcgccatatatggaa-gacgtcaccatag  Satsuma tubeworm
agccaatgaga...tgacttg---tcctcccaaatatggaa-ggagtaaccatgg  shingle tubeworm
agccaatagga...tgacacg---ttgtgccttataaggta-gatgttgtcatag  shamisen shell
agccaatagga...tgacatg---cggagccttatatggca-cgagttactatag  bootlace worm
-----------...tgacagg----cttaccatatatagaa-tcagttgctaggg  stony coral
-----------...tgacagc----cggaccatatatagca-gtggtttccacgg  lace coral
-----------...tgacagt---tattgccatatatggtg-cacgcttccatgg  starlet sea anemone
-----------...tgactct---tgctcccttatatggca-tg-gctgctatgg  pale anemone
```

```
gtcttttcaca----aaccagcagctggt   human GSX1
ttctattcagg---agttctgcagctggt   human GSX2
gtcttttcaca----aaccagcagctggt   chimaera gsx1
gtcttttcacg----agctagcagctggt   chimaera gsx2
ggtgggtttcc------acggcagatggt   lancelet
gttttttcacc----aaacggcagatggt   acorn worm
gtttttctacc----aaacggcagatggt   velvet worm
gttttttccacc----aaacgccagctggt  cactus worm
gtttttctacc----aaccggcagatggt   chiton
gtt------------------------     abalone
gtttttctacc----aaacggcagatggc   nautilus
gtttttctacc----aaacggcagatggc   octopus
ggtttttttgccttcaatacggcagatgtt  peanut worm
tttttttccgca----aaacggcagatgtt  shamisen shell
gttctttcgtc----atacggcagatggt   horseshoe worm
gcttttctggc----aaacggcagatgtc   bootlace worm
```

```
cccgcccccggagacccacaattggtcc   human TBX2
cacgcctatgcagcaacacaattggtcc   human TBX3
cacgcctatacgggcacacaattggtcc   chimaera tbx2
cacgcctatggaggcacacaattggtcc   chimaera tbx3
cacgcctctctggcgaggggattggctg   lancelet
cacgcctatactcaacgctgattggtcg   acorn worm
cccgcctatttagcgcccggattggtgg   velvet worm
cccgccccgctcgttaaatgattggttg   abalone
cacgcctatttctcaccctgattggccc   bootlace worm
```

```
cctttgat---caggaaaatccaat   human SP5
gctttgat---cagacaaatccaat   chimaera sp5
cctttgat---gaagtcaacccaat   chimaera sp5-like
cctttgatcaagagttggcccctat   lancelet
tctttgat---ggcacgaatccaat   acorn worm
cctttgat---aactgaaatccaat   sea urchin
ccctcaaa---ggtcggaatccaat   horseshoe crab
gctttgat---ctcctcaatccaat   oyster
cctttgat---ctcccgaatccaat   nautilus
tctttgat---gtgccaaatccaat   peanut worm
gctttgaa---gttgaaaatccaat   shingle tubeworm
cctttgat---ctcgtaaatccaat   shamisen shell
gctttgat---catagaaatccaat   stony coral
```

```
ttctgccct-ccgcgagccaatcg   human PTCH1
ttctgcct--tcctcggccaatga   chimaera ptch1
ttgcttcttccctttagccaatgg   chimaera ptch2
ttgtggag--cccgcggccaatca   lancelet
ttggccg---tgagtgaccaatca   acorn worm
tagaaag---ccacgagccaatcg   horseshoe crab
tagaaag---ccacgagccaatcg   horseshoe crab
ctgcgcta--taattgaccaatca   chiton
cctactga--taattgaccaatca   sea hare
cctactga--taattgaccaatca   abalone
ctgtgggc--ttaagaaccaatca   shamisen shell
```

Figure 2: Parts of alignments between regulatory DNA sequences. The full alignments are in the Supplement. The over-bars show: CCAAT boxes (in the *Nr2f*, *Sp5*, and *Ptch* promoters), reverse-strand CCAAT box ATTGG (*Tbx2/3* promoter), E-box CAnnTG (*Gsx* enhancer), Wnt response element CTTTG (*Sp5* promoter), and CArG box CC(A/T)$_6$GG (*Nr2f* promoter). The *Ptch* promoter matches two places in the horseshoe crab genome, both near genes annotated as "patched-like".

Figure 3: Position and orientation of conserved DNA segments (red arrows) relative to nearby genes (blue arrows). Within each column, all the genes are homologous to each other. Animals without gene annotations are not shown. A regulatory segment doesn't really have an orientation, but once an orientation is arbitrarily chosen for any one segment, that defines the orientations of all its homologs.

an intron, so is transcribed into RNA: it might function in the DNA or RNA or both. This RNA region, of the mouse *Coe* gene *EBF1*, interacts with PSPC1 protein during fat cell development, promoting export of the RNA from the nucleus and expression of EBF1 protein [21].

The *Znf503/703* genes encode transcriptional repressors that are related to Sp transcription factors, but are thought not to bind directly to DNA. They contribute to brain and eye development in vertebrates and flies. In vertebrates they define hindbrain territories and motor neuron subtypes, and contribute to development of the brain's striatum, and limbs [22]. The *Znf503* enhancer was previously found in tick [5]: here it is found in diverse protostomes (fig. 1). The *Znf503/703* enhancer is 2.5 kb further upstream: it directs expression, in mouse embryos, in branchial arches and their derivatives, apical ectodermal ridge of limb buds, and urogenital tissues [23].

The other genes also regulate development [24–27]. *Hmx* interacts with *Gsx* in development of the brain's hypothalamus [28]. *Ptch* genes encode the main receptors that recognize hedgehog signalling molecules, and Smad6 proteins inhibit response to BMP signalling molecules [29]. *Sp5* is a target of Wnt signalling in bilaterians, and also in cnidaria where it prevents development of multiple heads [30]. It has a promoter region conserved from human to coral, which includes an especially conserved sequence resembling a Wnt response element (fig. 2). The *Rfx1/3* genes control formation of cilia on cells, which are used for hedgehog signalling, establish left-right body asymmetry, and more [31]. The promoter is spectacularly conserved across bilateria (fig. 2). Ssbp2 protein partners with LIM homeodomain proteins, which specify motor neurons in the same way in vertebrates and flies, among other roles [32]. *Nr2f* participates in neural development of bilaterians and cnidaria [33]: its promoter has been conserved since early neural evolution (fig. 1, 2).

Interestingly, the three genes with conserved 3′-UTR segments encode proteins that bind to RNA. The Cpeb proteins control mRNA localization and translation, in neural development and memory formation [34]. They are regulated by miRNAs binding to their 3′-UTRs [35–37]. *Mex3* contributes to defining anterior territories in bilaterian embryos, by destabilizing mRNAs [38, 39]. Its 3′-UTR contains elements for translational enhancement and destabilization, including by its own protein or miRNA [38, 40]. The UTR was previously found conserved between vertebrates and lancelet [38]: here it is found in other phyla and even a sea anemone (fig. 1). The Znf207 protein is a bit of an outlier: it contributes to chromosome alignment during cell division, a function not specific to animal development. It also, however, contributes to early neural development by binding to DNA or RNA [41].

Do these particular developmental genes have a common theme? It is hard to say, but perhaps there is a tendency for roles in ventral CNS (e.g. subpallium, striatum, hypothalamus, neurohypophysis, motor neurons).

In the conserved DNA segments, conserved transcription factor binding sites were identified (see the Supplement), using the JASPAR database of DNA-binding tendencies [42]. The results are uncertain, because transcription factors have weak sequence preferences, often similar to those of other transcription factors. Perhaps different factors bind to overlapping sites at different times depending on molecule concentrations.

On the other hand, there are clearly-conserved "boxes", including E-boxes and many CCAAT boxes (fig. 2). The E-box, CAnnTG, binds to various transcription factors that have a bHLH (basic-helix-loop-helix) structure. bHLH factors initiate neural development in bilateria and cnidaria [11], perhaps by binding to these conserved E-boxes. CCAAT boxes bind to the transcription factor NF-Y, which controls many developmental genes [43]. One CArG box was found, conserved from human to sea anemone (though imperfect in coral), in the *Nr2f* promoter (fig. 2). In general, CArG boxes bind to the Srf transcription factor, which controls cell migration and neural outgrowth in development [44].

Another interesting result is that the conserved DNA segments rarely change position (upstream/downstream) or orientation relative to the nearby genes. Figure 3 shows just one change, for *Gsx* in shamisen shell. There are a few others. In chimaera, the *sox21* gene has been inverted relative to its neighboring genes, but the inverted region doesn't include the enhancer. So, the enhancer is upstream of *sox21* in chimaera but downstream in the other animals.

Some protostome animals have more detected DNA elements than others (fig. 1). This is partly artifactual, e.g. some genomes lacked gene annotations so had the stricter 0.0001 E-value threshold. With that said, these seem to have many conserved elements: abalone, shamisen shell, and velvet worm (which lacks gene annotations). Perhaps these animals have undergone less evolution of their development and body plan. All three have sometimes been regarded as "primitive" or "living fossils": the shamisen shell (*Lingula*) was noted in Darwin's Origin of Species as little changed from ancient fossils.

On the other hand, no conserved segments were found in fly (*Drosophila melanogaster*), roundworm (*Caenorhabditis elegans*), or the leech *Helobdella robusta*. Perhaps the developmental systems of these animals have evolved more drastically [45]. Beyond bilateria, no conserved segments were found in ctenophores (comb jellies), sponges, or *Trichoplax* (Supplement table 2). A recent study found three regulatory DNA elements conserved between mammals and flies [46]: we confirmed they are conserved between human and chimaera, but didn't find them in invertebrates. Even when comparing each pair of human and fly sequences, boosting sensitivity 10^{12}-fold over comparing the whole genomes, our methods found tiny similar fragments in just one out of three pairs. Another study found identical DNA segments ≥ 50 bp between human and fly, and even human and sponge [47]. We confirmed that such intriguing matches exist, though they overlap regions that encode protein, rRNA, tRNA, or snRNA, or pseudogenes. A third study found $6\,533$ non-protein-coding segments conserved across deuterostomes [48], vastly more than found here or elsewhere [4, 5]. Some possible reasons are that they used E-value ≤ 0.05 per 5 kb (which is 1 per 100 kb), and didn't exclude pseudogenes or RNA genes.

In summary, we begin to see a system of DNA elements controlling gene expression for animal development, conserved since ancestral bilaterians and beyond in the Precambrian. This conservation helps us understand how early animals developed, and how modern development is built on these ancient components.

Why have these DNA sequences been so conserved? Judging by present-day bilaterians and cnidaria, the common ancestor of these animals already had an exquisitely integrated multicellular body, with a nervous system, that developed from a single fertilized cell [11, 12]. This complex development must have evolved by accretion of developmental processes, often by redeploying existing pathways [28]. So, one regulatory DNA element could become used in many parts of development, or its outcomes might get built upon by many processes. This brings us to Hyrum's Law of software development: "With a sufficient number of users...all observable behaviors of your system will be depended on by somebody" [49]. Thus, DNA changes with even subtle effects on transcription factor interactions, e.g. though DNA shape [50], would likely be harmful. This resulted in strongly-conserved DNA controlling core animal development, upon which evolved arthropods, molluscs, chordates, and so on.

References

1. Vavouri, T., Walter, K., Gilks, W. R., Lehner, B. & Elgar, G. Parallel evolution of conserved non-coding elements that target a common set of developmental regulatory genes from worms to humans. *Genome biology* **8,** 1–14 (2007).

2. Maeso, I., Irimia, M., Tena, J. J., Casares, F. & Gómez-Skarmeta, J. L. Deep conservation of cis-regulatory elements in metazoans. *Philosophical Transactions of the Royal Society B: Biological Sciences* **368,** 20130020 (2013).

3. Harmston, N., Barešić, A. & Lenhard, B. The mystery of extreme non-coding conservation. *Philosophical Transactions of the Royal Society B: Biological Sciences* **368,** 20130021 (2013).

4. Royo, J. L. *et al.* Transphyletic conservation of developmental regulatory state in animal evolution. *Proceedings of the National Academy of Sciences* **108,** 14186–14191 (2011).

5. Clarke, S. L. *et al.* Human developmental enhancers conserved between deuterostomes and protostomes. *PLoS genetics* **8,** e1002852 (2012).

6. Wong, E. S. *et al.* Deep conservation of the enhancer regulatory code in animals. *Science* **370,** eaax8137 (2020).

7. Frith, M. C. How sequence alignment scores correspond to probability models. *Bioinformatics* **36,** 408–415 (2020).

8. Venkatesh, B. *et al.* Elephant shark genome provides unique insights into gnathostome evolution. *Nature* **505,** 174–179 (2014).

9. Pei, J. & Grishin, N. V. C2H2 zinc finger proteins of the SP/KLF, Wilms tumor, EGR, Huckebein, and Klumpfuss families in metazoans and beyond. *Gene* **573,** 91–99 (2015).

10. Tomer, R., Denes, A. S., Tessmar-Raible, K. & Arendt, D. Profiling by image registration reveals common origin of annelid mushroom bodies and vertebrate pallium. *Cell* **142,** 800–809 (2010).

11. Arendt, D., Tosches, M. A. & Marlow, H. From nerve net to nerve ring, nerve cord and brain—evolution of the nervous system. *Nature Reviews Neuroscience* **17,** 61–72 (2016).

12. De Robertis, E. M. & Tejeda-Muñoz, N. Evo-Devo of Urbilateria and its larval forms. *Developmental biology* **487,** 10–20 (2022).

13. Desmaris, E. *et al.* DMRT5, DMRT3, and EMX2 cooperatively repress *Gsx2* at the pallium–subpallium boundary to maintain cortical identity in dorsal telencephalic progenitors. *Journal of Neuroscience* **38,** 9105–9121 (2018).

14. Salomone, J. *et al.* Conserved Gsx2/Ind homeodomain monomer versus homodimer DNA binding defines regulatory outcomes in flies and mice. *Genes & Development* **35,** 157–174 (2021).

15. MacKenzie, A., Purdie, L., Davidson, D., Collinson, M. & Hill, R. E. Two enhancer domains control early aspects of the complex expression pattern of *Msx1*. *Mechanisms of development* **62,** 29–40 (1997).

16. Sato, S. *et al.* Regulation of *Six1* expression by evolutionarily conserved enhancers in tetrapods. *Developmental biology* **368,** 95–108 (2012).

17. Trowe, M.-O. *et al.* Inhibition of Sox2-dependent activation of *Shh* in the ventral diencephalon by Tbx3 is required for formation of the neurohypophysis. *Development* **140,** 2299–2309 (2013).

18. Papaioannou, V. E. The T-box gene family: emerging roles in development, stem cells and cancer. *Development* **141,** 3819–3833 (2014).

19. Liu, X. *et al.* A single-nucleotide mutation within the *TBX3* enhancer increased body size in Chinese horses. *Current Biology* **32,** 480–487 (2022).

20. Catela, C. *et al.* An ancient role for collier/Olf/Ebf (COE)-type transcription factors in axial motor neuron development. *Neural development* **14,** 1–13 (2019).

21. Wang, J. *et al.* RNA-binding protein PSPC1 promotes the differentiation-dependent nuclear export of adipocyte RNAs. *The Journal of clinical investigation* **127,** 987–1004 (2017).

22. Pereira, F., Duarte-Pereira, S., Silva, R. M., da Costa, L. T. & Pereira-Castro, I. Evolution of the NET (NocA, Nlz, Elbow, TLP-1) protein family in metazoans: insights from expression data and phylogenetic analysis. *Scientific reports* **6,** 38383 (2016).

23. Chang, S. L.-Y. *et al.* Identification of two evolutionarily conserved 5'cis-elements involved in regulating spatiotemporal expression of *Nolz-1* during mouse embryogenesis. *PLoS One* **8,** e54485 (2013).

24. Parker, H. J., De Kumar, B., Pushel, I., Bronner, M. E. & Krumlauf, R. Analysis of lamprey *meis* genes reveals that conserved inputs from Hox, Meis and Pbx proteins control their expression in the hindbrain and neural tube. *Developmental Biology* **479,** 61–76 (2021).

25. Guntur, A. R., Venkatanarayan, A., Gangula, S. & Lundell, M. J. Zfh-2 facilitates Notch-induced apoptosis in the CNS and appendages of *Drosophila melanogaster*. *Developmental Biology* **475,** 65–79 (2021).

26. Lorente-Sorolla, J., Truchado-Garcia, M., Perry, K. J., Henry, J. Q. & Grande, C. Molecular, phylogenetic and developmental analyses of Sall proteins in bilaterians. *EvoDevo* **9,** 1–13 (2018).

27. Robinton, D. A. *et al.* The Lin28/*let-7* pathway regulates the mammalian caudal body axis elongation program. *Developmental cell* **48,** 396–405 (2019).

28. Wang, W., Grimmer, J. F., Van De Water, T. R. & Lufkin, T. *Hmx2* and *Hmx3* homeobox genes direct development of the murine inner ear and hypothalamus and can be functionally replaced by *Drosophila Hmx*. *Developmental cell* **7,** 439–453 (2004).

29. Miyazawa, K. & Miyazono, K. Regulation of TGF-β family signaling by inhibitory Smads. *Cold Spring Harbor perspectives in biology* **9,** a022095 (2017).

30. Vogg, M. C. *et al.* An evolutionarily-conserved Wnt3/β-catenin/Sp5 feedback loop restricts head organizer activity in *Hydra. Nature communications* **10,** 312 (2019).

31. Choksi, S. P., Lauter, G., Swoboda, P. & Roy, S. Switching on cilia: transcriptional networks regulating ciliogenesis. *Development* **141,** 1427–1441 (2014).

32. Yasuoka, Y. & Taira, M. LIM homeodomain proteins and associated partners: Then and now. *Current Topics in Developmental Biology* **145,** 113–166 (2021).

33. Gauchat, D. *et al.* The orphan COUP-TF nuclear receptors are markers for neurogenesis from cnidarians to vertebrates. *Developmental biology* **275,** 104–123 (2004).

34. Kozlov, E., Shidlovskii, Y. V., Gilmutdinov, R., Schedl, P. & Zhukova, M. The role of CPEB family proteins in the nervous system function in the norm and pathology. *Cell & Bioscience* **11,** 1–14 (2021).

35. Morgan, M., Iaconcig, A. & Muro, A. F. CPEB2, CPEB3 and CPEB4 are coordinately regulated by miRNAs recognizing conserved binding sites in paralog positions of their 3′-UTRs. *Nucleic acids research* **38,** 7698–7710 (2010).

36. Vetere, G. *et al.* Selective inhibition of miR-92 in hippocampal neurons alters contextual fear memory. *Hippocampus* **24,** 1458–1465 (2014).

37. Margvelani, G., Meparishvili, M., Kiguradze, T., McCabe, B. J. & Solomonia, R. Micro-RNAs, their target proteins, predispositions and the memory of filial imprinting. *Scientific Reports* **8,** 17444 (2018).

38. Takada, H. *et al.* The RNA-binding protein Mex3b has a fine-tuning system for mRNA regulation in early *Xenopus* development (2009).

39. Yang, L. *et al.* The human RNA-binding protein and E3 ligase MEX-3C binds the MEX-3–recognition element (MRE) motif with high affinity. *Journal of Biological Chemistry* **292,** 16221–16234 (2017).

40. Umehara, T. *et al.* Body temperature-dependent MicroRNA expression analysis in rats: *Rno-MiR-374-5p* regulates apoptosis in skeletal muscle cells via *Mex3B* under hypothermia. *Scientific Reports* **10,** 15432 (2020).

41. Malla, S. *et al.* ZFP207 sustains pluripotency by coordinating OCT4 stability, alternative splicing and RNA export. *EMBO reports* **23,** e53191 (2022).

42. Castro-Mondragon, J. A. *et al.* JASPAR 2022: the 9th release of the open-access database of transcription factor binding profiles. *Nucleic acids research* **50,** D165–D173 (2022).

43. Stanney III, W. *et al.* Combinatorial action of NF–Y and TALE at embryonic enhancers defines distinct gene expression programs during zygotic genome activation in zebrafish. *Developmental biology* **459,** 161–180 (2020).

44. Miano, J. M. Role of serum response factor in the pathogenesis of disease. *Laboratory investigation* **90,** 1274–1284 (2010).

45. Kuo, D.-H. & Weisblat, D. A. A new molecular logic for BMP-mediated dorsoventral patterning in the leech *Helobdella. Current Biology* **21,** 1282–1288 (2011).

46. Bridi, J. C. *et al.* Ancestral regulatory mechanisms specify conserved midbrain circuitry in arthropods and vertebrates. *Proceedings of the National Academy of Sciences* **117,** 19544–19555 (2020).

47. Ryu, T., Seridi, L. & Ravasi, T. The evolution of ultraconserved elements with different phylogenetic origins. *BMC evolutionary biology* **12,** 1–11 (2012).

48. Simakov, O. *et al.* Hemichordate genomes and deuterostome origins. *Nature* **527,** 459–465 (2015).

49. Wright, H. *Hyrum's Law* `https://www.hyrumslaw.com/`.

50. Gordân, R. *et al.* Genomic regions flanking E-box binding sites influence DNA binding specificity of bHLH transcription factors through DNA shape. *Cell reports* **3,** 1093–1104 (2013).