The TOR signaling pathway regulates vegetative development, aflatoxin biosynthesis, and pathogenicity in *Aspergillus flavus*

Guoqi Li, Xiaohong Cao, Elisabeth Tumukunde, Qianhua Zeng, Shihua Wang[*]

Key Laboratory of Pathogenic Fungi and Mycotoxins of Fujian Province, Key Laboratory of Biopesticide and Chemical Biology of Education Ministry, and School of Life Sciences, Fujian Agriculture and Forestry University, Fuzhou, 350002, China.

Corresponding author:

Name: Shihua Wang Professor

School of Life Sciences,

Fujian Agriculture and Forestry University

Fuzhou, Fujian, 350002, China

Phone/Fax: +86-591-83789312

E-mail: wshyyl@sina.com

These authors contributed equally to this work.

Abstract

The target of rapamycin (TOR) signaling pathway is highly conserved and plays a crucial role in diverse biological processes in eukaryotes. However, the underlying mechanism of the TOR pathway in *A. flavus* remains elusive. In this study, we identified and characterized nine genes encoding various components of the TOR pathway in *A. flavus,* and investigated their biological function. The FK506-binding protein Fkbp3 and its lysine succinylation are important for aflatoxin production and rapamycin resistance. The Tor kinase plays a pivotal role in the regulation of growth, spore production, aflatoxin biosynthesis, and osmotic and rapamycin stress. As a significant downstream effector molecule of the Tor kinase, the Sch9 kinase regulates aflatoxin B_1 (AFB_1) synthesis, osmotic and calcium stress response in *A. flavus,* and this regulation is mediated through its S_TKc, S_TK_X domains, and the ATP binding site at K340. We also showed that the Sch9 kinase may have a regulatory impact on the high-osmolarity glycerol (HOG) signaling pathway. TapA and TipA, the other downstream components of the Tor kinase, play significant roles in regulating sclerotia formation and cell wall stress response in *A. flavus*. The members of the TapA-phosphatase complexes, SitA and Ppg1, are crucial for various biological processes in *A. flavus,* including vegetative growth, sclerotia formation, AFB_1 biosynthesis, and pathogenicity. Furthermore, we showed that SitA and Ppg1 are involved in regulating lipid droplets (LDs) biogenesis and cell wall integrity (CWI) signaling pathway. In addition, another phosphatase complex, Nem1/Spo7, plays

critical roles in hyphal development, conidiation, aflatoxin, and lipid droplets

biogenesis. This study provides an important insight into the regulatory network of

the TOR signaling pathway and the molecular mechanism of aflatoxin biosynthesis in

A. flavus.

Keywords: *Aspergillus flavus*; the TOR signaling pathway; aflatoxin; vegetative

development

Introduction

Aspergillus flavus is one of the most important phytopathogenic fungi that

frequently contaminates various plants, causing economic losses in agricultural

commodities and posing serious health concerns in developing countries [1]. *A. flavus*

is also an opportunistic human pathogenic fungus that can cause aspergillosis in

immune-compromised patients [2]. Aflatoxins (AFs), the toxic and carcinogenic

secondary metabolites produced by *A. flavus*, are considered the most potent

carcinogen present in nature, causing liver cancer in both humans and animals [3]. The

aflatoxin biosynthesis is a complex process involving a series of reactions. Although

the majority of the enzyme reactions and genes involved in aflatoxin biosynthesis

have been elucidated [4], the related signaling pathway network and specific regulatory

mechanisms governing aflatoxin biosynthesis remain elusive. Therefore, exploring the

regulatory mechanism of the aflatoxin biosynthesis signaling pathway would provide

new insights for preventing *A. flavus* and aflatoxins contamination.

In eukaryotes, the target of rapamycin (TOR) signaling pathway is highly

conserved and plays essential roles in various significant biological processes, including ribosome biosynthesis, cell growth, and autophagy [5]. The serine/threonine-protein kinase Tor serves as a crucial protein component in the TOR signaling pathway, which interacts with other proteins to form multi-protein complexes [6]. In *Saccharomyces cerevisiae*, TOR exists in two distinct multiprotein functional complexes, namely TORC1 and TORC2 [7]. TORC1-mediated signaling governs cellular growth by modulating various growth-related mechanisms and exhibits sensitivity to rapamycin. Conversely, TORC2-mediated signaling primarily regulates cytoskeletal remodeling but remains unaffected by rapamycin [7]. In yeast, there are two *tor* genes, *tor1* and *tor2*, whereas in higher eukaryotes such as plants, animals, and filamentous fungi, there is only one *tor* gene [8-10].

In the fungal kingdom, the TOR signaling pathway has been extensively investigated in three prominent species: the budding yeast *S. cerevisiae* [11-12], the fission yeast *Schizosaccharomyces pombe* [13-15], and the human pathogen *Candida albicans* [16-17]. In *S. cerevisiae*, macrolide antibiotic rapamycin interacts with the cytosolic protein Fkbp12 to form a complex, and the Fkbp12-rapamycin complex effectively inhibits the activity of the Tor kinase function by binding to the FRB domain [18]. The Tap42 phosphatase complexs serves as the primary target of the Tor kinase. Additionally, the Tap42 protein interacts with the Tip41 protein to collaboratively regulate phosphatases, including PP2A and Sit4 [19-20]. Phosphatase Sit4 can dephosphorylate the transcription factor Gln3, regulating the nitrogen source metabolism pathway [21]. In addition, another immediate effector of TORC1 is the

AGC kinase Sch9, which participates in various cellular functions processes, such as

ribosome biosynthesis, translation initiation, and entry into G_0 phase [22]. The TOR

signaling pathway has been documented in various filamentous fungi, including

Fusarium graminearum [23-24], *Magnaporthe oryzae* [25], *Phanerochaete*

chrysosporium [26], *Podospora anserine* [27], *A. nidulans* [28], *A. fumigatus* [29-30], *F.*

fujikuroi [31], *Botrytis cinerea* [32], *F. oxysporum* [33], and *Mucor circinelloides* [34].

However, the TOR signaling pathway has not yet been reported in *A. flavus*.

In eukaryotic cells, the TOR signaling pathway and the Mitogen Activated Protein

Kinase (MAPK) signaling pathways play an important role in regulating adaptive

responses to extra- and intracellular conditions [35]. Several studies have demonstrated

that the TOR signaling pathway interacts with various other signaling pathways, such

as MAPK and CWI. These pathways engage in crosstalk, forming a complex

metabolic network that regulates cellular processes [36-40]. However, the underlying

mechanisms of multiple crosstalks between the TOR and other signaling pathways in

A. flavus remain unclear. Therefore, we intended to identify the genes associated with

the TOR signaling pathway and elucidate their roles and contributions in regulating

vegetative development and aflatoxin biosynthesis in *A. flavus*. Our results

demonstrated that the TOR signaling pathway is involved in multiple cellular

processes in *A. flavus*. Our findings revealed a complex interplay between the TOR

pathway and other signaling pathways (such as HOG and CWI) in *A. flavus*, which

play a crucial role in regulating cellular growth and survival under environmental

stress conditions.

Results

Rapamycin exhibits inhibitory effects on the growth, sporulation, sclerotia formation, and aflatoxin production in *A. flavus*

Rapamycin, a secondary metabolite synthesized by *Streptomyces hygroscopicus*, exhibits efficacy in suppressing specific filamentous fungi. In *F. graminearum*, rapamycin has a significant inhibitory impact on growth and asexual reproduction [23]. To investigate the effects of rapamycin on vegetative development, sclerotia formation, and secondary metabolism in *A. flavus*, we conducted an assay to determine the sensitivity of the wild-type strain to rapamycin. We observed that the wild-type strain exhibited a high sensitivity to rapamycin. We found that 100 ng/mL of rapamycin significantly inhibited mycelial growth and conidia formation (Fig.1A, 1D, 1E). More importantly, the wild-type strain exhibited a significant reduction in sclerotia formation and aflatoxin synthesis (Fig.1B, 1C, 1F, 1G). Additionally, rapamycin resulted in a significant decrease in spore germination rate and sparser conidiophores (Fig. S1A, S1B). These results illustrated that the TOR signaling pathway may be involved in regulating the growth, development, and secondary metabolism in *A. flavus*. Calcofluor white is a chemifluorescent blue dye commonly employed for its nonspecific binding affinity to fungal chitin. Fluorescence microscopy analysis revealed a notable reduction in the blue fluorescence on the cell wall following treatment with rapamycin (Fig.S1C). The intracellular lipid droplets within the hyphae were labeled with BODIPY. We observed that the hyphae treated

with rapamycin contained more lipid droplets compared to the control group (Fig.

S1D). These findings suggest that the TOR pathway may play important roles in

maintaining cell wall integrity and facilitating the biogenesis of lipid droplets.

To investigate the potential regulatory role of the TOR pathway in modulating

the sporulation, sclerotia formation, and aflatoxin biosynthesis in *A. flavus*, we

examined the expression levels of conidia-related, sclerotia formation-related, and

aflatoxin-related genes after 3, 6, and 9 hours of treatment with rapamycin. To

determine the effects of rapamycin on conidiation, qRT-PCR was conducted to assess

the transcript levels of two conidia-related genes, *abaA* and *brlA*. We also conducted

qRT-PCR to evaluate the expression levels of three regulators (*nsdC*, *nsdD*, and *sclR*)

involved in sclerotia development, and genes involved in AFB_1 biosynthesis, such as

the regulatory gene *aflS*, as well as several structural genes (*aflC* and *aflQ*). The

strains treated with rapamycin exhibited a significant reduction in the expression

levels of genes related to sporulation, sclerotia formation, and aflatoxin synthesis (Fig.

1H, 1I, 1J). Overall, these results indicated that the TOR pathway plays a crucial role

in regulating the growth, sporulation, sclerotia formation, and aflatoxin biosynthesis

in *A. flavu*s by modulating the expression of these genes.

The impact of Fkbp3 and its lysine succinylation on AFB_1 biosynthesis and

rapamycin resistance in *A. flavus*

FK506-binding proteins (Fkbps) belong to a highly conserved immunophilin

family, which are involved in various cellular functions, such as protein folding,

immunosuppression, signaling transduction, and transcription [41]. Fkbps function as

molecular switches by binding to target proteins and inducing conformational changes

[41]. In *S. cerevisiae*, Fkbp12 and rapamycin interact to form a complex that exerts a

negative regulatory effect on the activity of the Tor kinase [42]. The *fkbp12* gene

homologue *fprA* in *A. nidulans* shares high levels of homology with that from *S.*

cerevisiae [28]. According to the protein sequence of FprA from *A. nidulans*, four genes

encoding putative proteins with FK506 binding domain, named *fkbp1*

(AFLA_126880), *fkbp2* (AFLA_087480), *fkbp3* (AFLA_010910), and *fkbp4*

(AFLA_128200), have been identified from the *A. flavus* genome. We generated four

fkbp knockout strains using homologous recombination, and all *fkbp* disruption strains

were confirmed by PCR and sequencing analysis (Fig.S2). Phenotypic analyses

showed that the mycelial growth and conidia formation of all the *fkbp* mutants were

comparable to the wild-type strain on the PDA medium (Fig. 2A), suggesting that

Fkbps have minimal influence on the process of vegetative development. We

evaluated AFB_1 synthesis utilizing thin layer chromatography (TLC), and the findings

demonstrated a noteworthy reduction in AFB_1 synthesis upon deletion of Fkbp3, in

comparison to the wild-type strain (Fig.2B, 2D). Additionally, the quantity of sclerotia

exhibited a significant decrease in all *fkbp* mutants compared to the wild-type strain

(Fig.S3A, S3B). The aforementioned data indicated that Fkbp3 plays a crucial role in

sclerotia formation and aflatoxin biosynthesis in *A. flavus*.

To determine the specific Fkbp responsible for rapamycin sensitivity, all mutant

strains were cultured on PDA solid plates amended with 100 ng/mL rapamycin. The

strain with disrupted *fkbp3* showed significant resistance to rapamycin, while the

deletion mutants of *fkbp1*, *fkbp2*, and *fkbp4* did not exhibit any alteration in resistance

to rapamycin (Fig.2A, 2C). These results demonstrated that Fkbp3 serves as the

principal target for rapamycin in *A. flavu*s. In the FK506-sensitivity analysis, we

observed a significant suppression of mycelial growth and conidia formation in *A.*

flavus with the addition of 10 ng/mL FK506 to the PDA medium. Additionally, the

deletion of *fkbp3* led to the heightened resistance to FK506 compared to the wild-type

strain (Fig.2A, 2C). These findings suggested that Fkbp3 is mainly involved in the

modulation of FK506 resistance in *A. flavus*.

Previous studies have shown that lysine succinylation plays a significant role in

aflatoxin production and pathogenicity in *A. flavus* [43]. Based on the analysis of our

succinylome data, we have identified six reliable succinylation sites (K5, K19, K40,

K42, K55, and K65) on the Fkbp3 protein. To identify the function of Fkbp3

succinylation sites, site-directed mutations were constructed according to the

homologous recombination strategy. Compared to the wild-type strain, all point

mutants exhibited a comparable phenotype in vegetative growth and conidiation.

Interestingly, we observed that the K19A mutation resulted in increased resistance to

rapamycin (Fig. 2E, 2F). Additionally, the K19A strain exhibited a significant

reduction in the production of AFB_1 compared to the wild-type strain (Fig. 2G, 2H).

These findings suggested that the succinylation of Fkbp3 at K19 plays a crucial role in

rapamycin resistance and aflatoxin biosynthesis. Furthermore, we found that various

point mutants, including K40A, K42A, and K55A, exhibited reduced levels of AFB_1

production in comparison to the wild-type strain (Fig. 2G, 2H), suggesting that the

succinylation of Fkbp3 at other sites also contributes to AFB$_1$ biosynthesis in *A. flavus*.

The above results indicated that the K19 site likely plays a more prominent role in

Fkbp3 succinylation compared to other sites.

The Tor kinase plays a critical role **in *A. flavus***

In *S. cerevisiae*, the Tor kinase can interact with a diverse range of proteins,

resulting in the formation of complex structures known as TORC1 and TORC2 [44].

TORC1 can detect various signals, including nutrients, growth factors, and

environmental stress. It plays a crucial role in regulating cellular processes such as

gene transcription, protein translation, ribosome synthesis, and autophagy [45]. To

identify the ortholog of the Tor protein in *A. flavus*, the *S. cerevisiae* Tor protein was

utilized as search queries in the *A. flavus* genome database using the Basic Local

Alignment Search Tool (https://blast.ncbi.nlm.nih.gov/Blast.cgi). The genome of *A.*

flavus contains a solitary ortholog of the *tor* gene (AFLA_044350), which encodes a

protein exhibiting 48.91% similarity to *S. cerevisiae* Tor2. We used the absolute

quantitative PCR technique to identify the copy number of the *tor* gene at the genome

level in *A. flavus* (Table S1). To elucidate the function of the Tor kinase in *A. flavus*,

we employed gene deletion techniques to target the *tor* gene. While our attempt to

obtain a null mutant was unsuccessful, we constructed a single copy deletion strain,

tor+/-, through homologous recombination, which contains one copy of the *tor* gene

(Table S1). Compared to the wild-type strain, the *tor+/-* strain exhibited a

significantly slower growth rate on PDA (Fig. 3C). Additionally, the *tor+/-* strain

produced fewer conidia and sclerotia compared to the wild-type strain (Fig. 3B, S4).

The results from thin-layer chromatography (TLC) revealed a noteworthy decrease in

AFB$_1$ synthesis in the *tor+/-* strain compared to the wild-type strain (Fig. 3D, 3F).

Interestingly, the *tor+/-* mutant exhibited a notable increase in sensitivity to

rapamycin and decreased sensitivity to osmotic stress (Fig. 3C, 3E, S5). Furthermore,

the qRT-PCR results demonstrated a significant downregulation in the expression

levels of the genes associated with sexual sporulation and aflatoxin synthesis in the

tor+/- mutant compared to the wild-type strain (Fig. S6). The transcriptional level of

the *tor* gene in the *tor+/-* mutant was downregulated compared to the level in the

wild-type strain (Fig. S6). These findings suggested that the Tor kinase plays a crucial

role in various biological processes in *A. flavus*, including vegetative growth, asexual

development, sclerotia formation, response to environmental stress, and biosynthesis

of aflatoxins.

The Tor protein is characterized by the presence of various conserved domains,

including the DUF3385, FAT, FKBP12-rapamycin binding (FRB), kinase catalytic,

and FATC domains, which are located at the N-terminus (Fig. 3A). To investigate the

functional significance of the FRB domain within the Tor kinase, the FRB domain

was eliminated using homologous recombination. Compared to the *tor+/-* strain, the

tor$^{\Delta FRB}$ mutant displayed comparable deficiencies in vegetative growth, conidiation,

sclerotia formation, osmotic stress, and rapamycin sensitivity (Fig. 3B, 3C, S4, S5).

Based on these observations, we speculated that the FRB domain plays a pivotal role

in the modulation of the Tor kinase activity, thereby influencing fungal development

and rapamycin resistance in *A. flavus*.

The Sch9 kinase is involved in aflatoxin biosynthesis and the HOG pathway

Sch9, a Ser/Thr kinase belonging to the AGC family, is directly phosphorylated by TORC1. The phosphorylation of Sch9 is inhibited by rapamycin, as well as by carbon or nitrogen starvation [22]. The putative *sch9* gene (AFLA_127440) in *A. flavus* encodes a protein consisting of 706 amino acids, exhibiting 80.31% identity to the Sch9 protein in *A. nidulans*. To investigate the role of Sch9 in *A. flavus*, we attempted to create a *sch9* deletion mutant using a homologous recombination strategy. The Δ*sch9* strain exhibited typical vegetative growth and conidiation similar to the wild-type strain (Fig. 4B). TLC assay and quantitative analysis showed a significantly decreased aflatoxin production in the Δ*sch9* strain compared to the wild-type strain (Fig. 4C, 4G). These results indicated that Sch9 regulates aflatoxin biosynthesis in *A. flavus*.

In yeast, Sch9 plays an important role in the transcriptional activation of osmostress inducible genes [46]. To characterize the roles of Sch9 in stress response, we examined the sensitivity of the *sch9* deletion strain to various stresses including osmotic stress and calcium stress. Compared with the wild-type strain, the Δ*sch9* mutant significantly increased sensitivity to calcium stress (Fig. S7), and decreased sensitivity to osmotic stress induced by NaCl and KCl (Fig.4D, 4H). In *A. flavus*, the Sch9 kinase consists of several domains, including the protein kinase C conserved region 2 (C2), protein kinase (S_TKc), AGC-kinase C-terminal (S_TK_X) domain, and an ATP binding site (Fig. 4A). To explore the role of these domains and the ATP binding site in Sch9 kinase, we constructed deletion strains for both domains and a

sch9^{K340A} mutant. The results revealed that the *sch9*^{ΔS_TKc} strain, *sch9*^{ΔS_TK_X} strain, and

sch9^{K340A} mutant exhibited similar phenotypes in osmotic stress and AFB$_1$ production

as the Δ*sch9* mutant (Fig.4B, 4C, 4D). Therefore, we speculate that the protein kinase

domain, kinase C-terminal domain, and the ATP binding site at K340 play a crucial

role in modulating the impact of Sch9 on aflatoxin biosynthesis and stress response in

A. flavus.

To investigate the potential function of Sch9 in the HOG and TOR pathways in *A.*

flavus, we conducted a sensitivity analysis of the Δ*sch9* strain to rapamycin.

Compared to the wild-type strain, the *sch9* deletion strain exhibited a slightly higher

sensitivity to rapamycin (Fig. 4B, 4F). The MAKP Hog1 is an element of the high

osmolarity glycerol (HOG) pathway. To verify the connection between Sch9 and the

HOG-MAPK pathway, the phosphorylation levels of Hog1 kinase were measured

under osmotic stress (1M NaCl). Western blot analysis showed that the

phosphorylation level of Hog1 increased significantly in the Δ*sch9* strain (Fig. 4E).

This finding suggested that Sch9 plays a pivotal role in modulating the response of

Hog1 to osmotic stress. Taken together, these results reflected that Sch9 regulates

osmotic stress response via the HOG pathway in *A. flavus*.

TapA and TipA participate in regulation of sclerotia development and cell wall

stress response

Tap42, a protein that associates with phosphatase 2A, serves as the direct target of

the Tor kinase in yeast. Tap42 plays a crucial role in various Tor functions,

particularly in the regulation of transcriptional processes [47]. The genome of *A. flavus*

contains a homolog of *tap42* named *tapA* (AFLA_092770), which is predicted to

encode a 355 amino-acid protein (sharing 31.9% identity with *S. cerevisiae* Tap42).

To further determine the function of the TapA regulator in the TOR signaling pathway

of *A. flavus*, we conducted targeted gene deletion experiments on *tapA*. We failed to

obtain a null mutant, therefore we generated the *tapA+/-* strain, which contains one

copy of the *tapA* gene (Table S2). Compared to the wild-type strain, the *tapA+/-* strain

exhibited diminished hyphal growth and conidiation (Fig. 5A, 5C), indicating that

TapA is important for vegetative growth and conidiation. The results of the rapamycin

sensitivity test indicated that the *tapA+/-* strain displayed reduced sensitivity to

rapamycin (Fig.5A, 5E). In addition, the *TapA+/-* strain was unable to generate any

sclerotia (Fig. 5B, 5F), and almost lost its capability in AFB_1 production compared to

the wild-type strain (Fig. 5D, 5G), these findings indicated that TapA is essential for

sclerotia formation and biosynthesis of AFB_1. The *tapA+/-* mutant was more

insensitive to cell wall stress, including CFW and CR (Fig. S8). Furthermore, the

levels of *nsdC*, *nsdD*, and *sclR* expression were significantly decreased in *tapA+/-*

strain compared to those in the wild-type strain (Fig. S9). The transcriptional level of

the *tapA* gene in the *tapA+/-* strain was drastically downregulated compared to the

wild-type strain (Fig. S9). These findings have demonstrated the significantly positive

impact of TapA on the growth, sporulation, sclerotia formation, cell wall stress, and

AF biosynthesis in *A. flavus*.

In *S. cerevisiae*, Tip41 was identified as a Tap42-interacting protein and has been

measured to function as a negative regulator of Tap42, which downregulates TORC1

signaling via activation of PP2A phosphatase [48]. The putative orthologue of *tip41* in

A. flavus named *tipA* (AFLA_047310). The amino acid sequence of TipA exhibits a

similarity of 40.49% to Tap41 in *S. cerevisiae*. To elucidate the function of the TOR

signaling pathway protein TipA in *A. flavus*, we generated the *tipA* deletion mutants.

The Δ*tipA* strain exhibited a slightly reduced number of conidia produced on PDA

compared to the wild-type strain (Fig. 5A). Meanwhile, the Δ*tipA* strain exhibited

decreased sensitivity to rapamycin (Fig. 5A). The Δ*tipA* strain was unable to produce

any sclerotia (Fig. 5B). In addition, the Δ*tipA* strain was more sensitive to cell wall

and cell membrane stress, including CR and SDS (Fig. S8). Additionally, the

expression levels of *nsdC* and *nsdD* exhibited a decrease in Δ*tipA* strain compared to

the wild-type strain (Fig. S9). These results indicated that TipA is crucial for the

sclerotial formation and cell wall stress in *A. flavus*.

Phosphatase SitA and Ppg1 are involved in growth, conidiation, and sclerotial

formation

In *S. cerevisiae*, Tap42 has been identified to interact with the catalytic subunit of

type 2A protein phosphatases, including Pph3, Pph21, and Pph22, as well as the type

2A-like phosphatases Sit4 and Ppg1[19]. Phosphatase *sitA* (AFLA_108450) and *ppg1*

(AFLA_132610) were identified in *A. flavu*s, encoding putative orthologs of *sit4* and

ppg1 in *S. cerevisiae*. To determine the biological functions of phosphatase SitA and

Ppg1 in *A. flavu*s, we disrupted the *sitA* and *ppg1* by homologous recombination

strategy. To further investigate the relationship between phosphatase SitA and Ppg1,

we additionally generated complemented and double deletion mutants. The vegetative

development phenotype analysis showed that the $\Delta sitA$, $\Delta ppg1$, and $\Delta sitA/\Delta ppg1$

strains grew significantly more slowly than the wild-type strain on PDA (Fig. 6A, 6D).

Additionally, the $\Delta sitA$, $\Delta ppg1$, and $\Delta sitA/\Delta ppg1$ strains were more sensitive to

rapamycin (Fig. 6A, 6E). In addition, the $\Delta sitA$, $\Delta ppg1$, and $\Delta sitA/\Delta ppg1$ strains

formed lower number of conidiophores (Fig. 6A). To gain further insight into the role

of SitA and Ppg1 in the process of conidiation, qRT-PCR was employed to determine

the transcript levels of conidial synthesis-related genes (*abaA* and *brlA*). The

expression levels of the *abaA and brlA* genes were significantly downregulated in the

$\Delta sitA$, $\Delta ppg1$, and $\Delta sitA/\Delta ppg1$ mutants when compared to the wild-type and

complemented strains (Fig. 6G). The analysis of sclerotial formation revealed that the

$\Delta sitA$, $\Delta ppg1$, and $\Delta sitA/\Delta ppg1$ mutants exhibited a complete inability to produce any

sclerotia (Fig. 6B). To further prove these findings, we conducted qRT-PCR to

determine the expression levels of three regulators (*nsdC*, *nsdD*, and *nclR*) of sclerotia

development. Our findings indicated that the expression levels of *nsdC*, *nsdD,* and

sclR were all significantly reduced in the $\Delta sitA$, $\Delta ppg1$, and $\Delta sitA/\Delta ppg1$ mutants (Fig.

6H), which is consistent with the observed decrease in sclerotia production in the

$\Delta sitA$, $\Delta ppg1$, and $\Delta sitA/\Delta ppg1$ mutants. These results showed that SitA and Ppg1 are

crucial for hyphal development, conidiation, and sclerotia formation in *A. flavus*.

SitA and Ppg1 play crucial roles in aflatoxin biosynthesis and pathogenicity

We assayed AFB_1 biosynthesis in the $\Delta sitA$, $\Delta ppg1$, and $\Delta sitA/\Delta ppg1$ mutants.

TLC analyses revealed that the $\Delta sitA$ and $\Delta sitA/\Delta ppg1$ strains exhibited a decrease in

AFB_1 production, whereas the $\Delta ppg1$ strain showed a significant increase in AFB_1

production compared to the wild-type and complemented strains (Fig. 6C, 6F). The expression levels of genes related to aflatoxin synthesis, including structural genes (*aflO* and *aflP*) and regulatory genes (*aflR* and *aflS*), were determined by qRT-PCR in the Δ*sitA*, Δ*ppg1*, and Δ*sitA*/Δ*ppg1* mutants. The results revealed a significant downregulation of these genes in the Δ*sitA* and Δ*sitA*/Δ*ppg1* mutants, while upregulated in the Δ*ppg1* mutants, compared to the wild-type and complemented strains (Fig. 6I). These results indicated that SitA and Ppg1 play different roles in regulating aflatoxin biosynthesis in *A. flavu*s.

We conducted pathogenicity tests on peanuts and maize to examine the pathogenicity of the phosphatases SitA and Ppg1 on crops. The results showed that the wild-type and complemented strains demonstrated complete virulence on all peanut and maize seeds. In contrast, the Δ*sitA*, Δ*ppg1*, and Δ*sitA*/Δ*ppg1* mutants displayed compromised colonization on peanut and maize seeds (Fig. 7A). In addition, the number of conidial productions in these mutants on the infected peanut seeds also dramatically decreased compared to those of the wild-type and complemented strains (Fig. 7C). The thin-layer chromatography analyses revealed that the Δ*ppg1* mutant exhibited increased production of AFB$_1$ in peanuts and maize. Conversely, the levels of aflatoxin produced by the Δ*sitA* and Δ*sitA*/Δ*ppg1* mutants on peanut and maize were significantly reduced compared to the the wild-type and complemented strains (Fig. 7B, 7D). This observation aligns with the above results of aflatoxin biosynthesis from all mutant strains in the YES medium. All these data showed that SitA and Ppg1 are crucial for crop seeds' pathogenicity.

SitA and Ppg1 are involved in regulating cell wall integrity

To explore the role of phosphatases SitA and Ppg1 in response to cell wall stress, all strains were cultivated on PDA amended with CFW, CR, and SDS. We found that the relative growth inhibition of Δ*sitA*, Δ*ppg1* and Δ*sitA*/Δ*ppg1* mutants induced by cell wall stress were significantly increased compared to the wild-type and complemented strains (Fig. 8A, 8B). Further investigations using qRT-PCR revealed a notable reduction in the expression levels of genes related to the synthesis of cell walls, specifically chitin synthase genes *chsA*, *chsD*, *chsE*, and *β-1,6*, in the Δ*sitA*, Δ*ppg1*, and Δ*sitA*/Δ*ppg1* strains (Fig. 8C). Western blot analysis also revealed that the phosphorylation level of Slt2 was significantly elevated in the Δ*sitA*, Δ*ppg1*, and Δ*sitA*/Δ*ppg1* strains under cell wall stress (200 μg/ml CFW) (Fig. 8D). In conclusion, the phosphatases SitA and Ppg1 play a crucial role in regulating cell wall integrity.

The phosphatase complex Nem1/Spo7 are important for growth, aflatoxin biosynthesis, and LD biogenesis

Lipid droplets (LDs) are highly dynamic spherical organelles, which appear in the cytosol of most eukaryotic cell types[49]. Rapamycin treatment led to the accumulation of lipid droplets in various filamentous fungi [24]. The FgTOR pathway regulates the phosphorylation status of FgNem1 mainly via the FgPpg1/Sit4-FgCak1/other kinase cascade in *F. graminearum* [24]. In this study, we observed that the treatment of hyphae with rapamycin resulted in the accumulation of LD biogenesis (Fig. 9A). This finding suggested that the TOR pathway may regulate the LD biogenesis. However, the underlying regulatory mechanism of LD

biosynthesis remain largely unknown in *A. flavu*s. To investigate the regulatory

mechanism of the TOR pathway in *A. flavus* that controls LD biogenesis, we

examined LD accumulation in Δ*sch9*, Δ*sitA*, and Δ*ppg1* mutants. BODIPY staining

assays revealed that LD biogenesis was not observed in the Δ*sitA* and Δ*ppg1* strains

upon treatment with rapamycin, in contrast to the wild-type and Δ*sch9* strains (Fig.

9A, 9B, 9C, 9D). These results indicated that LD biogenesis induced by rapamycin is

predominantly reliant on the phosphatases SitA and Ppg1 in *A. flavus*. In *F.*

graminearum, the phosphatase FgNem1/FgSpo7 complex was involved in the

induction of LD biogenesis by rapamycin. Phosphatase *nem1* (AFLA_002649) and

spo7 (AFLA_028590) were identified in *A. flavu*s, which encode putative orthologs of

nem1 and *spo7* in *F. graminearum.* To examine the role of the *nem1* and *spo7* genes in

A. flavus, we generated Δ*nem1* and Δ*spo7* mutants using homologous recombination.

The Δ*nem1* and Δ*spo7* mutants displayed a notable reduction in both vegetative

growth and conidiation compared to the wild-type strain (Fig. 10A, 10C, 10D). TLC

assay revealed a significantly decreased aflatoxin production in the Δ*nem1* and Δ*spo7*

compared to the wild-type strain (Fig. 10B). In addition, BODIPY staining showed no

visible lipid droplets in the hyphae of Δ*nem1* and Δ*spo7* after rapamycin treatment

(Fig. 10E). Taken together, these results indicated that Nem1/Spo7 are involved in

vegetative growth, conidiation, aflatoxin production, and LD biogenesis.

Discussion

Aflatoxins (AFs) are a class of toxic secondary metabolites synthesized by *A.*

flavus, including aflatoxin B_1, B_2, G_1, and G_2, among which AFB_1 has been regarded

as the most pathogenic and toxic aflatoxin[50]. Therefore, understanding the regulatory

mechanisms involved in fungal secondary metabolites biosynthesis would be

instrumental for effectively addressing aflatoxin control in *A. flavus*. In this study, we

demonstrated that the target of rapamycin (TOR) signaling pathway plays a pivotal

role in regulating growth, sporulation, sclerotia formation, aflatoxin production, and

various stress response in *A. flavus*. Through conducting functional studies on genes

within the TOR signaling pathway, we explored the conservation and complexity

exhibited by the TOR signaling pathway in *A. flavu*s. In addition, we constructed a

crosstalk network between the TOR and other signaling pathways (HOG, CWI) and

analyzed the specific regulatory process. We proposed that the TOR signaling

pathway interacts with multiple signaling pathways, including HOG and CWI

pathway, to modulate the cellular responses to diverse environmental stresses (Fig.

11).

We have observed that rapamycin has a significant impact on the growth and

sporulation of *A. flavus* (Fig. 1A). Additionally, it exhibits a strong inhibitory effect on

sclerotia production and aflatoxin biosynthesis (Fig. 1B, 1C). The inhibitory effect has

been documented in various fungal species, including *Candida albicans* [51],

Cryptococcus neoformans [51]*, Podospora anserine* [27]*, A. nidulans* [28]*, Magnaporthe*

oryzae [39]*,* and *F. graminearum* [23]. This suggested that the TOR signaling pathway

may have a conserved function in filamentous fungi. The number of FK506-binding

proteins varies among different species: there are four Fkbps protein in *S. cerevisiae*

and three Fkbps protein in *S. pombe* [52]. In *S. cerevisiae*, rapamycin can bind to Fkbp

protein, resulting in the irreversible inhibition of the G_1 phase of the cell cycle and the

regulation of cell growth [53]. In *A. nidulans,* the *fkbp12* gene homologue *fprA* displays

considerable sequence similarity with homologues identified in the *S. cerevisiae* [28].

In this study, based on the homologous alignment of the protein sequence of FprA, it

was observed that four *fkbp* genes present in *A. flavus*. Among these, deletion of *fkbp3*

exhibited enhanced tolerance to rapamycin (Fig. 2A). So we speculated that

rapamycin forms a complex with Fkbp3 in *A. flavus*, subsequently targeting

downstream target genes in order to exert its biological function. We found that Fkbp3

is involved in sclerotia and aflatoxin biosynthesis in *A. flavus* (Fig. 2A, S3)*,* and

subsequent analysis revealed that the underlying mechanism may be associated with

succinylation modification. The succinylation sites of Fkbp3 were identified through

the utilization of succinylation proteomics data. By constructing point mutant strains

and conducting phenotype experiments, we found that the succinylation of Fkbp3 at

residue K19 is involved in rapamycin resistance and aflatoxin biosynthesis (Fig. 2E,

2G). Previous studies have demonstrated that Fkbp12 has been identified as the

receptor for FK506 and rapamycin, which inhibits calcineurin and target of rapamycin

complex 1 (TORC1), respectively [53]. The deletion of Fkbp12 confers the resistance

to rapamycin in *C. albicans* [51] and *C. neoformans* [51]. The entomopathogenic fungus

Beauveria bassiana contains three putative *fkbp* genes, and disruption of *Bbfkbp12*

significantly increases the resistance to rapamycin and FK506 [54]. The plant

pathogenic fungus *F. graminearum* harbors three Fkbps proteins, among which

deletion of *Fgfkbp12* leads to resistance to rapamycin [23]. Thus, we propose that these

Fkbps may exhibit analogous functionalities in conferring resistance to rapamycin. In

Botrytis cinerea, deletion of *Bcfkbp12* cause a reduction of the virulence of strain T4,

while knockout of the *fkbp12* gene did not affect the pathogenic development of the

strain B05.10 [32]. These findings demonstrated that the *fkbp12* homologous genes

exhibited conservation in terms of rapamycin resistance. However, there is significant

variation in other biological functions among fungal species, which may be attributed

to species or strain evolutionary divergence.

The Tor kinase functions as a fundamental component of the TOR signaling

pathway, which exhibits evolutionary conservation across fungi, plants, and mammals

[55]. TOR assembles into two distinct multi-protein complexes known as TORC1 and

TORC2. A characteristic that distinguishes TORC1 is its interaction with KOG1

(Raptor in mammals) and its sensitivity to the rapamycin-FKBP12 complex [56].

Almost all other eukaryotes, including plants, worms, flies, and mammals, have a

single *tor* gene, whereas *S. cerevisiae* has two *tor* genes [45]. The Tor kinase is widely

conserved both in terms of its structural characteristics and its role as the target of

Fkbp-rapamycin. Typically, the Tor kinase exhibits comparable structural

characteristics, including the HEAT-like, FAT, FRB, kinase, and FATC domains. The

FRB domain is the binding region of the Fkbp-rapamycin complex, and rapamycin

resistance is usually caused by the disruption of the FRB domain [45]. The comparison

of homology and the analysis of phylogenetics revealed that there is only one

predicted *tor* gene in *A. flavus*. To ascertain its functionality, we generated the *tor*

single-copy deletion mutant by homologous recombination. We discovered that the

Tor kinase is involved in the vegetative growth and asexual development of *A. flavus*

(Fig. 3B, 3C). In addition, the *tor+/-* mutant shows a high sensitivity to rapamycin

(Fig. 3C). Our findings have demonstrated the significant involvement of Tor kinase

and its FRB domain in the processes of morphogenesis, aflatoxin biosynthesis, and

multiple stress responses in *A. flavus*. We speculate the common expectation that the

deletion of the *tor* and FRB domain should result in insensitivity and resistance to

rapamycin，as it disrupts the binding site for Fkbp-rapamycin. However, we observed

that the *tor+/-* and FRB domain-deleted mutant were more sensitive to rapamycin.

This intriguing result suggests that there are additional factors or complexities

involved in TOR signaling pathway regulation in *A. flavus*. We hypothesize that this

result is related to the double copy of the *tor* gene. Previous findings demonstrated

that the Tor kinase could function as a molecular switch connecting an iron cue to

defend against pathogen infection in *C. elegans* [57]. In *A. fumigatus*, the Tor kinase

represents a central regulatory node controlling genes and proteins involved in

nutrient sensing, stress response, cell cycle progression, protein biosynthesis, and

degradation, as well as adaptation to low iron conditions[58]. In summary, the Tor

kinase acts as a central regulator in the TOR signaling pathway, playing a crucial role

in the regulation of the cell cycle, protein synthesis, and cellular energy metabolism.

In *S. cerevisiae*, the protein kinase Sch9 is a major downstream effector of

TORC1, and this kinase is known to have crucial functions in stress resistance,

longevity, and nutrient sensing [46]. In addition, the TORC1-Sch9 pathway acts as a

crucial mediator of chronological lifespan [59]. In filamentous fungi, an intricate

network of interactions exists between the TOR and HOG pathways. In *F.*

graminearum [38], FgSch9 and FgHog1 mutants displayed heightened susceptibility to

osmotic and oxidative stresses. The phosphorylation level of FgHog1 increased

significantly in FgSch9 mutant. Additionally, the affinity capture-MS assay showed

that the Tor kinase was among the proteins co-purifying with FgSch9. These results

suggest that FgSch9 serves as an intermediary for the TOR and HOG signaling

pathways [38]. Furthermore, FgSch9 plays a crucial role in the regulation of various

biological processes such as hyphal differentiation, asexual development, virulence,

and DON biosynthesis[38]. In *A. fumigatus*, the SchA mutant was sensitive to

rapamycin, high concentrations of calcium and hyperosmotic stress, and SchA was

involved in iron metabolism. The SchA mutant exhibited enhanced phosphorylation

of SakA in response to osmotic stress [51]. In this study, we discovered that Sch9 is

involved in regulating aflatoxin biosynthesis (Fig.4C), and responses to osmotic stress

and rapamycin stress (Fig.4B, 4D). The S_TKc domain deletion strain (*sch9*$^{\Delta S\_TKc}$),

S_TK_X domain deletion strain (*sch9*$^{\Delta S\_TK\_X}$)**,** and *sch9*^{K340A} mutant displayed a

similar phenotype to the Δ*sch9* strain (Fig. 4B, 4C, 4D). These findings revealed that

Sch9 plays a significant role in the biological synthesis of aflatoxin and in responding

to various stresses. This role is primarily mediated through its kinase domain,

kinase-C domain**,** and ATP binding site k340. We also found that the phosphorylation

level of Hog1 increased significantly in the Δ*sch9* strain under osmotic stress (Fig.

4E). Based on the above results, we hypothesized that Sch9 was involved in the

osmotic stress response by modulating the HOG pathway.

In *S. pombe*, the protein Tip41, which interacts with Tap42, plays a significant

role in the cellular responses to nitrogen sources by regulating type 2A phosphatases

[60]. In *S. cerevisiae*, Tap42-phosphatase complexes associate with TORC1, whereas

Tip41 functions to attenuate TORC1 signaling by stimulating the PP2A phosphatase

[48]. Unlike its yeast counterpart Tip41, TIPRL, a mammalian Tip41-like protein,

exerts a positive influence on mTORC1 signaling by interacting with PP2Ac [61].

While the interaction between Tip41 and Tap42 was not observed in the filamentous

fungi. The yeast two-hybrid (Y2H) assay showed that FgTip41 interacts with the

phosphatase FgPpg1 rather than FgTap42, while the FgTap42 bind with phosphatases

FgPp2A, implying that FgTap42, FgPpg1, and FgTip41 form a heterotrimer in *F.*

graminearum [23]. In the rice blast fungus *M. oryzae*, MoTip41 does not interact with

the MoTap42. However, MoTip41 has the ability to bind with the phosphatase

MoPpe1, thereby facilitating crosstalk between the TOR pathway and the cell wall

integrity pathway [40]. In this study, we found that the TapA and TipA played a crucial

role in sclerotia production and cell wall stress in *A. flavus* (Fig. 5B, S8). The *tapA*

single-copy deletion mutant and the *tipA* deletion mutant exhibited decreased

sensitivity to rapamycin (Fig. 5A). However, the interaction between TapA and TipA

is not yet clear in *A. flavus*. We speculated that the interrelations and functions of

TapA and TipA in *A. flavus* are comparable to their ortholog in other filamentous

fungi.

In *S. cerevisiae*, Tap42 can bind and regulate various PP2A phosphatases,

including Pph3, Pph21, Pph22, Sit4, and Ppg1. This interaction plays a crucial role in

the TOR signaling pathway, particularly in response to rapamycin treatment or

nutrient deprivation [48]. The phosphatase Sit4 plays a crucial role in regulating cell

growth, morphogenesis, and virulence in *C. albicans* [62]. FgSit4 and FgPpg1 are

associated with various cellular processes, including mycelial growth, conidiation,

DON biosynthesis, and virulence in *F. graminearum* [23]. In addition, FgSit4 and

FgPpg1 are involved in the regulation of various signaling pathways, such as the CWI

pathway and lipid droplets biogenesis regulation pathway [23]. In this study, we found

that the phosphatase SitA and Ppg1 play critical role in vegetative growth, conidiation,

sclerotia formation, and aflatoxin biosynthesis in *A. flavus* (Fig.6A, 6B, 6C).

Additionally, the phosphatase SitA and Ppg1 also affect pathogenicity and LD

biogenesis (Fig. 7A, 9C, 9D). In addition, TapA/TipA and SitA/Ppg1 exhibits similar

functions in sclerotia production and cell wall stress response. Based on these

observations, we hypothesized that TapA-phosphatase complexes play a collaborative

role in the regulation of the sclerotia formation and cell wall stress response in *A.*

flavus.

After the deletion of the *sitA* and *ppg1* genes, the expression level of genes

related to cell wall synthesis has decreased significantly, resulting in damage to the

cell wall and showing a higher sensitivity to cell wall integrity stress (Fig. 8C).

Western blotting results showed that the phosphorylation level of MAPK kinase Slt2

has increased significantly in the ΔsitA and Δppg1 strains under cell wall stress,

indicating that SitA and Ppg1 exert a negative regulatory effect on the CWI pathway

(Fig. 8D). In *M. oryzae*, MoPpe1 and MoSap1 serve as an adaptor complex that

connects the CWI and TOR signaling pathways. Activation of the TOR pathway

results in the inhibition of the CWI pathway [39]. In addition, the phosphatase MoPpe1

interacts with the phosphatase-associated protein MoTip41, facilitating the exchange

of information between the TOR and CWI signaling pathways [39]. These results

suggest that the phosphatase Ppe1 engages in interactions with various proteins to

effectively coordinate the TOR and CWI signaling pathways, thereby regulating the

growth and pathogenicity of the rice blast fungus *M. oryzae* [39]. Taken together, we

speculated that the phosphatase SitA and Ppg1 might exhibit differential regulation in

the CWI and TOR pathways across species in order to fulfill their respective

functions.

In *F. graminearum,* the phosphatase complex FgNem1/Spo7-FgPah1 cascade

plays significant roles in fungal development, DON production, and lipid droplet (LD)

biogenesis [24]. In this study, our results indicated that Nem1 and Spo7 are important

for vegetative growth and secondary metabolism of *A. flavus* (Fig. 10A, 10B). We

have also discovered that LD biogenesis is facilitated by the Nem1/Spo7 and

SitA/Ppg1 pathways (Fig. 9C, 9D, 10E). These observations suggest that the PP2A

phosphatases play a central role as core phosphatases in the LD biogenesis signaling

network. However, it is likely that there are other kinases involved in the regulation of

LD biogenesis. To gain a comprehensive understanding of the precise molecular

mechanism, further research should focus on investigating the interplay between

phosphatases and other kinases.

In conclusion, we identified the complex regulatory network that provides insight into the interplay between the TOR, HOG, and CWI signaling pathways, and how they collectively regulate the development, aflatoxin biosynthesis, and pathogenicity of *A. flavus*. By elucidating the functions of genes associated with the TOR signaling pathway, our research could establish a solid foundation for comprehending the molecular mechanism underlying the aflatoxin biosynthesis of *A. flavus*. This, in turn, can provide novel insights into the management of aflatoxin contamination and the mitigation of *A. flavus* infection.

Materials and methods

Strains and cultural conditions

The *A. flavus* strains in this study were shown in Table S3. All strains were cultivated on three different types of agar media: potato dextrose agar (PDA), yeast extract-sucrose agar (YES), and yeast extract-glucose agar (YGT). The growth and conidiation assays were conducted at 37℃. Additionally, aflatoxin production was assessed using YES liquid medium at 29℃.

Constructions of gene deletion and complemented mutants

The construction of gene deletion and complemented strains was carried out using the SOE-PCR strategy, following previously established protocols [63]. The primers utilized for the amplification of the flanking sequences of each gene can be found in Table S3. The putative gene deletion mutants were validated through PCR

assays using the appropriate primers (Fig. S2), and their confirmation was

subsequently obtained through sequencing analysis.

Detection of *tor* and *tapA* genes copy number in strains

Absolute quantification PCR has been used as a common method for molecular

detection of gene copy numbers. According to previous reports, the *sumo* gene is a

single-copy gene in *A. flavus*[64]. Following previously established protocols[65], the

primers were used to design the amplification of the *sumo* reference gene and *tor* and

tapA gene. Using *sumo* gene as reference, the *tor* and *tapA* gene copy number was

calculated by standard curve.

Growth, conidiation, and sclerotia analysis

To evaluate mycelial growth and conidia formation, 1 μL of 10^6 spores/mL conidia

of all strains was point inoculated onto the center of PDA medium, and then cultured

at 37°C for 5 days in the dark. The conidia were washed by 0.07% Tween-20 and

counted using a hemocytometer and microscope [43]. For sclerotia formation analysis,

each strain was inoculated and cultivated on CM agar medium at 37°C in the dark for

7 days, and then 75% ethanol was used to wash away mycelia and conidia on the

surface of the medium [64]. These assays were repeated at least three times.

Extraction and quantitative analysis of aflatoxin

To extract and quantify aflatoxins (AFs) production, 1 μL of 10^6 spores/mL

conidia of all strains was point inoculated onto liquid YES medium and cultures were

then incubated at 29°C in the dark for 6 days. AFs were extracted from the media and

detected using thin layer chromatography (TLC) [65]. In addition, the standard AFB_1

was employed for visual comparison, and AFs were quantitatively analyzed using the

Image J tool. The assays were repeated at least three times.

Stress assay

To determine the roles of the gene in *A. flavus* responding to various

environmental stresses, 1 µL of 10^6 spores/mL conidia of all strains were point

inoculated onto these PDA solid plates supplemented with various agents. These

agents included 1M NaCl, 1M KCl, and 1M $CaCl_2$ for inducing osmotic stress, as

well as 100 µg/mL SDS, 200 µg/mL Calcofluor white (CFW), and 200 µg/mL Congo

red (CR) for inducing cell wall stress. After 5 days of incubation at 37°C in the dark,

the relative rates of inhibition were determined [43]. The assays were repeated at least

three times.

Seeds infection assays

To measure the pathogenicity of *A. flavus*, all peanut and maize seeds were

washed in 0.05% sodium hypochlorite, 75% ethanol, and sterile water, respectively.

Subsequently, all peanut and maize seeds were immersed in sterile water containing

10^6 spores/mL of all strains at 29°C in the dark for 60 min. After the conidia adhesion,

the seeds were cultured at 29°C for 6 days in the dark [64].

Quantitative RT-PCR

For the real-time quantitative PCR (qPCR) assays, the *A. flavus* strains were

cultivated in PDA medium at 29°C for 3 days in the dark. Total RNA was extracted

from the mycelia of all strains, and cDNA was synthesized with the First Strand

cDNA Synthesis Kit, and then the cDNA was used as the template for qRT-PCR

analysis with SYBR Green qPCR mix. The relative transcript level of each gene was

calculated by the $2^{-\Delta\Delta CT}$ method [66].

Western blot analysis

To extract the total protein, the mycelia of each strain were cultivated in YES

liquid medium for 48 h, then the sample was grinded under liquid nitrogen to a fine

powder and transferred to a 1.5 mL microcentrifuge tube added with RIPA lysis buffer

and 1 mM PMSF protease inhibitor. Proteins were separated by 12% SDS-PAGE and

transferred onto a polyvinylidene fluoride (PVDF) membrane. Phosphorylation levels

of Hog1 and Slt2 were detected by Phospho-p38 MAPK antibody (Cell Signaling

Technology, Boston, MA, USA) and Phospho-p44/42 MAPK (Erk 1/2) antibody (Cell

signaling Technology, MA, USA), respectively. The experiment was conducted three

times independently.

Statistical analysis

670   All data were measured with means ± standard deviation (SD) of three biological

671   replicates. Graph Pad Prism 7 software was used for conducting statistical and

672   significance analysis, and p-values < 0.05 was represented statistical significance.

673   Student's *t*-test was used to compare the two means for differences, whereas Tukey's

674   multiple comparisons were used for testing multiple comparisons.

Acknowledgments

676   The authors also thank Professor Zhenhong Zhuang, Jun Yuan, Yu Wang, Kunzhi

677   Jia, Xiuna Wang, and Xinyi Nie for their help in the experiments. This work was

678 supported by the National Natural Science Foundation of China (Grant No.

679 31972214).

Author contributions

681 Guoqi Li and Xiaohong Cao designed the experiments. Guoqi Li, Xiaohong Cao,

and Qianhua Zeng performed the experiments. Guoqi Li, Xiaohong Cao, and Shihua

Wang analyzed the data. Guoqi Li, Elisabeth Tumukunde, and Shihua Wang wrote the

manuscript. The manuscript was discussed and revised by all the authors.

References

[1] Klich MA. *Aspergillus flavus*: the major producer of aflatoxin. Mol Plant Pathol.

2007; 8(6): 713-22.

[2] Hedayati MT, Pasqualotto AC, Warn PA, et al. *Aspergillus flavus*: human

pathogen, allergen and mycotoxin producer. Microbiology (Reading). 2007;

153(Pt 6): 1677-1692.

[3] Khan R, Ghazali FM, Mahyudin NA, et al. Biocontrol of aflatoxins using

non-aflatoxigenic *Aspergillus flavus*: a literature review. J Fungi (Basel). 2021;

7(5): 381.

[4] Yabe K, Nakajima H. Enzyme reactions and genes in aflatoxin biosynthesis. Appl

Microbiol Biotechnol. 2004; 64(6): 745-55.

[5] Wullschleger S, Loewith R, Hall MN. TOR signaling in growth and metabolism.

Cell. 2006; 124(3): 471-84.

[6] De Virgilio C, Loewith R. The TOR signalling network from yeast to man. Int J

Biochem Cell Biol. 2006; 38(9): 1476-1481.

[7] Loewith R, Jacinto E, Wullschleger S, et al. Two TOR complexes, only one of

which is rapamycin sensitive, have distinct roles in cell growth control. Mol Cell.

2002; 10(3): 457-68.

[8] Beauchamp EM, Platanias LC. The evolution of the TOR pathway and its role in

cancer. Oncogene. 2013 Aug 22; 32(34): 3923-32.

[9] Fu L, Liu Y, Qin G, et al. The TOR-EIN2 axis mediates nuclear signalling to

modulate plant growth. Nature. 2021; 591(7849): 288-292.

[10] Teichert S, Wottawa M, Schönig B, et al. Role of the *Fusarium fujikuroi* TOR

kinase in nitrogen regulation and secondary metabolism. Eukaryot Cell. 2006;

5(10): 1807-1819.

[11] Crespo JL, Hall MN. Elucidating TOR signaling and rapamycin action: lessons

from *Saccharomyces cerevisiae*. Microbiol Mol Biol Rev. 2002; 66(4): 579-591.

[12] Rohde JR, Bastidas R, Puria R, et al. Nutritional control via Tor signaling in

Saccharomyces cerevisiae. Curr Opin Microbiol. 2008; 11(2): 153-60.

[13] Ishiguro J, Shibahara K, Ueda Y, et al. Fission yeast TOR signaling is essential

for the down-regulation of a hyperactivated stress-response MAP kinase under

salt stress. Mol Genet Genomics. 2013; 288(1-2): 63-75.

[14] Otsubo Y, Yamamato M. TOR signaling in fission yeast. Crit Rev Biochem Mol

Biol. 2008; 43(4): 277-83.

[15] Petersen J. TOR signalling regulates mitotic commitment through stress-activated

MAPK and Polo kinase in response to nutrient stress. Biochem Soc Trans. 2009;

37(Pt1): 273-7.

[16]Bastidas RJ, Heitman J, Cardenas ME. The protein kinase Tor1 regulates adhesin

gene expression in *Candida albicans*. PLoS Pathog. 2009; 5(2): e1000294.

[17]Liu NN, Flanagan PR, Zeng J, et al. Phosphate is the third nutrient monitored by

TOR in *Candida albicans* and provides a target for fungal-specific indirect TOR

inhibition. Proc Natl Acad Sci U S A. 2017; 114(24): 6346-6351.

[18]Hu K, Guo S, Yan G, et al. Ubiquitin regulates TORC1 in yeast *Saccharomyces*

cerevisiae. Mol Microbiol. 2016; 100(2): 303-314.

[19]Wang H, Wang X, Jiang Y. Interaction with Tap42 is required for the essential

function of Sit4 and type 2A phosphatases. Mol Biol Cell. 2003; 14(11): 4342-51.

[20]Yan G, Shen X, Jiang Y. Rapamycin activates Tap42-associated phosphatases by

abrogating their association with Tor complex 1. EMBO J. 2006; 25(15):

3546-55.

[21]Tate JJ, Tolley EA, Cooper TG. Sit4 and PP2A Dephosphorylate Nitrogen

Catabolite Repression-Sensitive Gln3 When TorC1 Is Up- as Well as

Downregulated. Genetics. 2019; 212(4): 1205-1225.

[22]Urban J, Soulard A, Huber A, et al. Sch9 is a major target of TORC1 in

Saccharomyces cerevisiae. Mol Cell. 2007; 26(5): 663-74.

[23]Yu F, Gu Q, Yun Y, et al. The TOR signaling pathway regulates vegetative

development and virulence in *Fusarium graminearum*. New Phytol. 2014; 203(1):

219-32.

[24]Liu N, Yun Y, Yin Y, et al. Lipid droplet biogenesis regulated by the

FgNem1/Spo7-FgPah1 phosphatase cascade plays critical roles in fungal development and virulence in *Fusarium graminearum*. New Phytol. 2019; 223(1): 412-429.

[25] Sun G, Elowsky C, Li G, et al. TOR-autophagy branch signaling via Imp1 dictates plant-microbe biotrophic interface longevity. PLoS Genet. 2018; 14(11): e1007814.

[26] Nguyen DV, Roret T, Fernandez-Gonzalez A, et al. Target of rapamycin pathway in the white-rot fungus *Phanerochaete chrysosporium*. PLoS One. 2020; 15(2): e0224776.

[27] Dementhon K, Paoletti M, Pinan-Lucarré B, et al. Rapamycin mimics the incompatibility reaction in the fungus *Podospora anserina*. Eukaryot Cell. 2003; 2(2): 238-46.

[28] Fitzgibbon GJ, Morozov IY, Jones MG, et al. Genetic analysis of the TOR pathway in *Aspergillus nidulans*. Eukaryot Cell. 2005; 4(9): 1595-8.

[29] Alves de Castro P, Dos Reis TF, et al. The *Aspergillus fumigatus* SchA[SCH9] kinase modulates SakA[HOG1] MAP kinase activity and it is essential for virulence. Mol Microbiol. 2016; 102(4): 642-671.

[30] Baldin C, Valiante V, Krüger T, et al. Comparative proteomics of a tor inducible *Aspergillus fumigatus* mutant reveals involvement of the Tor kinase in iron regulation. Proteomics. 2015; 15(13): 2230-43.

[31] Teichert S, Wottawa M, Schönig B, et al. Role of the *Fusarium fujikuroi* TOR kinase in nitrogen regulation and secondary metabolism. Eukaryot Cell. 2006;

5(10): 1807-19.

[32] Meléndez HG, Billon-Grand G, Fèvre M, et al. Role of the *Botrytis cinerea*

FKBP12 ortholog in pathogenic development and in sulfur regulation. Fungal

Genet Biol. 2009; 46(4): 308-20.

[33] López-Berges MS, Rispail N, Prados-Rosales RC, et al. A nitrogen response

pathway regulates virulence functions in *Fusarium oxysporum* via the protein

kinase TOR and the bZIP protein MeaB. Plant Cell. 2010; 22(7): 2459-75.

[34] Bastidas RJ, Shertz CA, Lee SC, et al. Rapamycin exerts antifungal activity *in*

vitro and *in vivo* against *Mucor circinelloides* via FKBP12-dependent inhibition

of Tor. Eukaryot Cell. 2012; 11(3): 270-81.

[35] Madrid, M, Vázquez-Marín, B, Franco, A, et al. Multiple crosstalk between TOR

and the cell integrity MAPK signaling pathway in fission yeast. Sci Rep. 2016

Nov 23; 6: 37515.

[36] Torres J, Di Como CJ, Herrero E, et al. Regulation of the cell integrity pathway

by rapamycin-sensitive TOR function in budding yeast. J Biol Chem. 2002 Nov 8;

277(45): 43495-504.

[37] Chen DD, Shi L, Yue SN, et al. The Slt2-MAPK pathway is involved in the

mechanism by which target of rapamycin regulates cell wall components in

Ganoderma lucidum. Fungal Genet Biol. 2019; 123: 70-77.

[38] Gu Q, Zhang C, Yu F, et al. Protein kinase FgSch9 serves as a mediator of the

target of rapamycin and high osmolarity glycerol pathways and regulates multiple

stress responses and secondary metabolism in *Fusarium graminearum*. Environ

Microbiol. 2015; 17(8): 2661-76.

[39] Qian B, Liu X, Jia J, et al. MoPpe1 partners with MoSap1 to mediate TOR and cell wall integrity signalling in growth and pathogenicity of the rice blast fungus *Magnaporthe oryzae*. Environ Microbiol. 2018; 20(11): 3964-3979.

[40] Qian B, Liu X, Ye Z, et al. Phosphatase-associated protein MoTip41 interacts with the phosphatase MoPpe1 to mediate crosstalk between TOR and cell wall integrity signalling during infection by the rice blast fungus *Magnaporthe oryzae*. Environ Microbiol. 2021; 23(2): 791-809.

[41] Tong M, Jiang Y. FK506-Binding Proteins and Their Diverse Functions. Curr Mol Pharmacol. 2015; 9(1): 48-65.

[42] Heitman J, Movva NR, Hall MN. Targets for cell cycle arrest by the immunosuppressant rapamycin in yeast. Science. 1991; 253(5022): 905-9.

[43] Ren S, Yang M, Yue Y, et al. Lysine Succinylation Contributes to Aflatoxin Production and Pathogenicity in *Aspergillus flavus*. Mol Cell Proteomics. 2018; 17(3): 457-471.

[44] Weisman R. Target of Rapamycin (TOR) Regulates Growth in Response to Nutritional Signals. Microbiol Spectr. 2016 Oct; 4(5).

[45] Loewith R, Hall MN. Target of rapamycin (TOR) in nutrient signaling and growth control. Genetics. 2011 Dec; 189(4): 1177-201.

[46] Pascual-Ahuir A, Proft M. The Sch9 kinase is a chromatin-associated transcriptional activator of osmostress-responsive genes. EMBO J. 2007; 26(13): 3098-108.

[47] Düvel K, Broach JR. The role of phosphatases in TOR signaling in yeast. Curr

Top Microbiol Immunol. 2004; 279: 19-38.

[48] Jacinto E, Guo B, Arndt KT, et al. TIP41 interacts with TAP42 and negatively

regulates the TOR signaling pathway. Mol Cell. 2001; 8(5): 1017-26.

[49] Sánchez-Álvarez M, Del Pozo MÁ, Bosch M, et al. Insights into the biogenesis

and emerging functions of lipid droplets from unbiased molecular profiling

approaches. Front Cell Dev Biol. 2022 Jun 8; 10: 901321.

[50] Gizachew D, Chang CH, Szonyi B, et al. Aflatoxin B1 (AFB$_1$) production by

Aspergillus flavus and *Aspergillus parasiticus* on ground Nyjer seeds: The effect

of water activity and temperature. Int J Food Microbiol. 2019; 296: 8-13.

[51] Cruz MC, Goldstein AL, Blankenship J, et al. Rapamycin and less

immunosuppressive analogs are toxic to *Candida albicans* and *Cryptococcus*

neoformans via FKBP12-dependent inhibition of TOR. Antimicrob Agents

Chemother. 2001; 45(11): 3162-70.

[52] Galat A. Functional diversity and pharmacological profiles of the FKBPs and

their complexes with small natural ligands. Cell Mol Life Sci. 2013; 70(18):

3243-75.

[53] Kasahara K. Physiological function of FKBP12, a primary target of

rapamycin/FK506: a newly identified role in transcription of ribosomal protein

genes in yeast. Curr Genet. 2021 Jun; 67(3): 383-388.

[54] Li Y, Ren H, Zhao Y, et al. Characterization of three FK506-binding proteins in

the entomopathogenic fungus *Beauveria bassiana*. J Invertebr Pathol. 2020; 171:

107334.

[55] Xiong Y, Sheen J. The role of target of rapamycin signaling networks in plant growth and metabolism. Plant Physiol. 2014; 164(2): 499-512.

[56] Adami A, García-Alvarez B, Arias-Palomo E, et al. Structure of TOR and its complex with KOG1. Mol Cell. 2007; 27(3): 509-16.

[57] Ma YC, Dai LL, Qiu BB, et al. TOR functions as a molecular switch connecting an iron cue with host innate defense against bacterial infection. PLoS Genet. 2021; 17(3): e1009383.

[58] Baldin C, Valiante V, Krüger T, et al. Comparative proteomics of a tor inducible *Aspergillus fumigatus* mutant reveals involvement of the Tor kinase in iron regulation. Proteomics. 2015; 15(13): 2230-43.

[59] Pascual-Ahuir A, Proft M. The Sch9 kinase is a chromatin-associated transcriptional activator of osmostress-responsive genes. EMBO J. 2007; 26(13): 3098-108.

[60] Fenyvuesvolgyi C, Elder RT, Benko Z, et al. Fission yeast homologue of Tip41-like proteins regulates type 2A phosphatases and responses to nitrogen sources. Biochim Biophys Acta. 2005; 1746(2): 155-62.

[61] Nakashima A, Tanimura-Ito K, Oshiro N, et al. A positive role of mammalian Tip41-like protein, TIPRL, in the amino-acid dependent mTORC1-signaling pathway through interaction with PP2A. FEBS Lett. 2013; 587(18): 2924-9.

[62] Han Q, Pan C, Wang Y, et al. PP2A-like protein phosphatase (Sit4) regulatory subunits, Sap155 and Sap190, regulate *Candida albicans*' cell growth,

morphogenesis, and virulence. Front Microbiol. 2019 Dec 20; 10: 2943.

[63] Tan L, Chen H, Yu S, et al. A SOE-PCR method of introducing multiple

mutations into Mycoplasma gallisepticum neuraminidase. J Microbiol Methods.

2013; 94(2): 117-120.

[64] Nie X, Yu S, Qiu M, et al. *Aspergillus flavus* SUMO contributes to fungal

virulence and toxin attributes. J Agric Food Chem. 2016 Sep 7; 64(35): 6772-82.

[65] Wang Y, Wang S, Nie X, et al. Molecular and structural basis of nucleoside

diphosphate kinase-mediated regulation of spore and sclerotia development in the

fungus *Aspergillus flavus*. J Biol Chem. 2019 Aug 16; 294(33): 12415-12431.

[66] Livak KJ, Schmittgen TD. Analysis of relative gene expression data using

real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. Methods. 2001; 25(4):

402–408.

Fig. 1 Impacts of rapamycin on the growth, sporulation, sclerotia formation,and aflatoxin production of *Aspergillus flavus*.

(A) Colony morphology of the WT strain cultured on PDA medium amended with different concentrations of rapamycin at 37°C for 5 days.

(B) Colony phenotype of the WT strain grown on CM medium amended with different concentrations of rapamycin at 37°C for 7 days.

(C) TLC assay of AFB$_1$ production by the WT strain cultured in YES liquid media amended with different concentrations of rapamycin at 29°C for 6 days.

(D) Statistical analysis of the colony diameter by the WT strain treated with different concentrations of rapamycin described in (A).

(E) Conidial quantification of the WT strain treated with different concentrations of rapamycin as mentioned in (A).

(F) Quantitative analysis of sclerotium formation by the WT strain treated with different concentrations of rapamycin described in (B).

(G) Relative quantification of AFB$_1$ production by the WT strain treated with different concentrations of rapamycin as mentioned in (C)

(H) Relative expression levels of conidia-related genes in WT strain treated with 100 ng/mL rapamycin after 3, 6, 9 h.

(I) Relative expression levels of sclerotia-related genes in WT strain treated with 100 ng/mL rapamycin after 3, 6, 9 h.

(J) Relative expression levels of aflatoxin biosynthesis regulatory and structural genes in WT strain treated with 100 ng/mL rapamycin after 3, 6, 9 h.

A

	WT	Δ*fkbp1*	Δ*fkbp2*	Δ*fkbp3*	Δ*fkbp4*
PDA					
Rapamycin					
FK506					

B

ST　WT　Δ*fkbp1*　Δ*fkbp2*　Δ*fkbp3*　Δ*fkbp4*

C

D

E

	WT	Δ*fkbp3*	*fkbp3*K5A	*fkbp3*K19A	*fkbp3*K40A	*fkbp3*K42A	*fkbp3*K55A	*fkbp3*K65A
PDA								
Rapamycin								

F

G

ST　WT　Δ*fkbp3*　*fkbp3*K5A　*fkbp3*K19A　*fkbp3*K40A

ST　WT　Δ*fkbp3*　*fkbp3*K42A　*fkbp3*K55A　*fkbp3*K65A

H

Fig. 2 Disruption of *fkbp3* significantly increases the resistance of *A. flavus* to rapamycin and FK506.

(A) Phenotype of the WT and all mutant strains (Δ*fkbp1*, Δ*fkbp2*, Δ*fkbp3*, and Δ*fkbp4*) grown on PDA amended with rapamycin and FK506 at 37°C for 5 days.

(B) TLC analysis of AFB_1 production by the WT and all mutant strains (Δ*fkbp1*, Δ*fkbp2*, Δ*fkbp3*, and Δ*fkbp4*) cultured in YES liquid medium at 29°C for 6 days.

(C) The growth inhibition rate of the WT and all mutant strains (Δ*fkbp1*, Δ*fkbp2*, Δ*fkbp3*, and Δ*fkbp4*) under rapamycin and FK506 stress.

(D) AFB_1 quantitative analysis of the WT and all mutant strains (Δ*fkbp1*, Δ*fkbp2*, Δ*fkbp3*, and Δ*fkbp4*) as described in (B).

(E) Phenotype of the WT and all mutant strains (Δ*fkbp3*, K5A, K19A, K40A, K42A, K55A, and K65A) grown on PDA amended with 100 ng/mL rapamycin at 37°C for 5 days.

(F) The growth inhibition rate of the WT and all mutant strains (Δ*fkbp3*, K5A, K19A, K40A, K42A, K55A, and K65A) under rapamycin stress.

(G) TLC assay of AFB_1 production by the WT and all mutant strains (Δ*fkbp3*, K5A, K19A, K40A, K42A, K55A, and K65A) cultured in YES liquid medium at 29°C for 6 days.

(H) Relative quantification of AFB_1 production in the WT and all mutant strains (Δ*fkbp3*, K5A, K19A, K40A, K42A, K55A, and K65A) as mentioned in (G).

A

low complexity (246-255)	Domain of unknown function (DUF3385) (735-895)	FAT domain (1371-1743)
Rapamycin binding domain (1850-1949)	PI3Kc domain (2018-2307)	FATC domain (2353-2384)

B

WT   *tor* +/-   *tor* $^{\Delta FRB}$

C

WT   *tor* +/-   *tor* $^{\Delta FRB}$

PDA

Rapamycin

D

ST   WT   *tor* +/-   *tor* $^{\Delta FRB}$

E

Growth repression (%)

Rapamycin

WT
tor +/-
tor $^{\Delta FRB}$

F

AFB$_1$ concentration (mg/ml)

YES

WT
tor +/-
tor $^{\Delta FRB}$

Fig. 3 The Tor kinase plays a vital role in *A. flavus*.

(A) The structure diagram of the Tor kinase.

(B) Morphology of conidiophores in the WT, *tor*+/-, and *tor*^{ΔFRB}strains observed by microscope.

(C) Phenotype of the WT, *tor*+/-, and *tor*^{ΔFRB}strains grown on PDA amended with 100 ng/mL rapamycin at 37°C for 5 days.

(D) TLC assay of AFB_1 production by the WT, *tor*+/-, and *tor*^{ΔFRB}strains cultured in YES liquid medium at 29°C for 6 days.

(E) The growth inhibition rate of the WT, *tor*+/-, and *tor*^{ΔFRB}strains under rapamycin stress.

(F) AFB_1 quantitative analysis of the WT, *tor*+/-, and *tor*^{ΔFRB}strains described in (D).

Fig. 4 The Sch9 kinase participates in aflatoxin biosynthesis and the HOG pathway.

(A) The structure diagram of the Sch9 kinase.

(B) Phenotype of the WT, $\Delta sch9$, $sch9^{\Delta C2}$, $sch9^{\Delta S\_TKc}$, $sch9^{\Delta S\_TK\_X}$, and $sch9^{K340A}$ strains grown on PDA medium amended with 100 ng/mL rapamycin at 37°C for 5 days.

(C) TLC assay of AFB_1 production from the WT, $\Delta sch9$, $sch9^{\Delta C2}$, $sch9^{\Delta S\_TKc}$, $sch9^{\Delta S\_TK\_X}$, and $sch9^{K340A}$ strains cultured in YES liquid medium at 29°C for 6 days.

(D) The phenotypes of the WT, $\Delta sch9$, $sch9^{\Delta C2}$, $sch9^{\Delta S\_TKc}$, $sch9^{\Delta S\_TK\_X}$, and $sch9^{K340A}$ strains on YGT media amended with 1 M NaCl and 1 M KCl for 3 days.

(E) The phosphorylation levels of Hog1 in the WT and $\Delta sch9$ strains were determined with or without osmotic stress.

(F) The growth inhibition rate of the WT, $\Delta sch9$, $sch9^{\Delta C2}$, $sch9^{\Delta S\_TKc}$, $sch9^{\Delta S\_TK\_X}$, and $sch9^{K340A}$ strains under rapamycin stress.

(G) Quantitative analysis of AFB_1 as shown in (C).

(H) The growth inhibition rate of the WT, $\Delta sch9$, $sch9^{\Delta C2}$, $sch9^{\Delta S\_TKc}$, $sch9^{\Delta S\_TK\_X}$, and $sch9^{K340A}$ strains under osmotic stress.

Fig. 5 TapA and TipA regulate sclerotia formation in *A. flavus*.

(A) Phenotype of the WT, *tapA* +/-, and Δ*tipA* strains grown on PDA amended with 100 ng/mL rapamycin at 37°C for 5 days.

(B) Phenotypic characterization of the WT, *tapA* +/-, and Δ*tipA* strains grown on CM medium at 37°C for 7 days.

(C) Morphology of conidiophores in the WT, *tapA* +/-, and Δ*tipA* strains observed by microscope.

(D) TLC assay of AFB$_1$ production by the WT, *tapA* +/-, and Δ*tipA* strains cultured in YES liquid medium at 29°C for 6 days.

(E) The growth inhibition rate of the WT, *tapA* +/-, and Δ*tipA* strains under rapamycin stress.

(F) Amount of sclerotia produced by the WT, *tapA* +/-, and Δ*tipA* strains.

(G) AFB$_1$ quantitative analysis of the WT, *tapA* +/-, and Δ*tipA* strains described in (D).

Fig. 6 The impact of the phosphatases SitA and Ppg1 on the growth, conidiation, and aflatoxin biosynthesis in *A. flavus*.

(A) Colony morphology of the WT, single knockout, and double knockout strains cultured on PDA medium amended with 100 ng/mL rapamycin at 37°C for 5 days.

(B) Colony morphology of the WT, single knockout, and double knockout strains cultured on WKM medium at 37°C for 7 days.

(C) TLC analysis of AFB_1 production from the WT, single knockout, and double knockout strains cultured in YES liquid medium at 29°C for 6 days.

(D) Growth diameter of the WT, single knockout, and double knockout strains on PDA media.

(E) The growth inhibition rate of the WT, single knockout, and double knockout strains under rapamycin stress.

(F) AFB_1 quantitative analysis of the WT, single knockout, and double knockout strains.

(G) Relative expression levels of conidia synthesis genes in the WT, single knockout, and double knockout strains.

(H) Relative expression levels of sclerotia synthesis genes in the WT, single knockout, and double knockout strains.

(I) Relative expression levels of aflatoxin biosynthesis genes in the WT, single knockout, and double knockout strains.

Fig. 7 Pathogenicity analysis of the phosphatase SitA and Ppg1 in *A. flavus*.

(A) Phenotypes of peanut and maize infected by the WT, single knockout, and double knockout strains.

(B) TLC was used to detect the AFB_1 from peanut and maize infected by the WT, single knockout, and double knockout strains.

(C) Quantitative analysis of conidia production from peanut and maize infected by the WT, single knockout, and double knockout strains.

(D) Quantitative analysis of AFB_1 from peanut and maize infected by the WT, single knockout, and double knockout strains.

A

B

C

D

Fig. 8 Sensitivity of phosphatase SitA and Ppg1 to cell wall damaging agents in *A. flavus*.

(A) Morphology of the WT, single knockout, and double knockout strains grown on YES media supplemented with 200 µg/mL CR, 200 µg/mL CFW, or 100 µg/mL SDS at 37°C for 5 days.

(B) The growth inhibition rate of the WT, single knockout, and double knockout strains under cell wall and cell membrane stress.

(C) Relative expression levels of cell wall synthesis genes (*chsA*, *chsD*, *chsE* and *β-1,6*) in the WT, single knockout, and double knockout strains.

(D) The phosphorylation level of Slt2 in the WT, single knockout, and double knockout strains was detected with or without cell wall stress.

Fig. 9 Phosphatase SitA and Ppg1 are involved in regulating lipid droplet biogenesis.

(A) The intracellular lipid droplets were stained by boron dipyrromethene difluoride (BODIPY) in hyphae of the WT strain and observed under fluorescence microscopy. The phenotype of lipid droplets accumulation in the mycelia of the WT strain treated with or without 100 ng/mL rapamycin for 6 h (Bar, 10 μm).

(B) Phenotype of lipid droplets accumulation in the mycelia of the Δ*sch9* strain treated with or without 100 ng/mL rapamycin for 6 h (Bar, 10 μm).

(C) Phenotype of lipid droplets accumulation in the mycelia of the Δ*sitA* strain treated with or without 100 ng/mL rapamycin for 6 h (Bar, 10 μm).

(D) Phenotype of lipid droplets accumulation in the mycelia of the Δ*ppg1* strain treated with or without 100 ng/mL rapamycin for 6 h (Bar, 10 μm).

Fig. 10 Phosphatase complex Nem1/Spo7 play significant role in growth, conidiation, aflatoxin, and lipid droplet biogenesis.

(A) Phenotype of the WT, Δ*nem1*, and Δ*spo7* strains grown on PDA amended with 100 ng/mL rapamycin at 37°C for 5 days.

(B) TLC assay of AFB$_1$ production from the WT, Δ*nem1*, and Δ*spo7* strains cultured in YES liquid medium at 29°C for 6 days.

(C) Growth diameter of the WT, Δ*nem1*, and Δ*spo7* strains on PDA media.

(D) Statistical analysis of the sporulation by the WT, Δ*nem1*, and Δ*spo7* strains.

(E) Phenotype of lipid droplets accumulation in the mycelia of the WT, Δ*nem1*, and Δ*spo7* strains treated with or without 100 ng/mL rapamycin for 6 h (Bar, 10 μm).

Vegetative development, stress responses, aflatoxin biosynthesis, and pathogenicity.

Fig. 11 The proposed model of the target of rapamycin (TOR) pathway in *A. flavus*.

Based on the above results, we propose a hypothesis regarding the TOR signaling pathway model in *A. flavus*. Rapamycin forms a complex with Fkbp3, and this particular complex can bind to the Tor kinase, thereby impeding its regular functionality. Sch9, functioning as a downstream element of the Tor kinase, regulates aflatoxin biosynthesis and HOG signaling pathway. As another important target of the Tor kinase, TapA-phosphatase complex is involved in the regulation of growth, sporulation, sclerotia, aflatoxin production, CWI signaling pathway, lipid droplet synthesis, and other processes. In conclusion, the TOR signaling pathway plays a crucial role in various aspects of *A. flavus*, including vegetative development, stress response, aflatoxin biosynthesis, and pathogenicity.