The TOR signaling pathway regulates vegetative development, aflatoxin biosynthesis, and pathogenicity in *Aspergillus flavus*

Guoqi Li, Xiaohong Cao, Elisabeth Tumukunde, Qianhua Zeng, Shihua Wang[*]

State Key Laboratory of Ecological Pest Control for Fujian and Taiwan Crops, Key Laboratory of

Pathogenic, Fungi and Mycotoxins of Fujian Province, School of Life Sciences, Fujian

Agriculture and Forestry University, Fuzhou 350002, China

Corresponding author:

Name: Shihua Wang Professor

School of Life Sciences,

Fujian Agriculture and Forestry University

Fuzhou, Fujian, 350002, China

Phone/Fax: +86-591-83789312

E-mail: wshyyl@sina.com

Abstract

The target of rapamycin (TOR) signaling pathway is highly conserved and plays a crucial role in diverse biological processes in eukaryotes. Despite its significance, the underlying mechanism of the TOR pathway in *A. flavus* remains elusive. In this study, we comprehensively analyzed the TOR signaling pathway in *A. flavus* by identifying and characterizing nine genes that encode distinct components of this pathway. The FK506-binding protein Fkbp3 and its lysine succinylation are important for aflatoxin production and rapamycin resistance. The TorA kinase plays a pivotal role in the regulation of growth, spore production, aflatoxin biosynthesis, and responses to rapamycin and cell membrane stress. As a significant downstream effector molecule of the TorA kinase, the Sch9 kinase regulates aflatoxin B_1 (AFB_1) synthesis, osmotic and calcium stress response in *A. flavus,* and this regulation is mediated through its S_TKc, S_TK_X domains, and the ATP binding site at K340. We also showed that the Sch9 kinase may have a regulatory impact on the high-osmolarity glycerol (HOG) signaling pathway. TapA and TipA, the other downstream components of the TorA kinase, play a significant role in regulating cell wall stress response in *A. flavus*. Moreover, the members of the TapA-phosphatase complexes, SitA and Ppg1, are important for various biological processes in *A. flavus*, including vegetative growth, sclerotia formation, AFB_1 biosynthesis, and pathogenicity. We also demonstrated that SitA and Ppg1 are involved in regulating lipid droplets (LDs) biogenesis and cell wall integrity (CWI) signaling pathways. In

addition, another phosphatase complex, Nem1/Spo7, plays critical roles in hyphal

development, conidiation, aflatoxin production, and lipid droplets biogenesis.

Collectively, our study has provided important insight into the regulatory network of

the TOR signaling pathway and has elucidated the underlying molecular mechanisms

of aflatoxin biosynthesis in *A. flavus*.

Keywords: *Aspergillus flavus*; the TOR signaling pathway; aflatoxin; vegetative

development

Introduction

Aspergillus flavus is a highly significant phytopathogenic fungus that frequently

contaminates a diverse array of agricultural crops, leading to substantial economic

losses in the food and agriculture industry and posing considerable health risks,

particularly in developing countries [1]. Moreover, *A. flavus* is recognized as an

opportunistic human pathogenic fungus that can cause aspergillosis in

immune-compromised patients [2]. The fungus is notorious for its production of

aflatoxins (AFs), which are potent secondary metabolites with toxic and carcinogenic

properties and are classified among the most powerful carcinogens known in nature,

with established links to liver cancer in both humans and animals [3]. Aflatoxin

biosynthesis is a multifaceted and intricate biochemical process encompassing a

cascade of enzymatic reactions. Although the majority of the enzyme reactions and

genes involved in aflatoxin biosynthesis have been elucidated [4], the related signaling

pathway network and specific regulatory mechanisms governing aflatoxin

biosynthesis remain elusive. Therefore, exploring the regulatory mechanism of the

aflatoxin biosynthesis signaling pathway would provide new insights for preventing *A.*

flavus and aflatoxin contamination.

In eukaryotes, the target of rapamycin (TOR) signaling pathway is highly

conserved and plays essential roles in various significant biological processes,

including ribosome biosynthesis, cell growth, and autophagy [5]. The

serine/threonine-protein kinase Tor serves as a crucial protein component in the TOR

signaling pathway, which interacts with other proteins to form multi-protein

complexes [6]. In *Saccharomyces cerevisiae*, TOR exists in two distinct multiprotein

functional complexes, namely TORC1 and TORC2 [7]. TORC1-mediated signaling

governs cellular growth by modulating various growth-related mechanisms and

exhibits sensitivity to rapamycin. Conversely, TORC2-mediated signaling primarily

regulates cytoskeletal remodeling but remains unaffected by rapamycin [7]. In yeast,

there are two *tor* genes, *tor1* and *tor2*, whereas in higher eukaryotes such as plants,

animals, and filamentous fungi, there is only one *tor* gene [8-10].

In the fungal kingdom, the TOR signaling pathway has been extensively

investigated in three prominent species: the budding yeast *S. cerevisiae* [11-12], the

fission yeast *Schizosaccharomyces pombe* [13-15], and the human pathogen *Candida*

albicans [16-17]. In *S. cerevisiae*, the macrolide antibiotic rapamycin forms a complex

with the cytosolic protein Fkbp12, which effectively inhibits the catalytic function of

the Tor kinase by interacting with its FRB domain [18]. The Tap42 phosphatase

complex serves as the primary target of the Tor kinase and plays a crucial role in the

regulation of downstream effectors. Additionally, the Tap42 protein interacts with the

Tip41 protein to collaboratively modulate the activities of phosphatases, including

PP2A and Sit4 [19-20]. Among these, the Sit4 phosphatase is known to

dephosphorylate the transcription factor Gln3, thereby regulating the nitrogen source

metabolism pathway [21]. In addition, the AGC kinase Sch9, an immediate effector of

TORC1, is involved in regulating various cellular functions processes, such as

ribosome biosynthesis, translation initiation, and entry into the G_0 phase [22]. The TOR

signaling pathway has been documented in various filamentous fungi, including

Fusarium graminearum [23-24], *Magnaporthe oryzae* [25], *Phanerochaete*

chrysosporium [26], *Podospora anserine* [27], *A. nidulans* [28], *A. fumigatus* [29-30], *F.*

fujikuroi [31], *Botrytis cinerea* [32], *F. oxysporum* [33], and *Mucor circinelloides* [34].

However, the TOR signaling pathway in *A. flavus* remains unexplored.

In eukaryotic cells, the TOR signaling pathway and the mitogen-activated protein

kinase (MAPK) signaling pathways play an important role in regulating adaptive

responses to extra- and intracellular conditions [35]. Several studies have demonstrated

that the TOR signaling pathway interacts with various other signaling pathways, such

as MAPK and cell wall integrity (CWI) pathway. These pathways are recognized to

participate in crosstalk, forming a complex metabolic network that regulates cellular

processes [36-40]. However, the underlying mechanisms of multiple crosstalks between

the TOR and other signaling pathways in *A. flavus* remain largely unknown. Therefore,

we intended to identify the genes associated with the TOR signaling pathway and

elucidate their roles and contributions in regulating vegetative development and

aflatoxin biosynthesis in *A. flavus*. Our results demonstrated that the TOR signaling

pathway is involved in multiple cellular processes in *A. flavus*. Notably, our findings

revealed a complex interplay between the TOR pathway and other signaling pathways

(such as HOG and CWI) in *A. flavus*, which play a crucial role in regulating cellular

growth and survival under environmental stress conditions.

Results

Rapamycin exhibits inhibitory effects on the growth, sporulation, sclerotia

formation, and aflatoxin production in *A. flavus*

Rapamycin, a secondary metabolite synthesized by *Streptomyces hygroscopicus*,

exhibits efficacy in suppressing specific filamentous fungi. In *F. graminearum*,

rapamycin has a significant inhibitory impact on growth and asexual reproduction [23].

To investigate the effects of rapamycin on vegetative development, sclerotia

formation, and secondary metabolism in *A. flavus*, we conducted an assay to

determine the sensitivity of the wild-type (WT) strain to rapamycin. We observed that

the WT strain exhibited a high sensitivity to rapamycin. We found that 100 ng/mL of

rapamycin significantly inhibited mycelial growth and conidia formation (Fig. 1A, 1D,

1E). More importantly, the WT strain exhibited a significant reduction in sclerotia

formation and aflatoxin synthesis (Fig. 1B, 1C, 1F, 1G). Additionally, rapamycin

resulted in a significant decrease in spore germination rate and sparser conidiophores

(Fig. S1A, S1B). These results illustrated that the TOR signaling pathway may be

involved in regulating the growth, development, and secondary metabolism of *A.*

flavus. Calcofluor white is a chemifluorescent blue dye commonly employed for its nonspecific binding affinity to fungal chitin. Fluorescence microscopy analysis revealed a notable reduction in the blue fluorescence on the cell wall following treatment with rapamycin (Fig. S1C). The intracellular lipid droplets within the hyphae were labeled with BODIPY. We observed that the hyphae treated with rapamycin contained more lipid droplets compared to the control group (Fig. S1D). These findings suggest that the TOR pathway may play important roles in maintaining cell wall integrity and facilitating the biogenesis of lipid droplets.

To investigate the potential regulatory role of the TOR pathway in modulating the sporulation, sclerotia formation, and aflatoxin biosynthesis in *A. flavus*, we examined the expression levels of conidia-related, sclerotia formation-related, and aflatoxin-related genes after 3, 6, and 9 hours of treatment with rapamycin. To determine the effects of rapamycin on conidiation, qRT-PCR was conducted to assess the transcript levels of two conidia-related genes, *abaA* and *brlA*. We also conducted qRT-PCR to evaluate the expression levels of three regulators (*nsdC*, *nsdD*, and *sclR*) involved in sclerotia development, and genes involved in AFB$_1$ biosynthesis, such as the regulatory gene *aflS*, as well as several structural genes (*aflC* and *aflQ*). The strains treated with rapamycin exhibited a significant reduction in the expression levels of genes related to sporulation, sclerotia formation, and aflatoxin synthesis (Fig. 1H, 1I, 1J). Overall, these results indicated that the TOR pathway plays a crucial role in regulating the growth, sporulation, sclerotia formation, and aflatoxin biosynthesis in *A. flavu*s by modulating the expression of these genes.

The impact of Fkbp3 and its lysine succinylation on AFB$_1$ biosynthesis and rapamycin resistance in *A. flavus*

FK506-binding proteins (Fkbps) belong to a highly conserved immunophilin family, which are involved in various cellular functions, such as protein folding, immunosuppression, signaling transduction, and transcription [41]. Fkbps function as molecular switches by binding to target proteins and inducing conformational changes [41]. In *S. cerevisiae*, Fkbp12 and rapamycin interact to form a complex that exerts a negative regulatory effect on the activity of the Tor kinase [42]. The *fkbp12* gene homologue *fprA* in *A. nidulans* shares high levels of homology with that from *S. cerevisiae* [28]. According to the protein sequence of FprA from *A. nidulans*, four genes encoding putative proteins with FK506 binding domain, named *fkbp1* (AFLA_126880), *fkbp2* (AFLA_087480), *fkbp3* (AFLA_010910), and *fkbp4* (AFLA_128200), have been identified from the *A. flavus* genome. We generated four *fkbp* knockout strains using homologous recombination, and all *fkbp* disruption strains were confirmed by PCR and sequencing analysis (Fig. S2). Phenotypic analyses showed that the mycelial growth and conidia formation of all the *fkbp* mutants were similar to the WT strain on the PDA medium (Fig. 2A), suggesting that Fkbps have minimal influence on the process of vegetative development. We evaluated AFB$_1$ synthesis utilizing thin layer chromatography (TLC), and the findings demonstrated a noteworthy reduction in AFB$_1$ synthesis upon deletion of Fkbp3, in comparison to the WT strain (Fig. 2B, 2D). Additionally, the quantity of sclerotia exhibited a significant decrease in all *fkbp* mutants compared to the WT strain (Fig. S3A, S3B). The

aforementioned data indicated that Fkbp3 plays a crucial role in sclerotia formation

and aflatoxin biosynthesis in *A. flavus*.

To determine the specific Fkbp responsible for rapamycin sensitivity, all mutant

strains were cultured on PDA solid plates amended with 100 ng/mL rapamycin. The

strain with disrupted *fkbp3* showed significant resistance to rapamycin, while the

deletion mutants of *fkbp1*, *fkbp2*, and *fkbp4* did not exhibit any alteration in resistance

to rapamycin (Fig. 2A, 2C). These results demonstrated that Fkbp3 serves as the

principal target for rapamycin in *A. flavu*s. In the FK506-sensitivity analysis, we

observed a significant suppression of mycelial growth and conidia formation in WT

strain with the addition of 10 ng/mL FK506 to the PDA medium. Additionally, the

deletion of *fkbp3* led to the heightened resistance to FK506 compared to the WT strain

(Fig. 2A, 2C). These findings suggested that Fkbp3 is mainly involved in the

modulation of FK506 resistance in *A. flavus*.

Previous studies have shown that lysine succinylation plays a significant role in

aflatoxin production and pathogenicity in *A. flavus* [43]. Based on the analysis of our

succinylome data, we have identified six reliable succinylation sites (K5, K19, K40,

K42, K55, and K65) on the Fkbp3 protein. To identify the function of Fkbp3

succinylation sites, site-directed mutations were constructed according to the

homologous recombination strategy. Compared to the WT strain, all point mutants

exhibited a similar phenotype in vegetative growth and conidiation. Interestingly, we

observed that the K19A mutation resulted in increased resistance to rapamycin (Fig.

2E, 2F). Additionally, the K19A strain exhibited a significant reduction in the

production of AFB_1 compared to the WT strain (Fig. 2G, 2H). These findings

suggested that the succinylation of Fkbp3 at K19 plays a crucial role in rapamycin

resistance and aflatoxin biosynthesis. Furthermore, we found that various point

mutants, including K40A, K42A, and K55A, exhibited reduced levels of AFB_1

production in comparison to the WT strain (Fig. 2G, 2H), suggesting that the

succinylation of Fkbp3 at other sites also contributes to AFB_1 biosynthesis in *A. flavus*.

The above results indicated that the K19 site likely plays a more prominent role in

Fkbp3 succinylation compared to other sites.

The TorA kinase plays a critical role in *A. flavus*

In *S. cerevisiae*, the Tor kinase is known to interact with a diverse range of

proteins, resulting in the formation of complex structures known as TORC1 and

TORC2 [44]. TORC1 can detect various signals, including nutrients, growth factors,

and environmental stress. It plays a crucial role in regulating cellular processes such

as gene transcription, protein translation, ribosome synthesis, and autophagy [45]. To

identify the ortholog of the Tor protein in *A. flavus*, the *S. cerevisiae* Tor protein was

utilized as search reference in the *A. flavus* genome database using the Basic Local

Alignment Search Tool (https://blast.ncbi.nlm.nih.gov/Blast.cgi). The genome of *A.*

flavus contains a solitary ortholog of the *tor* gene (AFLA_044350), which encodes a

protein exhibiting 48.91% similarity to *S. cerevisiae* Tor2. To elucidate the role of

torA in *A. flavus*, we attempted to obtain *torA* deletion strains by homologous

recombination, but not successful. As an alternative approach, a mutant strain with a

xylose promoter mutation, referred as $^{xylP}torA$, was constructed to validate the

function of *torA*. Subsequent confirmation via PCR analysis ensured the integrity of the mutant strain. Xylan and xylose are highly effective inducers, whereas glucose strongly represses the activity of the promoter. To ascertain the impact of *torA* on *A. flavus*, we conducted a comparative analysis of *torA* transcript levels between the $^{xylP}torA$ strain and the WT strain. Our findings revealed that the *torA* transcript levels in the $^{xylP}torA$ strain were markedly reduced compared to the WT strain (Figure S4A). Both the WT strain and $^{xylP}torA$ mutant were cultured in YXT medium supplemented with xylose and YGT medium devoid of xylose for a duration of 5 days at 37℃ under dark conditions. Remarkably, the $^{xylP}torA$ strain exhibited complete inability to thrive in the absence of xylose, with only partial restoration of growth observed upon xylose supplementation (Fig. 3A, 3D). Notably, the $^{xylP}torA$ strain also demonstrated a markedly reduced growth rate on YXT compared to the WT strain (Fig. 3A, 3C). Thin-layer chromatography (TLC) assays indicated a noteworthy decrease in AFB_1 synthesis in the $^{xylP}torA$ strain compared to the WT strain (Figure 3B, 3F). Additionally, the $^{xylP}torA$ strain produced fewer conidia and sclerotium formation was absent on YXT medium (Fig. 3E, S5). Interestingly, the $^{xylP}torA$ strain exhibited a notable increase in sensitivity to rapamycin and SDS stress (Figure 3G, 3H). Collectively, these findings suggested that the TorA kinase plays a crucial role in various biological processes in *A. flavus*, including vegetative growth, asexual development, sclerotia formation, response to environmental stress, and biosynthesis of aflatoxins.

The Sch9 kinase is involved in aflatoxin biosynthesis and the HOG pathway

Sch9, a Ser/Thr kinase belonging to the AGC family, is directly phosphorylated by

TORC1. The phosphorylation of Sch9 is inhibited by rapamycin, as well as by carbon

or nitrogen starvation [22]. The putative *sch9* gene (AFLA_127440) in *A. flavus*

encodes a protein consisting of 706 amino acids, exhibiting 80.31% identity to the

Sch9 protein in *A. nidulans*. To investigate the role of Sch9 in *A. flavus*, we attempted

to create a *sch9* deletion mutant using a homologous recombination strategy. The

Δ*sch9* strain exhibited typical vegetative growth and conidiation similar to the WT

strain (Fig. 4B). TLC assay and quantitative analysis showed a significantly decreased

aflatoxin production in the Δ*sch9* strain compared to the WT strain (Fig. 4C, 4G).

These results indicated that Sch9 regulates aflatoxin biosynthesis in *A. flavus*.

In yeast, Sch9 plays an important role in the transcriptional activation of

osmostress inducible genes [46]. To characterize the roles of Sch9 in stress response,

we examined the sensitivity of the *sch9* deletion strain to various stresses including

osmotic stress and calcium stress. Compared with the WT strain, the Δ*sch9* mutant

significantly increased sensitivity to calcium stress (Fig. S6A, S6B), and decreased

sensitivity to osmotic stress induced by NaCl and KCl (Fig. 4D, 4H). In *A. flavus*, the

Sch9 kinase consists of several domains, including the protein kinase C conserved

region 2 (C2), protein kinase (S_TKc), AGC-kinase C-terminal (S_TK_X) domain,

and an ATP binding site (Fig. 4A). To explore the role of these domains and the ATP

binding site in Sch9 kinase, we constructed deletion strains for both domains and a

sch9$^{\text{K340A}}$ mutant. The results revealed that the *sch9*$^{\Delta \text{S\_TKc}}$ strain, *sch9*$^{\Delta \text{S\_TK\_X}}$ strain, and

sch9$^{\text{K340A}}$ mutant exhibited similar phenotypes in osmotic stress and AFB$_1$ production

as the Δ*sch9* mutant (Fig. 4B, 4C, 4D). Therefore, we speculate that the protein kinase domain, kinase C-terminal domain, and the ATP binding site at K340 play a crucial role in modulating the impact of Sch9 on aflatoxin biosynthesis and stress response in *A. flavus*.

To investigate the potential function of Sch9 in the HOG and TOR pathways in *A. flavus*, we conducted a sensitivity analysis of the Δ*sch9* strain to rapamycin. Compared to the WT strain, the *sch9* deletion strain exhibited a slightly higher sensitivity to rapamycin (Fig. 4B, 4F). The MAKP Hog1 is an element of the high osmolarity glycerol (HOG) pathway. To verify the connection between Sch9 and the HOG-MAPK pathway, the phosphorylation levels of Hog1 kinase were measured under osmotic stress (1M NaCl). Western blot analysis showed that the phosphorylation level of Hog1 increased significantly in the Δ*sch9* strain (Fig. 4E). This finding suggested that Sch9 plays a pivotal role in modulating the response of Hog1 to osmotic stress. Taken together, these results reflected that Sch9 regulates osmotic stress response via the HOG pathway in *A. flavus*.

TapA and TipA participate in the regulation of cell wall stress response

Tap42, a protein that associates with phosphatase 2A, serves as the direct target of the Tor kinase in yeast. Tap42 plays a crucial role in various Tor functions, particularly in the regulation of transcriptional processes [47]. The genome of *A. flavus* contains a homolog of *tap42* named *tapA* (AFLA_092770), which is predicted to encode a 355 amino-acid protein (sharing 31.9% identity with *S. cerevisiae* Tap42). To further determine the function of the TapA regulator in the TOR signaling pathway

of *A. flavus*, we used homologous recombination to replace the original promoter with the *gpdA* promoter, a well-characterized strong promoter, to construct an overexpression mutant (*OE*::*tapA*). The transformants of the mutant were verified using PCR. Quantitative expression analysis of the *tapA* gene in the overexpressed strains revealed a substantial upregulation compared to the WT strain (Fig. S4B). Our results showed that the colony diameter of the *OE*::*tapA* mutant was not significantly changed compared to the WT strain (Fig. 5A). However, the rapamycin sensitivity assay indicated that the *OE*::*tapA* strain displayed increased sensitivity to rapamycin (Fig. 5A, 5D). We found no significant difference in aflatoxin production between the *OE*::*tapA* and the WT strains (Fig. 5C, 5E), suggesting that TapA may not play a direct role in the regulation of aflatoxin biosynthesis. In addition, the *OE*::*tapA* strain was more sensitive to cell wall and cell membrane stress, including SDS, CFW, and CR (Fig. 5B, 5F). These findings have demonstrated the significantly positive impact of TapA on cell wall stress in *A. flavus*.

In *S. cerevisiae*, Tip41 was identified as a Tap42-interacting protein and has been measured to function as a negative regulator of Tap42, which downregulates TORC1 signaling via activation of PP2A phosphatase [48]. The putative orthologue of *tip41* in *A. flavus* is named *tipA* (AFLA_047310). The amino acid sequence of TipA exhibits a similarity of 40.49% to Tip41 in *S. cerevisiae*. To elucidate the function of the TOR signaling pathway protein TipA in *A. flavus*, we generated the *tipA* deletion mutants. The Δ*tipA* strain exhibited a slightly reduced number of conidia produced on PDA compared to the WT strain (Fig. 6A, 6C). Meanwhile, the Δ*tipA* strain exhibited

decreased sensitivity to rapamycin (Fig. 6A, 6E). The Δ*tipA* strain was unable to

produce any sclerotia (Fig. 6B, 6F). Whereas Δ*tipA* strain do not affect aflatoxin

production(Fig. 6D, 6G). In addition, the Δ*tipA* strain was more sensitive to the cell

wall and cell membrane stress, including CR and SDS (Fig. 6H, 6I). Additionally, the

expression levels of *nsdC* and *nsdD* exhibited a decrease in Δ*tipA* strain compared to

the WT strain (Fig. 6J). These results indicated that TipA is crucial for the sclerotial

formation and cell wall stress in *A. flavus*.

Phosphatase SitA and Ppg1 are involved in growth, conidiation, and sclerotial

formation

In *S. cerevisiae*, Tap42 has been identified to interact with the catalytic subunit of

type 2A protein phosphatases, including Pph3, Pph21, and Pph22, as well as the type

2A-like phosphatases Sit4 and Ppg1[19]. Phosphatase *sitA* (AFLA_108450) and *ppg1*

(AFLA_132610) were identified in *A. flavu*s, encoding putative orthologs of *sit4* and

ppg1 in *S. cerevisiae.* To determine the biological functions of phosphatase SitA and

Ppg1 in *A. flavu*s, we disrupted the *sitA* and *ppg1* by homologous recombination

strategy. To further investigate the relationship between phosphatase SitA and Ppg1,

we additionally generated complemented and double deletion mutants. The vegetative

development phenotype analysis showed that the Δ*sitA*, Δ*ppg1*, and Δ*sitA*/Δ*ppg1*

strains grew significantly more slowly than the WT strain on PDA (Fig. 7A, 7D).

Additionally, the Δ*sitA*, Δ*ppg1*, and Δ*sitA*/Δ*ppg1* strains were more sensitive to

rapamycin (Fig. 7A, 7E). In addition, the Δ*sitA*, Δ*ppg1*, and Δ*sitA*/Δ*ppg1* strains

formed lower number of conidiophores (Fig. 7A). To gain further insight into the role

of SitA and Ppg1 in the process of conidiation, qRT-PCR was employed to determine

the transcript levels of conidiation-related genes (*abaA* and *brlA*). The expression

levels of the *abaA and brlA* genes were significantly downregulated in the $\Delta sitA$,

$\Delta ppg1$, and $\Delta sitA/\Delta ppg1$ mutants when compared to the WT and complemented

strains (Fig. 7G). The analysis of sclerotial formation revealed that the $\Delta sitA$, $\Delta ppg1$,

and $\Delta sitA/\Delta ppg1$ mutants exhibited a complete inability to produce any sclerotia (Fig.

7B, 7E). To further prove these findings, we conducted qRT-PCR to determine the

expression levels of three regulators (*nsdC*, *nsdD*, and *sclR*) of sclerotia development.

Our findings indicated that the expression levels of *nsdC*, *nsdD*, and *sclR* were all

significantly reduced in the $\Delta sitA$, $\Delta ppg1$, and $\Delta sitA/\Delta ppg1$ mutants (Fig. 7H), which

is consistent with the observed decrease in sclerotia production in the $\Delta sitA$, $\Delta ppg1$,

and $\Delta sitA/\Delta ppg1$ mutants. These results showed that SitA and Ppg1 are crucial for

hyphal development, conidiation, and sclerotia formation in *A. flavus*.

SitA and Ppg1 play crucial roles in aflatoxin biosynthesis and pathogenicity

We assayed AFB_1 biosynthesis in the $\Delta sitA$, $\Delta ppg1$, and $\Delta sitA/\Delta ppg1$ mutants.

TLC analyses revealed that the $\Delta sitA$ and $\Delta sitA/\Delta ppg1$ strains exhibited a decrease in

AFB_1 production, whereas the $\Delta ppg1$ strain showed a significant increase in AFB_1

production compared to the WT and complemented strains (Fig. 7C, 7F). The

expression levels of genes related to aflatoxin synthesis, including structural genes

(*aflO* and *aflP*) and regulatory genes (*aflR* and *aflS*), were determined by qRT-PCR in

the $\Delta sitA$, $\Delta ppg1$, and $\Delta sitA/\Delta ppg1$ mutants. The results revealed a significant

downregulation of these genes in the $\Delta sitA$ and $\Delta sitA/\Delta ppg1$ mutants, while

upregulated in the Δ*ppg1* mutants, compared to the WT and complemented strains

(Fig. 7I). These results indicated that SitA and Ppg1 play different roles in regulating

aflatoxin biosynthesis in *A. flavu*s.

We conducted pathogenicity tests on peanuts and maize to examine the

pathogenicity of the phosphatases SitA and Ppg1 on crops. The results showed that

the WT and complemented strains demonstrated complete virulence on all peanut and

maize seeds. In contrast, the Δ*sitA*, Δ*ppg1*, and Δ*sitA*/Δ*ppg1* mutants displayed

compromised colonization on peanut and maize seeds (Fig. 8A). In addition, the

number of conidial productions in these mutants on the infected peanut seeds also

dramatically decreased compared to those of the WT and complemented strains (Fig.

8C). The thin-layer chromatography analyses revealed that the Δ*ppg1* mutant

exhibited increased production of AFB_1 in peanuts and maize. Conversely, the levels

of aflatoxin produced by the Δ*sitA* and Δ*sitA*/Δ*ppg1* mutants on peanut and maize

were significantly reduced compared to the WT and complemented strains (Fig. 8B,

8D). This observation aligns with the above results of aflatoxin biosynthesis from all

mutant strains in the YES medium. All these data showed that SitA and Ppg1 are

crucial for crop seeds' pathogenicity.

SitA and Ppg1 are involved in regulating cell wall integrity

To explore the role of phosphatases SitA and Ppg1 in response to cell wall stress,

all strains were cultivated on PDA amended with CFW, CR, and SDS. We found that

the relative growth inhibition of Δ*sitA*, Δ*ppg1*, and Δ*sitA*/Δ*ppg1* mutants induced by

cell wall stress was significantly increased compared to the WT and complemented

strains (Fig. 9A, 9B). Further investigations using qRT-PCR revealed a notable

reduction in the expression levels of genes related to the synthesis of cell walls,

specifically chitin synthase genes *chsA*, *chsD*, and *chsE* in the Δ*sitA*, Δ*ppg1*, and

Δ*sitA*/Δ*ppg1* strains (Fig. 9C). Western blot analysis also revealed that the

phosphorylation level of Slt2 was significantly elevated in the Δ*sitA*, Δ*ppg1*, and

Δ*sitA*/Δ*ppg1* strains under cell wall stress (200 μg/ml CFW) (Fig. 9D). In conclusion,

the phosphatases SitA and Ppg1 play a crucial role in regulating cell wall integrity.

The phosphatase complex Nem1/Spo7 is important for growth, aflatoxin

biosynthesis, and LD biogenesis

Lipid droplets (LDs) are highly dynamic spherical organelles, which appear in

the cytosol of most eukaryotic cell types[49]. Rapamycin treatment led to the

accumulation of lipid droplets in various filamentous fungi [24]. The FgTOR pathway

regulates the phosphorylation status of FgNem1 mainly via the

FgPpg1/Sit4-FgCak1/other kinase cascade in *F. graminearum* [24]. In this study, we

observed that the treatment of hyphae with rapamycin resulted in the accumulation of

LD biogenesis (Fig. 10A). This finding suggested that the TOR pathway may regulate

the LD biogenesis. However, the underlying regulatory mechanism of LD

biosynthesis remain largely unknown in *A. flavu*s. To investigate the regulatory

mechanism of the TOR pathway in *A. flavus* that controls LD biogenesis, we

examined LD accumulation in Δ*sch9*, Δ*sitA*, and Δ*ppg1* mutants. BODIPY staining

assays revealed that LD biogenesis was not observed in the Δ*sitA* and Δ*ppg1* strains

upon treatment with rapamycin, in contrast to the WT and Δ*sch9* strains (Fig. 10A,

10B, 10C, 10D). These results indicated that LD biogenesis induced by rapamycin is

predominantly reliant on the phosphatases SitA and Ppg1 in *A. flavus*. In *F.*

graminearum, the phosphatase FgNem1/FgSpo7 complex was involved in the

induction of LD biogenesis by rapamycin[24]. Phosphatase *nem1* (AFLA_002649) and

spo7 (AFLA_028590) were identified in *A. flavu*s, which encode putative orthologs of

nem1 and *spo7* in *F. graminearum.* To examine the role of the *nem1* and *spo7* genes in

A. flavus, we generated Δ*nem1* and Δ*spo7* mutants using homologous recombination.

The Δ*nem1* and Δ*spo7* mutants displayed a notable reduction in both vegetative

growth and conidiation compared to the WT strain (Fig. 11A, 11C, 11D). TLC assay

revealed a significantly decreased aflatoxin production in the Δ*nem1* and Δ*spo7*

compared to the WT strain (Fig. 11B). In addition, BODIPY staining showed no

visible lipid droplets in the hyphae of Δ*nem1* and Δ*spo7* after rapamycin treatment

(Fig. 9E). Taken together, these results indicated that Nem1/Spo7 are involved in

vegetative growth, conidiation, aflatoxin production, and LD biogenesis.

Discussion

Aflatoxins (AFs) are a class of toxic secondary metabolites synthesized by *A.*

flavus, including aflatoxin B_1, B_2, G_1, and G_2, among which AFB_1 has been regarded

as the most pathogenic and toxic aflatoxin[50]. Therefore, understanding the regulatory

mechanisms involved in fungal secondary metabolites biosynthesis would be

instrumental for effectively addressing aflatoxin control in *A. flavus*. In this study, we

demonstrated that the target of rapamycin (TOR) signaling pathway plays a pivotal

role in regulating growth, sporulation, sclerotia formation, aflatoxin production, and

various stress responses in *A. flavus*. Through conducting functional studies on genes

within the TOR signaling pathway, we explored the conservation and complexity

exhibited by the TOR signaling pathway in *A. flavu*s. In addition, we constructed a

crosstalk network between the TOR and other signaling pathways (HOG, CWI) and

analyzed the specific regulatory process. We proposed that the TOR signaling

pathway interacts with multiple signaling pathways, including HOG and CWI

pathways, to modulate the cellular responses to diverse environmental stresses (Fig.

10).

Our findings indicated that the TOR signaling pathway is pivotal in *A. flavus*,

we have observed that rapamycin has a significant impact on the growth and

sporulation of *A. flavus* (Fig. 1A). Additionally, it exhibits a strong inhibitory effect on

sclerotia production and aflatoxin biosynthesis (Fig. 1B, 1C). The inhibitory effect has

been documented in various fungal species, including *Candida albicans* [51],

Cryptococcus neoformans [51], *Podospora anserine* [27], *A. nidulans* [28], *Magnaporthe*

oryzae [39], and *F. graminearum* [23]. This suggested that the TOR signaling pathway

may have a conserved function in filamentous fungi. The number of FK506-binding

proteins varies among different species: there are four Fkbps proteins in *S. cerevisiae*

and three Fkbps proteins in *S. pombe* [52]. In *S. cerevisiae*, rapamycin can bind to Fkbp

protein, resulting in the irreversible inhibition of the G_1 phase of the cell cycle and the

regulation of cell growth [53]. In *A. nidulans,* the *fkbp12* gene homologue *fprA* displays

considerable sequence similarity with homologues identified in the *S. cerevisiae* [28].

In this study, based on the homologous alignment of the protein sequence of FprA, it

was observed that four *fkbp* genes are present in *A. flavus*. Among these, deletion of

fkbp3 exhibited enhanced tolerance to rapamycin (Fig. 2A). So we speculated that

rapamycin forms a complex with Fkbp3 in *A. flavus*, subsequently targeting

downstream target genes to exert its biological function. We found that Fkbp3 is

involved in sclerotia formation and aflatoxin biosynthesis in *A. flavus* (Fig. 2A, S3),

and subsequent analysis revealed that the underlying mechanism may be associated

with succinylation modification. The succinylation sites of Fkbp3 were identified

through the utilization of succinylation proteomics data. By constructing point mutant

strains and conducting phenotype experiments, we found that the succinylation of

Fkbp3 at residue K19 is involved in rapamycin resistance and aflatoxin biosynthesis

(Fig. 2E, 2G). Our results provided new insights to the understanding of

post-translational modifications in the regulation of TOR pathway components.

Previous studies have demonstrated that Fkbp12 has been identified as the receptor

for FK506 and rapamycin, which inhibits calcineurin and target rapamycin complex 1

(TORC1), respectively [53]. The deletion of Fkbp12 confers the resistance to

rapamycin in *C. albicans* [51] and *C. neoformans* [51]. The entomopathogenic fungus

Beauveria bassiana contains three putative *fkbp* genes, and disruption of *Bbfkbp12*

significantly increases the resistance to rapamycin and FK506 [54]. The plant

pathogenic fungus *F. graminearum* harbors three Fkbps proteins, among which

deletion of *Fgfkbp12* leads to resistance to rapamycin [23]. Thus, we propose that these

Fkbps may exhibit analogous functionalities in conferring resistance to rapamycin. In

Botrytis cinerea, deletion of *Bcfkbp12* caused a reduction of the virulence of strain T4,

while knockout of the *fkbp12* gene did not affect the pathogenic development of strain

B05.10 [32]. These findings demonstrated that the *fkbp12* homologous genes exhibited

conservation in terms of rapamycin resistance. However, there is significant variation

in other biological functions among fungal species, which may be attributed to species

or strain evolutionary divergence. Our study indicated the pivotal role of Fkbp3 in

modulating the effects of rapamycin on *A. flavus*, Fkbp3 is also crucial for aflatoxin

biosynthesis. Targeted inhibition of Fkbp3 could potentially suppress aflatoxin

production, given its significance, Fkbp3 can be developed as a drug target to reduce

aflatoxin biosynthesis, thereby reducing the risk of aflatoxin contamination in

agricultural products.

The Tor kinase functions as a fundamental component of the TOR signaling

pathway, which exhibits evolutionary conservation across fungi, plants, and mammals

[55]. In *S. cerevisiae*, the Tor kinase can interact with a diverse range of proteins,

resulting in the formation of complex structures known as TORC1 and TORC2 [44].

TORC1 can detect various signals, including nutrients, growth factors, and

environmental stress. It plays a crucial role in regulating cellular processes such as

gene transcription, protein translation, ribosome synthesis, and autophagy [45]. A

characteristic that distinguishes TORC1 is its interaction with KOG1 (Raptor in

mammals) and its sensitivity to the rapamycin-FKBP12 complex [56]. The Tor kinase

is widely conserved both in terms of its structural characteristics and its role as the

target of Fkbp-rapamycin [57]. Typically, the Tor kinase exhibits similar structural

characteristics, including the HEAT-like, FAT, FRB, kinase, and FATC domains [57].

The FRB domain is the binding region of the Fkbp-rapamycin complex, and

rapamycin resistance is usually caused by the disruption of the FRB domain [45].

Homology analysis and phylogenetic studies have identified a single predicted *torA*

gene within the *A. flavus* genome. To elucidate the functional role of *torA* in *A. flavus*,

our initial approach involved an attempt to generate an *torA* deletion strain through

homologous recombination. However, these efforts were unsuccessful. We speculated

that the deletion of the *torA* gene may be lethal in *A. flavus*. This observation aligns

with the situation observed in *F. graminearum* [23]. Therefore, to investigate the

function of TorA in *A. flavus*, we constructed a mutant strain ($^{xylP}torA$) using an

xylose-inducible promoter, which allows for conditional induction with the addition

of xylose. We observed that the $^{xylP}torA$ mutant is completely unable to grow in YGT

medium without xylose, whereas partial growth is restored in YXT medium with the

addition of xylose (Fig. 3A). These results indicated that TorA is essential for the

growth of *A. flavus*. We also discovered that the Tor kinase is involved in the

vegetative growth, asexual development and aflatoxin production of *A. flavus* (Fig.

3A, 3B). In addition, the the $^{xylP}torA$ mutant shows a high sensitivity to rapamycin and

SDS (Fig. 3A). Our findings have demonstrated that the TorA kinase plays a crucial

role in the processes of morphogenesis, aflatoxin biosynthesis, and multiple stress

responses in *A. flavus*. Previous findings demonstrated that the Tor kinase could

function as a molecular switch connecting an iron cue to defend against pathogen

infection in *C. elegans* [58]. In *A. fumigatus*, the Tor kinase represents a central

regulatory node controlling genes and proteins involved in nutrient sensing, stress response, cell cycle progression, protein biosynthesis, and degradation, as well as adaptation to low iron conditions[59]. In summary, the Tor kinase acts as a central regulator in the TOR signaling pathway, playing a crucial role in the regulation of the cell cycle, protein synthesis, and cellular energy metabolism. Its influence on key cellular processes makes it a promising target for future research to control fungal growth and aflatoxin production.

The serine/threonine protein kinase Sch9, a critical downstream effector of TORC1 in *S. cerevisiae*, has been extensively studied for its roles in stress resistance, longevity, and nutrient sensing [46]. In addition, the TORC1-Sch9 pathway is also recognized as crucial mediator of chronological lifespan [46]. In filamentous fungi, an intricate network of interactions exists between the TOR and HOG pathways. In *F. graminearum* [38], FgSch9 and FgHog1 mutants displayed increased susceptibility to osmotic and oxidative stresses. The phosphorylation level of FgHog1 increased significantly in FgSch9 mutant. Additionally, the affinity capture-MS assay showed that the Tor kinase was among the proteins co-purifying with FgSch9. These results suggested that FgSch9 serves as an intermediary for the TOR and HOG signaling pathways [38]. Furthermore, FgSch9 plays a crucial role in the regulation of various biological processes such as hyphal differentiation, asexual development, virulence, and DON biosynthesis[38]. In *A. fumigatus*, the SchA mutant was sensitive to rapamycin, high concentrations of calcium and hyperosmotic stress, and SchA was involved in iron metabolism. The SchA mutant exhibited enhanced phosphorylation

of SakA in response to osmotic stress [29]. In this study, we discovered that Sch9 is involved in regulating aflatoxin biosynthesis (Fig. 4C), and responses to osmotic and rapamycin stress (Fig. 4B, 4D). The S_TKc domain deletion strain ($sch9^{\Delta S\_TKc}$), S_TK_X domain deletion strain ($sch9^{\Delta S\_TK\_X}$)**,** and $sch9^{K340A}$ mutant displayed a similar phenotype to the $\Delta sch9$ strain (Fig. 4B, 4C, 4D). These findings revealed that Sch9 plays a significant role in the biological synthesis of aflatoxin and in responding to various stresses. This role is primarily mediated through its kinase domain, kinase-C domain**,** and ATP binding site K340. We also found that the phosphorylation level of Hog1 increased significantly in the $\Delta sch9$ strain under osmotic stress (Fig. 4E). Based on the above results, we hypothesized that Sch9 was involved in the osmotic stress response by modulating the HOG pathway in *A. flavus*.

The phosphorylation state of Tap42 is crucial for its interaction with PP2A subunits and is regulated by a balance between Tor-mediated phosphorylation and Cdc55/Tpd3-mediated dephosphorylation in yeast [60]. The genome of *A. flavus* contains a homolog of *tap42* named *tapA* (AFLA_092770), which is predicted to encode a 355 amino-acid protein (sharing 31.9% identity with *S. cerevisiae* Tap42). To further determine the function of the TapA regulator in the TOR signaling pathway of *A. flavus*, we conducted targeted gene deletion experiments on *tapA*. We failed to obtain a null mutant, we hypothesised that *tapA* knockout was lethal in *A. flavus*. To investigate the function of TapA, we employed homologous recombination, to replace the native promoter with the strong *gpdA* promoter, thereby constructing an overexpression strain (*OE*::*tapA*) with the enhanced promoter. Compared with WT

strain, *OE*::*tapA* strain were more sensitive to rapamycin and SDS stress, indicating that *tapA* gene plays an important role in regulating cell wall stress (Fig. 5B, 5F). In *S. pombe*, the protein Tip41, which interacts with Tap42, plays a significant role in the cellular responses to nitrogen sources by regulating type 2A phosphatases [61]. In *S. cerevisiae*, Tap42-phosphatase complexes associate with TORC1, whereas Tip41 functions to attenuate TORC1 signaling by stimulating the PP2A phosphatase [48]. Unlike its yeast counterpart Tip41, TIPRL, a mammalian Tip41-like protein, exerts a positive influence on mTORC1 signaling by interacting with PP2Ac [62]. While some functions of Tap42 and its binding partners Tip41 may be conserved, the interaction between Tip41 and Tap42 was not observed in the filamentous fungi. The yeast two-hybrid (Y2H) assay showed that FgTip41 interacts with the phosphatase FgPpg1 rather than FgTap42, while the FgTap42 bind with phosphatases FgPp2A, implying that FgTap42, FgPpg1, and FgTip41 form a heterotrimer in *F. graminearum* [23]. In the rice blast fungus *M. oryzae*, MoTip41 does not interact with MoTap42. However, MoTip41 can bind with the phosphatase MoPpe1, thereby facilitating crosstalk between the TOR pathway and the cell wall integrity pathway [40]. In *A. flavus,* the *OE*::*tapA* and Δ*tipA* strains showed increased sensitivity to cell wall stress compared with the WT strain(Fig. 5B, 6H). Based on the conserved roles of Tap42 and Tip41 in yeast and other filamentous fungi, we hypothesize that TapA and TipA in *A. flavus* also participate in cell wall integrity pathway. In summary, our study provided important insights into the role of TapA and TipA in the TOR signaling pathway of *A. flavus*, particularly in relation to cell wall stress. However, their interaction and

implications for TOR and CWI signaling in *A. flavus* remain to be elucidated.

In *S. cerevisiae*, Tap42 can bind and regulate various PP2A phosphatases,

including Pph3, Pph21, Pph22, Sit4, and Ppg1. This interaction plays a crucial role in

the TOR signaling pathway, particularly in response to rapamycin treatment or

nutrient deprivation [48]. These phosphatases play important roles in other fungi and

are involved in a variety of cellular processes, including growth, morphogenesis, and

virulence. For example, the phosphatase Sit4 plays a crucial role in regulating cell

growth, morphogenesis, and virulence in *C. albicans* [63]. Similarly, FgSit4 and

FgPpg1 are associated with various cellular processes, including mycelial growth,

conidiation, DON biosynthesis, and virulence in *F. graminearum* [23]. These findings

indicated the conserved role of PP2A phosphatases in fungal biology. In addition to

their roles in growth and development, FgSit4 and FgPpg1 are involved in the

regulation of various signaling pathways, such as the CWI pathway and lipid droplets

biogenesis regulation pathway [23]. In this study, we found that the phosphatase SitA

and Ppg1 play critical role in vegetative growth, conidiation, sclerotia formation, and

aflatoxin biosynthesis in *A. flavus* (Fig. 7A, 7B, 7C). Additionally, the phosphatase

SitA and Ppg1 also affect pathogenicity and LD biogenesis (Fig. 8A, 10C, 10D). This

highlights the multifaceted influence of PP2A phosphatases on cellular signaling.

After the deletion of the *sitA* and *ppg1* genes, the expression level of genes related

to cell wall synthesis has decreased significantly, resulting in damage to the cell wall

and showing a higher sensitivity to cell wall integrity stress (Fig. 9C). Western

blotting results showed that the phosphorylation level of MAPK kinase Slt2 has

increased significantly in the Δ*sitA* and Δ*ppg1* strains under cell wall stress, indicating

that SitA and Ppg1 exert a negative regulatory effect on the CWI pathway (Fig. 9D).

In *M. oryzae*, MoPpe1 and MoSap1 serve as an adaptor complex that connects the

CWI and TOR signaling pathways. The activation of the TOR pathway leads to the

inhibition of the CWI pathway, highlighting a critical regulatory interplay between

these two pathways in fungal biology [39]. We speculated that SitA and Ppg1 might

exhibit similar regulation within the CWI and TOR pathways across different fungal

species. In addition, the phosphatase MoPpe1 interacts with the

phosphatase-associated protein MoTip41, facilitating the exchange of information

between the TOR and CWI signaling pathways [39]. These results suggested that the

phosphatase Ppe1 engages in interactions with various proteins to effectively

coordinate the TOR and CWI signaling pathways, thereby regulating the growth and

pathogenicity of the rice blast fungus *M. oryzae* [39]. Given the conserved functions of

PP2A phosphatases and the observed roles of SitA, Ppg1, TapA, and TipA in cell wall

stress response, we hypothesized that TapA-phosphatase complexes may collaborate

to regulate cell wall integrity processes in *A. flavus*. This hypothesis is supported by

the similar functions exhibited by these proteins.

In *F. graminearum,* the phosphatase complex FgNem1/Spo7-FgPah1 cascade

plays significant roles in fungal development, DON production, and lipid droplet (LD)

biogenesis [24]. Our study explored the conservation and functional significance of

Nem1 and Spo7 homologs in *A. flavus*, revealing their pivotal roles in vegetative

growth and secondary metabolism (Fig. 11A, 11B). Our results indicated that lipid

droplet biogenesis is facilitated by the Nem1/Spo7 and SitA/Ppg1 pathways in *A. flavus* (Fig. 10C, 10D, 10E). These observations suggested a conserved and central role for PP2A phosphatases as core regulators within the LD biogenesis signaling network, highlighting their importance in cellular lipid metabolism and energy storage. While our findings underscored the significance of phosphatases in LD biogenesis, it is also plausible that other kinases contribute to the regulation of this process. To gain a comprehensive understanding of the precise molecular mechanism, it is essential to investigate the interplay between phosphatases and other kinases.

In conclusion, we identified the complex regulatory network that provides important insight into the interplay between the TOR, HOG, and CWI signaling pathways, and how they collectively regulate the development, aflatoxin biosynthesis, and pathogenicity of *A. flavus*. By elucidating the functions of genes associated with the TOR signaling pathway, our research could establish a solid foundation for comprehending the molecular mechanism underlying the aflatoxin biosynthesis of *A. flavus*. Our results could also contribute to the development of targeted strategies to manage aflatoxin contamination and mitigate the impact of *A. flavus* infections.

Materials and methods

Strains and cultural conditions

The *A. flavus* strains in this study were shown in Table S1. All strains were cultivated on four different types of agar media: potato dextrose agar (PDA), yeast extract-sucrose agar (YES), yeast extract-glucose agar (YGT) and yeast extract-xylose

634 agar (YXT). The growth and conidiation assays were conducted at 37°C. Additionally,

635 aflatoxin production was assessed using YES and YXT liquid medium at 29°C.

Constructions of gene deletion and complemented mutants

The construction of gene deletion and complemented strains was carried out

using the SOE-PCR strategy, following previously established protocols [64]. To

generate the $^{xylP}torA$ mutant of *A. flavus* , an *torA* xylose promoter mutant cassette was

synthesized via overlapping extension PCR. This cassette was designed to incorporate

an upstream fragment of the *torA* gene, a selectable marker gene (*pyrG* from

Aspergillus fumigatus), the xyIP xylose-inducible promoter [65], and a fragment of the

torA coding sequence. Subsequently, this composite cassette was integrated into the *A.*

flavus genome via homologous recombination, specifically replacing the native *torA*

promoter with the inducible xylose promoter. To construct over-expression cassettes

for the *tapA* gene , we utilized a molecular cloning strategy that entailed the fusion of

PCR-amplified fragments from the *pyrG*-selectable marker and the *gpdA* promoter.

Subsequently, employing a homologous recombination approach, we inserted this

composite fragment upstream of the *tapA* gene, effectively replacing its native

promoter with the stronger *gpdA* promoter. The primers utilized for the amplification

of the flanking sequences of each gene can be found in Table S2. The putative gene

deletion mutants were validated through PCR assays using the appropriate primers

(Fig. S2), and their confirmation was subsequently obtained through sequencing

analysis.

Growth, conidiation, and sclerotia analysis

To evaluate mycelial growth and conidia formation, 1 μL of 10^6 spores/mL conidia

of all strains was point inoculated onto the center of the PDA medium, and then

cultured at 37°C for 5 days in the dark. The conidia were washed with 0.07%

Tween-20 and counted using a hemocytometer and microscope [43]. For sclerotia

formation analysis, each strain was inoculated and cultivated on CM and YXT agar

medium at 37°C in the dark for 7 days, and then 75% ethanol was used to wash away

mycelia and conidia on the surface of the medium [66]. These assays were repeated at

least three times.

Extraction and quantitative analysis of aflatoxin

To extract and quantify aflatoxins (AFs) production, 1 μL of 10^6 spores/mL

conidia of all strains was point inoculated onto liquid YES medium, and cultures were

then incubated at 29°C in the dark for 6 days. AFs were extracted from the media and

detected using thin layer chromatography (TLC) [66]. In addition, the standard AFB_1

was employed for visual comparison, and AFs were quantitatively analyzed using the

Image J tool. The assays were repeated at least three times.

Stress assay

To determine the roles of the gene in *A. flavus* responding to various

environmental stresses, 1 μL of 10^6 spores/mL conidia of all strains were point

inoculated onto these PDA solid plates supplemented with various agents. These

agents included 1M NaCl, 1M KCl, and 1M $CaCl_2$ for inducing osmotic stress, as

well as 100 μg/mL or 300 μg/mL SDS, 200 μg/mL Calcofluor white (CFW), and 200

μg/mL Congo red (CR) for inducing cell wall stress. After 5 days of incubation at

37°C in the dark, the relative rates of inhibition were determined [67]. The assays were

repeated at least three times.

Seeds infection assays

681   To measure the pathogenicity of *A. flavus*, all peanut and maize seeds were

682   washed in 0.05% sodium hypochlorite, 75% ethanol, and sterile water, respectively.

683   Subsequently, all peanut and maize seeds were immersed in sterile water containing

684   10^6 spores/mL of all strains at 29°C in the dark for 60 min. After the conidia adhesion,

685   the seeds were cultured at 29°C for 6 days in the dark [67].

Quantitative RT-PCR

For the real-time quantitative PCR (qPCR) assays, the *A. flavus* strains were

cultivated in the PDA medium at 29°C for 3 days in the dark. Total RNA was

extracted from the mycelia of all strains, and cDNA was synthesized with the First

Strand cDNA Synthesis Kit, and then the cDNA was used as the template for

qRT-PCR analysis with SYBR Green qPCR mix. The relative transcript level of each

gene was calculated by the $2^{-\Delta\Delta CT}$ method [68].

Western blot analysis

To extract the total protein, the mycelia of each strain were cultivated in YES

liquid medium for 48 h, then the sample was grinded under liquid nitrogen to a fine

powder and transferred to a 1.5 mL microcentrifuge tube added with RIPA lysis buffer

and 1 mM PMSF protease inhibitor. Proteins were separated by 12% SDS-PAGE and

transferred onto a polyvinylidene fluoride (PVDF) membrane. Phosphorylation levels

of Hog1 and Slt2 were detected by Phospho-p38 MAPK antibody (Cell Signaling

Technology, Boston, MA, USA) and Phospho-p44/42 MAPK (Erk 1/2) antibody (Cell signaling Technology, MA, USA), respectively. The experiment was conducted three times independently.

Statistical analysis

All data were measured with means ± standard deviation (SD) of three biological replicates. Graph Pad Prism 7 software was used for conducting statistical and significance analysis, and p-values < 0.05 represented statistical significance. Student's *t*-test was used to compare the two means for differences, whereas Tukey's multiple comparisons were used for testing multiple comparisons.

Acknowledgments

The authors also thank Zhenhong Zhuang, Jun Yuan, Yu Wang, Kunzhi Jia, Xiuna Wang, and Xinyi Nie for their help in the experiments. This work was supported by the National Natural Science Foundation of China (Grant No. 31972214).

Author contributions

The authors also Guoqi Li and Xiaohong Cao designed the experiments. Guoqi Li, Xiaohong Cao, and Qianhua Zeng performed the experiments. Guoqi Li, Xiaohong Cao, and Shihua Wang analyzed the data. Guoqi Li, Elisabeth Tumukunde, and Shihua Wang wrote the manuscript. The manuscript was discussed and revised by all the authors.

References

[1] Klich MA. *Aspergillus flavus*: the major producer of aflatoxin. Mol Plant Pathol.

721   2007; 8(6): 713-22.

[2] Hedayati MT, Pasqualotto AC, Warn PA, et al. *Aspergillus flavus*: human

pathogen, allergen and mycotoxin producer. Microbiology (Reading). 2007;

153(Pt 6): 1677-1692.

[3] Khan R, Ghazali FM, Mahyudin NA, et al. Biocontrol of aflatoxins using

non-aflatoxigenic *Aspergillus flavus*: a literature review. J Fungi (Basel). 2021;

7(5): 381.

[4] Yabe K, Nakajima H. Enzyme reactions and genes in aflatoxin biosynthesis. Appl

Microbiol Biotechnol. 2004; 64(6): 745-55.

[5] Wullschleger S, Loewith R, Hall MN. TOR signaling in growth and metabolism.

Cell. 2006; 124(3): 471-84.

[6] De Virgilio C, Loewith R. The TOR signalling network from yeast to man. Int J

Biochem Cell Biol. 2006; 38(9): 1476-1481.

[7] Loewith R, Jacinto E, Wullschleger S, et al. Two TOR complexes, only one of

which is rapamycin sensitive, have distinct roles in cell growth control. Mol Cell.

2002; 10(3): 457-68.

[8] Beauchamp EM, Platanias LC. The evolution of the TOR pathway and its role in

cancer. Oncogene. 2013 Aug 22; 32(34): 3923-32.

[9] Fu L, Liu Y, Qin G, et al. The TOR-EIN2 axis mediates nuclear signalling to

modulate plant growth. Nature. 2021; 591(7849): 288-292.

[10] Teichert S, Wottawa M, Schönig B, et al. Role of the *Fusarium fujikuroi* TOR

kinase in nitrogen regulation and secondary metabolism. Eukaryot Cell. 2006;

743   5(10): 1807-1819.

[11] Crespo JL, Hall MN. Elucidating TOR signaling and rapamycin action: lessons from *Saccharomyces cerevisiae*. Microbiol Mol Biol Rev. 2002; 66(4): 579-591.

[12] Rohde JR, Bastidas R, Puria R, et al. Nutritional control via Tor signaling in *Saccharomyces cerevisiae*. Curr Opin Microbiol. 2008; 11(2): 153-60.

[13] Ishiguro J, Shibahara K, Ueda Y, et al. Fission yeast TOR signaling is essential for the down-regulation of a hyperactivated stress-response MAP kinase under salt stress. Mol Genet Genomics. 2013; 288(1-2): 63-75.

[14] Otsubo Y, Yamamato M. TOR signaling in fission yeast. Crit Rev Biochem Mol Biol. 2008; 43(4): 277-83.

[15] Petersen J. TOR signalling regulates mitotic commitment through stress-activated MAPK and Polo kinase in response to nutrient stress. Biochem Soc Trans. 2009; 37(Pt1): 273-7.

[16] Bastidas RJ, Heitman J, Cardenas ME. The protein kinase Tor1 regulates adhesin gene expression in *Candida albicans*. PLoS Pathog. 2009; 5(2): e1000294.

[17] Liu NN, Flanagan PR, Zeng J, et al. Phosphate is the third nutrient monitored by TOR in *Candida albicans* and provides a target for fungal-specific indirect TOR inhibition. Proc Natl Acad Sci U S A. 2017; 114(24): 6346-6351.

[18] Hu K, Guo S, Yan G, et al. Ubiquitin regulates TORC1 in yeast *Saccharomyces cerevisiae*. Mol Microbiol. 2016; 100(2): 303-314.

[19] Wang H, Wang X, Jiang Y. Interaction with Tap42 is required for the essential function of Sit4 and type 2A phosphatases. Mol Biol Cell. 2003; 14(11): 4342-51.

[20] Yan G, Shen X, Jiang Y. Rapamycin activates Tap42-associated phosphatases by

abrogating their association with Tor complex 1. EMBO J. 2006; 25(15):

3546-55.

[21] Tate JJ, Tolley EA, Cooper TG. Sit4 and PP2A dephosphorylate nitrogen

catabolite repression-sensitive Gln3 when TorC1 is up- as well as downregulated.

Genetics. 2019; 212(4): 1205-1225.

[22] Urban J, Soulard A, Huber A, et al. Sch9 is a major target of TORC1 in

Saccharomyces cerevisiae. Mol Cell. 2007; 26(5): 663-74.

[23] Yu F, Gu Q, Yun Y, et al. The TOR signaling pathway regulates vegetative

development and virulence in *Fusarium graminearum*. New Phytol. 2014; 203(1):

219-32.

[24] Liu N, Yun Y, Yin Y, et al. Lipid droplet biogenesis regulated by the

FgNem1/Spo7-FgPah1 phosphatase cascade plays critical roles in fungal

development and virulence in *Fusarium graminearum*. New Phytol. 2019; 223(1):

412-429.

[25] Sun G, Elowsky C, Li G, et al. TOR-autophagy branch signaling via Imp1

dictates plant-microbe biotrophic interface longevity. PLoS Genet. 2018; 14(11):

e1007814.

[26] Nguyen DV, Roret T, Fernandez-Gonzalez A, et al. Target of rapamycin pathway

in the white-rot fungus *Phanerochaete chrysosporium*. PLoS One. 2020; 15(2):

e0224776.

[27] Dementhon K, Paoletti M, Pinan-Lucarré B, et al. Rapamycin mimics the

incompatibility reaction in the fungus *Podospora anserina*. Eukaryot Cell. 2003;

2(2): 238-46.

[28] Fitzgibbon GJ, Morozov IY, Jones MG, et al. Genetic analysis of the TOR

pathway in *Aspergillus nidulans*. Eukaryot Cell. 2005; 4(9): 1595-8.

[29] Alves de Castro P, Dos Reis TF, et al. The *Aspergillus fumigatus* SchA[SCH9] kinase

modulates SakA[HOG1] MAP kinase activity and it is essential for virulence. Mol

Microbiol. 2016; 102(4): 642-671.

[30] Baldin C, Valiante V, Krüger T, et al. Comparative proteomics of a tor inducible

Aspergillus fumigatus mutant reveals involvement of the Tor kinase in iron

regulation. Proteomics. 2015; 15(13): 2230-43.

[31] Teichert S, Wottawa M, Schönig B, et al. Role of the *Fusarium fujikuroi* TOR

kinase in nitrogen regulation and secondary metabolism. Eukaryot Cell. 2006;

5(10): 1807-19.

[32] Meléndez HG, Billon-Grand G, Fèvre M, et al. Role of the *Botrytis cinerea*

FKBP12 ortholog in pathogenic development and in sulfur regulation. Fungal

Genet Biol. 2009; 46(4): 308-20.

[33] López-Berges MS, Rispail N, Prados-Rosales RC, et al. A nitrogen response

pathway regulates virulence functions in *Fusarium oxysporum* via the protein

kinase TOR and the bZIP protein MeaB. Plant Cell. 2010; 22(7): 2459-75.

[34] Bastidas RJ, Shertz CA, Lee SC, et al. Rapamycin exerts antifungal activity *in*

vitro and *in vivo* against *Mucor circinelloides* via FKBP12-dependent inhibition

of Tor. Eukaryot Cell. 2012; 11(3): 270-81.

[35] Madrid, M, Vázquez-Marín, B, Franco, A, et al. Multiple crosstalk between TOR

and the cell integrity MAPK signaling pathway in fission yeast. Sci Rep. 2016

Nov 23; 6: 37515.

[36] Torres J, Di Como CJ, Herrero E, et al. Regulation of the cell integrity pathway

by rapamycin-sensitive TOR function in budding yeast. J Biol Chem. 2002 Nov 8;

277(45): 43495-504.

[37] Chen DD, Shi L, Yue SN, et al. The Slt2-MAPK pathway is involved in the

mechanism by which target of rapamycin regulates cell wall components in

Ganoderma lucidum. Fungal Genet Biol. 2019; 123: 70-77.

[38] Gu Q, Zhang C, Yu F, et al. Protein kinase FgSch9 serves as a mediator of the

target of rapamycin and high osmolarity glycerol pathways and regulates multiple

stress responses and secondary metabolism in *Fusarium graminearum*. Environ

Microbiol. 2015; 17(8): 2661-76.

[39] Qian B, Liu X, Jia J, et al. MoPpe1 partners with MoSap1 to mediate TOR and

cell wall integrity signalling in growth and pathogenicity of the rice blast fungus

Magnaporthe oryzae. Environ Microbiol. 2018; 20(11): 3964-3979.

[40] Qian B, Liu X, Ye Z, et al. Phosphatase-associated protein MoTip41 interacts

with the phosphatase MoPpe1 to mediate crosstalk between TOR and cell wall

integrity signalling during infection by the rice blast fungus *Magnaporthe oryzae*.

Environ Microbiol. 2021; 23(2): 791-809.

[41] Tong M, Jiang Y. FK506-binding proteins and their diverse functions. Curr Mol

Pharmacol. 2015; 9(1): 48-65.

[42] Heitman J, Movva NR, Hall MN. Targets for cell cycle arrest by the immunosuppressant rapamycin in yeast. Science. 1991; 253(5022): 905-9.

[43] Ren S, Yang M, Yue Y, et al. Lysine succinylation contributes to aflatoxin production and pathogenicity in *Aspergillus flavus*. Mol Cell Proteomics. 2018; 17(3): 457-471.

[44] Weisman R. Target of rapamycin (TOR) regulates growth in response to nutritional signals. Microbiol Spectr. 2016; 4(5).

[45] Loewith R, Hall MN. Target of rapamycin (TOR) in nutrient signaling and growth control. Genetics. 2011; 189(4): 1177-201.

[46] Pascual-Ahuir A, Proft M. The Sch9 kinase is a chromatin-associated transcriptional activator of osmostress-responsive genes. EMBO J. 2007; 26(13): 3098-108.

[47] Düvel K, Broach JR. The role of phosphatases in TOR signaling in yeast. Curr Top Microbiol Immunol. 2004; 279: 19-38.

[48] Jacinto E, Guo B, Arndt KT, et al. TIP41 interacts with TAP42 and negatively regulates the TOR signaling pathway. Mol Cell. 2001; 8(5): 1017-26.

[49] Sánchez-Álvarez M, Del Pozo MÁ, Bosch M, et al. Insights into the biogenesis and emerging functions of lipid droplets from unbiased molecular profiling approaches. Front Cell Dev Biol. 2022; 10: 901321.

[50] Gizachew D, Chang CH, Szonyi B, et al. Aflatoxin B1 (AFB$_1$) production by *Aspergillus flavus* and *Aspergillus parasiticus* on ground Nyjer seeds: the effect of water activity and temperature. Int J Food Microbiol. 2019; 296: 8-13.

[51] Cruz MC, Goldstein AL, Blankenship J, et al. Rapamycin and less

immunosuppressive analogs are toxic to *Candida albicans* and *Cryptococcus*

neoformans via FKBP12-dependent inhibition of TOR. Antimicrob Agents

Chemother. 2001; 45(11): 3162-70.

[52] Galat A. Functional diversity and pharmacological profiles of the FKBPs and

their complexes with small natural ligands. Cell Mol Life Sci. 2013; 70(18):

3243-75.

[53] Kasahara K. Physiological function of FKBP12, a primary target of

rapamycin/FK506: a newly identified role in transcription of ribosomal protein

genes in yeast. Curr Genet. 2021 Jun; 67(3): 383-388.

[54] Li Y, Ren H, Zhao Y, et al. Characterization of three FK506-binding proteins in

the entomopathogenic fungus *Beauveria bassiana*. J Invertebr Pathol. 2020; 171:

107334.

[55] Xiong Y, Sheen J. The role of target of rapamycin signaling networks in plant

growth and metabolism. Plant Physiol. 2014; 164(2): 499-512.

[56] Adami A, García-Alvarez B, Arias-Palomo E, et al. Structure of TOR and its

complex with KOG1. Mol Cell. 2007; 27(3): 509-16.

[57] Tatebe H, Shiozaki K. Evolutionary conservation of the components in the TOR

signaling pathways. Biomolecules. 2017 ;7(4):77.

[58] Ma YC, Dai LL, Qiu BB, et al. TOR functions as a molecular switch connecting

an iron cue with host innate defense against bacterial infection. PLoS Genet. 2021;

17(3): e1009383.

[59] Baldin C, Valiante V, Krüger T, et al. Comparative proteomics of a tor inducible *Aspergillus fumigatus* mutant reveals involvement of the Tor kinase in iron regulation. Proteomics. 2015; 15(13): 2230-43.

[60] Jiang Y, Broach JR. Tor proteins and protein phosphatase 2A reciprocally regulate Tap42 in controlling cell growth in yeast. EMBO J. 1999;18(10):2782-92.

[61] Fenyvuesvolgyi C, Elder RT, Benko Z, et al. Fission yeast homologue of Tip41-like proteins regulates type 2A phosphatases and responses to nitrogen sources. Biochim Biophys Acta. 2005; 1746(2): 155-62.

[62] Nakashima A, Tanimura-Ito K, Oshiro N, et al. A positive role of mammalian Tip41-like protein, TIPRL, in the amino-acid dependent mTORC1-signaling pathway through interaction with PP2A. FEBS Lett. 2013; 587(18): 2924-9.

[63] Han Q, Pan C, Wang Y, et al. PP2A-like protein phosphatase (Sit4) regulatory subunits, Sap155 and Sap190, regulate *Candida albicans*' cell growth, morphogenesis, and virulence. Front Microbiol. 2019; 10: 2943.

[64] Tan L, Chen H, Yu S, et al. A SOE-PCR method of introducing multiple mutations into Mycoplasma gallisepticum neuraminidase. J Microbiol Methods. 2013; 94(2): 117-120.

[65] Zadra I, Abt B, Parson W, Haas H. *xylP* promoter-based expression system and its use for antisense downregulation of the *Penicillium chrysogenum* nitrogen regulator NRE. Appl Environ Microbiol. 2000;66(11):4810-6.

[66] Nie X, Yu S, Qiu M, et al. *Aspergillus flavus* SUMO contributes to fungal virulence and toxin attributes. J Agric Food Chem. 2016 Sep 7; 64(35): 6772-82.

[67] Tan C, Deng J, Zhang F, et al. CWI pathway participated in vegetative growth

and pathogenicity through a downstream effector AflRlm1 in *Aspergillus flavus*.

iScience. 2021;24(10):103159.

[68] Livak KJ, Schmittgen TD. Analysis of relative gene expression data using

real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. Methods. 2001; 25(4):

402–408.

[69] Chang PK, Scharfenstein LL, Wei Q, Bhatnagar D. Development and refinement

of a high-efficiency gene-targeting system for *Aspergillus flavus*. J Microbiol

Methods. 2010;81(3):240-6.

Fig. 1 Impacts of rapamycin on the growth, sporulation, sclerotia formation,and aflatoxin production of *Aspergillus flavus*.

(A) Colony morphology of the WT strain cultured on PDA medium amended with different concentrations of rapamycin at 37°C for 5 days.

(B) Colony phenotype of the WT strain grown on CM medium amended with different concentrations of rapamycin at 37°C for 7 days.

(C) TLC assay of AFB_1 production by the WT strain cultured in YES liquid media amended with different concentrations of rapamycin at 29°C for 6 days.

(D) Statistical analysis of the colony diameter by the WT strain treated with different concentrations of rapamycin described in (A).

(E) Conidial quantification of the WT strain treated with different concentrations of rapamycin as mentioned in (A).

(F) Quantitative analysis of sclerotium formation by the WT strain treated with different concentrations of rapamycin described in (B).

(G) Relative quantification of AFB_1 production by the WT strain treated with different concentrations of rapamycin as mentioned in (C)

(H) Relative expression levels of conidia-related genes in WT strain treated with 100 ng/mL rapamycin after 3, 6, 9 h.

(I) Relative expression levels of sclerotia-related genes in WT strain treated with 100 ng/mL rapamycin after 3, 6, 9 h.

(J) Relative expression levels of aflatoxin biosynthesis regulatory and structural genes in WT strain treated with 100 ng/mL rapamycin after 3, 6, 9 h.

Fig. 2 Disruption of *fkbp3* significantly increases the resistance of *A. flavus* to rapamycin and FK506.

(A) Phenotype of the WT and all mutant strains (Δ*fkbp1*, Δ*fkbp2*, Δ*fkbp3*, and Δ*fkbp4*) grown on PDA amended with rapamycin and FK506 at 37°C for 5 days.

(B) TLC analysis of AFB_1 production by the WT and all mutant strains (Δ*fkbp1*, Δ*fkbp2*, Δ*fkbp3*, and Δ*fkbp4*) cultured in YES liquid medium at 29°C for 6 days.

(C) The growth inhibition rate of the WT and all mutant strains (Δ*fkbp1*, Δ*fkbp2*, Δ*fkbp3*, and Δ*fkbp4*) under rapamycin and FK506 stress.

(D) AFB_1 quantitative analysis of the WT and all mutant strains (Δ*fkbp1*, Δ*fkbp2*, Δ*fkbp3*, and Δ*fkbp4*) as described in (B).

(E) Phenotype of the WT and all mutant strains (Δ*fkbp3*, K5A, K19A, K40A, K42A, K55A, and K65A) grown on PDA amended with 100 ng/mL rapamycin at 37°C for 5 days.

(F) The growth inhibition rate of the WT and all mutant strains (Δ*fkbp3*, K5A, K19A, K40A, K42A, K55A, and K65A) under rapamycin stress.

(G) TLC assay of AFB_1 production by the WT and all mutant strains (Δ*fkbp3*, K5A, K19A, K40A, K42A, K55A, and K65A) cultured in YES liquid medium at 29°C for 6 days.

(H) Relative quantification of AFB_1 production in the WT and all mutant strains (Δ*fkbp3*, K5A, K19A, K40A, K42A, K55A, and K65A) as mentioned in (G).

Fig. 3 The TorA kinase plays a critical role in *A. flavus*

(A) Phenotype of the WT and the $^{xylP}torA$ strains in YGT and YXT medium amended with 100 ng/mL rapamycin and 300 mg/mL SDS at 37°C for 5 days.

(B) TLC assay of AFB_1 production from the WT and the $^{xylP}torA$ strains cultured in s on YXT medium containing 1 g/L $MgSO_4·7H_2O$ at 29°C for 6 days.

(C) Statistical analysis of the colony diameter by the WT and the $^{xylP}torA$ strains in YXT medium.

(D) Statistical analysis of the colony diameter by the WT and the $^{xylP}torA$ strains in YGT medium.

(E) Conidial quantification of theWT and the $^{xylP}torA$ strains in YXT medium.

(F) Quantitative analysis of AFB_1 as shown in (B).

(G) The growth inhibition rate of the WT and the $^{xylP}torA$ strains under rapamycin.

(H) The growth inhibition rate of the WT and the $^{xylP}torA$ strains under SDS stress.

Fig. 4 The Sch9 kinase participates in aflatoxin biosynthesis and the HOG pathway.

(A) The structure diagram of the Sch9 kinase.

(B) Phenotype of the WT, $\Delta sch9$, $sch9^{\Delta C2}$, $sch9^{\Delta S\_TKc}$, $sch9^{\Delta S\_TK\_X}$, and $sch9^{K340A}$ strains grown on PDA medium amended with 100 ng/mL rapamycin at 37°C for 5 days.

(C) TLC assay of AFB_1 production from the WT, $\Delta sch9$, $sch9^{\Delta C2}$, $sch9^{\Delta S\_TKc}$, $sch9^{\Delta S\_TK\_X}$, and $sch9^{K340A}$ strains cultured in YES liquid medium at 29°C for 6 days.

(D) The phenotypes of the WT, $\Delta sch9$, $sch9^{\Delta C2}$, $sch9^{\Delta S\_TKc}$, $sch9^{\Delta S\_TK\_X}$, and $sch9^{K340A}$ strains on YGT media amended with 1 M NaCl and 1 M KCl for 3 days.

(E) The phosphorylation levels of Hog1 in the WT and $\Delta sch9$ strains were determined with or without osmotic stress.

(F) The growth inhibition rate of the WT, $\Delta sch9$, $sch9^{\Delta C2}$, $sch9^{\Delta S\_TKc}$, $sch9^{\Delta S\_TK\_X}$, and $sch9^{K340A}$ strains under rapamycin stress.

(G) Quantitative analysis of AFB_1 as shown in (C).

(H) The growth inhibition rate of the WT, $\Delta sch9$, $sch9^{\Delta C2}$, $sch9^{\Delta S\_TKc}$, $sch9^{\Delta S\_TK\_X}$, and $sch9^{K340A}$ strains under osmotic stress.

Fig. 5 TapA regulates cell wall stress in *A. flavus*.

(A) Phenotype of the WT and *OE*::*tapA* strains grown on PDA amended with 100 ng/mL rapamycin at 37°C for 5 days.

(B) Colony morphology of the WT and *OE*::*tapA* strains grown on PDA media supplemented with 200 µg/mL CR, 200µg/mL CFW, or 300 µg/mL SDS at 37°C for 5 days.

(C) TLC assay of AFB_1 production by the WT and *OE*::*tapA* strains cultured in YES liquid medium at 29°C for 6 days.

(D) The growth inhibition rate of the WT and *OE*::*tapA* strains under rapamycin.

(E) Quantitative analysis of AFB_1 as shown in (C).

(F) The growth inhibition rate of the WT and *OE*::*tapA* strains under cell wall and cell membrane stress.

Fig. 6 TipA regulates sclerotia development and cell wall stress in *A. flavus*.

(A) Phenotype of the WT and Δ*tipA* strains grown on PDA amended with 100 ng/mL rapamycin at 37°C for 5 days.

(B) Phenotypic characterization of the WT and Δ*tipA* strains grown on CM medium at 37°C for 7 days.

(C) Morphology of conidiophores in the WT and Δ*tipA* strains observed by microscope.

(D) TLC assay of AFB_1 production by the WT and Δ*tipA* strains cultured in YES liquid medium at 29°C for 6 days.

(E) The growth inhibition rate of the WT and Δ*tipA* strains under rapamycin stress.

(F) Amount of sclerotia produced by the WT and Δ*tipA* strains.

(G) AFB_1 quantitative analysis of the WT and Δ*tipA* strains described in (D).

(H) Colony morphology of the WT and Δ*tipA* strains grown on PDA media supplemented with 200 µg/mL CR, 200µg/mL CFW, and 100 µg/mLSDS at 37°C for 5 days.

(I) The growth inhibition rate of the WT and Δ*tipA* strains under cell wall and cell membrane stress.

(J) Relative expression levels of sclerotia formation genes in the WT and Δ*tipA* strains.

Fig. 7 The impact of the phosphatases SitA and Ppg1 on the growth, conidiation, and aflatoxin biosynthesis in *A. flavus*.

(A) Colony morphology of the WT, single knockout, and double knockout strains cultured on PDA medium amended with 100 ng/mL rapamycin at 37°C for 5 days.

(B) Colony morphology of the WT, single knockout, and double knockout strains cultured on CM medium at 37°C for 7 days.

(C) TLC analysis of AFB_1 production from the WT, single knockout, and double knockout strains cultured in YES liquid medium at 29°C for 6 days.

(D) Growth diameter of the WT, single knockout, and double knockout strains on PDA media.

(E) The growth inhibition rate of the WT, single knockout, and double knockout strains under rapamycin stress.

(F) AFB_1 quantitative analysis of the WT, single knockout, and double knockout strains.

(G) Relative expression levels of conidia synthesis genes in the WT, single knockout, and double knockout strains.

(H) Relative expression levels of sclerotia synthesis genes in the WT, single knockout, and double knockout strains.

(I) Relative expression levels of aflatoxin biosynthesis genes in the WT, single knockout, and double knockout strains.

Fig. 8 Pathogenicity analysis of the phosphatase SitA and Ppg1 in *A. flavus*.

(A) Phenotypes of peanut and maize infected by the WT, single knockout, and double knockout strains.

(B) TLC was used to detect the AFB_1 from peanut and maize infected by the WT, single knockout, and double knockout strains.

(C) Quantitative analysis of conidia production from peanut and maize infected by the WT, single knockout, and double knockout strains.

(D) Quantitative analysis of AFB_1 from peanut and maize infected by the WT, single knockout, and double knockout strains.

A

	WT	ΔsitA	ΔsitA-C	Δppg1	Δppg1-C	ΔsitAΔppg1
Control						
200µg/ml CFW						
200µg/ml CR						
100µg/ml SDS						

B

C

D

| P-Slt2/Slt2 | 0.13 | 0.44 | 0.5 | 0.72 | 0.24 | 0.61 | 0.86 | 1.05 |

Fig. 9 Sensitivity of phosphatase SitA and Ppg1 to cell wall damaging agents in *A. flavus*.

(A) Morphology of the WT, single knockout, and double knockout strains grown on YES media supplemented with 200 µg/mL CR, 200 µg/mL CFW, or 100 µg/mL SDS at 37°C for 5 days.

(B) The growth inhibition rate of the WT, single knockout, and double knockout strains under cell wall and cell membrane stress.

(C) Relative expression levels of cell wall synthesis genes (*chsA*, *chsD* and *chsE*) in the WT, single knockout, and double knockout strains.

(D) The phosphorylation level of Slt2 in the WT, single knockout, and double knockout strains was detected with or without cell wall stress.

Fig. 10 Phosphatase SitA and Ppg1 are involved in regulating lipid droplet biogenesis.

(A) The intracellular lipid droplets were stained by boron dipyrromethene difluoride (BODIPY) in hyphae of the WT strain and observed under fluorescence microscopy. The phenotype of lipid droplets accumulation in the mycelia of the WT strain treated with or without 100 ng/mL rapamycin for 6 h (Bar, 10 μm).

(B) Phenotype of lipid droplets accumulation in the mycelia of the Δ*sch9* strain treated with or without 100 ng/mL rapamycin for 6 h (Bar, 10 μm).

(C) Phenotype of lipid droplets accumulation in the mycelia of the Δ*sitA* strain treated with or without 100 ng/mL rapamycin for 6 h (Bar, 10 μm).

(D) Phenotype of lipid droplets accumulation in the mycelia of the Δ*ppg1* strain treated with or without 100 ng/mL rapamycin for 6 h (Bar, 10 μm).

Fig. 11 Phosphatase complex Nem1/Spo7 play significant role in growth, conidiation, aflatoxin, and lipid droplet biogenesis.

(A) Phenotype of the WT, Δ*nem1*, and Δ*spo7* strains grown on PDA amended with 100 ng/mL rapamycin at 37°C for 5 days.

(B) TLC assay of AFB_1 production from the WT, Δ*nem1*, and Δ*spo7* strains cultured in YES liquid medium at 29°C for 6 days.

(C) Growth diameter of the WT, Δ*nem1*, and Δ*spo7* strains on PDA media.

(D) Statistical analysis of the sporulation by the WT, Δ*nem1*, and Δ*spo7* strains.

(E) Phenotype of lipid droplets accumulation in the mycelia of the WT, Δ*nem1*, and Δ*spo7* strains treated with or without 100 ng/mL rapamycin for 6 h (Bar, 10 μm).

Fig. 12 The proposed model of the target of rapamycin (TOR) pathway in *A. flavus*.

Based on the above results, we propose a hypothesis regarding the TOR signaling pathway model in *A. flavus*. Rapamycin forms a complex with Fkbp3, and this particular complex can bind to the TorA kinase, thereby impeding its regular functionality. Sch9, functioning as a downstream element of the TorA kinase, regulates aflatoxin biosynthesis and HOG signaling pathway. As another important target of the TorA kinase, TapA-phosphatase complex is involved in the regulation of growth, sporulation, sclerotia, aflatoxin production, CWI signaling pathway, lipid droplet synthesis, and other processes. In conclusion, the TOR signaling pathway plays a crucial role in various aspects of *A. flavus*, including vegetative development, stress response, aflatoxin biosynthesis, and pathogenicity.