Title: Efficient estimation for large-scale linkage disequilibrium patterns of the human genome

Authors: Xin Huang[1,2,5](huangxin0221@zju.edu.cn), Tian-Neng Zhu[1,5](zhutianneng@zju.edu.cn),

Ying-Chao Liu[1](22016031@zju.edu.cn), Jian-Nan Zhang[3](zjn364739@alibaba-inc.com), Guo-Bo

Chen[2,4,*](chenguobo@gmail.com; orcid 0000-0001-5475-8237)

Affiliations:

[1]Institute of Bioinformatics, Zhejiang University, Hangzhou, Zhejiang Province, China;

[2]Center for General Practice Medicine, Department of General Practice Medicine; Center for Reproductive

Medicine, Department of Genetic and Genomic Medicine, and Clinical Research Institute, Zhejiang

Provincial People's Hospital, People's Hospital of Hangzhou Medical College, Hangzhou, Zhejiang,

China;

[3]Alibaba Group, Hangzhou, Zhejiang, China;

[4]Key Laboratory of Endocrine Gland Diseases of Zhejiang Province, Hangzhou, Zhejiang, China.

[*]**Correspondence (GBC)**: chenguobo@gmail.com

[5]Equal contribution

Abstract

In this study, we proposed an efficient algorithm (X-LD) for estimating LD patterns for a genomic grid, which can be of inter-chromosomal scale or of a pair of small segments. Compared with conventional methods, the proposed method was significantly faster, and consequently we were permitted to explore in depth unknown or reveal long-anticipated LD features of the human genome. Having applied the algorithm as demonstrated in 1000 Genome Project (1KG), we found: **I**) The extended LD, driven by population structure, was universally existed, and the strength of inter-chromosomal LD was about 10% their respective intra-chromosomal LD in relatively homogeneous cohorts, such as FIN and to nearly 56% in admixed cohort, such as ASW. **II**) After splitting each chromosome into upmost more than a half million grids, we elucidated the LD of HLA region was nearly 42 folders higher than chromosome 6 in CEU and 11.58 in ASW; on chromosome 11, we observed that the LD of its centromere was nearly 94.05 folders higher than chromosome 11 in YRI and 42.73 in ASW. **III**) We uncovered the long-anticipated inversely proportional linear relationship between the length of a chromosome and the strength of chromosomal LD, and their Pearson's correlation was on average over 0.80 for 26 1KG cohorts. However, this linear norm was so far perturbed by chromosome 11 given its more completely sequenced centromere region. Uniquely chromosome 8 of ASW was found most deviated from the linear norm than any other autosomes. The proposed algorithm has been realized in C++ (called X-LD) and available at https://github.com/gc5k/gear2, and can be applied to explore LD features in any sequenced populations.

Introduction

Linkage disequilibrium (LD) is the association for a pair of loci and the metric of LD serves as the basis for developing genetic applications in agriculture, evolutionary biology, and biomedical researches (Weir, 2008; Hill and Robertson, 1966). The structure of LD of the human genome is shaped by many factors, mutation, recombination, population demography, epistatic fitness, and completeness of genomic data itself (Myers *et al.*, 2005; Nei and Li, 1973; Ardlie *et al.*, 2002). Due to its overwhelming cost, LD structure investigation is often compromised to a small genomic region (Chang *et al.*, 2015; Theodoris *et al.*, 2021), and their typical LD structure is as illustrated for a small segment (Barrett *et al.*, 2005). Now, given the availability of large-scale genomic data, such as millions of single nucleotide polymorphisms (SNPs), the large-scale LD patterns of the human genome play crucial roles in determining genomics studies, and many theories and useful algorithms upon large-scale LD structure, from genome-wide association studies, polygenic risk prediction for complex diseases, and choice for reference panels for genotype imputation (Vilhjálmsson *et al.*, 2015; Yang and Zhou, 2020; Bulik-Sullivan *et al.*, 2015; Yang *et al.*, 2011; Das *et al.*, 2016).

However, there are impediments, largely due to intensified computational cost, in both investigating large-scale LD and providing high-resolution illustration for their details. If we consider a genomic grid that is consisted of m^2 SNP pairs, given a sample of n individuals and m SNPs ($n \ll m$) – typically as observed in 1000 Genomes Project (1KG) (Lowy-Gallego *et al.*, 2019), its benchmark computational time cost for estimating all pairwise LD is $\mathcal{O}(nm^2)$, a burden that quickly drains computational resources given the volume of the genomic data. In practice, it is of interest to know the mean LD of the m_i^2 SNP pairs for a genomic grid, which covers $m_i \times m_j$ SNP pairs. Upon how a genomic grid is defined, a genomic grid consequently can be consisted of : **i)** the whole genome-wide m^2 SNP pairs, and we denote their mean LD as ℓ_g; **ii)** the intra-chromosomal mean LD for the i^{th} chromosome of m_i^2 SNP pairs, and denote as ℓ_i; **iii)** the inter-chromosomal mean LD i^{th} and j^{th} chromosomal $m_i m_j$ SNP pairs, and denoted as ℓ_{ij}.

In this study we propose an efficient algorithm that can estimate ℓ_g, ℓ_i, and ℓ_{ij}, the computational time of which can be reduced from $\mathcal{O}(nm_i^2)$ to $\mathcal{O}(n^2 m_i)$ for ℓ_i and $\mathcal{O}(nm_i m_j)$ to $\mathcal{O}(n^2 m_i + n^2 m_j)$ for ℓ_{ij}. The rationale of the proposed method relies on the connection between the genetic relationship matrix

(GRM) and LD (Chen, 2014; Goddard, 2009), and in this study a more general transformation from GRM

to LD can be established via Isserlis's theorem (Isserlis, 1918; Zhou, 2017). The statistical properties, such

as sampling variance, of the estimated LD have been derived too.

The proposed method can be analogously considered a more powerful realization for Haploview (Barrett *et*

al., 2005), but additional utility can be derived to bring out unprecedented survey of LD patterns of the

human genome. As demonstrated in 1KG, we consequently investigate how biological factors such as

population structure, admixture, or variable local recombination rates can shape large-scale LD patterns of

the human genomes.

1) The proposed method provides statistically unbiased estimates for large-scale LD patterns and

shows computational merits compared with the conventional methods **(Figure 2)**.

2) We estimated ℓ_g, and 22 autosomal ℓ_i and 231 inter-autosomal ℓ_{ij} for the 1KG cohorts. There

were ubiquitously existence of extended LD, which was associated with population structure or

admixture **(Figure 3)**.

3) We provided high-resolution illustration that decomposed a chromosome into upmost nearly a

million grids, each of which was consisted of 250×250 SNP pairs, the highest resolution that

has been realized so far at autosomal level **(Figure 4)**; tremendous variable recombination rates

led to regional strong LD as highlighted for the HLA region of chromosomes 6 and the centromere

region of chromosome 11.

4) Furthermore, a consequently linear regression constructed could quantify LD decay score

genome-widely, and in contrast LD decay was previously surrogated in a computational expensive

method. There was strong ethnicity effect that was associated with extended LD **(Figure 5)**.

5) We demonstrate that the strength of autosomal ℓ_i was inversely proportional to the SNP number,

an anticipated relationship that is consistent to genome-wide spread of recombination hotspots.

However, the chromosome 8 of ASW showed substantial deviation from the fitted linear

relationship **(Figure 6)**.

The proposed algorithm has been realized in C++ and is available at: https://github.com/gc5k/gear2. As

tested the software could handle sample size as large as more than 10,000 individuals.

Methods and Materials

The overall rationale for large-scale LD analysis

We assume LD for a pair of biallelic loci is measured by the squared Pearson's correlation, $\rho^2_{l_1 l_2} =$

$\frac{D^2_{l_1 l_2}}{p_{l_1} q_{l_1} p_{l_2} q_{l_2}}$, in which $D_{l_1 l_2}$ the LD of loci l_1 and l_2, p and q the reference and the alternative allele

frequencies. If we consider the averaged LD for a genomic grid over m_i^2 SNP pairs, the conventional

estimator is $\hat{\ell}_i = \frac{1}{m_i^2} \sum_{l_1, l_2}^{m_i} \rho^2_{l_1 l_2}$, and, if we consider the averaged LD for m_i and m_j SNP pairs between

two genomic segments, then $\hat{\ell}_{ij} = \frac{1}{m_i m_j} \sum_{l_1, l_2}^{m_i, m_j} \rho^2_{l_1 l_2}$. Now let us consider the 22 human autosomes

(**Figure 1A**). We naturally partition the genome into $\mathcal{C} = 22$ blocks, and its genomic LD, denoted as

ℓ_g, can be expressed as

$$\ell_g = \frac{1}{m^2} \sum_{l_1, l_2}^{m} \rho^2_{l_1 l_2} = \sum_i^{\mathcal{C}} \left(\frac{1}{m_i^2} \sum_{l_1, l_2}^{m_i} \rho^2_{l_1 l_2} \right) + \sum_{i \neq j}^{\mathcal{C}} \left(\frac{1}{m_i m_j} \sum_{l_1}^{m_i} \sum_{l_2}^{m_j} \rho^2_{l_1 l_2} \right) = \sum_i^{\mathcal{C}} \ell_i + \sum_{i \neq j}^{\mathcal{C}} \ell_{ij} \qquad \textbf{(Eq 1)}$$

103 So we can decompose ℓ_g into $\mathcal{C}$ ℓ_i and $\frac{\mathcal{C}(\mathcal{C}-1)}{2}$ unique ℓ_{ij}. Obviously, **Eq 1** can be also expressed in

104 the context for a single chromosome $\ell_i = \sum_u^{\mathcal{B}_i} \ell_u + \sum_{u \neq v}^{\mathcal{B}_i} \ell_{uv}$, in which $\mathcal{B}_i = \frac{m_i}{m}$ the number of SNP

105 segments, each of which has m SNPs. Geometrically it leads to $\mathcal{B}_i$ diagonal grids and $\frac{\mathcal{B}_i(\mathcal{B}_i - 1)}{2}$ unique

106 off-diagonal grids (**Figure 1B**).

LD-decay regression

109 As human genome can be boiled down to small LD blocks by genome-widely spread recombination

110 hotspots (Hinch *et al.*, 2019; Li *et al.*, 2022), mechanically there is self-similarity for each chromosome

111 that the relatively strong ℓ_i for juxtaposed grids along the diagonal but weak ℓ_{ij} for grids slightly

112 off-diagonal. So, for a chromosomal ℓ_i, we can further express it as

$$\ell_i = \frac{1}{\mathcal{B}_i^2} \left(\sum_u^{\mathcal{B}_i} \ell_u + \sum_{u \neq v}^{\mathcal{B}_i} \ell_{uv} \right) = E(\ell_u) \frac{1}{\mathcal{B}_i} + E(\ell_{uv}) \left(1 - \frac{1}{\mathcal{B}_i} \right) = \frac{1}{\mathcal{B}_i} [E(\ell_u) - E(\ell_{uv})] + E(\ell_{uv}) \qquad \textbf{(Eq 2)}$$

113 in which ℓ_u is the mean LD for a diagonal grid, ℓ_{uv} the mean LD for off-diagonal grids, and m_i the

114 number of SNPs on the i^{th} chromosome. Consider a linear model below,

$$\boldsymbol{\ell} = b_0 + b_1 \boldsymbol{x} + e \qquad \textbf{(Eq 3)}$$

115   in which $x_i = \frac{1}{m_i}$ the inversion of the SNP number of the i^{th} chromosome. After some algebra, if

116    $E(\ell_u) \gg E(\ell_{uv})$ – say if the former is one order greater than the latter, the interpretation of b_1 and b_0

117   can be

$$\begin{cases} E(b_1) = E(\ell_u - \ell_{uv})m \approx E(\ell_u)m \\ E(b_0) = E(\ell_{uv}) \end{cases} \tag{Eq 4}$$

It should be noticed that $E(b_1) \approx E(\ell_u)m$ quantifies the averaged LD decay of the genome.

Conventional LD decay is analysed via the well-known LD decay analysis, but **Eq 4** provides a direct

estimate of both LD decay and possible existence of extended LD. We will see the application of the

model in **Figure 5** that the strength of the long-distance LD is associated with population structure. Of note,

the underlying assumption of **Eq 3** and **Eq 4** is genome-wide spread of recombination hotspots, an

established result that has been revealed and confirmed (Hinch *et al.*, 2019).

Efficient estimation for ℓ_g, ℓ_i, and ℓ_{ij}

For the aforementioned analyses, the bottleneck obviously lies in the computational cost in estimating ℓ_i

and ℓ_{ij}. ℓ_i and ℓ_{ij} are used to be estimated via the current benchmark algorithm as implemented in

PLINK (Chang *et al.*, 2015), and the computational time complex is proportional to $\mathcal{O}(nm^2)$. We present

a novel approach to estimate ℓ_i and ℓ_{ij}. Given a genotypic matrix **X**, a $n \times m$ matrix, if we assume that

there are m_i and m_j SNPs on chromosomes i and j, respectively, we can construct $n \times n$ genetic

relatedness matrices as below

$$\begin{cases} \mathbf{G}_i = \dfrac{1}{m_i}\widetilde{\mathbf{X}}_i\widetilde{\mathbf{X}}_i^{\mathrm{T}} \\ \mathbf{G}_j = \dfrac{1}{m_j}\widetilde{\mathbf{X}}_j\widetilde{\mathbf{X}}_j^{\mathrm{T}} \end{cases} \tag{Eq 5}$$

in which $\widetilde{\mathbf{X}}_i$ is the standardized $\mathbf{X}_i$ and $\tilde{x}_{kl} = \frac{x_{kl}-2p_l}{\sqrt{2(1+F)p_l q_l}}$, where x_{kl} is the genotype for the k^{th}

individual at the l^{th} biallelic locus, F is the inbreeding coefficient having the value of 0 for random mating

population and 1 for an inbred population, p_l and q_l are the frequencies of the reference and the alternative

alleles ($p_l + q_l = 1$), respectively. When GRM is given, we can obtain some statistical characters of $\mathbf{G}_i$.

From $\mathbf{G}_i$, we extract lower-triangle off-diagonal matrix $\mathbf{G}_{i_o}$ and diagonal matrix $\mathbf{G}_{i_d}$, then we decompose

$\mathbf{G}_i = \mathbf{G}_{i_o} + \mathbf{G}_{i_o}^{\mathrm{T}} + \mathbf{G}_{i_d}$. The mathematical expectation of $\mathbf{G}_{i_o}^2$, in which $E(\mathbf{G}_{i_o}^2) = \frac{1}{n(n-1)}\sum_{k_1 \neq k_2}^n \mathbf{G}_{k_1,k_2}^2$, can

be established according to Isserlis's theorem in terms of the four-order moment (Isserlis, 1918),

$$E\left(\mathbf{G}_{i_o}^2\right) = \frac{1}{m_i^2 n(n-1)}\sum_{k_1 \neq k_2}^{n}\sum_{l_1,l_2}^{m_i}\left[\left(1 + \theta_{k_1 k_2}^2\right)\rho_{l_1 l_2}^2 + \theta_{k_1 k_2}^2\right] \tag{Eq 6}$$

139   in which $E\left(\theta_{k_1 k_2}\right) = \left(\frac{1}{2}\right)^r$ is the expected relatedness score. $r = 0$ for the same individual, and $r = 1$ for

140   first degree of relatives. Similarly, we can derive for $E\left(\mathbf{G}_{i_o}\mathbf{G}_{j_o}\right)$. **Eq 6** establishes the connection between

141   GRM and the aggregated LD estimation that $\ell_i = E\left(\mathbf{G}_{i_o}^2\right)$. According to Delta method (Lynch and Walsh,

142   1998), the means and the sampling variances for ℓ_i and ℓ_{ij} are,

$$\begin{cases} E\left(\mathbf{G}_{i_o}^2\right) = \ell_i = \frac{1}{m_i^2}\sum_{l_1,l_2}^{m_i}\rho_{l_1 l_2}^2 \\[2mm] var(\ell_i) = \frac{4\left[\widehat{var}\left(\mathbf{G}_{i_o}\right)\right]^2}{n(n-1)} \\[2mm] E\left(\mathbf{G}_{i_o}\mathbf{G}_{j_o}\right) = \ell_{ij} = \frac{1}{m_i m_j}\sum_{l_1,l_2=1}^{m_i,m_j}\rho_{l_1 l_2}^2 \\[2mm] var(\ell_{ij}) = \frac{2\left\{\widehat{var}\left(\mathbf{G}_{i_o}\right)\widehat{var}\left(\mathbf{G}_{j_o}\right) + \left[\widehat{cov}\left(\mathbf{G}_{i_o},\mathbf{G}_{j_o}\right)\right]^2\right\}}{n(n-1)} \end{cases} \tag{Eq 7}$$

143   in   which $var\left(\mathbf{G}_{i_o}\right) = E\left(\mathbf{G}_{i_o}^2\right) - \left[E\left(\mathbf{G}_{i_o}\right)\right]^2 = \ell_i - \frac{1}{(n-1)^2}$ and

144    $cov\left(\mathbf{G}_{i_o},\mathbf{G}_{j_o}\right) = E\left(\mathbf{G}_{i_o}\mathbf{G}_{j_o}\right) - E\left(\mathbf{G}_{i_o}\right)E\left(\mathbf{G}_{j_o}\right) = \ell_{ij} - \frac{1}{(n-1)^2}$, respectively. Of note, the properties of ℓ_g can

145   be derived similarly if we replace ℓ_i with ℓ_g in **Eq 7**. We can develop $\tilde{\ell}_{ij}$, a scaled version of ℓ_{ij}, as below

$$\tilde{\ell}_{ij} = \frac{\ell_{ij}}{\sqrt{\tilde{\ell}_i \tilde{\ell}_j}} \tag{Eq 8}$$

146   in which $\tilde{\ell}_i = \frac{m_i \ell_i - 1}{m_i - 1}$, a modification that removed the LD with itself. According to Delta method, the

147   sampling variance of $\tilde{\ell}_{ij}$ is

$$var(\tilde{\ell}_{ij}) = \frac{2\left(\widehat{\tilde{\ell}_{ij}}\right)^2}{n(n-1)}\left[\frac{\widehat{var}\left(\mathbf{G}_{i_o}\right)\widehat{var}\left(\mathbf{G}_{j_o}\right)}{\left(\widehat{cov}\left(\mathbf{G}_{i_o},\mathbf{G}_{j_o}\right)\right)^2} + \frac{\left(\widehat{cov}\left(\mathbf{G}_{i_o},\mathbf{G}_{j_o}\right)\right)^2}{\widehat{var}\left(\mathbf{G}_{i_o}\right)\widehat{var}\left(\mathbf{G}_{j_o}\right)} - 2\right] \tag{Eq 9}$$

Of note, when there is no LD between a pair of loci, ℓ yields zero and its counterpart PLINK estimate

yields $\frac{1}{n}$, a difference that can be reconciled in practice **(see Figure 2)**.

<u>Raise of LD due to population structure</u>

In this study, the connection between LD and population structure is bridged via two pathways below, in

terms of a pair of loci and of the aggregated LD for all pair of loci. For a pair of loci, their LD is often

simplified as $\rho_{l_1 l_2}^2 = \frac{D_{l_1 l_2}^2}{p_{l_1}q_{l_1}p_{l_2}q_{l_2}}$, but will be inflated if there are subgroups (Nei and Li, 1973). In addition,

it is well established the connection between population structure and eigenvalues, and in particular the

largest eigenvalue is associated with divergence of subgroups (Patterson *et al.*, 2006). In this study, the

existence of subgroups of cohort is surrogated by the largest eigenvalue λ_1 or $\bar{F}_{st} \approx \frac{\lambda_1}{n}$.

Data description and quality control

The 1KG (Auton *et al.*, 2015), which is launched to produce a deep catalogue of human genomic variation

by whole genome sequencing (WGS) or whole exome sequencing (WES), and 2,503 strategically selected

individuals of global diversity are included (containing 26 cohorts). We used the following criteria for SNP

inclusion for each of the 26 1KG cohorts: i) autosomal SNPs only; ii) SNPs with missing genotype rates

higher than 0.2 were removed, and missing genotypes were imputed; iii) Only SNPs with minor allele

frequencies higher than 0.05 were retained. Then 2,997,635 consensus SNPs that were present in each of

the 26 cohorts were retained. According to their origins, the 26 cohorts are grouped as African (AFR: MSL,

GWD, YRI, ESN, ACB, LWK, and ASW), European (EUR: TSI, IBS, CEU, GBR, and FIN), East Asian

(EA: CHS, CDX, KHV, CHB, and JPT), South Asian (SA: BEB, ITU, STU, PJL, and GIH), and American

(AMR: MXL, PUR, CLM, and PEL), respectively.

In addition, to test the capacity of the developed software (X-LD), we also included CONVERGE cohort

($n = 10,640$), which was used to investigate Major Depressive Disorder (MDD) in the Han Chinese

population (Cai *et al.*, 2015). We performed the same criteria for SNP inclusion as that of the 1KG cohorts,

and $m = 5,215,820$ SNPs were remained for analyses.

X-LD software implementation

The proposed algorithm has been realized in our X-LD software, which X-LD is written in C++ and reads

in binary genotype data as often used in PLINK. As multi-thread programming is adopted, the efficiency

of X-LD can be improved upon the availability of computational resources. We have tested X-LD in

various independent datasets for its reliability and robustness. Certain data management options, such as

flexible inclusion or exclusion of chromosomes, have been built into the commands of X-LD. In X-LD,

missing genotypes are naively imputed according to Hardy-Weinberg proportions.

The most time-consuming part of X-LD was the construction of GRM $\mathbf{G} = \frac{1}{m}\widetilde{\mathbf{X}}\widetilde{\mathbf{X}}^T$, and the established

computational time complex was $\mathcal{O}(n^2 m)$. However, if $\widetilde{\mathbf{X}}$ is decomposed into $\widetilde{\mathbf{X}} = [\widetilde{\mathbf{X}}_{[t_{1,}]} \vdots \widetilde{\mathbf{X}}_{[t_{2,}]} \vdots \cdots \vdots$

$\widetilde{\mathbf{X}}_{[t_{z,}]}]$, in which $\widetilde{\mathbf{X}}_{[t_{i,}]}$ has dimension of $n \times B$, using Mailman algorithm the computational time complex

for building $\mathbf{G}$ can be reduced to $\mathcal{O}(\frac{n^2 m}{\log_3 m})$ (Liberty and Zucker, 2009). This idea of embedding Mailman

algorithm into certain high throughput genomic studies has been successful, and our X-LD software is also

leveraged by absorbing its recent practice (Wu and Sankararaman, 2018).

Results

Statistical properties of the proposed method

As schematically illustrated in **Figure 1**, ℓ_g could be decomposed into C ℓ_i and $\frac{C(C-1)}{2}$ unique ℓ_{ij}

components. We compared the estimated ℓ_i and ℓ_{ij} in X-LD with those being estimated in PLINK

(known as "--r2"). Considering the substantial computational cost of PLINK, only 100,000 randomly

selected autosome SNPs were used for each 1KG cohort, and 22 $\hat{\ell}_i$ and 231 $\hat{\ell}_{ij}$ were estimated. After

regressing 22 $\hat{\ell}_i$ against those of PLINK, we found that the regression slope was close to unity and bore

an anticipated intercept a quantity of approximately $\frac{1}{n}$ (**Figure 2A and Figure 2B**). In other words,

PLINK gave $\frac{1}{n}$ even for SNPs of no LD. However, when regressing 231 $\hat{\ell}_{ij}$ estimates against those of

PLINK, it was found that largely because of tiny quantity of $\hat{\ell}_{ij}$ it was slightly smaller than 1 but

statistically insignificant from 1 in these 26 1KG cohorts (mean of 0.86 and s.d. of 0.10, and its 95 %

confidence interval was (0.664, 1.056)); when the entire 1KG samples were used, its much larger LD due

to subgroups, nearly no estimation bias was found (**Figure 2A and Figure 2B**). In contrast, because of

their much larger values, $\hat{\ell}_i$ components were always consistent with their corresponding estimates from

PLINK (mean of 1.03 and s.d. of 0.012, 95% confidence interval was (1.006, 1.053), bearing an ignorable

bias). Furthermore, we also combined the African cohorts together (MSL, GWD, YRI, ESN, LWK,

totaling 599 individuals), the East Asian cohorts together (CHS, CDX, KHV, CHB, and JPT, totaling 504

individuals), and the European cohorts together (EUR: TSI, IBS, CEU, GBR, and FIN, totaling 503

individuals), the resemblance pattern between X-LD and PLINK was similar as observed in each cohort

alone **(Figure S1)**. The empirical data in 1KG verified that the proposed method was sufficiently accurate.

To fairly evaluate the computational efficiency of our proposed method, the benchmark comparison was

conducted on the first chromosome of the entire 1KG dataset ($n = 2,503$ and $m = 225,967$), and 10

CPUs were used for multi-thread computing. Compared with PLINK, the calculation efficiency of X-LD

was nearly 30~40 times faster for the tested chromosome, and its computational time of X-LD was

proportional to $\mathcal{O}\left(\frac{n^2 m}{\log_3 m}\right)$ **(Figure S2)**. So, X-LD provided a feasible and reliable estimation of large-scale

complex LD patterns. More detailed computational time of the tested tasks would be reported in their

corresponding sections below; since each 1KG cohort has sample size around 100, otherwise specified the

computational time was only reported for CHB ($n = 103$) as a reference **(Table 1)**. In order to test the

capability of the software, the largest dataset tested was CONVERGE ($n = 10,640$, and $m = 5,215,820$),

and it took 77,508.00 seconds, about 22 hours, to estimate 22 autosomal $\hat{\ell}_i$ and 231 $\hat{\ell}_{ij}$ **(Figure 1A)**;

When zooming into chromosome 2 of CONVERGE, on which 420,949 SNP had been evenly split into

1,000 blocks and yielded 1000 $\hat{\ell}_u$ grids, and 499,500 $\hat{\ell}_{uv}$ LD grids, it took 45,125.00 seconds, about 12.6

hours, to finished the task **(Figure 1B)**.

<u>Ubiquitously extended LD and population structure/admixture</u>

We partitioned the 2,997,635 SNPs into 22 autosomes **(Figure 3A and Figure S3)**, and the general LD

patterns were as illustrated for CEU, CHB, YRI, ASW, and 1KG. As expected, $\hat{\ell}_{ij} < \hat{\ell}_g < \hat{\ell}_i$ for each

cohort **(Figure 3B)**. As observed in these 1KG cohorts, all these three LD measures were associated with

population structure, which was surrogated by $\bar{F}_{st} = \frac{\lambda_1}{n}$, and their squared correlation R^2 were greater

than 0.8. ACB, ASW, PEL, and MXL, which all showed certain admixture, tended to have much greater

$\hat{\ell}_g$, $\hat{\ell}_i$, and $\hat{\ell}_{ij}$ **(Table 2 and Figure 3B)**. In contrast, East Asian (EA) and European (EUR) orientated

cohorts, which showed little within cohort genetic differentiation – as their largest eigenvalues were

slightly greater than 1, had their aggregated LD relatively low and resembled each other **(Table 2)**.

Furthermore, for several European (TSI, IBS, and FIN) and East Asian (JPT) cohorts, the ratio between

$\hat{\ell}_{ij}$ and $\hat{\ell}_i$ components could be smaller than 0.1, and the smallest ratio was found to be about 0.091 in

FIN. The largest ratio was found in 1KG that $\hat{\ell}_{ij}$ =5.7e-3 and $\hat{\ell}_i$ =6.5e-3, and the ratio was 0.877

because of the inflated LD due to population structure. A more concise statistic to describe the ratio

between ℓ_{ij} and ℓ_i was $\tilde{\ell}_{ij}$, **Eq 8**, and the corresponding values for 231 scaled $\tilde{\ell}_{ij}$ for FIN was

$\hat{\bar{\tilde{\ell}}}_{ij} = 0.10$ (s.d. of 0.027) and for 1KG was $\hat{\bar{\tilde{\ell}}}_{ij} = 0.88$ (s.d. of 0.028).

In terms of computational time, for 103 CHB samples, it took about 101.34 seconds to estimate 22

autosomal $\hat{\ell}_i$ and 231 $\hat{\ell}_{ij}$; for all 1KG 2,503 samples, X-LD took about 3,008.29 seconds **(Table 1)**.

Conventional methods took too long to complete the analyses in this section, so no comparable

computational time was provided. For detailed 22 $\hat{\ell}_i$ and 231 $\hat{\ell}_{ij}$ estimates for each 1KG cohort, please

refer to **Extended Data 1 (Excel Sheet 1-27)**.

<u>Detecting exceedingly high LD grids shaped by variable recombination rates</u>

We further explored each autosome with high-resolution grid LD visualization. We set $m = 250$, so each

grid had the ℓ_{uv} for 250×250 SNP pairs. The computational time complex was $\mathcal{O}(n^2 \left(m_i + \frac{B_i^2}{4}\right))$, in

which $\mathcal{B}_i = \frac{m_i}{250}$, and with our proposed method in CHB it costed 66.86 seconds for chromosome 2, which

had the most 241,241 SNPs and was totaled 466,095 unique grids, and 3.22 seconds for chromosome 22,

which had the least 40,378 SNPs and was totaled 13,203 unique grids **(Table 1)**. In contrast, under

conventional methods those LD grids were not very likely to be exhaustively surveyed because of its

computational cost was $\mathcal{O}(nm_i^2)$: for CHB chromosome 2, it would have taken about 40 hours as

estimated. As the result was very similar for $m = 500$ **(Figure S4)**, we only reported the results under

$m = 250$ below.

As expected, chromosome 6 (206,165 SNPs, totaling 340,725 unique grids) had its HLA cluster showing

much higher LD than the rest of chromosome 6. In addition, we found very dramatic variation of HLA

cluster LD $\hat{\ell}_{HLA}$ (28,477,797-33,448,354 bp, totaling 3,160 unique grids) across ethnicities. For CEU,

CHB, YRI, and ASW, their $\hat{\ell}_6 = 0.0010$, 0.00090, 0.00064, and 0.0019, respectively, but their

corresponding HLA cluster grids had $\hat{\ell}_{HLA} = 0.042$, 0.029, 0.025, and 0.022, respectively **(Figure 4)**.

Consequently, the largest ratio for $\frac{\hat{\ell}_{HLA}}{\hat{\ell}_6}$ was of 42.00 in CEU, 39.06 in YRI, and 32.22 in CHB, but was

reduced to 11.58 in ASW. Before the release of CHM13 (Hoyt *et al.*, 2022), chromosome 11 had the most

completely sequenced centromere region, which had much rarer recombination events, all four cohorts

showed an strong LD $\hat{\ell}_{11.c}$ around the centromere (46,061,947-59,413,484 bp, totaling 1,035 unique grids)

regardless of their ethnicities **(Figure 4)**. $\hat{\ell}_{11} = 0.0012, 0.0012, 0.00084$, and 0.0022, respectively, and

$\hat{\ell}_{11.c} = 0.098, 0.10, 0.079$, and 0.094, respectively; the ratio for $\frac{\hat{\ell}_{11.c}}{\hat{\ell}_{11}} = 81.67, 83,33$, and $94,05$, for CEU,

CHB, and YRI, respectively; the lowest ratio was found in ASW of 42.73. In addition, removing the HLA

region of chromosome 6 or the centromere region of chromosome 11 would significantly reduce $\hat{\ell}_6$ or

$\hat{\ell}_{11}$ in comparison with the randomly removal of other regions **(Figure S5)**.

<u>Model-based LD decay regression revealed LD composition</u>

The real LD block size was not exact of $m = 250$ or $m = 500$, but an unknown parameter that should

be inferred in computational intensive "LD decay" analysis (Zhang *et al.*, 2019; Chang *et al.*, 2015). We

conducted the conventional LD decay for the 26 1KG cohorts **(Figure 5A)**, and the time cost was 1,491.94

seconds for CHB. For each cohort, we took the area under the LD decay curve in the LD decay plot, and it

quantified approximately the LD decay score for each cohort. The smallest score was 0.0421 for MSL and

the largest was 0.0598 for PEL **(Table 4)**. However, this estimation was not taken into account the real

extent of LD, so it was not precise enough to reflect the LD decay score. For example, for admixture

population, such as American cohorts, the extent of LD would be longer.

In contrast, we proposed a model-based method, as given in **Eq 3**, which could estimate LD decay score

(regression coefficient b_1) and long-distance LD score (intercept b_0) jointly. Given the estimated 22 $\hat{\ell}_i$

(Extended data 1; Table 3 for four representative cohorts), we regressed each autosomal $\hat{\ell}_i$ against its

correspondingly inversion of SNP number, and all yielded positive slopes (Pearson's correlation

$\mathcal{R} > 0.80$, **Table 4; Figure 5B**), an observation that was consistent with genome-wide spread of

recombination hotspots. This linear relationship could consequently be considered the norm for a relative

homogenous population as observed in most 1KG cohorts **(Figure S6)**, while for the all 2,503 1KG

samples $\mathcal{R} = 0.55$ only **(Table 4)**, indicating that the population structure and possible differentiated

recombination hotspots across ethnicities disturbed the assumption underlying **Eq 3** and smeared the

linearity. We extracted $\hat{b}_0$ and $\hat{b}_1$ for the 26 1KG cohorts for further analysis. The rates of LD decay

score, as indicated by $\hat{b}_1$, within the African cohorts (AFR) were significantly faster than other continents,

consistent with previous observation that African population had relative shorter LD (Gabriel *et al.*, 2002);

while subgroups within the American continent (AMR) tended to have extended LD range due to their

admixed genetic composition **(Table 4** and **Figure 5B)**. Notably, the correlation between $\hat{b}_1$ and the

approximated LD decay score was $\mathcal{R} = 0.88$. The estimated $\bar{F}_{st}$ were highly correlated with $\hat{b}_0$

$(\mathcal{R} = 0.94)$.

A common feature was universally relative high LD of chromosome 6 and 11 in the 26 1KG cohorts

(Figure S6). We quantified the impact of chromosome 6 and 11 by leave-one-chromosome-out test in

CEU, CHB, YRI, and ASW for details **(Figure 6A** and **6B)**, and found that chromosome 6 could lift $\mathcal{R}$ on

average by 0.017, and chromosome 11 by 0.046. One possible explanation was that the centromere regions

of chromosomes 6 and 11 have been assembled more completely than other chromosomes before the

completion of CHM13 (Hoyt *et al.*, 2022), whereas meiotic recombination tended to be reduced around the

centromeres (Hinch *et al.*, 2019). We estimated ℓ_i after having knocked out the centromere region

(46,061,947-59,413,484 bp, chr 11) in CEU, CHB, YRI, and ASW, and chromosome 11 then did not

deviate much from their respective fitted lines **(Figure 6C)**. A notable exceptional pattern was found in

ASW, the chromosome 8 of which had even more deviation than chromosome 11 ($\mathcal{R}$ was 0.83 and 0.87

with and without chromosome 8 in leave-one-chromosome out test) **(Figure 6B)**. The deviation of

chromosome 8 of ASW was consistent even more SNPs were added **(Figure S7)**. We also provided

high-resolution LD grids illustration for chromosome 8 (163,436 SNPs, totaling 214,185 grids) of the four

representative cohorts for more detailed virtualization **(Figure 6D)**. ASW had $\hat{\ell}_8 =0.0022$, but 0.00075,

0.00069 and 0.00043 for CEU, CHB, and YRI, respectively.

Discussion

In this study, we present a computationally efficient method to estimate mean LD of genomic grids of

many SNP pairs. Our LD analysis framework is based on GRM, which has been embedded in variance

component analysis for complex traits and genomic selection (Goddard, 2009; Visscher *et al.*, 2014; Chen,

2014). The key connection from GRM to LD was bridged via the transformation between $n \times n$ matrix

and $m \times m$ matrix, in particular here via Isserlis's theorem under the fourth-order moment (Isserlis, 1918).

With this connection, the computational cost for estimating the mean LD of $m \times m$ SNP pairs is reduced

from $\mathcal{O}(nm^2)$ to $\mathcal{O}(n^2 m)$, and the statistical properties of the proposed method are derived in theory and

validated in 1KG datasets. In addition, as the genotype matrix $\mathbf{X}$ is of limited entries $\{0, 1, 2\}$, assuming

missing genotypes are imputed first, using Mailman algorithm the computational cost of GRM can be further

reduced to $\mathcal{O}\left(\frac{n^2 m}{\log_3 m}\right)$ (Liberty and Zucker, 2009). The largest data tested so far for the proposed method

has the sample size of 10,640 and of more than 5 million SNPs, it can complete genomic LD analysis in

77,508.00 seconds **(Table 1)**. Obviously, with the availability of such as UK Biobank data (Bycroft *et al.*,

2018), the proposed method may not be adequate and other new methods are needed.

We also applied the proposed method into 1KG and revealed certain characteristics of the human genomes.

Firstly, we found the ubiquitously existence of extended LD, which was likely emerged because of

population structure, even very slightly, and admixture history. We quantified the $\hat{\ell}_i$ and $\hat{\ell}_{ij}$ in 1KG,

and as indicated by $\tilde{\ell}_{ij}$ we found the inter-chromosomal LD was nearly an order lower than

intra-chromosomal LD; for admixed cohorts, the ratio was much higher, even very close to each other such

as in all 1KG samples. Secondly, variable recombination rates shaped peak of local LD. For example, the

HLA region showed high LD in European and East Asian cohorts, but relatively low LD in such as YRI,

consistent with their much longer population history. Thirdly, it existed general linear correlation between

ℓ_i and the inversion of the SNP number, a long-anticipated result that is as predicted with genome-wide

spread of recombination hotspots (Hinch *et al.*, 2019). One outlier of this linear norm was chromosome 11,

which had so far most completely genotyped centromere and consequently had more elevated LD

compared with other autosomes. We anticipate that with the release of CHM13 the linear correlation

should be much closer to unity (Hoyt *et al.*, 2022). Of note, under the variance component analysis for

complex traits, it is often a positive correlation between the length of a chromosome (as surrogated by the

number of SNPs) and the proportion of heritability explained (Chen *et al.*, 2014).

In contrast, throughout the study recurrent outstanding observations were found in ASW. For example, in

ASW the ratio of $\hat{\ell}_{HLA}/\hat{\ell}_6$ was substantially dropped down compared with that of CEU, CHB, or YRI as

illustrated in **Figure 4**. Furthermore, chromosome 8 in ASW fluctuated upwards most from the linear

correlation (**Figure 6**), and even after various analyses, such as expanding SNP numbers. One possible

explanation may lay under the complex demographic history of ASW, which can be investigated and

tested in additional African American samples or possible existence for epistatic fitness (Ni *et al.*, 2020).

Data availability

Public genetic datasets used in this study can be freely downloaded from the following URLs.

1000 Genomes Project: https://wwwdev.ebi.ac.uk/eva/?eva-study=PRJEB30460

CONVERGE: https://wwwdev.ebi.ac.uk/eva/?eva-study=PRJNA289433

Acknowledgements

GBC conceived and initiated the study. XH, TNZ, and YL conducted simulation and analyzed data. GBC,

XH, TNZ, and JZ developed the software. GBC wrote the first draft of the paper. All authors contributed

to the writing, discussion of the paper, and validation of the results. This work was supported by National

Natural Science Foundation of China (31771392) and GZY-ZJ-KJ-23001. The funders played no role in

designing, preparation, and submission of the paper.

Conflict of Interests: None.

373   **Reference**

Ardlie KG, Kruglyak L, Seielstad M. Patterns of linkage disequilibrium in the human genome. *Nat Rev Genet*

2002;**3**:299–309.

Auton A, Brooks LD, Durbin RM *et al*. A global reference for human genetic variation. *Nature*

2015;**526**:68–74.

Barrett JC, Fry B, Maller J *et al.* Haploview: Analysis and visualization of LD and haplotype maps.

Bioinformatics 2005;**21**:263–5.

Bulik-Sullivan BK, Loh P-R, Finucane HK *et al.* LD Score regression distinguishes confounding from

polygenicity in genome-wide association studies. *Nat Genet* 2015;**47**:291–5.

Bycroft C, Freeman C, Petkova D *et al.* The UK Biobank resource with deep phenotyping and genomic data.

Nature 2018;**562**:203–9.

Cai N, Bigdeli T, Kretzschmar W *et al.* Sparse whole-genome sequencing identifies two loci for major

depressive disorder. *Nature* 2015;**523**:588–91.

Chang CC, Chow CC, Tellier LC *et al.* Second-generation PLINK: rising to the challenge of larger and richer

datasets. *Gigascience* 2015;**4**:7.

Chen GB. Estimating heritability of complex traits from genome-wide association studies using IBS-based

Haseman–Elston regression. *Front Genet* 2014;**5**:107.

Chen GB, Lee SH, Brion MJA *et al.* Estimation and partitioning of (co)heritability of inflammatory bowel

disease from GWAS and immunochip data. *Hum Mol Genet* 2014;**23**:4710–20.

Das S, Forer L, Schönherr S *et al.* Next-generation genotype imputation service and methods. *Nat Genet*

2016;**48**:1284-7.

Gabriel SB, Schaffner SF, Nguyen H *et al.* The structure of haplotype blocks in the human genome. *Science*

2002;**296**:2225–9.

Goddard M. Genomic selection: prediction of accuracy and maximisation of long term response. *Genetica*

2009;**136**:245–57.

Hill WG, Robertson A. The effect of linkage on limits to artificial selection. *Genet Res* 1966;**8**:269–94.

Hinch AG, Zhang G, Becker PW *et al.* Factors influencing meiotic recombination revealed by whole-genome

sequencing of single sperm. *Science* 2019;**363**:eaau8861.

Hoyt SJ, Storer JM, Hartley GA *et al.* From telomere to telomere: The transcriptional and epigenetic state of

human repeat elements. *Science* 2022;**376**:eabk3112.

Isserlis L. On a formula for the product-moment coefficient of any order of a normal frequency distribution in

any number of variables. *Biometrika* 1918;**12**:134–9.

Li Y, Chen SY, Rapakoulia T *et al.* Deep learning identifies and quantifies recombination hotspot determinants.

Bioinformatics 2022;**38**:2683–91.

Liberty E, Zucker SW. The Mailman algorithm: A note on matrix-vector multiplication. *Inf Process Lett*

2009;**109**:179–82.

Lowy-Gallego E, Fairley S, Zheng-Bradley X *et al.* Variant calling on the GRCh38 assembly with the data from

phase three of the 1000 Genomes Project. *Wellcome Open Res* 2019;**4**:50.

Lynch M, Walsh B. *Genetics and analysis of quantitative traits.* Sunderland, MA: Sinauer, 1998.

Myers S, Bottolo L, Freeman C *et al.* A fine-scale map of recombination rates and hotspots across the human

genome. *Science* 2005;**310**:321–4.

Nei M, Li W-H. Linkage disequilibrium in subdivided populations. *Genetics* 1973;**75**:213–9.

Ni XM, Zhou MS, Wang HM *et al.* Detecting fitness epistasis in recently admixed populations with

genome-wide data. *BMC Genomics* 2020;**21**:476.

Patterson N, Price AL, Reich D. Population structure and eigenanalysis. *PLoS Genet* 2006;**2**:e190.

Theodoris C, Low MT, Pavlidis P *et al.* quickLD: An efficient software for linkage disequilibrium analyses. *Mol*

Ecol Resour 2021;**21**:2580–7.

Vilhjálmsson BJ, Yang J, Finucane HK *et al.* Modeling linkage disequilibrium increases accuracy of polygenic

risk scores. *Am J Hum Genet* 2015;**97**:576–92.

Visscher PM, Hemani G, Vinkhuyzen AAE *et al.* Statistical power to detect genetic (co) variance of complex

traits using SNP data in unrelated samples. *PLoS Genet* 2014;**10**:e1004269.

Weir BS. Linkage disequilibrium and association mapping. *Annu Rev Genomics Hum Genet* 2008;**9**:129–42.

Wu Y, Sankararaman S. A scalable estimator of SNP heritability for biobank-scale data. *Bioinformatics*

2018;**34**:187–94.

Yang J, Weedon MN, Purcell S *et al.* Genomic inflation factors under polygenic inheritance. *Eur J Hum Genet*

2011;**19**:807–12.

Yang S, Zhou X. Accurate and scalable construction of polygenic scores in large Biobank data sets. *Am J Hum*

Genet 2020;**106**:679–93.

Zhang C, Dong SS, Xu JY *et al.* PopLDdecay: A fast and effective tool for linkage disequilibrium decay

analysis based on variant call format files. *Bioinformatics* 2019;**35**:1786–8.

Zhou X. A unified framework for variance component estimation with summary statistics in genome-wide

association studies. *Ann Appl Stat* 2017;**11**:2027–51.

Figure 1 Schematic illustration for large-scale LD analysis as exampled for CONVERGE cohort. **A)** The 22 human autosomes have consequently 22 $\hat{\ell}_i$ and 231 $\hat{\ell}_{ij}$, without (left) and with (right) scaling transformation; Scaling transformation is given in **Eq 8**. **B)** If zoom into chromosome 2 of 420,946 SNPs, a chromosome of relative neutrality is expected to have self-similarity structure that harbors many approximately strong $\hat{\ell}_u$ along the diagonal, and relatively weak $\hat{\ell}_{uv}$ off-diagonally. Here chromosome 2 of CONVERGE has been split into 1,000 blocks and yielded 1000 $\hat{\ell}_u$ LD grids, and 499,500 $\hat{\ell}_{uv}$ LD grids.

Figure 2 Reconciliation for LD estimators in the 26 1KG cohorts. A) Consistency examination for the 26

1KG cohorts for their $\hat{\ell}_i$ and $\hat{\ell}_{ij}$ estimated by X-LD and PLINK (--r2). In each figure, the 22 $\hat{\ell}_i$ fitting line is

in purple, whereas the 231 $\hat{\ell}_{ij}$ fitting line is in green. The gray solid line, $y = \frac{1}{n} + x$, in which n the sample

size of each cohort, represents the expected fit between PLINK and X-LD estimates, and the two estimated

regression models at the top-right corner of each plot shown this consistency. The sample size of each cohort is

in parentheses. **B)** Distribution of R^2 of $\hat{\ell}_i$ and $\hat{\ell}_{ij}$ fitting lines is based on X-LD and PLINK algorithms in

the 26 cohorts.

26 1KG cohorts: MSL (Mende in Sierra Leone), GWD (Gambian in Western Division, The Gambia), YRI

(Yoruba in Ibadan, Nigeria), ESN (Esan in Nigeria), ACB (African Caribbean in Barbados), LWK (Luhya in

Webuye, Kenya), ASW (African Ancestry in Southwest US); CHS (Han Chinese South), CDX (Chinese Dai in

Xishuangbanna, China), KHV (Kinh in Ho Chi Minh City, Vietnam), CHB (Han Chinese in Beijing, China),

JPT (Japanese in Tokyo, Japan); BEB (Bengali in Bangladesh), ITU (Indian Telugu in the UK), STU (Sri

Lankan Tamil in the UK), PJL (Punjabi in Lahore, Pakistan), GIH (Gujarati Indian in Houston, TX); TSI

(Toscani in Italia), IBS (Iberian populations in Spain), CEU (Utah residents (CEPH) with Northern and Western

European ancestry), GBR (British in England and Scotland), FIN (Finnish in Finland); MXL (Mexican Ancestry

in Los Angeles, California), PUR (Puerto Rican in Puerto Rico), CLM (Colombian in Medellin, Colombia),

PEL (Peruvian in Lima, Peru).

Figure 3 Various LD components for the 26 1KG cohorts. A) Chromosomal scale LD components for 5 representative cohorts (CEU, CHB, YRI, ASW, and 1KG). The upper parts of each figure represent $\hat{\ell}_i$ (along the diagonal) and $\hat{\ell}_{ij}$ (off-diagonal), and the lower part $\hat{\tilde{\ell}}_{ij}$ as in **Eq 8**. For visualization purposes, the quantity of LD before scaling is transformed to a -log10 scale, with smaller values (red hues) representing larger LD, and a value of 0 representing that all SNPs are in LD. **B)** The relationship between the degree of population structure (approximated by $\bar{F}_{st}$) and $\hat{\ell}_i$, $\hat{\ell}_g$, and $\hat{\ell}_{ij}$ in the 26 1KG cohorts.

Figure 4 High-resolution illustration for LD grids for CEU, CHB, YRI, and ASW ($m = 250$). For each

cohort, we partition chromosomes 6 and 11 into high-resolution LD grids (each LD grid contains 250×250

SNP pairs). The bottom half of each figure shows the LD grids for the entire chromosome. Further zooming into

HLA on chromosome 6 and the centromere region on chromosome 11, and their detailed LD in the relevant

regions are also provided in the upper half of each figure. For visualization purposes, LD is transformed to a

-log10-scale, with smaller values (red hues) representing larger LD, and a value of 0 representing that all SNPs

are in LD.

Figure 5 LD decay analysis for 26 1KG cohorts. A) Conventional LD decay analysis in PLINK for 26 cohorts.

To eliminate the influence of sample size, the inverse of sample size has been subtracted from the original LD

values. The YRI cohort, represented by the orange dotted line, is chosen as the reference cohort in each plot.

The top-down arrow shows the order of LDdecay values according to **Table 4**. **B)** Model-based LD decay

analysis for the 26 1KG cohorts. We regressed each autosomal   against its corresponding inversion of the

SNP number for each cohort. Regression coefficient   quantifies the averaged LD decay of the genome and

intercept   provides a direct estimate of possible existence of long-distance LD. The   values in the first

three plot indicate the correlation between   and LD decay score in three different physical distance and the

correlation between $\hat{b}_1$ (left-side vertical axis) and LD decay score (right-side vertical axis) and the correlation

between $\hat{b}_0$ (left-side vertical axis) and $\bar{F}_{st}$ (right-side vertical axis), respectively. The last plot assessed the

impact of centromere region of chromosome 11 on the linear relationship between chromosomal LD and the

inverse of the SNP number. The dark and light gray dashed lines represent the mean of the $\mathcal{R}$ with and without

the presence of centromere region of chromosome 11.

Figure 6 The correlation between the inversion of the SNP number and $\hat{\ell}_i$. A) The correlation between the inversion of the SNP number and $\hat{\ell}_i$ in CEU, CHB, YRI, and ASW. **B)** Leave-one-chromosome-out strategy is adopted to evaluate the contribution of a certain chromosome on the correlation between the inverse of the SNP number and $\hat{\ell}_i$. **C)** The correlation between the inversion of the SNP number and chromosomal LD in CEU, CHB, YRI, and ASW after removing the centromere region of chromosome 11. **D)** High-resolution illustration for LD grids for chromosome 8 in CEU, CHB, YRI, and ASW. For each cohort, we partition chromosome 8 into consecutive LD grids (each LD grid contains 250×250 SNP pairs). For visualization purposes, LD is transformed to a -log10-scale, with smaller values (red hues) representing larger LD, and a value of 0 representing that all SNPs are in LD.

504 **Table 1 Computational time for the demonstrated estimation tasks**

Cohort	Task description	Time cost	Computational time complex
CHB ($n = 103$, $m = 2,997,655$)	Estimation for 22 autosomal ℓ_i, and 231 inter-chromosomal ℓ_{ij}. Results see **Figure 3 and Table 2**.	101,34 secs	$\mathcal{O}(n^2 m)$
1KG ($n = 2,503$, $m = 2,997,655$)	Same as above.	3,008.29 secs	Same as above
CONVERGE ($n = 10,640$, $m = 5,215,820$)	Same as above. Result see **Figure 1A**.	77,508.00 secs	Same as above
	Estimation for high-resolution LD interaction given bin size of 250 SNPs		
CHB ($n = 103$, $m_2 = 241,241$)	Chromosome 2, estimation for 965 ℓ_i, and 465,130 ℓ_{ij}. Results see **Figure 4**.	66.86 secs	$\mathcal{O}\left(n^2 \left(m_i + \left(\frac{m_i}{250}\right)^2\right)\right)$
CHB ($n = 103$, $m_{22} = 40,378$)	Chromosome 22, estimation for 162 ℓ_i, and 13,041 ℓ_{ij}. Results see **Figure 4**.	3.22 secs	Same as above
CONVERGE ($n = 10,640$, $m_{22} = 71,407$)	Chromosome 22, estimation for 286 ℓ_i, and 40,755 ℓ_{ij}.	8,736.29 secs	Same as above
CONVERGE ($n = 10,640$, $m_2 = 420,949$)	Chromosome 2, estimation for 1,000 ℓ_i, and 499,500 ℓ_{ij}. Result see **Figure 1B**.	45,125.00 secs	Chromosome 2 was split into 1000 blocks, each of which had about 420 SNPs.

505 **Notes:** for the sake of fair comparison, 10 CPUs were used for multi-thread computing.

Table 2 X-LD estimation for complex LD components (2,997,635 SNPs)

Cohort (n)	Ancestry	λ_1 ($\bar{F}_{st}$)[1]	$\hat{\ell}_g$ (s.e.)[2]	$\hat{\ell}_i$ (s.d.)[3]	$\hat{\ell}_{ij}$ (s.d.)[3]	$\tilde{\ell}_{ij}$ (s.d.)[3]	Lower bound of LD[4]
MSL (85)	AFR	1.10 (0.013)	1.9e-4 (1.21e-6)	6.9e-4 (2.0e-4)	1.7e-4 (1.7e-5)	0.26 (0.053)	0.161971831
GWD (113)	AFR	1.07 (0.009)	1.1e-4 (5.61e-7)	6.0e-4 (2.0e-4)	8.7e-5 (8.1e-6)	0.16 (0.037)	0.247218789
YRI (107)	AFR	1.05 (0.010)	1.1e-4 (4.23e-7)	5.9e-4 (2.0e-4)	8.8e-5 (6.9e-6)	0.16 (0.04)	0.242001641
ESN (99)	AFR	1.09 (0.011)	1.4e-4 (7.67e-7)	7.0e-4 (2.2e-4)	1.2e-4 (1.2e-5)	0.19 (0.043)	0.217391304
ACB (96)	AFR	2.01 (0.021)	2.9e-4 (3.78e-6)	9.1e-4 (2.5e-4)	2.5e-4 (3.6e-5)	0.29 (0.070)	0.147727273
LWK (99)	AFR	1.35 (0.014)	2.2e-4 (2.38e-6)	8.4e-4 (2.5e-4)	1.9e-4 (3.2e-5)	0.24 (0.052)	0.173913043
ASW (61)	AFR	1.90 (0.031)	1.1e-3 (2.73e-5)	2.0e-3 (3.2e-4)	1.1e-3 (6.2e-5)	0.57 (0.059)	0.079681275
CHS (105)	EA	1.08 (0.010)	1.4e-4 (9.39e-7)	9.5e-4 (3.4e-4)	1.0e-4 (1.3e-5)	0.12 (0.030)	0.31147541
CDX (93)	EA	1.11 (0.012)	1.8e-4 (1.38e-6)	1.1e-3 (3.6e-4)	1.4e-4 (2.0e-5)	0.14 (0.040)	0.272277228
KHV (99)	EA	1.07 (0.011)	1.4e-4 (7.67e-7)	9.5e-4 (3.5e-4)	1.0e-4 (1.2e-5)	0.12 (0.031)	0.31147541
CHB (103)	EA	1.07 (0.010)	1.3e-4 (6.94e-7)	9.3e-4 (3.4e-4)	9.5e-5 (1.1e-5)	0.11 (0.030)	0.317948718
JPT (104)	EA	1.06 (0.010)	1.3e-4 (7.22e-7)	1.0e-3 (3.8e-4)	9.3e-5 (1.2e-5)	0.10 (0.028)	0.338638673
BEB (86)	SA	1.07 (0.012)	1.7e-4 (8.09e-7)	9.1e-4 (3.1e-4)	1.4e-4 (1.5e-5)	0.17 (0.042)	0.236363636
ITU (102)	SA	1.61 (0.016)	1.9e-4 (1.84e-6)	9.5e-4 (3.1e-4)	1.5e-4 (1.7e-5)	0.18 (0.044)	0.231707317
STU (102)	SA	1.56 (0.015)	2.6e-4 (3.21e-6)	1.0e-3 (3.3e-4)	2.3e-4 (3.1e-5)	0.23 (0.047)	0.171526587
PJL (96)	SA	1.67 (0.017)	2.4e-4 (2.74e-6)	1.1e-3 (3.4e-4)	2.0e-4 (2.2e-5)	0.21 (0.048)	0.20754717
GIH (103)	SA	1.73 (0.017)	2.7e-4 (3.41e-6)	1.1e-3 (3.4e-4)	2.4e-4 (1.9e-5)	0.23 (0.049)	0.179153094
TSI (107)	EUR	1.07 (0.010)	1.2e-4 (6.10e-7)	9.1e-4 (3.3e-4)	9.0e-5 (1.1e-5)	0.11 (0.029)	0.325
IBS (107)	EUR	1.07 (0.010)	1.2e-4 (6.10e-7)	9.1e-4 (3.3e-4)	8.8e-5 (1.1e-5)	0.11 (0.028)	0.329949239
CEU (99)	EUR	1.07 (0.011)	1.4e-4 (7.67e-7)	9.6e-4 (3.4e-4)	1.1e-4 (1.3e-5)	0.12 (0.030)	0.293577982
GBR (91)	EUR	1.11 (0.012)	1.7e-4 (1.08e-6)	1.0e-3 (3.6e-4)	1.4e-4 (1.8e-5)	0.15 (0.036)	0.253807107
FIN (99)	EUR	1.09 (0.011)	1.5e-4 (9.69e-7)	1.1e-3 (3.8e-4)	1.0e-4 (1.5e-5)	0.10 (0.027)	0.34375
MXL (64)	AMR	2.29 (0.036)	7.2e-4 (1.49e-5)	2.1e-3 (4.1e-4)	6.3e-4 (9.6e-5)	0.32 (0.072)	0.136986301
PUR (104)	AMR	1.43 (0.014)	1.6e-4 (1.30e-6)	1.2e-3 (4.2e-4)	1.2e-4 (1.7e-5)	0.11 (0.026)	0.322580645
CLM (94)	AMR	1.58 (0.017)	2.3e-4 (2.49e-6)	1.4e-3 (4.5e-4)	1.7e-4 (2.6e-5)	0.13 (0.035)	0.281690141
PEL (85)	AMR	2.38 (0.028)	4.5e-4 (7.33e-6)	1.9e-3 (5.1e-4)	3.7e-4 (8.5e-5)	0.21 (0.062)	0.196483971
1KG (2,503)	MIX	164.20 (0.066)	5.8e-3 (4.63e-6)	6.5e-3 (4.1e-4)	5.7e-3 (2.4e-4)	0.88 (0.028)	0.051505547

[1]Eigenvalue was estimated. In parentheses was the ratio between the listed largest eigenvalue and the sample

size. Since it exists an approximation that $\bar{F}_{st} \approx \frac{\lambda_1}{n}$, the ratio can be taken as an approximation of population

structure.

[2] Standard err was calculated as $\frac{2}{\sqrt{n(n-1)}}\left[\hat{l}_g - \frac{1}{(n-1)^2}\right]$, as **Eq 7**.

[3] Estimated empirically from $\mathcal{C}$ chromosomal $\hat{\ell}_i$; Estimated empirically from $\frac{\mathcal{C}(\mathcal{C}-1)}{2}$ inter-chromosomal $\hat{\ell}_{ij}$.

[4] It is estimated by $\frac{22\hat{l}_i}{22\hat{l}_i + 231\hat{\ell}_{ij}}$, indicating lower bound of true LD.

 Table 3 Estimates for 22 autosomal $\hat{\ell}_i$ in CEU, CHB, YRI, and ASW, respectively

Chromosome	SNP number	$\hat{\ell}_i$			
		CEU	**CHB**	**YRI**	**ASW**
1	225,967	5.0e-4 (8.2e-6)	0.00049 (7.8e-6)	0.00032 (4.3e-6)	0.0015 (4e-05)
2	241,241	5.0e-4 (8.1e-6)	5.0e-4 (7.9e-6)	3.0e-4 (4.1e-6)	0.0015 (4e-05)
3	212,670	6.0e-04 (1.0e-5)	0.00058 (9.5e-6)	0.00039 (5.7e-6)	0.0018 (5.1e-5)
4	222,241	0.00062 (1.0e-5)	0.00061 (1.0e-5)	0.00038 (5.4e-6)	0.0018 (5.0e-5)
5	193,632	0.00069 (1.2e-5)	7.0e-04 (1.2e-5)	0.00043 (6.5e-6)	0.0018 (4.9e-5)
6	206,165	0.0010 (1.9e-5)	9.0e-04 (1.6e-5)	0.00064 (1.0e-5)	0.0019 (5.4e-5)
7	177,414	0.00073 (1.3e-5)	0.00071 (1.2e-5)	0.00045 (6.8e-6)	0.0016 (4.3e-5)
8	163,436	0.00075 (1.3e-5)	0.00069 (1.2e-5)	0.00043 (6.5e-6)	0.0022 (6.4e-5)
9	129,440	0.00074 (1.3e-5)	0.00074 (1.3e-5)	0.00047 (7.2e-6)	0.0018 (5.0e-5)
10	152,251	0.00078 (1.4e-5)	8.0e-04 (1.4e-5)	0.00058 (9.3e-6)	0.0019 (5.6e-5)
11	151,751	0.0012 (2.3e-5)	0.0012 (2.2e-5)	0.00084 (1.4e-5)	0.0022 (6.2e-5)
12	139,684	8.0e-4 (1.4e-5)	0.00073 (1.2e-5)	0.00049 (7.5e-6)	0.0017 (4.8e-5)
13	113,390	0.0010 (1.8e-5)	0.00094 (1.6e-5)	0.00061 (9.8e-6)	0.0018 (4.9e-5)
14	97,335	0.0011 (2.0e-5)	0.0010 (1.8e-5)	0.00065 (1.1e-5)	0.0020 (5.6e-5)
15	85,307	0.0010 (1.8e-5)	0.00098 (1.7e-5)	6.0e-4 (9.6e-6)	0.0020 (5.8e-5)
16	92,007	0.00088 (1.6e-5)	0.00084 (1.5e-5)	0.00054 (8.4e-6)	0.0021 (6.2e-5)
17	79,478	0.0012 (2.3e-5)	0.0011 (2.0e-5)	0.00069 (1.1e-5)	0.0021 (6.0e-5)
18	87,105	0.0010 (1.8e-5)	0.00095 (1.7e-5)	0.00058 (9.2e-6)	0.0023 (6.8e-5)
19	72,794	0.0012 (2.3e-05)	0.0012 (2.1e-5)	0.00082 (1.4e-5)	0.0022 (6.2e-5)
20	68,881	0.0014 (2.6e-5)	0.0015 (2.7e-5)	0.00078 (1.3e-5)	0.0024 (7.0e-5)
21	45,068	0.0018 (3.4e-5)	0.0017 (3.2e-5)	0.00098 (1.7e-5)	0.0024 (7.1e-5)
22	40,378	0.0016 (3.1e-5)	0.0016 (2.9e-5)	0.0010 (1.8e-5)	0.0027 (8.1e-5)

516 **Note:** each $\hat{\ell}_i$ and its standard error in parentheses, as estimated in **Eq 7.**

518 **Table 4 LD decay regression analysis for 26 cohorts**

Cohort (n)	LD-decay regression[1]			Population parameters[2]				
	$\hat{b}_0$	$\hat{b}_1$	R		LD decay score	$\bar{F}_{st}$ (%)	Ancestry	True LD[3]
MSL (85)	0.00041	29.97	0.84		0.0421	0.013	AFR	0.62727273
GWD (113)	0.00031	30.17	0.83		0.0439	0.009	AFR	0.65934066
YRI (107)	0.00030	30.64	0.85		0.0436	0.010	AFR	0.66292135
ESN (99)	0.00037	34.82	0.87		0.0436	0.011	AFR	0.65420561
ACB (96)	0.00053	39.62	0.88		0.0451	0.021	AFR	0.63194444
LWK (99)	0.00046	40.52	0.92		0.0447	0.014	AFR	0.64615385
ASW (61)	0.0015	46.88	0.83		0.0472	0.031	AFR	0.57142857
CHS (105)	0.00046	52.36	0.87		0.0555	0.010	EA	0.67375887
CDX (93)	0.00055	53.77	0.83		0.0557	0.012	EA	0.66666667
KHV (99)	0.00044	53.79	0.87		0.0560	0.011	EA	0.68345324
CHB (103)	0.00041	54.90	0.90		0.0558	0.010	EA	0.69402985
JPT (104)	0.00045	57.75	0.85		0.0568	0.010	EA	0.68965517
BEB (86)	0.00045	48.84	0.88		0.0556	0.012	SA	0.66911765
ITU (102)	0.00048	49.58	0.89		0.0546	0.016	SA	0.66433566
STU (102)	0.00055	52.84	0.89		0.0546	0.015	SA	0.64516129
PJL (96)	0.00054	54.00	0.90		0.0546	0.017	SA	0.67073171
GIH (103)	0.00057	55.81	0.91		0.0562	0.017	SA	0.65868263
TSI (107)	0.00041	53.17	0.91		0.0558	0.010	EUR	0.68939394
IBS (107)	0.00039	54.22	0.92		0.0555	0.010	EUR	0.7
CEU (99)	0.00045	54.23	0.89		0.0559	0.011	EUR	0.68085106
GBR (91)	0.00047	58.23	0.91		0.0555	0.012	EUR	0.68027211
FIN (99)	0.00054	59.24	0.86		0.0579	0.011	EUR	0.67073171
MXL (64)	0.0014	66.13	0.89		0.0558	0.036	AMR	0.6
PUR (104)	0.00059	67.20	0.89		0.0571	0.014	AMR	0.67039106
CLM (94)	0.00069	75.97	0.95		0.0572	0.017	AMR	0.66985646
PEL (85)	0.0012	78.15	0.85		0.0598	0.028	AMR	0.61290323
1KG (2,503)	0.0061	40.65	0.55			0.066	Mixed	0.51587302

[1]The regression intercept $\hat{b}_0$ and the coefficients $\hat{b}_1$ is as represented in **Eq 3**.

[2]The column for LD decay score was taken the mean of the estimated $r^2 - \frac{1}{n}$ from PopLDdecay in a

physical distance of 1500kb, which was approximated to the area under the curve in **Figure 5A** for each

cohort; F_{st} was approximated by $\frac{\lambda_1}{n}$, in which λ_1 the largest eigenvalue for the cohort.

[3]True LD is defined as $\frac{\bar{\bar{\ell}}_{ij}}{\bar{\bar{\ell}}_{ij}+\bar{b}_0}$.

Figure S1 Reconciliation for LD estimators in AFR, EAS, and EUR. In each figure, the 22 $\hat{\ell}_i$ fit line is in

purple, whereas the 231 $\hat{\ell}_{ij}$ fit line is in green. The gray solid line, $y = \frac{1}{n} + x$, in which n the sample size,

represents the expected fit between PLINK and X-LD, and the two estimated regression models at the top-right

corner shown this consistency.

Figure S2 The computational efficiency of X-LD algorithm. Considering the high computational cost of

PLINK, only the first chromosome was chosen. In the process of evaluating computational efficiency, we kept

adding SNPs until the inclusion of entire chromosome. The bar chart and line chart show the actual calculation

time and theoretical calculation complexity, respectively.

537   **Figure S3 Chromosomal scale LD components for 26 cohorts in 1KG.** The upper and lower parts of each

538   figure represent the LD before and after scaling according to **Eq 8**. $\hat{\ell}_i$ and $\hat{\ell}_{ij}$ are represented by the diagonal

539   and the off-diagonal elements, respectively. For visualization purposes, LD before scaling is transformed to a

540   -log10-scale, with smaller values (red hues) representing larger LD, and a value of 0 representing that all SNPs

541   are in LD.

Figure S4 High-resolution illustration for LD grids for CEU, CHB, YRI, and ASW ($m = 500$). For each

cohort, we partitioned each chromosome into consecutive LD grids (each LD grid containing 500 SNPs). For

visualization purposes, LD is transformed to a -log10-scale, with smaller values (red hues) representing larger

LD, and a value of 0 representing that all SNPs are in LD.

Figure S5 Influence of HLA region on chromosome 6 and centromere region on chromosome 11 on

chromosomal LD in CEU, CHB, YRI, and ASW. When other region was removed, to avoid chance, the same

number of consecutive SNPs as HLA region or centromere region were randomly removed from the genomic

region, and this operation was repeated 100 times.

553   **Figure S6 The correlation between the inverse of the SNP number and chromosomal LD in 26 cohorts of**

554   **1KG.**

Figure S7 Influence of expanding of SNP numbers on the correlation between the inverse of the SNP

number and chromosomal LD in ASW. Randomly selected SNPs that were presented in ASW but were not

2,997,635 consensus SNPs were added to the ASW cohort to demonstrate the stable pattern of chromosome 8.