**Dysregulated Platelet Function in Patients with Post-Acute Sequelae of COVID-19**

Anu Aggarwal[1], Tamanna K. Singh[2], Michael Pham[2],

Matthew Godwin[1], Rui Chen[1], Thomas M. McIntyre[1], Alliefair Scalise[2], Mina K. Chung[1,2],

Courtney Jennings[1], Mariya Ali[1], Hiijun Park[1], Kristin Englund[3], Alok A. Khorana[1,4],

Lars G. Svensson[5], Samir Kapadia[2], Keith R. McCrae[1,2,3], Scott J. Cameron[1,4,5†]

**Running Title**: Decreased thrombosis in long COVID-19

**Affiliations:**
[1] Lerner Research Institute, Department of Cardiovascular and Metabolic Sciences, Cleveland,
Ohio
[2] Heart Vascular and Thoracic Institute, Department of Cardiovascular Medicine, Section of
Vascular Medicine, Cleveland Clinic Foundation, Cleveland, Ohio, USA
[3] Department of Infectious Disease, Cleveland Clinic Foundation, Cleveland, Ohio, USA
[4] Taussig Cancer Institute, Department of Hematology and Oncology, Cleveland, Ohio
[5] Heart Vascular and Thoracic Institute, Department of Cardiovascular Medicine, Department of
Cardiac Surgery, Cleveland Clinic Foundation, Cleveland, Ohio, USA
17

18
19 [†] **Address for Correspondence**: Section of Vascular Medicine, Heart Vascular and Thoracic
20 Institute, Department of Cardiovascular Medicine, Section of Vascular Medicine, Cleveland Clinic
21 Foundation, Cleveland, Ohio, USA
22 Email: cameros3@ccf.org

23
24 Word count: (abstract) 242, and (main manuscript) 2884
25
26
27 **Key words:** COVID-19, Post-acute sequelae of COVID-19 (PASC), Platelets, Thrombosis.
28
29

**Abstract**

**Background:** Post-acute sequelae of COVID-19 (PASC), also referred as Long-COVID, sometimes follows COVID-19, a disease caused by SARS-CoV-2. While SARS-CoV-2 is well-known to promote a prothrombotic state, less is known about the thrombosis risk in PASC.

**Aim:** Our objective was to evaluate the platelet function and thrombotic potential in patients following recovery from SARS-CoV-2 with clear symptoms of PASC.

**Methods:** PASC patients and matched healthy controls were enrolled in the study on average 15 months after documented SARS-CoV-2 infection. Platelet activation was evaluated by Light Transmission Aggregometry (LTA) and flow cytometry in response to platelet surface receptor agonists. Thrombosis in platelet-deplete plasma was evaluated by Factor Xa activity. A microfluidics system assessed thrombosis in whole blood under shear stress conditions.

**Results:** A mild increase in platelet aggregation in PASC patients through the thromboxane receptor was observed and platelet activation through the glycoprotein VI (GPVI) receptor was decreased in PASC patients compared to age- and sex-matched healthy controls. Thrombosis under shear conditions as well as Factor Xa activity were reduced in PASC patients. Plasma from PASC patients was an extremely potent activator of washed, healthy platelets – a phenomenon not observed when stimulating healthy platelets after incubation with plasma from healthy individuals.

**Conclusions:** PASC patients show dysregulated responses in platelets and coagulation in plasma, likely caused by a circulating molecule that promotes thrombosis. A hitherto

undescribed protective response appears to exists in PASC patients to counterbalance

ongoing thrombosis that is common to SARS-CoV-2 infection.

**Introduction:**

COVID-19, caused by the severe acute respiratory syndrome coronavirus 2

(SARS-CoV-2), may have residual effects following recovery. One third of infected

patients are asymptomatic, while others experience pulmonary and cardiovascular

symptoms leading to morbidity and mortality (1). The risk of both severe and persistent

symptoms of SARS-CoV-2 is higher males, in the advanced-age patient population, in

patients requiring immunosuppression, or with chronic kidney disease (2).  The risk of

severe and persistent symptoms of SARS-CoV-2 remains despite vaccination (3, 4).

Post-acute sequelae of COVID-19 (PASC) is an recognized multi-systemic

condition characterized by persistent symptoms four weeks beyond the initial SARS-CoV-

2 infection.  While acute respiratory distress syndrome (ARDS) is a feared consequence

of severe COVID-19, many acutely ill patients are reported to have thromboembolic

complications including deep vein thrombosis (DVT) and pulmonary embolism (PE). The

incidence of venous thromboembolic (VTE) complications may be as high as 40% (5, 6).

Observational data suggests that PASC patients develop or have persistent

symptoms long after recovery that were initially labeled Long COVID by the Centers for

Disease Control (CDC) and the National Institutes of Health (NIH). The National Institute

for Health and Care Excellence (NICE) now defines PASC as post-infection symptoms

beyond 4-12 weeks (7-9).  Patients blighted by the specter of PASC show structural and

functional organ derangements involving the respiratory, cardiovascular, neurological,

genitourinary, gastrointestinal, and musculoskeletal systems.  PASC symptoms typically

include fatigue, dyspnea, chest pain/tightness, headache, cough, difficulty concentrating,

and altered sensorium.  Poor memory and concentration are the most frequent symptoms

reported, closely followed by cognitive impairment, sleep disturbance, and post-traumatic

stress disorder (PTSD) (9-13).

Elevated  pro-thrombotic  markers  including  D-dimer,  C-reactive  protein,  and

neutrophil extracellular trap (NET) formation have been detected in acute SARS-CoV-2

infection.  In addition, many PASC patients complain of dyspnea on exertion, which leads

to physical inactivity and may further precipitate thromboembolic complications. Virus-

induced endothelial injury of severe COVID-19 patients initiates a state of dysregulated

coagulation.  Increased blood von Willebrand Factor (VWF), likely from both endothelial

cell and platelet activation, promotes hypercoagulability and thrombosis in patients with

COVID-19 (7, 8, 14).

Platelets have an important hemostatic function to prevent excessive bleeding

(hemostasis) alongside an important role in the host immune response to invading viral

and microbial infections, especially SARS-CoV-2 (15).  In a large observational study, no

additional risk for VTE after hospitalization with acute SARS-CoV-2 infection was noted

(16).  Contrasting this, the MICHELLE study revealed a protective effect of an oral Factor

Xa inhibitor on discharged patients by protecting them against DVT if they were previously

hospitalized,  even  without  an  initial  thrombotic  event  (17).  These  findings  raise  the

possibility that following SARS-CoV-2 recovery, patients may have ongoing functional

impairment in the coagulation cascade or abnormal platelet function. The Aim of this study

was to evaluate both platelet function and thrombotic potential in whole blood, in platelet-

deplete plasma, and in isolated platelets in patients following SARS-CoV-2 but with clear

PASC symptoms.

**Methods**

**Human Subjects:** 26 patients with Post-Acute Sequelae of SARS-CoV-2 infection (PASC) were enrolled based on a confirmed Covid-19 diagnosis and with persistent symptoms, including shortness of breath, fatigue, brain fog, dyspnea on exertion, mood disturbance, weakness, and memory impairment. PASC patients were compared to 19 matched healthy controls. Patients and healthy subjects were recruited only after informed consent by a research coordinator not involved in clinical care. All study procedures were conducted in accordance with the Declaration of Helsinki and approved by a local Institutional Review Board (IRB) protocol. Human platelet physiological studies must be conducted within 90 minutes in order for meaningful data to be obtained. These studies are labor-intensive and, in their entirety, cannot be conducted simultaneously for every patient within 90 minutes. The volume of blood required to conduct studies in whole blood, in platelet-rich plasma (PRP), in washed platelets, and in platelet deplete plasma is also substantial and exceeds the blood volume permitted to be drawn in the IRB protocol. Therefore, it is impossible to conduct every experiment in every enrolled PASC patient or healthy individual. Rather, the studies were conducted in a manner that was practically required as patients as PASC were identified.

**Cross-activation study using Flow-cytometry:**

To check whether plasma from PASC patients would modify platelet activation from healthy donors, we performed cross-activation tests. Platelet depleted plasma (20µl) from the PACS patients (n=18) as well as age- and sex-matched healthy controls (n=20) was added to the isolated washed platelets (80µl) from healthy controls (each performed in

triplicate) and incubated for 20 minutes at room temperature.  Platelets were stimulated

with agonists: Thrombin Receptor Activator Peptide 6 (TRAP6; 5µM), Thromboxane $A_2$

($TXA_2$) analogue U46619 (1µM), Adenosine diphosphate (ADP; 0.1µM), or Collagen

Related Peptide (CRP; 0.1µg/mL) for 15 minutes.  Samples were next stained with

phycoerythrin (PE)-conjugated anti-CD62P (p-selectin) for 30 minutes at room

temperature in darkness to prevent fluorophore bleaching before fixing with 4%

paraformaldehyde.  Fixed samples were read by flow cytometry (BD Accuri C6 Plus) to

acquire 20000 events and data was analyzed using FlowJo software.

**Antibody Profiling:**

Antibody profiling was performed by CDI labs using the HuProt 4.0 array. Briefly,

sequence-confirmed plasmids, which are used to make GST-purified recombinant

proteins in yeast.  After purification, GST-tagged proteins are piezoelectrically printed on

glass slides in duplicate, along with control proteins (GST, BSA, Histones, IgG, etc.). The

slides are barcoded for tracking and archiving. Each microarray batch is routinely

evaluated by anti-GST staining to demonstrate quality of expression and printing. The

new HuProt v4.0 consists of unique human proteins, isoform variants, and protein

fragments of the 19,613 canonical human proteins described in the Human Protein Atlas

(18-21).  We focused on protein exclusively expressed in platelets, highly expressed in

platelets, blood proteins known to be involved in the alterations in coagulation as noted.

Content includes major functional classes such as intracellular proteins, membrane

proteins, enzymes, secreted proteins, transcription factors, transporters, GPCRs,

cytokines, immune receptors, immune checkpoints, CD markers, ion channels, cytosolic

proteins, nuclear receptors. To measure antibody binding, slides are overlaid with a 1:100

dilution of serum, and stained with anti-IgG (red) and anti-IgM (green) secondary

antibodies. A heat map was generated to show each antibody concentration expressed

as standard deviations (SD) above or below the mean concentration found in healthy

individuals.

**Light Transmission Aggregometry (LTA):** Blood from healthy controls and PASC

patients was collected in sodium-citrate tubes after obtaining informed consent. Within 30

minutes to 1 hour of collection, anticoagulated whole blood was centrifuged at 200g for

15 minutes at room temperature (RT) to obtain platelet rich plasma (PRP). PRP was

collected in 15mL falcon tubes and Platelet poor plasma (PPP) was obtained from the

remaining sample by re-centrifugation at 2500 g for 10 minutes. PPP was used to initiate

baseline optical density. Platelet Aggregation was tested following stimulation with

agonists: ADP (1µM), TRAP-6 (5µM), U46619 (5µM) and CRP (1µg/mL). Optical density

changes were recorded after the addition of the agonists on Aggregometer (CHRONO-

LOG 700) for 6 minutes while stirring the sample at 1200 rpm at 37°C.

**Alpha granule exocytosis with Flow-cytometry:** Platelets were isolated from whole

blood collected from healthy controls and PASC patients in sodium-citrate tubes. Briefly,

whole blood was centrifuged at 200g for 15 minutes at RT and then PRP was collected

in the 15mL falcon tubes. Collected PRP was centrifuged again at 200g for 10 minutes to

remove remaining WBCs and RBCs. Platelets were washed using Tyrode's buffer in the

1:1 ratio with the addition of Prostaglandin I2 (PGI2; 10nM). Washed platelets were re-

suspended in 10mL of Tyrode's buffer. Washed platelets from PASC patients and healthy

controls were stimulated using different concentrations of agonist: TRAP-6 (1, 5, 10 and

20µM), U46619 (0.1, 0.5, 1.0, 10µM), ADP (0.01, 0.1, 1.0, 10µM) and CRP (0.05, 0.1,

0.5, 1.0µg/mL) for 15 minutes and then incubated with phycoerythrin (PE)-conjugated

anti-CD62P for 30 minutes at RT. Samples were fixed with 4% paraformaldehyde. P-

Selectin expressed on the surface on the platelets signifies an activated state. Samples

were read by flow cytometry (BD Accuri C6 Plus) to acquire 20000 events. Data were

analyzed using FloJo software.

**Total Thrombus formation Analysis System (T-TAS01)**: The T-TAS01 microfluidic

chamber system was used and the results were compared to healthy controls. T-TAS01

is a microchip flow-chamber system that measures thrombus formation under blood flow

conditions over time. Whole blood collected in the sodium-citrate tube was used for the

test. Briefly, before measurement 20µL of the CaCTI reagent (Corn trypsin inhibitor) was

added to 480µL of whole blood and then this blood was mixed gently with the pipette.

Then 450µL of mixed blood was loaded into the reservoir of the AR chip reservoir coated

with both collagen and thromboplastin (tissue factor) with a shear rate of 600 s$^{-1}$. As

coagulation and platelet activation progress, thrombosis is noted as pressure rises in the

channel following shear rate-mediated activation of blood in the microchip channel. These

pressure changes are monitored by a pressure sensor upstream of the chamber and are

defined as The Occlusion Start Time (OST), and the Occlusion Time (OT). The difference

between OT and OST was calculated for each subject. Furthermore, to check the pro-

thrombotic activity of the plasma of the PASC patients, 200µL plasma from the PASC

patients and healthy controls was incubated with 800µL whole blood from healthy control

for 30 minutes. Then this spiked whole blood with plasma was loaded into the reservoir

of the AR chip.

**Plasma Factor Xa activity:** Factor Xa (FXa) is the activated form of the coagulation factor

X. Factor X, a serine endopeptidase plays an important role in the coagulation pathway.

It converts prothrombin into active thrombin by complexing with activated co-factor V in

the prothrombinase complex. The Factor Xa Activity assay kit from Abcam (ab204711)

was used to measure the Factor Xa activity in the plasma of the PASC patients compared

to age- and sex-matched healthy controls. The assay was performed according to the

manufacturer's recommendations.

**Statistical Analysis:** All data were interrogated for normalcy using the Shapiro-Wilks

test. Normally distributed continuous variable group differences were assessed by the

student 2-tailed Student's t-test and skewed data assessed by the Mann-Whitney $U$ test.

Analyses were conducted using GraphPad Prism 7 (GraphPad Software). For Gaussian-

distributed data in 3 or more groups, 1-way ANOVA then the Bonferroni multiple

comparisons test was used, otherwise the Kruskal–Wallis test followed by Dunn post-test

was used. Significance was accepted as a $P$ value of less than 0.05.

**Results**

*Baseline Study Population:* 26 patients with PASC (mean age 49.9) were recruited from the pulmonary medicine clinic in Northeast Ohio and compared with 19 healthy volunteers (mean age 41.4). Healthy recruits or PASC patients were comparable, showing no significant difference between groups with respect to sex, BMI, and medical co-morbidities of PASC patients who were more commonly smokers (**Table 1)**. Patients with PASC presented on average 501 days $\pm$ 69 days after the last detection of a SARS-CoV-2 amplicon by polymerase chain reaction (PCR). Of the 26 common PASC symptoms in our patient population, most described fatigue (76.9%), brain fog (76.9%), shortness of breath at rest (69.2%), and exertional dyspnea (69.2%) (**Supplemental Table 1**).

*Platelet Reactivity in PASC patients*: When platelet-deplete plasma from healthy individuals or PASC patients was added to healthy, twice washed platelets, there was a surprisingly profound activation of platelets following stimulation with all platelet receptor agonists tested caused by PASC patient plasma incubation (**Fig. 1**). Given that immunothrombosis is common in COVID-19 and the complement system likely contributes to these responses (22-24), we heated and inactivated plasma from PASC donors at 55°C and repeated the experiments. We observed that platelets were persistently activated if spiked with room temperature plasma or heat-inactivated plasma prior to agonist stimulation from PASC patients suggesting that complement activation is unlikely to be a driver in the augmented platelet response observed (**Supplemental Fig. S1**).

Platelet aggregation in PRP was evaluated by LTA after stimulating platelets with surface

receptor agonists for the P2Y$_{12}$ receptor (ADP), the thromboxane receptor (U46619), the

GPVI receptor (CRP), and PAR1 (TRAP-6).  Overall, platelet aggregation by LTA in PASC

was similar to healthy individuals except through the thromboxane receptor which

demonstrated increased reactivity following stimulation with the agonist U46619 (87%±3

for PASC vs. 72%±4 for healthy subjects, p=0.03) (**Fig. 2 A-D**). To distinguish platelet

function that may be altered by mediators in PRP, these studies were repeated in a dose-

response manner after isolating and washing platelets twice, then assessing platelet

reactivity by alpha granule exocytosis by appearance of P-selectin on the surface of the

platelet as reported previously (25, 26).  When washed platelets from healthy volunteers

and PASC patients were stimulated with platelet surface receptor agonists, we found

similar reactivity to healthy individuals through the following receptors: P2Y$_{12}$,

thromboxane, PAR 1 (**Fig. 3A-C**). Curiously, platelet reactivity through GPVI was

markedly decreased in PASC patients compared with healthy controls (**Fig. 3D**).

To test the hypothesis that changes in platelet function as observed by alpha granule

exocytosis augmented in PASC following platelet thromboxane receptor stimulation, **Fig.**

**2B**) or platelet aggregation (decreased in PASC following platelet GPVI stimulation, **Fig.**

**3D**) are a consequence of changes in platelet surface receptor expression, we performed

flow cytometry using receptors tagged with a fluorophore, ultimately finding no difference

(**Supplemental Fig. S2**).

*Whole blood and platelet-independent coagulation in PASC patients*: Whole blood was drawn from patients who were healthy or with PASC, then passed through a 80 μm channel in a microfluidics system at 600 dynes (s$^{-1}$) shear stress.  This assay was specifically chosen because thrombosis in most patients with SARS-CoV-2 occurs in the venous vasculature, and the assay utilizes tissue factor and collagen to initiate thrombosis, which simulates endothelial damage common to SARS-CoV-2.  Overall, the time to pressure increase in the microchannel (kPa), which is a signature of *ex vivo* blood clotting as coagulated blood diminishes flow, was prolonged in PASC patients compared with healthy conditions (occlusion time-occlusion start time, 94±8 vs. 73±8 seconds, respectively, P=0.02) (**Fig. 4 A-B**). To test whether a mediator in plasma is responsible for the difference in coagulation observed in PASC, whole blood from a healthy individual was spiked with platelet-deplete plasma from a PASC patient was compared to spiking platelet-deplete plasma from a healthy individual.  This showed no difference between occlusion time and occlusion start time (62±4 healthy plasma vs. 66±8 PASC plasma, n=5 in each group, P=0.74).  We next assessed Factor Xa activity in platelet-deplete plasma which represents the point of convergence between the extrinsic and the intrinsic arms of the coagulation cascade. We observed Factor Xa activity was reduced in PASC vs. healthy individuals (23.6±5 vs. 50±8 ng/100μL, respectively; P=0.037) (**Fig. 4C**). While complement, which can increase platelet reactivity, is well-known to be denatured at 55˚C, antibodies targeted against platelet proteins or the coagulation cascade may retain some function (27-29).  We therefore performed a limited antibody array using serum from patients with PASC comparing this to healthy individuals for antibodies targeted against platelet proteins that could alter platelet function of antibodies targeted against circulating

Factor II, Factor II Receptor, or Factor X. We did not find a significant difference in either

IgM or IgG against those protein targets comparing PASC patients to healthy controls

(**Supplemental Fig. S3**).

These experiments suggest there is a change in PASC plasma with dysregulated

coagulation compared with healthy conditions that does not involve complement or

circulating antibodies. Overall, the observation that washed platelets from PASC patients

are relatively unchanged or less reactive than healthy platelets, and that platelet-deplete

plasma in PASC is less likely to clot, suggests a potential adaptive response in the host

long after COVID-19 infection, possibly to prevent thrombosis that is common in patients

with acute COVID-19 infection.

**Discussion**

SARS-CoV-2 is well-known to precipitate thrombotic events, especially DVT and PE (30, 31). Acute SARS-CoV-2 infection has also been shown to coincide with activated platelets by multiple independent groups (32-35). Our primary objective was to assess platelet function and thrombosis in patients with PASC given that rheologic manifestations of this syndrome are an under-developed area of investigation. Our major finding in patients more than one year following acute SARS-CoV-2 and with clear symptoms of PASC is dysregulated platelet function, with slight platelet activation through the thromboxane receptor, and attenuated activation through the GPVI receptor. We did not observe a change in platelet receptor signaling through PAR1 or $P2Y_{12}$ in PASC using two well-validated techniques to interrogate platelet reactivity.

This study may be the first to report a functional defect in the coagulation cascade given our finding of reduced activated Factor Xa in platelet-deplete plasma in patients with PASC (33). Curiously, using radioligand binding technology or a similar chromogenic substrate for Factor Xa activation as we employ, others have reported that platelets and platelet-derived microparticles may be a source of Factor Xa activity in the circulation (36-38). Moreover, a putative signaling mechanism between the platelet GPVI receptor and Factor Xa was previously reported (39). In addition, the profoundly under-reactive GPVI receptor identified by flow cytometry for alpha granule exocytosis using CRP as the agonist in our study complements the observation of prolonged time to thrombosis in human blood *ex vivo* under conditions of venous shear stress since collagen is present in the microfluidics channel which activates the GPVI receptor.

The most significant finding in our study is that platelet-deplete plasma in PASC

patients 'primes' and markedly activates healthy, washed platelets through every receptor

signaling pathway evaluated.  We did not see this when platelet-deplete plasma from

healthy controls was incubated with washed platelets before agonist stimulation.  The

hyper-reactive platelet effect caused by PASC plasma persisted even after heating

plasma to 55°C suggesting the complement cascade or circulating antibodies are likely

not responsible for augmented platelet activation.  Using an antibody array, we could not

find antibodies directed against platelet protein targets of coagulation proteins in the blood

that could explain the observation in patients with PASC.  Therefore, we offer the

suggestion that a putative pro-thrombotic factor exists in the blood of patients with PASC,

and its presence leads to an adaptive response in the host in which platelet reactivity and

thrombosis are paradoxically tipped in the opposite direction, presumably as a protective

response from thrombosis.

Our study in PASC patients differs from a recent letter by Martins-Gonçalves *et al.*

in patients reporting at least one persistent respiratory symptom after acute

hospitalization for COVID-19 and with persistent platelet hyperreactivity (40). Important

differences in the enrolled subjects were noted. Firstly, while the average age of patients

enrolled by Martins-Gonçalves *et al.* was similar to our cohort, the represented majority

were men while 73% of our patients and 79% of our healthy controls were women.

Platelet reactivity has been shown to differ between men and women in health and in

cardiovascular disorders (26, 36, 41).  Secondly, the control group employed by Martins-

Gonçalves *et al*. tested negative for SARS-CoV-2 weekly for almost two years prior to

enrollment, while several of our control comparators reported SARS-CoV-2 infection in

the preceding two years but without residual symptoms at the time of enrollment and this

was the reason why they were selected as controls.  Lastly, Martins-Gonçalves evaluated

platelet reactivity more in the convalescent setting 1-4 months after acute SARS-CoV-2

infection in 2021 whereas our PASC patients on average were evaluated >16 months

after the last documented acute SARS-CoV-2 infection in 2023. Clinical outcomes and

symptoms following SARS-Cov-2 infection appear to vary depending on vaccination

status and the viral strain of the infected individual (42). This means that interpretation of

data gained from patients evaluated during different waves of the COVID-19 pandemic

require careful attention. Given that SARS-CoV-2 was demonstrated to induce

morphological changes in platelets and promote apoptosis after viral internalization in an

ACE2-dendent and ACE2-independent receptor internalization pathway (15), platelet

reprogramming in patients with PASC may occur.

**Conclusion**

In summary, patients who have long recovered from acute SARS-CoV-2 but with

persistent symptoms under the constellation of PASC show evidence of dysregulated

platelet reactivity, alterations in normal coagulation of platelet-deplete plasma, and

diminished clotting in whole blood *ex vivo* under shear stress conditions. These events

are likely a consequence of adapting to a platelet-activating factor in plasma of patients

with PASC.  The stimulus behind the exhausted platelet phenomenon, particularly

through GPVI, in PASC patients, should be investigated and identification of the putative

pro-thrombotic mediator should be prioritized given the implications for patients with

PASC being considered as donors of blood and plasma products in future.

**Acknowledgements and Sources of Funding**

We are grateful for funding from NHLBI HL128856 to SJC and HL158669 to TMM. Dr. Khorana acknowledges research support from the Sondra and Stephen Hardis Chair in Oncology Research. The research was, in part, funded by the National Institutes of Health (NIH) Agreement 1OT2HL156812 through the National Heart, Lung, and Blood Institute (NHLBI) CONNECTS program. The views and conclusions contained in this document are those of the authors and should not be interpreted as representing the official policies, either expressed or implied, of the NIH.

**References**

1.      Oran DP, Topol EJ. The Proportion of SARS-CoV-2 Infections That Are Asymptomatic : A Systematic Review. Ann Intern Med. 2021;174(5):655-62.
2.      Peckham H, de Gruijter NM, Raine C, Radziszewska A, Ciurtin C, Wedderburn LR, et al. Male sex identified by global COVID-19 meta-analysis as a risk factor for death and ITU admission. Nat Commun. 2020;11(1):6317.
3.      Agrawal U, Bedston S, McCowan C, Oke J, Patterson L, Robertson C, et al. Severe COVID-19 outcomes after full vaccination of primary schedule and initial boosters: pooled analysis of national prospective cohort studies of 30 million individuals in England, Northern Ireland, Scotland, and Wales. Lancet. 2022;400(10360):1305-20.
4.      Lee SW, Lee J, Moon SY, Jin HY, Yang JM, Ogino S, et al. Physical activity and the risk of SARS-CoV-2 infection, severe COVID-19 illness and COVID-19 related mortality in South Korea: a nationwide cohort study. Br J Sports Med. 2022;56(16):901-12.
5.      Helms J, Tacquard C, Severac F, Leonard-Lorant I, Ohana M, Delabranche X, et al. High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. Intensive Care Med. 2020;46(6):1089-98.
6.      Klok FA, Kruip M, van der Meer NJM, Arbous MS, Gommers D, Kant KM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. Thromb Res. 2020;191:145-7.
7.      Libby P, Luscher T. COVID-19 is, in the end, an endothelial disease. Eur Heart J. 2020;41(32):3038-44.
8.      Ranucci M, Ballotta A, Di Dedda U, Baryshnikova E, Dei Poli M, Resta M, et al. The procoagulant pattern of patients with COVID-19 acute respiratory distress syndrome. J Thromb Haemost. 2020;18(7):1747-51.
9.      Crook H, Raza S, Nowell J, Young M, Edison P. Long covid-mechanisms, risk factors, and management. BMJ. 2021;374:n1648.
10.     Carfi A, Bernabei R, Landi F, Gemelli Against C-P-ACSG. Persistent Symptoms in Patients After Acute COVID-19. JAMA. 2020;324(6):603-5.
11.     Goertz YMJ, Van Herck M, Delbressine JM, Vaes AW, Meys R, Machado FVC, et al. Persistent symptoms 3 months after a SARS-CoV-2 infection: the post-COVID-19 syndrome? ERJ Open Res. 2020;6(4).
12.     Xiong Q, Xu M, Li J, Liu Y, Zhang J, Xu Y, et al. Clinical sequelae of COVID-19 survivors in Wuhan, China: a single-centre longitudinal study. Clin Microbiol Infect. 2021;27(1):89-95.
13.     Teuwen LA, Geldhof V, Pasut A, Carmeliet P. COVID-19: the vasculature unleashed. Nat Rev Immunol. 2020;20(7):389-91.
14.     Panigada M, Bottino N, Tagliabue P, Grasselli G, Novembrino C, Chantarangkul V, et al. Hypercoagulability of COVID-19 patients in intensive care unit: A report of thromboelastography findings and other parameters of hemostasis. J Thromb Haemost. 2020;18(7):1738-42.
15.     Koupenova M, Corkrey HA, Vitseva O, Tanriverdi K, Somasundaran M, Liu P, et al. SARS-CoV-2 Initiates Programmed Cell Death in Platelets. Circ Res. 2021;129(6):631-46.

16.　　Patel L, Stenzel A, Van Hove C, Sidebottom A, Kethireddy R, Ha N, et al.
Outcomes in patients discharged with extended venous thromboembolism prophylaxis
after hospitalization with COVID-19. Vasc Med. 2023:1358863X231159945.
17.　　Ramacciotti E, Barile Agati L, Calderaro D, Aguiar VCR, Spyropoulos AC, de
Oliveira CCC, et al. Rivaroxaban versus no anticoagulation for post-discharge
thromboprophylaxis after hospitalisation for COVID-19 (MICHELLE): an open-label,
multicentre, randomised, controlled trial. Lancet. 2022;399(10319):50-9.
18.　　Zhuang L, Huang C, Ning Z, Yang L, Zou W, Wang P, et al. Circulating tumor-
associated autoantibodies as novel diagnostic biomarkers in pancreatic
adenocarcinoma. Int J Cancer. 2023;152(5):1013-24.
19.　　Yang Q, Ye H, Sun G, Wang K, Dai L, Qiu C, et al. Human Proteome Microarray
identifies autoantibodies to tumor-associated antigens as serological biomarkers for the
diagnosis of hepatocellular carcinoma. Mol Oncol. 2023;17(5):887-900.
20.　　Xie X, Tian L, Zhao Y, Liu F, Dai S, Gu X, et al. BACH1-induced ferroptosis
drives lymphatic metastasis by repressing the biosynthesis of monounsaturated fatty
acids. Cell Death Dis. 2023;14(1):48.
21.　　Wang Y, Li J, Zhang X, Liu M, Ji L, Yang T, et al. Autoantibody signatures
discovered by HuProt protein microarray to enhance the diagnosis of lung cancer. Clin
Immunol. 2023;246:109206.
22.　　Aiello S, Gastoldi S, Galbusera M, Ruggenenti P, Portalupi V, Rota S, et al. C5a
and C5aR1 are key drivers of microvascular platelet aggregation in clinical entities
spanning from aHUS to COVID-19. Blood Adv. 2022;6(3):866-81.
23.　　Charitos P, Heijnen I, Egli A, Bassetti S, Trendelenburg M, Osthoff M. Functional
Activity of the Complement System in Hospitalized COVID-19 Patients: A Prospective
Cohort Study. Front Immunol. 2021;12:765330.
24.　　Rajamanickam A, Nathella PK, Venkataraman A, Dasan B, Putlibai S, Ahamed
SF, et al. Levels of Complement Components in Children With Acute COVID-19 or
Multisystem Inflammatory Syndrome. JAMA Netw Open. 2023;6(3):e231713.
25.　　Cameron SJ, Mix DS, Ture SK, Schmidt RA, Mohan A, Pariser D, et al. Hypoxia
and Ischemia Promote a Maladaptive Platelet Phenotype. Arterioscler Thromb Vasc
Biol. 2018;38(7):1594-606.
26.　　Soo Kim B, Auerbach DS, Sadhra H, Godwin M, Bhandari R, Ling FS, et al. Sex-
Specific Platelet Activation Through Protease-Activated Receptors Reverses in
Myocardial Infarction. Arterioscler Thromb Vasc Biol. 2021;41(1):390-400.
27.　　Gordon J, Turner GC. The intensifying effect of glucose on the protective action
of glycine against the heat-inactivation of complement. J Hyg (Lond). 1955;53(3):335-8.
28.　　Lai HM, Tang Y, Lau ZYH, Campbell RAA, Yau JCN, Chan CCY, et al. Antibody
stabilization for thermally accelerated deep immunostaining. Nat Methods.
2022;19(9):1137-46.
29.　　Jones FS. The Effect of Heat on Antibodies. J Exp Med. 1927;46(2):291-301.
30.　　Elbadawi A, Elgendy IY, Joseph D, Eze-Nliam C, Rampersad P, Ouma G, et al.
Racial Differences and In-Hospital Outcomes Among Hospitalized Patients with COVID-
19. J Racial Ethn Health Disparities. 2022;9(5):2011-8.
31.　　Elbadawi A, Elgendy IY, Sahai A, Bhandari R, McCarthy M, Gomes M, et al.
Incidence and Outcomes of Thrombotic Events in Symptomatic Patients With COVID-
19. Arterioscler Thromb Vasc Biol. 2021;41(1):545-7.

32.    Manne BK, Denorme F, Middleton EA, Portier I, Rowley JW, Stubben C, et al.
Platelet gene expression and function in patients with COVID-19. Blood.
2020;136(11):1317-29.
33.    Puhm F, Allaeys I, Lacasse E, Dubuc I, Galipeau Y, Zaid Y, et al. Platelet
activation by SARS-CoV-2 implicates the release of active tissue factor by infected
cells. Blood Adv. 2022;6(12):3593-605.
34.    Zaid Y, Puhm F, Allaeys I, Naya A, Oudghiri M, Khalki L, et al. Platelets Can
Associate with SARS-Cov-2 RNA and Are Hyperactivated in COVID-19. Circ Res.
2020;127(11):1404-18.
35.    Petito E, Franco L, Falcinelli E, Guglielmini G, Conti C, Vaudo G, et al. COVID-19
infection-associated platelet and neutrophil activation is blunted by previous anti-SARS-
CoV-2 vaccination. Br J Haematol. 2023;201(5):851-6.
36.    Godwin MD, Aggarwal A, Hilt Z, Shah S, Gorski J, Cameron SJ. Sex-Dependent
Effect of Platelet Nitric Oxide: Production and Platelet Reactivity in Healthy Individuals.
JACC Basic Transl Sci. 2022;7(1):14-25.
37.    Miletich JP, Jackson CM, Majerus PW. Interaction of coagulation factor Xa with
human platelets. Proc Natl Acad Sci U S A. 1977;74(9):4033-6.
38.    Walsh PN, Biggs R. The role of platelets in intrinsic factor-Xa formation. Br J
Haematol. 1972;22(6):743-60.
39.    Al-Tamimi M, Grigoriadis G, Tran H, Paul E, Servadei P, Berndt MC, et al.
Coagulation-induced shedding of platelet glycoprotein VI mediated by factor Xa. Blood.
2011;117(14):3912-20.
40.    Martins-Goncalves R, Campos MM, Palhinha L, Azevedo-Quintanilha IG, Abud
Mendes M, Ramos Temerozo J, et al. Persisting Platelet Activation and Hyperactivity in
COVID-19 Survivors. Circ Res. 2022;131(11):944-7.
41.    Faraday N, Goldschmidt-Clermont PJ, Bray PF. Gender differences in platelet
GPIIb-IIIa activation. Thromb Haemost. 1997;77(4):748-54.
42.    Robinson ML, Morris CP, Betz JF, Zhang Y, Bollinger R, Wang N, et al. Impact of
Severe Acute Respiratory Syndrome Coronavirus 2 Variants on Inpatient Clinical
Outcome. Clin Infect Dis. 2023;76(9):1539-49.

**Tables and Figures:**

**Table 1. Baseline Characteristics.** Patients with PASC (n=26, occurring at a mean duration of 501 ± 69 days after a diagnosis of COVID-19) were recruited for the study and compared with healthy volunteers (n=19). Data are reported as mean ± SEM. Differences between groups was calculated by the student's t-test for continuous variables or Chi-Squared for dichotomous variables. BMI=Body Mass Index. CVD=Cardiovascular Disease. BP=Blood Pressure. NSAID=non-steroidal anti-inflammatory medication.

**Fig. 1. Activation of Platelets by PASC but not Healthy Plasma.** Isolated, washed platelets were pre-incubated for 30 minutes with 20 µL of age- and sex-matched healthy plasma or PASC plasma, then stimulated with TRAP-6, 5 µM (PAR1), U46619, 1 µM (Thromboxane receptor), ADP, 0.1 µM (P2Y12 Receptor), or Collagen-Related Peptide, CRP, 0.1 µg/mL (GPVI Receptor) for 15 minutes. Platelet reactivity was assessed by FACS by surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by the Mann-Whitney U test, n=20 plasma isolated for healthy individuals and n=18 plasma isolates for PASC patients. Washed platelets from a healthy 25 year-old male.

**Fig. 2. Platelet Activation in PASC by Aggregometry**. **A-D:** Blood was drawn from healthy individual or PASC patients and PRP was isolated prior to stimulation with TRAP-6 (5µM), U466119 (5µM), ADP (1.0 µM), or CRP (1.0 µg/mL) and observed until platelet aggregation was noted. Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the Mann Whitney U test, n=5 for healthy individuals and n=5 for PASC or student's t-test, n=5 for healthy individuals and n=5 for PASC.

**Fig. 3. Platelet Activation in PASC by Flow Cytometry. A-D**: Blood was drawn from healthy individual or PASC patients and isolated, washed platelets were stimulated with TRAP-6 (0-20µM), U466119 (0-10 µM), ADP (0-10 µM), or CRP (0-1.0 µg/mL) for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-Selectin) using a tagged antibody. Data are represented as mean fold-change ± SEM. Group differences were assessed by the Kruskal-Wallis test, n=10 for healthy individuals and n=9 for PASC. NS=not significant.

**Fig. 4. Whole Blood and Plasma Coagulation in PASC by Microfluidics and Factor Xa Activity.** **A.** Blood was isolated and passed through a microfluidics chip using the Total Thrombus formation Analysis System (T-TAS01) at roughly venous shear stress (600 s-1). With clot generation, pressure in the microfluidics chamber (kPa) gradually increases until cessation of blood flow. Data are represented as mean ± SEM. **B.** The time from the start of occlusion to cessation of blood was calculated in seconds as mean ± SEM. Group differences were determined by the student's t-test for PASC patients (n=10) or or healthy volunteers (n=12). **C.** Citrate plasma devoid of platelets was evaluated for Factor Xa activity using a chromogenic substrate in PASC patients (n=7) or healthy volunteers (n=10). Data are represented as mean ± SEM. Group

598   differences were assessed by the student's t-test, n=12 for healthy individuals and n=10
599   for PASC.
600
601
602   **Supplementary Tables and Figures:**
603
604   **Table S1. Persistent symptoms of enrolled PASC patients.**
605   Data are expressed as frequency and % of population and mean ± SEM.
606
**Fig. S1. PASC Symptoms of Enrolled Subjects.** Platelet-deplete citrate plasma from
PASC
patients already demonstrated to activate healthy male platelets divided in half and then
either heat-treated (55 °C, 15 minutes) or not and incubated for 30 minutes prior to
stimulation with a surface receptor agonist TRAP-6, 5 µM (PAR1), U46619, 1 µM
(Thromboxane receptor), ADP, 0.1 µM (P2Y$_{12}$ Receptor), or Collagen-Related Peptide,
CRP, 5 µg/mL (GPVI Receptor). Platelet reactivity was then assessed by FACS by
surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ±
SEM. Group differences were assessed by Mann-Whitney *U* test for n=10.
616
**Fig. S2. Plasma Antibody Analysis.**  Blood was isolated from patients with PASC
(n=66, blue) or matched controls (n=10, green) and antibody profiling was performed
using the HuProt 4.0 array of unique human proteins known to be exclusively expressed
in platelets: Platelet Factor 4 (PF4), Glycoprotein VI (GP6), integrin alpha2 gene
(ITGA2). Other proteins highly expressed in platelets were also evaluated: von
Willebrand Factor (vWF), Platelet And Endothelial Cell Adhesion Molecule 1 (PECAM1),
E-selectin (SELE), P-selectin (SELP), Vascular Cell Adhesion Molecule 1 (VCAM1).
Coagulation cascade protein or receptors were also evaluated: Factor II (F2), factor II
receptor (FIIR), Factor X (F10). A heat map was generated to show each antibody
concentration expressed as standard deviations (SD) above or below the mean
concentration found in healthy individuals.
628
629   **Fig. S3. Platelet Surface Receptor Expression in PASC Patients.**  Blood was drawn
630   from healthy individuals (n=6) or PASC patients (n=6) and isolated, washed platelets
631   were assessed by FACS for expression the following receptors tagged with fluorescent
632   antibodies: P2Y$_{12}$, PAR, Thromboxane, GPVI. Data are represented as ± SEM. Group
633   differences were assessed by t-test with p-values as noted.

644    **Table 1.** Demographics for PASC Patients and Healthy Controls

| Demographics | Healthy (n=19) | | PASC (n=26) | | P-Value |
|---|---|---|---|---|---|
| | N | % | N | % | |
| ***Characteristics*** | | | | | |
| Mean Age (± SEM) | 41.4 ± 2.89 | | 49.9 ± 3.61 | | 0.090 |
| 18-27 years | 4 | 21.0% | 3 | 11.5% | 0.30 |
| 28-40 years | 4 | 21.0 % | 7 | 26.9 % | 0.76 |
| 41-60 years | 9 | 47.3 % | 5 | 19.2 % | 0.51 |
| 61-82 years | 2 | 10.5 % | 11 | 42.3 % | 0.62 |
| Male | 4 | 21.0 % | 7 | 35.0 % | 0.73 |
| Female | 15 | 78.9 % | 19 | 73.0 % | |
| BMI (± SEM) | 29.5 ± 1.29 | | 30.9 ± 2.17 | | 0.61 |
| Smoker | 1 | 7.7 % | 13 | 50.0 % | *0.002* |
| ***Medical History*** | | | | | |
| Hypertension | 4 | 21.0 % | 6 | 23.0 % | 0.99 |
| Hyperlipidemia | 2 | 10.5 % | 5 | 19.2 % | 0.68 |
| History of CVD | 1 | 5.2 % | 3 | 11.5 % | 0.62 |
| Diabetes | 2 | 10.5 % | 1 | 3.8 % | 0.56 |
| History of Pulmonary Embolism | 0 | 0.0 % | 1 | 3.8 % | 0.99 |
| History of Thrombosis | 1 | 5.2 % | 0 | 0.0 % | 0.42 |
| ***Medications*** | | | | | |
| Platelet inhibitors | 1 | 5.2 % | 6 | 23.0 % | 0.21 |
| Aspirin | 1 | 5.2 % | 5 | 19.2% | |
| DAPT (Dual Antiplatelet Therapy) | | | 1 | 3.8% | |
| Anticoagulants | 0 | 0.0 % | 2 | 7.6 % | 0.50 |
| Warfarin | | | 1 | 3.8% | |
| Apixaban | | | 1 | 3.8% | |
| Statins | 2 | 15.4 % | 5 | 19.2 % | 0.68 |
| NSAIDs | 6 | 31.5 % | 4 | 15.38 % | 0.28 |
| BP Medications | 4 | 21.0 % | 1 | 3.8 % | 0.14 |

656    **Supplemental Table 1.** Symptoms Experienced by PASC Patients

| Symptoms | PASC (n=26) | |
| --- | --- | --- |
| | N | % |
| Mean Time After Infection (± SEM) | 501 days ± 69 | |
| Fatigue | 20 | 76.9 % |
| Brain Fog | 20 | 76.9 % |
| Shortness of Breath | 18 | 69.2 % |
| Dyspnea on Exertion | 18 | 69.2 % |
| Dizziness | 17 | 65.3 % |
| Changes in Mood | 11 | 55.0 % |
| Insomnia | 14 | 53.8 % |
| Headaches | 13 | 50.0 % |
| Palpitations | 13 | 50.0 % |
| Chest Pain | 11 | 42.3 % |
| Joint Pain | 9 | 34.6 % |
| Cough | 7 | 26.9 % |
| Parosmia/Dysgeusia | 6 | 23.0 % |
| Memory Deficits | 6 | 23.0 % |
| Depression | 6 | 23.0 % |
| Syncope | 6 | 23.0 % |
| Nausea | 5 | 19.2 % |
| Body Aches | 4 | 15.3% |
| Incontinence | 3 | 11.5 % |
| Neck Pain | 3 | 11.5 % |
| HF Symptoms | 2 | 7.69 % |
| Diarrhea | 1 | 3.8 % |
| Erectile Dysfunction | 1 | 3.8 % |
| Fever | 1 | 3.8 % |
| Orthopnea | 1 | 3.8 % |
| Limb Paresthesia | 1 | 3.8 % |

**Fig. 1. Activation of Platelets by PASC but not Healthy Plasma.** Isolated, washed platelets were pre-incubated for 30 minutes with 5 µL of age- and sex-matched healthy plasma or PASC plasma, then stimulated with TRAP-6, 5 µM (PAR1), U46619, 1 µM (Thromboxane receptor), ADP, 0.1 µM (P2Y$_{12}$ Receptor), or Collagen-Related Peptide, CRP, 5 µg/mL (GPVI Receptor) for 15 minutes. Platelet reactivity was assessed by FACS by surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by the Mann-Whitney *U* test, n=20 plasma isolated for healthy individuals and n=18 plasma isolates for PASC patients. Washed platelets from a healthy 25 year-old male.

**Fig. 2A. Platelet Activation in PASC by Aggregometry.** Blood was drawn from healthy individual or PASC patients and PRP was isolated prior to stimulation with TRAP-6, 5µM until platelet aggregation was noted. Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the Mann Whitney *U* test, n=5 for healthy individuals and n=5 for PASC.

**Fig. 2B. Platelet Activation in PASC by Aggregometry.** Blood was isolated from healthy individual or PASC patients and PRP was isolated prior to stimulation with U466119, 5μM until platelet aggregation was noted. Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the student's t-test, n=5 for healthy individuals and n=5 for PASC.

**Fig. 2C. Platelet Activation in PASC by Aggregometry.** Blood was isolated from healthy individual or PASC patients and PRP was isolated prior to stimulation with ADP, 1 µM until platelet aggregation was noted.  Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the Mann Whitney *U* test, n=5 for healthy individuals and n=5 for PASC.

**Fig. 2D. Platelet Activation in PASC by Aggregometry.** Blood was isolated from healthy individual or PASC patients and PRP was isolated prior to stimulation with CRP, 1 μg/mL until platelet aggregation was noted. Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the Mann Whitney *U* test, n=5 for healthy individuals and n=5 for PASC.

**Fig. 3A. Platelet Activation in PASC by Flow Cytometry.** Isolated, washed platelets were stimulated with TRAP-6 for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-Selectin) using a tagged antibody. Data are represented as mean fold-change ± SEM. Group differences were assessed by the Kruskal-Wallis test, n=10 for healthy individuals and n=9 for PASC. NS=not significant.

**Fig. 3B. Platelet Activation in PASC by Flow Cytometry.** Isolated, washed platelets were stimulated with U46619 for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean fold-change ± SEM. Group differences were assessed by ANOVA, followed by the Bonferroni correction, n=10 for healthy individuals and n=10 for PASC.

**Fig. 3C. Platelet Activation in PASC by Flow Cytometry.** Isolated, washed platelets were stimulated with ADP for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean fold-change ± SEM. Group differences were assessed by ANOVA, followed by the Bonferroni correction, n=10 for healthy individuals and n=9 for PASC.

**Fig. 3D. Platelet Activation in PASC by Flow Cytometry.** Isolated, washed platelets were stimulated with collagen-related peptide (CRP) for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean fold-change ± SEM. Group differences were assessed by ANOVA, followed by the Bonferroni correction, n=10 for healthy individuals and n=9 for PASC.

**Fig. 4. Whole Blood and Plasma Coagulation in PASC by Microfluidics and Factor Xa Activity.** **A.** Blood was isolated and passed through a microfluidics chip using the Total Thrombus formation Analysis System (T-TAS01) at roughly venous shear stress ($600\ s^{-1}$). With clot generation, pressure in the microfluidics chamber (kPa) gradually increases until cessation of blood flow. Data are represented as mean ± SEM. **B.** The time from the start of occlusion to cessation of blood was calculated in seconds as mean ± SEM. Group differences were determined by the student's t-test for PASC patients (n=10) or or healthy volunteers (n=12). **C.** Citrate plasma devoid of platelets was evaluated for Factor Xa activity using a chromogenic substrate in PASC patients (n=7) or healthy volunteers (n=10). Data are represented as mean ± SEM. Group differences were assessed by the student's t-test, n=12 for healthy individuals and n=10 for PASC.

# Supplemental Data

**Fig. S1. Activation of Platelets by PASC plasma persists after heat treatment.** Platelet-deplete citrate plasma from PASC patients already demonstrated to activate healthy male platelets divided in half and one half was heat-treated (55 ˚C, 15 minutes).  Heat-treated and non-heat-treated plasma was incubated with washed platelets for 30 minutes prior to stimulation with a surface receptor agonist TRAP-6, 5 µM (PAR1), U46619, 1 µM (Thromboxane receptor), ADP, 0.1 µM (P2Y$_{12}$ Receptor), or Collagen-Related Peptide, CRP, 5 µg/mL (GPVI Receptor). Platelet reactivity was then assessed by FACS by surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by Mann-Whitney *U* test for n=10 (room temperature citrate plasma) or n=10 (heat-inactivated citrate plasma.

**Fig. S2. Platelet Surface Receptor Expression in PASC Patients.** Blood was drawn from healthy individuals (n=6) or PASC patients (n=6) and isolated, washed platelets were assessed by FACS for expression the following receptors tagged with fluorescent antibodies: $P2Y_{12}$, PAR1, Thromboxane Receptor (TR), GPVI, or 0 (control Immunoglobulin). Data are represented as MFI (Mean Fluorescence Intensity) of surface receptor expression $\pm$ SEM. Group differences were assessed by t-test with no differences noted between groups, ns=not significant)

**Fig. S3. Plasma Antibody Analysis.** Blood was isolated from patients with PASC (n=66, blue) or matched controls (n=10, green) and antibody profiling was performed using the HuProt 4.0 array of unique human proteins known to be exclusively expressed in platelets: Platelet Factor 4 (PF4), Glycoprotein VI (GP6), integrin alpha2 gene (ITGA2). Other proteins highly expressed in platelets were also evaluated: von Willebrand Factor (vWF), Platelet And Endothelial Cell Adhesion Molecule 1 (PECAM1), E-selectin (SELE), P-selectin (SELP), Vascular Cell Adhesion Molecule 1 (VCAM1). Coagulation cascade protein or receptors were also evaluated: Factor II (F2), factor II receptor (FIIR), Factor X (F10). A heat map was generated to show each antibody concentration expressed as standard deviations (SD) above or below the mean concentration found in healthy individuals.

**Fig. 1. Activation of Platelets by PASC but not Healthy Plasma.** Isolated, washed platelets were pre-incubated for 30 minutes with 5 µL of age- and sex-matched healthy plasma or PASC plasma, then stimulated with TRAP-6, 5 µM (PAR1), U46619, 1 µM (Thromboxane receptor), ADP, 0.1 µM (P2Y$_{12}$ Receptor), or Collagen-Related Peptide, CRP, 5 µg/mL (GPVI Receptor) for 15 minutes. Platelet reactivity was assessed by FACS by surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by the Mann-Whitney $U$ test, n=20 plasma isolated for healthy individuals and n=18 plasma isolates for PASC patients. Washed platelets from a healthy 25 year-old male.

**Fig. 2A. Platelet Activation in PASC by Aggregometry.** Blood was drawn from healthy individual or PASC patients and PRP was isolated prior to stimulation with TRAP-6, 5µM until platelet aggregation was noted. Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the Mann Whitney *U* test, n=5 for healthy individuals and n=5 for PASC.

**Fig. 2B. Platelet Activation in PASC by Aggregometry.** Blood was isolated from healthy individual or PASC patients and PRP was isolated prior to stimulation with U466119, 5µM until platelet aggregation was noted. Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the student's t-test, n=5 for healthy individuals and n=5 for PASC.

**Fig. 2C. Platelet Activation in PASC by Aggregometry.** Blood was isolated from healthy individual or PASC patients and PRP was isolated prior to stimulation with ADP, 1 µM until platelet aggregation was noted. Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the Mann Whitney *U* test, n=5 for healthy individuals and n=5 for PASC.

**Fig. 2D. Platelet Activation in PASC by Aggregometry.** Blood was isolated from healthy individual or PASC patients and PRP was isolated prior to stimulation with CRP, 1 μg/mL until platelet aggregation was noted. Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the Mann Whitney *U* test, n=5 for healthy individuals and n=5 for PASC.

**Fig. 3A. Platelet Activation in PASC by Flow Cytometry.** Isolated, washed platelets were stimulated with TRAP-6 for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-Selectin) using a tagged antibody. Data are represented as mean fold-change ± SEM. Group differences were assessed by the Kruskal-Wallis test, n=10 for healthy individuals and n=9 for PASC. NS=not significant.

**Fig. 3B. Platelet Activation in PASC by Flow Cytometry.** Isolated, washed platelets were stimulated with U46619 for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean fold-change ± SEM. Group differences were assessed by ANOVA, followed by the Bonferroni correction, n=10 for healthy individuals and n=10 for PASC.

**Fig. 3C. Platelet Activation in PASC by Flow Cytometry.** Isolated, washed platelets were stimulated with ADP for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean fold-change ± SEM. Group differences were assessed by ANOVA, followed by the Bonferroni correction, n=10 for healthy individuals and n=9 for PASC.

**Fig. 3D. Platelet Activation in PASC by Flow Cytometry.** Isolated, washed platelets were stimulated with collagen-related peptide (CRP) for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean fold-change ± SEM. Group differences were assessed by ANOVA, followed by the Bonferroni correction, n=10 for healthy individuals and n=9 for PASC.

**Fig. 4. Whole Blood and Plasma Coagulation in PASC by Microfluidics and Factor Xa Activity. A.** Blood was isolated and passed through a microfluidics chip using the Total Thrombus formation Analysis System (T-TAS01) at roughly venous shear stress (600 s$^{-1}$). With clot generation, pressure in the microfluidics chamber (kPa) gradually increases until cessation of blood flow. Data are represented as mean ± SEM. **B.** The time from the start of occlusion to cessation of blood was calculated in seconds as mean ± SEM. Group differences were determined by the student's t-test for PASC patients (n=10) or or healthy volunteers (n=12). **C.** Citrate plasma devoid of platelets was evaluated for Factor Xa activity using a chromogenic substrate in PASC patients (n=7) or healthy volunteers (n=10). Data are represented as mean ± SEM. Group differences were assessed by the student's t-test, n=12 for healthy individuals and n=10 for PASC.

**Fig. S1. Activation of Platelets by PASC plasma persists after heat treatment.** Platelet-deplete citrate plasma from PASC patients already demonstrated to activate healthy male platelets divided in half and one half was heat-treated (55 °C, 15 minutes). Heat-treated and non-heat-treated plasma was incubated with washed platelets for 30 minutes prior to stimulation with a surface receptor agonist TRAP-6, 5 μM (PAR1), U46619, 1 μM (Thromboxane receptor), ADP, 0.1 μM (P2Y$_{12}$ Receptor), or Collagen-Related Peptide, CRP, 5 μg/mL (GPVI Receptor). Platelet reactivity was then assessed by FACS by surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by Mann-Whitney *U* test for n=10 (room temperature citrate plasma) or n=10 (heat-inactivated citrate plasma.

**Fig. S2. Platelet Surface Receptor Expression in PASC Patients.** Blood was drawn from healthy individuals (n=6) or PASC patients (n=6) and isolated, washed platelets were assessed by FACS for expression the following receptors tagged with fluorescent antibodies: $P2Y_{12}$, PAR1, Thromboxane Receptor (TR), GPVI, or 0 (control Immunoglobulin). Data are represented as MFI (Mean Fluorescence Intensity) of surface receptor expression $\pm$ SEM. Group differences were assessed by t-test with no differences noted between groups, ns=not significant)

**Fig. S3. Plasma Antibody Analysis.** Blood was isolated from patients with PASC (n=66, blue) or matched controls (n=10, green) and antibody profiling was performed using the HuProt 4.0 array of unique human proteins known to be exclusively expressed in platelets: Platelet Factor 4 (PF4), Glycoprotein VI (GP6), integrin alpha2 gene (ITGA2). Other proteins highly expressed in platelets were also evaluated: von Willebrand Factor (vWF), Platelet And Endothelial Cell Adhesion Molecule 1 (PECAM1), E-selectin (SELE), P-selectin (SELP), Vascular Cell Adhesion Molecule 1 (VCAM1). Coagulation cascade protein or receptors were also evaluated: Factor II (F2), factor II receptor (FIIR), Factor X (F10). A heat map was generated to show each antibody concentration expressed as standard deviations (SD) above or below the mean concentration found in healthy individuals.