Dysregulated Platelet Function and Thrombosis in Patients with

Post-Acute Sequelae of COVID-19

Anu Aggarwal, PhD[1], Tamanna K. Singh MD[2], Michael Pham MS[2],

Matthew Godwin BS[1], Rui Chen MS[1], Thomas M. McIntyre PhD[1], Mina K. Chung MD[1,2],

Courtney Jennings MS[1], Mariya Ali BS[1], Huijun Park MS[1], Kristin Englund MD[3],

Alok A. Khorana MD[1,4], Lars G. Svensson MD PhD[5], Samir Kapadia[2],

Keith R. McCrae MD[1,2,3], and Scott J. Cameron, MD PhD [1,4,5†]

Running Title: Decreased thrombosis in long COVID-19

Affiliations:
[1] Lerner Research Institute, Department of Cardiovascular and Metabolic Sciences, Cleveland,
Ohio
[2]Heart Vascular and Thoracic Institute, Department of Cardiovascular Medicine, Section of
Vascular Medicine, Cleveland Clinic Foundation, Cleveland, Ohio, USA
[3]Department of Infectious Disease, Cleveland Clinic Foundation, Cleveland, Ohio, USA
[4] Taussig Cancer Institute, Department of Hematology and Oncology, Cleveland, Ohio
[5]Heart Vascular and Thoracic Institute, Department of Cardiovascular Medicine, Department of
Cardiac Surgery, Cleveland Clinic Foundation, Cleveland, Ohio, USA
20

21
22 [†]**Address for Correspondence**: Section of Vascular Medicine, Heart Vascular and Thoracic
23 Institute, Department of Cardiovascular Medicine, Section of Vascular Medicine, Cleveland
24 Clinic Foundation, Cleveland, Ohio, USA
25 Email: cameros3@ccf.org
26

27 Word count: (abstract) 228, and (main manuscript) 2747
28

29
30 **Key words:** COVID-19, Post-acute sequelae of COVID-19 (PASC), Platelets, Thrombosis.
31

Abstract

Objective: Post-acute sequelae of COVID-19 (PASC, also referred as Long-COVID) sometimes follows COVID-19, a disease caused by SARS-CoV-2. While SARS-CoV-2 is well-known to promote a prothrombotic state, and especially to activate platelets acutely, less is known about the thrombosis risk in PASC.

Approach and Results: PASC patients and age-matched healthy controls were enrolled in the study on average 15 months after documented SARS-CoV-2 infection. Platelet activation was evaluated by Light Transmission Aggregometry (LTA) and flow cytometry in response to platelet surface receptor agonists. Thrombosis in platelet-deplete plasma was evaluated by Factor Xa activity. A microfluidics system assessed thrombosis in whole blood under venous shear stress conditions. While only a mild increase in platelet aggregation in PASC patients through the thromboxane receptor was observed platelet activation through the glycoprotein VI (GPVI) receptor was markedly decreased in PASC patients compared to age- and sex-matched healthy controls. Thrombosis under venous shear conditions as well as Factor Xa activity were reduced in PASC patients. Plasma from PASC patients was an extremely potent activator of washed, healthy platelets – a phenomenon not observed using age- and sex-matched platelets from healthy individuals.

Conclusions: PASC patients demonstrate dysregulated responses in platelets and coagulation in plasma, likely caused by a circulating plasma-derived molecule that promotes thrombosis. A hitherto undescribed protective response appears to exists in PASC patients to counterbalance ongoing thrombosis that is common to SARS-CoV-2 infection.

Introduction:

COVID-19, caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), may have residual effects following recovery. One third of infected patients are asymptomatic, while others experience pulmonary and cardiovascular symptoms leading to morbidity and mortality [1]. The risk of both severe and persistent symptoms of SARS-CoV-2 is increased in males, in the advanced-age patient population, in individuals requiring immunosuppression, and in patients with chronic kidney disease [2]. The risk of severe and persistent symptoms of SARS-CoV-2 remains in spite of widespread vaccination [3,4].

Post-acute sequelae of COVID-19 (PASC, also referred as Long COVID) is an emerging multi-systemic condition characterized by persistent symptoms four weeks beyond the initial SARS-CoV-2 infection. While acute respiratory distress syndrome (ARDS) is a feared consequence of severe COVID-19, many patients in intensive care units are reported to have thromboembolic complications including deep vein thrombosis (DVT) and pulmonary embolism (PE). The incidence of venous thromboembolic (VTE) may be as high as 40% [5,6]. Observational data suggests that PASC patients continue to develop or have persistent symptoms long after recovery that were initially labeled Long COVID by the Centers for Disease Control (CDC) and the National Institutes of Health (NIH). The National Institute for Health and Care Excellence (NICE) defines PASC as symptoms 4-12 weeks post-infection [7-9]. Patients blighted by the specter of PASC show structural and functional organ derangements involving the respiratory, cardiovascular, neurological, genitourinary, gastrointestinal,

and musculoskeletal systems. PASC symptoms typically include fatigue, dyspnea,

chest pain/tightness, headache, cough, difficulty concentrating, and altered sensorium.

Poor memory and concentration are the most frequent symptoms reported by patients

with PASC, closely followed by cognitive impairment, sleep disturbance, and post-

traumatic stress disorder (PTSD) [9-13].

Elevated blood markers biomarker of a pro-thrombotic state including D-dimer, C-

reactive protein, and neutrophil extracellular traps formation have been detected in

acute SARS-CoV-2 infection. In addition, many PASC patients complain of dyspnea on

exertion, which leads to physical inactivity and may further increase thromboembolic

risk. Virus-induced endothelial injury of severe COVID-19 initiates dysregulated

coagulation in patients. Increased levels of von Willebrand Factor (VWF), likely from

both endothelial cell and platelet activation, lead to hypercoagulability and thrombosis in

COVID-19 patients [7,8,14]. Platelets have an important and established hemostatic

function by preventing excessive bleeding (hemostasis) as well as an important immune

response to invading viral and microbial infections, especially SARS-CoV-2 [15].

In a large observational study of COVID-19 patients who never experienced thrombosis,

no additional risk for VTE after hospitalization with acute SARS-CoV-2 infection was

noted after hospital discharge[16]. Contrasting this, the MICHELLE study suggested a

beneficial effect of an oral Factor Xa inhibitor on discharged patients by protecting

against DVT if the patient was were previously hospitalized, even without an initial

thrombotic event [17]. These findings raise the possibility that following SARS-CoV-2

recovery, patients may have ongoing functional impairment in the coagulation cascade

or abnormal platelet function. The Aim of this study was to evaluate both platelet

function and thrombotic potential in whole blood, in platelet-deplete plasma, and in

isolated platelets from patients following acute SARS-CoV-2 recovery but with clear

PASC symptoms.

Methods

Human Subjects: 20 patients with Post-Acute Sequelae of SARS-CoV-2 infection (PASC) were enrolled on the basis of positive Covid-19 diagnosis and with persistent symptoms, including shortness of breath, fatigue, brain fog, dyspnea on exertion, mood disturbance, weakness, and memory impairment. PASC patients were compared to 13 matched healthy controls. Patients and healthy subjects were recruited only after informed consent by a coordinator not involved in clinical care. All study procedures were conducted in accordance with the Declaration of Helsinki and our approved local Institutional Review Board protocol.

Light Transmission Aggregometry (LTA): Blood from healthy controls and PASC patients was collected in sodium-citrate tubes after obtaining informed consent. Within 30 minutes to 1 hour of collection, anticoagulated whole blood was centrifuged at 200g for 15 minutes at room temperature (RT) to obtain platelet rich plasma (PRP). PRP was collected in 15mL falcon tubes and Platelet poor plasma (PPP) was obtained from the remaining sample by re-centrifugation at 2500 g for 10 minutes. PPP was used to initiate baseline optical density. Platelet Aggregation was tested following stimulation with agonists: Adenosine diphosphate (ADP), Thrombin Receptor Activator Peptide 6 (TRAP-6), thromboxane A_2 (TXA$_2$) analogue U46619, and Collagen Related Peptide (CRP). Optical density changes were recorded after the addition of agonists (CHRONO-LOG 700 aggregometer) for 6 minutes while stirring the sample at 1200 rpm at a temperature of 37°C.

Alpha granule exocytosis with Flow-cytometry: Platelets were isolated from whole blood collected from healthy controls and PASC patients in sodium-citrate tubes. Briefly, whole blood was centrifuged at 200g for 15 minutes at RT and then PRP was collected in the 15mL falcon tubes. Collected PRP was centrifuged again at 200g for 10 minutes to remove remaining WBCs and RBCs. Platelets were washed using Tyrode's buffer in the 1:1 ratio with the addition of Prostaglandin I2 (PGI2; 10nM). Washed platelets were re-suspended in 10mL of Tyrode's buffer. Washed platelets from PASC patients and healthy controls were stimulated using different concentrations of agonist: TRAP-6 (1, 5, 10 and 20µM), U46619 (0.1, 0.5, 1.0, 10µM), ADP (0.01, 0.1, 1.0, 10µM) and CRP (0.05, 0.1, 0.5, 1.0µg/mL) for 15 minutes and then incubated with phycoerythrin (PE)-conjugated anti-CD62P for 30 minutes at RT. Samples were fixed with 4% paraformaldehyde. P-Selectin expressed on the surface on the platelets signifies an activated state. Samples were read on the flow cytometer (BD Accuri C6 Plus) to acquire 20000 events. Data were analyzed using FloJo software.

The Total Thrombus formation Analysis System (T-TAS01): The T-TAS01 microfluidic chamber system was use,d and the results were compared to healthy controls. T-TAS01 is a microchip flow-chamber system that measures thrombus formation under blood flow conditions over time. Whole blood collected in the sodium-citrate tube was used for the test. Before measurement 20µL of the CaCTI reagent (Corn trypsin inhibitor) was added to 480µL of whole blood and then this blood was mixed gently with the pipette. Then 450µL of mixed blood was loaded into the reservoir

of the AR chip reservoir coated with both collagen and thromboplastin (tissue factor)

with a shear rate of 600 s^{-1}. As coagulation and platelet activation progress, thrombosis

is noted as increased pressure in the channel following shear rate-mediated activation

of blood in the microchip channel. These pressure changes are monitored by a pressure

sensor upstream of the chamber and are defined as The Occlusion Start Time (OST),

and the Occlusion Time (OT). The difference between OT and OST was calculated for

each subject. Furthermore, to check the pro-thrombotic activity of the plasma of the

PASC patients, 200µL plasma from the PASC patients and healthy controls was

incubated with 800µL whole blood from healthy control for 30 minutes. Then this spiked

whole blood with plasma was loaded into the reservoir of the AR chip.

Plasma Factor Xa activity: Factor Xa (FXa) is the activated form of the coagulation

factor X. Factor X, a serine endopeptidase plays an important role in the coagulation

pathway. It covverts prothrombin into active thrombin by complexing with activated co-

factor V in the prothrombinase complex. Factor Xa Activity assay kit from Abcam

(ab204711) was used to measure the Factor Xa activity in the plasma of the PASC

patients compared to age- and sex-matched healthy controls. The assay was performed

according to the manufacturer's recommendations. The samples were read on the flow

cytometer (BD Accuri C6 Plus) to acquire 20000 events and data were analyzed using

FlowJo software. VICTOR Nivo multimode microplate reader (Pekin Elmer).

Statistical Analysis: All data were interrogated for normalcy using the Shapiro-Wilks

test. Normally distributed continuous variable group differences were assessed by the

student 2-tailed Student's t-test and skewed data assessed by the Mann-Whitney *U*

test. Analyses were conducted using GraphPad Prism 7 (GraphPad Software). For

Gaussian-distributed data in 3 or more groups, 1-way ANOVA then the Bonferroni

multiple comparisons test was used, otherwise the Kruskal–Wallis test followed by Dunn

post-test was used. Significance was accepted as a *P* value of less than 0.05.

Results

Baseline Study Population: 20 patients with PASC (mean age 46.5 years) were recruited from a pulmonary medicine clinic in Northeast Ohio and compared with 13 healthy volunteers (mean age 52.3). Healthy recruits or PASC patients had comparable demographics and co-morbidities, showing no significant difference between groups as indicated in **Table 1**. Patients with PASC were studied on average 471 days $\pm$ 77 days after the last detected SARS-CoV-2 amplicon by polymerase chain reaction (PCR). Of the 26 common symptoms in our PASC patient population, the majority described fatigue (80%), brain fog (80%), shortness of breath at rest (70%), and exertional dyspnea (70%) (**Supplemental Table 1**).

Platelet-dependent thrombosis in PASC: Platelet aggregation in PRP was evaluated by LTA after stimulating platelets with surface receptor agonists directed toward the $P2Y_{12}$ receptor (ADP), the thromboxane receptor (U46619), the GPVI receptor (CRP), and PAR1 (TRAP-6). Overall, platelet aggregation by LTA in PASC was similar to healthy individuals except through the thromboxane receptor, which demonstrated increased reactivity following stimulation with the platelet surface receptor agonist U46619 (87%±3 for PASC vs. 72%±4 for healthy subjects, p=0.03) (**Fig. 1 A-D**). To distinguish platelet function that may be altered by plasma mediators, these studies were repeated after washing platelets twice, then assessing platelet reactivity after dese-dependent agonist stimulation by alpha granule exocytosis and appearance of P-selectin on the surface of the platelet as reported previously[18,19] . When washed platelets from healthy volunteers

and PASC patients were stimulated with platelet surface receptor agonists, we found

similar reactivity to healthy individuals through the following receptors: $P2Y_{12}$,

thromboxane, PAR 1 (**Fig. 2A-C**). Curiously, platelet reactivity through GPVI was

markedly decreased in PASC patients compared with healthy controls (**Fig. 2D**).

Platelet-independent thrombosis in PASC: Whole blood was drawn from patients who

were healthy or with PASC, then passed through a 80 μm channel in a microfluidics

system at 600 dynes (s^{-1}) which approximates shear stress conditions in veins. This

assay was specifically chosen because thrombosis in most patients with SARS-CoV-2

occurs in the venous vasculature, and the assay utilizes tissue factor and collagen to

initiate thrombosis, which simulates endothelial damage common to SARS-CoV-2.

Overall, the time to pressure increase in the microchannel (kPa), which is a signature of

thrombosis, as coagulated blood demonstrates diminished flow, was prolonged in PASC

patients compared with healthy conditions (occlusion time-occlusion start time, 94±8 vs.

73±8 seconds, respectively, P=0.022) (**Fig. 3 A-B**). To test whether a mediator in

plasma is responsible for the difference in coagulation observed in PASC, whole blood

from a healthy individual was spiked with platelet-deplete plasma from a PASC patient

compared to spiking platelet-deplete plasma from a healthy individual. This experiment

showed no difference between occlusion time and occlusion start time (62±4 healthy

plasma vs. 66±8 PASC plasma, n=5 in each group, P=0.74). We next assessed

platelet-deplete plasma Factor Xa activity which represents the point of convergence

between the extrinsic and the intrinsic arms of the coagulation cascade. Factor Xa

activity was reduced in PASC patients compared to healthy individuals (23.6±5 vs. 50±8

ng/100µL, respectively, P=0.037). (**Fig. 3C**). These experiments suggest a change in

PASC plasma resulting in dysregulated coagulation compared with healthy conditions.

A plasma mediator in PASC patients activates platelets: We demonstrated that washed

platelets from PASC patients are relatively unchanged or less reactive than healthy

platelets (**Fig. 2**) and that platelet-deplete plasma in PASC is less likely to clot *ex vivo*

(**Fig. 3C**). These observations suggests a potential adaptive response long after

COVID-19 infection, possibly to prevent thrombosis that is common in patients with

acute COVID-19 infection. When platelet-deplete plasma from healthy individuals or

PASC patients was added to healthy, twice washed platelets (n=20 plasma samples for

PASC and n=18 plasma samples for healthy controls), there was a surprising and

profound activation of healthy platelets through all platelet receptor agonists tested after

PASC plasma inclubation (**Fig. 4**). Given that immunothrombosis is common in COVID-

19 and the complement system likely contributes to these responses [20-22], we heated

and therefore inactivated plasma from healthy donors or PASC patients at 55˚C and

repeated these experiments. We observed that platelets were persistently activated if

spiked with room temperature plasma or heat-inactivated plasma prior to agonist

stimulation from PASC patients. This suggests complement activation is unlikely to be a

key driver in the augmented platelet response observed by PASC plasma exposure to

healthy platelets (**Fig. S2**).

Discussion

SARS-CoV-2 is well-known to precipitate thrombotic events, especially DVT and PE [23,24]. Acute SARS-CoV-2 infection has been shown to coincide with activated platelets by multiple independent groups [25-28]. Our primary objective was to assess platelet function and thrombosis in human blood in patients with PASC after recovery from COVID-19 given that rheologic manifestations of this syndrome are unknown. Our major observation in patients more than one year following acute SARS-CoV-2 and with clear symptoms of PASC is one of dysregulated platelet function, with slight selective platelet activation through the thromboxane receptor, and attenuated activation through the GPVI receptor. We did not observe a change in platelet receptor signaling through PAR1 or $P2Y_{12}$ in PASC using two well-validated techniques to interrogate platelet activation.

This study may be the first to report a functional defect in the coagulation cascade given our finding of reduced activated Factor Xa in platelet-deplete plasma in patients with PASC and prolonged time to thrombosis of whole blood under conditions of biomechanical shear under flow. [26]. Using radioligand binding technology or a similar chromogenic substrate for Factor Xa activation as we employ, others have reported that platelets and platelet-derived microparticles may be a source of Factor Xa activity in the circulation [29-31]. Moreover, a putative signaling mechanism between the platelet GPVI receptor and Factor Xa was previously reported [32]. In addition, the profoundly under-reactive GPVI receptor identified by flow cytometry using CRP as the agonist in our study complements the observation of prolonged time to thrombosis in human blood *ex*

vivo under conditions of venous shear stress since collagen is present in the

microfluidics channel.

The most significant finding of our study is that platelet-deplete plasma in PASC

patients 'primes' and markedly activates healthy, washed platelets through every

receptor signaling pathway we evaluated. We did not see this when platelet-deplete

plasma from healthy controls was incubated with washed platelets prior to platelet

receptor agonist stimulation. The hyper-reactive platelet phenotype persisted even after

heating plasma to 55 °C suggesting the complement cascade is unlikely to be

responsible for the augmented platelet activation observed. Therefore, we offer the

suggestion that a putative pro-thrombotic factor exists in the blood of patients with

PASC, and its presence leads to an adaptive response in the host in which platelet

reactivity and thrombosis are paradoxically limited, presumably as a protective response

from thrombosis.

Our study in PASC patients differs from a recent research letter by Martins-Gonçalves

et al. in patients reporting at least one persistent respiratory symptom after acute

hospitalization for COVID-19 and with persistent platelet hyperreactivity[33]. Distinct

differences in the enrolled subjects is apparent. Firstly, while the average age of

patients enrolled by Martins-Gonçalves *et al.* was similar to our cohort, the represented

majority were men while 65% of our patients and 84% of our healthy controls were

women. Platelet reactivity has been shown to differ between men and women in health

and in cardiovascular disorders [19,29,34]. Secondly, the control group employed by

Martins-Gonçalves *et al.* tested negative for SARS-CoV-2 weekly for almost two years

prior to enrollment, while many of our control comparators reported SARS-CoV-2

infection in the preceding two years but without residual symptoms at the time of

enrollment.   This was the major reason why they were selected as controls for our

study. Lastly, Martins-Gonçalves evaluated platelet reactivity more in the convalescent

setting 1-4 months after acute SARS-CoV-2 infection in 2021 whereas our PASC

patients on average were evaluated 15 months after the last documented acute SARS-

CoV-2 infection in 2023.   Clinical outcomes and symptoms following SARS-Cov-2

infection appear to vary depending on vaccination status as well as the the viral strain of

the infected individual [35]. This means that interpretation of data gained from patients

evaluated during different waves of the COVID-19 pandemic require careful attention.

An important consideration is that different viral strains may manifest as different

symptoms in persons with PASC Given that SARS-CoV-2 was demonstrated to induce

morphological changes in platelets and promote apoptosis after viral internalization in

an ACE2-dendent and ACE2-independent receptor internalization pathway[15], platelet

reprogramming in patients with PASC may occur at the level of the megakaryocyte.

In summary, patients who since recovered from acute SARS-CoV-2 but with persistent

and long-standing symptoms under the constellation of PASC show evidence of

dysregulated platelet reactivity, alterations in normal coagulation in platelet-deplete

plasma, and diminished thrombosis in whole blood *ex vivo* under shear stress

conditions.   These events are likely to be a consequence of adapting to a platelet-

activating factor in plasma.   The stimulus behind the exhausted platelet phenomenon,

particularly through GPVI, in PASC patients, should be investigated and identification of

the putative pro-thrombotic mediator should be prioritized given the implications for

some patients with PASC being considered as donors of blood and plasma products in

future.

References

1. Oran DP, Topol EJ. The Proportion of SARS-CoV-2 Infections That Are Asymptomatic : A Systematic Review. *Ann Intern Med*. 2021;174:655-662. doi: 10.7326/M20-6976

2. Peckham H, de Gruijter NM, Raine C, Radziszewska A, Ciurtin C, Wedderburn LR, Rosser EC, Webb K, Deakin CT. Male sex identified by global COVID-19 meta-analysis as a risk factor for death and ITU admission. *Nat Commun*. 2020;11:6317. doi: 10.1038/s41467-020-19741-6

3. Agrawal U, Bedston S, McCowan C, Oke J, Patterson L, Robertson C, Akbari A, Azcoaga-Lorenzo A, Bradley DT, Fagbamigbe AF, et al. Severe COVID-19 outcomes after full vaccination of primary schedule and initial boosters: pooled analysis of national prospective cohort studies of 30 million individuals in England, Northern Ireland, Scotland, and Wales. *Lancet*. 2022;400:1305-1320. doi: 10.1016/S0140-6736(22)01656-7

4. Lee SW, Lee J, Moon SY, Jin HY, Yang JM, Ogino S, Song M, Hong SH, Abou Ghayda R, Kronbichler A, et al. Physical activity and the risk of SARS-CoV-2 infection, severe COVID-19 illness and COVID-19 related mortality in South Korea: a nationwide cohort study. *Br J Sports Med*. 2022;56:901-912. doi: 10.1136/bjsports-2021-104203

5. Helms J, Tacquard C, Severac F, Leonard-Lorant I, Ohana M, Delabranche X, Merdji H, Clere-Jehl R, Schenck M, Fagot Gandet F, et al. High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med*. 2020;46:1089-1098. doi: 10.1007/s00134-020-06062-x

6. Klok FA, Kruip M, van der Meer NJM, Arbous MS, Gommers D, Kant KM, Kaptein FHJ, van Paassen J, Stals MAM, Huisman MV, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res*. 2020;191:145-147. doi: 10.1016/j.thromres.2020.04.013

7. Libby P, Luscher T. COVID-19 is, in the end, an endothelial disease. *Eur Heart J*. 2020;41:3038-3044. doi: 10.1093/eurheartj/ehaa623

8. Ranucci M, Ballotta A, Di Dedda U, Baryshnikova E, Dei Poli M, Resta M, Falco M, Albano G, Menicanti L. The procoagulant pattern of patients with COVID-19 acute respiratory distress syndrome. *J Thromb Haemost*. 2020;18:1747-1751. doi: 10.1111/jth.14854

9. Crook H, Raza S, Nowell J, Young M, Edison P. Long covid-mechanisms, risk factors, and management. *BMJ*. 2021;374:n1648. doi: 10.1136/bmj.n1648

10. Carfi A, Bernabei R, Landi F, Gemelli Against C-P-ACSG. Persistent Symptoms in Patients After Acute COVID-19. *JAMA*. 2020;324:603-605. doi: 10.1001/jama.2020.12603

11. Goertz YMJ, Van Herck M, Delbressine JM, Vaes AW, Meys R, Machado FVC, Houben-Wilke S, Burtin C, Posthuma R, Franssen FME, et al. Persistent symptoms 3 months after a SARS-CoV-2 infection: the post-COVID-19 syndrome? *ERJ Open Res*. 2020;6. doi: 10.1183/23120541.00542-2020

12.   Xiong Q, Xu M, Li J, Liu Y, Zhang J, Xu Y, Dong W. Clinical sequelae of COVID-19
survivors in Wuhan, China: a single-centre longitudinal study. *Clin Microbiol Infect*.
2021;27:89-95. doi: 10.1016/j.cmi.2020.09.023
13.   Teuwen LA, Geldhof V, Pasut A, Carmeliet P. COVID-19: the vasculature unleashed.
Nat Rev Immunol. 2020;20:389-391. doi: 10.1038/s41577-020-0343-0
14.   Panigada M, Bottino N, Tagliabue P, Grasselli G, Novembrino C, Chantarangkul V,
Pesenti A, Peyvandi F, Tripodi A. Hypercoagulability of COVID-19 patients in intensive
care unit: A report of thromboelastography findings and other parameters of hemostasis.
J Thromb Haemost. 2020;18:1738-1742. doi: 10.1111/jth.14850
15.   Koupenova M, Corkrey HA, Vitseva O, Tanriverdi K, Somasundaran M, Liu P, Soofi S,
Bhandari R, Godwin M, Parsi KM, et al. SARS-CoV-2 Initiates Programmed Cell Death
in Platelets. *Circ Res*. 2021;129:631-646. doi: 10.1161/CIRCRESAHA.121.319117
16.   Patel L, Stenzel A, Van Hove C, Sidebottom A, Kethireddy R, Ha N, Beddow D,
Manunga J, Qadri G, Kirven J, et al. Outcomes in patients discharged with extended
venous thromboembolism prophylaxis after hospitalization with COVID-19. *Vasc Med*.
2023:1358863X231159945. doi: 10.1177/1358863X231159945
17.   Ramacciotti E, Barile Agati L, Calderaro D, Aguiar VCR, Spyropoulos AC, de Oliveira
CCC, Lins Dos Santos J, Volpiani GG, Sobreira ML, Joviliano EE, et al. Rivaroxaban
versus no anticoagulation for post-discharge thromboprophylaxis after hospitalisation for
COVID-19 (MICHELLE): an open-label, multicentre, randomised, controlled trial.
Lancet. 2022;399:50-59. doi: 10.1016/S0140-6736(21)02392-8
18.   Cameron SJ, Mix DS, Ture SK, Schmidt RA, Mohan A, Pariser D, Stoner MC, Shah P,
Chen L, Zhang H, et al. Hypoxia and Ischemia Promote a Maladaptive Platelet
Phenotype. *Arterioscler Thromb Vasc Biol*. 2018;38:1594-1606. doi:
10.1161/ATVBAHA.118.311186
19.   Soo Kim B, Auerbach DS, Sadhra H, Godwin M, Bhandari R, Ling FS, Mohan A, Yule
DI, Wagner L, 2nd, Rich DQ, et al. Sex-Specific Platelet Activation Through Protease-
Activated Receptors Reverses in Myocardial Infarction. *Arterioscler Thromb Vasc Biol*.
2021;41:390-400. doi: 10.1161/ATVBAHA.120.315033
20.   Aiello S, Gastoldi S, Galbusera M, Ruggenenti P, Portalupi V, Rota S, Rubis N, Liguori
L, Conti S, Tironi M, et al. C5a and C5aR1 are key drivers of microvascular platelet
aggregation in clinical entities spanning from aHUS to COVID-19. *Blood Adv*.
2022;6:866-881. doi: 10.1182/bloodadvances.2021005246
21.   Charitos P, Heijnen I, Egli A, Bassetti S, Trendelenburg M, Osthoff M. Functional
Activity of the Complement System in Hospitalized COVID-19 Patients: A Prospective
Cohort Study. *Front Immunol*. 2021;12:765330. doi: 10.3389/fimmu.2021.765330
22.   Rajamanickam A, Nathella PK, Venkataraman A, Dasan B, Putlibai S, Ahamed SF,
Selvaraj N, Sadasivam K, Sundaram B, Nutman TB, et al. Levels of Complement
Components in Children With Acute COVID-19 or Multisystem Inflammatory
Syndrome. *JAMA Netw Open*. 2023;6:e231713. doi:
10.1001/jamanetworkopen.2023.1713
23.   Elbadawi A, Elgendy IY, Joseph D, Eze-Nliam C, Rampersad P, Ouma G, Bhandari R,
Kirksey L, Chaudhury P, Chung MK, et al. Racial Differences and In-Hospital Outcomes
Among Hospitalized Patients with COVID-19. *J Racial Ethn Health Disparities*.
2022;9:2011-2018. doi: 10.1007/s40615-021-01140-2

24.   Elbadawi A, Elgendy IY, Sahai A, Bhandari R, McCarthy M, Gomes M, Bishop GJ,
Bartholomew JR, Kapadia S, Cameron SJ. Incidence and Outcomes of Thrombotic
Events in Symptomatic Patients With COVID-19. *Arterioscler Thromb Vasc Biol*.
2021;41:545-547. doi: 10.1161/ATVBAHA.120.315304
25.   Manne BK, Denorme F, Middleton EA, Portier I, Rowley JW, Stubben C, Petrey AC,
Tolley ND, Guo L, Cody M, et al. Platelet gene expression and function in patients with
COVID-19. *Blood*. 2020;136:1317-1329. doi: 10.1182/blood.2020007214
26.   Puhm F, Allaeys I, Lacasse E, Dubuc I, Galipeau Y, Zaid Y, Khalki L, Belleannee C,
Durocher Y, Brisson AR, et al. Platelet activation by SARS-CoV-2 implicates the release
of active tissue factor by infected cells. *Blood Adv*. 2022;6:3593-3605. doi:
10.1182/bloodadvances.2022007444
27.   Zaid Y, Puhm F, Allaeys I, Naya A, Oudghiri M, Khalki L, Limami Y, Zaid N, Sadki K,
Ben El Haj R, et al. Platelets Can Associate with SARS-Cov-2 RNA and Are
Hyperactivated in COVID-19. *Circ Res*. 2020;127:1404-1418. doi:
10.1161/CIRCRESAHA.120.317703
28.   Petito E, Franco L, Falcinelli E, Guglielmini G, Conti C, Vaudo G, Paliani U, Becattini
C, Mencacci A, Tondi F, et al. COVID-19 infection-associated platelet and neutrophil
activation is blunted by previous anti-SARS-CoV-2 vaccination. *Br J Haematol*.
2023;201:851-856. doi: 10.1111/bjh.18726
29.   Godwin MD, Aggarwal A, Hilt Z, Shah S, Gorski J, Cameron SJ. Sex-Dependent Effect
of Platelet Nitric Oxide: Production and Platelet Reactivity in Healthy Individuals. *JACC*
Basic Transl Sci. 2022;7:14-25. doi: 10.1016/j.jacbts.2021.10.009
30.   Miletich JP, Jackson CM, Majerus PW. Interaction of coagulation factor Xa with human
platelets. *Proc Natl Acad Sci U S A*. 1977;74:4033-4036. doi: 10.1073/pnas.74.9.4033
31.   Walsh PN, Biggs R. The role of platelets in intrinsic factor-Xa formation. *Br J Haematol*.
1972;22:743-760. doi: 10.1111/j.1365-2141.1972.tb05719.x
32.   Al-Tamimi M, Grigoriadis G, Tran H, Paul E, Servadei P, Berndt MC, Gardiner EE,
Andrews RK. Coagulation-induced shedding of platelet glycoprotein VI mediated by
factor Xa. *Blood*. 2011;117:3912-3920. doi: 10.1182/blood-2010-08-301523
33.   Martins-Goncalves R, Campos MM, Palhinha L, Azevedo-Quintanilha IG, Abud Mendes
M, Ramos Temerozo J, Toledo-Mendes J, Rosado-de-Castro PH, Bozza FA, Souza
Rodrigues R, et al. Persisting Platelet Activation and Hyperactivity in COVID-19
Survivors. *Circ Res*. 2022;131:944-947. doi: 10.1161/CIRCRESAHA.122.321659
34.   Faraday N, Goldschmidt-Clermont PJ, Bray PF. Gender differences in platelet GPIIb-IIIa
activation. *Thromb Haemost*. 1997;77:748-754.
35.   Robinson ML, Morris CP, Betz JF, Zhang Y, Bollinger R, Wang N, Thiemann DR, Fall
A, Eldesouki RE, Norton JM, et al. Impact of Severe Acute Respiratory Syndrome
Coronavirus 2 Variants on Inpatient Clinical Outcome. *Clin Infect Dis*. 2023;76:1539-
1549. doi: 10.1093/cid/ciac957
510

Tables

518   **Table 1.** Demographics Enrolled PASC Patients and Healthy Individuals

Demographics	Healthy (n=13)		PASC (n=20)		P-Value
	N	%	N	%	
Characteristics					
Mean Age (± SEM)	46.5 ± 2.95		52.3 ± 4.03		0.26
28-40 years	3	23.1 %	6	30.0 %	0.660
41-60 years	8	61.5 %	3	15.0 %	*0.006*
61-82 years	2	15.4 %	9	45.0 %	0.08
Male	2	15.4 %	7	35.0 %	0.22
Female	11	84.6 %	13	65.0 %	
BMI (± SEM)	31.8 ± 1.37		29.4 ± 1.86		0.34
Smoker	1	7.7 %	7	35.0 %	0.07
Medical History					
Hypertension	4	30.8 %	4	20.0 %	0.48
Hyperlipidemia	2	15.4 %	4	20.0 %	0.73
Diabetes	1	7.7 %	1	5.0 %	0.75
History of CVD	1	7.7%	2	10.0 %	0.82
History of Thrombosis	1	7.7 %	1	5.0 %	0.75
History of Pulmonary Embolism	0	0.0 %	0	0.0 %	1.0
Medications					
Platelet inhibitors	1	7.7 %	4	20.0 %	0.34
Anticoagulants	1	0.0 %	3	15.0 %	0.52
Statins	2	15.4 %	4	20.0 %	0.73
NSAIDs	6	46.1 %	4	20.0 %	0.11
BP Medications	4	30.8 %	1	5.0 %	*0.04*

Table 1. Baseline Characteristics. Patients with PASC (n=20, occurring at a mean duration of
471 ± 77 days after a diagnosis of COVID-19) were recruited for the study and compared with
healthy volunteers (n=13). Data are reported as mean ± SEM. Differences between groups was
calculated by the student's t-test for continuous variables or Chi-Squared for dichotomous
variables. BMI=Body Mass Index. CVD=Cardiovascular Disease. BP=Blood Pressure.
NSAID=non-steroidal anti-inflammatory medication.

532
533
534
535
536 **Supplemental Table 1.** Symptoms Experienced by PASC Patients

Symptoms	PASC (n=20)	
	N	%
Mean Time After Infection (± SEM)	471.4 days ± 77.33	
Fatigue	16	80.0 %
Brain Fog	16	80.0 %
Shortness of Breath	14	70.0 %
Dyspnea on Exertion	14	70.0 %
Dizziness	13	65.0 %
Headaches	12	60.0 %
Changes in Mood	11	55.0 %
Insomnia	9	45.0 %
Palpitations	9	45.0 %
Chest Pain	8	40.0 %
Joint Pain	8	40.0 %
Cough	6	30.0 %
Parosmia/Dysgeusia	6	30.0 %
Diarrhea	1	15.0 %
Incontinence	3	15.0 %
Memory Deficits	3	15.0 %
Body Aches	2	10.0%
HF Symptoms	2	10.0 %
Nausea	2	10.0 %
Depression	1	5.0 %
Erectile Dysfunction	1	5.0 %
Fever	1	5.0 %
Neck Pain	1	5.0 %
Orthopnea	1	5.0 %
Limb Paresthesia	1	5.0 %
Syncope	1	5.0 %

537
538 **Fig. S1. Persistent symptoms of enrolled PASC patients.**
539 Data are expressed as frequency and % of population and mean ± SEM.

550
551
552
553
554   **Figures**
555
Fig. 1. Platelet Activation in PASC by Aggregometry. Blood was drawn from healthy
individual or PASC patients and PRP was isolated prior to stimulation with **A.** TRAP-6, **B.**
U46619, **C.** ADP, **D.** GPVI. Platelet aggregation was evaluated by LTA and data reported as
mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group
differences were assessed by the Mann Whitney U test, n=5 for healthy individuals and n=5 for
PASC.
562
563   **Fig. 2. Platelet Activation in PASC by Flow Cytometry.** Isolated, washed platelets were
564   stimulated with **A.** TRAP-6, **B.** U46619, **C.** ADP, **D.** CRP for 15 minutes, then reactivity was
565   assessed by FACS by fold-change in surface CD62 (P-Selectin) using a tagged antibody. Data
566   are represented as mean ± SEM. Group differences were assessed by the Kruskal-Wallis test,
567   n=10 for healthy individuals and n=9 for PASC. NS=not significant. CRP=C=reactive protein.
568
Fig. 3. Platelet-independent Thrombosis in PASC. A. Blood was isolated and passed through
a microfluidics chip using the Total Thrombus formation Analysis System (T-TAS01) at roughly
venous shear stress (600 s^{-1}). With thrombus generation, pressure in the microfluidics chamber
(kPa) gradually increases until cessation of flow. Data are represented as mean ± SEM. **B.** The
time from the start of occlusion to cessation of blood was calculated in seconds as mean ± SEM.
Group differences were determined by the student's t-test. **C.** Citrate plasma devoid of platelets
was evaluated for Factor Xa activity using a chromogenic substrate in PASC patients (n=7) or
healthy volunteers (n=10). Data are represented as mean ± SEM. Group differences were
assessed by the student's t-test, n=12 for healthy individuals and n=10 for PASC.
578
Fig. 4. Activation of Platelets by PASC but not Healthy Plasma. Isolated, washed platelets
were pre-incubated for 30 minutes with 5 µL of age- and sex-matched healthy plasma or PASC
plasma, then stimulated with TRAP-6, 5 µM (PAR1), U46619, 1 µM (Thromboxane receptor),
ADP, 0.1 µM (P2Y$_{12}$Receptor), or Collagen-Related Peptide, CRP, 5 µg/mL (GPVI Receptor)
for 15 minutes. Platelet reactivity was assessed by FACS by surface CD62P (P-selectin) using a
tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by the
Mann-Whitney U test, n=20 plasma isolated for healthy individuals and n=18 plasma isolates for
PASC patients. Washed platelets from a healthy 25 year-old male.

Supplementary Figures:

Fig. S2. PASC Symptoms of Enrolled Subjects. Platelet-deplete citrate plasma from PASC patients already demonstrated to activate healthy male platelets divided in half and then either heat-treated (55 ˚C, 15 minutes) or not and incubated for 30 minutes prior to stimulation with a surface receptor agonist TRAP-6, 5 µM (PAR1), U46619, 1 µM (Thromboxane receptor), ADP, 0.1 µM (P2Y$_{12}$Receptor), or Collagen-Related Peptide, CRP, 5 µg/mL (GPVI Receptor). Platelet reactivity was then assessed by FACS by surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by Mann-Whitney *U* test for n=10;

642
643
644
645
Acknowledgements and Sources of Funding

We are grateful for funding from NHLBI HL128856 to SJC and HL158669 to TMM. Dr.
Khorana acknowledges research support from the Sondra and Stephen Hardis Chair in Oncology
Research. The research was, in part, funded by the National Institutes of Health (NIH)
Agreement 1OT2HL156812 through the National Heart, Lung, and Blood Institute (NHLBI)
CONNECTS program. The views and conclusions contained in this document are those of the
authors and should not be interpreted as representing the official policies, either expressed or
implied, of the NIH.

A.

Fig. 1. Platelet Activation in PASC by Aggregometry. Blood was drawn from healthy individual or PASC patients and PRP was isolated prior to stimulation with TRAP-6. Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the Mann Whitney *U* test, n=5 for healthy individuals and n=5 for PASC.

B.

Fig. 1. Platelet Activation in PASC by Aggregometry. Blood was isolated from healthy individual or PASC patients and PRP was isolated prior to stimulation with ADP. Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the student's t-test, n=5 for healthy individuals and n=5 for PASC.

C.

Fig. 1. Platelet Activation in PASC by Aggregometry. Blood was isolated from healthy individual or PASC patients and PRP was isolated prior to stimulation with CRP. Platelet aggregation was evaluated by LTA and data reported as mean % maximum aggregation ± SEM. A representative aggregometry tracing is shown. Group differences were assessed by the Mann Whitney *U* test, n=5 for healthy individuals and n=5 for PASC.

D.

Fig. 2. Platelet Activation in PASC by Flow Cytometry. Isolated, washed platelets were stimulated with TRAP-6 for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62 (P-Selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by the Kruskal-Wallis test, n=10 for healthy individuals and n=9 for PASC. NS=not significant.

A.

Fig. 2. Platelet Activation in PASC by Flow Cytometry. Isolated, washed platelets were stimulated with TRAP-6 for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-Selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by the Kruskal-Wallis test, n=10 for healthy individuals and n=9 for PASC. NS=not significant.

Fig. 2. Platelet Activation in PASC by Flow Cytometry. Isolated, washed platelets were stimulated with U46619 for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by ANOVA, followed by the Bonferroni correction, n=10 for healthy individuals and n=10 for PASC.

Fig. 2. Platelet Activation in PASC by Flow Cytometry. Isolated, washed platelets were stimulated with ADP for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by ANOVA, followed by the Bonferroni correction, n=10 for healthy individuals and n=9 for PASC.

Fig. 2. Platelet Activation in PASC by Flow Cytometry. Isolated, washed platelets were stimulated with collagen-related peptide (CRP) for 15 minutes, then reactivity was assessed by FACS by fold-change in surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by ANOVA, followed by the Bonferroni correction, n=10 for healthy individuals and n=9 for PASC.

Fig. 3. Platelet-independent Thrombosis in PASC. A. Blood was isolated and passed through a microfluidics chip using the Total Thrombus formation Analysis System (T-TAS01) at roughly venous shear stress (600 s^{-1}). With thrombus generation, pressure in the microfluidics chamber (kPa) gradually increases until cessation of flow. Data are represented as mean ± SEM. **B.** The time from the start of occlusion to cessation of blood was calculated in seconds as mean ± SEM. Group differences were determined by the student's t-test. **C.** Citrate plasma devoid of platelets was evaluated for Factor Xa activity using a chromogenic substrate in PASC patients (n=7) or healthy volunteers (n=10). Data are represented as mean ± SEM. Group differences were assessed by the student's t-test, n=12 for healthy individuals and n=10 for PASC.

Fig. 4. Activation of Platelets by PASC but not Healthy Plasma. Isolated, washed platelets were pre-incubated for 30 minutes with 5 µL of age- and sex-matched healthy plasma or PASC plasma, then stimulated with TRAP-6, 5 µM (PAR1), U46619, 1 µM (Thromboxane receptor), ADP, 0.1 µM (P2Y$_{12}$ Receptor), or Collagen-Related Peptide, CRP, 5 µg/mL (GPVI Receptor) for 15 minutes. Platelet reactivity was assessed by FACS by surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by the Mann-Whitney *U* test, n=20 plasma isolated for healthy individuals and n=18 plasma isolates for PASC patients. Washed platelets from a healthy 25 year-old male.

Fig. S2. PASC Symptoms of Enrolled Subjects. Platelet-deplete citrate plasma from PASC patients already demonstrated to activate healthy male platelets divided in half and then either heat-treated (55 °C, 15 minutes) or not and incubated for 30 minutes prior to stimulation with a surface receptor agonist TRAP-6, 5 μM (PAR1), U46619, 1 μM (Thromboxane receptor), ADP, 0.1 μM (P2Y$_{12}$Receptor), or Collagen-Related Peptide, CRP, 5 μg/mL (GPVI Receptor). Platelet reactivity was then assessed by FACS by surface CD62P (P-selectin) using a tagged antibody. Data are represented as mean ± SEM. Group differences were assessed by Mann-Whitney *U* test for n=10 (room temperature citrate plasma) or n=10 (heat-inactivated citrate plasma.