1   **Manuscript title:** The dependence of forecasts on sampling frequency as a guide to optimizing

2   monitoring in community ecology

3   **Authors:** Uriah Daugaard[1], Stefanie Merkli[2], Ewa Merz[2], Francesco Pomati[2], Owen L. Petchey[1]

4   **Affiliations:**

    5   1) Department of Evolutionary Biology and Environmental Studies, University of Zurich,

    6   Winterthurerstrasse 190, 8057 Zurich, Switzerland

    7   2) Department of Aquatic Ecology, Eawag, Swiss Federal Institute of Aquatic Science and

    8   Technology, Überlandstrasse 133, Dübendorf, 8600 Switzerland

Corresponding author: Uriah Daugaard, uriah.daugaard@ieu.uzh.ch

Manuscript type: article

Open research statement: Data and code used for the analyses and figures are available on

Zenodo: https://doi.org/10.5281/zenodo.7970886. Environmental (lake) data (Merkli et al. 2022)

are available from ERIC: https://doi.org/10.25678/00066D.

Keywords: high-frequency data, ecological forecasting, prediction, sampling design, sampling

interval, species interactions, interaction strength, empirical dynamic modelling, phytoplankton,

zooplankton, food webs, complexity.

Abstract

Facing climate change and biodiversity loss, it is critical that ecology advances such that processes, such as species interactions and dynamics, can be correctly estimated and skillfully forecasted. As different processes occur on different time scales, the sampling frequency used to record them should intuitively match these scales. Yet, the effect of data sampling frequency on forecasting accuracy is understudied. Using a simple simulated dataset as a baseline and a more complex high-frequency plankton dataset, we tested how different sampling frequencies impacted abundance forecasts of different plankton classes and the estimation of their interactions. We then investigated whether plankton growth rates and body sizes could be used to select the most appropriate sampling frequency. The simple simulated dataset showed that the optimal sampling frequency scaled positively with growth rate. This finding was not repeated in the analyses of the plankton time series, however. There, we found that a reduction in sampling frequency worsened forecasts and led us to both over- and underestimate plankton interactions. This suggests that forecasting can be used to determine the ideal sampling frequency in scientific and monitoring programs. A better study design will improve theoretical understanding of ecology and advance policy measures dealing with current global challenges.

Introduction

Ecological processes, such as species abundances and interactions, are temporal scale dependent and can range from short-term responses to perturbations (e.g. Medeiros *et al.* 2023) to long-term adaptations to changed environmental conditions (Crozier and Hutchings 2014). A central aim of ecology is not only to infer such processes from population dynamics, but also to forecast them to advance and test ecological theory and to inform decision-making (Lewis et al. 2023),

for example in conservation ecology (Tulloch, Hagger, and Greenville 2020). It is thus

intuitively important to record population sizes with sampling frequencies that match the

temporal scale of the focal processes. Despite this, little is known about how sampling

frequencies affect forecast skill, or how to select appropriate sampling frequencies.

In general, sampling infrequently can yield time series that are too sparse to adequately

infer the processes of interest (Estes et al. 2018). Infrequent sampling can cause under- or

overestimations (e.g. Queiroz and Ferreira 2009), imprecise estimates (e.g. Taylor and Howes

1994), and non-detection of focal dynamics (Lehtiniemi et al. 2022). Ultimately, this can lead to

the non-detection of relations between variables or to spurious associations (Cabella, Meloni, and

Martinez 2019), hindering also the reproducibility of results (Estes et al. 2018).

The few studies that investigated how forecasting is affected by sampling frequency

found that sampling more often could both improve and worsen forecasts (Wauchope et al. 2019;

Derot, Yajima, and Schmitt 2020). However, to our knowledge these studies did not account for

the fact that higher sampling frequencies result in bigger sample sizes, which needs to be

controlled for as it might have acted as a confounding variable by also yielding better forecasts.

While sampling sufficiently often remains a challenge in some systems, in others

automated data collection approaches that create high-frequency data have become available

(e.g. Kays et al. 2015; Besson et al. 2022). Nevertheless, also in cases like these it is of interest to

select appropriate sampling frequencies that avoid unnecessary samplings and associated costs.

Yet, sampling frequencies are commonly chosen based on experience and logistics, but remain

otherwise often unjustified and potentially unoptimized (Ma, McKindsey, and Johnson 2022).

Generally speaking, the sampling frequency should be high enough that all system-

relevant signals are recorded (Isles and Pomati 2021), but not higher. We hypothesize that a

candidate sampling frequency for this is the lowest frequency that still yields the highest forecast

skill, as this might suggest that all relevant signals have been captured by the data. However, the

high-frequency data necessary to test this is often not available and thus alternative criteria to

decide the sampling frequency are needed. For instance, in abundances time series the sampling

frequency could potentially be selected based on the growth rate and body size of the focal

species. Indeed, species with lower intrinsic growth rates (i.e. longer generation times) are

generally larger in size (Gillooly 2000; Bonner 2015), and their abundances tend to be forecasted

better (Petchey et al. 2015; Anderson and Gillooly 2020).

In studies where multiple species are present, however, the different growth rates

complicate the process of finding the optimal sampling frequency. Further, a higher species

richness implies more species interactions (Borrett and Patten 2003) and abundance forecasts

depend on how connected the focal species is, with forecasts being better for species that have

many but on average weak interactions (Daugaard et al. 2022). Consequentially, the question

becomes whether growth rates can guide the selection of sampling frequencies also in multi-

species studies. Moreover, species growth rates and the sampling frequency might not only

impact the achieved forecast skills, but also the estimation of species interactions, which is a

central aim of community ecology but crucially also strongly data-dependent (Marquez et al.

2022). It follows that if both forecasts and interaction estimates are affected by the sampling

frequency, relations between them might be undetectable if the sampling design is inadequate.

Here, we aimed at clarifying the influence of sampling design choices on the outcome of

quantitative analyses, which ultimately can help the design of experiments and field

observations. In a first step, we carried out a simple simulation of abundance time series to test

the possibility of selecting the sampling frequency based on species growth rates and to clarify

the relation between sampling frequency and abundance forecast skill. We then extended the

analysis by using field data and explored whether the results found *in silico* hold in natural

systems and estimated the impacts of sampling frequency, number of time points, growth rates

and body sizes first on abundance forecasting and in a second part also on the estimation of

species interactions. Lastly, we combined the species interaction estimates and abundance

forecast skills found in the previous step. We investigated whether the relation between these

two quantities found in a controlled laboratory setting is also found in an observational dataset

with greater system complexity and whether the sampling design affected this detection.

Alongside the simulations we used a high-frequency lake plankton dataset. Plankton

population dynamics represent an ideal framework for our study. They show variable and fast-

paced time series, driven by their short generation times and their interactions with their system

components (e.g. climate and nutrients, Philippart et al. 2000; Merz et al. 2023). Plankton is of

great importance in marine and freshwater ecosystems because it is at the base of aquatic food

webs regulating biomass transfer and element cycles, and their dynamics affect many ecosystem

services (Falkowski 2012). There is therefore a strong interest in forecasting plankton dynamics,

including algal blooms (e.g. Rousso *et al.* 2020; Woelmer *et al.* 2022).

Using the field data, we separately (1) sub-sampled the time series using different

frequencies while keeping the number of time points constant and (2) reduced the number of

time points while keeping the sampling frequency constant. We then estimated the interactions

between plankton classes and forecasted their abundances. We hypothesized that growth rates

could aid the selection of adequate sampling frequencies in the simulations, but that the greater

complexity of field data might prevent this. We also anticipated that fewer or sparser samplings

would generally result in worse abundance forecasts, and potentially in under- or overestimations

of field plankton interactions. Lastly, we also expected that the non-optimization of frequency

and number of samplings would result in insufficient statistical power to detect expected

associations between variables, such as the one between forecast skill and species interactions.

Material and Methods

Forecast error as a function of sampling design and functional traits

Field data collection and processing

We recorded plankton abundance time series at three meters depth in the eutrophic Lake Greifen

(northern Switzerland) with an automated darkfield microscope based on the Scripps Plankton

Camera system (Orenstein et al. 2020). The camera takes images at two magnifications (5.0

aimed at phytoplankton and ciliates and 0.5 for zooplankton species) every hour for ten minutes

with a frame rate of one frame per second. The instrument has previously been calibrated and its

performance compared to traditional microscopy methods has been validated (Merz et al. 2021).

For this study, we used images from April 2019 until December 2021 (994 days). We

processed these images as described by Merz *et al.* (2021) to extract regions of interest (ROIs,

i.e. the imaged individuals). It has been shown that ROIs/sec is a valid proxy of plankton

densities and that ROI area is a robust estimate of their body size (Merz et al. 2021). We

classified the ROIs with previously trained convolutional neural networks into zooplankton and

phytoplankton classes (Kyathanahally et al. 2021; Kyathanahally 2022). For the zooplankton we

had the classes Calanoid Copepods, Cyclopoid Copepods, Ciliates, Daphnids, Nauplii and

Rotifers. As is commonly done, we grouped the phytoplankton species into six bins of equal

width on the $\log_{10}$-scale based on their cell size (i.e ROI area). Size-based phytoplankton bins

can be studied as a function of environmental conditions (Yvon-Durocher et al. 2010; Marañón

2015). We calculated daily abundances by summing the hourly abundances (ROIs/sec) per size-

bin and per zooplankton class (Appendix S1: Figure S3). We further used lake properties

available from Merkli et al. (2022), specifically the daily epilimnetic temperature, the mixed

layer depth and irradiance and the weekly concentrations of ammonium, nitrate and phosphate.

See Appendix S1: Section S1.2 for details regarding the data collection and processing.

Processing of recorded time series

The field data included 1.61% missing data for the phytoplankton and 3.72% for the zooplankton

groups. We imputed the missing data by using cubic hermite splines to create complete times

series with equidistant spaced data points. From the weekly nutrient chemistry data, we imputed

daily values using loess (span=0.15). We added noise to all imputed values to avoid statistical

artifacts caused by the imputation (Appendix S1: Section S1.2.3). Following commonplace

practices (e.g. Beninca *et al.* 2008), we further processed the time series by carrying out a fourth-

root power transformation of the data to dampen population spikes and by detrending and

standardizing the time series. We detrended the data by regressing the time series against time

and henceforth using the standardized residuals as the new time series (Appendix S1: Figure S4).

Estimation of maximum net growth rates

Using the untransformed data, we estimated the daily growth rates of the plankton classes as

$r_T = \log(N_{T,t}/N_{T,t-\tau})/\tau$, with N being the abundance, T the target, t the time and τ the time

step (here $\tau = 1$ day). To avoid overestimations of growth rates caused by noise, we selected the

90[th] percentile of the positive values of r_T as the maximum net growth rate of each target.

Subsampled time series: reducing sampling frequency and number of time points

To have time series of different resolutions we subsampled the daily data using different

frequencies as visualized in Figure 1. A constraint was that at the lowest sampling frequency the

recorded time series still contained sufficient time points. We set the lowest sampling frequency

to 1/12 (i.e. one sampling every 12 days, resulting in 82 time points). So that we could forecast

the same time ahead across all sampling frequencies, the other sampling frequencies were: 1/6,

1/3 and 1/1 (i.e. daily sampling). We further constrained all the subsampled time series to also

contain 82 time points. To achieve this while also ensuring that we used all time points exactly

once per sampling frequency, we created 12 datasets at each sampling frequency (Figure 1). For

the daily sampling the 12 datasets were sequential, while for the sampling frequency of once

every 12 days their start dates each differed by one day. Accordingly, and as necessary, at the

two intermediate sampling frequencies the datasets were both sequential and shifted by one day.

In summary, with this subsampling method we varied the sampling frequency while

controlling the number of data points in the time series. This necessarily resulted in varying

absolute time series lengths. For instance, the daily sampled times series are 82 days long, while

those with samples every 12 days are 973 days long. This created the possibility that the less

frequently sampled times series contained stronger longer-term dynamics (see discussion).

We created time series datasets containing different amounts of time points by keeping

the following proportions of the daily data: 9/12, 6/12, 3/12, 2/12 and 1/12 (Figure 1). To use

every recorded time point exactly once regardless of the proportion of data retained, we created

several datasets at each of the desired lengths (see Figure 1). The exceptions to this are the two

datasets that kept 9/12 of the time points as they partially overlapped (Figure 1).

Forecasting of species abundances

We forecasted the abundances of taxa from simulated and field data using empirical dynamic

modeling (EDM, Ye et al. 2015) using the R-package "rEDM" (Park et al. 2021). In EDM the

state of a variable (e.g. the abundance of a taxa) is predicted based on how the variable behaved

when it was in a similar state at other times. The similarity of states can be determined using

multiple (time-lagged) variables (Takens 1981) and the number of variables used for this is the

embedding dimension E (Appendix S1: Section S1.3).

Forecasting of simulated dynamics

We carried out a simulation study to investigate the relation between sampling frequency,

number of time points, growth rate and forecast error. The aim was to have a simple baseline

model (i.e. with minimal assumptions and complexity) to inform our research hypotheses under

controlled settings and which we then tested in the more complex setting of the field data. We

simulated single species time series by using the R-package "odin" (FitzJohn 2022) and the

delayed logistic equation (Ruan 2006): $\frac{dN}{dt} = rN(1 - \frac{N_{t-\tau}}{K})$. In this equation, the instantaneous

rate of change dN/dt of the abundance N of a taxon depends on the abundance at time point t

minus τ (the time delay), the growth rate r and the carrying capacity K. Stable population cycles

emerge when $r\tau > \pi/2$. We varied the number of time points, sampling frequency (between 2

and 1/8 samplings per day), measurement noise and growth rate (between 0.3 and 1.2, with

constant $r\tau$). We repeated the simulations 100 times for each combination of parameter values.

We used simplex EDM (a single time series technique) with optimized E for the forecasts

(Sugihara and May 1990). We split the simulated time series into training and evaluation data.

We used the former to train the eight days-ahead forecast models (matching the lowest sampling

frequency). We then forecasted the abundances in the evaluation data with the models. To assess

the forecasts, we calculated the standardized root mean square error (RMSE, i.e. the forecast

error, with a value below one indicating a better forecast than one based on the average

abundance). See Appendix S1: Sections S1.1 and S1.3.1 for more information.

Forecasting of field abundances

We forecasted the abundances of the phyto- and zooplankton classes (henceforth referred to as

targets). As predictors we used the targets, the mean epilimnion temperature, the mean mixed

layer depth and irradiance, and the concentrations of ammonium, phosphate, and nitrate. We

used the EDM technique multiview embedding (Ye and Sugihara 2016). This approach is

capable of dealing with high-dimensional systems by fitting all possible low-dimensional (i.e.

small E) forecast models of which the best ones are then used in an ensemble forecast.

For all the datasets we made 12-days ahead forecasts (matching the lowest sampling

frequency) varying the number of steps forecasted ahead based on sampling frequency

(Appendix S1: Table S1). With the reduced sampling frequency datasets, we forecasted target

abundances using leave-one-out cross-validation (CV). To reduce computational load, for the

complete daily dataset and all other datasets we used k-fold CV, by dividing the time series into

training data and evaluation data (21 time points, Appendix S1: Table S1). We evaluated

forecasts based on the standardized RMSE as a measure of forecast error. We summarized across

the k-fold CV and across datasets of the same sampling frequency and of the same number of

time points by calculating the median RMSE (Appendix S1: Section S1.3.2).

Forecast error regressions

We used the calculated forecast errors as the response variable in separate regressions for each of

the explanatory variables: (1) sampling frequency; (2) number of time points; (3) target

maximum net growth rates; (4) target body size. For regressions (1) and (2) we included an

interaction term with a variable indicating whether the targets were zoo- or phytoplankton and

we included random intercepts and slopes for each target. For regressions (3) and (4) we used the

complete data and $\log_{10}$-transformed the explanatory variables to meet the model assumptions.

Interaction estimates as a function of sampling design and functional traits

Estimation of number and strength of interactions

We determined which field targets were causally linked with a test of causation (convergent cross mapping CCM, Sugihara *et al.* 2012). We followed the recommendations of Deyle et al (2016) and added a convergence test (Appendix S1: Section S1.4). In this way we determined the number of interactions of the targets. We did this in every dataset separately.

To estimate the interaction strengths between causally linked state variables we used regularized Smap EDM (Deyle et al. 2016; Cenci, Sugihara, and Saavedra 2019). This method calculates the interaction time series between interacting variables. This is again done by considering how the system reacted when it was in a comparable state at other times, but more weight is assigned to more similar states. The regularized version of Smap further handles process noise. More details regarding CCM and Smap is given in Appendix S1: Section S1.4.

To calculate the mean interaction strength of a target, we averaged the absolute values of estimated interaction time series over both time and interacting state variables. To have a single estimate of number and mean strength of interactions per target at each sampling frequency and proportion of time points retained, we averaged over the respective datasets.

Interaction estimates regressions

As for the forecast error, we used the number and the average strength of interactions of the targets as response variables in separate regressions with the explanatory variables: (1) sampling frequency; (2) number of time points; (3) target maximum net growth rate; (4) target body size. The used covariates, interactions, random effects and covariate transformations were identical to when the response variable was the forecast error. Models (3) and (4) were Quasi-Poisson regressions for the number of interactions. This was not the case in models (1) and (2) because

the number of interactions was not a count but an average calculated over the reduced datasets.

Effect of sampling design on the detection of correlations between variables

In a final step, we combined the abundance forecasts and the interaction estimates. A recent

result shows that forecasts are better for targets with many but on average weak interactions

(Daugaard et al. 2022). We tested whether we found this relation also in the field data and

whether the sampling frequency and the number of time points influenced the detection.

Accordingly, for each sampling frequency and proportion of time points retained we fitted two

regressions with the forecast error as the response variable, separately for the two explanatory

variables number of interactions and mean interaction strength.

Results

Forecast error as a function of sampling design and functional traits

In the simulations, the shape of the relation between sampling frequency and forecast error

depended on the growth rate of the target (Figure 2a). The optimal sampling frequency (i.e. the

one resulting in the lowest RMSE) increased with target growth rate (Figure 2b). Neither the

measurement error nor the number of time points changed this pattern (Appendix S1: Figure S5).

We found no relation between the maximum net growth rates of the field targets and the

optimal sampling frequency, as we achieved the best forecasts with the daily sampling for all but

one target (1/12 samplings per day were best for ciliate forecasts, Figure 2b). Across targets,

forecast errors decreased linearly by 0.026 and 0.016 for every increase of 0.1 samplings per

day ($F_{1,10} = 42.91, p < 0.0001$, Figure 2c, Appendix S1: Table S2), respectively for the phyto-

and zooplankton targets with no difference between the two groups ($F_{1,10} = 2.15, p = 0.1734$).

We found weak evidence in the complete data that the forecast error increased by 0.337

for every tenfold increase in target maximum net growth rate ($t_{10} = 2.22, p = 0.0509$, Figure

2d, Appendix S1: Table S2). Targets with larger body sizes showed bigger maximum net growth

rates ($\log_{10}$-$\log_{10}$ scale, slope $= 5.168$, $t_{10} = 2.94$, $p = 0.0147$, Figure 2e, Appendix S1: Table

S3). Target body size and forecast error were positively correlated (Appendix S1: Figure S6).

Across targets, increasing the proportion of retained time points resulted in lower forecast

errors (phytoplankton slope: -0.087; zooplankton slope: -0.050; $F_{1,10} = 5.67$, $p = 0.0386$,

Figure 2f, Appendix S1: Table S2). While there was no significant difference in slope between

the two plankton groups ($F_{1,10} = 0.418$, $p = 0.5323$), the forecast errors were lower for the

phytoplankton targets (intercept difference: -0.069, $F_{1,10} = 5.20$, $p = 0.0457$).

Interaction estimates as a function of sampling design and functional traits

Increasing the sampling frequency by 0.1 samplings per day increased the estimated number of

interactions by 0.225 across the phytoplankton targets ($t_{10.8} = 2.76$, $p = 0.0190$, Figure 3a,

Appendix S1: Table S4), but not for the zooplankton targets ($t_{10.8} = 0.03$, $p = 0.9793$). The

sampling frequency had no effect on the estimated mean interaction strengths of the targets

($F_{1,10} = 0.13$, $p = 0.7245$, Figure 3b, Appendix S1: Table S5), regardless of the plankton group.

For the phytoplankton, decreasing the proportion of time points by 10% decreased the

estimated number of interactions by 0.469 ($t_{10} = 2.69$, $p = 0.0227$, Figure 3c, Appendix S1:

Table S4) and there was weak evidence that it increased the estimated mean interaction strengths

by 0.015 ($t_{10} = 1.85$, $p = 0.0943$, Figure 3d, Appendix S1: Table S5). Across the zooplankton

targets, the proportion of time points used did not affect the interactions estimates ($t_{10} = 0.08$,

$p = 0.9376$ and $t_{10} = 0.25$, $p = 0.8069$, respectively for the number and the mean strength of

interactions), but we detected target-specific trends (Appendix S1: Figure S7).

The maximum net growth rates were unrelated with the number of interactions ($t_{10} =$

1.41, $p = 0.1890$, Figure 3e, Appendix S1: Table S4) and their mean strength ($t_{10} = 1.17$,

$p = 0.2686$, Figure 3f, Appendix S1: Table S5), as was the body size (Appendix S1: Figure S6).

Effect of sampling design on the detection of correlations between variables

We found no relation between the interaction estimates and the forecast error in the reduced

sampling frequency data (Figures 4a and 4b, Appendix S1: Table S6). With the data containing

9/12 of the time points we found a negative relation between the number of interactions and the

abundance forecast error (Figure 4c). With the data containing 9/12 and 6/12 of the time points

we found a positive relation between the mean interaction strength and the abundance forecast

error (Figure 4d). For the complete data and the other data containing a reduced number of time

points the estimated slopes between the interaction estimates and the abundance forecast errors

also matched in sign with the results of Daugaard *et al.* (2022, Figures 4e and 4f), but with

confidence intervals overlapping zero (Figures 4c and 4d; exception: data containing 1/12 of the

points in Figure 4d). We further confirmed these relations with the complete data when we made

one day ahead forecasts, as done by Daugaard *et al.* (2022, Appendix S1: Figure S10). The

number and mean strength of interactions were negatively correlated (Appendix S1: Figure S9).

Discussion

We show that abundance forecasts are negatively affected by a reduction in sampling frequency

across almost all targets. Further, our forecasts were better for targets with smaller maximal net

growth rates and for targets with smaller body sizes. Despite this, we found that growth rates are

useful indicators of the optimal sampling frequency only in the simulated single-species time

series, but not in the field data. The estimation of interactions also depended on sampling design,

as we estimated the phytoplankton targets to interact more the denser the time series were.

Sampling frequency has widely been recognized as having significant impacts on various

analyses (e.g. Lehtiniemi et al. 2022; Ma, McKindsey, and Johnson 2022), and yet sampling

frequencies are commonly too low (Estes et al. 2018). The implications for forecasting are not

well known, as we are aware of few and contrasting findings regarding the effects of sampling

frequency on forecasting (Wauchope et al. 2019; Derot, Yajima, and Schmitt 2020), and which

potentially were influenced by sample sizes. We found that lowering the sampling frequency

worsened the abundance forecasts for 11 out of the 12 targets. Because forecasting is one of the

central aims of ecology (Dietze et al. 2018), its dependence on sampling frequency exemplifies

the importance of thoughtful sampling design. Failure to carefully calibrate the sampling

frequency to match the process of interest could result in inadequate estimations and forecasts,

which ultimately could lead to wrong conclusions and inappropriate policies.

Conversely, because forecasts can be evaluated quantitatively their dependence on

sampling frequency shows their potential to guide sampling designs. Ecological forecasting is

not only an objective, but it can also advance theoretical understanding (Lewis et al. 2023).

Building on this, we argue that besides improving theory and informing decision-makers,

forecasting can guide the design of scientific studies and monitoring programs. Adjusting the

sampling frequency based on forecast skill is likely to improve the time series so that the process

relevant signals are captured, which is crucial in community ecology (Isles and Pomati 2021).

We found that we estimated fewer interactions for the phytoplankton targets when we

decreased the frequency of the samplings. As we achieved the best forecasts at the highest

sampling frequency, we expect the estimated number of interactions to be most accurate at this

frequency and underestimated otherwise. While the true target interactions are unknown and thus

the correctness of the estimations cannot be evaluated, this may become possible thanks to recent

advances in the manipulation of species interactions in microcosm experiments (Hu et al. 2022).

The high-frequency data needed to select the sampling design based on forecasts is not

always available. In such cases, alternative criteria to decide the sampling frequency are needed.

We found that selecting based on growth rates was possible in simulated single-species dynamics

but not in the field data, although we achieved better forecasts for targets with smaller growth

rates (which also had smaller body sizes, see discussion in Appendix S1: Section S2.1). Thus, in

natural systems more frequent sampling is needed than what would be expected based on growth

rates. While this could be because of a mismatch between the true target division rates and the

estimated maximum net growth rates, the found relation between the latter and the abundance

forecasts makes this less likely. Instead, we hypothesize that the reason is the greater complexity

of the natural system. Indeed, the Nyquist condition used in signal processing theory states that

to correctly record a time series the sampling frequency needs to be at least twice as high as the

highest frequency present in the true time series (e.g. Amidror 2015). With interacting variables

leaving imprints in each other's time series (Takens 1981), to adequately capture the dynamics of

a target it may be necessary to select the sampling frequency not based on the target itself, but on

the interacting variable with the fastest time series (e.g. the species with the highest growth rate).

The need for high-frequency data is further corroborated by our result that the effects of

sampling design on forecasts and interaction estimations were compounded. We were not able to

reproduce the laboratory result that the forecasting of a target depends on the number and

strength of its interactions (Daugaard et al. 2022), unless we used the high-frequency field data

containing enough time points. Auspiciously, the need for high-frequency data in ecology is

increasingly being covered (e.g. Pomati et al. 2011; Besson et al. 2022).

Reducing the number of time points worsened forecasts and led to target-specific over-

and underestimations of interactions. The smaller the sample size was the less we were able to

estimate interactions, with estimates becoming more similar for all targets (Appendix S1: Figure

S8). This further shows the importance of adequate sampling design for quantitative analyses.

Varying the sampling frequency necessarily results in time series that either differ in their

time range or in number of time points. We controlled for the latter as we considered it to be a

stronger confounder. To still try to control for the different time ranges, we used every time point

once per sampling frequency so that the entire time range was covered. Yet, it is possible that the

different time ranges influenced the results, for instance because the species network may have

changed over time (Merz et al. 2023). However, because we only considered a relatively short

time and used grouped data which is less likely to change, we do not expect this to be the case.

In conclusion, we clarify the role of sampling frequency in ecology for the forecasting

and estimation of processes. As we face climate change and biodiversity loss (Bellard et al.

2012; Cardinale et al. 2012), ecological forecasting is a field of increasing importance (Dietze

2017). Our results have the potential to improve the design of experiments and field

observations, and the estimation of species interactions (an important aspect of biodiversity).

Especially the use of forecasts not only as an aim but also as a tool shows promise in this regard.

Ultimately, better study designs will improve ecological inference and forecasting, which is

fundamental for a better theoretical understanding of ecology and for the implementation of

better performing policies and measures that deal with current global challenges.

Author contributions

377 SM, EM & FP collected, curated and processed the data. UD & OLP conceived the research

378 questions and designed the analyses. UD analyzed the data and wrote the manuscript. All authors

379 contributed critically to the drafts and gave final approval for publication.

Conflict of interest statement

381 The authors declare no conflicts of interest.

References

Amidror, Isaac. 2015. 'Sub-Nyquist Artefacts and Sampling Moiré Effects'. *Royal Society Open*

Science 2 (3): 140550. https://doi.org/10.1098/rsos.140550.

Anderson, David M., and James F. Gillooly. 2020. 'Allometric Scaling of Lyapunov Exponents

in Chaotic Populations'. *Population Ecology* 62 (3): 364–69. https://doi.org/10.1002/1438-

390X.12053.

Bellard, Céline, Cleo Bertelsmeier, Paul Leadley, Wilfried Thuiller, and Franck Courchamp.

2012. 'Impacts of Climate Change on the Future of Biodiversity'. *Ecology Letters* 15 (4):

365–77. https://doi.org/10.1111/j.1461-0248.2011.01736.x.

Benincà, Elisa, Jef Huisman, Reinhard Heerkloss, Klaus D. Jöhnk, Pedro Branco, Egbert H. Van

Nes, Marten Scheffer, and Stephen P. Ellner. 2008. 'Chaos in a Long-Term Experiment with

a Plankton Community'. *Nature* 451 (7180): 822–25. https://doi.org/10.1038/nature06512.

Besson, Marc, Jamie Alison, Kim Bjerge, Thomas E. Gorochowski, Toke T. Høye, Tommaso

Jucker, Hjalte M. R. Mann, and Christopher F. Clements. 2022. 'Towards the Fully

Automated Monitoring of Ecological Communities'. *Ecology Letters* 25 (12): 2753–75.

https://doi.org/10.1111/ele.14123.

Bonner, John Tyler. 2015. *Size and Cycle: An Essay on the Structure of Biology*. Princeton

University Press.

Borrett, Stuart R, and Bernard C Patten. 2003. 'Structure of Pathways in Ecological Networks:

Relationships between Length and Number'. *Ecological Modelling*, ISEM The third

European Ecological Modelling Conference, 170 (2): 173–84.

https://doi.org/10.1016/S0304-3800(03)00224-2.

Cabella, Brenno, Fernando Meloni, and Alexandre S. Martinez. 2019. 'Inadequate Sampling

Rates Can Undermine the Reliability of Ecological Interaction Estimation'. *Mathematical*

and Computational Applications 24 (2): 48. https://doi.org/10.3390/mca24020048.

Cardinale, Bradley J., J. Emmett Duffy, Andrew Gonzalez, David U. Hooper, Charles Perrings,

Patrick Venail, Anita Narwani, et al. 2012. 'Biodiversity Loss and Its Impact on Humanity'.

Nature 486 (7401): 59–67. https://doi.org/10.1038/nature11148.

Cenci, Simone, George Sugihara, and Serguei Saavedra. 2019. 'Regularized S-Map for Inference

and Forecasting with Noisy Ecological Time Series'. *Methods in Ecology and Evolution* 10

(5): 650–60. https://doi.org/10.1111/2041-210X.13150.

Crozier, Lisa G., and Jeffrey A. Hutchings. 2014. 'Plastic and Evolutionary Responses to

Climate Change in Fish'. *Evolutionary Applications* 7 (1): 68–87.

https://doi.org/10.1111/eva.12135.

Daugaard, Uriah, Stephan B. Munch, David Inauen, Frank Pennekamp, and Owen L. Petchey.

2022. 'Forecasting in the Face of Ecological Complexity: Number and Strength of Species

Interactions Determine Forecast Skill in Ecological Communities'. *Ecology Letters* 25 (9):

1974–85. https://doi.org/10.1111/ele.14070.

Derot, Jonathan, Hiroshi Yajima, and François G. Schmitt. 2020. 'Benefits of Machine Learning

and Sampling Frequency on Phytoplankton Bloom Forecasts in Coastal Areas'. *Ecological*

Informatics 60 (November): 101174. https://doi.org/10.1016/j.ecoinf.2020.101174.

Deyle, Ethan R., Robert M. May, Stephan B. Munch, and George Sugihara. 2016. 'Tracking and

Forecasting Ecosystem Interactions in Real Time'. *Proceedings of the Royal Society B:*

Biological Sciences 283 (1822): 20152258. https://doi.org/10.1098/rspb.2015.2258.

Dietze, Michael. 2017. *Ecological Forecasting. Ecological Forecasting*. Princeton University

Press. https://doi.org/10.1515/9781400885459.

Dietze, Michael C., Andrew Fox, Lindsay M. Beck-Johnson, Julio L. Betancourt, Mevin B.

Hooten, Catherine S. Jarnevich, Timothy H. Keitt, et al. 2018. 'Iterative Near-Term

Ecological Forecasting: Needs, Opportunities, and Challenges'. *Proceedings of the National*

Academy of Sciences 115 (7): 1424–32. https://doi.org/10.1073/pnas.1710231115.

Estes, Lyndon, Paul R. Elsen, Timothy Treuer, Labeeb Ahmed, Kelly Caylor, Jason Chang,

Jonathan J. Choi, and Erle C. Ellis. 2018. 'The Spatial and Temporal Domains of Modern

Ecology'. *Nature Ecology & Evolution* 2 (5): 819–26. https://doi.org/10.1038/s41559-018-

0524-4.

Falkowski, Paul. 2012. 'Ocean Science: The Power of Plankton'. *Nature* 483 (7387): S17–20.

https://doi.org/10.1038/483S17a.

FitzJohn, Rich. 2022. *Odin: ODE Generation and Integration*. Version 1.2.4. https://CRAN.R-

project.org/package=odin.

Gillooly, James F. 2000. 'Effect of Body Size and Temperature on Generation Time in

Zooplankton'. *Journal of Plankton Research* 22 (2): 241–51.

https://doi.org/10.1093/plankt/22.2.241.

Hu, Jiliang, Daniel R. Amor, Matthieu Barbier, Guy Bunin, and Jeff Gore. 2022. 'Emergent

Phases of Ecological Diversity and Dynamics Mapped in Microcosms'. *Science* 378 (6615):

85–89. https://doi.org/10.1126/science.abm7841.

Isles, Peter DF, and Francesco Pomati. 2021. 'An Operational Framework for Defining and

Forecasting Phytoplankton Blooms'. *Frontiers in Ecology and the Environment* 19 (8): 443–

50. https://doi.org/10.1002/fee.2376.

Kays, Roland, Margaret C. Crofoot, Walter Jetz, and Martin Wikelski. 2015. 'Terrestrial Animal

Tracking as an Eye on Life and Planet'. *Science* 348 (6240): aaa2478.

https://doi.org/10.1126/science.aaa2478.

Kyathanahally, Sreenath P. 2022. 'Plankiformer'. Python.

https://github.com/kspruthviraj/Plankiformer.

Kyathanahally, Sreenath P., Thomas Hardeman, Ewa Merz, Thea Bulas, Marta Reyes, Peter

Isles, Francesco Pomati, and Marco Baity-Jesi. 2021. 'Deep Learning Classification of Lake

Zooplankton'. *Frontiers in Microbiology* 12.

https://www.frontiersin.org/articles/10.3389/fmicb.2021.746297.

Lehtiniemi, Maiju, Elaine Fileman, Heidi Hällfors, Harri Kuosa, Sirpa Lehtinen, Inga Lips, Outi

Setälä, Sanna Suikkanen, Jarno Tuimala, and Claire Widdicombe. 2022. 'Optimising

Sampling Frequency for Monitoring Heterotrophic Protists in a Marine Ecosystem'. *ICES*

Journal of Marine Science 79 (3): 925–36. https://doi.org/10.1093/icesjms/fsab132.

Lewis, Abigail S. L., Christine R. Rollinson, Andrew J. Allyn, Jaime Ashander, Stephanie

Brodie, Cole B. Brookson, Elyssa Collins, et al. 2023. 'The Power of Forecasts to Advance

Ecological Theory'. *Methods in Ecology and Evolution* 14 (3): 746–56.

https://doi.org/10.1111/2041-210X.13955.

Ma, Kevin C. K., Christopher W. McKindsey, and Ladd E. Johnson. 2022. 'Detecting Rare

Species With Passive Sampling Tools: Optimizing the Duration and Frequency of Sampling

for Benthic Taxa'. *Frontiers in Marine Science* 9.

https://www.frontiersin.org/articles/10.3389/fmars.2022.809327.

Marañón, Emilio. 2015. 'Cell Size as a Key Determinant of Phytoplankton Metabolism and

Community Structure'. *Annual Review of Marine Science* 7 (1): 241–64.

https://doi.org/10.1146/annurev-marine-010814-015955.

Marquez, Jonatan, Stefan Vriend, Emily G. Simmonds, Marie V. Henriksen, Lisa Sandal,

Marlène Gamelon, Christophe Coste, Knut Hovstad, and Aline Magdalena Lee. 2022.

'Multispecies Models for Population Dynamics: Progress, Challenges and Future

Directions'. Preprint. Preprints. https://doi.org/10.22541/au.165045606.66517681/v1.

Medeiros, Lucas P., Stefano Allesina, Vasilis Dakos, George Sugihara, and Serguei Saavedra.

2023. 'Ranking Species Based on Sensitivity to Perturbations under Non-Equilibrium

Community Dynamics'. *Ecology Letters* 26 (1): 170–83. https://doi.org/10.1111/ele.14131.

Merkli, Stefanie, Marta Reyes, Christian Ebi, Ewa Merz, Thea Bulas, Stuart R. Dennis, and

Francesco Pomati. 2022. 'Dataset: Greifensee Chlorophyll-a Forecasting Based on Deep

Learning 2019-2021 (Version 1.0)'. Eawag: Swiss Federal Institute of Aquatic Science and

Technology. https://doi.org/10.25678/00066D.

Merz, Ewa, Thea Kozakiewicz, Marta Reyes, Christian Ebi, Peter Isles, Marco Baity-Jesi, Paul

Roberts, et al. 2021. 'Underwater Dual-Magnification Imaging for Automated Lake

Plankton Monitoring'. *Water Research* 203 (September): 117524.

https://doi.org/10.1016/j.watres.2021.117524.

Merz, Ewa, Erik Saberski, Luis J. Gilarranz, Peter D. F. Isles, George Sugihara, Christine

Berger, and Francesco Pomati. 2023. 'Disruption of Ecological Networks in Lakes by

Climate Change and Nutrient Fluctuations'. *Nature Climate Change* 13 (4): 389–96.

https://doi.org/10.1038/s41558-023-01615-6.

Orenstein, Eric C., Devin Ratelle, Christian Briseño-Avena, Melissa L. Carter, Peter J. S. Franks,

Jules S. Jaffe, and Paul L. D. Roberts. 2020. 'The Scripps Plankton Camera System: A

Framework and Platform for in Situ Microscopy'. *Limnology and Oceanography: Methods*

18 (11): 681–95. https://doi.org/10.1002/lom3.10394.

Park, Joseph, Cameron Smith, George Sugihara, and Ethan Deyle. 2021. 'rEDM: Empirical

Dynamic Modeling ("EDM")'. Version 1.13.1. https://CRAN.R-project.org/package=rEDM

Petchey, Owen L., Mikael Pontarp, Thomas M. Massie, Sonia Kéfi, Arpat Ozgul, Maja

Weilenmann, Gian Marco Palamara, et al. 2015. 'The Ecological Forecast Horizon, and

Examples of Its Uses and Determinants'. *Ecology Letters* 18 (7): 597–611.

https://doi.org/10.1111/ele.12443.

Philippart, Catharina J. M., Gerhard C. Cadée, Wim van Raaphorst, and Roel Riegman. 2000.

'Long-Term Phytoplankton-Nutrient Interactions in a Shallow Coastal Sea: Algal

Community Structure, Nutrient Budgets, and Denitrification Potential'. *Limnology and*

Oceanography 45 (1): 131–44. https://doi.org/10.4319/lo.2000.45.1.0131.

Pomati, Francesco, Jukka Jokela, Marco Simona, Mauro Veronesi, and Bas W. Ibelings. 2011.

'An Automated Platform for Phytoplankton Ecology and Aquatic Ecosystem Monitoring'.

Environmental Science & Technology 45 (22): 9658–65. https://doi.org/10.1021/es201934n.

Queiroz, Rose Emília Macedo De, and Renata Gonçalves Ferreira. 2009. 'Sampling Interval for

Measurements of Estuarine Dolphins' (Sotalia Guianensis) Behaviour'. *Marine Biodiversity*

Records 2 (March): e53. https://doi.org/10.1017/S1755267209000700.

Rousso, Benny Zuse, Edoardo Bertone, Rodney Stewart, and David P. Hamilton. 2020. 'A

Systematic Literature Review of Forecasting and Predictive Models for Cyanobacteria

Blooms in Freshwater Lakes'. *Water Research* 182 (September): 115959.

https://doi.org/10.1016/j.watres.2020.115959.

Ruan, S. 2006. 'Delay Differential Equations in Single Species Dynamics'. In *Delay Differential*

Equations and Applications, edited by O. Arino, M.L. Hbid, and E. Ait Dads, 477–517.

NATO Science Series. Dordrecht: Springer Netherlands. https://doi.org/10.1007/1-4020-

3647-7_11.

Sugihara, George, and Robert M. May. 1990. 'Nonlinear Forecasting as a Way of Distinguishing

Chaos from Measurement Error in Time Series'. *Nature* 344 (6268): 734–41.

https://doi.org/10.1038/344734a0.

Sugihara, George, Robert May, Hao Ye, Chih-hao Hsieh, Ethan Deyle, Michael Fogarty, and

Stephan Munch. 2012. 'Detecting Causality in Complex Ecosystems'. *Science* 338 (6106):

496–500. https://doi.org/10.1126/science.1227079.

Takens, Floris. 1981. 'Detecting Strange Attractors in Turbulence'. In *Dynamical Systems and*

Turbulence, Warwick 1980, edited by David Rand and Lai-Sang Young, 366–81. Lecture

Notes in Mathematics. Berlin, Heidelberg: Springer. https://doi.org/10.1007/BFb0091924.

Taylor, Cd, and Bl Howes. 1994. 'Effect of Sampling Frequency on Measurements of Seasonal

Primary Production and Oxygen Status in Near-Shore Coastal Ecosystems'. *Marine Ecology*

Progress Series 108: 193–203. https://doi.org/10.3354/meps108193.

Tulloch, Ayesha I. T., Valerie Hagger, and Aaron C. Greenville. 2020. 'Ecological Forecasts to

Inform Near-Term Management of Threats to Biodiversity'. *Global Change Biology* 26

(10): 5816–28. https://doi.org/10.1111/gcb.15272.

Wauchope, Hannah S., Tatsuya Amano, William J. Sutherland, and Alison Johnston. 2019.

'When Can We Trust Population Trends? A Method for Quantifying the Effects of Sampling

Interval and Duration'. *Methods in Ecology and Evolution* 10 (12): 2067–78.

https://doi.org/10.1111/2041-210X.13302.

Woelmer, Whitney M., R. Quinn Thomas, Mary E. Lofton, Ryan P. McClure, Heather L.

Wander, and Cayelan C. Carey. 2022. 'Near-Term Phytoplankton Forecasts Reveal the

Effects of Model Time Step and Forecast Horizon on Predictability'. *Ecological*

Applications 32 (7): e2642. https://doi.org/10.1002/eap.2642.

Ye, Hao, Richard J. Beamish, Sarah M. Glaser, Sue C. H. Grant, Chih-hao Hsieh, Laura J.

Richards, Jon T. Schnute, and George Sugihara. 2015. 'Equation-Free Mechanistic

Ecosystem Forecasting Using Empirical Dynamic Modeling'. *Proceedings of the National*

Academy of Sciences 112 (13): E1569. https://doi.org/10.1073/pnas.1417063112.

Ye, Hao, and George Sugihara. 2016. 'Information Leverage in Interconnected Ecosystems:

Overcoming the Curse of Dimensionality'. *Science* 353 (6302): 922–25.

https://doi.org/10.1126/science.aag0863.

Yvon-Durocher, Gabriel, Jose Maria Montoya, Mark Trimmer, and Guy Woodward. 2010.

'Warming Alters the Size Spectrum and Shifts the Distribution of Biomass in Aquatic

Ecosystems'. *Global Change Biology* 17 (4): 1681. https://doi.org/10.1111/j.1365-

2486.2010.02321.x.

Figure captions

Figure 1: Schematic view of the field datasets used. The middle column shows the complete

data. This data is manipulated by reducing the sampling frequency (left column) and by reducing

the proportion of kept time points (right column). Each numbered rectangle represents a reduced

dataset. In the left column, the red arrows indicate that the sampling start dates of the

corresponding time series were shifted by one day so that none of the 12 datasets per sampling

frequency used the same time points. In the right column the red arrow indicates by how many

days the start date was shifted so that all time points are used at least once.

Figure 2: The forecast analyses. (a) The relation between sampling frequency and forecast error

for the simulated abundance dynamics with different growth rates (75 time points; 60 individuals

std. dev. noise; dots: mean forecast errors; shaded region: std. dev. of forecast errors). (b) The

relation between growth rate and the optimal sampling frequency for abundance forecasting.

Connected black dots (mean values) and the shaded region (standard deviations) show the results

for the simulations, with parameter values as in (a). The colored points show the relation for the

targets in the field data. (c, d and f) The relations between abundance forecast error (field data)

and the (c) sampling frequency, (d) target maximum net growth rate (complete data), and (f)

proportion of time points used. (e) The relation between target maximum net growth rates and

their median body sizes (complete data). In subfigures (c - f) the dots are colored by target and

the lines are the fixed effects of the corresponding regressions (shaded regions: 95% CI).

Figure 3: The interaction analyses. (a - b) The effect of sampling frequency on the estimated

number of interactions (a) and the mean interaction strengths (b). (c - d) The effect of the

proportion of time points used on the number of interactions (c) and the mean interaction

strengths (d). (e - f) the relation between the target maximum net growth rate (complete data) and

the number of interactions (e) and the mean interaction strengths (f). The dots are colored by

target and in (c) they are jittered horizontally to avoid overplotting. The lines are the fixed effects

of the corresponding regressions (shaded regions: 95% CI), if a significant relation was found.

Figure 4: Estimated slopes between the number of interactions of the targets and their abundance

forecast error (left column) and between their mean interaction strength and forecast error (right

column). (a - b) Reduced sampling frequency analysis. (c - d) Reduced time points analysis. (e -

f) Results reported in Daugaard et al. (2022, constant temperature case). The error-bars denote

95% CIs. The datasets differed in the number of targets (i.e. regression sample size): Lake

Greifen data had 12 targets, and Daugaard et al. (2022) had eight targets with nine replicates.

Figures

Figure 1

 Figure 2

 Figure 3

590   **Figure 4**