GTPase activating protein DLC1 spatio-temporally regulates Rho signaling

Lucien Hinderling[1,2,†], Max Heydasch[1,†], Giliane Rochat[1], Laurent Dubied[1], Jakobus van Unen[1], Maciej Dobrzynski[1], Olivier Pertz[1,*]

[1]Institute of Cell Biology, University of Bern, 3012 Bern, Switzerland

[2]Graduate School for Cellular and Biomedical Sciences, University of Bern, Switzerland

[†] These authors contributed equally to this work

[*] Corresponding author: olivier.pertz@unibe.ch

Abstract

Tightly regulated spatio-temporal Rho GTPase activity patterns regulate morphogenetic processes such as cell migration. Emerging evidence suggests that binding of Guanine nucleotide exchange factors (GEFs) and GTPase activating proteins (GAPs) to the cytoskeleton or adhesions mediate feedback regulation to spatio-temporal Rho GTPase activation. To explore such feedback regulation, we study the Rho specific GAP Deleted in Liver Cancer 1 (DLC1) which binds to focal adhesions (FAs) through mechanosensitive interactions. Using a FRET biosensor, we show that DLC1 loss of function leads to global increase in Rho activity and contractility throughout the cell without affecting a striking lamellar RhoA activity band in fibroblasts. Although RhoA activity may appear steady at the population level, individual molecules are continually activated by GEFs, inactivated by GAPs, and diffusing between subcellular regions, a dynamic turnover referred to as "signaling flux". To directly measure this flux, we built a genetic circuit combining an optogenetic actuator to control Rho activity with a Rho activity biosensor. In spreading cells at steady state, optogenetic manipulation of Rho activity reveals that DLC1 controls the rate of Rho activation rather than duration, both at FAs and at the plasma membrane (PM). Local and reversible optogenetic control of contractility shows that DLC1 associates/dissociates with FAs during their reinforcement/relaxation. This might provide positive feedback that locally increases the rate of Rho activation at FAs that experience local tension to facilitate FA disassembly. Our results indicate that DLC1 operates both at the PM and at FAs to regulate global Rho activity levels at steady state, or to amplify local Rho activity at FAs experiencing a strong mechanical input, presumably to induce robust FA disassembly. This provides new insights in the complexity of spatio-temporal Rho GTPase signaling.

Introduction

Rho GTPases regulate the cytoskeletal dynamics that power morphogenetic processes at the cell- and tissue level (Etienne-Manneville and Hall, 2002). Their activity is tightly controlled in time and space by GEFs and GAPs, which respectively activate and inhibit Rho GTPases. In the GTP-loaded, active state, Rho GTPases bind downstream effectors that regulate cytoskeletal dynamics and other cellular processes. Based on classic experiments that include measuring population-averaged Rho GTPase activity levels, as well as overexpressing dominant positive/negative Rho GTPase mutants, it was proposed that Rac1 controls lamellipodial protrusion, Cdc42 regulates filopodium formation, and RhoA controls myosin-based contractility.

Recent studies in which spatio-temporal Rho GTPase activation dynamics were measured using fluorescence resonance energy transfer (FRET)-based biosensors, revealed a much higher signaling complexity. During cell migration, RhoA, Rac1 and Cdc42 are all activated in membrane protrusions at the leading edge (Itoh et al., 2002; Kraynov et al., 2000; Nalbant et al., 2004; Pertz et al., 2006). Further, the three Rho GTPases are activated in specific spatio-temporal sequences during the protrusion/retraction cycles that fluctuate on time and length scales of seconds and single micrometers (Machacek et al., 2009). A further level of complexity is that these Rho GTPase activity sequences vary depending on the cell system (Hu et al., 2022; Machacek et al., 2009; Martin et al., 2016). This indicates that complex spatio-temporal Rho GTPase signaling programs regulate processes such as cell migration. Similarly as for Ras GTPases (Schmick et al., 2015), it is now clear that these Rho GTPase signaling patterns most likely result from a precise spatio-temporal regulation of Rho GTPase signaling fluxes controlled by GEFs and GAPs (Bement and von Dassow, 2014; Fritz and Pertz, 2016a; Miller and Bement, 2009; Pertz, 2010). In this signaling flux, specific GEFs will be locally activated to load Rho GTPases with GTP depending on different cellular inputs. These GTP-loaded GTPases will then subsequently diffuse in the plasma membrane (PM) by virtue of their C-terminal lipid moiety until they encounter a locally activated GAP which eventually deactivates them. The interplay of GEFs, GAPs and diffusion of Rho GTPases in the PM will ultimately lead to the formation of a Rho GTPase activity pattern. Modelling studies have suggested that sophisticated dynamic signaling behaviors can emerge from the interplay of GEFs, GAPs, and GTPases within a signaling network (Tsyganov et al., 2012). Steady-state images of Rho GTPase activity patterns do not provide information about the whole Rho GTPase flux mentioned above.

This signaling complexity is consistent with the large amount of GEFs and GAPs that are ubiquitously expressed in cells (Fusco et al., 2016, 2016; Moon and Zheng, 2003; Mosaddeghzadeh and Ahmadian, 2021; Müller et al., 2020; Rossman et al., 2005). Recently, a system-wide screen has revealed that many GEFs and GAPS localize to

cytoskeletal structures as well as adhesion complexes such as focal adhesions (FAs)
(Müller et al., 2020). This strongly suggests the existence of feedback mechanisms from
the cytoskeleton and FAs to Rho GTPase signaling. In this study, it was shown that Rac1-
specific GEFs bind to FAs at the front, while Rac1-specific GAPs bind to FAs and at the
back of the lamella of fibroblasts. This precise, asymmetric positioning of GEFs and GAPs
might then regulate the Rho GTPase flux that produces the wide gradient of Rac activity
observed at the leading edge of fibroblasts (Itoh et al., 2002; Kraynov et al., 2000; Martin
et al., 2016). Such feedback regulation might allow leading edge Rac1 activity to constantly
integrate mechanosensitive inputs from FAs, allowing dynamic regulation of Rac1 activity
required to fine tune cell migration. Another prominent example of such spatio-temporal
feedback regulation are the excitable RhoA activity patterns observed in the cortex during
cytokinesis in *Xenopus Laevis* (Bement and von Dassow, 2014), that have also been
observed in mammalian cells (Graessl et al., 2017). Here, RhoA activity waves that control
F-actin wave patterns are spatially regulated by a RhoGAP that is locally regulated by F-
actin (Bement et al., 2015). Thus, cytoskeletal feedback to Rho GTPase regulation might
be crucial to generate dynamic Rho GTPase signaling patterns. Understanding such
spatio-temporal feedback regulation is not accessible with classic genetic approaches in
which long term perturbation of Rho GTPases, their regulators and effectors rapidly result
in a new mechanochemical state of the cell, that will not be informative about an initial
signaling/mechanical state of interest (Isogai and Danuser, 2018). Further, it is currently
unknown which specific features of spatio-temporal Rho GTPase signaling are regulated
to produce signaling patterns. For example, does negative regulation by a GAP solely
control Rho GTPase signal termination, or could it also modulate, less intuitive parameters
such as the rate of activation? Tackling such questions requires new tools to acutely and
transiently perturb the Rho GTPase flux to understand its spatio-temporal regulation,
directly in single living cells. Note that this experimental paradigm was successful at
analyzing the Mitogen activated protein kinase (MAPK network) (Blum et al., 2019;
Dessauges et al., 2022; Ryu et al., 2015).

A promising candidate to investigate how mechanical feedback from FAs can spatio-
temporally shape Rho GTPase activity is the RhoA-specific GAP Deleted in Liver Cancer
1 (DLC1). DLC1 has been shown to both bind the PM (Erlmann et al., 2009), as well as
FAs (Haining et al., 2018; Kaushik et al., 2014). At the PM, interaction with
phosphatidylinositol-4,5-bisphosphate (PI(4,5)P2) can regulate GAP activity (Erlmann et
al., 2009). At the FA, DLC1 interacts with the r7-r8 domains of the FA protein talin in a
mechanosensitive fashion (Gingras et al., 2008; Goult et al., 2021; Haining et al., 2018;
Zhao et al., 2022), as well as with Focal adhesion kinase (Li et al., 2011). When talin is in
a state of low mechanical strain, it binds to DLC1 and presumably leads to local RhoA
inactivation. However, upon force application, the r7-r8 domains unfold leading to
dissociation of DLC1 from talin, leading to loss of negative RhoA regulation. This suggests

that DLC1 is regulated at FAs by mechanosensitive interactions, which can relay
information about the mechanical state of the cell to control RhoA activity.

In this work, we explore the spatio-temporal regulation of RhoA activity by DLC1. We show
that knocking out DLC1 in REF52 fibroblasts leads to increased amplitude in RhoA
throughout the cell. This is accompanied by an increased contractility that augments FAs
and stress fibers (SFs). To explore how DLC1 contributes to Rho GTPase activity fluxes,
we built a genetic circuit consisting of an optogenetic actuator to activate Rho with light,
and a spectrally compatible biosensor to measure Rho activity. Transient optogenetic
recruitment of a Rho GEF domain at FAs or at the PM led to an increase in the rate of
activation of Rho in *DLC1* KO versus WT cells at both subcellular localizations. However,
we found that in both control and DLC1 null cells, RhoA activation was more efficient at
FAs compared to the PM. Further, acute and local manipulation of contractility with the
optogenetic actuator revealed that DLC1 dynamically associates/dissociates with FAs
under acute mechanical tension/relaxation. Our results suggest a complex mechanism in
which DLC1-dependent Rho regulation can occur both at the PM and at FAs, with a
dependence on mechanosensitive signals on the latter.

Results

DLC1 deficient cells show increased formation of SFs and FAs

To investigate the function of DLC1 in cytoskeletal and adhesion dynamics, we knocked
down (KD) DLC1 in rat REF52 fibroblasts via small interfering RNA (siRNA). We also
created a knockout (KO) cell line via Crispr/Cas9, that has a frameshift deletion at position
841 of Exon 5 of the gene, resulting in an early in frame stop codon 100 bp downstream of
the cut. Gene-editing was validated by sequencing the endogenous locus. We were unable
to test expression levels using western blot because of the absence of commercially
available antibodies specific for rat DLC1. We then also rescued the *DLC1* KO cells with
an mCherry-labeled DLC1 construct using stable transfection with a piggyBac transposase
system. REF52 fibroblasts were seeded at subconfluent density onto fibronectin-coated
glass bottom wells and allowed to spread for one hour before fixation, leading to an
isotropic spreading state in control, wild-type (WT) cells. To assess the impact of DLC1
deficiency on the formation of cytoskeletal and adhesion structures, we stained for F-actin
(phalloidin) and paxillin (Figure 1A) or phospho-myosin light chain (pMLC) (Figure 1B).
Consistent with an expected increase in RhoA activity, we observed a phenotype of
increased contractility as documented by increased lamella size, SF content, number of
FAs, and increased pMLC. This increased contractility led to the loss of the isotropic
spreading observed in WT cells. Rescue with the DLC1 construct reverted the cells to a
less contractile phenotype that displayed a lamella of identical size as WT cells, enabling
the characteristic isotropic spreading observed in WT cells. Quantification of these images
revealed both an increase in lamella size (Figure 1C) and FA area (Figure 1D) in DLC1 KD

and KO cells versus WT and rescue cells (Figure1C). As previously documented in
fibroblasts (Kaushik et al., 2014), these results indicate that DLC1 feeds into the regulation
of contractility, SF and FA formation in fibroblasts. The less penetrant phenotype observed
in KO versus KD suggests that the KO cells might be able to adapt to the long-term absence
of DLC1, while KD cells had less than 48 hours to adjust to the new mechanical
state. These experiments also validate the *DLC1* KO cells and *DLC1* KO rescued with
mCherry-DLC1 that will be used in subsequent experiments.

158 DLC1 deficient cells display altered spreading dynamics, lack of polarization
159 and efficient migration

We timelapsed REF52 WT and *DLC1* KO cells for multiple hours in presence of platelet
derived growth factor (PDGF), a potent stimulator of fibroblast motility (Martin et al., 2014),
and evaluated their ability to migrate. We found that *DLC1* KO cells displayed a reduction
in both their speed of migration and their ability to migrate directionally (Figure S1A-C). We
then investigated cytoskeletal dynamics during spreading using REF52 fibroblasts stably
expressing the F-actin marker LifeAct (Riedl et al., 2008). Imaging early spreading (e.g. 20
minutes after plating) revealed no clear differences in edge dynamics in WT versus *DLC1*
KO cells (Figure 2A). However, kymograph analysis clearly showed that the lamella that
characteristically starts directly at the leading edge of the highly contractile REF52 cells
(Martin et al., 2016, 2014), was wider in *DLC1* KO versus WT cells. Later, 1 hour after
plating, when the cells are still spreading, lamellipodial protrusion/retraction cycles of wider
amplitude were observed in *DLC1* KO versus WT cells (Figure 2B, Movie S1), with again
a large increase in SF content (Figure 2B, right panel). After initial spreading, REF52
fibroblasts break their symmetry and eventually display short episodes of polarized cell
migration. Later, contractility builds up leading cells with spindle-like morphology. This
process occurs on timescales of multiple hours. To capture these behaviors, we
investigated cytoskeletal dynamics over the next 10 hours after reseeding at a slower
timescale that precludes observation of lamellipodial dynamics. While WT cells efficiently
broke their symmetry, and could display short stretches of polarized migration, *DLC1* KO
cells displayed robust stress fibers that hampered symmetry breaking and polarization
(Figure 2C, Movie S2). *DLC1* KO cells therefore rapidly adopted a spindle-like shape.
Quantification of the adoption of such a spindle shape phenotype showed that WT cells will
assume the spindle-like shape much later than *DLC1* KO cells and often fail to do so
entirely within the 10 hours of imaging (Figure 2D). Further highlighting a role for DLC1 in
RhoA-mediated control of contractility, we found that DLC1 overexpression leads to strong
reduction in the number of SFs (Figure S2A). Together, these results show that DLC1 loss
of function leads to increased contractility at the onset of spreading, which later impedes
polarization and directional cell migration.

DLC1 deficient cells display an increase in global RhoA activity, without any marked differences in RhoA activity pattern

To explore how DLC1 spatio-temporally controls RhoA activity, we used RhoA2G (Fritz et al., 2013), a fluorescence resonance energy transfer (FRET)-based biosensor that reports on RhoA activity together with the Lifeact F-actin marker during spreading. As previously observed (Martin et al., 2014), a RhoA activity band that correlates with the lamellar contractile actomyosin network of the cell can be observed at the cell edge of WT REF52 fibroblast (Figure 3A, Movie S3). In *DLC1* KO cells, we observed global elevation of RhoA activity throughout the cell. Quantification of both the average activity per cell and the local edge RhoA activity revealed increased RhoA activity (Figure 3B). Kymograph analysis revealed that the lamellar RhoA activity band remained of constant size during spreading (Figure 3A). Examining cells approximately 10 hours after spreading when they had fully spread, and WT cells had adopted a more contractile state, the DLC1 *KO* cells still exhibited a global increase in RhoA activity (Figure 3C,D). This increase was however less strong than in spreading cells, most likely because of a higher state of contractility in WT well-spread versus spreading cells. We also evaluated the effect of overexpression of an mCherry-tagged DLC1 on RhoA activity (Figure S2B). We observed that mild DLC1 expression, that remains mostly bound to FAs and to the PM and does not compromise cell morphology, leads to a decrease in global RhoA activity. Strong DLC1 expression that leads to a large cytosolic pool of DLC1 further diminishes global RhoA activity but then compromises cell morphology. Together, these results suggest that DLC1 signaling at the PM and at FAs controls global levels of RhoA activity within cells, rather than controlling the spatio-temporal RhoA activity pool observed at the lamella.

Design of an optogenetic actuator - Rho biosensor to interrogate the Rho GTPase flux

While FRET measurements as shown above provide insight about a steady-state RhoA activity pattern, they cannot probe in detail how a GEF/GAP/GTPase signaling network might affect the Rho GTPase flux which most likely emerges from the regulation of multiple GEFs and GAPs. For example, GAP-mediated negative regulation might control Rho GTPase signal duration, but might also be involved in adjusting the rate of activation. We reasoned that transient, acute optogenetic perturbations of the Rho GTPase flux, followed by measuring Rho activity dynamics might provide new insights about its regulation. To manipulate Rho activity in single living cells, we re-engineered an iLID-based optogenetic actuator based on a DH/PH domain of the GEF LARG (Oakes et al., 2017) (Figure 4A). We used a high affinity nano SspB domain for efficient light-dependent recruitment of the LARG GEF domain to the iLID anchor. To better focus optogenetic activation, we also fused the iLID anchor to a stargazin membrane anchor that slows down its diffusion (Natwick and Collins, 2021). Both the stargazin-iLID anchor and the SspB-LARG domains were

separated by a P2A self-cleaving peptide, allowing for equimolar expression of both units
from a single operon. We refer to this construct as optoLARG. To interrogate the Rho
GTPase flux, we engineered a REF52 line that stably expressed the optogenetic construct
as well as a rhotekin-based G protein binding domain (rGBD) that reports on Rho activation
(Mahlandt et al., 2021). The rGBD probe is labelled with a tandem dimeric Tomato
(tdTomato) fluorophore that is spectrally orthogonal to optogenetic activation. We used a
truncated CMV promoter to warrant low expression level of the rGBD probe to avoid
dominant negative effects of the construct through inhibition of Rho signaling. We also
expressed a spectrally orthogonal far-red Lifeact-miRFP construct that reports on F-actin
dynamics. Because rGBD cannot distinguish between the activity of the RhoA, RhoB and
RhoC isoforms, we refer to any rGBD measurement as Rho activity. This is in marked
contrast with RhoA2G, the FRET biosensor we used above that is specific for RhoA (Fritz
et al., 2013), but is not spectrally compatible with optoLARG. Note that both biosensors are
able to measure active Rho GTPase pools during robust morphogenetic events such as
tail retraction (Fritz et al., 2013; Mahlandt et al., 2021), while rGBD misses leading edge
Rho GTPase activity pools widely observed using FRET probes (Machacek et al., 2009;
Martin et al., 2016; Pertz et al., 2006).

We then used a digital micromirror device (DMD) to spatially shine repetitive pulses of blue
light delivered at different frequencies on 2 distinct regions of interest (ROIs) at the edge
of fibroblasts. When light pulses were delivered at high frequency, we observed robust Rho
activation at the edge, which correlated with assembly of contractile F-actin structures and
potent edge retraction (Figure 4B, kymograph shown in 4C, Movie S4). Removal of the
blue light pulses led to a subsequent decrease in Rho activity, and reversion of the edge
to a protrusive state. Further, stimulating a ROI at the opposite edge of the cell with a lower
light pulse frequency led to a lower Rho activation, and less robust edge retraction. These
results indicate that depending on the strength of the light input, optoLARG can locally,
reversibly, and quantitatively control Rho activity in single living cells on timescales of
seconds, allowing us to acutely manipulate the Rho GTPase flux.

Optogenetic interrogation of the Rho GTPase flux in WT and *DLC1* KO cells

Because DLC1 might both regulate the Rho GTPase flux both at the PM (Erlmann et al.,
2009) and at FAs (Haining et al., 2018), we performed a series of experiments using
optoLARG to transiently activate RhoA with the expectation of observing different
perturbations in Rho activity fluxes in control versus *DLC1* KO cells. For that purpose, we
produced WT and *DLC1* KO REF52 lines expressing optoLARG, the tdTomato-rGBD
biosensor and miRFP-paxillin as a readout for FAs. We first sought to locally activate RhoA
by illuminating large ROIs at edges and evaluate any difference in Rho activity patterns
using the spectrally compatible rGBD biosensor in WT versus *DLC1* KO cells. We however
were not able to identify a threshold optogenetic light input that would induce different RhoA
activity patterns and edge dynamics in WT versus the more contractile *DLC1* KO cells. We

observed that the same light input could induce edge retraction in both WT and *DLC1* KO
cells in a non-predictable manner. This might be due to the spatial heterogeneity of the
mechanical states within cells, as well as small differences in optoLARG and rGBD
expression levels in the stable cell lines. To address this caveat, we reasoned that we
should activate Rho using optoLARG with a more subtle modality by illuminating only small
ROIs and measuring small local fluctuations of Rho activity using rGBD without inducing
large changes in mechanical states. Further, to dissect potential regulation of *DLC1* GAP
activity by PI(4,5)P2 at the PM (Erlmann et al., 2009) versus mechanosensitive interactions
at the FA (Haining et al., 2018), we decided to make measurements on FAs but use ROIs
outside of FAs (non-FAs ROIs) to probe the PM. For that purpose, we built an image
analysis pipeline to first segment a cell in a field of view and to then segment FAs based
on their fluorescent intensity, size and localization at the cell periphery using the paxillin-
miRFP signal (Figure 5A). We then used these FA segmentations to illuminate small ROIs
within FAs or in neighboring non-FA ROIs with one pulse of blue light using the DMD and
evaluated rGBD fluorescence fluctuations at this location for a couple of minutes until Rho
activity returned to baseline (Figure 5B). This optogenetic stimulation had no visible effect
on edge retraction, or FA morphodynamics, as observed when stimulating larger ROIs.
Quality control of our automated segmentation revealed high versus low paxillin signals in
FA versus non-FA ROIs (Figure 5C). This experiment was performed on multiple cells,
allowing to average out any experimental noise due to small differences in expression
levels of optoLARG or rGBD. These data were then averaged for FA and non-FA ROIs.
We observed that this pulsed optoLARG light input led to a local transient of Rho activity
for approximately 2.5 minutes in WT cells when FAs or the PM were stimulated (Figure
5D). To quantify observed activation ($k_{\mathrm{ON}}^{\mathrm{obs.}}$) and deactivation ($k_{\mathrm{OFF}}^{\mathrm{obs.}}$) rates, we defined a 10-
90% time window of the activation and deactivation phases and computed the mean
derivative over each window. The identical optoLARG input led to an increased observed
rate of Rho activation leading to higher amplitude when an FA versus a non-FA ROI was
stimulated ($k_{\mathrm{ON}}^{\mathrm{obs.}}$: 0.0132 (non-FA) < 0.0176 (FA)). In *DLC1* KO cells, we observed
increased Rho activation rate that also led to augmented Rho activity amplitude in
comparison with WT cells in both FA and non-FA ROIs ($k_{\mathrm{ON}}^{\mathrm{obs.}}$: 0.0195 (non-FA) < 0.0221
(FA)). This was also accompanied with faster deactivation kinetics of RhoA ($k_{\mathrm{OFF}}^{\mathrm{obs.}}$: 0.0060
(non-FA KO) < 0.0068 (FA KO), $k_{\mathrm{OFF}}^{\mathrm{obs.}}$: 0.0042 (non-FA WT) < 0.0054 (FA WT)). These data
suggest that under weak transient GEF input, DLC1 regulates the rate of Rho activation
and peak amplitude rather than signal duration. Note that $k_{\mathrm{ON}}^{\mathrm{obs.}}$ and $k_{\mathrm{OFF}}^{\mathrm{obs.}}$ here refer to the
rates observed in the biosensor signal, which reflect the combined effects of underlying
activation and deactivation processes rather than parameters of GEF/GAP activity. This
can occur both at the PM and at FAs. The finding that *DLC1* KO also leads to faster
deactivation suggests the existence of compensatory negative feedback that points to
complex modalities of spatio-temporal regulation that most likely involves additional
GEFs/GAPs. In both WT and *DLC1* KO cell lines, RhoA activation rates and peak levels

were higher at FAs than in non-FA regions. To exclude that this difference could be an artefact from differential recruitment of the optogenetic tool, we directly measured optoLARG localization using a cell line expressing mScarlet3-optoLARG and miRFP-paxillin, which allowed imaging of optoLARG recruitment without activation. We repeated the experiment from Figure 5, this time quantifying optoLARG intensity instead of RhoA activity, and found no significant difference in recruitment kinetics between FAs and non-FAs (Figure S3A,B), excluding tool recruitment as a confounding factor. These results instead suggest that FAs provide a subcellular environment that favors GEF-mediated activation of Rho activity compared to non-FA regions. The recruitment and dissociation dynamics of optoLARG were well described by an inverse exponential function during association and an exponential decay function during dissociation (Figure S3C). The measured deactivation half-life was 10.8 s, which is faster than previously reported values 2016) (t½ OFF = 18 ± 2 s in vitro; 32.0 ± 3.7 s for membrane localization in cells) (Hallett et al.). Our experimental design differs from previous reports in that activation was restricted to small regions of the cell, whereas prior studies used whole-cell illumination. The accelerated decay we observed may therefor result from additional diffusion of optoLARG out of the locally stimulated regions.

The experimental data revealed distinct activation and deactivation kinetics between WT and *DLC1* KO cells, suggesting altered regulatory feedback within the RhoA signaling network. We next sought to investigate the mechanistic basis of these differences by mathematical modeling.

Modeling suggests loss of negative RhoA autoregulation in *DLC1* KO cells

To quantitatively describe the regulatory dynamics of RhoA signaling following transient, localized optogenetic stimulation at FAs and the PM in WT and *DLC1* KO cell lines, we constructed a system of ordinary differential equations (scheme in Figure 6A). This modeling framework allows us to disentangle the underlying activation and deactivation rates from the apparent rates observed with the biosensor and infer qualitative regulatory differences between WT and KO cells. The experimental measurements of optoLARG recruitment were first used to fit a simple linear model of protein activation and deactivation, enabling estimation of the kinetic parameters governing optoLARG dynamics. Based on this fit, the active GEF profile (GEF^*) was treated as a fixed, time-dependent input to the downstream model of RhoA activity. The remaining regulatory interactions, including RhoA activation by GEFs, RhoA inactivation by GAPs, and RhoA-dependent GAP activation, are captured in the following system using loosely constrained parameters:

$$\frac{d\mathrm{LARG}^*}{dt} \approx k_{\mathrm{on}}^{\mathrm{LARG}} \cdot f_{\mathrm{signal}}(t) - k_{\mathrm{off}}^{\mathrm{LARG}} \cdot \mathrm{LARG}^*$$

$$\frac{d\mathrm{GAP}^*}{dt} \approx \frac{V_{\mathrm{on}}^{\mathrm{GAP}} \cdot \mathrm{Rho}^*}{Km^{\mathrm{GAP}} + \mathrm{Rho}^*} - k_{\mathrm{off}}^{\mathrm{GAP}} \cdot \mathrm{GAP}^*$$

$$341 \qquad \frac{d\text{Rho}^*}{dt} \approx \frac{V_{\text{on}}^{\text{Rho}} \cdot \text{LARG}^*}{Km_{\text{on}}^{\text{Rho}} + \text{LARG}^*} - \frac{V_{\text{off}}^{\text{Rho}} \cdot \text{GAP}^* \cdot \text{Rho}^*}{Km_{\text{on}}^{\text{Rho}} + \text{Rho}^*}$$

$$342 \qquad k_{\text{on}}^{\text{LARG}} = 0.0123462$$

$$343 \qquad k_{\text{off}}^{\text{LARG}} = 0.0700499$$

Michaelis-Menten kinetics were used to model RhoA activation by active optoLARG

(LARG^*), inactivation by active GAPs (GAP^*), and a RhoA-dependent feedback on GAP

activation (Rho^*) (De Seze et al., 2023). GAP inactivation was approximated as a first-order

process. Given the transient and low-amplitude nature of the optogenetic stimulus, we

assumed that the inactive forms of optoLARG, GAP, and Rho remained in large excess

throughout the simulation. This allowed us to simplify the model by treating the

concentrations of the inactive species as effectively constant over time. The resulting

system of ODEs was solved and fitted to the experimental data using gradient-based

optimization in Julia (Figure 6B,C). We additionally assessed the structural identifiability of

all parameters. Full implementation details and the results of the structural identifiability

analysis are provided in the Methods section.

In WT cells, the model provides a strong fit to the experimental data and supports the use

of canonical Michaelis-Menten kinetics for RhoA activation and inactivation, as well as the

negative feedback on GAP. The results indicate that RhoA activity is dynamically regulated

by active GAPs, which are themselves activated by RhoA, forming a negative feedback

loop. This feedback ensures that RhoA activation remains transient and self-limiting. The

fitted parameters reflect saturable enzyme-substrate interactions, consistent with standard

biochemical expectations.

In contrast, the behavior observed in *DLC1* KO cells reveals a disruption of this feedback

mechanism. When fitted with the same model structure, the parameter for RhoA-dependent

GAP activation converges to zero, indicating loss of negative feedback. Instead, the system

appears to rely on a constitutively active or pre-localized GAP-like component that

inactivates RhoA at a fixed rate. This configuration eliminates feedback modulation and

produces RhoA dynamics that are governed by constitutive GAP activity, possibly due to

the global GAP activity of multiple RhoA-specific GAPs in the cell.

A limitation we encountered in our modeling is that the fitted parameter magnitudes

happened to depend on the bounds imposed during optimization. Structural identifiability

analysis confirmed that all parameters are theoretically globally identifiable given the model

structure and observables. Yet in practice, some fitted values were found to vary with the

parameter bounds chosen. While some parameters (e.g., responsible for RhoA activation

or RhoA-dependent GAP activation) were robust across fits, others varied without

significantly affecting model performance. This reflects the distinction between structural

and practical identifiability (Gutenkunst et al., 2007; Lam et al., 2022) where the robustness

and uniqueness of fitted parameter can be limited by the experimental data. We therefore
limited our attention on qualitative differences in regulation (feedback present vs. absent),
rather than absolute numerical parameter values.

DLC1 differently interact with FAs depending on their mechanical states

Our finding that we did not observe striking patterns of RhoA activity in the vicinity of FAs
in spread cells challenges the idea that mechanical inputs regulate Rho activity in this
specific regime of mechanical forces. We speculated that the DLC1/talin system in FAs is
only mechanosensitive to stronger mechanical inputs. To evaluate FA-DLC1 interactions,
we imaged fibroblasts rescued with mCherry-DLC1 and miRFP-paxillin as a marker for
FAs. We observed that DLC1 binds to FAs at the onset of FA assembly, and that the local
DLC1 density augmented with the paxillin density (Figure S4A-C). During FA disassembly,
DLC1 density diminished concomitantly with decreasing paxillin density until the FA was
fully disassembled (Figure S4D-F).

To characterize force-dependent interactions of FAs with DLC1 in regimes of strong
mechanical perturbations, we engineered our mCherry-*DLC1* rescue cells to stably express
the optoLARG system, as well as the miRFP-paxillin construct. We then used the DMD to
activate Rho-mediated contractility locally and transiently in small ROIs containing FAs with
a standardized light input. The transient light pulse induced either of the two FA behaviors:
light-mediated FA reinforcement as evidenced by a local increase in paxillin density, was
followed by 1. FA relaxation (and decrease in paxillin density) upon light removal (Figure
7A-C, quantified in Figure 7D, Movie S5), or 2. FA rupture (Figure S5A-C). This was
accompanied by a striking simultaneous decrease in DLC1 intensity at the FA during FA
reinforcement, and a reassociation of DLC1 during FA relaxation upon light input removal.
If the light input led to FA rupture, then the DLC1 signal gradually decreased along FA
reinforcement until the FA ultimately ruptured in an all-or-nothing manner. FAs in the same
cell that were not subjected to the light input displayed dynamics as in normal cells, as
shown in Figure 7 (Figure S5D-F). These results indicate that DLC1 dynamically
dissociates and associates during acute FA reinforcement and relaxation in response to a
strong mechanical input.

Discussion

A key question in Rho GTPase biology is how multiple GEFs and GAPs control spatio-
temporal Rho GTPase signaling patterns to regulate cytoskeletal dynamics that power
morphogenetic processes such as cell migration. The recent finding that many GEFs and
GAPs themselves bind cytoskeletal and adhesion structures (Müller et al., 2020) strongly
suggests direct feedback from the cytoskeleton to Rho GTPase signaling. Self-
organizational properties of cytoskeletal and adhesions structures might therefore
contribute to spatio-temporal control of Rho GTPase signaling. Because DLC1 has been

proposed to bind to FAs in a mechanosensitive manner (Haining et al., 2018), we decided
to study it as a prototypical example of an adhesion feedback to Rho GTPase signaling.
Using novel technologies that allow for acute spatio-temporal perturbations, we provide
some new insights about how Rho GTPases are spatio-temporally controlled by a RhoGAP
that integrates mechanosensitive inputs.

We found that in spreading fibroblasts, *DLC1* KO globally augments contractility, as
illustrated by a robust increase in SFs and FAs (Figure 1). Evaluation of F-actin dynamics
immediately after spreading, when global contractility levels are still low, reveal that *DLC1*
KO fibroblasts rapidly assemble a wider contractile lamella than WT cells, suggesting
aberrant increase of myosin-based contractility already at an early stage of spreading
(Figure 2A). One hour after cell plating, when cells are well spread but still retain an
isotropic shape, we observed a global increase in contractility in *DLC1* KO versus WT cells
(Figure 3B). On a timescale of multiple hours after spreading, when WT cells break
symmetry, display polarized cell migration episodes, and finally adopt a contractile state
leading to a spindle morphology, we find that *DLC1* KO cells transition directly to the
contractile phenotype without being able to polarize (Figure 2C,D). Thus, DLC1 loss of
function leads to a global increase of aberrant contractility.

We had previously shown that REF52 fibroblasts display a wide band of RhoA activity that
correlates with the lamellar contractile myosin network during spreading (Martin et al.,
2016, 2014). We found that *DLC1* KO leads to a global increase in RhoA activity throughout
the cell, without however markedly modifying the spatial band of RhoA activity at the
lamellum (Figure 3A,B). Similar results were observed when cells were allowed to further
polarize and assemble a robust actomyosin cytoskeleton (Figure 3C,D). Further, we
observed that mild DLC1 expression, that can still mostly be titrated by FAs and the PM
without spilling out into the cytosol, is already able to strongly downregulate global RhoA
activity levels (Figure S2B). These findings suggest that in spread cells, DLC1 mainly sets
the overall baseline of RhoA activity, while the spatial distribution of RhoA activation, such
as observed at the lamellar band observed in spreading cells, appears to be maintained by
other, spatially restricted GEFs/GAPs that control localized RhoA pools at protrusive edges
and filopodia (Fritz and Pertz, 2016b; Machacek et al., 2009; Pertz et al., 2006), within the
lamella (Martin et al., 2016, 2014). Similarly, additional GEFs/GAPs might be specifically
spatio-temporally regulating the cytokinetic furrow (Basant and Glotzer, 2018; Bement et
al., 2015). Note that the classic genetic perturbation paradigm is only of limited value to
study the dynamic regulation of Rho by DLC1 at FAs because long-term loss of DLC1
function results in cells being locked in a state of excessive contractility.

To address this question, we used optogenetic control of Rho activation to probe different
aspects of DLC1-mediated regulation of Rho activity at FAs, sites where DLC1 is highly
enriched (Figure 7), and regulates Rho through talin- and tensin-dependent interactions,
as well as at control regions on the PM, where DLC1 can bind lipids but is less abundant

(Erlmann et al., 2009; Haining et al., 2018). In a first experimental modality, we evaluated
the Rho activity flux at FAs as well as the PM in a mechanical regime that does not involve
FA reinforcement (Figure 5). In the 2nd experimental modality, we evaluated force-
dependent DLC1-FA interaction in a mechanical regime in which FA reinforcement is
clearly occurring (Figure 5).

In the first experimental modality, we probed the Rho GTPase flux under conditions that
did not trigger FA reinforcement (Figure 5D). The goal was to probe DLC1-dependent
regulation of Rho in absence of strong mechanical strain. To achieve this, we applied small
light inputs that induced only subtle fluctuations in Rho activity. Due to the weak signal-to-
noise ratio (SNR) of these subtle GEF inputs, meaningful measurements of Rho activity
response dynamics could only be obtained by averaging signals across hundreds of FAs.
We observed that a transient synthetic optoLARG input applied to a FA yielded a Rho
activation transient indicating the presence of tonic Rho GAP activity in the cell (Figure
5E,F). Interestingly, we observed that the identical synthetic optoLARG input applied to the
PM in absence of FA led to a Rho activity transient of lower amplitude. This suggests that
FAs are intrinsically more permissive for GEF-mediated Rho activation, potentially due to
local Src-dependent release of Rho from RhoGDI (DerMardirossian et al., 2006).

Since Rho activity may be regulated by DLC1 at FAs, we initially expected to observe
altered Rho dynamics at these sites in *DLC1* KO cells, specifically that transient optoLARG
input would lead to prolonged Rho activity. We found however that RhoA activation
occurred faster, reached a higher peak amplitude, and returned to baseline more rapidly in
both FA and non-FA PM regions compared to control cells (Figure 5E,F). Given that DLC1
is strongly enriched at FAs relative to the PM, the observation that similar RhoA flux
changes occur at both subcellular locations suggests that these effects do not arise solely
from local DLC1 activity. Rather, they may also reflect compensatory expression or
activation of other GEFs and GAPs triggered by DLC1 loss. This interpretation is consistent
with our experimental observation that *DLC1* KO cells engage compensatory mechanisms
over time, resulting in weaker phenotypic differences than those observed in acute
knockdown (Figure 1). However, such compensation likely lacks spatial specificity, as it
does not adapt to local RhoA concentrations. This behavior mirrors findings in T cells,
where Cdc42 deletion triggered compensatory transcriptional programs that restored
cellular function (Rochussen et al., 2025), illustrating how Rho GTPase networks can
rapidly adapt to gene loss and thereby mask the impact of individual regulators.

Our mechanistic model of dynamic RhoA modulation, fitted to experimental measurements
of GEF input and resulting RhoA activity, further supports this view. The model revealed
qualitative differences between WT and *DLC1* KO cells: WT cells exhibited Rho-dependent
negative feedback that dampens activation dynamics, whereas this feedback was lost in
the KO condition. The absence of this negative autoregulation explains the higher Rho
activation amplitude observed in KO cells and implies that, in the absence of DLC1,

alternative Rho-regulated GAPs or regulatory proteins act independently of RhoA levels.
Such negative feedback architectures are a hallmark of small GTPase signaling networks
(Bement et al., 2024; Tsyganov et al., 2012; Wu and Lew, 2013), where they confer
robustness and dynamic stability to signaling outputs. Altogether, our results uncover an
unexpected complexity in the spatiotemporal regulation of Rho GTPase signaling at FAs,
likely involving multiple GEFs and GAPs. They also highlight how long-term genetic
perturbations can activate compensatory mechanisms, complicating the interpretation of
single-gene knockouts. Future experiments will require high-resolution, dynamic
measurements of multiple GEFs, GAPs, and RhoA activity in unperturbed cells, or
alternatively, acute perturbations that precede the onset of compensatory responses.

In the second experimental modality, we investigated mechanosensitive interactions
between FAs and DLC1 under optoLARG stimulation conditions that promote robust FA
reinforcement. We transiently increased actomyosin contractility within a small ROI, which
triggered FA reinforcement followed by relaxation (Figure 7). Under such acute mechanical
perturbations, we observed distinct DLC1 dynamics at FAs, likely involving its known
mechanosensitive interactions with talin (Haining et al., 2018). In response to the induced
strain, DLC1 rapidly dissociated from FAs exhibiting strong reinforcement (Figure 7;
schematic in Figure 7F). This behavior correlates with in vitro findings showing that DLC1
dissociates from stretched talin due to unfolding of its R7-R8 domain (Haining et al., 2018).
We also observed that FA relaxation promotes re-binding of DLC1 to FAs (Figure 7),
suggesting that the R7-R8 talin domain might refold in these conditions. In this context,
strong local mechanical input leading to FA reinforcement and talin stretching may cause
rapid unbinding of DLC1 from FAs, transiently reducing local RhoGAP activity and
providing a positive feedback loop that increases RhoA activation and local contractility.
This may, in turn, promote FA disassembly as observed experimentally. Conversely, when
the mechanical input ceases, DLC1 re-binding to relaxing FAs may rapidly suppress RhoA
activity, restoring a low-contractility steady state. These nonlinear feedback mechanisms
may enable amplification of local mechanical inputs to regulate all-or-nothing cytoskeletal
events such as tail retraction during migration. Tail retraction is characterized by brief bursts
of localized RhoA activity followed by rapid relaxation once retraction is complete (Martin
et al., 2014; Pertz et al., 2006). In this model, stress fiber pulling on FAs at the cell rear
could trigger local DLC1 release, transiently elevate RhoA activity, and drive strong
localized contraction leading to tail retraction. Testing this hypothesis experimentally is
challenging, as long-term DLC1 perturbation will induce global hypercontractility,
preventing cell polarization, directional migration and thus the production of tail retraction
events.

Summarizing the results from these two experimental regimes, we propose that DLC1
provides two distinct levels of Rho regulation. Under low mechanical input, DLC1 may help
control global RhoA activity by positioning localized hotspots of RhoGAP activity throughout

the cell, most likely together with other Rho GAPs. In contrast, under strong mechanical
input leading to FA reinforcement, the DLC1-talin mechanosensitive interaction may permit
rapid, local positive feedback on RhoA activity through DLC1 dissociation from FAs. This
positive feedback could enable efficient FA turnover in response to mechanical stimuli and
regulate processes such as tail retraction.

Our results illustrate how spatially controlled optogenetics with optoLARG enables
spatiotemporal, reversible control of contractility, revealing mechanosensitive signaling
interactions that are otherwise obscured by classical global perturbations of RhoA, such as
nocodazole-induced microtubule depolymerization (Krendel et al., 2002) or
lysophosphatidic acid-mediated RhoA activation (Pertz et al., 2006) which trigger broad
and nonspecific contractile responses. These findings provide new insight into the network
circuitry governing spatiotemporal Rho GTPase signaling and point toward a new
experimental modality for dissecting these mechanisms. By combining precise, reversible,
and spatially confined optogenetic perturbations with automated optogenetic targeting, we
can now locally modulate Rho activity and probe signaling fluxes across different regimes,
using gentle perturbations to nudge the system out of equilibrium and reveal its relaxation
back to steady state, or stronger inputs to engage mechanical feedback and adhesion
reinforcement. Although this study focused on a GAP localized to FAs, the same approach
can be extended to other cytoskeletal structures by targeting actin, myosin, or microtubule
markers to guide light stimulation in real time. Such strategies will allow systematic mapping
of how localized biochemical and mechanical feedback, mediated by multiple GEFs and
GAPs, combine to generate emergent signaling patterns. Future studies systematically
perturbing individual and combined GEFs and GAPs under controlled optogenetic
stimulation will be essential to identify their specific contributions to Rho GTPase flux and
feedback architecture. Ultimately, achieving this level of quantitative, structure-specific
control will be key to understanding how dynamic Rho GTPase signaling under the control
of multiple GEFs and GAPs orchestrates cell morphogenesis.

558 Materials and methods

559 Cell culture

560 Rat Embryo Fibroblast 52 (REF52) were cultured in Dubelcco's Modified Eagle Medium
561 with 4.5 g/l glucose, 10% 4mM L-Glutamine, and 100 U/ml penicillin/streptomycin. Cells
562 were grown at 37 °C and 5% CO_2.

564 SiRNA knockdown

4µl lipofectamine RNAi max was mixed with a total of 500µl of OPTIMEM medium and a
pool of SiRNAs against the gene of interest at a final concentration of 60 nM (SiTools

Biotech). The mix was added to a 6 well plate containing 100000 cells in a 2 ml medium
volume medium.

CrispR/Cas9 Knockout generation

Two guide sequences targeting DNA within exon 5 of the DLC1 gene were selected using
the CRISPOR tool for predicted high-specificity protospacer adjacent motif target sites in
the rat genome [32] . Two complementary oligos each containing the DLC1 guide sequence
and Bbsl ligation adapters were synthesized (Microsynth) (sense sequence: 5'-
CACCGAACCGAGAGAGCTACCCGG-3', antisense sequence: 5'-
AAACCCGGGTAGCTCTCTCGGTTC-3'). The guide sequences were annealed and
ligated into a pSpCas9(BB)-2A-GFP vector. REF52 cells were grown in six-well plates to
60% confluency and transfected with 1µg of the vector together with 1 µl of Lipofectamine
3000 and 5 µl of P3000 solution (ThermoFisher Scientific). Two days post transfection cells
were detached, suspended in PBS +1% FBS and sorted into 96 well-plates using
fluorescence-activated cell sorting. After expansion, individual clones were detached and
suspended in the lysis buffer (10mM TRIS, 50mM KCL, 2.5mM MgCl2, 0.45% Tween-20,
0.05% Gelatine, 0.12 mg/ml of proteinase K). The cell solution was frozen at -80 °C and
subsequently kept at 63 °C for one hour. Proteinase K was inactivated by heating the
solution to 95 °C for 15 minutes. PCR primers were designed to amplify a 500bp region
around the CRISPR cut site (forward primer: 5'-
AAGGAGTGTGTCTAACTCCACGCAGACCAG-3', reverse primer: 5'-
CTCCTTAGGACTGTCGCTGCTGTTTTCTCT-3'). Genomic sequences were amplified by
PCR and sequenced by Sanger sequencing.

DNA constructs

A "pB3.0" piggybac vector was created by adapting the pPBbsr2 (Matasci et al., 2011) to
make it smaller in size. pPBbsr2 was digested with AscI and PacI, yielding 3922 bp and
2864 bp fragments. Full size pPBbsr2 was used as template for a PCR with forward primer
5'-TTAGCATTAATTAAGCGGCCGCGTTGCTGGCGTTTTTCC-3' and reverse primer 5'-
GTGCCTTTACAACTTATGAGTAACCCCGCGCGGACGATT-3', yielding a 1671bp insert
that was ligated back into the receiving 3922bp part of the digested pPBbsr2 vector,
creating "pB3.0-BLAST" (Blasticidine resistance). To introduce different resistance
cassettes, we performed an overlap extension PCR with pB3.0-BLAST as template for
PCR1 (primers A+B) and PCR2 (primers C+D) with the following primers:

primer A: 5'-AAGGATGCCCAGAAGGTACCCCATTGTATGGGATCTGATCTGGG-3',
primer B: 5'-GGAAACTTTTTGTGCTATTATGGTGGCCATTCAGCTCTACGTAGCTACT-

3', primer C 5'-
CCTTTGAAAAACACGATAATACCACCGGTAAGTCGAGATGCATCGATGA-3', primer D:
5'-CTCCGCCTTTCTTGGACGTCGGGTTCGAACCGCATTAGT-3'. Primers A and D, the
product of PCR1 and PCR2 were used for fusing the two fragments by PCR. The resulting
PCR product was ligated into the pB3.0-BLAST vector, digested with Kpn1 and Pst1, by
fusion cloning, yielding a pB3.0-noAB vector with a single Age1 site to introduce antibiotic
resistance cassettes. To produce pB3.0-HYGRO and pB3.0-PURO, PCRs were performed
on template vectors pHygro-PB and pPuro-PB, respectively. For pB3.0-HYGRO forward
primer 5'-AACACGATAATACCACCATGAAAAAGCCTGAACTCACCGC-3' and reverse
primer
5'-GCAGGCTCCCGTTTCCTTATCGGCCATTCAGCTCTA-3' were used. For pB3.0-
PURO forward primer 5'-AACACGATAATACCACCATGACCGAGTACAAGCCCACG-3'
and reverse primer 5'-GCGTTCGGGCCACGGACTGGCCATTCAGCTCTA-3' were used.
The respective inserts were inserted via fusion cloning in the pB3.0-noAB vector [33],
digested with AgeI.

Construction of the pB3.0-optoLARG-mVenus-SspB-p2A-stargazin-mtq2-iLID optoLARG
system was as follows. The cDNA for optoLARG-mVenus-SspB(nano)-p2A-iLID-CAAX
was synthesized by custom gene synthesis (Genewiz, Azenta life sciences, US). The
sequence contained the catalytical DH-PH domain of human RhoGEF12 (LARG, aa 766-
1138), connected via a linker (GSGSGSGS) to full-length mVenus fused to SSPB(nano).
This is followed by a p2A sequence (GSGATNFSLLKQAGDVEENPGP), the iLID module
(Guntas et al., 2015) and a CAAX box (KRAS). The construct was cloned into pB3.0-BLAST
containing a CAG promotor, yielding pB3.0-optoLARG-mVenus-SspB-p2A-iLID-CAAX.
After publication of an improved PM anchor for the iLID-module (Natwick and Collins,
2021), we changed the part after the p2A sequence. The pB3.0-optoLARG-mVenus-SspB-
p2A-iLID-CAAX vector was digested with BsrGI and AflII, and a custom gene synthesized
stretch of cDNA containing SspB(nano)-p2A, full-length stargazin, mTurqoise2 and iLID
was inserted. This yielded pB3.0-optoLARG-mVenus-SspB(nano)-p2A-stargazin-mtq2-
iLID (exact sequences of the gene synthesis products are available on request). We also
developed a version with the recruitable SSpB domain tagged in red (mScarlet3) instead
of yellow (mVenus), which allowed to image optoLARG recruitment without activating it.

An eGFP DLC1 plasmid containing the 1091aa long isoform originating from mouse (kindly
provided by Monilola A. Olayioye) was used to produce mCherry-DLC1. The DLC1 gene
was cut out of this plasmid with BamHI and, N-terminally fused to FKBP12-mCherry and
inserted into pB3.0-BLAST containing a CAG-promotor, yielding pB3.0-FKBP12-mCherry-
DLC1.

Spreading Assay

Glass bottom well plates (Celvis) were coated with 5 µg/ml of human plasma fibronectin
purified protein (Merck) for one hour at room temperature. REF52 cells were seeded at a
density of 7000 cells per well (using 24 well plates) in FluoroBrite DMEM medium
containing 1% FBS, 0.1% BSA, 4mM L-Glutamine and 100 U/ml penicillin/streptomycin.
Imaging was done with a Nikon Eclipse Ti-E inverted microscope with an automatic stage.
Temperature was kept at 37ºC with a temperature control system, humidity (100%) and
CO_2 (~5%) with a gas mixer (Life Imaging Services). Focus drift was prevented by the
equipped Perfect Focus System (Nikon). The microscope was controlled with Metamorph
software (Universal Imaging).

Optogenetic Stimulation experiments

Glass bottom well plates (Celvis) were coated with 5 µg/ml of human plasma fibronectin
purified protein (Merck) for one hour at room temperature. REF52 cells were seeded at
6000 cells/well into 12 well plates and allowed to attach for 12 hours. For both RhoA activity
and mCherry-DLC1/miRFP-paxillin optogenetic stimulation experiments cells were
incubated in FluoroBrite DMEM medium overnight. The microscope was controlled with
NIS-Elements (Nikon) for observing DLC1 dynamics and with open-source micromanager
software (Edelstein et al., 2014) for edge stimulation. Optogenetic stimulation was localized
to specific regions using an Andor mosaic 3 DMD. For mCherry-DLC1/miRFP-paxillin
experiments, ROIs were user-defined. For RhoA activity experiments, cells in different
fields of view were selected by a user, cells were then segmented with a custom pixel-
classifier based on a pre-trained VGG16 convolutional neural network (Hinderling et al.,
2024), which was trained on a field of view of the experimental data. ROIs for stimulation
masks were then automatically defined at the top and bottom of the cell for consecutive
illumination with different light pulse regimes using our method for automated optogenetic
targeting (Hinderling et al., 2025). Fields of views were imaged one after the other on
multiple cells. For data extraction the stimulation mask was expanded by gaussian
smoothing and then thresholded to cover a wider region, reducing artifacts caused by
membrane dynamics. Image intensities were calculated for the overlap of this mask with
the segmented cells and the full segmented cells respectively.

Immuno-histochemistry

One hour after seeding, cells were fixed for 10 minutes with 0.2% paraformaldehyde. A
0.1% Triton X solution was used for permeabilization, after which cells were incubated
overnight at 4C with the blocking buffer, containing 5% Bovine-Serum-Albumin and 0.05%
Tween20 in PBS. After each step of the following protocol cells were washed three times
with Phosphate-Buffered-Saline (PBS). Cells were incubated overnight with the primary

antibody (1:250 dilution for paxillin (Abcam ab32084), 1:50 for pMLC (Cell Signaling
Antibody #3671). Secondary antibody incubation (1:1000) lasted one hour. Phalloidin
incubation (1:200 Phalloidin-Atto647N) was performed for 20 minutes and DAPI incubation
for 10 hours. Cells were imaged immediately after the last step.

Image analysis

Lamella size, focal adhesion areas (Figure 1) and RhoA activity ratio measurements
(Figure 3) were computed with CellProfiler in conjunction with a pixel classifier trained with
Ilastik (Berg et al., 2019). Spreading dynamics (Figure 2D) were manually analyzed. FRET
ratio images were computed and analyzed with custom python scripts. DLC1 and paxillin
dynamics (Figure S4D) were calculated with manually drawn ROIs in Fiji (Schindelin et al.,
2012).

ODE model and data fitting

For data fitting and parameters optimisation, we used the Julia programming language and
tools made available by the Scientific Machine Learning (SciML) organization as
framework. The packages ModelingToolkit.jl (Ma et al., 2022) and DifferentialEquations.jl
(Rackauckas and Nie, 2017) allowed to symbolically define and solve the ODE system.
The ODE solver algorithm was specified as Rodas4P method. In addition, we used the
automatic switching algorithm AutoVern9 (Verner, 2010) for stiffness detection. For the
parameter optimization, we defined the problem with the package Optimization.jl (Dixit and
Rackauckas, 2023). The gradient-based method L-BFGS (Wright et al., 2006) was used
as optimization solver from the packages Optim.jl (K Mogensen and N Riseth, 2018).
Additionally, the packages SciMLSensitivity.jl (Rackauckas et al., 2021) and ForwardDiff.jl
(Revels et al., 2016) were required for an efficient and automatic computation of the
gradient with forward propagation. Initial conditions were set to zero for all model species
($\text{Rho}^*_0, \text{GAP}^*_0 \text{ GEF}^*_0$). Total protein concentrations were assumed to be conserved between
active and inactive forms. Given the low concentrations at which the system operates, and
the relatively small fraction of protein that becomes activated, changes in the inactive pool
were considered negligible. As a result, the inactive species were assumed to remain
constant throughout the simulation.

To estimate the activation and deactivation rates of optoLARG, we fitted a simple linear
model to experimental measurements of membrane recruitment. In the case of RhoA
signaling model, Michaelis-Menten kinetics were used to model the interactions between
active species, apart from GAP inactivation which was treated as a first-order (linear)
deactivation process.

In the *DLC1* KO condition, the Michaelis-Menten constant (Km) of RhoA-dependent GAP
activation consistently converged toward zero during optimization, indicating a loss of
negative feedback regulation. To reflect this and simplify the model, we explicitly set this
parameter to zero, resulting in a reduced model in which GAP activity is independent of
RhoA levels. This simplified configuration still produced a strong fit to the experimental
data, supporting the hypothesis that the feedback loop regulating GAP activity is disrupted
in the KO condition.

We also assessed whether model parameters are uniquely recoverable in principle from
the observation scheme using StructuralIdentifiability.jl (Dong et al., 2023). For both the
WT and *DLC1* KO ODE model presentations, we ran the package's known-initial-conditions
workflow with our initial states treated as known and fixed (here set to 0) and default
probabilistic settings (probability threshold 0.99). The analysis returns whether each
parameter (and selected parameter functions) is globally identifiable given the model
equations and measured outputs, i.e., independent of data noise or sampling.

The following table summarizes the results of the structural identifiability analysis:

Parameter	Membrane (WT)	Membrane (*DLC1* KO)	FAs (WT)	FAs (*DLC1* KO)	Comment
$V_{\text{ON}}^{\text{GAP}}$	25.2255	21.51569	33.706	15.6152	*alias $k_{\text{ON}}^{\text{GAP}}$ for the DLC1 KO adapted model*
Km^{GAP}	0.093982	$\to 0$	0.205939	$\to 0$	*consistent across fits*
$k_{\text{OFF}}^{\text{GAP}}$	66.0697	68.84138	99.9969	59.1818	*varied with upper parameter bound*
$V_{\text{ON}}^{\text{Rho}}$	0.0061944	0.00997069	0.00457155	0.00759578	*consistent across fits*
$Km_{\text{ON}}^{\text{Rho}}$	0.0118618	0.031674	0.0108763	0.0382543	*consistent across fits*
$V_{\text{OFF}}^{\text{Rho}}$	25.2258	21.5156	26.7483	15.6151	*varied with upper parameter bound*
$Km_{\text{OFF}}^{\text{Rho}}$	60.6878	94.83013	76.8346	98.2568	*varied with upper parameter bound*

Code availability

The code used to obtain the presented results can be found in the GitHub repository
RhoAModelling (https://github.com/girochat/RhoAModelling/tree/main) as Pluto and
Jupyter notebooks. It contains the model and data fitting of RhoA dynamics at FAs and
membrane as well as the parameter estimation of optoLARG dynamics. The Pluto.jl
package offers a Julia programming environment analog to Jupyter notebook. For
reproducibility, the repository describes how to set a similar coding environment and
references all packages that were used.

Acknowledgements

This work was supported by the Swiss National Science Foundation IZSAZ3_173462
Argentinian-Swiss Joint Research Programme and by the Sinergia CRSII5_183550 grants
to Olivier Pertz. We are grateful to the Microscopy Imaging Center of the University of Bern
for support (https://www.mic.unibe.ch). We thank Kazuhiro Aoki (NIBB, Japan) for providing
the pPBbsr2 vector, David L. Hacker (EPFL, Switzerland), for the pPuro-PB and pHygro-
PB plasmids, Judith Trüb (UniBE, Switzerland) for the cloning of pB3.0-BLAST, Monilola
A. Olayioye (University of Stuttgart, Germany) for the eGFP DLC1 plasmid, Dean E.
Natwick and Sean R. Collins (University of California Davis) for sharing the Stargazin-iLID
constructs. We are grateful to Miguel Vicente-Manzanares, Bernhard Wehrle-Haller and
Daniel Riveline for constructive comments on the manuscript

Author contributions

750   O.P. supervised the research. O.P., M.H. and L.H. wrote the paper. M.H. designed and
751   performed experiments. L.H. designed and performed optogenetic experiments. G.R.
752   developed and implemented the models. J.v.U. built the optoLARG/rGBD circuit. L.D.
753   developed the optoLARG-mScarlet3 construct. M.D. designed data analysis tools.

Additional Information

755   The authors declare no competing interests.

Figure legends.

Figure 1. Cytoskeletal structures in WT and DLC1 deficient cells.

A,B. Representative immunofluorescence images of paxillin and F-actin (phalloidin) (**A**), and pMLC and F-actin (phalloidin) (**B**) immunostains are shown in inverted black and white (ibw) contrast as well as color composites (DAPI signal also included). Scale bar = 10µm.

C. Compared to WT cells (n=243) lamella size is significantly increased in *DLC1* KD cells (n=290, p-adjusted < 0.001) and *DLC1* KO cells (n=242, p-adjusted < 0.001), while *DLC1* rescue cells (n=57) show significantly lower lamella sizes than both, WT (p-adjusted = 0.0098), as well as KO cells (p-adjusted < 0.001).

D. *DLC1* KD cells (n=290) display a significant increase in the total area of focal adhesions per cell (p-adjusted <0.001) compared to WT cells (n=243), while *DLC1* KO cells (n=242) do not show a significant difference (p-adjusted's = 0.09). Rescue cells show a significant decrease in the total area of focal adhesions (p-adjusted < 0.001). ANOVA plus Tukey's honestly significant difference test.

Figure 2. F-actin dynamics and cell morphodynamics in WT and *DLC1* KO cells.

All time series are representative ibw contrast images of WT and *DLC1* KO cells expressing Lifeact-mCherry. Dotted lines mark the area used for the kymographs shown in the right panels.

A. Cells were imaged for 10 minutes in 10 second intervals during spreading immediately after replating. Dotted red lines display differences in lamella size in WT and *DLC1* KO cells. Time

B. One hour after reseeding, when cells are still isotropically spreading, *DLC1* KO cells display increased lamellipodia size and more prominent edge protrusion retraction cycles. High magnification insets are shown for time point 10 minutes and show the robust increase of contractility in *DLC1* KO cells.

C. Cells were imaged for 10 hours with 15 minutes intervals starting from spreading. WT cells break symmetry and display episodes of polarized motility. KO cells are much less dynamic and reach a contractile phenotype much faster, without being able to polarize.

D. Box plot showing at which time point cells exhibited the elongated phenotype described in (**C**). *DLC1* KO cells (n=42) exhibited this phenotype a mean of 90 minutes earlier than WT cells (n=22) p-adjusted = 0.002, Student's t-test. 17 additional WT cells failed to develop that phenotype entirely.

791   Scale bars = 20 μm for all images.

Figure 3. RhoA activation dynamics in WT and *DLC1* KO cells

Cells are stably expressing Lifeact-mCherry, and the RhoA2G FRET sensor. RhoA
localization images show RhoA2G localization that is identical to RhoA. RhoA activity
images display the computed FRET ratio. Images are color-coded according to the
normalized scales shown below the panels.

A. Representative images of WT and *DLC1* KO cells during spreading. Kymographs for the
violet lines are shown in the right panels. Note the RhoA activity band maintains constant
width during spreading at the periphery in WT cells. Note increased global RhoA activity in
DLC1 KO cells, with maintenance of a similar RhoA band pattern at the cell edge.

B. Box plots of FRET ratio averaged over the whole cell (right panel) or a ROI placed at
the cell edge (left panel). This shows that during spreading *DLC1* KO cells (n = 149) have
an increased total RhoA activity compared to WT cells (n = 114, p = 0.005). In addition, the
FRET ratio at just the cell edge is increased as well (p=0.011).

C. Representative images of WT cells and DLC1 KO cells that have transitioned in a
contractile state 12 hours after plating. No difference in RhoA activity pattern can be
observed between WT and *DLC1* KO cells, although the latter still display slightly higher
global RhoA activity levels.

D. Box plots of FRET ratio averaged over the whole cell (right panel) or a ROI placed at
the cell edge (left panel). This shows that contractile *DLC1* KO cells (n=103) have an
increased total RhoA activity (p < 0.001) and edge FRET ratio (p < 0.001) compared to WT
cells (n = 82) compared to WT cells (n = 82). Students t-test.

(A,B) Scale bars = 20 μm.

Figure 4. An optogenetic actuator- Rho biosensor circuit to probe Rho GTPase flux

A. Schematics of the optogenetic actuator - Rho biosensor to measure Rho GTPase flux.
OptoLarg is based on an iLID system which does not interact with Ssb-LARG in the dark
state. The iLID module is anchored to the PM by a Stargazin anchor that displays slow
diffusion, allowing better focussing of optogenetic activation. Upon light exposure ssb-
LARG is locally recruited to the plasma membrane, activating Rho. Rho activity is
measured by a rGBD effector binding domain.

B. Time-series of REF52 cells locally stimulated with light first in a ROI at the cell top with
a high stimulation frequency, and then with a ROI at the cell bottom with a low stimulation
frequency. Light pulses have the same intensity. Thick and thin blue thunder symbols
represent high and low optogenetic stimulation. Red dotted line is used for the kymograph
shown in (C).

C. Kymographs of cells in (B). Light pulse stimulation regimes of top and bottom ROIs are
shown in the upper box. Blue dotted boxes indicate the region and length of the stimulation.

Note how intense optogenetic stimulation in the top ROI leads to Rho activity, assembly of
contractile F-actin structures, and robust edge retraction. Upon removal of the light input,
the Rho activity and F-actin resume, and edge protrusion occurs again. Lower optogenetic
stimulation in the bottom ROI leads to much lower Rho activity and F-actin structures, as
well as lower edge retraction.

(**B**,**C**) Scale bars = 10 µm

Figure 5. Rho GTPase activation kinetics in WT versus DLC1 KO cells

A. Computer vision pipeline and experiment: 1) a cell is segmented using the rGBD-dTomato
channel. 2) FAs and non-FAs are detected using the paxillin channel (see Material and
Methods). ROIs on FAs (blue dots, shown in the bottom left panel) and ROIs on non-FA
regions in between (orange dots, shown in the upper left panel) are selected for stimulation.
3) Image of the stimulation pattern (green channel) shows that the calibrated DMD can
stimulate the regions with high spatial precision (image is overexposed to show the
diffraction pattern of the mirrors).

B. The DMD is used to stimulate ROIs (FAs or non-FAs) with a pulse of blue light (blue line).
rGBD signal fluctuations are then measured in the ROIs.

C. Distribution of paxillin-miRFP intensities in the FA and non-FA ROIs normalized to the
mean paxillin-miRFP intensity of the whole cell shows that our segmentation pipeline
accurately identifies FA and non-FA ROIs.

D. Paxillin-miRFP fluorescene of stimulated ROIs, normalized to the mean paxillin intensity
of the whole cell. Median and 99% CI are shown. Intensity of paxillin does not change upon
stimulation, indicating that the stimulation is too weak to trigger reinforcement of focal
adhesions (compare to Figure 7A-D).

E. Normalized and averaged rGBD fluorescence fluctuations upon ROI optogenetic
stimulation. For each stimulated ROIs, the fold change to the baseline (average activity from
0-150 seconds before optogenetic stimulation) is calculated. Median and 99% CI are shown.
Regions on top of focal adhesions have a larger fold change in rGBD activity than regions

between focal adhesions. *DLC1* KO cells have a larger rGBD fold change in the initial time
after stimulation (150-200 seconds), but then also fall back down to the baseline quicker.
(KO FA: N = 3643, KO non-FA: N = 3643, WT FA: N = 2144, WT non-FA: N = 2144. Mean
number of regions per cell: ~16.90, Cells WT = 321, *DLC1* KO = 431. The same cell can
appear multiple times in the experiment, but with a relaxation time in between and new
stimulation regions).

F. The different dynamics described in (**E**) can be robustly observed in technical replicates
of the experiment. In all replicates the rGBD recruitment is higher in FAs vs non-FAs, and
we see the trend of faster accumulation and faster return to baseline of rGBD in *DLC1* KO
vs WT cells.

Figure 6. Mathematical Model of RhoA differential activation/deactivation
dynamics in WT vs. *DLC1* KO cells
871

A. Schematics of RhoA activation/deactivation system that correspond to the ODE model
of RhoA dynamics. The light pulse increases GEF activation rate, which in turn increases
RhoA activation. Active RhoA activates GAP, negatively auto-regulating itself. RhoA
deactivation is mediated by active GAPs. Detailed description of modeling approach is
provided in the Methods section.

B. Results after fitting the ODE model to the data for both RhoA dynamics at the membrane
(left panel) and at focal adhesions (right panel). WT and *DLC1* KO conditions were fitted
separately in each case. The fold-change to the baseline of averaged and normalized rGBD
intensity was taken as datapoint for each acquisition frame.

C. Illustration of the dynamics of the other model species that are active GEF and active
GAP after the light pulse trigger. The left panel shows the dynamics at the membrane, right
panel at focal adhesions.

Figure 7. Optogenetic control of force-dependent DLC1 interactions with FAs

A. Color-coded fluorescence micrographs of REF52 fibroblast expressing miRFP-paxillin
(left) and mCherry-DLC1 (right) and the optoLARG construct (not shown). The black boxes
indicate the area used for close up images in (B). The white boxes indicate the ROI for
optogenetic illumination. Selected FAs denoted by the pink and green arrowheads. The
black dotted lines were used for the kymograph in B.

B. Kymographs showing two selected FAs, the grey box indicates the time at which
optogenetic stimulation has been applied. Optogenetic stimulation is applied on select
ROIs placed over FAs with 50 ms light pulses per frame (every 15 seconds) for a duration
of 7.5 minutes.

C. Closeups time series of paxillin and DLC1 signals at a single FA denoted by the respective arrowheads. Scale bars = 10 μm.

D. Quantification of mcherry-DLC1 and miRFP-paxillin fluorescence signals during optoLARG mediated control of FA reinforcement and relaxation. Normalized miRFP-paxilllin and mCherry-DLC1 from 2 FAs shown in panels A-C.

E. Model of Rho GTPase activity modulation by DLC1 at FAs and at the plasma membrane relevant to Figure 5. Left and right panels show schematics of Rho activation dynamics in response to optoLARG optogenetic input at FAs (left panel) and plasma membrane (right panels). Top and bottom panel show schematics for Rho activation dynamics in response to optoLARG optogenetic input in control (top) and DLC1 KO (bottom) panels.

F. Model of force dependent regulation of DLC1 at FAs relevant to this and Figure S4. Left panel, in absence of acute mechanical input, DLC1 increases with FA assembly and decreases with FA disassembly. Central panel, upon acute local increase of mechanical stress in response to application of an optoLARG optogenetic input, DLC1 unbinds from FA in a reinforcement regime, and rebinds FA in a relaxing regime when the optoLARG input is removed. Right panel, upon acute local increase of mechanical stress in response to application of an optoLARG optogenetic input some FAs rupture after DLC1 dissociation and FA rupture.

Supplementary Figure legends

Figure S1: Cell motility properties in WT versus DLC1 KO cells

A. Example fields of view of tracked WT and DLC1 KO cells. 20000 REF fibroblasts expressing a histone H2B-miRFP marker were seeded in a 24 well-plate well coated with fibronectin and stimulated with 20 ng/ml PDGF, imaged for 12 hours at 15 minutes interval. Tracking was performed using stardist 37 on the H2B image). DLC1 KO cells display a large subpopulation of cells that remain extremely stationary.

B. Mean Square displacements (MSDs) for different time intervals, WT move more than *DLC1* KO cells for all possible lag times. Plot shows directionality (exponential bit between 0.25 and 1.5 hours). For times longer than 2 hours the curve is flat indicating that migration has characteristics of random walk at these timescales. Thick lines: mean, thin lines: standard deviation. Dotted grey line shows delta t chosen for figure C.

C. Distribution of velocities calculated as μm/hour from Root-MSD. We again observe the bimodal distribution in the *DLC1* KO cell with a big subgroup of the cells move extremely slowly (5 μm/hour). Thick lines are mean and extrema. Two sided t-test: (statistic=25.1, p-value=1.6e-131)

Figure S2. Effect of DLC1 overexpression on F-actin cytoskeleton and RhoA activity

A. Effect of DLC1 overexpression of F-actin. REF52 cells stably expressing Lifeact-mCherry were transfected with mCherry-DLC1 or mCherry plasmids, allowed to spread for 12 hours on fibronectin-coated coverslips and imaged. Images are shown in ibw contrast. Large mCherry-DLC1 pool in the cytosol levels documents its overexpression. Note how DLC1 overexpression leads to loss of contractile F-actin structures.

B. Effect of DLC1 overexpression of RhoA. Cells stably transfected with the RhoA2G biosensor were transfected with mCherry-paxillin or mCherry-DLC1. mCherry signals are shown in ibw contrast. RhoA2G FRET ratio, and expression levels are color-coded according to the scale. Note how low mCherry-DLC1 expression that remains associated with FA already lowers RhoA activity, while high mCherry-DLC1 expression leads to even lower RhoA activity and aberrant cell morphology.

Figure S3: Quantification of optoLARG recruitment dynamics to subcellular locations.

A. optoLARG recruitment over time shows similar dynamics to FA ROIs (N=190) and non-FA ROIs (N=184) regions (background subtracted, divided by mean optoLARG expression in the cell and normalized to baseline of first 10 frames / 100 s). Mean of all ROIs and 99% CI are shown.

B. Comparing optoLARG recruitment at peak response (t = 200 s), averaging all FA/non-FA ROIs within same cell. Wilcoxon paired test shows no significant difference between the two conditions (cells N = 39, p = 0.419, mean FA: 1.1450 ± 0.23, mean NON-FA: 1.1884 ± 0.22).

C. Characterization of optoLARG recruitment dynamics. As FA and non-FA regions shows similar dynamics, here we average all ROIs. To quantify the kinetics of recruitment and dissociation, we fit the mean normalized and drift-corrected time series with piecewise exponential models. The rising phase during stimulation (150-200 s) was well described by an inverse exponential function, with an amplitude of 0.166 and a half-life of 7.9 seconds (R^2 = 0.991, RSS = $1.88*10^{-4}$). The decay phase (200-300 s) was well described by a single exponential decay with a half-life of 10.8 seconds and a slightly lower amplitude of 0.164 (R^2 = 0.989, RSS = $3.24*10^{-4}$).

Figure S4. DLC1 dynamics at FAs in unperturbed cells

Panels **A-C** document FAs in an assembly state. This shows that the DLC1 signal
augments concomitantly with paxillin signal during FA assembly.

Panels **D-F** document FAs in a disassembly state. This shows that the DLC1 signal
decreases concomitantly with paxillin signal during FA disassembly.

A,D. Color-coded fluorescence micrographs of REF52 fibroblast expressing miRFP-paxillin
(top) and mCherry-DLC1 (bottom). The white boxes indicate the area used for close up
images in (B,E).

B,E. Closeups of the ROIs (left panel) and kymographs (right panel) of selected FAs
denoted by the red arrowheads. White dotted lines were used for the kymograph. In the
kymograph, the grey box indicates the time at which optogenetic stimulation has been
applied.

C,F. Closeups time series of paxillin and DLC1 signals at a single FA denoted by the
respective red arrowheads.

Scale bars = 10 µm.

Figure S5. DLC1 dynamics in an optoLARG stimulated FA that undergoes
reinforcement followed by disassembly, as well as FA behavior in absence
of optoLARG stimulus

Panels **A-C** document FAs that when subjected to a nearby pulse of optogenetic Rho-
mediated contractility display a behavior of FA reinforcement followed by disassembly.
Optogenetic stimulation is applied on select ROIs placed over FAs with 50 ms light pulses
per frame (every 15 seconds) for a duration of 8 minutes.

Panels **D-F** document the dynamics of DLC1 in a ROI not stimulated with light in the same
cell as in shown in Figure 6A-C.

A. Color-coded fluorescence micrographs of REF52 fibroblast expressing miRFP-paxillin
(top) and mCherry-DLC1 (bottom) and the optoLARG construct (not shown). The black
boxes indicate the area used for close up images in (B). The white boxes indicate the ROI
for optogenetic illumination.

B. Closeups of the ROIs (left panel) and kymographs (right panel) of selected FAs denoted
by the arrowheads. The pink and green arrowheads indicate the FAs of interest in (B). The
black arrowhead indicates the FA of interest in (C). The black dotted lines were used for
the kymograph. In the kymograph, the grey box indicates the time at which optogenetic
stimulation has been applied.

C. Closeups time series of paxillin and DLC1 signals at a single FA denoted by the respective arrowheads. Scale bars = 10 μm.

D. Color-coded fluorescence micrographs of REF52 fibroblast expressing miRFP-paxillin (top) and mCherry-DLC1 (bottom) and the optoLARG construct (not shown). The black boxes indicate the area used for close up images in (B,E). The white boxes indicate the ROI for optogenetic illumination.

E. Closeups of the ROIs (left panel) and kymographs (right panel) of selected FAs denoted by the arrowheads. The pink and green arrowheads indicate the FAs of interest in (B). The black arrowhead indicates the FA of interest in (D). The black dotted lines were used for the kymograph. In the kymograph, the grey box indicates the time at which optogenetic stimulation has been applied.

F. Closeups time series of paxillin and DLC1 signals at a single FA denoted by the respective arrowheads.

Supplementary Movie legends

Movie S1: F-actin dynamics of WT and DLC1 KO REF52 cells during late spreading (relevant to Figure 2B).

Representative ibw contrast movie of WT and *DLC1* KO cells expressing Lifeact-mCherry during late spreading. Time: Minutes:seconds. Scale bar = 20 mm.

Movie S2: F-actin dynamics and edge dynamics of WT and DLC1 KO REF52 cells during acquisition of a polarized cell migration phenotype over a period of 15 hours (relevant to Figure 2C).

Representative ibw contrast movie of WT and *DLC1* KO cells expressing Lifeact-mCherry for 15. Edge dynamics were color-coded with respect to time. Time: Minutes:seconds. Scale bar = 20 mm.

Movie S3: F-actin and RhoA activity dynamics WT and DLC1 KO REF52 cells during late spreading (relevant to Figure 3A).

Representative movie of WT and *DLC1* KO cells expressing Lifeact-mCherry and the RhoA2G FRET biosensor during late spreading.

1027 Left panels: F-actin channel in ibw contrast. Right panels: RhoA activity emission ratio color
1028 coded according to color scale.

1029 Time: Minutes:seconds. Scale bar = 20 mm.

1030

Movie S4: Characterisation of the optoLARG/rGBD genetic circuit (relati to
Figure 5)

REF52 cells stably expressing the optoLARG/rGBD circuit as well as Lifeact-miRFP are
locally stimulated with blue light of identical intensity, first in a ROI at the cell top with a high
stimulation frequency, and then with a ROI at the cell bottom with a lower stimulation
frequency. The stimulation frequencies are indicated by the appearance/disappearance
frequency of the ROIs.

Left panel: rGBD signal, Right panel: F-actin lifeact-miRFP. Note the more robust edge
contraction, as well as increased RhoA activity in response to high versus low light
frequency stimulation.

Time: Minutes:seconds. Scale bar = 10 mm.

1042

Movie S5: DLC1 dynamics at FAs during an optoLARG-induced contractility
pulse (relevant to Figure 6A-C, cell denoted by pink arrow)

REF52 KO cells rescued with mCherry-DLC1 and expressing miRFP-paxillin, and the
optoLARG/rGBD circuit were imaged before, during and after a transient optogenetic
stimulation in the whole field of view shown in the movie.

Time: Minutes:seconds. Scale bar = 5 mm.

References

Basant A, Glotzer M. 2018. Spatiotemporal Regulation of RhoA during Cytokinesis. *Curr Biol* **28**:R570–R580. doi:10.1016/j.cub.2018.03.045

Bement WM, Goryachev AB, Miller AL, Von Dassow G. 2024. Patterning of the cell cortex by Rho GTPases. *Nat Rev Mol Cell Biol* **25**:290–308. doi:10.1038/s41580-023-00682-z

Bement WM, Leda M, Moe AM, Kita AM, Larson ME, Golding AE, Pfeuti C, Su K-C, Miller AL, Goryachev AB, von Dassow G. 2015. Activator-inhibitor coupling between Rho signalling and actin assembly makes the cell cortex an excitable medium. *Nat Cell Biol* **17**:1471–1483.

Bement WM, von Dassow G. 2014. Single cell pattern formation and transient cytoskeletal arrays. *Curr Opin Cell Biol* **26**:51–59.

Berg S, Kutra D, Kroeger T, Straehle CN, Kausler BX, Haubold C, Schiegg M, Ales J, Beier T, Rudy M, Eren K, Cervantes JI, Xu B, Beuttenmueller F, Wolny A, Zhang C, Koethe U, Hamprecht FA, Kreshuk A. 2019. ilastik: interactive machine learning for (bio) image analysis. *Nat Methods* **16**:1–7.

Blum Y, Mikelson J, Dobrzyński M, Ryu H, Jacques M-A, Jeon NL, Khammash M, Pertz O. 2019. Temporal perturbation of ERK dynamics reveals network architecture of FGF2/MAPK signaling. *Mol Syst Biol* **15**:e8947. doi:10.15252/msb.20198947

De Seze J, Gatin J, Coppey M. 2023. RhoA regulation in space and time. *FEBS Lett* **597**:836–849. doi:10.1002/1873-3468.14578

DerMardirossian C, Rocklin G, Seo J-Y, Bokoch GM. 2006. Phosphorylation of RhoGDI by Src regulates Rho GTPase binding and cytosol-membrane cycling. *Mol Biol Cell* **17**:4760–4768. doi:10.1091/mbc.e06-06-0533

Dessauges C, Mikelson J, Dobrzyński M, Jacques M, Frismantiene A, Gagliardi PA, Khammash M, Pertz O. 2022. Optogenetic actuator – ERK biosensor circuits identify MAPK network nodes that shape ERK dynamics. *Mol Syst Biol* **18**. doi:10.15252/msb.202110670

Dixit VK, Rackauckas C. 2023. Optimization.jl: A Unified Optimization Package. doi:10.5281/ZENODO.7738525

Dong R, Goodbrake C, Harrington HA, Pogudin G. 2023. Differential Elimination for Dynamical Models via Projections with Applications to Structural Identifiability. *SIAM J Appl Algebra Geom* **7**:194–235. doi:10.1137/22M1469067

Edelstein AD, Tsuchida MA, Amodaj N, Pinkard H, Vale RD, Stuurman N. 2014. Advanced methods of microscope control using µManager software. *J Biol Methods* **1**. doi:10.14440/jbm.2014.36

Erlmann P, Schmid S, Horenkamp FA, Geyer M, Pomorski TG, Olayioye MA. 2009. DLC1 activation requires lipid interaction through a polybasic region preceding the RhoGAP domain. *Mol Biol Cell* **20**:4400–4411. doi:10.1091/mbc.e09-03-0247

Etienne-Manneville S, Hall A. 2002. Rho GTPases in cell biology. *Nature* **420**:629–635.

Fritz RD, Letzelter M, Reimann A, Martin K, Fusco L, Ritsma L, Ponsioen B, Fluri E, Schulte-Merker S, van Rheenen J, Pertz O. 2013. A versatile toolkit to produce sensitive FRET biosensors to visualize signaling in time and space. *Sci Signal* **6**:rs12. doi:10.1126/scisignal.2004135

Fritz RD, Pertz O. 2016a. The dynamics of spatio-temporal Rho GTPase signaling: formation of signaling patterns. *F1000Research* **5**. doi:10.12688/f1000research.7370.1

Fritz RD, Pertz O. 2016b. The dynamics of spatio-temporal Rho GTPase signaling: formation of signaling patterns. *F1000Research* **5**:F1000 Faculty Rev-749. doi:10.12688/f1000research.7370.1

Fusco L, Lefort R, Smith K, Benmansour F, Gonzalez G, Barillari C, Rinn B, Fleuret F, Fua P, Pertz O. 2016. Computer vision profiling of neurite outgrowth dynamics reveals spatiotemporal modularity of Rho GTPase signaling. *J Cell Biol* **212**:91–111. doi:10.1083/jcb.201506018

Gingras AR, Bate N, Goult BT, Hazelwood L, Canestrelli I, Grossmann JG, Liu H, Putz NSM, Roberts GCK, Volkmann N, Hanein D, Barsukov IL, Critchley DR. 2008. The structure of the C-terminal actin-binding domain of talin. *EMBO J* **27**:458–469. doi:10.1038/sj.emboj.7601965

Goult BT, Brown NH, Schwartz MA. 2021. Talin in mechanotransduction and mechanomemory at a glance. *J Cell Sci* **134**:jcs258749. doi:10.1242/jcs.258749

Graessl M, Koch J, Calderon A, Kamps D, Banerjee S, Mazel T, Schulze N, Jungkurth JK, Patwardhan R, Solouk D, Hampe N, Hoffmann B, Dehmelt L, Nalbant P. 2017. An excitable Rho GTPase signaling network generates dynamic subcellular contraction patterns. *J Cell Biol* **17**:jcb.201706052-15.

Guntas G, Hallett RA, Zimmerman SP, Williams T, Yumerefendi H, Bear JE, Kuhlman B. 2015. Engineering an improved light-induced dimer (iLID) for controlling the localization and activity of signaling proteins. *Proc Natl Acad Sci U S A* **112**:112–117.

Gutenkunst RN, Waterfall JJ, Casey FP, Brown KS, Myers CR, Sethna JP. 2007. Universally Sloppy Parameter Sensitivities in Systems Biology Models. *PLoS Comput Biol* **3**:e189. doi:10.1371/journal.pcbi.0030189

Haining AWM, Rahikainen R, Cortes E, Lachowski D, Rice A, von Essen M, Hytönen VP, del Río Hernández A. 2018. Mechanotransduction in talin through the interaction of the R8 domain with DLC1. *PLOS Biol* **16**:e2005599-20.

Hallett RA, Zimmerman SP, Yumerefendi H, Bear JE, Kuhlman B. 2016. Correlating in Vitro and in Vivo Activities of Light-Inducible Dimers: A Cellular Optogenetics Guide. *ACS Synth Biol* **5**:53–64. doi:10.1021/acssynbio.5b00119

Hinderling L, Landolt AE, Grädel B, Dubied L, Zahni C, Kwasny M, Frismantiene A, Lambert T, Dobrzyński M, Pertz O. 2025. Real-time feedback control microscopy for automation of optogenetic targeting. doi:10.1101/2025.08.17.670729

Hinderling L, Witz G, Schwob R, Stojiljkovic A, Dobrzynski M, Vladymyrov M, Frei J, Graedel B, Frismantiene A, Pertz O. 2024. Convpaint - Universal framework for interactive pixel classification using pretrained neural networks. doi:10.1101/2024.09.12.610926

Hu J, Gong X, Strömblad S. 2022. Local temporal Rac1-GTP nadirs and peaks restrict cell protrusions and retractions. *Sci Adv* **8**:eabl3667. doi:10.1126/sciadv.abl3667

Isogai T, Danuser G. 2018. Discovery of functional interactions among actin regulators by analysis of image fluctuations in an unperturbed motile cell system. *Philos Trans R Soc B Biol Sci* **373**:20170110. doi:10.1098/rstb.2017.0110

Itoh RE, Kurokawa K, Ohba Y, Yoshizaki H, Mochizuki N, Matsuda M. 2002. Activation of Rac and Cdc42 Video Imaged by Fluorescent Resonance Energy Transfer-Based Single-Molecule Probes in the Membrane of Living Cells. *Mol Cell Biol* **22**:6582–6591. doi:10.1128/MCB.22.18.6582-6591.2002

K Mogensen P, N Riseth A. 2018. Optim: A mathematical optimization package for Julia. *J Open Source Softw* **3**:615. doi:10.21105/joss.00615

Kaushik S, Ravi A, Hameed FM, Low BC. 2014. Concerted modulation of paxillin dynamics at focal adhesions by deleted in liver cancer-1 and focal adhesion kinase during early cell spreading: DLC1 Regulates Paxillin Dynamics at Focal Adhesions via FAK. *Cytoskeleton* **71**:677–694. doi:10.1002/cm.21201

Kraynov VS, Chamberlain C, Bokoch GM, Schwartz MA, Slabaugh S, Hahn KM. 2000. Localized Rac activation dynamics visualized in living cells. *Science* **290**:333–337.

Krendel M, Zenke FT, Bokoch GM. 2002. Nucleotide exchange factor GEF-H1 mediates cross-talk between microtubules and the actin cytoskeleton. *Nat Cell Biol* **4**:294–301. doi:10.1038/ncb773

Lam NN, Docherty PD, Murray R. 2022. Practical identifiability of parametrised models: A review of benefits and limitations of various approaches. *Math Comput Simul* **199**:202–216. doi:10.1016/j.matcom.2022.03.020

Li G, Du X, Vass WC, Papageorge AG, Lowy DR, Qian X. 2011. Full activity of the deleted in liver cancer 1 (DLC1) tumor suppressor depends on an LD-like motif that binds talin and focal adhesion kinase (FAK). *Proc Natl Acad Sci* **108**:17129–17134. doi:10.1073/pnas.1112122108

Ma Y, Gowda S, Anantharaman R, Laughman C, Shah V, Rackauckas C. 2022. ModelingToolkit: A Composable Graph Transformation System For Equation-Based Modeling. doi:10.48550/arXiv.2103.05244

Machacek M, Hodgson L, Welch C, Elliott H, Pertz O, Nalbant P, Abell A, Johnson GL, Hahn KM, Danuser G. 2009. Coordination of Rho GTPase activities during cell protrusion. *Nature* **461**:99–103. doi:10.1038/nature08242

Mahlandt EK, Arts JJG, van der Meer WJ, van der Linden FH, Tol S, van Buul JD, Gadella TWJ, Goedhart J. 2021. Visualizing endogenous Rho activity with an improved localization-based, genetically encoded biosensor. *J Cell Sci* **134**:jcs258823. doi:10.1242/jcs.258823

Martin K, Reimann A, Fritz RD, Ryu H, Jeon NL, Pertz O. 2016. Spatio-temporal co-ordination of RhoA, Rac1 and Cdc42 activation during prototypical edge protrusion and retraction dynamics. *Sci Rep* **6**:21901. doi:10.1038/srep21901

Martin K, Vilela M, Jeon NL, Danuser G, Pertz O. 2014. A growth factor-induced, spatially organizing cytoskeletal module enables rapid and persistent fibroblast migration. *Dev Cell* **30**:701–716. doi:10.1016/j.devcel.2014.07.022

Matasci M, Baldi L, Hacker DL, Wurm FM. 2011. The PiggyBac transposon enhances the frequency of CHO stable cell line generation and yields recombinant lines with superior productivity and stability. *Biotechnol Bioeng* **108**:2141–2150. doi:10.1002/bit.23167

Miller AL, Bement WM. 2009. Regulation of cytokinesis by Rho GTPase flux. *Nat Cell Biol* **11**:71–77. doi:10.1038/ncb1814

Moon SY, Zheng Y. 2003. Rho GTPase-activating proteins in cell regulation. *Trends Cell Biol* **13**:13–22. doi:10.1016/s0962-8924(02)00004-1

Mosaddeghzadeh N, Ahmadian MR. 2021. The RHO Family GTPases: Mechanisms of Regulation and Signaling. *Cells* **10**:1831. doi:10.3390/cells10071831

Müller PM, Rademacher J, Bagshaw RD, Wortmann C, Barth C, van Unen J, Alp KM, Giudice G, Eccles RL, Heinrich LE, Pascual-Vargas P, Sanchez-Castro M, Brandenburg L, Mbamalu G, Tucholska M, Spatt L, Czajkowski MT, Welke R-W, Zhang S, Nguyen V, Rrustemi T, Trnka P, Freitag K, Larsen B, Popp O, Mertins P, Gingras A-C, Roth FP, Colwill K, Bakal C, Pertz O, Pawson T, Petsalaki E, Rocks O. 2020. Systems analysis of RhoGEF and RhoGAP regulatory proteins reveals spatially organized RAC1 signalling from integrin adhesions. *Nat Cell Biol* **22**:498–511. doi:10.1038/s41556-020-0488-x

Nalbant P, Hodgson L, Kraynov V, Toutchkine A, Hahn KM. 2004. Activation of endogenous Cdc42 visualized in living cells. *Science* **305**:1615–1619.

Natwick DE, Collins SR. 2021. Optimized iLID Membrane Anchors for Local Optogenetic Protein Recruitment. *ACS Synth Biol* **10**:1009–1023. doi:10.1021/acssynbio.0c00511

Oakes PW, Wagner E, Brand CA, Probst D, Linke M, Schwarz US, Glotzer M, Gardel ML. 2017. Optogenetic control of RhoA reveals zyxin-mediated elasticity of stress fibres. *Nat Commun* **8**:15817–12.

Pertz O. 2010. Spatio-temporal Rho GTPase signaling - where are we now? *J Cell Sci* **123**:1841–1850. doi:10.1242/jcs.064345

Pertz O, Hodgson L, Klemke RL, Hahn KM. 2006. Spatiotemporal dynamics of RhoA activity in migrating cells. *Nature* **440**:1069–1072. doi:10.1038/nature04665

Rackauckas C, Ma Y, Martensen J, Warner C, Zubov K, Supekar R, Skinner D, Ramadhan A, Edelman A. 2021. Universal Differential Equations for Scientific Machine Learning. doi:10.48550/arXiv.2001.04385

Rackauckas C, Nie Q. 2017. DifferentialEquations.jl – A Performant and Feature-Rich Ecosystem for Solving Differential Equations in Julia. *J Open Res Softw* **5**:15. doi:10.5334/jors.151

Revels J, Lubin M, Papamarkou T. 2016. Forward-Mode Automatic Differentiation in Julia. doi:10.48550/arXiv.1607.07892

Riedl J, Crevenna AH, Kessenbrock K, Yu JH, Neukirchen D, Bista M, Bradke F, Jenne D, Holak TA, Werb Z, Sixt M, Wedlich-Soldner R. 2008. Lifeact: a versatile marker to visualize F-actin. *Nat Methods* **5**:605–607. doi:10.1038/nmeth.1220

Rochussen AM, Ma CY, Griffiths GM. 2025. Transcriptional adaptation after deletion of *Cdc42* in primary T cells. *J Cell Sci* **138**:jcs263826. doi:10.1242/jcs.263826

Rossman KL, Der CJ, Sondek J. 2005. GEF means go: turning on RHO GTPases with guanine nucleotide-exchange factors. *Nat Rev Mol Cell Biol* **6**:167–180.

Ryu H, Chung M, Dobrzyński M, Fey D, Blum Y, Lee SS, Peter M, Kholodenko BN, Jeon NL, Pertz O. 2015. Frequency modulation of ERK activation dynamics rewires cell fate. *Mol Syst Biol* **11**:838. doi:10.15252/msb.20156458

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**:676–682. doi:10.1038/nmeth.2019

Schmick M, Kraemer A, Bastiaens PIH. 2015. Ras moves to stay in place. *Trends Cell Biol* **25**:190–197.

Tsyganov MA, Kolch W, Kholodenko BN. 2012. The topology design principles that determine the spatiotemporal dynamics of G-protein cascades. *Mol Biosyst* **8**:730–743.

Wu C-F, Lew DJ. 2013. Beyond Symmetry Breaking: Competition and Negative Feedback in GTPase regulation. *Trends Cell Biol* **23**:476–483. doi:10.1016/j.tcb.2013.05.003

Zhao Y, Lykov N, Tzeng C. 2022. Talin-1 interaction network in cellular mechanotransduction (Review). *Int J Mol Med* **49**:1–12. doi:10.3892/ijmm.2022.5116

Figure 1. Cytoskeletal structures in WT and DLC1 deficient cells.

Figure 2. F-actin dynamics and cell morphodynamics in WT and *DLC1* KO cells.

Figure 3. RhoA activation dynamics in WT and *DLC1* KO cells.

Figure 4. An optogenetic actuator- Rho biosensor circuit to probe Rho GTPase flux.

Figure 5. Rho GTPase activation kinetics in WT versus DLC1 KO cells.

Figure 6. Mathematical Model of RhoA differential activation/deactivation dynamics in WT vs. *DLC1* KO cells.

Figure 7. Optogenetic control of force-dependent DLC1 interactions with FAs.

Figure S1: Cell motility properties in WT versus DLC1 KO cells.

Figure S2. Effect of DLC1 overexpression on F-actin cytoskeleton and RhoA activity.

Figure S3: Quantification of optoLARG recruitment dynamics to subcellular locations.

Figure S4. DLC1 dynamics at FAs in unperturbed cells.

Figure S5. DLC1 dynamics in an optoLARG stimulated FA that undergoes reinforcement followed by disassembly, as well as FA behavior in absence of optoLARG stimulus.