Light-exercise-induced dopaminergic and noradrenergic

stimulation in the dorsal hippocampus:

using a rodent physiological exercise model

Taichi Hiraga[a], Toshiaki Hata[a], Masahiro Okamoto[a,b], and Hideaki Soya[a,b]

[a] Laboratory of Exercise Biochemistry and Neuroendocrinology, Faculty of Health and Sport Sciences,

University of Tsukuba, Tsukuba, Ibaraki 305-8574, Japan

[b] Division of Sport Neuroscience, Department of Mind, Advanced Research Initiative for Human High

Performance (ARIHHP), Faculty of Health and Sport Sciences, University of Tsukuba, Tsukuba,

Ibaraki 305-8574, Japan

Corresponding author: Hideaki Soya, Ph.D.

Email: soya.hideaki.gt@u.tsukuba.ac.jp

Running title: Light exercise activates hippocampus via NA and DA systems

Abbreviations:

5-HT: serotonin

AAS: ascending arousal system

ACTH: adrenocorticotropic hormone

AD: Alzheimer's disease

BDNF: brain-derived neurotrophic factor

BSA: bovine serum albumin

CA: cornu ammonis

DA: dopamine

DAPI: 4',6-diamidino-2-phenylindole

DG: dentate gyrus

DRN: dorsal raphe nucleus

HPLC-ECD: high-performance liquid chromatography with an electrochemical detector

LC: locus coeruleus

LTD: long-term depression

LTP: long-term potentiation

LT: lactate threshold

MCI: mild cognitive impairment

MRN: median raphe nucleus

NA: noradrenaline

PB: phosphate buffer

PBT: phosphate buffer with Triton X-100

PFA: paraformaldehyde

PRP: plasticity-related protein

PVN: paraventricular nucleus

RM: repeated measures

RRF: retrorubral field

SNpc: substantia nigra pars compacta

TH: tyrosine hydroxylase

VTA: ventral tegmental area

Graphical abstract

Possible mechanism underlying hippocampal activation

Rat brain

dorsal hippocampus

VTA

LC

Light Exercise

dorsal hippocampus

dopamine   noradrenaline

VTA terminal   LC terminal

CONCLUSION

Light exercise could activate the dorsal hippocampus through LC-NA and VTA-DA stimulation.

Our previous research with both animals and humans has demonstrated that even light exercise below the lactate threshold (or 50% $\dot{V}O_{2peak}$) can boost neuronal activity in the dorsal hippocampus and enhance learning and memory. While accumulating evidence highlights the significance of cellular and molecular basis of exercise benefits, the basic circuit mechanism underlying this boost remains largely undetermined. It has been proposed that exercise engages the ascending monoaminergic projections, which facilitates dorsal hippocampus-associated functions. Thus, we examined this hypothesis using our animal exercise model, and found a possible contribution of noradrenergic neurons in the locus coeruleus and dopaminergic neurons in the ventral tegmental area to dorsal hippocampal activation during light exercise, implying a circuit mechanism for light-exercise-induced benefits to the dorsal hippocampus.

Abstract

Growing evidence shows that exercise acts on the dorsal hippocampus and promotes the

learning process and memory formation. For decades, these beneficial effects of exercise

have been based on a lot of research using demanding and stress-response-inducing

exercise at moderate-to-vigorous intensities. In contrast, our translational research with

animals and humans has focused on light exercise, which almost anyone can safely

perform with minimal stress. Using our special exercise model, which is based on

physiological responses such as lactate threshold (LT), we found that even light exercise

can boost dorsal hippocampal activity and enhance memory performance. Although the

circuit mechanism underlying this boost remains unclear, arousal promotion even with

light-intensity exercise implies the involvement of the monoaminergic ascending arousal

system, which is crucial for memory enhancement. We thus aimed to examine this

hypothesis using our physiological exercise model based on the LT of rats. First, we

immunohistochemically assessed the neuronal activation of the dorsal hippocampal sub-

regions, as well as tyrosine hydroxylase-positive (TH^+) neurons and serotonin-positive

($5\text{-}HT^+$) neurons in the brainstem monoaminergic nuclei. Second, we monitored the

dynamics of monoamine release at the dorsal hippocampus using *in vivo* microdialysis

combined with the high-performance liquid chromatography (HPLC) system. We found

that even light exercise increased neuronal activity in the dorsal hippocampal sub-regions

and induced the release of noradrenaline (NA) and dopamine (DA). Additionally, we

found that TH^+ neurons in the locus coeruleus (LC) and the ventral tegmental area (VTA)

were activated even by light exercise and were both positively correlated with the dorsal

hippocampal activation. Meanwhile, the serotonergic system was not affected by

exercise. Overall, these results suggest that light exercise stimulates the dorsal

hippocampal neurons probably through the noradrenergic neurons in the LC and

dopaminergic neurons in the VTA. Our findings provide insight into the underlying circuit

mechanisms of light-exercise-enhanced memory.

Keywords Light exercise; Dorsal hippocampus; Locus coeruleus; Ventral tegmental area;

Noradrenaline; Dopamine

Introduction

There is accumulating evidence showing that physical exercise has beneficial effects on

the brain and its functions, including hippocampus-dependent learning and memory (van

Praag *et al.* 1999; Erickson *et al.* 2011; Vargas *et al.* 2017; van Dongen *et al.* 2016). The

hippocampus is well-known for being involved in several processes related to emotion,

stress resilience, and cognitive function. In particular, the dorsal part of the hippocampus

plays a key role in the learning process and memory formation (Fanselow and Dong 2010).

Physical exercise has been found to increase the activity of the dorsal hippocampal

neurons (Oladehin and Waters 2001; Lee *et al.* 2003; Nishijima and Soya 2006), which

induces the production of plasticity-related proteins (PRPs) such as the brain-derived

neurotrophic factor (BDNF) (Soya *et al.* 2007b; Neeper *et al.* 1995; Vaynman *et al.* 2003).

This is, in turn, thought to trigger neuronal and synaptic plasticity and memory

enhancement (Vaynman *et al.* 2004; Li *et al.* 2008; van Praag *et al.* 1999; Lee *et al.* 2012).

For decades, the benefits of exercise to learning and memory have been based on

evidence using exercise at moderate-to-vigorous intensities (Erickson *et al.* 2011; Vargas

et al. 2017; van Dongen *et al.* 2016; Pereira *et al.* 2007), which is generally recommended

in exercise therapy. However, such demanding exercise can cause physiological stress

and reduce exercise adherence (Soya *et al.* 2007a; Ekkekakis *et al.* 2008; Saito and Soya

2004; Hill *et al.* 2008; Perri *et al.* 2002). In contrast, we have focused on light-intensity

exercise (light exercise), which can be safely performed with minimum physiological

stress, and have revealed its beneficial effects on the hippocampus. We have been

employing a rodent treadmill running model based on physiological responses such as

lactate threshold (LT). LT is the signpost for exercise-induced stress, which begins with

the onset of lactate accumulation in the blood and the accumulation of adrenocorticotropic

hormone (ACTH) (Soya *et al.* 2007a; Saito and Soya 2004). This physiological exercise

model enables us to define running below the LT as light exercise with minimum

physiological stress and running above the LT as vigorous, stress-inducing exercise

stimuli. Using this model, we found that even light exercise was sufficient to acutely

stimulate hippocampal activity and BDNF induction (Soya *et al.* 2007b), and to promote

hippocampal neurogenesis and spatial memory (Inoue *et al.* 2015a; Inoue *et al.* 2015b;

Okamoto *et al.* 2012). Conversely, excessive ACTH secretion induced by vigorous

exercise cancels out these beneficial effects on hippocampal plasticity and memory due

to the stress-induced hippocampal vulnerability (Inoue *et al.* 2015a; Inoue *et al.* 2015b;

Okamoto *et al.* 2012; Conrad 2010; Schoenfeld and Gould 2012). Additionally, our

research with human participants found that acute light exercise increases hippocampal

activity and enhances pattern separation, which is a hippocampus-related function

(Suwabe *et al.* 2018). Taken together, our translational research demonstrates the

importance of exercise intensity for determining the effect of exercise on the

hippocampus and highlights the advantages of light exercise.

However, prior studies, including our own findings, have yet to address the circuit

mechanism underlying the benefits of light exercise to the dorsal hippocampus.

Classically, it is known that exercise, even light exercise, can promote arousal (Suwabe

et al. 2018; Kuwamizu *et al.* 2022; McMorris 2016; Cooper 1973; Yamazaki *et al.* 2023),

which is thought to originate from the activation of the ascending arousal system (AAS)

(Saper *et al.* 2001; Jones 2003). The AAS mainly consists of cholinergic neurons and

monoaminergic neurons and sends out widespread projections in order to increase global

brain activity. In particular, it has been proposed that exercise can stimulate the release of

neuromodulatory factor monoamine in several brain regions, including the dorsal

hippocampus (Meeusen *et al.* 2001). Monoamines, including noradrenaline (NA),

dopamine (DA), and serotonin (5-HT), are essential for multiple functions such as arousal,

attention, reward, motivation, and mood regulation (Carter *et al.* 2010; Bari *et al.* 2020;

Saunders *et al.* 2018; Zhu *et al.* 2016; Kim *et al.* 2013; Warden *et al.* 2012; Teissier *et al.*

2015). At the dorsal hippocampus, they regulate learning and memory processes by

modulating the expression of PRPs and neuronal plasticity, such as neurogenesis and

long-term potentiation (LTP) (Palacios-Filardo and Mellor 2019; Poe *et al.* 2020; Wang

et al. 2010; Mahar *et al.* 2014; Coradazzi *et al.* 2016; Salvi *et al.* 2016). The hippocampal

monoamines are mainly synthesized and provided by the specific nuclei located in the

pons noradrenergic locus coeruleus (LC), the midbrain dopaminergic nuclei, such as the

ventral tegmental area (VTA), the substantia nigra pars compacta (SNpc), and the

retrorubral field (RRF), and the midbrain serotonergic raphe nuclei (the dorsal raphe:

DRN and the median raphe: MRN) (Gasbarri *et al.* 1996; Schwarz *et al.* 2015; Broussard

et al. 2016; Muzerelle *et al.* 2016; Luchetti *et al.* 2020). Previous research demonstrates

that single bouts of exercise activate brainstem monoaminergic neurons (Morikawa *et al.*

2021; Tsai *et al.* 2019; Rodovalho *et al.* 2020) and affect the hippocampal monoamine

transmissions (Vargas *et al.* 2017; Goekint *et al.* 2012; Gomez-Merino *et al.* 2001),

thereby contributing to the enhancement of hippocampus-related functions (Vargas *et al.*

2017; Vargas *et al.* 2020; Morikawa *et al.* 2021). These findings lead us to the hypothesis

that the monoamines provided by the brainstem nuclei could be involved in the beneficial

effects of light exercise on the dorsal hippocampus.

To examine this hypothesis, we employed a treadmill running model based on the

LT of rats to investigate the involvement of the brainstem monoaminergic system on

dorsal hippocampal activation during light exercise. First, we analyzed the neuronal

activation of the dorsal hippocampus and of the brainstem monoaminergic nuclei using

immunohistochemistry and then we monitored the dorsal hippocampal monoamine

transmission with *in vivo* microdialysis combined with the high-performance liquid

chromatography (HPLC) system.

Materials and Methods

Animals

All animal care and experimental procedures were performed in accordance with

protocols approved by the University of Tsukuba Animal Experiment Committee (19-378,

20-405, and 21-373) based on the NIH Guidelines for the Care and Use of Laboratory

Animals (NIH publication, 1996). Male adult Long-Evans rats ($n = 72$; 380-470 g; 10-12

weeks old at the beginning of experiments) from the TH-Cre rat colony (i.e., Cre-negative

littermate control of TH-Cre Long-Evans rats) were housed with 2–3 rats in each cage

and singly housed after surgery. Rats were kept in a controlled environment at $22 \pm 1°C$

with a 12/12-h light/dark cycle (lights on 7 a.m. to 7 p.m.) and given *ad libitum* access to

water and food (MF, Oriental Yeast Co., Ltd., Tokyo, Japan). All experiments were done

during the light cycle.

Experimental design

Four experiments were performed independently: measurement of LT (Experiment 1: n = 6), validation of exercise intensity (Experiment 2: n = 5-7/group, 3 groups), immunohistochemical analysis of neuronal activation in dorsal hippocampus and brainstem monoaminergic nuclei (Experiment 3: n = 6-8/group, 3 groups), and analysis of dorsal hippocampal monoamine release with *in vivo* microdialysis (Experiment 4: n = 6-7/group, 3 groups). All rats were handled for 1 week before the start of all procedures and habituated to treadmill running (6 days, Table 1). Then, they were randomly allocated to three groups based on LT as follows: 1) sedentary on treadmill (Control: 0 m/min), 2) below-LT light-intensity (low speed) running (Light: 15 m/min), and 3) above-LT vigorous-intensity (high speed) running (Vigorous: 25 m/min). Exercise and control conditions were maintained for 30 minutes.

Surgery to insert jugular catheter

In Experiments 1 and 2, 1 day after the habituation to treadmill running periods, the rats were anesthetized with isoflurane, and a silicone catheter was inserted into the jugular vein and fixed with a silk thread suture (37 mm). The external distal end of the catheter was fixed at the nape of the neck.

Stereotaxic surgery

In Experiment 4, after a week of handling, all rats were anesthetized with isoflurane and placed in a stereotaxic frame (Model 51900, Muromachi Kikai Co., Ltd., Tokyo, Japan). A guide cannula (AG-6, Eicom, Kyoto, Japan) was implanted above the dorsal hippocampus (AP: -3.6 mm, ML: 3.8 mm, DV: -1.8 mm, 20˚ angle, from bregma). The

cannula was secured to the skull with three anchoring screws and dental cement. To

prevent occlusion, a dummy cannula (AD-6, Eicom) was inserted into the guide cannula.

After the surgery, the rats were housed individually and were allowed to recover for at

least 2 days before habituation to treadmill running.

Habituation to treadmill running

All rats were habituated to running on a treadmill (KN-73, Natsume Seisakusho Co, Ltd.,

Tokyo, Japan) for a total of five sessions over 6 days. The running duration was 30

min/day, and the running speed was gradually increased from 5 to 25 m/min with no

incline (Nishijima and Soya 2006; Soya *et al.* 2007a). Electrical shock grids were placed

at the rear end of the treadmill and provided mild but aversive foot shocks (30 V) to rats

to encourage running at the set treadmill speed. To minimize any stress derived from the

shocks, electric shocks were limited and a gentle tail touch was given when rats did not

start running again. After the habituation period, rats were able to run at the appropriate

speeds even when the electric grid was turned off.

Treadmill running tests

On the test day, all rats fasted for 2 h before running to obtain stable metabolic conditions.

In Experiments 1 and 2, treadmill running tests were conducted 2-3 days after the surgery

for inserting the jugular catheter. In Experiment 1, an incremental exercise test was

performed to assess the running speed at LT. Rats were forced to run to exhaustion; the

running speed was started at 5 m/min and increased by 2.5 m/min every 3 minutes. The

speed at which the LT was reached was determined according to the method by Beaver *et*

al. (Beaver *et al.* 1985). In Experiment 2, a single 30-minute bout of running was

conducted to verify the validation of exercise intensity based on the speed at LT as

determined in Experiment 1. Rats were randomly subjected to run at 15 m/min (Light) or

25 m/min (Vigorous), or they were placed on a treadmill as the at-rest Control for 30

minutes. In both experiments, serial blood samples (0.1 ml) were taken at designated time

points. Blood lactate and glucose were measured using an automated glucose–lactate

analyzer (2300 Stat Plus, Yellow Springs Instruments, OH, USA).

In Experiment 3, rats were returned to their home cage after the test, rested for 90 min,

and then were transcardially perfused with 0.9% saline, followed by 4%

paraformaldehyde (PFA) in a 0.1 M phosphate buffer (PB; pH 7.4) under deep isoflurane

anesthesia. The brains were sampled for immunohistochemical analysis.

In Experiment 4, a microdialysis probe (2 mm membrane: FX-I-6-02, Eicom) was

inserted into the dorsal hippocampus using a guide cannula 1 day before the test. The

probe was connected on a treadmill and perfused with Ringer's solution (147 mM NaCl,

4 mM KCl, and 2.3 mM CaCl2) at 1.0 μl/min 2-3 h before the start of the test because a

stable dialysate monoamine level is usually obtained 2 h after insertion of the probe.

Using a fraction collector (EFC-82, Eicom), 15 μl of dialysate was collected every

15 minutes and 5 μl of 0.1M PB (pH 3.5) was added to ensure stable monoamine detection

using HPLC with an electrochemical detector (HPLC-ECD). At the end of the

experiments, rats were transcardially perfused with 0.9% saline, followed by 4% PFA in

a 0.1 M PB under deep isoflurane anesthesia and the brain was collected to verify the

position of the probe in coronal sections with DAPI (4',6-diamidino-2-phenylindole).

Please note that 7 rats in Experiment 4 were excluded from the analysis for the following

reasons: failed insertion of the probe into the dorsal hippocampus (1 rat), detachment of

the fixed probe during running (3 rats), and failed detection of peaks in chromatography

(3 rats).

Immunohistochemistry

The brain was post-fixed for 24 h in 4% paraformaldehyde (PFA, 02890-45, Nacalai

Tesque, Tokyo, Japan) in a 0.1 M PB and cryoprotected by immersion in a 0.1 M PB

containing 20% sucrose for 1 day and 30% sucrose for 2-3 days. Frozen coronal brain

sections of 30 μm thickness were cut with a microtome (REM-710, Yamato Kohki

Industrial Co., Ltd., Saitama. Japan). Sections were washed with a 0.1M PB and blocked

with a 0.1 M PB containing 1% Triton X-100 (35501-15, Nacalai Tesque) (PBT) plus 1%

bovine serum albumin (01863-77, BSA; Nacalai Tesque). Then, slices were incubated

with the designated primary antibodies in PBT-BSA for 24 h at 4 °C. The primary

antibodies used in this study were rabbit polyclonal antibody against c-Fos (1:5,000; Cat#

226-003, RRID: AB_2231974, Synaptic Systems, GAU, Germany), mouse monoclonal

antibody against NeuN (1:1,000; Cat# MAB377, RRID: AB_2298772, Millipore, MA,

USA), mouse monoclonal antibody against tyrosine hydroxylase (1:2,000; Cat# F11:sc-

25269, RRID: AB_628422, Santa Cruz, CA, USA), and goat polyclonal antibody against

5-HT (1:1,000; Cat# 20079, RRID: AB_572262, Immunostar Inc., WI, USA). After

repeated washings with PBT-BSA, sections were incubated in a dark space with

designated secondary antibodies diluted in PBT-BSA for >16 h at 4 °C. The secondary

antibodies used in this study were Alexa Fluor 488-conjugated goat anti-rabbit IgG

(1:5,000; Cat# A-11008, RRID: AB_143165, Invitrogen; Thermo Fisher Scientific Inc.,

MA, USA), Alexa Fluor 488-conjugated donkey anti-rabbit IgG (1:5,000; Cat# ab150073,

RRID: AB_2636877, abcam, Cambridge, UK), Alexa Fluor 594-conjugated donkey anti-

goat IgG (1:1,000; Cat# A-11058, RRID: AB_142540, Invitrogen; Thermo Fisher

Scientific Inc., MA, USA), and AMCA AffiniPure donkey anti-mouse IgG (1:1,000; Cat#

715-155-150, RRID: AB_2340806, Jackson ImmunoResearch Laboratories, Inc., PA,

USA). For DAPI staining to verify the location of microdialysis probes, slices were

washed three times in a 0.1 M PB and then incubated with DAPI (1:1,000; 28718-90-3,

Dojindo Laboratory, Kumamoto, Japan) in PBT-BSA for 10 minutes in the dark. Slices

were washed three times in a 0.1 M PB, mounted on subbed slides, air dried, and

coverslipped using a mounting medium.

Counts and quantification

All fluorescence z-stacked images were examined on an all-in-one microscope (BZ-X710,

Keyence, Osaka, Japan) with a 20× objective lens. Manual counts of neurons were

performed by an observer blinded to the group allocations. The number of c-Fos and

NeuN double-positive neurons ($c\text{-Fos}^+/\text{NeuN}^+$) in the dorsal hippocampal dentate gyrus

(DG), cornu ammonis (CA) 1, CA2, and CA3 sub-regions were counted hemilaterally in

4 coronal sections (from Bregma: -3.6, -4.2, -5.2, -5.8). Total neuron numbers were

normalized based on NeuN^+ region volume (mm^3), which was calculated by the area

(mm^2) multiplied by the thickness of sections (30 μm) using a BZ-X analyzer (Keyence).

The numbers of c-Fos and TH or 5-HT double-positive neurons ($c\text{-Fos}^+/\text{TH}^+$ or $c\text{-Fos}^+/5\text{-}$

HT^+) were bilaterally counted 120 μm apart throughout the LC and 360 μm apart

throughout the VTA, SNpc, RRF, DRN, and MRN, all of which are known as

monoaminergic nuclei projecting to the dorsal hippocampus.

HPLC-ECD for monoamine detection in dorsal hippocampus

Dialysate samples collected from the probe at the dorsal hippocampus were injected into

an HPLC-ECD system (HTEC-500; Eicom) using an autosampler (M-510; Eicom) and

analyzed for NA, DA, and 5-HT concentration using the HPLC system with an

EICOMPAK CAX column (2.0 mm, i.d. × 200 mm; Eicom) and a graft electrode (WE-

3G; Eicom) set at 450 mV (vs Ag/AgCl reference electrode). The flow rate and injection

volume were set to 250 μl/min and 17 μl, respectively. The mobile phase contained a

0.1 M ammonium acetate buffer, 0.05 mg/l sodium sulfate, 50 mg/l EDTA, and methanol

(7:3, v/v), with a pH of 6.0. Data analysis was performed using Eicom EPC-710 software.

Basal concentrations of the three monoamines (i.e., NA, DA, and 5-HT) were estimated

from the averages at four HPLC time points before the beginning of running.

Statistical analyses

All data are expressed as mean ± standard error (SEM) and were analyzed using R

software (version 4.2.3). Assessment of the normality (Shapiro-Wilk test) of the data,

equality of variance (Levene's test), and sphericity (Mendoza's test of sphericity) for

repeated measures (RM) were performed. Statistical significance for parametric data was

analyzed using one-way analysis of variance (ANOVA) followed by Shaffer's *post-hoc*

test, but Welch's ANOVA was performed if equality of variance was not assumed. For the

analysis of non-parametric data, the Kruskal-Wallis test with the Mann-Whitney U test

adjusted by Holm's post-hoc test was applied. For RM, two-way ANOVA followed by

Shaffer's post-hoc test was applied and the degree of freedom was adjusted using Huynh-

Feldt-Lecoutre's epsilon for non-spherical data. Pearson's correlation coefficient was

used for analyzing the strength of relationships. No tests for outliers were performed.

Statistical significance was assumed at $p < 0.05$. No statistical methods were used to

predetermine sample size, but the sample sizes were comparable to those reported in

previous studies (Soya *et al.* 2007a; Tsai *et al.* 2019; Rodovalho *et al.* 2020; Goekint *et*

al. 2012; Lee *et al.* 2003).

Results

Measurement of LT and validation of exercise intensity

Initially, we developed a varied-intensity treadmill running model with adult male Long-

Evans rats using LT as a signpost for exercise-induced stress. Our analysis revealed that

the running speed at the LT of rats was 21.87 m/min ± 1.02 (*n* = 6; Fig. 1B, C). Thus, we

defined 15 m/min, which is below the LT, as the speed for light-intensity running and 25

m/min, which is above the LT, as the speed for vigorous-intensity running. The

observation of a significant increase in blood lactate concentration after a 30-minute

session of vigorous-intensity running (two-way RM ANOVA, Shaffer's *post hoc* test,

group and time interaction, $F_{(2, 15)} = 66.036$, $p < 0.001$, Fig 1E) was sufficient to validate

the varied-intensity running model based on LT.

Light-intensity-running-induced neuronal activation of dorsal hippocampus

We assessed dorsal hippocampal neuronal activation during different intensities of

running using c-Fos immunostaining. Representative images in Figure 2B demonstrate

that c-Fos expression in dorsal hippocampal sub-regions, including the DG, CA1, and

CA3, significantly increased regardless of running speed. Conversely, the CA2 region

was not affected by running (DG: Welch's one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)}$

$_{18)} = 12.876$, $p < 0.001$, Fig. 2C; CA1: one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)}$

$= 9.994$, $p < 0.001$, Fig. 2D; CA2: one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)} =$

0.173, $p = 0.843$, Fig. 2E; CA3: one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)} = 11.256$,

$p < 0.001$, Fig. 2F).

Light-intensity running increased dorsal hippocampal NA and DA release

To assess the monoaminergic neurotransmission to the dorsal hippocampus, we measured

extracellular monoamine levels in the dorsal hippocampus using *in vivo* microdialysis and

HPLC methods. We found that treadmill running induced a significant increase of both

NA and DA levels in the dorsal hippocampus depending on running speed (NA: two-way

RM ANOVA, Shaffer's *post hoc* test, group and time interaction, $F_{(7.01, 59.58)} = 5.393$ $p <$

0.001; DA: two-way RM ANOVA, Shaffer's *post hoc* test, group and time interaction, F

$_{(8.03, 68.23)} = 4.959$, $p < 0.001$, Fig. 3B, C). Notably, DA release was significantly sustained

into the post-running time period. Dorsal hippocampal 5-HT levels, on the other hand,

were not affected by running (two-way RM ANOVA, Shaffer's *post hoc* test, interaction,

$F_{(5.58, 47.46)} = 1.16$, $p = 0.343$, Fig. 3D).

Running stimulated TH$^+$ neurons in LC and VTA, but not raphe 5-HT$^+$ neurons

To examine the responsiveness of brainstem monoaminergic nuclei, we performed

double-immunostaining of c-Fos$^+$/TH$^+$ neurons for the activation of NA- and DA-

producing neurons, and c-Fos$^+$/5-HT$^+$ neurons for the activation of serotonergic neurons,

as shown in Figure 4A-C. TH$^+$ neurons in the LC (LCTH) and VTA (VTATH), which are

known as the main sources of dorsal hippocampal NA and DA, respectively, were

significantly stimulated by both light- and vigorous-intensity running (LC: one-way

ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)} = 20.908$, $p < 0.001$, Fig. 4D; VTA: one-way

ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)} = 4.992$, $p < 0.05$, Fig. 4E). However, running

did not have any impact on dopaminergic neurons in the SNpc or RRF (SNpcTH neurons

and RRFTH neurons, respectively), or on serotonergic neurons in the DRN and MRN

(DRN$^{5\text{-}HT}$ neurons and MRN$^{5\text{-}HT}$ neurons, respectively) (SNpc: Kruskal-Wallis test,

Mann-Whitney U test adjusted by Holm's *post-hoc* test, $X^2 = 4.653$, df = 2, $p = 0.097$, Fig.

4F; RRF: Kruskal-Wallis test, Mann-Whitney U test adjusted by Holm's *post-hoc* test, X^2

$= 4.494$, df = 2, $p = 0.106$, Fig. 4G; DRN: one-way ANOVA, Shaffer's *post hoc* test, $F_{(2,}$

$_{18)} = 0.832$, $p = 0.454$, Fig. 4H; MRN: one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)}$

$= 1.640$, $p = 0.222$, Fig. 4I). Note that TH$^+$ neurons were also observed in the rostral

region of the DRN (DRNTH neurons), which are reported to promote wakefulness through

the release of DA (Cho *et al.* 2017). In this study, however, running did not affect the c-

Fos expression in DRNTH neurons (Fig. S2A).

Neuronal associations between dorsal hippocampus and brainstem monoaminergic

nuclei

To investigate whether the neuronal activities of brainstem monoaminergic nuclei are

associated with dorsal hippocampal activation, we performed Pearson's correlation

analysis between c-Fos expression in the brainstem monoaminergic nuclei and in the

dorsal hippocampus. A correlation matrix heatmap that summarizes the correlation results

is shown in Figure 5. c-Fos expression in LCTH neurons exhibited a significant and strong

positive correlation within all examined dorsal hippocampal sub-regions except CA2 (*n*

$= 21$, DG: $r = 0.672$, $p < 0.001$; CA1: $r = 0.768$, $p < 0.001$; CA2: $r = 0.212$, $p = 0.357$;

CA3: $r = 0.700$, $p < 0.001$). Additionally, a significant positive correlation was found

between c-Fos expression in VTATH, SNpcTH, and RRFTH neurons and in the dorsal CA1

(VTA: *n* $= 21$, DG: $r = 0.429$, $p = 0.0521$; CA1: $r = 0.566$, $p < 0.01$; CA2: $r = 0.033$, $p =$

0.886; CA3: $r = 0.378$, $p = 0.091$; SNpc: *n* $= 21$, DG: $r = 0.257$, $p = 0.261$; CA1: $r = 0.624$,

$p < 0.01$; CA2: $r = 0.198$, $p = 0.389$; CA3: $r = 0.321$, $p = 0.155$; RRF: $n = 21$, DG: $r =$

0.251, $p = 0.273$; CA1: $r = 0.713$, $p < 0.001$; CA2: $r = 0.173$, $p = 0.454$; CA3: $r = 0.352$,

$p = 0.155$). The higher level of c-Fos expression in DRN$^{5\text{-HT}}$ neurons was significantly

correlated with lower levels of DG, CA2, and CA3 c-Fos expression ($n = 21$, DG: $r = $-

0.463, $p < 0.05$; CA1: $r = -0.204$, $p = 0.375$; CA2: $r = -0.583$, $p < 0.01$; CA3: $r = -0.573$,

$p < 0.01$), but no correlation was found in the c-Fos expression between MRN$^{5\text{-HT}}$ neurons

and dorsal hippocampal sub-regions ($n = 21$, DG: $r = -0.275$, $p = 0.228$; CA1: $r = 0.055$,

$p = 0.814$; CA2: $r = 0.076$, $p = 0.743$; CA3: $r = -0.307$, $p = 0.176$).

Discussion

Our present study provides evidence of possible noradrenergic and dopaminergic

stimulation of dorsal hippocampal activation during light exercise. These findings may

help to elucidate the neural circuit mechanisms underlying the positive effects of light

exercise on dorsal hippocampal plasticity and function. Our results show that even light

exercise increases c-Fos expression in the dorsal hippocampal subregions and induces the

release of NA and DA at the dorsal hippocampus. Additionally, LCTH neurons and VTATH

neurons were activated even under the light exercise condition, and both were positively

correlated with dorsal hippocampal activation. Meanwhile, the serotonergic system was

not affected by exercise.

In this study, we first established a varied-intensity treadmill running model for male

adult Long-Evans rats based on LT as a physiological index of running intensity and stress.

The speed at the LT was similar to that in our previous work using male adult Wistar rats

(Long-Evans: 21.87 m/min ± 1.02 vs Wistar: 20.7 m/min ± 2.1; Fig. 1B, C; (Soya *et al.*

2007a). In addition, a significant increase in blood lactate concentration was observed

following a single bout of vigorous-intensity running, but not with light-intensity running

(Fig. 1E), which is consistent with our previous reports (Soya *et al.* 2007a; Saito and Soya

2004). These results confirm the validity of our experimental model (LT-based varied-

intensity running) and demonstrate that the two running groups exhibited different

metabolic and stress-related responses.

Previous studies have reported that acute varied-intensity treadmill running activates

the dorsal hippocampus, including the DG, CA1, and CA3 sub-regions, even at low-

speeds (Oladehin and Waters 2001; Lee *et al.* 2003; Soya *et al.* 2007b). Our current results

support these findings since both light- and vigorous-intensity running increased c-Fos

expression in the dorsal DG, CA1, and CA3 sub-regions (Fig. 2C, D, F). In contrast, the

dorsal CA2 region was not activated with running (Fig. 2E). The CA2 is known as a small

hippocampal sub-region that modulates social memory and that is stimulated by novel

encounters (Chen *et al.* 2020). Since all rats were well familiarized with the experimental

conditions and there were no novel or social stimuli, it is reasonable that activity in the

CA2 region was not affected. Additionally, we examined the activation of the ventral

hippocampus following running and found an increase of c-Fos expression across all sub-

regions of the ventral hippocampus in both running groups (Fig. S1). These results

suggest that light-intensity running is sufficient to elevate the neuronal activity in the

hippocampus across the dorsal to ventral axis, which may help to explain the positive

effects of exercise on dorsal-hippocampus-related functions, such as memory, as well as

ventral-hippocampus-related anxiolytic and antidepressant effects (Otsuka *et al.* 2016;

Inoue *et al.* 2015a; Vargas *et al.* 2017).

Our *in vivo* microdialysis experiment provides the first evidence that even light-

intensity running increases the release of NA and DA at the dorsal hippocampus (Fig. 3B,

C). Given that NA and DA both perform critical roles in dorsal hippocampal neuronal

activity, plasticity, and memory process (Takeuchi *et al.* 2016; Mello-Carpes *et al.* 2016;

Coradazzi *et al.* 2016; Salvi *et al.* 2016; Takamura *et al.* 2014), our results suggest that

the dorsal hippocampal NA and DA release contributes to the light-exercise-induced

hippocampal activation and beneficial effects, such as hippocampal neurogenesis and

spatial memory performance (Soya *et al.* 2007b; Inoue *et al.* 2015b; Inoue *et al.* 2015a).

Notably, NA and DA have been recognized as important factors in memory consolidation

(Takeuchi *et al.* 2016; Mello-Carpes *et al.* 2016), as shown by the synaptic tagging and

capture hypothesis (Frey and Morris 1997; Redondo and Morris 2010), raising a

hypothesis that a single-bout of light-intensity running could facilitate memory retention

through noradrenergic and dopaminergic dorsal hippocampal activation. Intriguingly,

prior studies demonstrate that a single session of vigorous-intensity exercise elevates NA

and DA concentrations in the hippocampus and promotes novel object recognition, which

is a dorsal-hippocampus-related function, through the β adrenergic receptor and

dopamine D1 receptor, respectively (Vargas *et al.* 2017; Vargas *et al.* 2020). Although we

did not directly examine this in the present study, these findings support our hypothesis.

We confirmed that treadmill running, even at light intensity, activates LCTH and

VTATH neurons (Fig. 4D, E). Furthermore, our correlation analysis observed that LCTH

activation positively correlates with dorsal hippocampal activation, and the activities of

midbrain dopaminergic nuclei, including VTATH, were positively associated with those

of the dorsal CA1 region (Fig. 5). These results suggest that running stimulates the dorsal

hippocampal neurons through noradrenergic projection from the LC and dopaminergic

projection from the VTA. Although both nuclei are known as major sources of dorsal

hippocampal NA and DA (Kempadoo *et al.* 2016; Weitemier and McHugh 2017), recent

480 studies have shown that LC terminals co-release DA and NA to the dorsal hippocampus,

481 and that LC-DA, but not LC-NA, plays a significant role in spatial memory and context

482 learning (Kempadoo *et al.* 2016; Takeuchi *et al.* 2016; Wagatsuma *et al.* 2017;

483 Chowdhury *et al.* 2022). On the other hand, VTA[TH] neurons have sparse input to the

484 dorsal hippocampus and do not have a significant impact on spatial memory (Kempadoo

485 *et al.* 2016; Takeuchi *et al.* 2016). However, hippocampal DA from VTA[TH] neurons is still

486 reported to be involved in hippocampus-related functions, such as reward-related spatial

487 memory, aversive memory, and associative memory (McNamara *et al.* 2014; Kramar *et*

488 *al.* 2021; Lee *et al.* 2021; Tsetsenis *et al.* 2021). The significance of LC[TH] and VTA[TH]

489 neurons for the dorsal hippocampal functions remains elusive; thus, future studies using

490 viral-tracing techniques and chemogenetic/optogenetic manipulation are needed to

491 investigate the noradrenergic and dopaminergic mechanisms underlying the beneficial

492 effects of light exercise.

493   Unlike the noradrenergic and the dopaminergic systems, 5-HT release was not

494 increased at the dorsal hippocampus, nor were DRN[5-HT] or MRN[5-HT] neurons found to be

495 stimulated by treadmill running (Fig. 3D; Fig. 4H, I). However, we found a negative

496 correlation between c-Fos expression in the DG, CA2, and CA3 sub-regions and that in

497 DRN[5-HT] neurons (Fig. 4). Previous studies have shown that running increases

498 hippocampal neuronal activity and also facilitates hippocampal LTP induction (Miladi-

499 Gorji *et al.* 2014; Radahmadi *et al.* 2016; Farmer *et al.* 2004). Conversely, the

500 hippocampal 5-HT can exert inhibitory effects on basal neural activity and LTP induction

501 (Villani and Johnston 1993; Shakesby *et al.* 2002; Kulla and Manahan-Vaughan 2002).

502 Considering these findings, our results imply that running enhances hippocampal

503 neuronal activity and LTP induction through the inhibitory control of the serotonergic

inhibitory projection from DRN$^{5\text{-}HT}$ neurons to the dorsal hippocampus.

From a clinical perspective, light exercise, such as yoga, tai-chi, and slow running,

is a simple and feasible intervention accessible by most people, including patients with

mild cognitive impairment (MCI) and Alzheimer's disease (AD). Recent studies have

shown that the hippocampal NA and DA loss and degeneration of LCTH and VTATH

neurons and axons are found in AD model rodents and AD patients (Nobili *et al.* 2017;

Sala *et al.* 2021; Francis *et al.* 2012; Rorabaugh *et al.* 2017; Mehla *et al.* 2019). Notably,

selective accumulation of amyloid β and tau proteins are observed in LC neurons before

the onset or at the early stage of AD symptoms (Matchett *et al.* 2021; Betts *et al.* 2019;

Mehla *et al.* 2019; Rorabaugh *et al.* 2017). Light exercise has been reported to have a

positive effect on MCI and AD (Yoon *et al.* 2021; Meng *et al.* 2020; Vasconcelos-Filho

et al. 2021). Therefore, our findings may provide insight into understanding the

mechanisms underlying the preventive or therapeutic role of light exercise in cognitive

dysfunction conditions, such as AD.

Conclusion

In conclusion, our results reveal that even light-intensity running activates the dorsal

hippocampus and LC-noradrenergic and VTA-dopaminergic system but that it does not

affect the serotonergic system. These findings suggest a possible contribution of LC-

noradrenergic and VTA-dopaminergic projections to dorsal hippocampal activation

during light exercise, which may provide insight into the circuit mechanism that underlies

light-exercise-enhanced learning and memory.

Acknowledgments

The authors are grateful to colleagues at the Laboratory of Exercise Biochemistry and

Neuroendocrinology (Faculty of Health and Sport Sciences, University of Tsukuba,

Japan), particularly R. Shimoda for technical support and R. Kuwamizu, H. Koizumi, M.

Soya, F. Grenier, T. Ferenc, and M.J. Ikemoto for insightful scientific discussions. We

thank J.P. Johansen (RIKEN CBS) for providing the experimental animals (TH-Cre Long-

Evans rats) and discussion. We would like to express our gratitude to M. Noguchi (ELCS

English Language Consultation, Japan) for helping with the manuscript.

Funding

This research was supported in part by KAKENHI Grants-in-Aid for Scientific Research

(A) (18H04081; 21H04858) (to H.S.); KAKENHI Grants-in-Aid for Scientific Research

on Innovative Areas: Next Generation Exercise Program for Developing Motivation,

Body and Mind Performance (16H06405) (to H.S.); Japan Science and Technology

Agency (JST)-Mirai Program (JPMJMI19D5) (to H.S.); Grant-in-Aid for Japan Society

for the Promotion of Science Fellowships (20J20887) (to T.H.).

Conflicts of Interest disclosure

The authors declare no competing financial interests.

References

Bari A., Xu S., Pignatelli M., Takeuchi D., Feng J., Li Y., Tonegawa S. (2020)

Differential attentional control mechanisms by two distinct noradrenergic coeruleo-

frontal cortical pathways. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 29080–29089.

Beaver W. L., Wasserman K., Whipp B. J. (1985) Improved detection of lactate

threshold during exercise using a log-log transformation. *J. Appl. Physiol.* **59**,

1936–1940.

Betts M. J., Kirilina E., Otaduy M. C. G., Ivanov D., Acosta-Cabronero J., Callaghan

M. F., Lambert C., et al. (2019) Locus coeruleus imaging as a biomarker for

noradrenergic dysfunction in neurodegenerative diseases. *Brain* **142**, 2558–2571.

Broussard J. I., Yang K., Levine A. T., Tsetsenis T., Jenson D., Cao F., Garcia I., et al.

(2016) Dopamine Regulates Aversive Contextual Learning and Associated In Vivo

Synaptic Plasticity in the Hippocampus. *Cell Rep.* **14**, 1930–1939.

Carter M. E., Yizhar O., Chikahisa S., Nguyen H., Adamantidis A., Nishino S.,

Deisseroth K., Lecea L. De (2010) Tuning arousal with optogenetic modulation of

locus coeruleus neurons. *Nat. Neurosci.* **13**, 1526–1535.

Chen S., He L., Huang A. J. Y., Boehringer R., Robert V., Wintzer M. E., Polygalov D.,

et al. (2020) A hypothalamic novelty signal modulates hippocampal memory.

Nature, 1–5.

Cho J. R., Treweek J. B., Robinson J. E., Xiao C., Bremner L. R., Greenbaum A.,

Gradinaru V. (2017) Dorsal Raphe Dopamine Neurons Modulate Arousal and

Promote Wakefulness by Salient Stimuli. *Neuron* **94**, 1205-1219.e8.

Chowdhury A., Luchetti A., Fernandes G., Filho D. A., Kastellakis G., Tzilivaki A.,

Ramirez E. M., Tran M. Y., Poirazi P., Silva A. J. (2022) A locus coeruleus-dorsal

CA1 dopaminergic circuit modulates memory linking. *Neuron* **110**, 3374-3388.e8.

Conrad C. D. (2010) A critical review of chronic stress effects on spatial learning and

memory. *Prog. Neuro-Psychopharmacology Biol. Psychiatry* **34**, 742–755.

Cooper C. J. (1973) Anatomical and physiological mechanisms of arousal, with special

reference to the effects of exercise. *Ergonomics* **16**, 601–609.

Coradazzi M., Gulino R., Fieramosca F., Falzacappa L. V., Riggi M., Leanza G. (2016)

Selective noradrenaline depletion impairs working memory and hippocampal

neurogenesis. *Neurobiol. Aging* **48**, 93–102.

Dongen E. V. van, Kersten I. H. P., Wagner I. C., Morris R. G. M., Fernández G. (2016)

Physical Exercise Performed Four Hours after Learning Improves Memory

Retention and Increases Hippocampal Pattern Similarity during Retrieval. *Curr.*

Biol. **26**, 1722–1727.

Ekkekakis P., Hall E. E., Petruzzello S. J. (2008) The Relationship Between Exercise

Intensity and Affective Responses Demystified: To Crack the 40-Year-Old Nut,

Replace the 40-Year-Old Nutcracker! *Ann. Behav. Med.* **35**, 136–149.

Erickson K. I., Voss M. W., Prakash R. S., Basak C., Szabo A., Chaddock L., Kim J. S.,

et al. (2011) Exercise training increases size of hippocampus and improves

memory. *Proc. Natl. Acad. Sci.* **108**, 3017–3022.

Fanselow M. S., Dong H. W. (2010) Are the Dorsal and Ventral Hippocampus

Functionally Distinct Structures? *Neuron* **65**, 7–19.

Farmer J., Zhao X., Praag H. Van, Wodtke K., Gage F. H., Christie B. R. (2004) Effects

of voluntary exercise on synaptic plasticity and gene expression in the dentate

gyrus of adult male sprague-dawley rats in vivo. *Neuroscience* **124**, 71–79.

Francis B. M., Yang J., Hajderi E., Brown M. E., Michalski B., McLaurin J.,

Fahnestock M., Mount H. T. J. (2012) Reduced tissue levels of noradrenaline are

associated with behavioral phenotypes of the TgCRND8 mouse model of

Alzheimer's disease. *Neuropsychopharmacology* **37**, 1934–1944.

Frey U., Morris R. G. M. (1997) Synaptic tagging and long-term potentiation. *Nat. 1997*

3856616 **385**, 533–536.

Gasbarri A., Packard M., Sulli A., Pacitti C., Innocenzi R., Perciavalle V. (1996) The

projections of the retrorubral field A8 to the hippocampal formation in the rat. *Exp.*

Brain Res. **112**, 244–252.

Goekint M., Bos I., Heyman E., Meeusen R., Michotte Y., Sarre S. (2012) Acute

running stimulates hippocampal dopaminergic neurotransmission in rats, but has

no influence on brain-derived neurotrophic factor. *J. Appl. Physiol.* **112**, 535–541.

Gomez-Merino D., Béquet F., Berthelot M., Chennaoui M., Guezennec C. Y. (2001)

Site-dependent effects of an acute intensive exercise on extracellular 5-HT and 5-

HIAA levels in rat brain. *Neurosci. Lett.* **301**, 143–146.

Hill E. E., Zack E., Battaglini C., Viru M., Viru A., Hackney A. C. (2008) Exercise and

circulating cortisol levels: The intensity threshold effect. *J. Endocrinol. Invest.* **31**,

587–591.

Inoue K., Hanaoka Y., Nishijima T., Okamoto M., Chang H., Saito T., Soya H. (2015a)

Long-term mild exercise training enhances hippocampus-dependent memory in

rats. *Int. J. Sports Med.* **36**, 280–285.

Inoue K., Okamoto M., Shibato J., Lee M., Matsui T., Rakwal R., Soya H. (2015b)

Long-Term Mild, rather than Intense, Exercise Enhances Adult Hippocampal

Neurogenesis and Greatly Changes the Transcriptomic Profile of the

Hippocampus. *PLoS One* **10**, e0128720.

Jones B. E. (2003) Arousal systems. *Front. Biosci.* **8**, 438–451.

Kempadoo K. A., Mosharov E. V., Choi S. J., Kandel E. R., Sulzer D. (2016) Dopamine

release from the locus coeruleus to the dorsal hippocampus promotes spatial

learning and memory. *Proc. Natl. Acad. Sci.* **113**, 14835–14840.

Kim S. Y., Adhikari A., Lee S. Y., Marshel J. H., Kim C. K., Mallory C. S., Lo M., et

al. (2013) Diverging neural pathways assemble a behavioural state from separable

features in anxiety. *Nature* **496**, 219–223.

Kramar C. P., Castillo-Díaz F., Gigante E. D., Medina J. H., Barbano M. F. (2021) The

late consolidation of an aversive memory is promoted by VTA dopamine release in

the dorsal hippocampus. *Eur. J. Neurosci.* **53**, 841–851.

Kulla A., Manahan-Vaughan D. (2002) Modulation by serotonin 5-HT4 receptors of

long-term potentiation and depotentiation in the dentate gyrus of freely moving

rats. *Cereb. Cortex* **12**, 150–162.

Kuwamizu R., Yamazaki Y., Aoike N., Ochi G., Suwabe K., Soya H. (2022) Pupil-

linked arousal with very light exercise: pattern of pupil dilation during graded

exercise. *J. Physiol. Sci.* **72**, 1–9.

Lee J. Y., Jun H., Soma S., Nakazono T., Shiraiwa K., Dasgupta A., Nakagawa T., et al.

(2021) Dopamine facilitates associative memory encoding in the entorhinal cortex.

Nat. 2021 5987880 **598**, 321–326.

Lee M. C., Okamoto M., Liu Y. F., Inoue K., Matsui T., Nogami H., Soya H. (2012)

Voluntary resistance running with short distance enhances spatial memory related

to hippocampal BDNF signaling. *J. Appl. Physiol.* **113**, 1260–1266.

Lee T.-H., Jang M.-H., Shin M.-C., Lim B.-V., Kim Y.-P., Kim H., Choi H.-H., Lee K.-

S., Kim E.-H., Kim C.-J. (2003) Dependence of rat hippocampal c-Fos expression

on intensity and duration of exercise. *Life Sci.* **72**, 1421–1436.

Li Y., Luikart B. W., Birnbaum S., Chen J., Kwon C. H., Kernie S. G., Bassel-Duby R.,

Parada L. F. (2008) TrkB Regulates Hippocampal Neurogenesis and Governs

Sensitivity to Antidepressive Treatment. *Neuron* **59**, 399–412.

Luchetti A., Bota A., Weitemier A., Mizuta K., Sato M., Islam T., McHugh T. J.,

Tashiro A., Hayashi Y. (2020) Two functionally distinct serotonergic projections

into hippocampus. *J. Neurosci.* **40**, JN-RM-2724-19.

Mahar I., Bambico F. R., Mechawar N., Nobrega J. N. (2014) Stress, serotonin, and

hippocampal neurogenesis in relation to depression and antidepressant effects.

Neurosci. Biobehav. Rev. **38**, 173–192.

Matchett B. J., Grinberg L. T., Theofilas P., Murray M. E. (2021) The mechanistic link

between selective vulnerability of the locus coeruleus and neurodegeneration in

Alzheimer's disease. *Acta Neuropathol. 2021 1415* **141**, 631–650.

McMorris T. (2016) Developing the catecholamines hypothesis for the acute exercise-

cognition interaction in humans: Lessons from animal studies. *Physiol. Behav.* **165**,

291–299.

McNamara C. G., Tejero-Cantero Á., Trouche S., Campo-Urriza N., Dupret D. (2014)

Dopaminergic neurons promote hippocampal reactivation and spatial memory

persistence. *Nat. Neurosci.* **17**, 1658–1660.

Meeusen R., Piacentini M. F., Meirleir K. De (2001) *Brain microdialysis in exercise*

research.

Mehla J., Lacoursiere S. G., Lapointe V., McNaughton B. L., Sutherland R. J.,

McDonald R. J., Mohajerani M. H. (2019) Age-dependent behavioral and

biochemical characterization of single APP knock-in mouse (APPNL-G-F/NL-G-

F) model of Alzheimer's disease. *Neurobiol. Aging* **75**, 25–37.

Mello-Carpes P. B., Silva de Vargas L. da, Gayer M. C., Roehrs R., Izquierdo I. (2016)

Hippocampal noradrenergic activation is necessary for object recognition memory

consolidation and can promote BDNF increase and memory persistence.

Neurobiol. Learn. Mem. **127**, 84–92.

Meng Q., Lin M. S., Tzeng I. S. (2020) Relationship Between Exercise and Alzheimer's

Disease: A Narrative Literature Review. *Front. Neurosci.* **14**.

Miladi-Gorji H., Rashidy-Pour A., Fathollahi Y., Semnanian S., Jadidi M. (2014)

Effects of voluntary exercise on hippocampal long-term potentiation in morphine-

dependent rats. *Neuroscience* **256**, 83–90.

Morikawa R., Kubota N., Amemiya S., Nishijima T., Kita I. (2021) Interaction between

intensity and duration of acute exercise on neuronal activity associated with

depression-related behavior in rats. *J. Physiol. Sci.* **71**, 1.

Muzerelle A., Scotto-Lomassese S., Bernard J. F., Soiza-Reilly M., Gaspar P. (2016)

Conditional anterograde tracing reveals distinct targeting of individual serotonin

cell groups (B5–B9) to the forebrain and brainstem. *Brain Struct. Funct.* **221**, 535–

561.

Neeper S. A., Góauctemez-Pinilla F., Choi J., Cotman C. (1995) Exercise and brain

neurotrophins. *Nature* **373**, 109.

Nishijima T., Soya H. (2006) Evidence of functional hyperemia in the rat hippocampus

during mild treadmill running. *Neurosci. Res.* **54**, 186–191.

Nobili A., Latagliata E. C., Viscomi M. T., Cavallucci V., Cutuli D., Giacovazzo G.,

Krashia P., et al. (2017) Dopamine neuronal loss contributes to memory and

reward dysfunction in a model of Alzheimer's disease. *Nat. Commun. 2017 81* **8**,

1–14.

Okamoto M., Hojo Y., Inoue K., Matsui T., Kawato S., McEwen B. S., Soya H. (2012)

Mild exercise increases dihydrotestosterone in hippocampus providing evidence

for androgenic mediation of neurogenesis. *Proc. Natl. Acad. Sci. U. S. A.* **109**,

13100–5.

Oladehin A., Waters R. S. (2001) Location and distribution of Fos protein expression in

rat hippocampus following acute moderate aerobic exercise. *Exp. Brain Res.* **137**,

26–35.

Otsuka T., Nishii A., Amemiya S., Kubota N., Nishijima T., Kita I. (2016) Effects of

acute treadmill running at different intensities on activities of serotonin and

corticotropin-releasing factor neurons, and anxiety- and depressive-like behaviors

in rats. *Behav. Brain Res.* **298**, 44–51.

Palacios-Filardo J., Mellor J. R. (2019) Neuromodulation of hippocampal long-term

synaptic plasticity. *Curr. Opin. Neurobiol.* **54**, 37–43.

Pereira A. C., Huddleston D. E., Brickman A. M., Sosunov A. A., Hen R., McKhann G.

M., Sloan R., Gage F. H., Brown T. R., Small S. A. (2007) An in vivo correlate of

exercise-induced neurogenesis in the adult dentate gyrus. *Proc. Natl. Acad. Sci. U.

S. A.* **104**, 5638–5643.

Perri M. G., Anton S. D., Durning P. E., Ketterson T. U., Sydeman S. J., Berlant N. E.,

Kanasky W. F., Newton R. L., Limacher M. C., Daniel Martin A. (2002)

Adherence to exercise prescriptions: Effects of prescribing moderate versus higher

levels of intensity and frequency. *Heal. Psychol.* **21**, 452–458.

Poe G. R., Foote S., Eschenko O., Johansen J. P., Bouret S., Aston-Jones G., Harley C.

W., et al. (2020) Locus coeruleus: a new look at the blue spot. *Nat. Rev. Neurosci.

2020 2111* **21**, 644–659.

Praag H. van, Christie B. R., Sejnowski T. J., Gage F. H. (1999) Running enhances

neurogenesis, learning, and long-term potentiation in mice. *Proc. Natl. Acad. Sci.

U. S. A.* **96**, 13427–13431.

Radahmadi M., Hosseini N., Alaei H. (2016) Effect of exercise, exercise withdrawal,

and continued regular exercise on excitability and long-term potentiation in the

dentate gyrus of hippocampus. *Brain Res.* **1653**, 8–13.

Redondo R. L., Morris R. G. M. (2010) Making memories last: the synaptic tagging and

capture hypothesis. *Nat. Rev. Neurosci. 2011 121* **12**, 17–30.

Rodovalho G. V., Drummond L. R., Coimbra C. C. (2020) Involvement of brainstem

noradrenergic system in cutaneous heat loss during exercise. *Brain Res. Bull.* **164**,

372–379.

Rorabaugh J. M., Chalermpalanupap T., Botz-Zapp C. A., Fu V. M., Lembeck N. A.,

Cohen R. M., Weinshenker D. (2017) Chemogenetic locus coeruleus activation

restores reversal learning in a rat model of Alzheimer's disease. *Brain* **140**, 3023–

3038.

Saito T., Soya H. (2004) Delineation of responsive AVP-containing neurons to running

stress in the hypothalamus. *Am. J. Physiol. Integr. Comp. Physiol.* **286**, R484–

R490.

Sala A., Caminiti S. P., Presotto L., Pilotto A., Liguori C., Chiaravalloti A., Garibotto

V., et al. (2021) In vivo human molecular neuroimaging of dopaminergic

vulnerability along the Alzheimer's disease phases. *Alzheimer's Res. Ther.* **13**, 1–

12.

Salvi R., Steigleder T., Schlachetzki J. C. M., Waldmann E., Schwab S., Winner B.,

Winkler J., Kohl Z. (2016) Distinct effects of chronic dopaminergic stimulation on

hippocampal neurogenesis and striatal doublecortin expression in adult mice.

Front. Neurosci. **10**, 77.

Saper C. B., Chou T. C., Scammell T. E. (2001) Review. *TRENDS Neurosci.* **24**.

Saunders B. T., Richard J. M., Margolis E. B., Janak P. H. (2018) Dopamine neurons

create Pavlovian conditioned stimuli with circuit-defined motivational properties.

Nat. Neurosci. 2018 218 **21**, 1072–1083.

Schoenfeld T. J., Gould E. (2012) Stress, Stress Hormones, and Adult Neurogenesis.

Exp. Neurol. **233**, 12.

Schwarz L. A., Miyamichi K., Gao X. J., Beier K. T., Weissbourd B., Deloach K. E.,

Ren J., et al. (2015) Viral-genetic tracing of the input-output organization of a

central noradrenaline circuit. *Nature* **524**, 88–92.

Shakesby A. C., Anwyl R., Rowan M. J. (2002) Overcoming the Effects of Stress on

Synaptic Plasticity in the Intact Hippocampus: Rapid Actions of Serotonergic and

Antidepressant Agents. *J. Neurosci.* **22**, 3638.

Soya H., Mukai A., Deocaris C. C., Ohiwa N., Chang H., Nishijima T., Fujikawa T.,

Togashi K., Saito T. (2007a) Threshold-like pattern of neuronal activation in the

hypothalamus during treadmill running: Establishment of a minimum running

stress (MRS) rat model. *Neurosci. Res.* **58**, 341–348.

Soya H., Nakamura T., Deocaris C. C., Kimpara A., Iimura M., Fujikawa T., Chang H.,

McEwen B. S., Nishijima T. (2007b) BDNF induction with mild exercise in the rat

hippocampus. *Biochem. Biophys. Res. Commun.* **358**, 961–967.

Suwabe K., Byun K., Hyodo K., Reagh Z. M., Roberts J. M., Matsushita A., Saotome

K., et al. (2018) Rapid stimulation of human dentate gyrus function with acute

mild exercise. *Proc. Natl. Acad. Sci.* **115**, 10487–10492.

Takamura N., Nakagawa S., Masuda T., Boku S., Kato A., Song N., An Y., et al. (2014)

The effect of dopamine on adult hippocampal neurogenesis. *Prog. Neuro-*

Psychopharmacology Biol. Psychiatry **50**, 116–124.

Takeuchi T., Duszkiewicz A. J., Sonneborn A., Spooner P. A., Yamasaki M., Watanabe

M., Smith C. C., et al. (2016) Locus coeruleus and dopaminergic consolidation of

everyday memory. *Nature* **537**, 357–362.

Teissier A., Chemiakine A., Inbar B., Bagchi S., Ray R. S., Palmiter R. D., Dymecki S.

M., Moore H., Ansorge M. S. (2015) Activity of Raphé Serotonergic Neurons

Controls Emotional Behaviors. *Cell Rep.* **13**, 1965–1976.

Tsai S.-F., Liu Y.-W., Kuo Y.-M. (2019) Acute and long-term treadmill running

differentially induce c-Fos expression in region- and time-dependent manners in

mouse brain. *Brain Struct. Funct.*, 1–13.

Tsetsenis T., Badyna J. K., Wilson J. A., Zhang X., Krizman E. N., Subramaniyan M.,

Yang K., Thomas S. A., Dani J. A. (2021) Midbrain dopaminergic innervation of

the hippocampus is sufficient to modulate formation of aversive memories. *Proc.*

Natl. Acad. Sci. U. S. A. **118**.

Vargas L. S., Neves B. H. S. das, Roehrs R., Izquierdo I., Mello-Carpes P. B. (2017)

One-single physical exercise session after object recognition learning promotes

memory persistence through hippocampal noradrenergic mechanisms. *Behav.*

Brain Res. **329**, 120–126.

Vargas L. S., Ramires Lima K., Piaia Ramborger B., Roehrs R., Izquierdo I., Mello-

Carpes P. B. (2020) Catecholaminergic hippocampal activation is necessary for

object recognition memory persistence induced by one-single physical exercise

session. *Behav. Brain Res.* **379**, 112356.

Vasconcelos-Filho F. S. L., Rocha Oliveira L. C. da, Freitas T. B. C. de, Pontes P. A. D.

S. de, Rocha-e-Silva R. C. da, Chaves E. M. C., Silva C. G. L. da, Soares P. M.,

Ceccatto V. M. (2021) Neuroprotective mechanisms of chronic physical exercise

via reduction of β-amyloid protein in experimental models of Alzheimer's disease:

A systematic review. *Life Sci.* **275**, 119372.

Vaynman S., Ying Z., Gomez-Pinilla F. (2003) Interplay between brain-derived

neurotrophic factor and signal transduction modulators in the regulation of the

effects of exercise on synaptic-plasticity. *Neuroscience* **122**, 647–657.

Vaynman S., Ying Z., Gomez-Pinilla F. (2004) Hippocampal BDNF mediates the

efficacy of exercise on synaptic plasticity and cognition. *Eur. J. Neurosci.* **20**,

2580–2590.

Villani F., Johnston D. (1993) Serotonin inhibits induction of long-term potentiation at

commissural synapses in hippocampus. *Brain Res.* **606**, 304–308.

Wagatsuma A., Okuyama T., Sun C., Smith L. M., Abe K., Tonegawa S. (2017) Locus

coeruleus input to hippocampal CA3 drives single-trial learning of a novel context.

Proc. Natl. Acad. Sci. U. S. A. **115**, E310–E316.

Wang S. H., Redondo R. L., Morris R. G. M. (2010) Relevance of synaptic tagging and

capture to the persistence of long-term potentiation and everyday spatial memory.

Proc. Natl. Acad. Sci. U. S. A. **107**, 19537–19542.

Warden M. R., Selimbeyoglu A., Mirzabekov J. J., Lo M., Thompson K. R., Kim S. Y.,

Adhikari A., Tye K. M., Frank L. M., Deisseroth K. (2012) A prefrontal cortex–

brainstem neuronal projection that controls response to behavioural challenge. *Nat.*

2012 4927429 **492**, 428–432.

Weitemier A. Z., McHugh T. J. (2017) Noradrenergic modulation of evoked dopamine

release and pH shift in the mouse dorsal hippocampus and ventral striatum. *Brain*

Res. **1657**, 74–86.

Yamazaki Y., Suwabe K., Nagano-Saito A., Saotome K., Kuwamizu R., Hiraga T.,

Torma F., et al. (2023) A possible contribution of the locus coeruleus to arousal

enhancement with mild exercise: evidence from pupillometry and neuromelanin

imaging. *Cereb. Cortex Commun.* **4**, 1–8.

Yoon M., Yang P. S., Jin M. N., Yu H. T., Kim T. H., Jang E., Uhm J. S., Pak H. N.,

Lee M. H., Joung B. (2021) Association of Physical Activity Level With Risk of

Dementia in a Nationwide Cohort in Korea. *JAMA Netw. Open* **4**, e2138526–

e2138526.

Zhu X., Ottenheimer D., DiLeone R. J. (2016) Activity of D1/2 Receptor Expressing

Neurons in the Nucleus Accumbens Regulates Running, Locomotion, and Food

Intake. *Front. Behav. Neurosci.* **10**, 1–10.

Fig. 1 Measurement of lactate threshold and validation of exercise intensity

(A) Protocol for incremental running test. (B) A representative LT profile from the

incremental running test. (C) Speed of rats at LT was 21.87 m/min ± 1.02 ($n = 6$). (D)

Protocol for a single bout of the running test. We defined 15 m/min for light-intensity

running and 25 m/min for vigorous-intensity running. (E) Blood lactate concentration

was significantly increased only after a 30-minute session of vigorous-intensity running.

All data are expressed as mean ± SEM. $^{***}p < 0.001$ vs Control, $^{###}p < 0.001$ vs Light, n

= 5-7/group. Control: sedentary control; Light: 15 m/min running; Vigorous: 25 m/min

running; hab.: habituation.

Fig. 2 Running-induced neuronal activation at the dorsal hippocampus during

varied-intensity treadmill running.

(A) Protocol for a single bout of the running test for immunohistochemical analysis. (B)

Representative double-fluorescence images of c-Fos (green) and NeuN (blue) in the

DG, CA1, and CA3 regions of the dorsal hippocampus. Scale bars show 100 μm. c-Fos

expressions in the DG (C), CA1 (D), CA2 (E), and CA3(F) were significantly increased

with running exercise regardless of running speed (light or vigorous). All data are

expressed as mean ± SEM. $^{**}p < 0.01$, $^{***}p < 0.001$ vs Control, n = 6-8 / group. Control:

sedentary control; Light: 15 m/min running; Vigorous: 25 m/min running; hab.:

habituation.

Fig. 3 Running increases dorsal hippocampal NA and DA release, but not 5-HT

release

(A) Protocol for a single bout of the running test for dorsal hippocampal monoamine

detection with *in vivo* microdialysis. Treadmill running induced a significant increase of

both NA (B) and DA (C) levels in the dorsal hippocampus, depending on running speed,

but not 5-HT (D). (E) Representative chromatograms showing NA (peak 1), DA (peak

2), and 5-HT (peak 3) from Control (light gray), Light (blue), and Vigorous (dark gray).

(F) Representative DAPI fluorescence image of microdialysis probe position in the

dorsal hippocampus. Scale bars show 500 μm. (G) Schematic representations indicating

the location of the microdialysis probe. Each bar represents the probe position of an

individual rat, and the color of the bar corresponds to the experimental group (Control:

light gray, Light: blue, and Vigorous: dark gray). All data are expressed as mean ± SEM.

$^{*}p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$ vs Sed, $^{\#}p < 0.05$, $^{\#\#}p < 0.01$ vs Light, $n = 6$-7/group.

Control: sedentary control; Light: 15 m/min running; Vigorous: 25 m/min running; hab.:

habituation.

Fig. 4 LCTH and VTATH neurons, but not raphe 5-HT$^+$ neurons, stimulated by

running

Representative double-fluorescence images of c-Fos (green), TH (blue), and 5-HT (red)

neurons in LC (A); VTA, SNpc, and RRF (B); and DRN and MRN (C). Scale bars show

100 μm. Yellow heads show c-Fos$^+$/TH$^+$ neurons. LCTH and VTATH neurons were

significantly stimulated in both running groups (D, E). However, running did not have

an impact on SNpcTH and RRFTH neurons (F, G) or DRN^{5-HT} and MRN^{5-HT} neurons (H,

I). All data are expressed as mean ± SEM. $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$ vs Control,

$^{\#\#}p < 0.01$ vs Light, $n = 6\text{-}8$ / group. Control: sedentary control; Light: 15 m/min

running; Vigorous: 25 m/min running.

Fig. 5 Neuronal associations between dorsal hippocampus and brainstem

monoaminergic nuclei

A correlation matrix heatmap is shown to summarize the correlation analysis between c-

Fos expression in dorsal-hippocampal sub-regions and that in LC[TH], VTA[TH], SNpc[TH],

RRF[TH], DRN[5-HT], and MRN[5-HT] neurons. Colors in the scale show correlation strength.

$^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$. $n = 21$.

936   Table 1 Protocol for habituation to treadmill running

Day	Running speed (duration)
1	Rest (10 min) → 5 m/min (10 min) → 10 m/min (10 min)
2	Rest (5 min) → 5 m/min (10 min) → 10 m/min (10 min) → 15 m/min (10 min)
3	Rest
4	Rest (5 min) → 10 m/min (10 min) → 15 m/min (10 min) → 20 m/min (10 min)
5	Rest (5 min) → 15 m/min (10 min) → 20 m/min (10 min) → 25 m/min (10 min)
6	Rest (5 min) → 15 m/min (10 min) → 20 m/min (10 min) → 25 m/min (10 min)

Supporting information

Light exercise-induced dopaminergic and noradrenergic stimulation in the dorsal

hippocampus: using a rodent physiological exercise model

Taichi Hiraga, Toshiaki Hata, Masahiro Okamoto, and Hideaki Soya

Corresponding author: Hideaki Soya, Ph.D.

Email: soya.hideaki.gt@u.tsukuba.ac.jp

This PDF file includes:

・Figure S1

・Figure S2

Fig. S1. Running induces neuronal activation of the ventral hippocampus

(A) Representative double-fluorescence images of NeuN (blue) and c-Fos (green) in the

DG, CA1, and CA3. Scale bars show 100 μm. c-Fos expression in ventral hippocampal

sub-regions, including the DG (B), CA1 (C), and CA3 (D), was also significantly

increased regardless of running speed (DG: Kruskal-Wallis test, Mann-Whitney U test

adjusted by Holm's *post-hoc* test, X^2 = 10.245, df = 2, $p < 0.01$; CA1: one-way ANOVA,

Shaffer's *post hoc* test, $F_{(2, 18)}$ = 11.443, $p < 0.001$; CA3: Welch's one-way ANOVA,

Shaffer's *post hoc* test, $F_{(2, 18)}$ = 11.758, $p < 0.001$). (E) A correlation matrix heatmap is

shown to summarize the correlation results between c-Fos expression in monoaminergic

nuclei and that in ventral hippocampal sub-regions. Colors in the scale show correlation

strength. Data were analyzed using Pearson's correlation. c-Fos expression in LC[TH] and

VTA[-TH] neurons had a significant positive correlation with that in all ventral hippocampal

sub-regions (LC: n = 21, DG: r = 0.435, $p < 0.05$; CA1: r = 0.658, $p < 0.01$; CA3: r =

0.631, $p < 0.01$; VTA: n = 21, DG: r = 0.599, $p < 0.05$; CA1: r = 0.610, $p < 0.01$; CA3: r

= 0.499, $p < 0.05$). Additionally, the higher level of c-Fos expression in DRN[5-HT] neurons

was significantly correlated with the lower level of ventral CA3 c-Fos expression ($n = 21$;

DG: $r = -0.085$, $p = 0.715$; CA1: $r = -0.090$, $p = 0.697$; CA3: $r = -0.522$, $p < 0.05$). No

correlation was found in c-Fos expression between SNpc[TH], RRF[TH], or MRN[5-HT] neurons

or in ventral hippocampal sub-regions (SNpc: $n = 21$, DG: $r = 0.064$, $p = 0.783$; CA1: r

$= 0.366$, $p = 0.102$; CA3: $r = 0.286$, $p = 0.208$; RRF: $n = 21$, DG: $r = 0.389$, $p = 0.082$;

CA1: $r = 0.412$, $p = 0.063$; CA3: $r = 0.429$, $p = 0.052$; MRN: $n = 21$, DG: $r = -0.255$, $p =$

0.265; CA1: $r = -0.159$, $p = 0.492$; CA3: $r = -0.181$, $p = 0.429$). All data are expressed as

mean ± SEM. [*]$p < 0.05$, [**]$p < 0.01$, [***]$p < 0.001$ vs Control; $n = $ 6-8/group. Control:

sedentary control; Light: 15 m/min running; Vigorous: 25 m/min running.

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

Fig. S2 DRN^TH neurons not activated by running and not associated with hippocampal neuronal activation

(A) DRN^TH neurons were not activated by running (Welch's one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)}$ = 0.384, *p* =0.687). (B) A correlation matrix heatmap is shown to summarize the correlation results between c-Fos expression in DRN^TH neurons and that in hippocampal sub-regions. Colors in the scale show correlation strength. There was no significant correlation between c-Fos expression in hippocampal subregions and c-Fos expression in DRN^TH neurons (dorsal DG: *r* = -0.364, *p* = 0.105; dorsal CA1: *r* = -0.174, *p* = 0.451; dorsal CA2: *r* = -0.381, *p* = 0.088; dorsal CA3: *r* = -0.409, *p* = 0.066; ventral DG: *r* = -0.063, *p* = 0.786; ventral CA1: *r* = -0.073, *p* = 0.754; ventral CA3: *r* = -0.222, *p* = 0.333). All data are expressed as mean ± SEM. *n* = 6-8/group.