Light-exercise-induced dopaminergic and noradrenergic stimulation in the dorsal hippocampus: using a rat physiological exercise model

Taichi Hiraga[a], Toshiaki Hata[a], Shingo Soya[b], Ryo Shimoda[a],

Kanako Takahashi[a,c], Mariko Soya[a,d], Koshiro Inoue[a,e],

Joshua P. Johansen[f], Masahiro Okamoto[a,c], and Hideaki Soya[a,c]

[a] Laboratory of Exercise Biochemistry and Neuroendocrinology, Institute of Health and Sport Sciences, University of Tsukuba, Tsukuba, Ibaraki 305-8574, Japan
[b] International Institute for Integrative Sleep Medicine (WPI-IIIS), University of Tsukuba, Tsukuba, Ibaraki 305-8575 Japan
[c] Division of Sport Neuroscience, *Kokoro* Division, Advanced Research Initiative for Human High Performance (ARIHHP), Institute of Health and Sport Sciences, University of Tsukuba, Tsukuba, Ibaraki 305-8574, Japan
[d] Department of Anatomy and Neuroscience, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Fukuoka 812-8582, Japan
[e] Center for Education in Liberal Arts and Sciences, Health Sciences University of Hokkaido, Ishikari, Hokkaido 061-0293, Japan
[f] Laboratory for Neural Circuitry of Memory, RIKEN Center for Brain Science, Wako, Saitama 351-0198, Japan

ORCID ID: Taichi Hiraga https://orcid.org/0000-0002-4790-044X
ORCID ID: Hideaki Soya https://orcid.org/0000-0002-3061-4375

Running title: Light exercise activates hippocampus via NA and DA systems

Corresponding author: Hideaki Soya, Ph.D.

Address: Laboratory of Exercise Biochemistry and Neuroendocrinology; Sport Neuroscience Division, Advanced Research Initiative for Human High Performance (ARIHHP), Institute of Health and Sport Sciences, University of Tsukuba, 1-1-1 Tennoudai, Tsukuba, Ibaraki 305-8574, Japan

Telephone/Fax: +81-(0)29-853-2620

Email: soya.hideaki.gt@u.tsukuba.ac.jp

Abbreviations:

5-HT: serotonin

AAS: ascending arousal system

ACTH: adrenocorticotropic hormone

AD: Alzheimer's disease

BDNF: brain-derived neurotrophic factor

BSA: bovine serum albumin

CA: cornu ammonis

DA: dopamine

DAPI: 4',6-diamidino-2-phenylindole

DG: dentate gyrus

DRN: dorsal raphe nucleus

HPLC-ECD: high-performance liquid chromatography with an electrochemical detector

IGF-I: insulin-like growth factor I

LC: locus coeruleus

LTD: long-term depression

LTP: long-term potentiation

LT: lactate threshold

MCI: mild cognitive impairment

MRN: median raphe nucleus

NA: noradrenaline

PB: phosphate buffer

PBT: phosphate buffer with Triton X-100

PFA: paraformaldehyde

PRP: plasticity-related protein

PVN: paraventricular nucleus

RM: repeated measures

RRF: retrorubral field

SNpc: substantia nigra pars compacta

TH: tyrosine hydroxylase

VTA: ventral tegmental area

67 **Graphical abstract**

Possible circuit mechanism of light-exercise-induced hippocampal activation

CONCLUSION
Light exercise could activate the dorsal hippocampus through LC-NA and VTA-DA stimulation.

Our previous research with animals and humans has demonstrated that even light exercise

can boost neuronal activity in the dorsal hippocampus and improve memory. While the

mechanism underlying this remains undetermined, recent studies suggest the involvement

of the ascending monoaminergic system. Here, we examined this hypothesis and found a

possible contribution of noradrenergic neurons in the locus coeruleus and dopaminergic

neurons in the ventral tegmental area to dorsal hippocampal activation during light

exercise, implying a circuit mechanism for light-exercise-enhanced memory.

(80 / 80 words)

Abstract

Exercise activates the dorsal hippocampus, which triggers the synaptic and cellar

plasticity and ultimately promotes memory formation. For decades, these benefits have

been explored using demanding and stress-response-inducing exercise at moderate-to-

vigorous intensities. In contrast, our translational research with animals and humans has

focused on light exercise below the lactate threshold (LT), which almost anyone can

safely perform with minimal stress, and found that even light exercise can stimulate

hippocampal activity and enhance memory performance. Although the circuit mechanism

of this boost remains unclear, arousal promotion even with light exercise implies the

involvement of the ascending monoaminergic system, which is essential to modulate

hippocampal activity and impact memory. To examine this hypothesis, we employed our

physiological exercise model based on the LT of rats that can be applied to human and

immunohistochemically assessed the neuronal activation of the dorsal hippocampal sub-

regions and brainstem monoaminergic neurons. Also, we monitored the dynamics of

monoamine release at the dorsal hippocampus using *in vivo* microdialysis. We found that

even light exercise increased neuronal activity in the dorsal hippocampal sub-regions and

induced noradrenaline and dopamine release. Furthermore, we found that tyrosine

hydroxylase-positive neurons in the locus coeruleus (LC) and the ventral tegmental area

(VTA) were activated even by light exercise and were both positively correlated with the

dorsal hippocampal activation. In conclusion, our findings demonstrate that light exercise

stimulates hippocampal neurons, possibly through the LC-noradrenergic and/or VTA-

dopaminergic neurons. This sheds light on the circuit mechanisms responsible for

hippocampal neural activation during exercise, consequently enhancing memory function.

(250/250 words)

Keywords Light exercise; Dorsal hippocampus; Locus coeruleus; Ventral tegmental area;

Noradrenaline; Dopamine

1 Introduction

There is accumulating evidence showing that physical exercise has beneficial effects on the brain and its functions, including hippocampus-dependent learning and memory.[1–4] The hippocampus is well-known for being involved in several processes related to emotion, stress resilience, and cognitive function. In particular, the dorsal part of the hippocampus plays a key role in the learning process and memory formation.[5] Physical exercise has been found to stimulate the activity of the dorsal hippocampal neurons.[6–8] This subsequently triggers the production of plasticity-related proteins (PRPs) such as the brain-derived neurotrophic factor (BDNF) in the hippocampus.[9–11] Also, this activation increases cerebral blood flow[8,12] and facilitates the entrance of bioactive hormones "exerkines" secreted from peripheral organs, such as the insulin-like growth factor I (IGF-I), cathepsin B, and irisin, across the blood-brain barrier into the hippocampus.[13–18] In turn, these boosts ultimately contribute to cellar and synaptic plasticity and memory enhancement.[1,15,16,19–21]

For decades, the benefits of exercise to learning and memory have been based on evidence using exercise at moderate-to-vigorous intensities,[2–4,22] which is generally recommended in exercise therapy. However, such demanding exercise can cause physiological stress and reduce exercise adherence.[23–27] Besides, while the chronic intervention of voluntary wheel running is widely used in animal research,[1,15,16,18] this exercise model provides less information about effective exercise intensity, frequency, and duration for memory enhancement. Moreover, voluntary running rodents run at a relatively high intensity (45 m/min) and exhibit intermittent running patterns in the dark cycle,[28] suggesting that wheel running is not an appropriate translational exercise model for human exercise. We thus have been employing a rodent treadmill running model based

on physiological responses such as lactate threshold (LT). LT is the signpost for exercise-

induced stress, which begins with the onset of lactate accumulation in the blood and the

accumulation of adrenocorticotropic hormone (ACTH).[23,25] This physiological exercise

model enables us to define running below the LT as light exercise with minimum

physiological stress and running above the LT as vigorous, stress-inducing exercise

stimuli. Using this model, we found that even light exercise was sufficient to acutely

stimulate hippocampal activity and BDNF induction,[9] and to promote hippocampal

neurogenesis and spatial memory.[29–31] Conversely, excessive ACTH secretion induced by

vigorous exercise cancels out these beneficial effects on hippocampal plasticity and

memory due to the stress-induced hippocampal vulnerability.[29–33] Additionally, our

research with human participants and functional magnetic resonance imaging (fMRI)

found that acute light exercise increases hippocampal activity and enhances pattern

separation, which is a hippocampus-related function.[34] Taken together, our translational

research demonstrates the importance of exercise intensity for determining the effect of

exercise on the hippocampus and highlights the advantages of light exercise.

However, prior studies, including our own findings, have yet to address the circuit

mechanism underlying dorsal hippocampal activation by light exercise and its beneficial

effect on memory performance. Classically, it is known that exercise can promote

arousal,[35–37] which is thought to originate from the activation of the ascending arousal

system (AAS).[38,39] This is supported by our recent studies showing that arousal promotion

can occur by exercise even at very light-intensity accompanied by pupil dilation,[34,40–43]

which reflects the activity of AAS.[44–48] The AAS sends out widespread projections in

order to increase global brain activity, including the dorsal hippocampus, which largely

overlaps the monoaminergic neuromodulatory system. Monoamines like noradrenaline

(NA), dopamine (DA), and serotonin (5-HT) are essential for multiple functions such as arousal, attention, reward, motivation, and mood regulation.[49–55] At the dorsal hippocampus, monoaminergic stimulation facilitates learning and memory processes by regulating the expression of PRPs and synaptic and cellar plasticity, such as long-term potentiation (LTP) and neurogenesis.[56–61] The hippocampal monoamines are mainly synthesized and provided by the specific nuclei located in the pons noradrenergic locus coeruleus (LC),[62–66] the midbrain dopaminergic nuclei, such as the ventral tegmental area (VTA) and the substantia nigra pars compacta (SNpc),[67–71] and the midbrain serotonergic raphe nuclei (the dorsal raphe: DRN and the median raphe: MRN).[72,73] Previous research demonstrate that single bouts of exercise activate these brainstem monoaminergic neurons[74–76] and affect the hippocampal monoamine transmissions,[3,77,78] thereby contributing to the enhancement of hippocampus-related functions.[3,74,79] These findings lead us to the hypothesis that the monoamines provided by the brainstem nuclei could be involved in the light exercise-induced dorsal hippocampal activation, as the neural basis of memory enhancement.

To test this hypothesis, the involvement of the brainstem monoaminergic system in dorsal hippocampal activation during light exercise was investigated using a rat LT-based treadmill running model that can be applied to humans. We analyzed the neuronal activation of the dorsal hippocampus and of the brainstem monoaminergic neurons using immunohistochemistry. Also, we monitored the dorsal hippocampal monoamine transmission with *in vivo* microdialysis combined with the high-performance liquid chromatography (HPLC) system.

2 Materials and Methods

2.1 Animals

All animal care and experimental procedures were performed in accordance with protocols approved by the University of Tsukuba Animal Experiment Committee (19-378, 20-405, and 21-373) based on the NIH Guidelines for the Care and Use of Laboratory Animals (NIH publication, 1996). Male adult Long-Evans rats ($n = 72$; 380-470 g; 10-12 weeks old at the beginning of experiments) from the TH-Cre rat colony (i.e., Cre-negative littermate control of TH-Cre Long-Evans rats) were housed with 2–3 rats in each cage and singly housed after surgery. Rats were kept in a controlled environment at $22 \pm 1°C$ with a 12/12-h light/dark cycle (lights on 7 a.m. to 7 p.m.) and given *ad libitum* access to water and food (MF, Oriental Yeast Co., Ltd., Tokyo, Japan). All experiments were done during the light cycle.

2.2 Experimental design

Four experiments were performed independently: measurement of LT (Experiment 1: $n = 6$), validation of exercise intensity (Experiment 2: $n = $ 5-7/group, 3 groups), immunohistochemical analysis of neuronal activation in dorsal hippocampus and brainstem monoaminergic nuclei (Experiment 3: $n = $ 6-8/group, 3 groups), and analysis of dorsal hippocampal monoamine release with *in vivo* microdialysis (Experiment 4: $n = $ 6-7/group, 3 groups). All rats were handled for 1 week before the start of all procedures and habituated to treadmill running (6 days, Table 1). Then, they were randomly allocated to three groups based on LT as follows: 1) sedentary on treadmill (Control: 0 m/min), 2) below-LT light-intensity (low speed) running (Light: 15 m/min), and 3) above-LT vigorous-intensity (high speed) running (Vigorous: 25 m/min). Exercise and control conditions were maintained for 30 minutes.

204 Table 1 Protocol for habituation to treadmill running

Day	Running speed (duration)
1	Rest (10 min) → 5 m/min (10 min) → 10 m/min (10 min)
2	Rest (5 min) → 5 m/min (10 min) → 10 m/min (10 min) → 15 m/min (10 min)
3	Rest
4	Rest (5 min) → 10 m/min (10 min) → 15 m/min (10 min) → 20 m/min (10 min)
5	Rest (5 min) → 15 m/min (10 min) → 20 m/min (10 min) → 25 m/min (10 min)
6	Rest (5 min) → 15 m/min (10 min) → 20 m/min (10 min) → 25 m/min (10 min)

2.3 Surgery to insert jugular catheter

In Experiments 1 and 2, 1 day after the habituation to treadmill running periods, the rats

were anesthetized with isoflurane, and a silicone catheter was inserted into the jugular

vein and fixed with a silk thread suture (37 mm). The external distal end of the catheter

was fixed at the nape of the neck. After the surgery, the rats were injected with antibiotics

(Mycillin Sol; Meiji Seika, Tokyo, Japan). Then, they were housed individually and were

allowed to recover for 2 or 3 days before the treadmill running test.

2.4 Stereotaxic surgery

In Experiment 4, after a week of handling, all rats were anesthetized with isoflurane and

placed in a stereotaxic frame (Model 51900, Muromachi Kikai Co., Ltd., Tokyo, Japan).

A guide cannula (AG-6, Eicom, Kyoto, Japan) was implanted above the dorsal

hippocampus (AP: -3.6 mm, ML: 3.8 mm, DV: -1.8 mm, 20° angle, from bregma). The

cannula was secured to the skull with three anchoring screws and dental cement. To

prevent occlusion, a dummy cannula (AD-6, Eicom) was inserted into the guide cannula.

After the surgery, the rats were injected with antibiotics (Mycillin Sol; Meiji Seika, Tokyo,

Japan). Then, they were housed individually and were allowed to recover for at least 2

days before habituation to treadmill running.

2.5 Habituation to treadmill running

All rats were habituated to running on a treadmill (KN-73, Natsume Seisakusho Co, Ltd.,

Tokyo, Japan) for a total of five sessions over 6 days. The running duration was 30

min/day, and the running speed was gradually increased from 5 to 25 m/min with no

incline [8,23]. Electrical shock grids were placed at the rear end of the treadmill and provided

mild but aversive foot shocks (30 V) to rats to encourage running at the set treadmill

speed. To minimize any stress derived from the shocks, electric shocks were limited and

a gentle tail touch was given when rats did not start running again. After the habituation

period, rats were able to run at the appropriate speeds even when the electric grid was

turned off.

2.6 Treadmill running tests

On the test day, all rats fasted for 2 h before running to obtain stable metabolic conditions.

In Experiments 1 and 2, treadmill running tests were conducted 2-3 days after the surgery

for inserting the jugular catheter. In Experiment 1, an incremental exercise test was

performed to assess the running speed at LT. Rats were forced to run to exhaustion; the

running speed was started at 5 m/min and increased by 2.5 m/min every 3 minutes. The

speed at which the LT was reached was determined according to the method by Beaver *et*

al..[80] In Experiment 2, a single 30-minute bout of running was conducted to verify the

validation of exercise intensity based on the speed at LT as determined in Experiment 1.

Rats were randomly subjected to run at 15 m/min (Light) or 25 m/min (Vigorous), or they

were placed on a treadmill as the at-rest Control for 30 minutes. In both experiments,

serial blood samples (0.1 ml) were taken at designated time points. Blood lactate and

glucose were measured using an automated glucose–lactate analyzer (2300 Stat Plus,

Yellow Springs Instruments, OH, USA).

In Experiment 3, rats were returned to their home cage after the test, rested for 90 min,

and then were transcardially perfused with 0.9% saline, followed by 4%

paraformaldehyde (PFA) in a 0.1 M phosphate buffer (PB; pH 7.4) under deep isoflurane

anesthesia. The brains were sampled for immunohistochemical analysis.

In Experiment 4, a microdialysis probe (2 mm membrane: FX-I-6-02, Eicom) was

inserted into the dorsal hippocampus using a guide cannula 1 day before the test. The

probe was connected to the microsyringe pump (ESP-32, Eicom) using a Teflon tubing

(JT-10, Eicom) and perfused with Ringer's solution (147 mM NaCl, 4 mM KCl, and

2.3 mM CaCl2) at 1.0 μl/min 2-3 h before the start of the test because a stable dialysate

monoamine level is usually obtained 2 h after insertion of the probe. Using a fraction

collector (EFC-82, Eicom), 15 μl of dialysate was collected every 15 minutes and 5 μl of

0.1M PB (pH 3.5) was added to ensure stable monoamine detection using HPLC with an

electrochemical detector (HPLC-ECD). At the end of the experiments, rats were

transcardially perfused with 0.9% saline, followed by 4% PFA in a 0.1 M PB under deep

isoflurane anesthesia and the brain was collected to verify the position of the probe in

coronal sections with DAPI (4',6-diamidino-2-phenylindole). Please note that 7 rats in

Experiment 4 were excluded from the analysis for the following reasons: failed insertion

of the probe into the dorsal hippocampus (1 rat), detachment of the fixed probe during

running (3 rats), and failed detection of peaks in chromatography (3 rats).

2.7 Immunohistochemistry

The brain was post-fixed for 24 h in 4% paraformaldehyde (PFA, 02890-45, Nacalai

Tesque, Tokyo, Japan) in a 0.1 M PB and cryoprotected by immersion in a 0.1 M PB

containing 20% sucrose for 1 day and 30% sucrose for 2-3 days. Frozen coronal brain

sections of 30 μm thickness were cut with a microtome (REM-710, Yamato Kohki

Industrial Co., Ltd., Saitama. Japan). Sections were washed with a 0.1M PB and blocked

with a 0.1 M PB containing 1% Triton X-100 (35501-15, Nacalai Tesque) (PBT) plus 1%

bovine serum albumin (01863-77, BSA; Nacalai Tesque). Then, slices were incubated

with the designated primary antibodies in PBT-BSA for 24 h at 4 °C. The primary

antibodies used in this study were rabbit polyclonal antibody against c-Fos (1:5,000; Cat#

226-003, RRID: AB_2231974, Synaptic Systems, GAU, Germany), mouse monoclonal

antibody against NeuN (1:1,000; Cat# MAB377, RRID: AB_2298772, Millipore, MA,

USA), mouse monoclonal antibody against tyrosine hydroxylase (1:2,000; Cat# F11:sc-

25269, RRID: AB_628422, Santa Cruz, CA, USA), and goat polyclonal antibody against

5-HT (1:1,000; Cat# 20079, RRID: AB_572262, Immunostar Inc., WI, USA). After

repeated washings with PBT-BSA, sections were incubated in a dark space with

designated secondary antibodies diluted in PBT-BSA for >16 h at 4 °C. The secondary

antibodies used in this study were Alexa Fluor 488-conjugated goat anti-rabbit IgG

(1:5,000; Cat# A-11008, RRID: AB_143165, Invitrogen; Thermo Fisher Scientific Inc.,

MA, USA), Alexa Fluor 488-conjugated donkey anti-rabbit IgG (1:5,000; Cat# ab150073,

RRID: AB_2636877, abcam, Cambridge, UK), Alexa Fluor 594-conjugated donkey anti-

goat IgG (1:1,000; Cat# A-11058, RRID: AB_142540, Invitrogen; Thermo Fisher

Scientific Inc., MA, USA), and AMCA AffiniPure donkey anti-mouse IgG (1:1,000; Cat#

715-155-150, RRID: AB_2340806, Jackson ImmunoResearch Laboratories, Inc., PA,

USA). For DAPI staining to verify the location of microdialysis probes, slices were

washed three times in a 0.1 M PB and then incubated with DAPI (1:1,000; 28718-90-3, Dojindo Laboratory, Kumamoto, Japan) in PBT-BSA for 10 minutes in the dark. Slices were washed three times in a 0.1 M PB, mounted on subbed slides, air dried, and coverslipped using a mounting medium.

2.8 Counts and quantification

All fluorescence z-stacked images were examined on an all-in-one microscope (BZ-X710, Keyence, Osaka, Japan) with a 20× objective lens. Manual counts of neurons were performed by an observer blinded to the group allocations. The number of c-Fos and NeuN double-positive neurons (c-Fos$^+$/NeuN$^+$) in the dorsal hippocampal dentate gyrus (DG), cornu ammonis (CA) 1, CA2, and CA3 sub-regions were counted hemilaterally in 4 coronal sections (from Bregma -3.60 mm, -4.16 mm, -5.20 mm, -5.80 mm). Total neuron numbers were normalized based on NeuN$^+$ region volume (mm^3), which was calculated by the area (mm^2) multiplied by the thickness of sections (30 μm) using a BZ-X analyzer software (https://www.keyence.co.jp/products/microscope/fluorescence-microscope/bz-x700/models/bz-h3a/, version 1.4.1.1, RRID:SCR_017375). The numbers of c-Fos and TH or 5-HT double-positive neurons (c-Fos$^+$/TH$^+$ or c-Fos$^+$/5-HT$^+$) were bilaterally counted 120 μm apart throughout the LC (around Bregma -9.16 mm ~ 10.52 mm, 10 ~ 16 coronal sections) and 360 μm apart throughout the VTA, SNpc (around Bregma -4.80 mm ~ -6.72 mm, 5 coronal sections), DRN, and MRN (around Bregma -7.30 mm ~ -8.80 mm, 5 coronal sections), all of which are known as monoaminergic nuclei projecting to the dorsal hippocampus.

2.9 HPLC-ECD for monoamine detection in the dorsal hippocampus

Dialysate samples collected from the probe at the dorsal hippocampus were injected into

an HPLC-ECD system (HTEC-500; Eicom) using an autosampler (M-510; Eicom) and

analyzed for NA, DA, and 5-HT concentration using the HPLC system with an

EICOMPAK CAX column (2.0 mm, i.d. × 200 mm; Eicom) and a graft electrode (WE-

3G; Eicom) set at 450 mV (vs Ag/AgCl reference electrode). The flow rate and injection

volume were set to 250 μl/min and 17 μl, respectively. The mobile phase contained a

0.1 M ammonium acetate buffer, 0.05 mg/l sodium sulfate, 50 mg/l EDTA, and methanol

(7:3, v/v), with a pH of 6.0. Data analysis was performed using Eicom EPC-710 software.

Basal concentrations of the three monoamines (i.e., NA, DA, and 5-HT) were estimated

from the averages at four HPLC time points before the beginning of running.

2.10 Statistical analyses

All data are expressed as mean ± standard error (SEM) and figures are generated using

GraphPad Prism (https://www.graphpad.com/, version 8, RRID:SCR_002798, GraphPad

Software, San Diego, CA, USA). All data were analyzed using R software (https://www.r-

project.org/, version 4.2.3, RRID:SCR_001905). Assessment of the normality (Shapiro-

Wilk test) of the data, equality of variance (Levene's test), and sphericity (Mendoza's test

of sphericity) for repeated measures (RM) were performed. Statistical significance for

parametric data was analyzed using one-way analysis of variance (ANOVA) followed by

Shaffer's *post-hoc* test, but Welch's ANOVA was performed if equality of variance was

not assumed. For the analysis of non-parametric data, the Kruskal-Wallis test with the

Mann-Whitney U test adjusted by Holm's post-hoc test was applied. For RM, two-way

ANOVA followed by Shaffer's post-hoc test was applied and the degree of freedom was

adjusted using Huynh-Feldt-Lecoutre's epsilon for non-spherical data. Pearson's

correlation coefficient was used for analyzing the strength of relationships. No tests for

outliers were performed. Statistical significance was assumed at $p < 0.05$. No statistical

methods were used to predetermine sample size, but the sample sizes were comparable to

those reported in previous studies.[7,23,75–77]

3 Results

3.1 Measurement of LT and validation of exercise intensity

Initially, we developed a varied-intensity treadmill running model with adult male Long-

Evans rats using LT as a signpost for exercise-induced stress. Our analysis revealed that

the running speed at the LT of rats was 21.87 m/min ± 1.02 ($n = 6$; Fig. 1B, C). Thus, we

defined 15 m/min, which is below the LT, as the speed for light-intensity running and 25

m/min, which is above the LT, as the speed for vigorous-intensity running. The

observation of a significant increase in blood lactate concentration after a 30-minute

session of vigorous-intensity running (two-way RM ANOVA, Shaffer's *post hoc* test,

group and time interaction, $F_{(2, 15)} = 66.036$, $p < 0.001$, Fig 1E) was sufficient to validate

the varied-intensity running model based on LT.

3.2 Light-intensity-running-induced neuronal activation of dorsal hippocampus

We assessed dorsal hippocampal neuronal activation during different intensities of

running using c-Fos immunostaining. Representative images in Figure 2B demonstrate

that c-Fos expression in dorsal hippocampal sub-regions, including the DG, CA1, and

CA3, significantly increased regardless of running speed. Conversely, the CA2 region

was not affected by running (DG: Welch's one-way ANOVA, Shaffer's *post hoc* test, $F_{(2,}$

$_{18)} = 12.876$, $p < 0.001$, Fig. 2C; CA1: one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)}$

= 9.994, $p < 0.001$, Fig. 2D; CA2: one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)} =$

0.173, $p = 0.843$, Fig. 2E; CA3: one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)} = 11.256$,

$p < 0.001$, Fig. 2F).

3.3 Light-intensity running increased dorsal hippocampal NA and DA release

To assess the monoaminergic neurotransmission to the dorsal hippocampus, we measured

extracellular monoamine levels in the dorsal hippocampus using *in vivo* microdialysis and

HPLC methods. We found that treadmill running induced a significant increase of both

NA and DA levels in the dorsal hippocampus depending on running speed (NA: two-way

RM ANOVA, Shaffer's *post hoc* test, group, $F_{(2, 17)} = 11.545$, $p < 0.001$, time, $F_{(3.5, 59.58)}$

$= 16.236$, $p < 0.001$, group and time interaction, $F_{(7.01, 59.58)} = 5.393$, $p < 0.001$; DA: two-

way RM ANOVA, Shaffer's *post hoc* test, group, $F_{(2, 17)} = 18.720$, $p < 0.001$, time, $F_{(4.01,}$

$_{68.23)} = 14.019$, $p < 0.001$, group and time interaction, $F_{(8.03, 68.23)} = 4.959$, $p < 0.001$, Fig.

3E, F). Notably, DA release was significantly sustained into the post-running time period.

Dorsal hippocampal 5-HT levels, on the other hand, were not affected by running (two-

way RM ANOVA, Shaffer's *post hoc* test, group, $F_{(2, 17)} = 0.046$, $p = 0.955$, time, $F_{(2.79,}$

$_{47.46)} = 16.236$, $p < 0.05$, group and time interaction, $F_{(5.58, 47.46)} = 1.16$, $p = 0.343$, Fig.

3G).

3.4 Running stimulated TH$^+$ neurons in LC and VTA, but not raphe 5-HT$^+$ neurons

To examine the responsiveness of brainstem monoaminergic nuclei, we performed

double-immunostaining of c-Fos$^+$/TH$^+$ neurons for the activation of NA- and DA-

producing neurons, and c-Fos$^+$/5-HT$^+$ neurons for the activation of serotonergic neurons,

as shown in Figure 4A-C. TH$^+$ neurons in the LC (LCTH) and VTA (VTATH), which are

known as the main sources of dorsal hippocampal NA and DA, respectively, were

significantly stimulated by both light- and vigorous-intensity running (LC: one-way

ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)} = 20.908$, $p < 0.001$, Fig. 4D; VTA: one-way

ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)} = 4.992$, $p < 0.05$, Fig. 4E). However, running

did not have any impact on dopaminergic neurons in the SNpc (SNpc[TH] neurons), or on

serotonergic neurons in the DRN and MRN (DRN[5-HT] neurons and MRN[5-HT] neurons,

respectively) (SNpc: Kruskal-Wallis test, Mann-Whitney U test adjusted by Holm's *post-*

hoc test, $X^2 = 4.653$, df = 2, $p = 0.097$, Fig. 4F; DRN: one-way ANOVA, Shaffer's *post*

hoc test, $F_{(2, 18)} = 0.832$, $p = 0.454$, Fig. 4G; MRN: one-way ANOVA, Shaffer's *post hoc*

test, $F_{(2, 18)} = 1.640$, $p = 0.222$, Fig. 4H). Note that TH[+] neurons in the retrorubral field

(RRF) is also reported to have projections to the hippocampus,[81] which were not activated

by running (Fig. S2B). Additionally, TH[+] neurons were also observed in the rostral region

of the DRN (DRN[TH] neurons), which are reported to contribute to wakefulness by

salience stimuli, opioid addiction, and pain modulation through the release of DA.[82] In

this study, however, running did not affect the c-Fos expression in DRN[TH] neurons (Fig.

S2D).

3.5   Neuronal   associations   between   dorsal   hippocampus   and   brainstem

monoaminergic nuclei

To explore the potential association between light-exercise-induced neuronal activities in

brainstem monoaminergic nuclei and dorsal hippocampal activation, we conducted

Pearson's correlation analysis using the dataset from the control group and the light-

intensity running group. This analysis assessed the correlation between c-Fos expression

in the brainstem monoaminergic nuclei and the dorsal hippocampus. A correlation matrix

heatmap that summarizes the correlation results is shown in Figure 5. c-Fos expression in LCTH neurons exhibited a significant and positive correlation within all examined dorsal hippocampal sub-regions except CA2 ($n = 14$, DG: $r = 0.607$, $p < 0.05$, Fig 5B; CA1: $r = 0.691$, $p < 0.01$, Fig 5C; CA2: $r = 0.132$, $p = 0.652$; CA3: $r = 0.643$, $p < 0.05$, Fig 5D). Additionally, a significant positive correlation was found between c-Fos expression in VTATH neurons and in the dorsal CA1 (VTA: $n = 14$, DG: $r = 0.482$, $p = 0.0816$; CA1: $r = 0.570$, $p < 0.05$, Fig 5E; CA2: $r = 0.004$, $p = 0.989$; CA3: $r = 0.391$, $p = 0.167$). The higher level of c-Fos expression in DRN^{5-HT} neurons was significantly correlated with lower levels of CA2 and CA3 c-Fos expression ($n = 14$, DG: $r = -0.532$, $p = 0.0502$; CA1: $r = -0.506$, $p = 0.0649$; CA2: $r = -0.675$, $p < 0.01$, Fig 5F; CA3: $r = -0.704$, $p < 0.01$, Fig 5G). However, no correlation was found in the c-Fos expression between SNpcTH or MRN^{5-HT} neurons and dorsal hippocampal sub-regions (SNpc: $n = 14$, DG: $r = -0.0362$, $p = 0.902$; CA1: $r = 0.196$, $p = 0.502$; CA2: $r = -0.0477$, $p = 0.871$; CA3: $r = 0.0374$, $p = 0.899$; MRN: $n = 14$, DG: $r = -0.252$, $p = 0.385$; CA1: $r = -0.207$, $p = 0.478$; CA2: $r = 0.0693$, $p = 0.814$; CA3: $r = -0.402$, $p = 0.155$).

4. Discussion

To probe the neural circuitry responsible for activating the dorsal hippocampus during light exercise and its subsequent contribution to enhanced memory functions, this study examined the involvement of the ascending monoaminergic system—a critical neuromodulatory mechanism for memory—in the activation of the dorsal hippocampus induced by light exercise. Employing a rat exercise model based on the LT, we assessed neuronal activity in the dorsal hippocampus and brainstem monoaminergic neurons and observed real-time dynamics of monoamine release in the dorsal hippocampus using *in*

vivo microdialysis. The results demonstrate that even light exercise led to increased c-Fos

expression in specific subregions of the dorsal hippocampus and triggered the release of

both NA and DA within this brain region. Furthermore, the activity of LCTH neurons and

VTATH neurons elevated during exercise even at the light-intensity, and both were

positively correlated with the activation of the dorsal hippocampus. In contrast, the

serotonergic system did not appear to be influenced by exercise. These findings imply

LC-noradrenergic and VTA-dopaminergic contributions to dorsal hippocampal activation

during light exercise (Fig. 6), offering insights into the circuitry behind light-exercise-

enhanced learning and memory.

Here, we first established a varied-intensity treadmill running model for male adult

Long-Evans rats based on LT as a physiological index of running intensity and stress. The

speed at the LT was similar to that in our previous work using male adult Wistar rats

(Long-Evans: 21.87 m/min ± 1.02 vs Wistar: 20.7 m/min ± 2.1; Fig. 1B, C).[23] In addition,

a significant increase in blood lactate concentration was observed following a single bout

of vigorous-intensity running, but not with light-intensity running (Fig. 1E), which is

consistent with our previous reports.[23,25] These results confirm the validity of our

experimental model (LT-based varied-intensity running) and demonstrate that the two

running groups exhibited different metabolic and stress-related responses.

Previous studies have reported that acute varied-intensity treadmill running activates

the dorsal hippocampus, including the DG, CA1, and CA3 sub-regions, even at low-

speeds.[6,7,9] Our current results support these findings since both light- and vigorous-

intensity running increased c-Fos expression in the dorsal DG, CA1, and CA3 sub-regions

(Fig. 2C, D, F). In contrast, the dorsal CA2 region was not activated with running (Fig.

2E). The CA2 is known as a small hippocampal sub-region that modulates social memory

and that is stimulated by novel encounters.[83] Since all rats were well familiarized with the

experimental conditions and there were no novel or social stimuli, it is reasonable that

activity in the CA2 region was not affected. Additionally, we examined the activation of

the ventral hippocampus following running and found an increase of c-Fos expression

across all sub-regions of the ventral hippocampus in both running groups (Fig. S1). These

results suggest that light-intensity running is sufficient to elevate the neuronal activity in

the hippocampus across the dorsal-to-ventral axis, which may help to explain the positive

effects of exercise on dorsal-hippocampus-related functions, such as memory, as well as

ventral-hippocampus-related anxiolytic and antidepressant effects.[3,29,84]

Our *in vivo* microdialysis experiment provides the first evidence that even light-

intensity running increases the release of NA and DA at the dorsal hippocampus (Fig. 3E,

F). Given that NA and DA both perform critical roles in dorsal hippocampal neuronal

activity, plasticity, and memory process,[60,61,63,64,85–87] our results suggest that the dorsal

hippocampal NA and DA release contributes to the light-exercise-induced hippocampal

activation and beneficial effects, such as hippocampal neurogenesis and spatial memory

performance.[9,29,30] Notably, NA and DA have been recognized as important factors in

memory consolidation,[63,86] as shown by the synaptic tagging and capture hypothesis,[88,89]

raising a hypothesis that a single-bout of light-intensity running could facilitate memory

retention through noradrenergic and dopaminergic dorsal hippocampal activation.

Intriguingly, prior studies demonstrate that a single session of vigorous-intensity exercise

elevates NA and DA concentrations in the hippocampus and promotes novel object

recognition, which is a dorsal-hippocampus-related function, through the β adrenergic

receptor and dopamine D1 receptor, respectively.[3,79] Although we did not directly

examine this in the present study, these findings support our hypothesis.

We confirmed that treadmill running, even at light intensity, activates LC[TH] and

VTA[TH] neurons (Fig. 4D, E). Furthermore, our correlation analysis observed that LC[TH]

activation positively correlates with dorsal hippocampal activation, and the activities of

VTA[TH] were positively associated with those of the dorsal CA1 region (Fig. 5). These

results suggest that running stimulates the dorsal hippocampal neurons through

noradrenergic projection from the LC and dopaminergic projection from the VTA. While

our study didn't investigate the projections from light-exercise-activated TH[+] neurons in

the LC and the VTA to the dorsal hippocampus, interestingly, a recent study with *in vivo*

two-photon calcium imaging provides robust support for our hypothesis, which

demonstrates an elevation in the activities of both the LC and the VTA axons within the

dorsal CA1 in head-fixed mice during voluntary running.[90] Both nuclei are known as

major sources of dorsal hippocampal NA and DA,[64,65,91] but recent studies have shown

that LC terminals co-release DA and NA to the dorsal hippocampus, and that LC-DA, but

not LC-NA, plays a significant role in spatial memory and learning of novel context.[63–

66,85,92] On the other hand, VTA[TH] neurons have sparse input to the dorsal hippocampus

and do not have a significant impact on spatial memory.[63,64] However, hippocampal DA

from VTA[TH] neurons is still reported to be involved in hippocampus-related functions,

such as spatial memory and aversive memory.[70,71,93] The significance of LC[TH] and VTA[TH]

neurons for the dorsal hippocampal functions remains elusive; thus, future studies using

viral-tracing techniques and chemogenetic/optogenetic manipulation are needed to

investigate the noradrenergic and dopaminergic mechanisms underlying the beneficial

effects of light exercise.

Unlike the noradrenergic and the dopaminergic systems, 5-HT release was not

increased at the dorsal hippocampus, nor were DRN[5-HT] or MRN[5-HT] neurons found to be

stimulated by treadmill running (Fig. 3G; Fig. 4G, H). However, we found a negative

correlation between c-Fos expression in the DG, CA2, and CA3 sub-regions and that in

DRN[5-HT] neurons (Fig. 5F, G). Previous studies have shown that running increases

hippocampal neuronal activity and also facilitates hippocampal LTP induction.[94–96]

Conversely, the hippocampal 5-HT can exert inhibitory effects on basal neural activity

and LTP induction.[97–99] Considering these findings, our results imply that running

enhances hippocampal neuronal activity and LTP induction through the inhibitory control

of the serotonergic inhibitory projection from DRN[5-HT] neurons to the dorsal

hippocampus.

From a clinical perspective, light exercise, such as yoga, tai-chi, and slow running,

is a simple and feasible intervention accessible by most people, including patients with

mild cognitive impairment (MCI) and Alzheimer's disease (AD). Recent studies have

shown that the hippocampal NA and DA loss and degeneration of LC[TH] and VTA[TH]

neurons and axons are found in AD model rodents and AD patients.[100–104] Notably,

selective accumulation of amyloid β and tau proteins are observed in LC neurons before

the onset or at the early stage of AD symptoms.[103–107] Light exercise has been reported to

have a positive effect on MCI and AD.[108–110] Our findings using a translational animal

exercise model, therefore, could provide insight into understanding the mechanisms

underlying the preventive or therapeutic role of light exercise in cognitive dysfunction

conditions, such as AD.

5. Conclusion

Collectively, our findings using a translational exercise model revealed that even light-

intensity running can stimulate the dorsal hippocampal neuronal activity and the release

535   of NA and DA probably from LC and VTA, respectively but that it does not affect the

536   serotonergic system. These findings suggest a possible contribution of LC-noradrenergic

537   and VTA-dopaminergic projections to dorsal hippocampal activation during light exercise

538   (Fig. 6), which may provide insight into the circuit mechanism that underlies light-

539   exercise-enhanced learning and memory.

Author Contributions

Taichi Hiraga, Toshiaki Hata, Shingo Soya, Kanako Takahashi, Masahiro Okamoto, and

Hideaki Soya conceived and designed the experiments; Joshua P. Johansen provided the

experimental animals; Taichi Hiraga and Kanako Takahashi performed the treadmill

running experiments for LT measurement and validation of exercise intensity; Taichi

Hiraga performed the treadmill running experiments for immunohistochemistry and

microdialysis; Toshiaki Hata and Ryo Shimoda performed experimental assistances;

Taichi Hiraga and Ryo Shimoda analyzed the data; Taichi Hiraga wrote the original

manuscript, and all authors were involved in reviewing and editing the manuscript.

Acknowledgments

The authors are grateful to colleagues at the Laboratory of Exercise Biochemistry and

Neuroendocrinology (Faculty of Health and Sport Sciences, University of Tsukuba,

Japan), particularly R. Kuwamizu, H. Koizumi, T. Ferenc, F. Grenier, and M.J. Ikemoto

for insightful scientific discussions. We thank J.P. Johansen (RIKEN CBS) for providing

the experimental animals (TH-Cre Long-Evans rats line) and discussion. We would like

to express our gratitude to M. Noguchi (ELCS English Language Consultation, Japan)

for helping with the manuscript. This research was supported in part by KAKENHI

Grants-in-Aid for Scientific Research (A) (18H04081; 21H04858) (to H.S.); KAKENHI

Grants-in-Aid for Scientific Research on Innovative Areas: Next Generation Exercise

Program for Developing Motivation, Body and Mind Performance (16H06405) (to H.S.);

Japan Science and Technology Agency (JST)-Mirai Program (JPMJMI19D5) (to H.S.);

Grant-in-Aid for Japan Society for the Promotion of Science Fellowships (20J20887) (to

T.H.).

Disclosures

The authors declare no competing financial interests.

References

1.   van Praag H, Christie BR, Sejnowski TJ, Gage FH. Running enhances

neurogenesis, learning, and long-term potentiation in mice. *Proc Natl Acad Sci U*

S A. 1999;96(23):13427-13431. doi:10.1073/pnas.96.23.13427

2.   Erickson KI, Voss MW, Prakash RS, et al. Exercise training increases size of

hippocampus and improves memory. *Proc Natl Acad Sci*. 2011;108(7):3017-3022.

doi:10.1073/pnas.1015950108

3.   Vargas LS, Neves BHS das, Roehrs R, Izquierdo I, Mello-Carpes PB. One-single

physical exercise session after object recognition learning promotes memory

persistence through hippocampal noradrenergic mechanisms. *Behav Brain Res*.

2017;329:120-126. doi:10.1016/j.bbr.2017.04.050

4.   van Dongen E V., Kersten IHP, Wagner IC, Morris RGM, Fernández G. Physical

Exercise Performed Four Hours after Learning Improves Memory Retention and

Increases Hippocampal Pattern Similarity during Retrieval. *Curr Biol*.

2016;26(13):1722-1727. doi:10.1016/J.CUB.2016.04.071

5.   Fanselow MS, Dong HW. Are the Dorsal and Ventral Hippocampus Functionally

Distinct Structures? *Neuron*. 2010;65(1):7-19. doi:10.1016/j.neuron.2009.11.031

6.   Oladehin A, Waters RS. Location and distribution of Fos protein expression in rat

hippocampus following acute moderate aerobic exercise. *Exp Brain Res*.

2001;137(1):26-35. doi:10.1007/s002210000634

7.   Lee TH, Jang MH, Shin MC, et al. Dependence of rat hippocampal c-Fos

expression on intensity and duration of exercise. *Life Sci*. 2003;72(12):1421-1436.

doi:10.1016/S0024-3205(02)02406-2

8.   Nishijima T, Soya H. Evidence of functional hyperemia in the rat hippocampus

during mild treadmill running. *Neurosci Res*. 2006;54(3):186-191.

doi:10.1016/J.NEURES.2005.11.005

9.   Soya H, Nakamura T, Deocaris CC, et al. BDNF induction with mild exercise in

the rat hippocampus. *Biochem Biophys Res Commun*. 2007;358(4):961-967.

doi:10.1016/j.bbrc.2007.04.173

10.   Neeper SA, Góauctemez-Pinilla F, Choi J, Cotman C. Exercise and brain

neurotrophins. *Nature*. 1995;373(6510):109. doi:10.1038/373109a0

11.   Vaynman S, Ying Z, Gomez-Pinilla F. Interplay between brain-derived

neurotrophic factor and signal transduction modulators in the regulation of the

effects of exercise on synaptic-plasticity. *Neuroscience*. 2003;122(3):647-657.

doi:10.1016/J.NEUROSCIENCE.2003.08.001

12.   Nishijima T, Okamoto M, Matsui T, Kita I, Soya H. Hippocampal functional

hyperemia mediated by NMDA receptor/NO signaling in rats during mild exercise.

J Appl Physiol. 2012;112(1):197-203. doi:10.1152/japplphysiol.00763.2011

13.   Trejo JL, Carro E, Torres-Alemán I. Circulating Insulin-Like Growth Factor I
Mediates Exercise-Induced Increases in the Number of New Neurons in the Adult
Hippocampus. *J Neurosci*. 2001;21(5):1628-1634. doi:10.1523/JNEUROSCI.21-
05-01628.2001

14.   Wrann CD, White JP, Salogiannnis J, et al. Exercise Induces Hippocampal BDNF
through a PGC-1α/FNDC5 Pathway. *Cell Metab*. 2013;18(5):649-659.
doi:10.1016/J.CMET.2013.09.008

15.   Islam MR, Valaris S, Young MF, et al. Exercise hormone irisin is a critical
regulator of cognitive function. *Nat Metab*. 2021;3(8):1058. doi:10.1038/S42255-
021-00438-Z

16.   Lourenco M V., Frozza RL, de Freitas GB, et al. Exercise-linked FNDC5/irisin
rescues synaptic plasticity and memory defects in Alzheimer's models. *Nat Med*.
2019;25(1):165. doi:10.1038/S41591-018-0275-4

17.   Nishijima T, Piriz J, Duflot S, et al. Neuronal Activity Drives Localized Blood-
Brain-Barrier Transport of Serum Insulin-like Growth Factor-I into the CNS.
Neuron. 2010;67(5):834-846. doi:10.1016/j.neuron.2010.08.007

18.   Moon HY, Becke A, Berron D, et al. Running-Induced Systemic Cathepsin B
Secretion Is Associated with Memory Function. *Cell Metab*. 2016;24(2):332-340.
doi:10.1016/J.CMET.2016.05.025

19.   Vaynman S, Ying Z, Gomez-Pinilla F. Hippocampal BDNF mediates the efficacy
of exercise on synaptic plasticity and cognition. *Eur J Neurosci*.
2004;20(10):2580-2590. doi:10.1111/J.1460-9568.2004.03720.X

20.   Li Y, Luikart BW, Birnbaum S, et al. TrkB Regulates Hippocampal Neurogenesis
and Governs Sensitivity to Antidepressive Treatment. *Neuron*. 2008;59(3):399-

412. doi:10.1016/J.NEURON.2008.06.023

21.   Lee MC, Okamoto M, Liu YF, et al. Voluntary resistance running with short

distance enhances spatial memory related to hippocampal BDNF signaling. *J Appl*

Physiol. 2012;113(8):1260-1266. doi:10.1152/japplphysiol.00869.2012

22.   Pereira AC, Huddleston DE, Brickman AM, et al. An in vivo correlate of exercise-

induced neurogenesis in the adult dentate gyrus. *Proc Natl Acad Sci U S A*.

2007;104(13):5638-5643.

doi:10.1073/PNAS.0611721104/SUPPL_FILE/11721FIG5.PDF

23.   Soya H, Mukai A, Deocaris CC, et al. Threshold-like pattern of neuronal activation

in the hypothalamus during treadmill running: Establishment of a minimum

running stress (MRS) rat model. *Neurosci Res*. 2007;58(4):341-348.

doi:10.1016/j.neures.2007.04.004

24.   Ekkekakis P, Hall EE, Petruzzello SJ. The Relationship Between Exercise

Intensity and Affective Responses Demystified: To Crack the 40-Year-Old Nut,

Replace the 40-Year-Old Nutcracker! *Ann Behav Med*. 2008;35(2):136-149.

doi:10.1007/S12160-008-9025-Z

25.   Saito T, Soya H. Delineation of responsive AVP-containing neurons to running

stress in the hypothalamus. *Am J Physiol Integr Comp Physiol*. 2004;286(3):R484-

R490. doi:10.1152/ajpregu.00453.2003

26.   Hill EE, Zack E, Battaglini C, Viru M, Viru A, Hackney AC. Exercise and

circulating cortisol levels: The intensity threshold effect. *J Endocrinol Invest*.

2008;31(7):587-591. doi:10.1007/BF03345606/METRICS

27.   Perri MG, Anton SD, Durning PE, et al. Adherence to exercise prescriptions:

Effects of prescribing moderate versus higher levels of intensity and frequency.

Heal Psychol. 2002;21(5):452-458. doi:10.1037/0278-6133.21.5.452

28. Rodnick KJ, Reaven GM, Haskell WL, Sims CR, Mondon CE. Variations in

running activity and enzymatic adaptations in voluntary running rats.

https://doi.org/101152/jappl19896631250.   1989;66(3):1250-1257.

doi:10.1152/JAPPL.1989.66.3.1250

29. Inoue K, Hanaoka Y, Nishijima T, et al. Long-term mild exercise training enhances

hippocampus-dependent memory in rats. *Int J Sports Med*. 2015;36(4):280-285.

doi:10.1055/S-0034-1390465/ID/R4165-0031

30. Inoue K, Okamoto M, Shibato J, et al. Long-Term Mild, rather than Intense,

Exercise Enhances Adult Hippocampal Neurogenesis and Greatly Changes the

Transcriptomic Profile of the Hippocampus. *PLoS One*. 2015;10(6):e0128720.

doi:10.1371/JOURNAL.PONE.0128720

31. Okamoto M, Hojo Y, Inoue K, et al. Mild exercise increases dihydrotestosterone

in hippocampus providing evidence for androgenic mediation of neurogenesis.

Proc   Natl   Acad   Sci   U   S   A.   2012;109(32):13100-13105.

doi:10.1073/pnas.1210023109

32. Conrad CD. A critical review of chronic stress effects on spatial learning and

memory. *Prog Neuro-Psychopharmacology Biol Psychiatry*. 2010;34(5):742-755.

doi:10.1016/J.PNPBP.2009.11.003

33. Schoenfeld TJ, Gould E. Stress, Stress Hormones, and Adult Neurogenesis. *Exp*

Neurol. 2012;233(1):12. doi:10.1016/J.EXPNEUROL.2011.01.008

34. Suwabe K, Byun K, Hyodo K, et al. Rapid stimulation of human dentate gyrus

function with acute mild exercise. *Proc Natl Acad Sci*. 2018;115(41):10487-10492.

doi:10.1073/pnas.1805668115

35.   McMorris T. Developing the catecholamines hypothesis for the acute exercise-

cognition interaction in humans: Lessons from animal studies. *Physiol Behav*.

2016;165:291-299. doi:10.1016/J.PHYSBEH.2016.08.011

36.   Cooper CJ. Anatomical and physiological mechanisms of arousal, with special

reference to the effects of exercise. *Ergonomics*. 1973;16(5):601-609.

doi:10.1080/00140137308924551

37.   Dietrich A, Audiffren M. The reticular-activating hypofrontality (RAH) model of

acute exercise. *Neurosci Biobehav Rev*. 2011;35(6):1305-1325.

doi:10.1016/J.NEUBIOREV.2011.02.001

38.   Saper CB, Chou TC, Scammell TE. Review. *TRENDS Neurosci*. 2001;24(12).

Accessed June 8, 2023. http://tins.trends.com0166

39.   Jones BE. Arousal systems. *Front Biosci*. 2003;8(SUPPL.):438-451.

doi:10.2741/1074/PDF

40.   Kuwamizu R, Yamazaki Y, Aoike N, Ochi G, Suwabe K, Soya H. Pupil-linked

arousal with very light exercise: pattern of pupil dilation during graded exercise. *J*

Physiol Sci. 2022;72(1):1-9. doi:10.1186/S12576-022-00849-X/FIGURES/5

41.   Yamazaki Y, Suwabe K, Nagano-Saito A, et al. A possible contribution of the

locus coeruleus to arousal enhancement with mild exercise: evidence from

pupillometry and neuromelanin imaging. *Cereb Cortex Commun*. 2023;4(2):1-8.

doi:10.1093/TEXCOM/TGAD010

42.   Kuwamizu R, Yamazaki Y, Aoike N, et al. Pupil dynamics during very light

exercise predict benefits to prefrontal cognition. *Neuroimage*. Published online

June 21, 2023:120244. doi:10.1016/J.NEUROIMAGE.2023.120244

43.   Suwabe K, Kuwamizu R, Hyodo K, et al. Improvement of mnemonic

discrimination with acute light exercise is mediated by pupil-linked arousal in

healthy older adults. *Neurobiol Aging*. 2024;133:107-114.

doi:10.1016/J.NEUROBIOLAGING.2023.09.006

44.   Reimer J, McGinley MJ, Liu Y, et al. Pupil fluctuations track rapid changes in

adrenergic and cholinergic activity in cortex. *Nat Commun*. 2016;7(1):1-7.

doi:10.1038/ncomms13289

45.   Cazettes F, Reato D, Morais JP, Renart A, Mainen ZF. Phasic Activation of Dorsal

Raphe Serotonergic Neurons Increases Pupil Size. *Curr Biol*. 2021;31(1):192-

197.e4. doi:10.1016/J.CUB.2020.09.090

46.   Zerbi V, Floriou-Servou A, Markicevic M, et al. Rapid Reconfiguration of the

Functional Connectome after Chemogenetic Locus Coeruleus Activation. *Neuron*.

2019;103(4):702-718.e5. doi:10.1016/J.NEURON.2019.05.034

47.   Balderas I, Moreno-Castilla P, Bermudez-Rattoni F. Dopamine D1 receptor

activity modulates object recognition memory consolidation in the perirhinal

cortex but not in the hippocampus. *Hippocampus*. 2013;23(10):873-878.

doi:10.1002/hipo.22143

48.   de Gee JW, Colizoli O, Kloosterman NA, Knapen T, Nieuwenhuis S, Donner TH.

Dynamic modulation of decision biases by brainstem arousal systems. *Elife*.

2017;6. doi:10.7554/ELIFE.23232

49.   Carter ME, Yizhar O, Chikahisa S, et al. Tuning arousal with optogenetic

modulation of locus coeruleus neurons. *Nat Neurosci*. 2010;13(12):1526-1535.

doi:10.1038/nn.2682

50.   Bari A, Xu S, Pignatelli M, et al. Differential attentional control mechanisms by

two distinct noradrenergic coeruleo-frontal cortical pathways. *Proc Natl Acad Sci*

U S A. 2020;117(46):29080-29089. doi:10.1073/pnas.2015635117

51.   Saunders BT, Richard JM, Margolis EB, Janak PH. Dopamine neurons create

729   Pavlovian conditioned stimuli with circuit-defined motivational properties. *Nat*

730   *Neurosci 2018 218*. 2018;21(8):1072-1083. doi:10.1038/s41593-018-0191-4

731   52.   Zhu X, Ottenheimer D, DiLeone RJ. Activity of D1/2 Receptor Expressing

732   Neurons in the Nucleus Accumbens Regulates Running, Locomotion, and Food

733   Intake.   *Front   Behav   Neurosci*.   2016;10(April):1-10.

734   doi:10.3389/fnbeh.2016.00066

735   53.   Kim SY, Adhikari A, Lee SY, et al. Diverging neural pathways assemble a

736   behavioural state from separable features in anxiety. *Nature*. 2013;496(7444):219-

737   223. doi:10.1038/nature12018

738   54.   Warden MR, Selimbeyoglu A, Mirzabekov JJ, et al. A prefrontal cortex–brainstem

739   neuronal projection that controls response to behavioural challenge. *Nat 2012*

740   *4927429*. 2012;492(7429):428-432. doi:10.1038/nature11617

741   55.   Teissier A, Chemiakine A, Inbar B, et al. Activity of Raphé Serotonergic Neurons

742   Controls   Emotional   Behaviors.   *Cell   Rep*.   2015;13(9):1965-1976.

743   doi:10.1016/J.CELREP.2015.10.061

744   56.   Palacios-Filardo J, Mellor JR. Neuromodulation of hippocampal long-term

745   synaptic   plasticity.   *Curr   Opin   Neurobiol*.   2019;54:37-43.

746   doi:10.1016/J.CONB.2018.08.009

747   57.   Poe GR, Foote S, Eschenko O, et al. Locus coeruleus: a new look at the blue spot.

748   *Nat Rev Neurosci 2020 2111*. 2020;21(11):644-659. doi:10.1038/s41583-020-

749   0360-9

58.   Wang SH, Redondo RL, Morris RGM. Relevance of synaptic tagging and capture

to the persistence of long-term potentiation and everyday spatial memory. *Proc*

Natl   Acad   Sci   U   S   A.   2010;107(45):19537-19542.

doi:10.1073/PNAS.1008638107/SUPPL_FILE/PNAS.201008638SI.PDF

59. Mahar I, Bambico FR, Mechawar N, Nobrega JN. Stress, serotonin, and

hippocampal neurogenesis in relation to depression and antidepressant effects.

Neurosci   Biobehav   Rev.   2014;38:173-192.

doi:10.1016/J.NEUBIOREV.2013.11.009

60. Coradazzi M, Gulino R, Fieramosca F, Falzacappa LV, Riggi M, Leanza G.

Selective noradrenaline depletion impairs working memory and hippocampal

neurogenesis.   *Neurobiol   Aging.*   2016;48:93-102.

doi:10.1016/J.NEUROBIOLAGING.2016.08.012

61. Salvi R, Steigleder T, Schlachetzki JCM, et al. Distinct effects of chronic

dopaminergic stimulation on hippocampal neurogenesis and striatal doublecortin

expression   in   adult   mice.   *Front   Neurosci.*   2016;10(MAR):77.

doi:10.3389/FNINS.2016.00077/BIBTEX

62. Schwarz LA, Miyamichi K, Gao XJ, et al. Viral-genetic tracing of the input-output

organization of a central noradrenaline circuit. *Nature.* 2015;524(7563):88-92.

doi:10.1038/nature14600

63. Takeuchi T, Duszkiewicz AJ, Sonneborn A, et al. Locus coeruleus and

dopaminergic consolidation of everyday memory. *Nature.* 2016;537(7620):357-

362. doi:10.1038/nature19325

64. Gálvez-Márquez DK, Salgado-Ménez M, Moreno-Castilla P, et al. Spatial

contextual recognition memory updating is modulated by dopamine release in the

dorsal hippocampus from the locus coeruleus. *Proc Natl Acad Sci U S A.*

2022;119(49):e2208254119.

doi:10.1073/PNAS.2208254119/SUPPL_FILE/PNAS.2208254119.SAPP.PDF

65.   Kempadoo KA, Mosharov E V., Choi SJ, Kandel ER, Sulzer D. Dopamine release

778   from the locus coeruleus to the dorsal hippocampus promotes spatial learning and

779   memory.   *Proc   Natl   Acad   Sci*.   2016;113(51):14835-14840.

780   doi:10.1073/pnas.1616515114

781   66.   Wagatsuma A, Okuyama T, Sun C, Smith LM, Abe K, Tonegawa S. Locus

782   coeruleus input to hippocampal CA3 drives single-trial learning of a novel context.

783   *Proc Natl Acad Sci U S A*. 2017;115(2):E310-E316. doi:10.1073/pnas.1714082115

784   67.   Gasbarri A, Packard MG, Campana E, Pacitti C. Anterograde and retrograde

785   tracing of projections from the ventral tegmental area to the hippocampal

786   formation in the rat. *Brain Res Bull*. 1994;33(4):445-452. doi:10.1016/0361-

787   9230(94)90288-7

788   68.   Gasbarri A, Verney C, Innocenzi R, Campana E, Pacitti C. Mesolimbic

789   dopaminergic neurons innervating the hippocampal formation in the rat: a

790   combined retrograde tracing and immunohistochemical study. *Brain Res*.

791   1994;668(1-2):71-79. doi:10.1016/0006-8993(94)90512-6

792   69.   Swanson LW. The projections of the ventral tegmental area and adjacent regions-

793   a combined fluorescent retrograde tracer and immunofluorescence study in the rat.

794   1982;9:321-353.

795   70.   Tsetsenis T, Badyna JK, Wilson JA, et al. Midbrain dopaminergic innervation of

796   the hippocampus is sufficient to modulate formation of aversive memories. *Proc*

797   *Natl   Acad   Sci   U   S   A*.   2021;118(40).

798   doi:10.1073/PNAS.2111069118/SUPPL_FILE/PNAS.2111069118.SM01.MP4

71. McNamara CG, Tejero-Cantero Á, Trouche S, Campo-Urriza N, Dupret D.

Dopaminergic neurons promote hippocampal reactivation and spatial memory

persistence. *Nat Neurosci*. 2014;17(12):1658-1660. doi:10.1038/nn.3843

72. Muzerelle A, Scotto-Lomassese S, Bernard JF, Soiza-Reilly M, Gaspar P.

Conditional anterograde tracing reveals distinct targeting of individual serotonin

cell groups (B5–B9) to the forebrain and brainstem. *Brain Struct Funct*.

2016;221(1):535-561. doi:10.1007/S00429-014-0924-4/FIGURES/13

73. Luchetti A, Bota A, Weitemier A, et al. Two functionally distinct serotonergic

projections into hippocampus. *J Neurosci*. 2020;40(25):JN-RM-2724-19.

doi:10.1523/jneurosci.2724-19.2020

74. Morikawa R, Kubota N, Amemiya S, Nishijima T, Kita I. Interaction between

intensity and duration of acute exercise on neuronal activity associated with

depression-related behavior in rats. *J Physiol Sci*. 2021;71(1):1.

doi:10.1186/s12576-020-00788-5

75. Tsai SF, Liu YW, Kuo YM. Acute and long-term treadmill running differentially

induce c-Fos expression in region- and time-dependent manners in mouse brain.

Brain Struct Funct. Published online July 27, 2019:1-13. doi:10.1007/s00429-019-

01926-5

76. Rodovalho GV, Drummond LR, Coimbra CC. Involvement of brainstem

noradrenergic system in cutaneous heat loss during exercise. *Brain Res Bull*.

2020;164:372-379. doi:10.1016/J.BRAINRESBULL.2020.08.029

77. Goekint M, Bos I, Heyman E, Meeusen R, Michotte Y, Sarre S. Acute running

stimulates hippocampal dopaminergic neurotransmission in rats, but has no

influence on brain-derived neurotrophic factor. *J Appl Physiol*. 2012;112(4):535-

541. doi:10.1152/japplphysiol.00306.2011

78.   Gomez-Merino D, Béquet F, Berthelot M, Chennaoui M, Guezennec CY. Site-

dependent effects of an acute intensive exercise on extracellular 5-HT and 5-HIAA

levels in rat brain. *Neurosci Lett*. 2001;301(2):143-146. doi:10.1016/S0304-

3940(01)01626-3

79.   Vargas LS, Ramires Lima K, Piaia Ramborger B, Roehrs R, Izquierdo I, Mello-

Carpes PB. Catecholaminergic hippocampal activation is necessary for object

recognition memory persistence induced by one-single physical exercise session.

Behav Brain Res. 2020;379:112356. doi:10.1016/j.bbr.2019.112356

80.   Beaver WL, Wasserman K, Whipp BJ. Improved detection of lactate threshold

during exercise using a log-log transformation. *J Appl Physiol*. 1985;59(6):1936-

1940. doi:10.1152/jappl.1985.59.6.1936

81.   Gasbarri A, Packard M, Sulli A, Pacitti C, Innocenzi R, Perciavalle V. The

projections of the retrorubral field A8 to the hippocampal formation in the rat. *Exp*

Brain Res. 1996;112(2):244-252. doi:10.1007/BF00227643

82.   Lin R, Liang J, Luo M. The Raphe Dopamine System: Roles in Salience Encoding,

Memory Expression, and Addiction. *Trends Neurosci*. 2021;44(5):366-377.

doi:10.1016/J.TINS.2021.01.002

83.   Chen S, He L, Huang AJY, et al. A hypothalamic novelty signal modulates

hippocampal memory. *Nature*. Published online September 30, 2020:1-5.

doi:10.1038/s41586-020-2771-1

84.   Otsuka T, Nishii A, Amemiya S, Kubota N, Nishijima T, Kita I. Effects of acute

treadmill running at different intensities on activities of serotonin and

corticotropin-releasing factor neurons, and anxiety- and depressive-like behaviors

in rats. *Behav Brain Res*. 2016;298:44-51. doi:10.1016/j.bbr.2015.10.055

85.   Tse D, Privitera L, Norton AC, et al. Cell-type-specific optogenetic stimulation of

the locus coeruleus induces slow-onset potentiation and enhances everyday

memory in rats. *Proc Natl Acad Sci U S A*. 2023;120(46):e2307275120.

doi:10.1073/PNAS.2307275120/SUPPL_FILE/PNAS.2307275120.SD01.XLSX

86.   Mello-Carpes PB, da Silva de Vargas L, Gayer MC, Roehrs R, Izquierdo I.

Hippocampal noradrenergic activation is necessary for object recognition memory

consolidation and can promote BDNF increase and memory persistence. *Neurobiol*

Learn Mem. 2016;127:84-92. doi:10.1016/J.NLM.2015.11.014

87.   Takamura N, Nakagawa S, Masuda T, et al. The effect of dopamine on adult

hippocampal neurogenesis. *Prog Neuro-Psychopharmacology Biol Psychiatry*.

2014;50:116-124. doi:10.1016/J.PNPBP.2013.12.011

88.   Frey U, Morris RGM. Synaptic tagging and long-term potentiation. *Nat 1997*

3856616. 1997;385(6616):533-536. doi:10.1038/385533a0

89.   Redondo RL, Morris RGM. Making memories last: the synaptic tagging and

capture hypothesis. *Nat Rev Neurosci 2011 121*. 2010;12(1):17-30.

doi:10.1038/nrn2963

90.   Heer CM, Sheffield MEJ. Distinct catecholaminergic pathways projecting to

hippocampal CA1 transmit contrasting signals during behavior and learning.

bioRxiv. Published online January 8, 2023:2023.11.29.569214.

doi:10.1101/2023.11.29.569214

91.   Weitemier AZ, McHugh TJ. Noradrenergic modulation of evoked dopamine

release and pH shift in the mouse dorsal hippocampus and ventral striatum. *Brain*

Res. 2017;1657:74-86. doi:10.1016/J.BRAINRES.2016.12.002

92.   Chowdhury A, Luchetti A, Fernandes G, et al. A locus coeruleus-dorsal CA1

dopaminergic circuit modulates memory linking. *Neuron*. 2022;110(20):3374-

3388.e8. doi:10.1016/J.NEURON.2022.08.001

93.   Kramar CP, Castillo-Díaz F, Gigante ED, Medina JH, Barbano MF. The late

consolidation of an aversive memory is promoted by VTA dopamine release in the

dorsal   hippocampus.   *Eur   J   Neurosci*.   2021;53(3):841-851.

doi:10.1111/EJN.15076

94.   Miladi-Gorji H, Rashidy-Pour A, Fathollahi Y, Semnanian S, Jadidi M. Effects of

voluntary exercise on hippocampal long-term potentiation in morphine-dependent

rats.   *Neuroscience*.   2014;256:83-90.

doi:10.1016/J.NEUROSCIENCE.2013.09.056

95.   Radahmadi M, Hosseini N, Alaei H. Effect of exercise, exercise withdrawal, and

continued regular exercise on excitability and long-term potentiation in the dentate

gyrus   of   hippocampus.   *Brain   Res*.   2016;1653:8-13.

doi:10.1016/j.brainres.2016.09.045

96.   Farmer J, Zhao X, Van Praag H, Wodtke K, Gage FH, Christie BR. Effects of

voluntary exercise on synaptic plasticity and gene expression in the dentate gyrus

of adult male sprague-dawley rats in vivo. *Neuroscience*. 2004;124(1):71-79.

doi:10.1016/j.neuroscience.2003.09.029

97.   Villani F, Johnston D. Serotonin inhibits induction of long-term potentiation at

commissural synapses in hippocampus. *Brain Res*. 1993;606(2):304-308.

doi:10.1016/0006-8993(93)90998-3

98.   Shakesby AC, Anwyl R, Rowan MJ. Overcoming the Effects of Stress on Synaptic

Plasticity in the Intact Hippocampus: Rapid Actions of Serotonergic and

Antidepressant Agents. *J Neurosci*. 2002;22(9):3638.

doi:10.1523/JNEUROSCI.22-09-03638.2002

99. Kulla A, Manahan-Vaughan D. Modulation by serotonin 5-HT4 receptors of long-

term potentiation and depotentiation in the dentate gyrus of freely moving rats.

Cereb Cortex. 2002;12(2):150-162. doi:10.1093/cercor/12.2.150

100. Nobili A, Latagliata EC, Viscomi MT, et al. Dopamine neuronal loss contributes

to memory and reward dysfunction in a model of Alzheimer's disease. *Nat*

Commun 2017 81. 2017;8(1):1-14. doi:10.1038/ncomms14727

101. Sala A, Caminiti SP, Presotto L, et al. In vivo human molecular neuroimaging of

dopaminergic vulnerability along the Alzheimer's disease phases. *Alzheimer's Res*

Ther. 2021;13(1):1-12. doi:10.1186/S13195-021-00925-1/FIGURES/5

102. Francis BM, Yang J, Hajderi E, et al. Reduced tissue levels of noradrenaline are

associated with behavioral phenotypes of the TgCRND8 mouse model of

Alzheimer's disease. *Neuropsychopharmacology*. 2012;37(8):1934-1944.

doi:10.1038/npp.2012.40

103. Rorabaugh JM, Chalermpalanupap T, Botz-Zapp CA, et al. Chemogenetic locus

coeruleus activation restores reversal learning in a rat model of Alzheimer's

disease. *Brain*. 2017;140(11):3023-3038. doi:10.1093/brain/awx232

104. Mehla J, Lacoursiere SG, Lapointe V, et al. Age-dependent behavioral and

biochemical characterization of single APP knock-in mouse (APPNL-G-F/NL-G-

F) model of Alzheimer's disease. *Neurobiol Aging*. 2019;75:25-37.

doi:10.1016/j.neurobiolaging.2018.10.026

105. Matchett BJ, Grinberg LT, Theofilas P, Murray ME. The mechanistic link between

selective vulnerability of the locus coeruleus and neurodegeneration in

919 Alzheimer's disease. *Acta Neuropathol 2021 1415*. 2021;141(5):631-650.

920 doi:10.1007/S00401-020-02248-1

921 106. Betts MJ, Kirilina E, Otaduy MCG, et al. Locus coeruleus imaging as a biomarker

922 for noradrenergic dysfunction in neurodegenerative diseases. *Brain*.

923 2019;142(9):2558-2571. doi:10.1093/BRAIN/AWZ193

924 107. Braak H, Thal DR, Ghebremedhin E, Del Tredici K. Stages of the Pathologic

925 Process in Alzheimer Disease: Age Categories From 1 to 100 Years. *J Neuropathol*

926 *Exp Neurol*. 2011;70(11):960-969. doi:10.1097/NEN.0B013E318232A379

927 108. Yoon M, Yang PS, Jin MN, et al. Association of Physical Activity Level With Risk

928 of Dementia in a Nationwide Cohort in Korea. *JAMA Netw Open*.

929 2021;4(12):e2138526-e2138526.

930 doi:10.1001/JAMANETWORKOPEN.2021.38526

931 109. Meng Q, Lin MS, Tzeng IS. Relationship Between Exercise and Alzheimer's

932 Disease: A Narrative Literature Review. *Front Neurosci*. 2020;14.

933 doi:10.3389/FNINS.2020.00131/FULL

934 110. Vasconcelos-Filho FSL, da Rocha Oliveira LC, de Freitas TBC, et al.

935 Neuroprotective mechanisms of chronic physical exercise via reduction of β-

936 amyloid protein in experimental models of Alzheimer's disease: A systematic

937 review. *Life Sci*. 2021;275:119372. doi:10.1016/J.LFS.2021.119372

Figure 1

Measurement of lactate threshold and validation of exercise intensity.

(A) Experimental design for incremental running test. (B) A representative LT profile

from the incremental running test. (C) Speed of rats at LT was 21.87 ± 1.02 m/min ($n =$

6). (D) Experimental design for a single bout of the running test. We defined 15 m/min

for light-intensity running and 25 m/min for vigorous-intensity running. (E) Blood

lactate concentration was significantly increased only after a 30-minute session of

vigorous-intensity running. All data are expressed as mean ± SEM. ***$p < 0.001$ vs

Control, ###$p < 0.001$ vs Light, $n =$ 5-7/group. Control: sedentary control; Light: 15

m/min running; Vigorous: 25 m/min running; hab.: habituation.

Figure 2

Running-induced neuronal activation at the dorsal hippocampus during varied-

intensity treadmill running.

(A) Experimental design for a single bout of the running test for immunohistochemical

analysis. (B) Representative double-fluorescence images of c-Fos (green) and NeuN

(blue) in the DG, CA1, and CA3 regions of the dorsal hippocampus. Scale bars show

100 μm. c-Fos expressions in the DG (C), CA1 (D), CA2 (E), and CA3(F) were

significantly increased with running exercise regardless of running speed (light or

vigorous). All data are expressed as mean ± SEM. $^{**}p < 0.01$, $^{***}p < 0.001$ vs Control, n

= 6-8 / group. Control: sedentary control; Light: 15 m/min running; Vigorous: 25 m/min

running; hab.: habituation.

Figure 3

Running increases dorsal hippocampal NA and DA release, but not 5-HT release.

(A) Experimental design for a single bout of the running test for dorsal hippocampal

monoamine detection with *in vivo* microdialysis. (B) Representative DAPI fluorescence

image of microdialysis probe position in the dorsal hippocampus. Scale bars show 500

μm. (C) Schematic representations indicating the location of the microdialysis probe.

Each bar represents the probe position of an individual rat, and the color of the bar

corresponds to the experimental group (Control: light gray, Light: blue, and Vigorous:

dark gray). (D) Representative chromatograms showing NA (peak 1), DA (peak 2), and

5-HT (peak 3) from Control (light gray), Light (blue), and Vigorous (dark gray).

Treadmill running induced a significant increase of both NA (E) and DA (F) levels in

983   the dorsal hippocampus, depending on running speed, but not 5-HT (G). All data are

984   expressed as mean $\pm$ SEM. $^{*}p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$ vs Sed, $^{\#}p < 0.05$, $^{\#\#}p <$

985   0.01 vs Light, $n = 6\text{-}7$/group. Control: sedentary control; Light: 15 m/min running;

986   Vigorous: 25 m/min running; hab.: habituation.

1000

1001

1002

1003

1004

1005

1006

1007

Figure 4

Running stimulates LCTH and VTATH neurons, but not raphe 5-HT$^+$ neurons.

Representative double-fluorescence images of c-Fos (green), TH (blue), and 5-HT (red)

neurons in LC (A); VTA and SNpc (B); and DRN and MRN (C). Scale bars show 100

μm. Yellow heads show c-Fos$^+$/TH$^+$ neurons. LCTH and VTATH neurons were

significantly stimulated in both running groups (D, E). However, running did not have

an impact on SNpcTH, DRN^{5-HT}, and MRN^{5-HT} neurons (F, G, H). All data are expressed

as mean ± SEM. $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$ vs Control, $^{##}p < 0.01$ vs Light, $n =$

6-8 / group. Control: sedentary control; Light: 15 m/min running; Vigorous: 25 m/min

running.

1018

Figure 5

Neuronal associations between dorsal hippocampus and brainstem monoaminergic

nuclei.

(A) A correlation matrix heatmap is shown to summarize the correlation analysis

between c-Fos expression in dorsal-hippocampal sub-regions and that in LCTH, VTATH,

SNpcTH, DRN^{5-HT}, and MRN^{5-HT} neurons. Colors in the scale show correlation strength.

c-Fos expression in LCTH neurons exhibited a significant and positive correlation within

the dorsal DG (B), CA1 (C), and CA3 (D). A significant positive correlation was also

found between c-Fos expression in VTATH neurons and in the dorsal CA1 (E). c-Fos

expression in DRN^{5-HT} neurons was significantly and negatively correlated with that in

CA2 (F) and CA3 (G). The colored line represents linear regression, and the colored

band represents 95% confidence bands. $^*p < 0.05$, $^{**}p < 0.01$. $n = 14$.

1031

1032

1033

1034

Figure 6

Schematic diagram showing possible circuit mechanism for light-exercise-induced

dorsal hippocampal activation.

Light exercise activates TH[+] neurons at the LC and the VTA. At the dorsal

hippocampus, subsequently, axon terminals originating from the LC[TH] and the VTA[TH]

increase the release of NA and DA, respectively. These, in turn, upregulate the neuronal

activity of the dorsal hippocampal DG, CA1, and CA3 sub-regions. HPC, hippocampus;

DG, dentate gyrus; NA, noradrenaline; DA, dopamine; LC, locus coeruleus; VTA,

ventral tegmental area.

1044

1045

1046

1047 **Supplemental data**

1048 Light-exercise-induced dopaminergic and noradrenergic stimulation in the dorsal

1049 hippocampus: using a rat physiological exercise model

1050

1051 Taichi Hiraga, Toshiaki Hata, Shingo Soya, Ryo Shimoda, Kanako Takahashi, Mariko Soya, Koshiro

1052 Inoue, Joshua P. Johansen, Masahiro Okamoto, and Hideaki Soya

1053

1054 Corresponding author: Hideaki Soya, Ph.D.

1055 Email: soya.hideaki.gt@u.tsukuba.ac.jp

1056

1057

1058 This PDF file includes:

1059 ・Figure S1 Running induces neuronal activation of the ventral hippocampus

1060 ・Figure S2 RRFTH neurons and DRNTH neurons are not activated by running.

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

Figure S1

Running induces neuronal activation of the ventral hippocampus.

(A) Representative double-fluorescence images of NeuN (blue) and c-Fos (green) in the DG, CA1, and CA3. Scale bars show 100 μm. c-Fos expression in ventral hippocampal sub-regions, including the DG (B), CA1 (C), and CA3 (D), was also significantly increased regardless of running speed (DG: Kruskal-Wallis test, Mann-Whitney U test adjusted by Holm's *post-hoc* test, X^2 = 10.245, df = 2, $p < 0.01$; CA1: one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)}$ = 11.443, $p < 0.001$; CA3: Welch's one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)}$ = 11.758, $p < 0.001$). (E) A correlation matrix heatmap is shown to summarize the correlation results between c-Fos expression in monoaminergic nuclei and that in ventral hippocampal sub-regions. Colors in the scale show correlation strength. Data were analyzed using Pearson's correlation. c-Fos expression in the LC[TH] had a significant positive correlation with that in all ventral hippocampal sub-regions (n = 14, DG: $r = 0.543$, $p < 0.05$; CA1: $r = 0.587$, $p < 0.05$; CA3: $r = 0.654$, $p < 0.05$). Additionally, a significant positive correlation was found between c-Fos expression in the

VTATH neurons and in the ventral DG and CA1 (n = 14, DG: r = 0.550, p < 0.05; CA1: r

= 0.564, p < 0.01; CA3: r = 0.507, p = 0.064). The higher level of c-Fos expression in the

DRN^{5-HT} neurons was significantly correlated with the lower level of ventral CA3 c-Fos

expression (n = 14; DG: r = -0.085, p = 0.715; CA1: r = -0.090, p = 0.697; CA3: r = -

0.522, p < 0.05). No correlation was found between c-Fos expression in SNpcTH or MRN^{5-}

^{HT} neurons and in ventral hippocampal sub-regions (SNpc: n = 14, DG: r = -0.234, p =

0.422; CA1: r = -0.051, p = 0.864; CA3: r = 0.039, p = 0.895; MRN: n = 14, DG: r = -

0.238, p = 0.412; CA1: r = -0.283, p = 0.328; CA3: r = -0.253, p = 0.382). All data are

expressed as mean ± SEM. $^{*}p$ < 0.05, $^{**}p$ < 0.01, $^{***}p$ < 0.001 vs Control; n = 6-8/group.

Control: sedentary control; Light: 15 m/min running; Vigorous: 25 m/min running.

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

Figure S2

RRF^{TH} neurons and DRN^{TH} neurons are not activated by running.

(A) Representative double-fluorescence images of c-Fos (green) and TH (blue) neurons in the RRF. Scale bars show 100 μm. Yellow heads show c-Fos$^+$/TH$^+$ neurons. (B) RRFTH neurons were not activated by running (Kruskal-Wallis test, Mann-Whitney U test adjusted by Holm's *post-hoc* test, $X^2 = 4.494$, df = 2, $p = 0.106$). (C) Representative triple-fluorescence images of c-Fos (green), TH (blue), and 5-HT (red) neurons in the DRN. Scale bars show 50 μm. (D) DRNTH neurons were not activated by running (Welch's one-way ANOVA, Shaffer's *post hoc* test, $F_{(2, 18)} = 0.384$, $p = 0.687$). All data are expressed as mean ± SEM. $n = 6$-8/group. Control: sedentary control; Light: 15 m/min running; Vigorous: 25 m/min running.