SARS-CoV-2 variants of concern exhibit differential gastro-intestinal tropism and pathogenesis in the Syrian golden hamster model.

Authors: Santhosh K. Nagaraj*[1,2], Christy M. Joy*[1,2], Rohan Narayan[1,2], Rishad Shiraz[1,2], Sumandeep Kaur[1,2], Oyahida Khatun[1,2], Sagar Dubey[1,2], Shashank Tripathi[1,2]

Correspondence: shashankt@iisc.ac.in

Affiliation:

[1]Emerging Viral Pathogens Laboratory, Infosys Wing, Centre for Infectious Disease Research, Indian Institute of Science; Bengaluru, 560012, India.

[2]Department of Microbiology and Cell Biology, Division of Biological Sciences, Indian Institute of Science; Bengaluru, 560012, India.

Abstract

The Coronavirus Disease 2019 (COVID-19) pandemic caused by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) has taken its toll on worldwide public health infrastructure. SARS-CoV-2 is reported to exhibit wide tissue tropism, contributing to its severe pathogenicity that often culminates in multiple-organ failure. The onslaught of this disease has intensified due to the emergence of variants of concern (VOC), such as Delta and Omicron. These variants have been linked to gastrointestinal (GI) symptoms, suggesting a potential fecal-oral route of viral transmission. Here we compared the broad tissue tropism of ancestral Hong-Kong SARS-CoV-2 (SARS-CoV-2 HK) against Delta and Omicron VOCs in aa hamster model by analyzing tissue samples collected from the upper and lower respiratory system and the GI tract. We observed an overall increase in vRNA load and pro-inflammatory cytokines, especially in GI tracts of animals infected with Delta virus, indicating selective virus tropism and pathology in these tissues. However, no apparent spike in Delta viral load was observed in the large intestine and fecal matter. Overall, our research investigates the wide range of tissues that various SARS-CoV-2 strains can infect in hamsters and presents evidence supporting the increased preference of Delta VOCs for infecting the GI tract.

Keywords: SARS-CoV-2, GI infection, hamster, VOCs, tissue tropism, feces

Introduction

Coronavirus Disease-19 (COVID-19) is a respiratory infection caused by Severe Acute Respiratory Syndrome Virus-2 (SARS-CoV-2), which is enveloped by positive-sense single-stranded RNA viruses of genus Betacoronavirus, family Coronaviridae (1). The virions are decorated with surface glycoproteins called Spike, which give the appearance of a crown (2). The SARS-CoV-2 spike glycoprotein binds to angiotensin-converting enzyme 2 (ACE2) receptors, and entry into permissive cells is facilitated by cleavage of spike by host proteases like transmembrane protease serine 2 (TMPRSS2) at the cell surface and cathepsin L in endosomes (3) (4).

The virus was first detected in patients epidemiologically linked to the Huanan wet market in Wuhan, China in December 2019 (5, 6). Since then, SARS-CoV-2 has spread throughout the globe, causing ~7 million deaths worldwide as of May 2023 (https://covid19.who.int/). The inherent airborne transmission capability of SARS-CoV-2 and high mutation rates associated with genome plasticity has favored high rates of human-human transmission of the virus on a global scale (7). This has also led to the emergence and spread of novel variants of concern (VOC) which have acquired heightened immune escape, transmission, and fitness advantages. These VOCs, designated Alpha, Beta, Delta, and Omicron variants have emerged independent from one another and throughout the pandemic, become dominant regionally or globally, outperforming previous variants (8). The large-scale use of effective vaccines and antivirals has brought the virus under control, but positive cases still occur sporadically (9).

SevereSevere SARS-CoV-2 infection often results in extrapulmonary manifestations that include gastrointestinal, cardiovascular, hematologic, renal, neurologic, ophthalmologic, and dermatologic systems (10, 11). The governing factors for such broad cell tropism of SARS-CoV-2 mainly include ACE2 distribution across tissues (11) and comorbidities or immunocompromised status of individuals who are more prone to severe infection (12). The pathogenicity of a virus can vary depending on multiple factors, including the host species, immune response, and other genetic and environmental factors. Studies involving animal models, such as hamsters, are valuable for understanding viral pathogenesis and assessing the effectiveness of

potential therapeutics or vaccines against COVID-19 (13). HereHere we compared

the tissue tropism of the most clinically relevant strains of SARS-CoV-2 in the golden

hamster model, which is a highly susceptible model that phenocopies several clinical

aspects of COVID-19 (14, 15). We compared infection and tissue tropism of parental

SARS-COV-2 HK strain with Delta and Omicron VOCs in hamsters and found the

Delta virus to be more prone to infect the GI tract and possibly contribute to fecal

transmission.

MATERIAL AND METHODS

Virus stocks

SARS-CoV-2 Isolate Wuhan (hCov-19/Hong Kong/VM20001061/2020, Cat No.: NR-

52282), Delta variant (USA/PHC658/2021, Lineage B.1.617.2; Cat No.: NR-55611,)

and Omicron variant (USA/PHC658/2021, Cat No.: NR-56461) were obtained from

BEI Resources, NIAID, NIH. All viruses were propagated and titrated by plaque

assay in VEROERO-E6 cells as previously described (16).

Ethics statement

This study was conducted in compliance with institutional biosafety guidelines,

(IBSC/IISc/ST/17/2020; IBSC/IISc/ST/18/2021), following the Indian Council of

Medical Research and Department of Biotechnology recommendations. All

experiments involving animals were reviewed and approved by the Institutional

Animal Ethics Committee (Ref: IAEC/IISc/ST/784/2020) at the Indian Institute of

Science. The experiments were performed according to CPCSEA (The Committee

for Control and Supervision of Experiments on Animals) guidelines.

Animals

Male and Female Syrian Golden Hamsters (*Mesocricetus auratus*) of 6-7 weeks old

(60-70 g) were obtained from Biogen Laboratory Animal Facility, Bengaluru, India. All

the animal experiments were conducted in the viral Biosafety level-3 Laboratory at,

the Indian Institute of Science, Bangalore. Animals (n=3) were maintained in

individually ventilated cages at 23±1°C temperature and 50±10% relative humidity

with 12 h day/night light cycle with food and water *ab libitum*. An overdose of

Ketamine (Bharat Parenterals Limited) and Xylazine (Indian Immunologicals Ltd)
was used to sacrifice animals upon completion of the experiment.

Grouping and virus infection

A total of 28 animals were divided into 7 different groups with 2 males and 2 females
per group. The animals from respective groups were intraperitoneally anesthetized
(Ketamine-150mg/kg; Bharat Parenterals Limited and Xylazine-10mg/kg; Indian
Immunologicals Ltd.) and infected intranasally with either Hongkong, Delta, or
Omicron virus at the dose of 10^5 and 10^6 Plaque Forming Units (PFU)/animal in 100
µL PBS (50µL/nostril). Fresh Feces (in Trizol) and Nasal washes were collected (in
100 µL PBS) before infection, on 2 and 4 days post-infection (Fig.1B). All animals
were sacrificed on day-4 post-infection, and blood was collected by cardiac puncture.
Sera were separated and stored at -80ºC until further use. Organs such as lungs,
esophagus, stomach, small intestine, and large intestine were stored in Trizol (for
RT-PCR) or DMEM (for plaque assay). A part of each organ was used to estimate
the viral RNA load by RT-PCR or plaque assay and another portion for
histopathological analysis.

Clinical signs

111 After infection, hamsters were daily observed for the following clinical signs till day 4
112 p.i. and scored based on severity; Lethargy (none=0, mild=1, moderate=2,
113 severe=3); piloerection (none=0, mild=1, moderate=2, severe=3); abdominal
114 respiration (absent=0, present=1); hunched back (absent=0, present=1) and body
115 weight loss (1-5 %= 1; 6-10%=2; 11-15%=3).

Scoring of lung gross pathology

117 The lungs were scored for focal and diffused hyperemia. Based on the severity of
118 hyperemia, scoring was given on a scale of 1 to 3 (mild=1, moderate=2, severe=3).
119 The total lung weight was recorded and images were captured. The lung-body
120 weight ratio was calculated for each animal on the day of dissection.

Lung viral plaque assay

The plaque assay was performed to determine the viral titer for the tissue
homogenates on VERO-E6 cells in 12-well plates. Briefly, plates were seeded with
0.2×10^6 cells/mL and incubated for 48h at 37ºC. All the samples were serially
diluted 10-fold in DMEM with 2 % FBS. A 100 µL of the diluted sample was added to
each well in duplicates and incubated at 37ºC for 1 h with shaking every 10-15 min.
After incubation, the inoculum was removed completely, overlayed with 1 mL of 0.6
% Avicel and kept at 37ºC with 5 % CO2 for 48 h. The overlay was removed
completely post-incubation and cells were fixed by adding 1 mL/well of 4%
formaldehyde in PBS. Cells were stained with 1% crystal violet solution to visualize
the plaques.

RNA extraction and viral load by Quantitative Real-Time PCR (qRT-PCR)

All the organs were processed using a FastPrep-24™ stainless-steel bead
homogenizer (MP Biomedicals, USA) and total RNA was extracted using TRIzol
(15596018, Thermo Fisher) as per the manufacturer's instructions. RNA from fecal
matter was extracted using the QiaAMP viral RNA mini kit (Cat No. 52906, Qiagen,
US) according to the manufacturer's instructions. A 10µL reaction mixture with 100ng
of RNA per sample, in a 384-well block was used to quantify viral RNA using AgPath-
ID™ One-Step RT-PCR kit (AM1005, Applied Biosystems). The following primers
and probes targeting the SARS CoV-2 N-1 gene were used. Forward primer:
5'GACCCCAAAATCAGCGAAAT3'   and   Reverse   primer: 5'
TCTGGTTACTGCCAGTTGAATCTG3',   Probe: (6-FAM/BHQ-1)
ACCCCGCATTACGTTTGGTGGACC. The Ct values were used to determine viral
copy numbers by generating a standard curve using SARS CoV-2 genomic RNA
standard.

Cytokine gene expression by qRT-PCR

The host mRNA expression of IFN-γ, IL-6, and TNF-α was quantified using suitable
primers with 18s RNA as an internal reference gene. Briefly, cDNA was prepared
from 1µg RNA using PrimeScript™ RT reagent Kit with gDNA Eraser (TaKaRa). RT-
qPCR was performed using 2.5 µL of cDNA, 2.5 µL of the primer mix, and 5 µL of
SYBR™ Green PCR Master Mix (Applied Biosystems). The following primers for
hamster IFN-γ, IL-6, and TNF-α genes were used. **IFN-γ:** Forward primer:
5'TGTTGCTCTCTCTCTCTCAGG3'   and   Reverse   primer:

5'AAGACGAGGTCCCCTCCATTC3'. **IL-6:** Forward primer: 5'AGGCCATCCTGATGGAGAAG3' and Reverse primer: 5' GGTATGCTAAGGCACAGCAC3'. **TNF-α:** Forward primer: 5'TGAGCCATCGTGCCAATG3' and Reverse primer: 5'AGCCCGTCTGCTGGTATC3'. The comparative threshold (Ct) values were used to calculate the mean Log_2 fold change compared to 18S RNA by the delta-delta Ct method.

Cytokine ELISA

The sera obtained from hamsters on day 4 p.i. were used to evaluate the level of IFN-☐ and IL-4 cytokines by sandwich ELISA method using a commercially available kit (ImmunoTag™, USA). The sensitivity of both IFN-☐ and IL-4 kit was 9.375 pg/mL.

Histopathology and Immunohistochemistry

Tissue specimens were fixed in 4% paraformaldehyde in PBS and embedded in paraffin blocks. Tissue sections of 4-6 µm thickness were stained with Hematoxylin and Eosin (H&E). Slides were examined by light microscopy for 3 histological criteria in the lung (Alveolar infiltration and exudation, vasculature inflammation, and peribronchiolar infiltration with epithelial desquamation). Based on the severity, each criterion was scored on a scale of 1 to 3.

The protocol for Immunohistochemistry (IHC) was modified from a previous report (17). Briefly, IHC of the duodenum samples was performed on 5µm paraffin-embedded tissue sections placed on the poly L-lysine coated slides. The sections were deparaffinized and hydrated using xylene and descending grades of alcohol respectively. Antigen retrieval was carried out by heating in sodium citrate buffer (10 mM, pH 6.0) for 15 minutes using a domestic pressure cooker in a steam setting. The sections were quenched for endogenous peroxidase using 3% hydrogen peroxide and protein blocked using 3% Bovine Serum Albumin (BSA) for 20 min each at RT. Slides were then incubated with a primary antibody for SARS-CoV-2 nucleocapsid protein (Novus Biological, NB 100-56576) at 4 °C overnight. Detection was performed using a goat anti-rabbit HRP secondary antibody (Invitrogen, 31460) for 60 min at RT and visualized using 3,3-diaminobenzidine (DAB) (Sigma,

186 11718096001) for 10 min at RT. Counterstaining was done with Harry's
187 Haematoxylin for 1 min. After dehydration and air drying, the sections were cleared
188 in xylene and mounted using Dibutylphthalate polystyrene xylene (DPX) mountant.
189 The mounted slides were viewed under the microscope and representative images
190 were recorded.

Statistical analysis

The data were analyzed using GraphPad Prism v 8.4.3 and represented as mean
±SEM. Statistical variations were determined by one-way ANOVA (Lung/bodyweight
ratio, tissue viral RNA load, and cytokine mRNA level) or two-way ANOVA (Body
weight, clinical signs) with Dunnett's multiple comparisons tests or by paired T-test
(Lung histopathology). Values were significant when *P < 0.05, **P < 0.01 or ***P <
0.001.

Results

Comparison of clinical signs and lung viral load in Syrian Hamsters infected with Delta, Omicron VOCs and ancestral SARS-CoV-2 strain.

Here, we compared the pathogenicity of SARS-CoV-2 HK against Delta (lineage
B.1.617.2) and Omicron (lineage B.1.1.529) VOCs. SARS-CoV-2 HK was originally
isolated from a nasopharyngeal aspirate and throat swab collected from an adult
male patient on January 22, 2020, in Hong Kong, China. The strain is assigned to
lineage A and has been used as part of vaccine preparations
(https://www.beiresources.org/Catalog/animalviruses/NR-
52282.aspxhttps://www.beiresources.org/Catalog/animalviruses/NR-52282.aspx)
(16). The Delta variant emerged in India in late 2020 and the Omicron in Botswana
in November 2021 (18, 19). Delta variant was associated with higher pathogenicity
compared to ancestral or other strains causing more severe illness and
hospitalization rates (18). On the other hand, the Omicron variant was
phylogenetically distinct and possessed over 30 mutations in the spike protein (9,
20). Nucleotide mapping comparing these three viruses is shown in **(Fig. 1A).**

To understand the pathogenicity and tissue tropism of SARS-CoV-2 HKHK in
comparison with Delta and Omicron strains, the body weight, clinical signs, and viral
replication in lungs and gastrointestinal tract were assessed post-infection at two

different doses of 10^5 and 10^6 PFU in hamsters. The viral shedding was also assessed in the nasal route or fecal matter collected at days 0, 2, and 4 post-infections **(Fig. 1B).** The body weight was reduced until 4 days post-infection (dpi) among all the three strains used. The Delta-infected hamsters had more drop in body weight and Omicron infection caused the least reduction in body weight in comparison with control-uninfected animals **(Fig. 1C).** The clinical signs in virus-infected hamsters also showed a similar pattern as of body weight among the strains used. Only Omicron-infected hamsters remained subclinical and started recovering from the drop in body weight **(Fig. 1D).** The viral RNA load as examined by RT-PCR was higher in both SARS HK and Delta-infected hamsters when compared to Omicron. The infectious viral titer as observed by plaque assay also showed a similar pattern of viral load **(Fig. 1F).** The nasal wash samples collected on day 2 showed that Omicron infection resulted in a similar viral load as Delta and Hongkong strain. However, on 4 dpi, the viral burden was lesser in the nasal region of Omicron-infected hamsters **(Sup Fig 1 A, B).**

Figure1. (A) SARS-CoV-2 nucleotide mutation mapping of Hongkong, Delta, and Omicron compared to the Wuhan reference sequence. X-axis represents nucleotide position in the genome and Y-axis shows the different sequence isolates used. Single nucleotide polymorphisms and deleted nucleotide positions are marked using "|", with black indicating a nucleotide deletion and different colors denoting different nucleotide substitutions. Insertions are marked using "v" in dark grey color. The 5'

and 3' untranslated Regions are marked in grey and the different ORFs and
structural proteins are represented in different colors. **(B)** Outline of the work plan
followed in this study. **(C)** Percentage bodyweight of hamsters (n=3 to 4) monitored
over 4 dpi, normalized to mean weight measured at Day 0. **(D)** Clinical signs in
hamsters (n=3 to 4) including piloerection, abdominal respiration, hunched back, and
lethargy were monitored over 4 dpi and scored from 1-4 based on severity. **(E)** The
total SARS CoV-2 viral RNA copy number in the lung was estimated by qRT PCR.
(F) Infectious SARS-CoV-2 titer was quantified by plaque assay. For statistics, data
corresponding to the two doses of Delta were compared respectively against Hong
Kong and Omicron viruses. * $p<0.05$ by paired t-test. Error bars indicate Mean ±
SEM.

Supplementary Figure 1. (A-B) Hamsters were infected with either 10^5 or 10^6
PFU/animal of SARS-CoV-2 Hong-kong strain, or VOCs Delta or Omicron. Nasal
washes were collected from animals at **(A)** 2 and **(B)** 4 dpi and viral RNA was
estimated by qRT PCR.

Comparison of lung pathology and inflammatory cytokine expression in Syrian
Hamsters infected with Delta, Omicron VOCs and Ancestral SARS-CoV-2
strain.

The hamsters infected with Delta showed more gross pathological changes than
SARS-CoV-2 HK and HK Omicron variants at 4 dpi. which was consistent with the
viral load data obtained from RT-PCR and plaque assay. Further, at 10^5 PFU dose,

Omicron infected hamsters didn't show any visible pathology **(Fig. 2A and 2B).**
Additionally, the lung-bodyweight ratio was higher in Delta-infected hamsters with
Hongkong and Omicron-infected hamsters having a lesser but similar lung-body
weight ratio (**Fig. S2**). The hamsters infected with Delta had relatively higher levels
of lung inflammatory cytokines such as TNF-α **(Fig. 2C)**, IFN-□ **(Fig. 2D),** and IL-6
(Fig. 2E) than Hongkong and Omicron variants at both doses studied. It's interesting
to note that, Omicron infection led to a similar level of cytokines expression when
compared to Hongkong infected hamsters. Further, IL-6 levels were consistently
lower among all the strains studied at 10^6 PFU in comparison with 10^5 PFU as
inoculum.

Figure 2. (A-E) Hamsters were infected with either 10^5 or 10^6 PFU/animal of SARS-CoV-2 Hong-kong strain, or VOCs Delta or Omicron. **(A)** Hamsters were sacrificed at 4 dpi and images of whole lungs were captured. **(B)** Lung gross pathology was scored 1-3 based on the increasing severity of focal or diffused hyperemia. Animals were sacrificed 4 dpi and a portion of the lungs was stored In TriZol. Total RNA was

extracted and used to quantify expression levels of **(C)** TNF-α, **(D)** IFN-☐, and **(E)** IL-
6 by qRT PCR.

Supplementary Figure 2. (A-C) Hamsters were infected with either 10^5 or 10^6
PFU/animal of SARS-CoV-2 Hong-kong strain, or VOCs Delta or Omicron. **(A)** The
total weight of the lungs was used to plot the lung-body weight ratio. Animals were
sacrificed 4 dpi and serum was collected from hamsters and used to quantify **(B)**
IFN-gamma and **(C)** IL-4 using a commercially available ELISA kit.

Comparison of lung histopathology and viral antigen localization in Syrian Hamsters infected with Delta, Omicron VOCs and Ancestral SARS-CoV-2 strain.

The histopathological assessment of SARS-CoV-2 infected hamster lungs collected on 4 dpi revealed bronchointerstitial pneumonia including vascular inflammation, alveolar infiltration, and Peribronchiolar infiltration with necrosis. Mononuclear cell infiltrations were also observed in peribronchial and perivascular areas with occasional neutrophils. These lesions, mainly vascular inflammation were more severe in Delta-infected hamsters, especially at 10^6 PFU dose **(Fig. 3A).** The cumulative histopathological lung scores were highest in Delta-infected hamsters, with significant difference between SARS-CoV-2 HK and Delta-infected groups. Consistent with other data, Omicron-infected hamsters showed lung pathological changes similar to as observed in Hongkong infected hamsters **(Fig. 3B).**

Figure 3 (A-B). (A) Histopathology sections of 10^6 PFU/animal group showed (1) bronchial epithelial necrosis and inflammatory changes (2) vascular inflammation (3)

alveolar wall thickening and inflammatory cell infiltration. Bronchial and vascular changes are represented in insets 1 and 2 respectively (200x magnification), while alveolar thickening is in the first column of the figure set (100x magnification). White arrows indicate denuded necrotized bronchiolar epithelial cells admixed with inflammatory cells in the airway lumen, whereas inflammatory cell infiltration in peribronchiolar and perivascular areas are indicated as black arrows (lymphocytes) and black arrowheads (neutrophils). **(B)** Cumulative histopathology scoring was plotted for each group with the selected criteria.

Comparison of viral RNA load and inflammatory marker expression in the GI tract of Syrian Hamsters infected with Delta, Omicron VOCs and Ancestral SARS-CoV-2 strain.

At both the 10^5 and 10^6 PFU doses tested, the viral RNA load was consistently higher in the esophagus, stomach, and small and large intestines of Hongkong and Delta-infected hamsters, with lower viral RNA load being detected in these tissues of Omicron-infected hamsters **(Fig. 4A-D).** Interestingly, the viral RNA load in the small intestine was highest in the Delta-infected hamsters (5.24 log) when compared to Hongkong or Omicron-infected hamsters. Also, viral RNA copy numbers were similar in the large intestine among the Hongkong and Delta-infected hamsters. Surprisingly, the fecal matter collected on 2 and 4 dpi showed a relatively higher viral RNA load in Hongkong-infected hamsters rather than in Delta-infected hamsters **(Fig. S3).**

Figure 4 (A-D). Hamsters were infected with 10^5 and 10^6 PFU/animal of SARS-CoV-2 Hong-Kong strain, or VOCs Delta or Omicron. Animals were sacrificed at 4 dpi and a portion of **(A)** esophagus, **(B)** stomach, **(C)** small, and **(D)** large intestines were stored In TriZol. Total RNA was extracted and viral RNA copy number was quantified by qRT-PCR.

Supplementary Figure 3. (A-B) Hamsters were infected with either 10^5 or 10^6
PFU/animal of SARS-CoV-2 Hong-kong strain, or VOCs Delta or Omicron. The fecal
matter was collected freshly from each animal group at **(A)** 2 and **(B)** 4 dpi and the
viral RNA copy number was estimated by qRT PCR.

**Delta VOC of SARS-CoV-2 infects and causes pathology in the small intestine
of Syrian hamsters.**

To understand the SARS-CoV-2 strain-induced pathology in the regions of the small
intestine, their epithelial morphometric analysis was performed. Here, the villus
length and Cyrpt depth were measured and their ratio was determined. The small
intestines harvested from hamsters infected with Delta variant at both 10^5 and 10^6
PFU doses had higher crypt depth and lesser villi length to crypt depth ratio than
Hongkong or Omicron-infected hamsters which were also statistically significant
compared to the uninfected control hamsters **(Fig. 5A-D).** The Villous length of all
groups was comparable and didn't show any significant difference. In Hongkong and
Omicron-infected hamsters, even though they had relatively similar villi length to
crypt depth ratio, Omicron-infected hamsters had higher epithelial crypt depth **(Fig.
5B-D).).**
The immunohistochemical staining also showed the higher presence of SARS-CoV-2
nucleocapsid protein in the duodenum epithelial cells of the small intestine harvested
from Delta-infected hamsters **(Fig 5E).** Overall, these data emphasize the small
intestine mucosal damage and subsequent regenerative changes possibly caused by
the Delta strain. Further correlating with these data, cytokine levels such as IFN-□,

TNF-α, and IL-6 were relatively higher in the small intestines harvested from Delta-
infected hamsters with similar levels observed among the Hongkong and Omicron-
infected hamsters **(Sup Fig. 4 A-C).** Notably, IFN-□ level was very high (≈ 3 fold)
during Delta infection when 10^5 PFU was used as inoculum **(Sup Fig. 4 A).).**

Figure 5. (A-E) Hamsters were infected with 10^6 PFU/animal of SARS-CoV-2 Hong-
Kong strain, or VOCs Delta or Omicron. Animals were sacrificed at 4 dpi and small
intestine was collected and processed for histology. **(A)** Representative H & E
stained sections of duodenum of normal and different infected groups at 40x

magnification. The black arrow shows the crypt and the white arrow indicates villi.

(B-D) Villi and crypt length were quantified using ImageJ/Fiji. Data shown indicate

(B) Villi length, **(C)** Crypt length, and **(D)** ratio of villi to crypt length. Data from 8-10

villi-crypt units per animal were measured (n=3-4 per group). **(E)** H&E staining of the

duodenum of uninfected and SARS CoV-2 delta VOC challenged hamsters (left) and

Immunohistochemistry analysis (right) with the red arrow indicating the presence of

SARS-CoV-2 N protein. Statistical measurements were done by comparing Delta to

Hong Kong and Omicron. * $p < 0.05$, ** $p < 0.01$ by paired t test. Error bars indicate

mean ±SEM.

Supplementary Figure 4. (A-C) Hamsters were infected with either 10^5 or 10^6

PFU/animal of SARS-CoV-2 Hong-kong strain, or VOCs Delta or Omicron. Animals

were sacrificed at 4 dpi and a portion of small intestines was stored In TriZol. Total

RNA was extracted and used to quantify expression levels of **(A)** IFN-□, **(B)** TNF-α,
and **(C)** IL-6 by qRT PCR.

Discussion

In this study, the SARS-CoV-2 ancestral strain (HK) and Delta VOC infection in
hamsters displayed enhanced clinical signs and higher viral load in the upper
respiratory tract, while the Omicron variant caused attenuated clinical signs. The viral
burden also started dropping down in the nasal regions at 4 dpi, causing less
penetrable to the lungs in the case of Omicron infection at 4 dpi which further
explains its lower replication efficiency as opposed to the Delta variant. These data
were consistent with earlier observations studied on hamsters (21). The earlier *in*
vitro comparative studies on growth kinetics of the Delta and Omicron variants in
Calu3 cells, a lung epithelial cell line, have shown that the Omicron has a poor
replication rate which correlates with our findings of low lung viral copy number
observed in Omicron infected hamsters (22). We observed severe and diffused lung
histopathological lesions with higher inflammatory cytokine levels in Delta-infected
groups when compared to Hongkong and Omicron-infected groups. An earlier report
on lung pathology during Omicron-infection in the hamster model showed that lung
disease is milder in the Omicron group and highest in the Delta groups when
compared to ancestral SARS-CoV-2 (23, 24). Also, contrary to our findings, a study
reported moderate to severe changes in the lungs of Omicron-infected hamsters
which were almost comparable to their Delta-infected group (25). Coronaviruses are
known causative agents of both respiratory and gastrointestinal tract infections in a
wide range of mammals and birds (26). Also, SARS-CoV-2 cell tropism was shown in
multiple organs of human post-mortem samples including lungs, trachea, small
intestines, and other tissues (11). SARS-CoV-2 has been shown earlier to infect and
replicate in both bat and human intestinal organoids, leading to viral replication within
these cells (22). Besides that, *in vitro* studies have shown the higher replicative
ability of the Delta strain in human colon epithelial cell lines such as Caco2 than
Omicron or other strains (27).

ACE2 is known to be an essential receptor for virus entry into the host cells and its
expression level differs among organs and tissues which might determine the
symptoms and outcome of COVID-19. ACE2 expression levels are highest in the

small intestine which suggests that the gastrointestinal tract could be a potential site of viral replication and transmission (28). Along with ACE2, SARS-CoV-2 entry was shown to be dependent on transmembrane serine protease 2 (TMPRSS2), a host protease, which cleaves the viral spike protein once it binds to the ACE2 receptor. The TMPRSS2 expression is also spread across the GI tract. Indeed, the co-expression of ACE2 and TMPRSS2 is highest in human nasal and intestinal epithelial cells with relatively lesser levels in lung and esophageal cells (29). In alignment with this data, the viral RNA was detected in hamsters at a high copy number in the esophagus, stomach, and small intestine, indicating that the virus is directly interacting by binding to the cells of the GI tract. The Delta variant being more replicative and further elevating the levels of inflammatory markers such as IFN-□, TNF-α, and IL-6 shows permissiveness of the intestinal cells and higher pathogenicity of the specific SARS-CoV-2 strain.

The reduced viral load in the large intestine and fecal samples could be mostly specific to the hamster infection model. While GI symptoms are predominantly seen in the severe form of COVID-19 patients (30), hamsters won't show such severe symptoms of SARS-CoV-2 infection, which could be a possible reason for us not being able to detect viral RNA in hamster fecal samples collected on 4 dpi. Additionally, earlier reports showed better ACE2 expression levels in the esophagus, duodenum, and ileum regions of Syrian hamsters with no expression in the large intestine (31). This correlates with our data, as the large intestine had less viral burden which might have reduced the fecal transmission in hamsters. Nevertheless, based on the meta-analysis data, the rate of GI infection and detection of viral RNA in stool samples is under 53 % (32). Also, though the viral RNA gets detected in the respiratory tract at earlier time points, it takes additional 4-5 days to show up in the human stool samples (33).

The villous blunting and increased epithelial regeneration in the duodenal samples of COVID-19 patients have been reported which is consistent with our study where regenerative changes were observed in the duodenum, especially among the Delta-infected hamsters (34). Even though no apparent inflammation was present, viral nucleocapsid protein could be detected in the duodenal epithelial cells of animals infected with the Delta strain (35). This resembles the detection of SARS-CoV-2 protein in intestinal samples of human covid patients without apparent architectural

disruption (33). Overall, the experimental data suggest that the Omicron strain is less
pathogenic and less transmissible when compared to the other two strains. Even
though, Delta and Hongkong strains showed similar viral burdens in many organs,
the Delta strain distinguished itself by causing more severe clinical signs, viral
burden, cellular inflammation, and pathology, especially in the small intestine of
hamsters. The Syrian hamster was also shown here to be permissive to SARS-CoV-
2 in the gastrointestinal tract, where the Delta strain exhibited higher tissue tropism.

The limitations of the study are that, firstly, the gap between the inoculum dose is
relatively lesser which could be a reason for not observing a significant difference in
the parameters studied between the two doses of inoculums used. Secondly, the use
of more animals per group could improve the higher correlation among the different
data generated. Nevertheless, this study warrants further exploration of host factors
required for GI tropism and associated disease by SARS-CoV-2 and other human
coronaviruses.

Acknowledgments

This work is supported by DBT-BIRAC (BT/CS0007/CS/02/20) and DBT-Wellcome
Trust India Alliance (IA/I/18/1/503613) grant to STacknowledge infrastructure and
research support to IISc from the Crypto Relief Fund, L & T Trust, DST-FIST
program, Institute of Eminence Fund, Ministry of Education, and DBT-IISc
partnership program (Phase II). OK, SK and RS acknowledge support from the
MHRD-IISc fellowship and Prime Minister's Research Fellowship (PMRF) and FICCI-
PMRF respectively.

Conflict of Interest

Authors have no conflicts of interest to declare.

References

1.   Gorbalenya AE, Baker SC, Baric RS, de Groot RJ, Drosten C, Gulyaeva AA, et al. The species
Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-
CoV-2. 2020;5(4):536-44.
2.   V'kovski P, Kratzel A, Steiner S, Stalder H, Thiel V. Coronavirus biology and replication:
implications for SARS-CoV-2. Nature Reviews Microbiology. 2021;19(3):155-70.
3.   Letko M, Marzi A, Munster V. Functional assessment of cell entry and receptor usage for
SARS-CoV-2 and other lineage B betacoronaviruses. Nature microbiology. 2020;5(4):562-9.

4.   Koch J, Uckeley ZM, Doldan P, Stanifer M, Boulant S, Lozach PY. TMPRSS2 expression dictates
the entry route used by SARS-CoV-2 to infect host cells. Embo j. 2021;40(16):e107821.

5.   Hui DS, Azhar EI, Madani TA, Ntoumi F, Kock R, Dar O, et al. The continuing 2019-nCoV
epidemic threat of novel coronaviruses to global health—The latest 2019 novel coronavirus outbreak
in Wuhan, China. 2020;91:264-6.

6.   Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. The proximal origin of SARS-CoV-2.
Nat Med. 2020;26(4):450-2.

7.   Phan T. Genetic diversity and evolution of SARS-CoV-2. Infection, genetics and evolution :
journal of molecular epidemiology and evolutionary genetics in infectious diseases. 2020;81:104260.

8.   Carabelli AM, Peacock TP, Thorne LG, Harvey WT, Hughes J, de Silva TI, et al. SARS-CoV-2
variant biology: immune escape, transmission and fitness. Nature Reviews Microbiology.
2023;21(3):162-77.

9.   Tegally H, Moir M, Everatt J, Giovanetti M, Scheepers C, Wilkinson E, et al. Emergence of
SARS-CoV-2 Omicron lineages BA.4 and BA.5 in South Africa. Nature Medicine. 2022;28(9):1785-90.

10.   Gupta A, Madhavan MV, Sehgal K, Nair N, Mahajan S, Sehrawat TS, et al. Extrapulmonary
manifestations of COVID-19. Nature Medicine. 2020;26(7):1017-32.

11.   Liu J, Li Y, Liu Q, Yao Q, Wang X, Zhang H, et al. SARS-CoV-2 cell tropism and multiorgan
infection. Cell Discovery. 2021;7(1):17.

12.   Carrau L, Frere JJ, Golynker I, Fajardo A, Rivera CF, Horiuchi S, et al. Delayed engagement of
host defenses enables SARS-CoV-2 viremia and productive infection of distal organs in the hamster
model of COVID-19. Science signaling. 2023;16(789):eadg5470.

13.   Rosenke K, Meade-White K, Letko M, Clancy C, Hansen F, Liu Y, et al. Defining the Syrian
hamster as a highly susceptible preclinical model for SARS-CoV-2 infection. Emerging microbes &
infections. 2020;9(1):2673-84.

14.   Imai M, Iwatsuki-Horimoto K, Hatta M, Loeber S, Halfmann PJ, Nakajima N, et al. Syrian
hamsters as a small animal model for SARS-CoV-2 infection and countermeasure development.
Proceedings of the National Academy of Sciences of the United States of America.
2020;117(28):16587-95.

15.   Boudewijns R, Thibaut HJ, Kaptein SJF, Li R, Vergote V, Seldeslachts L, et al. STAT2 signaling
restricts viral dissemination but drives severe pneumonia in SARS-CoV-2 infected hamsters. Nat
Commun. 2020;11(1):5838.

16.   Case JB, Bailey AL, Kim AS, Chen RE, Diamond MS. Growth, detection, quantification, and
inactivation of SARS-CoV-2. Virology. 2020;548:39-48.

17.   Tostanoski LH, Wegmann F, Martinot AJ, Loos C, McMahan K, Mercado NB, et al. Ad26
vaccine protects against SARS-CoV-2 severe clinical disease in hamsters. Nature Medicine.
2020;26(11):1694-700.

18.   European Centre for Disease Prevention and Control. Threat assessment brief: emergence of
SARS-CoV-2 B.1.617 variants in India and situation in the EU/EEA

2021.

19.   Viana R, Moyo S, Amoako DG, Tegally H, Scheepers C, Althaus CL, et al. Rapid epidemic
expansion of the SARS-CoV-2 Omicron variant in southern Africa. 2022;603(7902):679-86.

20.   Chakraborty C, Bhattacharya M, Sharma AR, Dhama K, Lee S-S. The rapid emergence of
multiple sublineages of Omicron (B.1.1.529) variant: Dynamic profiling via molecular phylogenetics
and mutational landscape studies. Journal of Infection and Public Health. 2022;15(11):1234-58.

21.   Toomer G, Burns W, Garcia L, Henry G, Biancofiori A, George A, et al. Characterization of
Three Variants of SARS-CoV-2 In Vivo Shows Host-Dependent Pathogenicity in Hamsters, While Not in
K18-hACE2 Mice. Viruses. 2022;14(11).

22.   Zhou J, Li C, Liu X, Chiu MC, Zhao X, Wang D, et al. Infection of bat and human intestinal
organoids by SARS-CoV-2. Nat Med. 2020;26(7):1077-83.

23.   Yuan S, Ye ZW, Liang R, Tang K, Zhang AJ, Lu G, et al. Pathogenicity, transmissibility, and
fitness of SARS-CoV-2 Omicron in Syrian hamsters. Science. 2022;377(6604):428-33.

24.   Suzuki R, Yamasoba D, Kimura I, Wang L, Kishimoto M, Ito J, et al. Attenuated fusogenicity
and pathogenicity of SARS-CoV-2 Omicron variant. Nature. 2022;603(7902):700-5.
25.   Mohandas S, Yadav PD, Sapkal G, Shete AM, Deshpande G, Nyayanit DA, et al. Pathogenicity
of SARS-CoV-2 Omicron (R346K) variant in Syrian hamsters and its cross-neutralization with different
variants of concern. EBioMedicine. 2022;79:103997.
26.   Chan JF, To KK, Tse H, Jin DY, Yuen KY. Interspecies transmission and emergence of novel
viruses: lessons from bats and birds. Trends in microbiology. 2013;21(10):544-55.
27.   Mautner L, Hoyos M, Dangel A, Berger C, Ehrhardt A, Baiker A. Replication kinetics and
infectivity of SARS-CoV-2 variants of concern in common cell culture models. Virol J. 2022;19(1):76.
28.   Zhang H, Li HB, Lyu JR, Lei XM, Li W, Wu G, et al. Specific ACE2 expression in small intestinal
enterocytes may cause gastrointestinal symptoms and injury after 2019-nCoV infection. International
journal of infectious diseases : IJID : official publication of the International Society for Infectious
Diseases. 2020;96:19-24.
29.   Sungnak W, Huang N, Bécavin C, Berg M, Queen R, Litvinukova M, et al. SARS-CoV-2 entry
factors are highly expressed in nasal epithelial cells together with innate immune genes. Nat Med.
2020;26(5):681-7.
30.   Arjmand B, Ghorbani F, Koushki M, Rezai-Tavirani M. Gastrointestinal symptoms in patients
with mild and severe COVID-19: a scoping review and meta-analysis. Gastroenterology and
hepatology from bed to bench. 2020;13(4):321-30.
31.   Suresh V, Parida D, Minz AP, Sethi M, Sahoo BS, Senapati S. Tissue Distribution of ACE2
Protein in Syrian Golden Hamster (Mesocricetus auratus) and Its Possible Implications in SARS-CoV-2
Related Studies. Frontiers in pharmacology. 2020;11:579330.
32.   Tian Y, Rong L, Nian W, He Y. Review article: gastrointestinal features in COVID-19 and the
possibility of faecal transmission. Alimentary pharmacology & therapeutics. 2020;51(9):843-51.
33.   Xiao F, Tang M, Zheng X, Liu Y, Li X, Shan H. Evidence for Gastrointestinal Infection of SARS-
CoV-2. Gastroenterology. 2020;158(6):1831-3.e3.
34.   Lehmann M, Allers K, Heldt C, Meinhardt J, Schmidt F, Rodriguez-Sillke Y, et al. Human small
intestinal infection by SARS-CoV-2 is characterized by a mucosal infiltration with activated CD8(+) T
cells. Mucosal immunology. 2021;14(6):1381-92.
35.   Sia SF, Yan LM, Chin AWH, Fung K, Choy KT, Wong AYL, et al. Pathogenesis and transmission
of SARS-CoV-2 in golden hamsters. Nature. 2020;583(7818):834-8.

A

B

Parameters studied/Days post-infection	Wuhan/Delta/Omicron				
	Day 0	Day 1	Day 2	Day 3	Day 4
Body weight					
Clinical signs					
Nasal wash					
Fecal matter					
Lung (Gross pathology, viral RNA, plaque assay, IFN-Ƴ/IL-6/TNF-α levels and Histology)					
Small Intestine (Viral RNA, IFN-Ƴ/IL-6/TNF-α levels and and Histology)					
Esophagus, stomach, large Intestine (Viral RNA)					
Sera (IFN-Ƴ/IL-4 ELISA)					

C

D

E

Lung Viral RNA

F

Lung Viral Plaque assay

A

Control
uninfected

Wuhan
10^6 PFU

Delta
10^6 PFU

Omicron
10^6 PFU

B

A

Control uninfected | Wuhan 10⁶ PFU | Delta 10⁶ PFU | Omicron 10⁶ PFU

B

Villi length (μm)

Uninfected | Hong-kong | Delta | Omicron

Virus inoculum(10⁶ PFU/animal)

ns | ns

C

Crypt depth (μm)

Uninfected | Hong-kong | Delta | Omicron

Virus inoculum(10⁶ PFU/animal)

** | ns

D

Villi/crypt length ratio

Uninfected | Hong-kong | Delta | Omicron

Virus inoculum(10⁶ PFU/animal)

* | *

E

H&E | IHC

Uninfected

Delta