Comparative population genomics of manta rays has global implications

for management

Emily Humble*[1,2], Jane Hosegood*[2,3], Gary Carvalho[3], Mark de Bruyn[3,4], Simon Creer[3], Guy

M W Stevens[2], Amelia Armstrong[5], Ramon Bonfil[6,7,8], Mark Deakos[9], Daniel Fernando[2,10], Niv

Froman[2], Lauren R Peel[2,11,12], Stephen Pollett[2], Alessandro Ponzo[13], Allan L Poquita[14], Joshua

D Stewart[2,15], Sabine Wintner[16,17] & Rob Ogden[1]

[1]Royal (Dick) School of Veterinary Studies and the Roslin Institute, University of Edinburgh,

EH25 9RG, Edinburgh, UK

[2]The Manta Trust, Catemwood House, Norwood Lane, Corscombe, Dorset, DT2 0NT, UK

[3]Molecular Ecology and Evolution Group, Bangor University, Bangor, UK

[4]Australian Research Centre for Human Evolution, Griffith University, Nathan, QLD, Australia

[5]School of Biomedical Sciences, The University of Queensland, St Lucia, QLD, Australia

[6]Océanos Vivientes AC, Mexico City, Mexico

[7]Consejo Nacional de Humanidades Ciencia y Tecnología (CONAHCyT), Mexico City, Mexico

[8]El Colegio de la Frontera Sur, Unidad Chetumal, Chetumal, Q. Roo, Mexico.

[9]Hawai'i Association for Marine Education and Research, Lahaina, HI, USA

[10]Blue Resources Trust, Colombo, Sri Lanka

[11]Save Our Seas Foundation – D'Arros Research Centre, Geneva, Switzerland

[12]School of Biological Sciences, Oceans Institute and Oceans Graduate School, The

University of Western Australia, Crawley, WA, Australia

[13]Large Marine Vertebrates Research Institute Philippines, Jagna, Philippines

[14]Bureau of Fisheries and Aquatic Resources, Arellano Boulevard, Cebu City, Cebu,

Philippines

[15]Ocean Ecology Lab, Marine Mammal Institute, Department of Fisheries, Wildlife &

Conservation Sciences, Oregon State University, Newport, OR, United States

[16]KwaZulu-Natal Sharks Board, Umhlanga Rocks, South Africa

[17]University of KwaZulu-Natal, School of Life Sciences, Durban, South Africa

*Joint first authors

Corresponding author: Emily Humble, Royal (Dick) School of Veterinary Studies and the

Roslin Institute, University of Edinburgh, EH25 9RG, UK. Email: emily.humble[at]ed.ac.uk

Abstract

Understanding population connectivity and genetic diversity is of fundamental importance to
conservation. However, in globally threatened marine megafauna, challenges remain due to
their elusive nature and wide-ranging distributions. As overexploitation continues to threaten
biodiversity across the globe, such knowledge gaps compromise both the suitability and
effectiveness of management actions. Here, we use a comparative framework to investigate
genetic differentiation and diversity of manta rays, one of the most iconic yet vulnerable groups
of elasmobranchs on the planet. Despite their recent divergence, we show how oceanic manta
rays (*Mobula birostris*) display significantly higher heterozygosity than reef manta rays
(*Mobula alfredi*) and that *M. birostris* populations display higher connectivity worldwide.
Through inferring modes of colonisation, we reveal how both contemporary and historical
forces have likely influenced these patterns, with important implications for population
management. Our findings highlight the potential for fisheries to disrupt population dynamics
at both local and global scales and therefore have direct relevance for international
conservation of marine species.

Running title

Comparative population genomics of manta rays

Main text

Introduction

Understanding the extent to which populations are connected is key to exploring population dynamics, predicting extinction risk and informing conservation management (Hanski & Gilpin, 1991; Lowe & Allendorf, 2010; Mills & Allendorf, 1996). In species with isolated populations characterised by limited dispersal, the risk of extirpation from local depletion is high (Reed, 2004). In such cases, local and regional scale management will be most appropriate for preventing and reversing population declines (Palumbi, 2003). In contrast, species with high rates of gene flow are potentially demographically and genetically more resilient to extrinsic factors (Lowe & Allendorf, 2010; Pascual et al., 2017). However, in order to maintain connectivity and mitigate genetic diversity loss in these taxa, management measures must be coordinated and encompass migratory corridors. As overexploitation and habitat destruction threaten to disrupt population dynamics at a global scale, characterising genetic variation and connectivity has become more important than ever before (Funk et al., 2012; Kardos et al., 2021; Palsbøll et al., 2007).

In widely distributed marine species with high dispersal potential, genetic differentiation is often found to be subtle or non-existent (Palumbi, 2003; Waples, 1998; Ward et al., 1994). Such patterns can arise from a range of mechanisms – from high contemporary gene flow to recent divergence of historically large populations (Palumbi, 2003; Waples et al., 2008; Waples & Gaggiotti, 2006) – and can therefore be difficult to interpret. The latter scenario reflects a disconnect between demographic and genetic connectivity and has important implications for species resilience (Bailleul et al., 2018; Lowe & Allendorf, 2010; Waples, 1998). This is because populations that appear genetically connected may not operate as single demographic units, making them more vulnerable to overexploitation. High-resolution SNP datasets go some way to addressing this problem by providing greater power to detect subtle differences at both neutral and adaptive loci (Gagnaire et al., 2015; Hauser & Carvalho, 2008). However, since population genetic differentiation can be affected by past, as well as contemporary patterns, parallel inference of historical relationships and genetic variation can allow the relative contribution of historical processes to be explicitly evaluated (Foote & Morin, 2016; Liu et al., 2022; Louis et al., 2021). Furthermore, when carried out within a comparative framework, such an approach can provide powerful insights into the drivers of population divergence and therefore improve recommendations for conservation management (Gagnaire, 2020).

Manta rays are large, mobile elasmobranchs inhabiting tropical and sub-tropical oceans
(Couturier et al., 2012) (Figure 1A, C) and provide an excellent opportunity to evaluate the
genomic consequences of historical and contemporary population processes within a
comparative framework. They comprise two described species estimated to have diverged
less than 0.5 Mya as a result of distinct habitat preferences (Kashiwagi et al., 2012). The reef
manta ray (*Mobula alfredi*) frequents near-shore tropical reef environments, such as coral
atolls and barrier reefs (Kashiwagi *et al.* 2011), with a high degree of residency (Deakos *et al.*
2011; Jaine *et al.* 2014; Braun *et al.* 2015; Setyawan *et al.* 2018; Peel *et al.* 2019; Knochel *et*
al. 2022b; Germanov *et al.* 2022). In contrast, while the oceanic manta ray (*Mobula birostris*)
also inhabits near-shore environments, it is often found ranging into sub-tropical habitats along
continental coastlines and at oceanic islands, usually adjacent to productive deep-water
upwellings (Kashiwagi *et al.* 2011; Andrzejaczek *et al.* 2021). As a result of these differences
in habitat use, *M. alfredi* and *M. birostris* have long been considered to display marked
differences in their migratory abilities and levels of gene flow. Yet, only a handful of long-
distance movements have ever been recorded in *M. birostris* (Andrzejaczek et al., 2021;
Knochel, Cochran, et al., 2022) alongside observations of site-fidelity (Cabral et al., 2023;
Garzon et al., 2023; Gordon & Vierus, 2022), raising questions about the extent to which
population structure and genetic variation may differ across species. To date, assessments of
genetic differentiation in *M. alfredi* have focussed on local and regional patterns (Lassauce et
al., 2022; Venables et al., 2021; Whitney et al., 2023) and we have little understanding of how
genetic variation is distributed across the species' range. In, *M. birostris*, the situation is even
less clear, with studies reporting both widespread connectivity and population differentiation
(Hosegood et al., 2020; López et al., 2022; Stewart et al., 2016). Critically, these differences
and uncertainties exist against a background of ongoing global exploitation and uncertain
implications for management.

Targeted and incidental fisheries, driven in part by increasing demand for mobulid gill plates
(Couturier et al., 2012; O'Malley et al., 2017), have led to widespread population declines in
manta rays (Carpenter et al., 2023; Croll et al., 2016; Moazzam, 2018; Rohner et al., 2017;
Ward-Paige et al., 2013). Currently, both species are managed through a patchwork of local,
regional and international measures with varying levels of implementation and enforcement
(Fernando & Stewart, 2021; Lawson et al., 2017; Lawson & Fordham, 2018). To determine
the appropriateness of management measures and assess population vulnerability, a global
assessment of management units is urgently required (Lawson et al., 2017; Stewart, Jaine, et
al., 2018). Here, we undertake a comparative genomic analysis of manta ray populations from
across their global distribution to investigate connectivity, genetic variation, and historical

relationships with an aim to guide effective fisheries management and emphasise the value
of genomic research for advancing knowledge of understudied elasmobranchs.

Materials and Methods

Sample collection

Tissue samples were opportunistically collected from 12 geographic locations to represent the
global distribution of each species (Figure 1A, C). For *M. alfredi* (total *n* = 120), these
originated from the Chagos Archipelago (n = 5), the Maldives (n = 48), Seychelles (n = 24),
Australia Pacific (n = 4), Fiji (n = 9) and Hawaii (n = 30). For *M. birostris* (total *n* = 112), these
originated from Sri Lanka (n = 43), the Philippines (n = 37), South Africa (n = 3), Mexico
Caribbean (n = 4), Mexico Pacific (n = 13) and Peru (n = 12). Samples from Mexico Caribbean,
where a third putative manta ray species occurs in sympatry (*Mobula cf. birostris,* Hinojosa-
Alvarez *et al.* 2016; Hosegood *et al.* 2020), were visually and genetically confirmed as *M.*
birostris. For both species, samples were collected from a combination of live animals and
fisheries specimens (See Supplementary Files for further information).

DNA extraction and ddRAD sequencing

Genomic DNA was extracted using the Qiagen DNeasy Blood and Tissue Kit and quantified
using a Qubit 3.0 Broad Range Assay. Double digest restriction-site associated DNA (ddRAD)
libraries were prepared following the Peterson *et al.* (2012) protocol with modifications
described in Palaiokostas *et al.* (2015) and were 125 bp paired-end sequenced on an Illumina
HiSeq. Sequencing reads from both species were assessed for quality using FastQC and
processed together using the Stacks v2.54 *de novo* assembly pipeline (Catchen et al., 2013).
The three main assembly parameters were chosen following the optimisation procedure
outlined in Rochette and Catchen (2017) (Figure S1 and S2). Initial quality filters were applied
to the resulting genotypes before generating three high-quality datasets for use in downstream
analysis: two species-specific datasets; and one dataset comprising both species. For the
species-specific datasets, we extracted either *M. birostris* or *M. alfredi* individuals, removed
individuals with high relatedness coefficients (Korneliussen & Moltke, 2015; Waples et al.,
2019, Figure S3) and filtered out SNPs with a minor allele count of less than 3, a genotyping
rate less than 90% and that were in linkage disequilibrium using PLINK. For the dataset
comprising both species, we first removed closely related individuals and then filtered out
SNPs with a minor allele count of less than 3 and a genotyping rate less than 90%. Population
genetic summary statistics were calculated using the R package diversity (Keenan et al., 2013,
Table S1). See Supplementary Material for further information on library preparation, read
processing and SNP and individual filtering.

Population structure

To investigate population structure we used the species-specific datasets and three complementary approaches. First, we carried out a discriminant analysis of principal components (DAPC) using the R package adegenet (Jombart, 2008). This approach initially transforms the SNP data using a principal components analysis (PCA) and then performs a discriminant analysis on the retained PCs. This serves to maximise discrimination of individuals between groups while minimising variation within (Jombart et al., 2010). Following the recommendations outlined in Thia (2023), the number of PCs retained as predictors was determined based on the $K–1$ criterion, where K is equal to the number of effective populations. For *M. alfredi*, this was set to 5, under the assumption that each sample site reflects a separate population. For *M. birostris*, this was set to 4 under the assumption that Mexico Pacific and Peru may represent a single population given their close geographic proximity. Second, we estimated admixture proportions for the individuals in each dataset using ADMIXTURE. Admixture runs were performed for ancestry clusters ranging from $K = 1–8$, with 10 runs for each K. The optimal K was identified based on the lowest cross-validation error. The runs with the highest likelihood were visualised. Third, we estimated pairwise genetic differentiation between populations within each species using the Weir and Cockerham F_{ST} value (Weir & Cockerham, 1984) calculated in the R package dartR (Gruber et al., 2018). Confidence intervals and *p*-values were estimated based on bootstrap resampling of individuals within each population 1000 times. *Mobula alfredi* samples from Australia Pacific and *M. birostris* samples from South Africa were excluded from this analysis due to low sample sizes.

Isolation by distance

To investigate patterns of isolation by distance, we examined the relationship between genetic and geographic distance between all pairs of populations in each species. Genetic distances were based on pairwise F_{ST} estimates calculated above. Geographic distances were determined based on a least-cost path analysis implemented using the R package marmap (Pante & Simon-Bouhet, 2013) with a minimum depth constraint of -10 metres in order to prevent paths overland. The significance of associations between genetic and geographic distance matrices was inferred using distance-based Moran's eigenvector maps (dbMEM) by redundancy analysis (RDA, Legendre et al., 2015). For this, geographical distances were transformed into dbMEMs using the R package adespatial, and genetic distances were decomposed into principal components using the R function prcomp. RDA was then performed using the R package vegan, with significance tested using 1000 permutations.

Contemporary gene flow

To infer the strength and directionality of contemporary gene flow between populations we
used the program BA3-SNPs BayesAss v1.1 (Mussmann et al., 2019) which estimates the
proportion of immigrants in a given population using Bayesian inference. This analysis was
restricted to *M. alfredi* as it assumes low levels of connectivity and imposes an upper-bound
on the proportion of non-migrants in a population. We first performed initial runs of BayesAss
to determine optimal mixing parameters (dM = migration rate, dA = allele frequency and dF =
inbreeding coefficient) using the autotune function in BA3-SNPs. We than ran BayesAss-3
with 10,000,000 iterations, a burn-in of 1,000,000 and a sampling interval of 1000. Mixing
parameters were set to dM = 0.21, dA = 0.44 and dF = 0.08. Results were averaged across
five replicate runs and migration rates were considered significant if 95% credible sets (mean
migration rate ± 1.96 x mean standard deviation) did not overlap zero. Chain convergence
was assessed, and migration rates visualised using R (Figure S4).

Historical relationships among populations

To explore historical relationships among populations of *M. alfredi* and *M. birostris* we used
the program TreeMix (Pritchard et al., 2000). TreeMix uses population allele frequencies to
estimate a bifurcating maximum likelihood tree with which to infer historical population splits,
admixture events and the degree of genetic drift. We first supplemented the *M. alfredi* dataset
with one randomly selected *M. birostris* individual and the *M. birostris* dataset with one
randomly selected *M. alfredi* to act as outgroups when rooting the trees. Both datasets were
then filtered for linkage, a minor allele count of less than 3, genotyping rate of less than 90%
and related individuals using PLINK v1.9 (Purcell et al., 2007). Allele frequencies for each
population were then calculated using the –freq and –within arguments in PLINK. For both the
M. birostris and *M. alfredi* datasets we then performed 10 initial runs of TreeMix for each
migration event (M) ranging from 0 to 10. The number of migration edges that explained 99.8%
of the variance was selected as the best model for each species (*M. birostris:* M= 0; *M. alfredi:*
M = 2, Figure S5). We then re-ran TreeMix 100 times using the optimal number of migration
edges. Consensus trees and bootstrap values were estimated and visualised using code
modified from the BITE R package (Milanesi et al., 2017).

Genome-wide heterozygosity

To assess levels of genetic variation both within and between species we used the high-quality
SNP dataset comprising both species. We then calculated multi-locus heterozygosity for each
individual using the R package inbreedR (Stoffel et al., 2016).

Results

High-quality SNPs were genotyped in 173 individuals from 12 locations representing the global
distribution of each species (Figure 1A, C). The species-specific datasets contained a total of

1,553 SNPs in 91 *M. alfredi* individuals, and 6,278 SNPs in 82 *M. birostris* individuals, while

the full dataset contained a total of 15,312 SNPs called across both species. See Materials

and Methods and Supplementary Information for details.

Contrasting patterns of population structure at a global scale

To investigate population differentiation within each species we used four complementary

approaches: discriminant analysis of principal components (DAPC), admixture, pairwise F_{ST}

and isolation by distance analysis. In *M. alfredi*, all methods supported the presence of strong

population structure at both global and regional scales. Populations inhabiting different ocean

basins displayed the highest degree of differentiation in the DAPC, with Pacific and Indian

Ocean populations forming distinct clusters along PC1 (Figure 1B). Regional differentiation

was also detected, with Seychelles, Chagos and the Maldives clustering apart along PC2, and

Hawaii separating from Australia Pacific and Fiji along PC3 (Figure 1B and Figure S6A). These

patterns were reinforced in the admixture analysis which highlighted two major ancestral

source populations, inferred an optimal value of $K = 4$ and resolved hierarchical structure up

to $K = 7$ (Figure S7 and Figure S8A). Interestingly, only weak separation was observed

between Australia Pacific and Fiji, however, this pattern may be confounded by the small

sample size of the former, which can lead to spurious merging of distinct populations

(Puechmaille, 2016). Pairwise F_{ST} estimates between ocean basins were on average over two

times higher than those within (mean pairwise F_{ST} between ocean basins = 0.30, mean

pairwise F_{ST} within ocean basins = 0.13, Figure 2A) yet all population comparisons were found

to be significant (Figure S9A, mean = 0.23, min = 0.08, max = 0.43). Finally, we detected a

significant relationship between pairwise F_{ST} and geographic distance (adjusted $R^2 = 0.65$, *P*

= 0.03) indicating an effect of isolation by distance (Figure 2B).

Figure 1. Contrasting patterns of population structure in manta rays. (A, C) Geographic distributions of (A) *Mobula alfredi* and (C) *Mobula birostris* visualised together with the locations of samples used in this study. Dark shaded distributions denote the confirmed species range and light shaded distributions denote the expected species range. Sampling location points are distinguished by colour and scaled by the number of samples. Further details are provided in the Supplementary Material. (B, D) Scatterplots showing individual variation in principal components (PC) one and two derived from discriminant analysis of principal components analysis for (B) *M. alfredi* and (D) *M. birostris* individuals. The amount of variance explained by each PC is shown in parentheses. Population abbreviations: AP = Australia Pacific, CHAG = Chagos, FIJI = Fiji, HAW = Hawaii, MAL = Maldives, SEY = Seychelles, MC = Mexico Caribbean, MP = Mexico Pacific, PERU = Peru, SA = South Africa, SL = Sri Lanka and PHI = the Philippines.

In stark contrast, *M. birostris* displayed little evidence for strong population structure across all methods. Individuals from different ocean basins clustered closely together along each axis in the DAPC (Figure 1D and Figure S6B). Admixture identified $K = 1$ as the optimal number of clusters, with increasing values of K merely introducing additional mixing (Figure S7 and Figure S8B). Pairwise F_{ST} estimates were two-fold lower than in *M. alfredi*, with no pairwise comparison falling above 0.007 (mean = 0.002, min = -0.005, max = 0.006, Figure 2C). Nevertheless, despite these broad patterns, several lines of evidence indicate the presence

of subtle geographic differentiation in this species. First, individuals from Mexico Pacific, Peru and Mexico Caribbean clustered separately from those sampled in South Africa, Sri Lanka, and the Philippines along PC1 (Figure 1D). Second, despite pairwise F_{ST} estimates being low, comparisons between Eastern-Pacific and Indo-Pacific populations, and between Sri Lanka and the Philippines were statistically significant (Figure S9B). Small F_{ST} values are expected when minor allele frequencies are low and therefore do not necessarily reflect an absence of differentiation (Jakobsson et al., 2013). Furthermore, it is possible that overall levels of population structure were underestimated due to the small sample size of two of our six populations (Puechmaille, 2016). Finally, while no significant relationship was observed between pairwise F_{ST} and geographic distance (adjusted R^2 = 0.50, P = 0.10), there was a tendency for populations separated by greater distances to display higher differentiation (Figure 2D).

Figure 2. Population genetic differentiation, isolation by distance and contemporary migration in manta rays. (A, C) Pairwise F_{ST} estimates between sampling locations for (A) *M. alfredi* and (C) *M. birostris*. Samples from Australia Pacific and South Africa were excluded from this analysis due to low sample sizes. (B, D) Relationship between genetic (F_{ST}) and geographic distance as calculated by least-cost path analysis for all pairwise population comparisons in (B) *M. alfredi* and (D) *M. birostris*. Solid lines and shaded areas reflect the regression slopes and standard errors respectively based on a linear model. Samples from Australia Pacific and South Africa were excluded from this analysis due to low sample sizes. (E) Contemporary gene flow estimates between populations of *M. alfredi*. The direction of each arrow represents the direction of gene flow, and the width of each ribbon reflects the relative amount of gene flow. Population abbreviations: AP = Australia Pacific, CHAG = Chagos, FIJI = Fiji, HAW = Hawaii, MAL = Maldives, SEY = Seychelles, MC = Mexico Caribbean, MP = Mexico Pacific, PERU = Peru, SL = Sri Lanka and PHI = the Philippines.

Contemporary gene flow

To characterise the strength and direction of gene flow between populations we used the program BA3-SNPs (Mussmann et al., 2019) to estimate recent migration. As this method assumes low levels of connectivity and imposes an upper-bound on the proportion of non-migrants in a population, this analysis was restricted to *M. alfredi*. As expected, contemporary gene flow was low (Figure 2E); the average migration rate between populations, measured as the estimated number of migrants per generation, was 0.029 (min = 0.008, max = 0.15), with this figure falling to 0.018 (min = 0.008 , max = 0.03) when considering gene flow between populations in different ocean basins. Migration into both Hawaii and the Maldives was lowest, indicating these populations are the most isolated of those sampled (Table S2). Migration rates were only deemed significant between Seychelles and Chagos (0.15) and between Fiji and Australia Pacific (0.15), in line with these populations being last to separate in the admixture analysis. These patterns highlight that while *M. alfredi* may have the propensity to travel over large distances, restricted movement likely dominates.

Historical relationships among populations

To place patterns of genetic differentiation into a historical context, we investigated population origins and colonisation patterns using TreeMix (Pritchard et al., 2000). This program uses allele frequency data to infer patterns of population splits and admixture events through the construction of a maximum likelihood tree. In *M. alfredi*, internal branch lengths were relatively long, with an initial split clearly separating populations in the Indian and Pacific Oceans (Figure 3A). The Maldives and Australia Pacific were the first to separate within each locality and displayed the lowest levels of genetic drift overall. Hawaii was among the last populations to split and displayed the highest amount of drift, in line with its geographic isolation. The best supported model inferred two migration events (Figure S5A); one from the *M. alfredi* population in Seychelles into *M. birostris*, and one from Chagos into Hawaii. However, because not all geographic regions are represented in our data set, the true sources and sinks of these admixture events may originate from related ghost populations. In contrast to *M. alfredi*, the addition of migration events led to no substantial improvement in the model for *M. birostris* (Figure S5B) and therefore the tree without migration is presented here. Interestingly, internal branch lengths were considerably shorter in *M. birostris*, indicating rapid radiation from a shared ancestral source population (Figure 3B). External branch lengths were also short, consistent with larger populations displaying marginal drift and low divergence. Nevertheless, despite these patterns, some geographic signal was detected in the *M. birostris* tree, with individuals originating from the Eastern Pacific and the Caribbean (Peru, Mexico Caribbean, and Mexico Pacific) grouping separately from those originating from the Atlantic and Indo-

322   Pacific (South Africa, the Philippines, and Sri Lanka), although bootstrap support was overall

323   low.

Figure 3. Historical relationships and heterozygosity in manta rays. (A–B) TreeMix maximum likelihood
consensus tree displaying the historical relationships among (A) *M. alfredi* and (B) *M. birostris*
populations. Horizontal branch lengths reflect the amount of genetic drift that has occurred along each
branch. Bootstrap support values for each node are indicated. Migration edges inferred using TreeMix
are represented as arrows and coloured according to their migration weight. The scale bar reflects 10
times the average standard error of the entries in the sample covariance matrix. (C–E) Variation in
individual multi-locus heterozygosity between (C) species and among populations of (D) *M. alfredi* and
(E) *M. birostris*. Note that Y-axis scales differ for (D) and (E). Centre lines of boxplots reflect the
median, bounds of the boxes extend from the first to the third quartiles, and upper and lower
whiskers reflect the largest and smallest values but no further than 1.5 * the interquartile range
from the hinge. Population abbreviations: AP = Australia Pacific, CHAG = Chagos, FIJI = Fiji, HAW =
Hawaii, MAL = Maldives, SEY = Seychelles, MC = Mexico Caribbean, MP = Mexico Pacific, PERU =
Peru, SA = South Africa, SL = Sri Lanka and PHI = the Philippines.

Heterozygosity landscape across species and populations

To explore how patterns of population structure and colonisation are associated with genome-
wide variation, we compared individual multi-locus heterozygosity between species and
among populations. Strikingly, heterozygosity was on average over three times higher in *M.*
birostris (mean = 0.10, min = 0.053, max = 0.12) than in *M. alfredi* (mean = 0.03, min = 0.01,
max = 0.051), with every individual displaying a higher value than any *M. alfredi* (ß = 0.07,
95% CI = 0.06–0.07, P = <2.2 X 10^{-16}, Figure 3C). This finding is in line with the patterns of
population structure and historical splits we observed in each species. Variation in
heterozygosity was also observed at a population level (Figure 3D–E). In *M. alfredi*, the
Maldives and Australia Pacific had the highest levels of heterozygosity within each ocean
basin, in line with these populations being first to split in the TreeMix analysis. Indian Ocean
populations displayed higher overall heterozygosity than Pacific populations and had mean
values that were overall similar. Interestingly, heterozygosity within Pacific populations
declined steeply from west to east, with heterozygosity in Hawaiian individuals being around
half that of the Australian animals (Figure 3D), in line with this population being last to split in
the TreeMix analysis and displaying the highest amount of drift. In contrast, *M. birostris*
populations displayed less extreme variation in heterozygosity, with mean values differing by
less than 0.02 and having no clear geographical pattern (Figure 3E). Furthermore, variance
within populations was an order of magnitude greater in *M. birostris* than in *M. alfredi*, and was
particularly large in Sri Lanka, the Philippines, and Mexico Pacific populations.

Discussion

Manta rays are iconic megafauna with cultural, socio-economic and ecological significance.
Due to targeted and bycatch fisheries operating across their broad-ranging distributions,
populations are declining worldwide. Elucidating levels of connectivity and genetic variation
remains a crucial priority for conservation management. We use reduced-representation
sequencing on a global set of samples and reveal striking differences in the population genetic
landscape of two recently diverged manta ray species. By considering the influence of both
contemporary and historical processes, our study provides a precautionary framework for
assessing conservation units in widely distributed marine species.

We first demonstrate the presence of strong genetic differentiation in *M. alfredi* at a global and
regional scale. From a total of six sampling locations, we found evidence for at least five
genetically distinct, and by extension, demographically independent populations. Two of these
were separated by a distance of ~1,200 km, which is close to the maximum recorded
movement in the species (Armstrong et al., 2019; Jaine et al., 2014), indicating that long
distance migrations are likely rare. Indeed, contemporary gene flow was low – especially
between geographically distant locations – with only a small proportion of individuals in any

population being identified as first- or second-generation migrants. Furthermore, when gene
flow was observed, it tended to be unidirectional. These results are in line with recent studies
demonstrating population differentiation between Western Australia and Mozambique
(Venables et al., 2021) and between Eastern Australia and New Caledonia (Lassauce et al.,
2022), together highlighting how large ocean basins form significant barriers to dispersal in
coastal elasmobranchs (Hirschfeld et al., 2021). High site-fidelity has been widely reported in
M. alfredi based on tagging and mark-recapture studies (Braun et al., 2015; Deakos et al.,
2011; Germanov et al., 2022; Jaine et al., 2014; Knochel, Hussey, et al., 2022; Peel et al.,
2019; Setyawan et al., 2018). However, the degree of residency has been shown to vary, with
movements rarely exceeding a few hundred kilometres in some locations (Braun et al., 2015;
Deakos et al., 2011; Kessel et al., 2017; Knochel, Hussey, et al., 2022; Setyawan et al., 2018)
yet reaching over 1,000 km in others (Armstrong et al., 2019; Germanov & Marshall, 2014).
Our study presents a comparatively broad-scale analysis relevant for regional and global
management planning. Further work on local patterns of population structure will shed light on
the nuances and drivers of fine-scale movement patterns in this species (Whitney et al., 2023).

To explore the mechanism by which manta rays colonised their distribution, we reconstructed
historical relationships and assessed levels of heterozygosity. In *M. alfredi*, we found strong
evidence for an initial split between the Indian and Pacific Oceans followed by further
separation within each locality. Furthermore, populations in the Pacific displayed a signal of
declining heterozygosity from west to east, together suggesting that *M. alfredi* underwent a
stepping-stone pattern of range expansion from an Indo-Pacific Ocean origin, involving
opportunistic long-range movements and associated founder events. This is consistent with a
recent observation of a pregnant *M. alfredi* individual at Cocos Island, Costa Rica (Arauz et
al., 2019), almost 6,000 km east of the nearest confirmed sighting, and the first record of *M.
alfredi* in the Eastern Pacific. Range expansion inherently impacts genetic variation, with a
stepping-stone model of colonisation predicted to result in the strongest cumulative effect of
founder events (Le Corre & Kremer, 1998). Among our sampled populations, Hawaii is the
most geographically isolated, situated at the edge of the *M. alfredi* distribution. Interestingly,
not only was Hawaii the most genetically differentiated from all populations in our study, but it
displayed the longest external branch lengths in the TreeMix analysis and the lowest levels of
heterozygosity. This could be suggestive of a single founder event by a small population.
Genetic variation is fundamental for enabling populations to adapt in response to selection
(Bonnet et al., 2022; Kardos et al., 2021; Lai et al., 2019). Our findings therefore expose how
isolated *M. alfredi* populations at the periphery of their distribution may be intrinsically more
vulnerable to changing environmental conditions and the genetic impacts of population
decline.

In stark contrast to the patterns observed in *M. alfredi*, *M. birostris* displayed markedly higher
levels of heterozygosity and only subtle genetic differentiation across ocean basins. Weak
population structure is common in highly mobile marine species (Leslie & Morin, 2018; Nikolic
et al., 2023; Vignaud et al., 2014), yet warrants careful interpretation, particularly considering
management recommendations (Younger et al., 2017). On the one hand, these findings may
be an indication of high contemporary gene flow and low natal philopatry, in line with the
species' occurrence at remote oceanic islands, tendency to range into sub-tropical habitats
and lower overall re-sight rates than *M. alfredi* (Couturier et al., 2014; Harty et al., 2022;
Rambahiniarison et al., 2023). To date, our understanding of the movement behaviour in *M.*
birostris has largely been based on coastal aggregations of adult individuals over relatively
short timeframes (Beale et al., 2019; Harty et al., 2022; Rohner et al., 2013; Stewart et al.,
2016). Such studies have a tendency to capture seasonal migrations as opposed to dispersal
events and may explain why only a handful of long-distance (~1,000 km) movements have
been recorded in the species (Andrzejaczek et al., 2021; Knochel, Cochran, et al., 2022). In
an infinite island model, only a few migrants per generation are required to obscure strong
population structure when N_e is large (Wright, 1931) and therefore it is possible the patterns
we observe translate to infrequent dispersal events. Furthermore, dispersal could be
segregated by age and/or sex (McClain et al., 2022; Phillips et al., 2021), and may vary among
individuals (Papastamatiou et al., 2013; Perryman et al., 2022; Thorburn et al., 2019). While
challenging, there is benefit in extending future tagging efforts to transient individuals away
from known aggregation sites (Garzon et al., 2023), as well as previously underrepresented
age classes – such as juveniles – to capture what may be infrequent yet evolutionarily relevant
movements.

An alternative explanation for the patterns we observe in *M. birostris* is that insufficient time
has elapsed to reliably identify recent genetic divergence among localities. In contrast to *M.*
alfredi, our TreeMix analysis indicated that *M. birostris* rapidly radiated from a large ancestral
source, with only marginal genetic drift occurring between regions. This was further evidenced
by substantially higher levels of genetic variation that differed little across sampling locations.
In addition, little differentiation was observed between Mexico Pacific and Mexico Caribbean,
two regions that have been geographically separated since the emergence of the Isthmus of
Panama. These findings are consistent with a recent mark-resight analysis that estimated the
population of *M. birostris* in coastal Ecuador to number at least 22,000 individuals (Harty et
al., 2022). Large effective population sizes and high genetic variation increase the time taken
for populations to diverge due to genetic drift (Bailleul et al., 2018; Taylor & Dizon, 1996;
Wright, 1931). This is further compounded in species with long and overlapping generations

(Hoffman et al., 2017) as is the case for manta rays (Dulvy et al., 2014). Taken together,
genetic similarities among *M. birostris* localities may be partially confounded by recent shared
ancestry and large effective population size.

On the basis of these considerations, we propose that a combination of large historical
population size and contemporary gene flow have contributed to the comparatively high levels
of heterozygosity and genetic homogeneity in *M. birostris*. The subtle population differentiation
we observe between the Indian Ocean, South-East Asia and the Eastern Pacific is likely best
explained by the geographic limits of dispersal as opposed to complete geographic isolation.
Yet, unlike in *M. alfredi* where genetic clusters almost certainly reflect discrete demographic
units relevant for conservation management, the extent to which genetic connectivity in *M.*
birostris reflects demographic connectivity is less clear. For example, in extreme cases, the
number of migrants required to eliminate signals of population structure will not be enough to
demographically link populations, and more importantly, replenish those that have been
depleted (Waples, 1998). Interestingly, while re-sight rates are typically lower in *M. birostris*
than *M. alfredi*, demographic independence has been implicated in several mark-recapture
studies where re-sightings follow predictable patterns (Beale et al., 2019; Cabral et al., 2023).
Furthermore, a population genetic analysis based on F_{ST} outliers uncovered allele frequency
differences between two Mexican locations and Sri Lanka (Stewart et al., 2016), suggesting
recent divergence against a background of ongoing gene flow. Taken together, we highlight
the potential for further work investigating adaptive divergence between *M. birostris*
populations and emphasise the need to combine molecular measures of connectivity with
empirical demographic data in this species (Cayuela et al., 2018; Lowe & Allendorf, 2010;
Younger et al., 2017).

Conservation implications
The remarkable differences we observe in the population genetics of manta rays directly
inform likely response to continued exploitation and respective conservation measures. At
present, *M. alfredi* is among the most protected mobulid species worldwide, with some
management frameworks in place at local, national, and international levels (Lawson et al.,
2017; Stevens et al., 2018). Our findings of global population structure underline how local
initiatives recognising populations as distinct management units will be most appropriate for
this species. However, we also demonstrate the consequence of geographic isolation on
genetic variation and reveal how *M. alfredi* likely faces a greater risk from local depletion. This
is especially true for populations at the edge of the species range and in regions with high
coastal fishing pressure. Prioritising these populations in conservation action plans and

maintaining local connectivity will therefore be crucial for boosting resilience and preventing
local extinction in this vulnerable species.

The implications of our findings for *M. birostris* are more nuanced. Despite detecting only
subtle population genetic differentiation, we cannot rule out the possibility that historical
processes and large effective population size are obscuring a higher degree of contemporary
demographic separation. Together with studies reporting high site-fidelity and restricted
movement patterns, our findings strongly suggests that local and national management action
should be considered essential for protecting resident aggregations of *M. birostris*.
Nevertheless, we expect that weak population structure and high genetic variation are
simultaneously being driven by some degree of contemporary dispersal. Consequently, any
fishing activity taking place along migratory corridors threatens to disrupt a mode of gene flow
that may be fundamental for long-term resilience of the species. Similarly, although we have
limited understanding of the number and distribution of breeding and nursery grounds
(Knochel, Cochran, et al., 2022; Pate & Marshall, 2020; Stewart, Nuttall, et al., 2018),
significant reduction of local stocks may impact long-term recruitment at oceanic and even
global scales. We therefore emphasise the escalating need to improve the implementation of
regional and international measures that seek to protect taxa in the high seas. Together with
local scale management, appropriate evidence-based actions will contribute to maintaining
large, connected and genetically diverse populations of manta rays into the future.

Acknowledgements

We are grateful to the following people and organisations for their support sourcing and
collecting tissue samples: Kerstin Forsberg, Jon Slayer, John Nevill, Dan Bowling, Heather
Pacey, Karen Fuentes, Manta Caribbean Project, Annie Murray, Tam Sawers, Beth Taylor,
Adam Thol'hath, Isabel Ender, Musa Mohamed, the Barefoot Collection, Four Seasons
Resorts Maldives, Six Senses Laamu and all Manta Trust and LAMAVE staff, volunteers and
field team. We are grateful to Helen Senn and Jennifer Kaden from the Royal Zoological
Society of Scotland for support during library preparation. ddRAD sequencing was carried out
by Edinburgh Genomics. Tissue sample collection in Seychelles was approved by the
Seychelles Bureau of Standards, the Seychelles Ministry of Environment, Energy and Climate
Change, and The University of Western Australia (RA/3/100/1480). Blue Resources Trust
(BRT) would like to thank the Department of Wildlife Conservation and the Department of
Fisheries and Aquatic Resources for enabling the fieldwork to be conducted in Sri Lanka.
Sample collection in the Maldives was made possible thanks to the permission and support
provided by the Ministry of Fisheries and Agriculture ([OTHR]30-D/INDIV/ 2016/220) and the

Ministry of Environment and Energy (EPA/2016/PSR-M02). LAMAVE would like to thank the
Bohol Environmental Management Office, the DA-Bureau of Fishery Region 7, the local
communities and Mayors of Jagna and Balcayon (Bohol) and the LAMAVE field staff and
volunteers including Joshua Rambahiniarison, Christina Pahang and Maita Verdote. Sampling
in Mexico Caribbean was made possible thanks to CONAPESCA (PPF/DGOPA-126/16 and
PPF/DGOPA/315717)   and   SEMARNAT   (SGPA/DGVS/07618/15   and
SGPA/DGVS/10067/17). We would like to acknowledge Richard Coleman for helpful
comments on the manuscript. For the purpose of open access, the author has applied a
Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version
arising from this submission.

Funding

We are grateful to the Save Our Seas Foundation and The People's Trust for Endangered
Species who provided funding for this work. JH was supported by a Natural Environment
Research Council CASE studentship through the ENVISION DTP (case partner: Royal
Zoological Society of Scotland) and received additional grants from the Fisheries Society of
the British Isles and the Genetics Society. Fieldwork in Seychelles was supported by the
SOSF-D'Arros Research Centre. RB was funded by the Save Our Seas Foundation and the
Marine Conservation Action Fund of the New England Aquarium. BRT acknowledges the
generous support provided by the Save Our Seas Foundation and the Marine Conservation
and Action Fund (MCAF) for field work and sample collection.

Author contributions

EH, JH, GC, MdB, SC, GMWS and RO conceived and designed the study. GMWS, AA, RB,
MD, DF, NF, LRP, SP, AP, JDS and SW provided samples. JH and JK carried out laboratory
work. EH analysed the data with input from JH. EH wrote the paper with input from all other
co-authors.

Competing interests

The authors declare no competing interests.

Data accessibility statement

Sequencing data have been deposited to the European Nucleotide Archive under study
accession   number   PRJEB66437.   Analysis   code   is   available   at
https://github.com/elhumble/manta_pop_gen_2022.

Benefit-sharing statement

A research collaboration was developed with scientists from regions providing genetic samples to advance the conservation of manta rays and their relatives through evidence based research, international collaboration and institutional capacity building. Data and analysis pipelines associated with this work have been shared on public databases for the benefit of this wider cause.

References

Andrzejaczek, S., Schallert, R. J., Forsberg, K., Arnoldi, N. S., Cabanillas-Torpoco, M., Purizaca, W., & Block, B. A. (2021). Reverse diel vertical movements of oceanic manta rays off the northern coast of Peru and implications for conservation. *Ecological Solutions and Evidence*, *2*(1), e12051. https://doi.org/10.1002/2688-8319.12051

Arauz, R., Chávez, E. J., Hoyos-Padilla, E. M., & Marshall, A. D. (2019). First record of the reef manta ray, Mobula alfredi, from the eastern Pacific. *Marine Biodiversity Records*, *12*(1), 3. https://doi.org/10.1186/s41200-019-0162-9

Armstrong, A. O., Armstrong, A. J., Bennett, M. B., Richardson, A. J., Townsend, K. A., & Dudgeon, C. L. (2019). Photographic identification and citizen science combine to reveal long distance movements of individual reef manta rays *Mobula alfredi* along Australia's east coast. *Marine Biodiversity Records*, *12*(1), 14. https://doi.org/10.1186/s41200-019-0173-6

Bailleul, D., Mackenzie, A., Sacchi, O., Poisson, F., Bierne, N., & Arnaud-Haond, S. (2018). Large-scale genetic panmixia in the blue shark (*Prionace glauca*): A single worldwide population, or a genetic lag-time effect of the "grey zone" of differentiation? *Evolutionary Applications*, *11*(5), 614–630. https://doi.org/10.1111/eva.12591

Beale, C. S., Stewart, J. D., Setyawan, E., Sianipar, A. B., & Erdmann, M. V. (2019). Population dynamics of oceanic manta rays (*Mobula birostris*) in the Raja Ampat Archipelago, West Papua, Indonesia, and the impacts of the El Niño–Southern Oscillation on their movement ecology. *Diversity and Distributions*, *25*(9), 1472–1487. https://doi.org/10.1111/ddi.12962

Bonnet, T., Morrissey, M. B., de Villemereuil, P., Alberts, S. C., Arcese, P., Bailey, L. D., Boutin, S., Brekke, P., Brent, L. J. N., Camenisch, G., Charmantier, A., Clutton-Brock, T. H., Cockburn, A., Coltman, D. W., Courtiol, A., Davidian, E., Evans, S. R., Ewen, J. G.,

Festa-Bianchet, M., … Kruuk, L. E. B. (2022). Genetic variance in fitness indicates rapid

contemporary adaptive evolution in wild animals. *Science*, *376*(6596), 1012–1016.

https://doi.org/10.1126/science.abk0853

Braun, C. D., Skomal, G. B., Thorrold, S. R., & Berumen, M. L. (2015). Movements of the reef

manta ray (*Manta alfredi*) in the Red Sea using satellite and acoustic telemetry. *Marine*

Biology, *162*(12), 2351–2362. https://doi.org/10.1007/s00227-015-2760-3

Cabral, M. M. P., Stewart, J. D., Marques, T. A., Ketchum, J. T., Ayala-Bocos, A., Hoyos-Padilla,

E. M., & Reyes-Bonilla, H. (2023). The influence of El Niño Southern Oscillation on the

population dynamics of oceanic manta rays in the Mexican Pacific. *Hydrobiologia*, *850*(2),

257–267. https://doi.org/10.1007/s10750-022-05047-9

Carpenter, M., Parker, D., Dicken, M. L., & Griffiths, C. L. (2023). Multi-decade catches of manta

rays (*Mobula alfredi, M. birostris*) from South Africa reveal significant decline. *Frontiers in*

Marine Science, *10*. https://www.frontiersin.org/articles/10.3389/fmars.2023.1128819

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: An

analysis tool set for population genomics. *Molecular Ecology*, *22*(11), 3124–3140.

Cayuela, H., Rougemont, Q., Prunier, J. G., Moore, J.-S., Clobert, J., Besnard, A., & Bernatchez,

L. (2018). Demographic and genetic approaches to study dispersal in wild animal

populations: A methodological review. *Molecular Ecology*, *27*(20), 3976–4010.

https://doi.org/10.1111/mec.14848

Couturier, L. I. E., Dudgeon, C. L., Pollock, K. H., Jaine, F. R. A., Bennett, M. B., Townsend, K.

A., Weeks, S. J., & Richardson, A. J. (2014). Population dynamics of the reef manta ray

Manta alfredi in eastern Australia. *Coral Reefs*, *33*(2), 329–342.

https://doi.org/10.1007/s00338-014-1126-5

Couturier, L. I. E., Marshall, A. D., Jaine, F. R. A., Kashiwagi, T., Pierce, S. J., Townsend, K. A.,

Weeks, S. J., Bennett, M. B., & Richardson, A. J. (2012). Biology, ecology and

conservation of the Mobulidae. *Journal of Fish Biology*, *80*(5), 1075–1119.

https://doi.org/10.1111/j.1095-8649.2012.03264.x

Croll, D. A., Dewar, H., Dulvy, N. K., Fernando, D., Francis, M. P., Galván-Magaña, F., Hall, M.,

Heinrichs, S., Marshall, A., Mccauley, D., Newton, K. M., Notarbartolo di Sciara, G.,

O'Malley, M., O'Sullivan, J., Poortvliet, M., Roman, M., Stevens, G., Tershy, B. R., &

White, W. T. (2016). Vulnerabilities and fisheries impacts: The uncertain future of manta

and devil rays. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *26*(3), 562–

575. https://doi.org/10.1002/aqc.2591

Deakos, M. H., Baker, J. D., & Bejder, L. (2011). Characteristics of a manta ray *Manta alfredi*

-population off Maui, Hawaii, and implications for management. *Marine Ecology Progress*

Series, *429*, 245–260. https://doi.org/10.3354/meps09085

Dulvy, N. K., Pardo, S. A., Simpfendorfer, C. A., & Carlson, J. K. (2014). Diagnosing the

dangerous demography of manta rays using life history theory. *PeerJ*, *2*, e400.

https://doi.org/10.7717/peerj.400

Fernando, D., & Stewart, J. D. (2021). High bycatch rates of manta and devil rays in the "small-

scale" artisanal fisheries of Sri Lanka. *PeerJ*, *9*, e11994.

https://doi.org/10.7717/peerj.11994

Foote, A. D., & Morin, P. A. (2016). Genome-wide SNP data suggest complex ancestry of

sympatric North Pacific killer whale ecotypes. *Heredity*, *117*(5), Article 5.

https://doi.org/10.1038/hdy.2016.54

Funk, W. C., McKay, J. K., Hohenlohe, P. A., & Allendorf, F. W. (2012). Harnessing genomics for

delineating conservation units. *Trends in Ecology & Evolution*, *27*(9), 489–496.

https://doi.org/10.1016/j.tree.2012.05.012

Gagnaire, P.-A. (2020). Comparative genomics approach to evolutionary process connectivity.

Evolutionary Applications, *13*(6), 1320–1334. https://doi.org/10.1111/eva.12978

Gagnaire, P.-A., Broquet, T., Aurelle, D., Viard, F., Souissi, A., Bonhomme, F., Arnaud-Haond,

S., & Bierne, N. (2015). Using neutral, selected, and hitchhiker loci to assess connectivity

of marine populations in the genomic era. *Evolutionary Applications*, *8*(8), 769–786.

https://doi.org/10.1111/eva.12288

Garzon, F., Seymour, Z. T. A., Monteiro, Z. L., & Graham, R. T. (2023). Spatial ecology of a

newly described oceanic manta ray population in the Atlantic Ocean. *Marine Biology*,

170(6), 68. https://doi.org/10.1007/s00227-023-04219-y

Germanov, E. S., & Marshall, A. D. (2014). Running the gauntlet: Regional movement patterns of

Manta alfredi through a complex of parks and fisheries. *PLOS ONE*, *9*(10), e110071.

https://doi.org/10.1371/journal.pone.0110071

Germanov, E. S., Pierce, S. J., Marshall, A. D., Hendrawan, I. G., Kefi, A., Bejder, L., &

Loneragan, N. (2022). Residency, movement patterns, behavior and demographics of

reef manta rays in Komodo National Park. *PeerJ*, *10*, e13302.

https://doi.org/10.7717/peerj.13302

Gordon, L., & Vierus, T. (2022). First photographic evidence of oceanic manta rays (*Mobula*

birostris) at two locations in the Fiji islands. *PeerJ*, *10*, e13883.

https://doi.org/10.7717/peerj.13883

Gruber, B., Unmack, P. J., Berry, O. F., & Georges, A. (2018). dartr: An r package to facilitate

analysis of SNP data generated from reduced representation genome sequencing.

Molecular Ecology Resources, *18*(3), 691–699. https://doi.org/10.1111/1755-0998.12745

Hanski, I., & Gilpin, M. (1991). Metapopulation dynamics: Brief history and conceptual domain.

Biological Journal of the Linnean Society, *42*(1–2), 3–16. https://doi.org/10.1111/j.1095-

8312.1991.tb00548.x

Harty, K., Guerrero, M., Knochel, A. M., Stevens, G. M. W., Marshall, A., Burgess, K., & Stewart,

J. D. (2022). Demographics and dynamics of the world's largest known population of

oceanic manta rays *Mobula birostris* in coastal Ecuador. *Marine Ecology Progress*

Series, *700*, 145–159. https://doi.org/10.3354/meps14189

Hauser, L., & Carvalho, G. R. (2008). Paradigm shifts in marine fisheries genetics: Ugly

hypotheses slain by beautiful facts. *Fish and Fisheries*, *9*(4), 333–362.

https://doi.org/10.1111/j.1467-2979.2008.00299.x

Hinojosa-Alvarez, S., Walter, R. P., Diaz-Jaimes, P., Galván-Magaña, F., & Paig-Tran, E. M.

(2016). A potential third Manta Ray species near the Yucatán Peninsula? Evidence for a

recently diverged and novel genetic Manta group from the Gulf of Mexico. *PeerJ*, *4*,

e2586. https://doi.org/10.7717/peerj.2586

Hirschfeld, M., Dudgeon, C., Sheaves, M., & Barnett, A. (2021). Barriers in a sea of

elasmobranchs: From fishing for populations to testing hypotheses in population

genetics. *Global Ecology and Biogeography*, *30*(11), 2147–2163.

https://doi.org/10.1111/geb.13379

Hoffman, J. R., Willoughby, J. R., Swanson, B. J., Pangle, K. L., & Zanatta, D. T. (2017).

Detection of barriers to dispersal is masked by long lifespans and large population sizes.

Ecology and Evolution, *7*(22), 9613–9623. https://doi.org/10.1002/ece3.3470

Hosegood, J., Humble, E., Ogden, R., de Bruyn, M., Creer, S., Stevens, G. M. W., Abudaya, M.,

Bassos-Hull, K., Bonfil, R., Fernando, D., Foote, A. D., Hipperson, H., Jabado, R. W.,

Kaden, J., Moazzam, M., Peel, L. R., Pollett, S., Ponzo, A., Poortvliet, M., … Carvalho,

G. (2020). Phylogenomics and species delimitation for effective conservation of manta

and devil rays. *Molecular Ecology*, *29*(24), 4783–4796.

https://doi.org/10.1111/mec.15683

Jaine, F. R. A., Rohner, C. A., Weeks, S. J., Couturier, L. I. E., Bennett, M. B., Townsend, K. A.,

& Richardson, A. J. (2014). Movements and habitat use of reef manta rays off eastern

Australia: Offshore excursions, deep diving and eddy affinity revealed by satellite

telemetry. *Marine Ecology Progress Series*, *510*, 73–86.

https://doi.org/10.3354/meps10910

Jakobsson, M., Edge, M. D., & Rosenberg, N. A. (2013). The relationship between *Fst* and the

frequency of the most frequent allele. *Genetics*, *193*(2), 515–528.

https://doi.org/10.1534/genetics.112.144758

Jombart, T. (2008). adegenet: A R package for the multivariate analysis of genetic markers.

Bioinformatics, *24*(11), 1403–1405.

Jombart, T., Devillard, S., & Balloux, F. (2010). Discriminant analysis of principal components: A

new method for the analysis of genetically structured populations. *BMC Genetics*, *11*(1),

94. https://doi.org/10.1186/1471-2156-11-94

Kardos, M., Armstrong, E. E., Fitzpatrick, S. W., Hauser, S., Hedrick, P. W., Miller, J. M.,

Tallmon, D. A., & Funk, W. C. (2021). The crucial role of genome-wide genetic variation

in conservation. *Proceedings of the National Academy of Sciences*, *118*(48),

e2104642118. https://doi.org/10.1073/pnas.2104642118

Kashiwagi, T., Marshall, A. D., Bennett, M. B., & Ovenden, J. R. (2012). The genetic signature of
recent speciation in manta rays (Manta alfredi and M. birostris). *Molecular Phylogenetics*
and Evolution, 64(1), 212–218.

Keenan, K., McGinnity, P., Cross, T. F., Crozier, W. W., & Prodöhl, P. A. (2013). diveRsity: An R
package for the estimation and exploration of population genetics parameters and their
associated errors. *Methods in Ecology and Evolution, 4*(8), 782–788.
https://doi.org/10.1111/2041-210X.12067

Kessel, S. T., Elamin, N. A., Yurkowski, D. J., Chekchak, T., Walter, R. P., Klaus, R., Hill, G., &
Hussey, N. E. (2017). Conservation of reef manta rays (*Manta alfredi*) in a UNESCO
World Heritage Site: Large-scale island development or sustainable tourism? *PLOS*
ONE, 12(10), e0185419. https://doi.org/10.1371/journal.pone.0185419

Knochel, A. M., Cochran, J. E. M., Kattan, A., Stevens, G. M. W., Bojanowksi, E., & Berumen, M.
L. (2022). Crowdsourced data reveal multinational connectivity, population
demographics, and possible nursery ground of endangered oceanic manta rays in the
Red Sea. *Aquatic Conservation: Marine and Freshwater Ecosystems, 32*(11), 1774–
1786. https://doi.org/10.1002/aqc.3883

Knochel, A. M., Hussey, N. E., Kessel, S. T., Braun, C. D., Cochran, J. E. M., Hill, G., Klaus, R.,
Checkchak, T., Elamin El Hassen, N. M., Younnis, M., & Berumen, M. L. (2022). Home
sweet home: Spatiotemporal distribution and site fidelity of the reef manta ray (*Mobula*
alfredi) in Dungonab Bay, Sudan. *Movement Ecology, 10*(1), 22.
https://doi.org/10.1186/s40462-022-00314-9

Korneliussen, T., & Moltke, I. (2015, December 15). *NgsRelate: A software tool for estimating*
pairwise relatedness from next-generation sequencing data. Bioinformatics (Oxford,
England); Bioinformatics. https://doi.org/10.1093/bioinformatics/btv509

Lai, Y.-T., Yeung, C. K. L., Omland, K. E., Pang, E.-L., Hao, Y., Liao, B.-Y., Cao, H.-F., Zhang,
B.-W., Yeh, C.-F., Hung, C.-M., Hung, H.-Y., Yang, M.-Y., Liang, W., Hsu, Y.-C., Yao, C.-
T., Dong, L., Lin, K., & Li, S.-H. (2019). Standing genetic variation as the predominant
source for adaptation of a songbird. *Proceedings of the National Academy of Sciences,*
116(6), 2152–2157. https://doi.org/10.1073/pnas.1813597116

Lassauce, H., Dudgeon, C. L., Armstrong, A. J., Wantiez, L., & Carroll, E. L. (2022). Evidence of

fine-scale genetic structure for reef manta rays Mobula alfredi in New Caledonia.

Endangered Species Research, *47*, 249–264. https://doi.org/10.3354/esr01178

Lawson, J. M., & Fordham, S. V. (2018). *Sharks ahead: Realizing the potential of the convention*

on migratory species to conserve elasmobranchs (Shark Advocates International, The

Ocean Foundation, pp. 1–76) [Report]. Shark Advocates International, The Ocean

Foundation. https://www.bmis-bycatch.org/references/c8zv4pcf

Lawson, J. M., Fordham, S. V., O'Malley, M. P., Davidson, L. N. K., Walls, R. H. L., Heupel, M.

R., Stevens, G., Fernando, D., Budziak, A., Simpfendorfer, C. A., Ender, I., Francis, M.

P., Sciara, G. N. di, & Dulvy, N. K. (2017). Sympathy for the devil: A conservation

strategy for devil and manta rays. *PeerJ*, *5*, e3027. https://doi.org/10.7717/peerj.3027

Le Corre, V., & Kremer, A. (1998). Cumulative effects of founding events during colonisation on

genetic diversity and differentiation in an island and stepping-stone model. *Journal of*

Evolutionary Biology, *11*(4), 495–512. https://doi.org/10.1046/j.1420-

9101.1998.11040495.x

Legendre, P., Fortin, M.-J., & Borcard, D. (2015). Should the Mantel test be used in spatial

analysis? *Methods in Ecology and Evolution*, *6*(11), 1239–1247.

https://doi.org/10.1111/2041-210X.12425

Leslie, M. S., & Morin, P. A. (2018). Structure and phylogeography of two tropical predators,

spinner (*Stenella longirostris*) and pantropical spotted (*S. attenuata*) dolphins, from SNP

data. *Royal Society Open Science*, *5*(4), 171615. https://doi.org/10.1098/rsos.171615

Liu, X., Rønhøj Schjøtt, S., Granquist, S. M., Rosing-Asvid, A., Dietz, R., Teilmann, J., Galatius,

A., Cammen, K., O'Corry-Crowe, G., Harding, K., Härkönen, T., Hall, A., Carroll, E. L.,

Kobayashi, Y., Hammill, M., Stenson, G., Kirstine Frie, A., Lydersen, C., Kovacs, K. M.,

… Tange Olsen, M. (2022). Origin and expansion of the world's most widespread

pinniped: Range-wide population genomics of the harbour seal (Phoca vitulina).

Molecular Ecology, *31*(6), 1682–1699. https://doi.org/10.1111/mec.16365

López, K. E. R., Guadalupe, J. J., Gordillo-Romero, M., Montero-Oleas, A., Pazmiño, D. A.,

Guerrero, M., & Torres, M. de L. (2022). Exploring the genetic diversity and population

structure of *Mobula birostris* in two key aggregation zones in the Eastern Tropical Pacific.

Marine Ecology Progress Series, *699*, 75–89. https://doi.org/10.3354/meps14171

Louis, M., Galimberti, M., Archer, F., Berrow, S., Brownlow, A., Fallon, R., Nykänen, M., O'Brien,

J., Roberston, K. M., Rosel, P. E., Simon-Bouhet, B., Wegmann, D., Fontaine, M. C.,

Foote, A. D., & Gaggiotti, O. E. (2021). Selection on ancestral genetic variation fuels

repeated ecotype formation in bottlenose dolphins. *Science Advances*, *7*(44), eabg1245.

https://doi.org/10.1126/sciadv.abg1245

Lowe, W. H., & Allendorf, F. W. (2010). What can genetics tell us about population connectivity?

Molecular Ecology, *19*(15), 3038–3051. https://doi.org/10.1111/j.1365-

294X.2010.04688.x

McClain, M. A., Hammerschlag, N., Gallagher, A. J., Drymon, J. M., Grubbs, R. D., Guttridge, T.

L., Smukall, M. J., Frazier, B. S., & Daly-Engel, T. S. (2022). Age-dependent dispersal

and relatedness in tiger sharks (*Galeocerdo cuvier*). *Frontiers in Marine Science*, *9*.

https://www.frontiersin.org/articles/10.3389/fmars.2022.900107

Milanesi, M., Capomaccio, S., Vajana, E., Bomba, L., Garcia, J. F., Ajmone-Marsan, P., & Colli,

L. (2017). *BITE: An R package for biodiversity analyses* (p. 181610). bioRxiv.

https://doi.org/10.1101/181610

Mills, L. S., & Allendorf, F. W. (1996). The one-migrant-per-generation rule in conservation and

management. *Conservation Biology*, *10*(6), 1509–1518.

Moazzam, M. (2018). *Unprecedented decline in the catches of mobulids: An important*

component of tuna gillnet fisheries of the Northern Arabian Sea (IOTC - 14th Working

Party on Ecosystems and Bycatch). IOTC-2018-WPEB14-30.

Mussmann, S. M., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2019). BA3-SNPs:

Contemporary migration reconfigured in BayesAss for next-generation sequence data.

Methods in Ecology and Evolution, *10*(10), 1808–1813. https://doi.org/10.1111/2041-

210X.13252

Nikolic, N., Devloo-Delva, F., Bailleul, D., Noskova, E., Rougeux, C., Delord, C., Borsa, P.,

Liautard-Haag, C., Hassan, M., Marie, A. D., Feutry, P., Grewe, P., Davies, C., Farley, J.,

Fernando, D., Biton-Porsmoguer, S., Poisson, F., Parker, D., Leone, A., … Arnaud-

Haond, S. (2023). Stepping up to genome scan allows stock differentiation in the

worldwide distributed blue shark *Prionace glauca*. *Molecular Ecology*, *32*(5), 1000–1019.

https://doi.org/10.1111/mec.16822

O'Malley, M., Townsend, K. A., Hilton, P., Heinrichs, S., & Stewart, J. D. (2017). Characterization

of the trade in manta and devil ray gill plates in China and South-east Asia through trader

surveys. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *27*(2), 394–413.

https://doi.org/10.1002/aqc.2670

Palaiokostas, C., Bekaert, M., Khan, M. G., Taggart, J. B., Gharbi, K., McAndrew, B. J., &

Penman, D. J. (2015). A novel sex-determining QTL in Nile tilapia (Oreochromis

niloticus). *BMC Genomics*, *16*(1), 171. https://doi.org/10.1186/s12864-015-1383-x

Palsbøll, P. J., Bérubé, M., & Allendorf, F. W. (2007). Identification of management units using

population genetic data. *Trends in Ecology & Evolution*, *22*(1), 11–16.

https://doi.org/10.1016/j.tree.2006.09.003

Palumbi, S. R. (2003). Population genetics, demographic connectivity, and the design of marine

reserves. *Ecological Applications*, *13*(sp1), 146–158. https://doi.org/10.1890/1051-

0761(2003)013[0146:PGDCAT]2.0.CO;2

Pante, E., & Simon-Bouhet, B. (2013). marmap: A package for importing, plotting and analyzing

bathymetric and topographic data in R. *PLOS ONE*, *8*(9), e73051.

https://doi.org/10.1371/journal.pone.0073051

Papastamatiou, Y. P., Meyer, C. G., Carvalho, F., Dale, J. J., Hutchinson, M. R., & Holland, K. N.

(2013). Telemetry and random-walk models reveal complex patterns of partial migration

in a large marine predator. *Ecology*, *94*(11), 2595–2606. https://doi.org/10.1890/12-

2014.1

Pascual, M., Rives, B., Schunter, C., & Macpherson, E. (2017). Impact of life history traits on

gene flow: A multispecies systematic review across oceanographic barriers in the

Mediterranean Sea. *PLOS ONE*, *12*(5), e0176419.

https://doi.org/10.1371/journal.pone.0176419

Pate, J. H., & Marshall, A. D. (2020). Urban manta rays: Potential manta ray nursery habitat

along a highly developed Florida coastline. *Endangered Species Research*, *43*, 51–64.

https://doi.org/10.3354/esr01054

Peel, L. R., Stevens, G. M. W., Daly, R., Daly, C. A. K., Lea, J. S. E., Clarke, C. R., Collin, S. P.,

& Meekan, M. G. (2019). Movement and residency patterns of reef manta rays *Mobula*

alfredi in the Amirante Islands, Seychelles. *Marine Ecology Progress Series*, *621*, 169–

184. https://doi.org/10.3354/meps12995

Perryman, R. J. Y., Mourier, J., Venables, S. K., Tapilatu, R. F., Setyawan, E., & Brown, C.

(2022). Reef manta ray social dynamics depend on individual differences in behaviour.

Animal Behaviour, *191*, 43–55. https://doi.org/10.1016/j.anbehav.2022.06.010

Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double Digest

RADseq: An inexpensive method for de novo snp discovery and genotyping in model and

non-model species. *PLoS ONE*, *7*(5), e37135.

Phillips, N. M., Devloo-Delva, F., McCall, C., & Daly-Engel, T. S. (2021). Reviewing the genetic

evidence for sex-biased dispersal in elasmobranchs. *Reviews in Fish Biology and*

Fisheries, *31*(4), 821–841. https://doi.org/10.1007/s11160-021-09673-9

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using

multilocus genotype data. *Genetics*, *155*(2), 945–959.

Puechmaille, S. J. (2016). The program structure does not reliably recover the correct population

structure when sampling is uneven: Subsampling and new estimators alleviate the

problem. *Molecular Ecology Resources*, *16*(3), 608–627.

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J.,

Sklar, P., de Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: a tool set for

whole-genome association and population-based linkage analyses. *The American*

Journal of Human Genetics, *81*(3), 559–575.

Rambahiniarison, J., Agustines, A., Alexopoulos, K., Araujo, G., Armstrong, A. O., Arnold, S.,

Barruga, A., Cañete, T., Conales Jr., S., Delijero, K., Enolva, N. P., Flam, A. L., Keane,

E., Labaja, J., Legaspi, C. G., Murie, C., Murray, R., Oliver, S. P., Pierce, S. J., … Barr,

Y. (2023). Distribution of the reef manta ray *Mobula alfredi* and the oceanic manta ray

Mobula birostris in the Philippines: A collaborative effort for conservation. *Journal of Fish Biology*, *102*(2), 492–503. https://doi.org/10.1111/jfb.15283

Reed, D. H. (2004). Extinction risk in fragmented habitats. *Animal Conservation Forum*, *7*(2), 181–191. https://doi.org/10.1017/S1367943004001313

Rochette, N. C., & Catchen, J. M. (2017). Deriving genotypes from RAD-seq short-read data using Stacks. *Nature Protocols*, *12*(12), Article 12. https://doi.org/10.1038/nprot.2017.123

Rohner, C. A., Flam, A. L., Pierce, S. J., & Marshall, A. D. (2017). Steep declines in sightings of manta rays and devilrays (Mobulidae) in southern Mozambique. *PeerJ Preprints*, *5*(e3051v1). https://doi.org/10.7287/peerj.preprints.3051v1

Rohner, C. A., Pierce, S. J., Marshall, A. D., Weeks, S. J., Bennett, M. B., & Richardson, A. J. (2013). Trends in sightings and environmental influences on a coastal aggregation of manta rays and whale sharks. *Marine Ecology Progress Series*, *482*, 153–168. https://doi.org/10.3354/meps10290

Setyawan, E., Sianipar, A. B., Erdmann, M. V., Fischer, A. M., Haddy, J. A., Beale, C. S., Lewis, S. A., & Mambrasar, R. (2018). Site fidelity and movement patterns of reef manta rays (*Mobula alfredi*: Mobulidae) using passive acoustic telemetry in northern Raja Ampat, Indonesia. *Nature Conservation Research*, *3*(4). https://doi.org/10.24189/ncr.2018.043

Stevens, G., Fernando, D., & Sciara, G. N. D. (2018). *Guide to the Manta and Devil Rays of the World*. Princeton University Press.

Stewart, J. D., Beale, C. S., Fernando, D., Sianipar, A. B., Burton, R. S., Semmens, B. X., & Aburto-Oropeza, O. (2016). Spatial ecology and conservation of *Manta birostris* in the Indo-Pacific. *Biological Conservation*, *200*, 178–183. https://doi.org/10.1016/j.biocon.2016.05.016

Stewart, J. D., Jaine, F. R. A., Armstrong, A. J., Armstrong, A. O., Bennett, M. B., Burgess, K. B., Couturier, L. I. E., Croll, D. A., Cronin, M. R., Deakos, M. H., Dudgeon, C. L., Fernando, D., Froman, N., Germanov, E. S., Hall, M. A., Hinojosa-Alvarez, S., Hosegood, J. E., Kashiwagi, T., Laglbauer, B. J. L., … Stevens, G. M. W. (2018). Research priorities to support effective manta and devil ray conservation. *Frontiers in Marine Science*, *5*. https://doi.org/10.3389/fmars.2018.00314

Stewart, J. D., Nuttall, M., Hickerson, E. L., & Johnston, M. A. (2018). Important juvenile manta

ray habitat at Flower Garden Banks National Marine Sanctuary in the northwestern Gulf

of Mexico. *Marine Biology*, *165*(7), 111. https://doi.org/10.1007/s00227-018-3364-5

Stoffel, M. A., Esser, M., Kardos, M., Humble, E., Nichols, H., David, P., & Hoffman, J. I. (2016).

InbreedR: an R package for the analysis of inbreeding based on genetic markers.

Methods in Ecology and Evolution, *7*(11), 1331–1339.

Taylor, B. L., & Dizon, A. E. (1996). The need to estimate power to link genetics and

demography for conservation. *Conservation Biology*, *10*(2), 661–664.

Thia, J. A. (2023). Guidelines for standardizing the application of discriminant analysis of

principal components to genotype data. *Molecular Ecology Resources*, *23*(3), 523–538.

https://doi.org/10.1111/1755-0998.13706

Thorburn, J., Neat, F., Burrett, I., Henry, L.-A., Bailey, D. M., Jones, C. S., & Noble, L. R. (2019).

Ontogenetic variation in movements and depth use, and evidence of partial migration in a

benthopelagic elasmobranch. *Frontiers in Ecology and Evolution*, *7*.

https://www.frontiersin.org/articles/10.3389/fevo.2019.00353

Venables, S. K., Marshall, A. D., Armstrong, A. J., Tomkins, J. L., & Kennington, W. J. (2021).

Genome-wide SNPs detect no evidence of genetic population structure for reef manta

rays (Mobula alfredi) in southern Mozambique. *Heredity*, *126*(2), Article 2.

https://doi.org/10.1038/s41437-020-00373-x

Vignaud, T. M., Maynard, J. A., Leblois, R., Meekan, M. G., Vázquez-Juárez, R., Ramírez-

Macías, D., Pierce, S. J., Rowat, D., Berumen, M. L., Beeravolu, C., Baksay, S., &

Planes, S. (2014). Genetic structure of populations of whale sharks among ocean basins

and evidence for their historic rise and recent decline. *Molecular Ecology*, *23*(10), 2590–

2601. https://doi.org/10.1111/mec.12754

Waples, R. (1998). Separating the wheat from the chaff: Patterns of genetic differentiation in high

gene flow species. *Journal of Heredity*, *89*(5), 438–450.

https://doi.org/10.1093/jhered/89.5.438

Waples, R., & Gaggiotti, O. (2006). INVITED REVIEW: What is a population? An empirical

evaluation of some genetic methods for identifying the number of gene pools and their

degree of connectivity. *Molecular Ecology*, *15*(6), 1419–1439.

https://doi.org/10.1111/j.1365-294X.2006.02890.x

Waples, R. K., Albrechtsen, A., & Moltke, I. (2019). Allele frequency-free inference of close

familial relationships from genotypes or low-depth sequencing data. *Molecular Ecology*,

28(1), 35–48. https://doi.org/10.1111/mec.14954

Waples, R., Punt, A. E., & Cope, J. M. (2008). Integrating genetic data into management of

marine resources: How can we do it better? *Fish and Fisheries*, *9*(4), 423–449.

https://doi.org/10.1111/j.1467-2979.2008.00303.x

Ward, R. D., Woodwark, M., & Skibinski, D. O. F. (1994). A comparison of genetic diversity levels

in marine, freshwater, and anadromous fishes. *Journal of Fish Biology*, *44*(2), 213–232.

https://doi.org/10.1111/j.1095-8649.1994.tb01200.x

Ward-Paige, C. A., Davis, B., & Worm, B. (2013). Global population trends and human use

patterns of manta and mobula rays. *PLOS ONE*, *8*(9), e74835.

https://doi.org/10.1371/journal.pone.0074835

Weir, B. S., & Cockerham, C. C. (1984). Estimating f-statistics for the analysis of population

structure. *Evolution*, *38*(6), 1358–1370. https://doi.org/10.2307/2408641

Whitney, J. L., Coleman, R. R., & Deakos, M. H. (2023). Genomic evidence indicates small

island-resident populations and sex-biased behaviors of Hawaiian reef Manta Rays. *BMC*

Ecology and Evolution, *23*(1), 31. https://doi.org/10.1186/s12862-023-02130-0

Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, *16*(2), 97–159.

Younger, J. L., Clucas, G. V., Kao, D., Rogers, A. D., Gharbi, K., Hart, T., & Miller, K. J. (2017).

The challenges of detecting subtle population structure and its importance for the

conservation of emperor penguins. *Molecular Ecology*, *26*(15), 3883–3897.

912