Comparative population genomics of manta rays has global implications for management

Emily Humble*[1,2], Jane Hosegood*[2,3], Gary Carvalho[3], Mark de Bruyn[3,4], Simon Creer[3], Guy M W Stevens[2], Amelia Armstrong[5], Ramon Bonfil[6,7,8], Mark Deakos[9], Daniel Fernando[2,10], Niv Froman[2], Lauren R Peel[2,11,12], Stephen Pollett[2], Alessandro Ponzo[13], Joshua D Stewart[2,14], Sabine Wintner[15,16] & Rob Ogden[1]

[1]Royal (Dick) School of Veterinary Studies and the Roslin Institute, University of Edinburgh, EH25 9RG, Edinburgh, UK

[2]The Manta Trust, Catemwood House, Norwood Lane, Corscombe, Dorset, DT2 0NT, UK

[3]Molecular Ecology and Evolution Group, Bangor University, Bangor, UK

[4]Australian Research Centre for Human Evolution, Griffith University, Nathan, QLD, Australia

[5]School of Biomedical Sciences, The University of Queensland, St Lucia, QLD, Australia

[6]Océanos Vivientes AC, Mexico City, Mexico

[7]Consejo Nacional de Humanidades Ciencia y Tecnología (CONAHCyT), Mexico City, Mexico

[8]El Colegio de la Frontera Sur, Unidad Chetumal, Chetumal, Q. Roo, Mexico.

[9]Hawai'i Association for Marine Education and Research, Lahaina, HI, USA

[10]Blue Resources Trust, Colombo, Sri Lanka

[11]Save Our Seas Foundation – D'Arros Research Centre, Geneva, Switzerland

[12]School of Biological Sciences, Oceans Institute and Oceans Graduate School, The University of Western Australia, Crawley, WA, Australia

[13]Large Marine Vertebrates Research Institute Philippines, Jagna, Philippines

[14]Ocean Ecology Lab, Marine Mammal Institute, Department of Fisheries, Wildlife & Conservation Sciences, Oregon State University, Newport, OR, United States

[15]KwaZulu-Natal Sharks Board, Umhlanga Rocks, South Africa

[16]University of KwaZulu-Natal, School of Life Sciences, Durban, South Africa

*Joint first authors

Corresponding author: Emily Humble, Royal (Dick) School of Veterinary Studies and the Roslin Institute, University of Edinburgh, EH25 9RG, UK. Email: emily.humble[at]ed.ac.uk

Abstract

Understanding population connectivity and genetic diversity is of fundamental importance to
conservation. However, in globally threatened marine megafauna, challenges remain due to
their elusive nature and wide-ranging distributions. As overexploitation continues to threaten
biodiversity across the globe, such knowledge gaps compromise both the suitability and
effectiveness of management actions. Here, we use a comparative framework to investigate
genetic differentiation and diversity of manta rays, one of the most iconic yet vulnerable groups
of elasmobranchs on the planet. Despite their recent divergence, we show how oceanic manta
rays *(Mobula birostris)* display significantly higher genetic diversity than reef manta rays
(Mobula alfredi) and that *M. birostris* populations display higher connectivity worldwide.
Through reconstructing modes of colonisation, we reveal how both contemporary and
historical forces have likely influenced these patterns, with important implications for
population management. Our findings highlight the potential for fisheries to disrupt population
dynamics at both local and global scales and therefore have direct relevance for international
marine conservation.

Teaser

Population genomics of manta rays reveals striking differences in differentiation and diversity
between two recently diverged species.

Main text

Introduction

Understanding the extent to which populations are connected is key to exploring population dynamics, predicting extinction risk and informing conservation management (*1–3*). In species with isolated populations characterised by limited dispersal, the risk of extirpation from local depletion is high (*4*). In such cases, local and regional scale management will be most appropriate for preventing and reversing population declines (*5*). In contrast, species with high rates of gene flow are potentially demographically and genetically more resilient to extrinsic factors (*3, 6*). However, in order to maintain connectivity and mitigate genetic diversity loss in these taxa, management measures must be coordinated and encompass migratory corridors. As overexploitation and habitat destruction threaten to disrupt population dynamics at a global scale, characterising genetic variation and connectivity has become more important than ever before (*7–9*).

In widely distributed marine species with high dispersal potential, genetic differentiation is often found to be subtle or non-existent (*5, 10, 11*). Such patterns can arise from a range of mechanisms – from high contemporary gene flow through to recent divergence of historically large populations (*5, 12, 13*) – and can therefore be difficult to interpret. The latter scenario reflects a disconnect between demographic and genetic connectivity and has important implications for species resilience (*3, 11, 14*). This is because populations that appear genetically connected may not operate as single demographic units, making them more vulnerable to overexploitation. High-resolution SNP datasets go some way to addressing this problem by providing greater power to detect subtle differences at both neutral and adaptive loci (*15, 16*). However, since population genetic differentiation can be affected by past, as well as contemporary patterns, parallel inference of historical relationships and genetic diversity can allow the relative contribution of historical processes to be explicitly evaluated (*17–19*). Furthermore, when carried out within a comparative framework, such an approach can provide powerful insights into the drivers of population divergence and therefore improve recommendations for conservation management (*20*).

Manta rays are large, mobile elasmobranchs inhabiting tropical and sub-tropical oceans (*21*) (Figure 1A, C) and provide an excellent opportunity to evaluate the genomic consequences of historical and contemporary population processes within a comparative framework. They comprise two described species estimated to have diverged less than 0.5 Mya as a result of

distinct habitat preferences (*22*). The reef manta ray (*Mobula alfredi)* frequents near-shore

tropical reef environments, such as coral atolls and barrier reefs (Kashiwagi *et al.* 2011), with

a high degree of residency (Deakos *et al.* 2011; Jaine *et al.* 2014; Braun *et al.* 2015; Setyawan

et al. 2018; Peel *et al.* 2019; Knochel *et al.* 2022b; Germanov *et al.* 2022). In contrast, the

oceanic manta ray (*Mobula birostris)* can also be found ranging into sub-tropical habitats along

continental coastlines and at oceanic islands, often adjacent to productive deep-water

upwellings areas (Kashiwagi *et al.* 2011; Andrzejaczek *et al.* 2021). As a result of these

differences in habitat use, *M. alfredi* and *M. birostris* have long been considered to display

marked differences in their migratory abilities and levels of gene flow. Yet, only a handful of

long-distance movements have ever been recorded in *M. birostris* (*23*, *24*) alongside

observations of site-fidelity (*25–27*), raising questions about the extent to which population

structure and genetic diversity may differ across species. To date, assessments of genetic

differentiation in *M. alfredi* have focussed on local and regional patterns (*28–30*) and we have

little understanding of how genetic diversity is distributed across the species' range. In, *M.*

birostris, the situation is even less clear, with studies reporting both widespread connectivity

and population differentiation (*31–33*). Critically, these differences and uncertainties exist

against a background of ongoing global exploitation and uncertain implications for

management.

Targeted and incidental fisheries, driven in part by increasing demand for mobulid gill plates

(*21*, *34*), have led to widespread population declines in manta rays (*35–39*). Currently, both

species are managed through a patchwork of local, regional and international measures with

varying levels of implementation and enforcement (*40–42*). To determine the appropriateness

of management measures and assess population vulnerability, a global assessment of

management units is urgently required (*40*, *43*). Here, we undertake a comparative genomic

analysis of manta ray populations from across their global distribution to investigate

connectivity, genetic diversity, and historical relationships with an aim to guide effective

fisheries management.

Results

We generated double digest restriction-site associated DNA (ddRAD) sequence data for 173

individuals from 12 locations to represent the global distribution of each species (Figure 1A,

C). For *M. alfredi* (*n* = 91), these comprised the Chagos Archipelago (n = 5), the Maldives (n

= 36), Seychelles (n = 22), Australia Pacific (n = 3), Fiji (n = 8) and Hawaii (n = 17). For *M.*

birostris (*n* = 82), these comprised Sri Lanka (n = 37), the Philippines (n = 22), South Africa (n

= 3), Mexico Caribbean (n = 4), Mexico Pacific (n = 9) and Peru (n = 7). Quality controlled

sequencing reads were *de novo* assembled using the STACKS v2.64 pipeline and a total of

15,312 high-quality SNPs were called across both species. See Materials and Methods and

Supplementary Information for details.

116   Contrasting patterns of population structure at a global scale

To investigate population differentiation within each species we used four complementary

approaches: discriminant analysis of principal components (DAPC), admixture, pairwise F_{ST}

and isolation by distance analysis. In *M. alfredi*, all methods supported the presence of strong

population structure at both global and regional scales. Populations inhabiting different ocean

basins displayed the highest degree of differentiation in the DAPC, with Pacific and Indian

Ocean populations forming distinct clusters along PC1 (Figure 1B). Regional differentiation

was also detected, with Seychelles, Chagos and the Maldives clustering apart along PC2, and

Hawaii separating from Australia Pacific and Fiji along PC3 (Figure 1B and Figure S1A). These

patterns were reinforced in the admixture analysis which highlighted two major ancestral

source populations, inferred an optimal value of $K = 4$ and resolved hierarchical structure up

to $K = 7$ (Figure S2 and Figure S3A). Pairwise F_{ST} estimates between ocean basins were on

average over two times higher than those within (mean pairwise F_{ST} between ocean basins =

0.30, mean pairwise F_{ST} within ocean basins = 0.13, Figure 2A) yet all population comparisons

were found to be significant (Figure S4A, mean = 0.23, min = 0.08, max = 0.43). Finally, we

detected a significant relationship between pairwise F_{ST} and geographic distance (Mantel's *r*

= 0.84, *P* = 0.02) indicating a strong effect of isolation by distance (Figure 2B).

Figure 1. Contrasting patterns of population structure in manta rays. (A, C) Geographic distributions of (A) *Mobula alfredi* and (C) *Mobula birostris* visualised together with the locations of samples used in this study. Dark shaded distributions denote the confirmed species range and light shaded distributions denote the expected species range. Sampling location points are distinguished by colour and scaled by the number of samples. Further details are provided in the Supplementary Material. (B, D) Scatterplots showing individual variation in principal components (PC) one and two derived from discriminant analysis of principal components analysis for (B) *M. alfredi* and (D) *M. birostris* individuals. The amount of variance explained by each PC is shown in parentheses. Population abbreviations: AP = Australia Pacific, CHAG = Chagos, FIJI = Fiji, HAW = Hawaii, MAL = Maldives, SEY = Seychelles, MC = Mexico Caribbean, MP = Mexico Pacific, PERU = Peru, SA = South Africa, SL = Sri Lanka and PHI = the Philippines.

In stark contrast, *M. birostris* displayed little evidence for strong population structure across all methods. Individuals from different ocean basins clustered closely together along each axis in the DAPC (Figure 1D and Figure S1B). Admixture identified $K = 1$ as the optimal number of clusters, with increasing values of K merely introducing additional mixing (Figure S2 and Figure S3B). Pairwise F_{ST} estimates were two-fold lower than in *M. alfredi*, with no pairwise comparison falling above 0.007 (mean = 0.002, min = -0.005, max = 0.006, Figure 2C). Nevertheless, despite these broad patterns, several lines of evidence indicate the presence

of subtle geographic differentiation in this species. First, individuals from Mexico Pacific, Peru, and Mexico Caribbean clustered apart from those sampled in South Africa, Sri Lanka, and the Philippines along PC1 (Figure 1D). Second, despite pairwise F_{ST} estimates being low, comparisons between Eastern-Pacific and Indo-Pacific populations, and between Sri Lanka and the Philippines were statistically significant (Figure S4B). Small F_{ST} values are expected when minor allele frequencies are low and therefore do not necessarily reflect an absence of differentiation (*44*). Finally, while no significant relationship was observed between pairwise F_{ST} and geographic distance (Mantel's $r = 0.45$, $P = 0.10$), there was a tendency for populations separated by greater distances to display higher differentiation (Figure 2D).

Figure 2. Population genetic differentiation, isolation by distance and contemporary migration in manta rays. (A, C) Pairwise F_{ST} estimates between sampling locations for (A) *M. alfredi* and (C) *M. birostris*. Samples from Australia Pacific and South Africa were excluded from this analysis due to low sample sizes. (B, D) Relationship between genetic (F_{ST}) and geographic distance as calculated by least-cost path analysis for all pairwise population comparisons in (B) *M. alfredi* and (D) *M. birostris*. Solid lines and shaded areas reflect the regression slopes and standard errors respectively. (E) Contemporary gene flow estimates between populations of *M. alfredi*. The direction of each arrow represents the direction of gene flow, and the width of each ribbon reflects the relative amount of gene flow.

Contemporary gene flow

To characterise the strength and direction of gene flow between populations we used the program BA3-SNPs (*45*) to estimate recent migration. As this method assumes low levels of connectivity and imposes an upper-bound on the proportion of non-migrants in a population,

we restricted the analysis to *M. alfredi* (see Materials and Methods for details). As expected,

contemporary gene flow was low (Figure 2E); the average migration rate between populations,

measured as the estimated number of migrants per generation, was 0.029 (min = 0.008, max

= 0.15), with this figure falling to 0.018 (min = 0.008 , max = 0.03) when considering gene flow

between populations in different ocean basins. Migration into both Hawaii and the Maldives

was lowest, indicating these populations are the most isolated of those sampled (Table S1).

Migration rates were only deemed significant between Seychelles and Chagos (0.15) and

between Fiji and Australia Pacific (0.15), in line with these populations being last to separate

in the admixture analysis. These patterns highlight that while *M. alfredi* have the propensity to

travel over large distances, restricted movement likely dominates.

Historical relationships among populations

To place patterns of genetic differentiation into a historical context, we investigated population

origins and colonisation patterns using TreeMix (*46*). This program uses allele frequency data

to infer patterns of population splits and admixture events through the construction of a

maximum likelihood tree. In *M. alfredi*, internal branch lengths were relatively long, with an

initial split clearly separating populations in the Indian and Pacific Oceans (Figure 3A). The

Maldives and Australia Pacific were the first to separate within each locality and displayed the

lowest levels of genetic drift overall. Hawaii was among the last populations to split and

displayed the highest amount of drift, in line with its geographic isolation. The best supported

model inferred two migration events (Figure S5A); one from the *M. alfredi* population in

Seychelles into *M. birostris*, and one from Chagos into Hawaii. However, because not all

geographic regions are represented in our data set, the true sources and sinks of these

admixture events may originate from related ghost populations. In contrast to *M. alfredi*, the

addition of migration events led to no substantial improvement in the model for *M. birostris*

(Figure S5B) and therefore the tree without migration is presented here. Interestingly, internal

branch lengths were considerably shorter in *M. birostris*, indicating rapid radiation from a

shared ancestral source population (Figure 3B). External branch lengths were also short,

consistent with larger populations displaying marginal drift and low divergence. Nevertheless,

despite these patterns, geographic signal could be detected in the *M. birostris* tree, with

populations from the Eastern Pacific and the Caribbean (Peru, Mexico Caribbean, and Mexico

Pacific) grouping separately from those in the Atlantic and Indo-Pacific (South Africa, the

Philippines, and Sri Lanka).

Figure 3. Historical relationships and genetic diversity in manta rays. (A–B) TreeMix maximum likelihood consensus tree displaying the historical relationships among (A) *M. alfredi* and (B) *M. birostris* populations. Horizontal branch lengths reflect the amount of genetic drift that has occurred along each branch. Bootstrap support values for each node are indicated. Migration edges inferred using TreeMix are represented as arrows and coloured according to their migration weight. The scale bar reflects 10 times the average standard error of the entries in the sample covariance matrix. (C–E) Variation in individual multi-locus heterozygosity between (C) species and among populations of (D) *M. alfredi* and (E) *M. birostris*. Centre lines of boxplots reflect the median, bounds of the boxes extend from the first to the third quartiles, and upper and lower whiskers reflect the largest and smallest values but no further than 1.5 * the interquartile range from the hinge.

Heterozygosity landscape across species and populations

To explore how patterns of population structure and colonisation are associated with genome-wide variation, we compared individual multi-locus heterozygosity between species and among populations. Strikingly, heterozygosity was on average over three times higher in *M. birostris* (mean = 0.10, min = 0.053, max = 0.12) than in *M. alfredi* (mean = 0.03, min = 0.01, max = 0.051), with every individual displaying a higher value than any *M. alfredi* (ß = 0.07, 95% CI = 0.06–0.07, *P* = <2.2 X 10^{-16}, Figure 3C). This finding is in line with the patterns of

population structure and historical splits we observed in each species. Variation in genetic
diversity was also observed at a population level (Figure 3D–E). In *M. alfredi*, the Maldives
and Australia Pacific had the highest levels of heterozygosity within each ocean basin, in line
with these populations being first to split in the TreeMix analysis. Indian Ocean populations
displayed higher overall levels of diversity than Pacific populations and while a weak negative
trend in variation was observed, mean values were overall similar. A steeper decline in
diversity was observed in the Pacific populations, with heterozygosity in Hawaiian individuals
being around half that of the Australian animals (Figure 3D), in line with this population being
last to split in the TreeMix analysis and displaying the highest amount of drift. In contrast, *M.*
birostris populations displayed less extreme variation in heterozygosity, with mean values
differing by less than 0.02 (Figure 3E). Furthermore, variance within populations was an order
of magnitude greater in *M. birostris* than in *M. alfredi*, and was particularly large in Sri Lanka,
the Philippines, and Mexico Pacific populations.

Discussion

Manta rays are iconic megafauna with cultural and socio-economic significance. Due to
targeted and bycatch fisheries operating across their broad-ranging distributions, populations
are declining worldwide. Elucidating levels of connectivity and genetic variation remains a
crucial priority for conservation management. We use reduced-representation sequencing on
a global set of samples and reveal striking differences in the population genetic landscape of
two recently diverged manta ray species. By considering the influence of both contemporary
and historical processes, our study takes a precautionary approach to assessing conservation
units, with important implications for management.

We first demonstrate the presence of strong genetic differentiation in *M. alfredi* at a global and
regional scale. From a total of six sampling locations, we found evidence for at least five
genetically distinct, and by extension, demographically independent populations. Two of these
were separated by a distance of ~1,200 km, which is close to the maximum recorded
movement in the species (*47, 48*), indicating that long distance migrations are likely rare.
Indeed, contemporary gene flow was low – especially between geographically distant
locations – with only a small proportion of individuals in any population being identified as first-
or second-generation migrants. Furthermore, when gene flow was observed, it tended to be
unidirectional. These results are in line with recent studies demonstrating population
differentiation between Western Australia and Mozambique (*28*) and between Eastern
Australia and New Caledonia (*29*), together highlighting how large ocean basins form

significant barriers to dispersal in coastal elasmobranchs (*49*). High site-fidelity has been
widely reported in *M. alfredi* based on tagging and mark-recapture studies (*47, 50–55*).
However, the degree of residency has been shown to vary, with movements rarely exceeding
a few hundred kilometres in some locations (*50–52, 54, 56*) yet reaching over 1,000 km in
others (*48, 57*). Our study presents a comparatively broad-scale analysis relevant for regional
and global management planning. Further work on local patterns of population structure will
shed light on the nuances and drivers of fine-scale movement patterns in this species (*30*).

To explore the mechanism by which manta rays colonised their distribution, we reconstructed
historical relationships and assessed levels of heterozygosity. In *M. alfredi*, we found strong
evidence for an initial split between the Indian and Pacific Oceans, followed by range
expansion within each. Population splits were associated with increasing genetic drift and
reduced heterozygosity, particularly in the Pacific, indicating that *M. alfredi* underwent a
stepping-stone pattern of colonisation involving opportunistic long-range movements and
associated founder events. This is consistent with a recent observation of a pregnant *M. alfredi*
individual at Cocos Island, Costa Rica (*58*), almost 6,000 km east of the nearest confirmed
sighting, and the first record of *M. alfredi* in the Eastern Pacific. Range expansion inherently
impacts genetic diversity, with a stepping-stone model of colonisation predicted to result in the
strongest cumulative effect of founder events (*59*). Among our sampled populations, Hawaii
is the most geographically isolated, situated at the edge of the *M. alfredi* distribution.
Interestingly, not only was Hawaii the most genetically differentiated from all populations in
our study, but it displayed the longest external branch lengths in the TreeMix analysis and the
lowest levels of heterozygosity. Genetic variation is fundamental for enabling populations to
adapt in response to selection (*9, 60, 61*). Our findings therefore expose how isolated *M.
alfredi* populations at the periphery of their distribution may be intrinsically more vulnerable to
changing environmental conditions and the genetic impacts of population decline.

In stark contrast to the patterns observed in *M. alfredi*, *M. birostris* displayed markedly higher
levels of heterozygosity and with only subtle genetic differentiation across ocean basins. Weak
population structure is common in highly mobile marine species (*62–64*), yet warrants careful
interpretation, particularly considering management recommendations (*65*). On the one hand,
these findings may be an indication of high contemporary gene flow and low natal philopatry,
in line with the species' occurrence at remote oceanic islands, tendency to range into sub-
tropical habitats and lower overall re-sight rates than *M. alfredi* (*66–68*). To date, our
understanding of the movement behaviour in *M. birostris* has largely been based on coastal

aggregations of adult individuals over relatively short timeframes (*32, 67, 69, 70*). Such studies

have a tendency to capture seasonal migrations as opposed to dispersal events and may

explain why very few long-distance (~1,000 km) movements have been recorded in the

species (*23, 24*). Indeed, with only a few migrants per generation required to obscure strong

population structure when N_e is large (*71*), it is possible that the patterns we observe translate

to infrequent dispersal events. Furthermore, dispersal could be segregated by age and/or sex

(*72, 73*), and may vary among individuals (*74–76*). While challenging, there is benefit in

extending future tagging efforts to transient individuals away from known aggregation sites

(*27*), as well as previously underrepresented age classes – such as juveniles – to capture

what may be infrequent yet evolutionarily relevant movements.

An alternative explanation for the patterns we observe in *M. birostris* is that insufficient time

has elapsed to reliably identify recent genetic divergence among localities. In contrast to *M.*

alfredi, our TreeMix analysis indicated that *M. birostris* rapidly radiated from a large ancestral

source, with only marginal genetic drift occurring between regions. This was further evidenced

by substantially higher levels of genetic variation that differed little across sampling locations.

In addition, little differentiation was observed between Mexico Pacific and Mexico Caribbean,

two regions that have been geographically separated since the emergence of the Isthmus of

Panama. These findings are consistent with a recent mark-resight analysis that estimated the

population of *M. birostris* in coastal Ecuador to number at least 22,000 individuals (*67*). Large

effective population sizes and high genetic variation increase the time taken for populations

to diverge due to genetic drift (*14, 71, 77*). This is further compounded in species with long

and overlapping generations (*78*) as is the case for manta rays (*79*). Taken together, genetic

similarities among *M. birostris* localities may be partially confounded by recent shared

ancestry and large effective population size.

On the basis of these considerations, we propose that a combination of large historical

population size and contemporary gene flow have contributed to the comparatively high levels

of diversity and genetic homogeneity in *M. birostris*. The subtle population differentiation we

observe between the Indian Ocean, South-East Asia and the Eastern Pacific is likely best

explained by the geographic limits of dispersal as opposed to complete geographic isolation.

Yet, unlike in *M. alfredi* where genetic clusters almost certainly reflect discrete demographic

units relevant for conservation management, the extent to which genetic connectivity in *M.*

birostris reflects demographic connectivity is less clear. For example, in extreme cases, the

number of migrants required to eliminate signals of population structure will not be enough to

demographically link populations, and more importantly, replenish those that have been
depleted (*11*). Interestingly, while re-sight rates are typically lower in *M. birostris* than *M.*
alfredi, demographic independence has been implicated in several mark-recapture studies
where re-sightings follow predictable patterns (*26*, *70*). Furthermore, a population genetic
analysis based on F_{ST} outliers uncovered allele frequency differences between two Mexican
locations and Sri Lanka (*32*), suggesting recent divergence against a background of ongoing
gene flow. Taken together, we highlight the potential for further work investigating adaptive
divergence between *M. birostris* populations and emphasise the need to combine molecular
measures of connectivity with empirical demographic data in this species (*3*, *65*, *80*).

Conservation implications
The remarkable differences we observe in the population genetics of manta rays directly
inform likely response to continued exploitation and respective conservation measures. At
present, *M. alfredi* is among the most protected mobulid species worldwide, with some
management frameworks in place at local, national, and international levels (*40*, *81*). Our
findings of global population structure underline how local initiatives recognising populations
as distinct management units will be most appropriate for this species. However, we also
demonstrate the consequence of geographic isolation on genetic variation and reveal how *M.*
alfredi likely faces a greater risk from local depletion. This is especially true for populations at
the edge of the species range and in regions with high coastal fishing pressure. Prioritising
these populations in conservation action plans and maintaining local connectivity will therefore
be crucial for boosting resilience and preventing local extinction in this vulnerable species.

The implications of our findings for *M. birostris* are more nuanced. Despite detecting only
subtle population genetic differentiation, we cannot rule out the possibility that historical
processes and large effective population size are obscuring a higher degree of contemporary
demographic separation. Together with studies reporting high site-fidelity and restricted
movement patterns, our findings strongly suggests that local and national management action
should be considered essential for protecting resident aggregations of *M. birostris*.
Nevertheless, we expect that weak population structure and high genetic variation are
simultaneously being driven by some degree of contemporary dispersal. Consequently, any
fishing activity taking place along migratory corridors threatens to disrupt a mode of gene flow
that is likely fundamental for long-term resilience of the species. Similarly, although we have
limited understanding of the number and distribution of breeding and nursery grounds (*24*, *82*,
83), significant reduction of local stocks may impact long-term recruitment at oceanic and even

global scales. We therefore emphasise the escalating need to improve the implementation of

regional and international measures that seek to protect taxa in the high seas. Together with

local scale management, appropriate evidence-based actions will contribute to maintaining

large, connected and genetically diverse populations of manta rays into the future.

Materials and Methods

Sample collection

Tissue samples were opportunistically collected from 12 geographic locations to represent the

global distribution of each species (Figure 1A, C). For *M. alfredi* (total *n* = 119), these

originated from the Chagos Archipelago (n = 5), the Maldives (n = 48), Seychelles (n = 23),

Australia Pacific (n = 4), Fiji (n = 9) and Hawaii (n = 30). For *M. birostris* (total *n* = 111), these

originated from Sri Lanka (n = 43), the Philippines (n = 36), South Africa (n = 3), Mexico

Caribbean (n = 4), Mexico Pacific (n = 13) and Peru (n = 12). Samples from Mexico Caribbean,

where a third putative manta ray species occurs in sympatry (*Mobula cf. birostris,* Hinojosa-

Alvarez *et al.* 2016; Hosegood *et al.* 2020), were visually and genetically confirmed as *M.*

birostris. For both species, samples were collected from a combination of live animals and

fisheries specimens (See Supplementary Files for further information).

DNA extraction and ddRAD sequencing

Genomic DNA was extracted using the Qiagen DNeasy Blood and Tissue Kit and quantified

using a Qubit 3.0 Broad Range Assay. Double digest restriction-site associated DNA (ddRAD)

libraries were prepared following the Peterson *et al.* (2012) protocol with modifications

described in Palaiokostas *et al.* (2015) and were 125 bp paired-end sequenced on an Illumina

HiSeq. Sequencing reads from both species were assessed for quality using FastQC and

processed together using the Stacks v2.54 pipeline (*87*). The three main assembly parameters

were chosen following the optimisation procedure outlined in Rochette and Catchen (2017)

(Figure S7). Initial quality filters were applied to the resulting genotypes before generating

three high-quality datasets for use in downstream analysis: two species-specific datasets; and

one dataset comprising both species. For the species-specific datasets, we extracted either

M. birostris or *M. alfredi* individuals, removed individuals with high relatedness coefficients

(*89, 90*) and filtered out SNPs with a minor allele count of less than 3, a genotyping rate less

than 90% and that were in linkage disequilibrium using PLINK. This left a total of 1,553 SNPs

in 91 *M. alfredi* individuals, and 6,278 SNPs in 82 *M. birostris* individuals. For the dataset

comprising both species, we first removed closely related individuals and then filtered out

SNPs with a minor allele count of less than 3 and a genotyping rate less than 90%. This left a

total of 15,312 SNPs called in 91 *M. alfredi* and 82 *M. birostris* individuals. See Supplementary

Material for further information on library preparation, read processing and SNP and individual

filtering.

Population structure

To investigate population structure we used the species-specific datasets and three
complementary approaches. First, we carried out a discriminant analysis of principal
components (DAPC) using the R package adegenet (*91*). This approach initially transforms
the SNP data using a principal components analysis (PCA) and then performs a discriminant
analysis on the retained PCs. This serves to maximise discrimination of individuals between
groups while minimising variation within (*92*). Following the recommendations outlined in Thia
(2023), the number of PCs retained as predictors was determined based on the *K*–*1* criterion,
where *K* is equal to the number of effective populations. For *M. alfredi*, this was set to 5, under
the assumption that each sample site reflects a separate population. For *M. birostris*, this was
set to 4 under the assumption that Mexico Pacific and Peru may represent a single population
given their close geographic proximity. Second, we estimated admixture proportions for the
individuals in each dataset using ADMIXTURE. Admixture runs were performed for ancestry
clusters ranging from $K = 1$–8, with 10 runs for each K. The optimal K was identified based on
the lowest cross-validation error. The runs with the highest likelihood were visualised. Third,
we estimated pairwise genetic differentiation between populations within each species using
the Weir and Cockerham F_{ST} value (*94*) calculated in the R package dartR (*95*). Confidence
intervals and *p*-values were estimated based on bootstrap resampling of individuals within
each population 1000 times. *Mobula alfredi* samples from Australia Pacific and *M. birostris*
samples from South Africa were excluded from this analysis due to low sample sizes.

Isolation by distance

To investigate patterns of isolation by distance, we examined the relationship between genetic
and geographic distance between all pairs of populations in each species. Genetic distances
were based on pairwise F_{ST} estimates calculated above. Geographic distances were
determined based on a least-cost path analysis implemented using the R package marmap
(*96*) with a minimum depth constraint of -10 metres in order to prevent paths overland. The
significance of associations between genetic and geographic distance matrices was inferred
using Mantel tests with the R package ade4 (*97*) .

Contemporary gene flow

To infer the strength and directionality of contemporary gene flow between populations we
used the program BA3-SNPs BayesAss v1.1 (*45*) which estimates the proportion of
immigrants in a given population using Bayesian inference. This analysis was restricted to *M.*
alfredi as it assumes low levels of connectivity and imposes an upper-bound on the proportion
of non-migrants in a population. We first performed initial runs of BayesAss to determine
optimal mixing parameters (dM = migration rate, dA = allele frequency and dF = inbreeding

coefficient) using the autotune function in BA3-SNPs. We than ran BayesAss-3 with
10,000,000 iterations, a burn-in of 1,000,000 and a sampling interval of 1000. Mixing
parameters were set to dM = 0.21, dA = 0.44 and dF = 0.08. Results were averaged across
five replicate runs and migration rates were considered significant if 95% credible sets (mean
migration rate ± 1.96 x mean standard deviation) did not overlap zero. Chain convergence
was assessed, and migration rates visualised using R (Figure S9).

Historical relationships among populations

To explore historical relationships among populations of *M. alfredi* and *M. birostris* we used
the program TreeMix (*46*). TreeMix uses population allele frequencies to estimate a bifurcating
maximum likelihood tree with which to infer historical population splits, admixture events and
the degree of genetic drift. We first supplemented the *M. alfredi* dataset with one randomly
selected *M. birostris* individual and the *M. birostris* dataset with one randomly selected *M.*
alfredi to act as outgroups when rooting the trees. Both datasets were then filtered for linkage,
a minor allele count of less than 3, genotyping rate of less than 90% and related individuals
using PLINK v1.9 (*98*). Allele frequencies for each population were then calculated using the
–freq and –within arguments in PLINK. For both the *M. birostris* and *M. alfredi* datasets we
then performed 10 initial runs of TreeMix for each migration event (M) ranging from 0 to 10.
The number of migration edges that explained 99.8% of the variance was selected as the best
model for each species (*M. birostris:* M= 0; *M. alfredi:* M = 2, Figure S5). We then re-ran
TreeMix 100 times using the optimal number of migration edges. Consensus trees and
bootstrap values were estimated and visualised using code modified from the BITE R package
(*99*).

Genome-wide heterozygosity

449 To assess levels of genetic diversity both within and between species we used the high-quality
450 SNP dataset comprising both species. Multi-locus heterozygosity was then calculated for each
451 individual using the R package inbreedR (*100*).

Acknowledgements

We are grateful to the following people and organisations for their support sourcing and
collecting tissue samples: Kerstin Forsberg, Jon Slayer, John Nevill, Dan Bowling, Heather
Pacey, Karen Fuentes, Manta Caribbean Project, Annie Murray, Tam Sawers, Beth Taylor,
Adam Thol'hath, Isabel Ender, Musa Mohamed, the Barefoot Collection, Four Seasons
Resorts Maldives, Six Senses Laamu and all Manta Trust and LAMAVE staff, volunteers and
field team. We are grateful to Helen Senn and Jennifer Kaden from the Royal Zoological

Society of Scotland for support during library preparation. ddRAD sequencing was carried out
by Edinburgh Genomics. Tissue sample collection in Seychelles was approved by the
Seychelles Bureau of Standards, the Seychelles Ministry of Environment, Energy and Climate
Change, and The University of Western Australia (RA/3/100/1480). Blue Resources Trust
(BRT) would like to thank the Department of Wildlife Conservation and the Department of
Fisheries and Aquatic Resources for enabling the fieldwork to be conducted in Sri Lanka.
Sample collection in the Maldives was made possible thanks to the permission and support
provided by the Ministry of Fisheries and Agriculture ([OTHR]30-D/INDIV/ 2016/220) and the
Ministry of Environment and Energy (EPA/2016/PSR-M02). Sampling in Mexico Caribbean
was made possible thanks to CONAPESCA (PPF/DGOPA-126/16 and PPF/DGOPA/315717)
and SEMARNAT (SGPA/DGVS/07618/15 and SGPA/DGVS/10067/17). We would like to
acknowledge Richard Coleman for helpful comments on the manuscript. For the purpose of
open access, the author has applied a Creative Commons Attribution (CC BY) licence to any
Author Accepted Manuscript version arising from this submission.

Funding

We are grateful to the Save Our Seas Foundation and The People's Trust for Endangered
Species who provided funding for this work. JH was supported by a Natural Environment
Research Council CASE studentship through the ENVISION DTP (case partner: Royal
Zoological Society of Scotland) and received additional grants from the Fisheries Society of
the British Isles and the Genetics Society. Fieldwork in Seychelles was supported by the
SOSF-D'Arros Research Centre. RB was funded by the Save Our Seas Foundation and the
Marine Conservation Action Fund of the New England Aquarium. BRT acknowledges the
generous support provided by the Save Our Seas Foundation and the Marine Conservation
and Action Fund (MCAF) for field work and sample collection.

Author contributions

EH, JH, GC, MdB, SC, GMWS and RO conceived and designed the study. GMWS, AA, RB,
MD, DF, NF, LRP, SP, AP, JDS and SW provided samples. JH and JK carried out laboratory
work. EH analysed the data with input from JH. EH wrote the paper with input from all other
co-authors.

Competing interests

The authors declare no competing interests.

Data and materials availability

Sequencing data will be available on the European Nucleotide Archive and scripts will be available on GitHub.

References

1. I. Hanski, M. Gilpin, Metapopulation dynamics: brief history and conceptual domain. *Biological Journal of the Linnean Society*. **42**, 3–16 (1991).

2. L. S. Mills, F. W. Allendorf, The one-migrant-per-generation rule in conservation and management. *Conservation Biology*. **10**, 1509–1518 (1996).

3. W. H. Lowe, F. W. Allendorf, What can genetics tell us about population connectivity? *Molecular Ecology*. **19**, 3038–3051 (2010).

4. D. H. Reed, Extinction risk in fragmented habitats. *Animal Conservation Forum*. **7**, 181–191 (2004).

5. S. R. Palumbi, Population genetics, demographic connectivity, and the design of marine reserves. *Ecological Applications*. **13**, 146–158 (2003).

6. M. Pascual, B. Rives, C. Schunter, E. Macpherson, Impact of life history traits on gene flow: A multispecies systematic review across oceanographic barriers in the Mediterranean Sea. *PLOS ONE*. **12**, e0176419 (2017).

7. P. J. Palsbøll, M. Bérubé, F. W. Allendorf, Identification of management units using population genetic data. *Trends in Ecology & Evolution*. **22**, 11–16 (2007).

8. W. C. Funk, J. K. McKay, P. A. Hohenlohe, F. W. Allendorf, Harnessing genomics for delineating conservation units. *Trends in Ecology & Evolution*. **27**, 489–496 (2012).

9. M. Kardos, E. E. Armstrong, S. W. Fitzpatrick, S. Hauser, P. W. Hedrick, J. M. Miller, D. A. Tallmon, W. C. Funk, The crucial role of genome-wide genetic variation in conservation. *Proceedings of the National Academy of Sciences*. **118**, e2104642118 (2021).

10. R. D. Ward, M. Woodwark, D. O. F. Skibinski, A comparison of genetic diversity levels in marine, freshwater, and anadromous fishes. *Journal of Fish Biology*. **44**, 213–232 (1994).

11. R. Waples, Separating the wheat from the chaff: patterns of genetic differentiation in high gene flow species. *Journal of Heredity*. **89**, 438–450 (1998).

12. R. S. Waples, A. E. Punt, J. M. Cope, Integrating genetic data into management of marine resources: how can we do it better? *Fish and Fisheries*. **9**, 423–449 (2008).

13. R. S. Waples, O. Gaggiotti, INVITED REVIEW: What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology*. **15**, 1419–1439 (2006).

14. D. Bailleul, A. Mackenzie, O. Sacchi, F. Poisson, N. Bierne, S. Arnaud-Haond, Large-scale genetic panmixia in the blue shark (*Prionace glauca*): A single worldwide population, or a genetic lag-time effect of the "grey zone" of differentiation? *Evolutionary Applications*. **11**, 614–630 (2018).

15. L. Hauser, G. R. Carvalho, Paradigm shifts in marine fisheries genetics: ugly hypotheses slain by beautiful facts. *Fish and Fisheries*. **9**, 333–362 (2008).

16. P.-A. Gagnaire, T. Broquet, D. Aurelle, F. Viard, A. Souissi, F. Bonhomme, S. Arnaud-Haond, N. Bierne, Using neutral, selected, and hitchhiker loci to assess connectivity of marine populations in the genomic era. *Evolutionary Applications*. **8**, 769–786 (2015).

17. A. D. Foote, P. A. Morin, Genome-wide SNP data suggest complex ancestry of sympatric North Pacific killer whale ecotypes. *Heredity*. **117**, 316–325 (2016).

18. M. Louis, M. Galimberti, F. Archer, S. Berrow, A. Brownlow, R. Fallon, M. Nykänen, J. O'Brien, K. M. Roberston, P. E. Rosel, B. Simon-Bouhet, D. Wegmann, M. C. Fontaine, A. D. Foote, O. E. Gaggiotti, Selection on ancestral genetic variation fuels repeated ecotype formation in bottlenose dolphins. *Science Advances*. **7**, eabg1245 (2021).

19. X. Liu, S. Rønhøj Schjøtt, S. M. Granquist, A. Rosing-Asvid, R. Dietz, J. Teilmann, A. Galatius, K. Cammen, G. O'Corry-Crowe, K. Harding, T. Härkönen, A. Hall, E. L. Carroll, Y. Kobayashi, M. Hammill, G. Stenson, A. Kirstine Frie, C. Lydersen, K. M. Kovacs, L. W. Andersen, J. I. Hoffman, S. J. Goodman, F. G. Vieira, R. Heller, I. Moltke, M. Tange Olsen, Origin and expansion of the world's most widespread pinniped: Range-wide population genomics of the harbour seal (Phoca vitulina). *Molecular Ecology*. **31**, 1682–1699 (2022).

20. P.-A. Gagnaire, Comparative genomics approach to evolutionary process connectivity. *Evolutionary Applications*. **13**, 1320–1334 (2020).

21. L. I. E. Couturier, A. D. Marshall, F. R. A. Jaine, T. Kashiwagi, S. J. Pierce, K. A. Townsend, S. J. Weeks, M. B. Bennett, A. J. Richardson, Biology, ecology and conservation of the Mobulidae. *Journal of Fish Biology*. **80**, 1075–1119 (2012).

22. T. Kashiwagi, A. D. Marshall, M. B. Bennett, J. R. Ovenden, The genetic signature of recent speciation in manta rays (Manta alfredi and M. birostris). *Molecular Phylogenetics and Evolution*. **64**, 212–218 (2012).

23. S. Andrzejaczek, R. J. Schallert, K. Forsberg, N. S. Arnoldi, M. Cabanillas-Torpoco, W. Purizaca, B. A. Block, Reverse diel vertical movements of oceanic manta rays off the northern coast of Peru and implications for conservation. *Ecological Solutions and Evidence*. **2**, e12051 (2021).

24. A. M. Knochel, J. E. M. Cochran, A. Kattan, G. M. W. Stevens, E. Bojanowksi, M. L. Berumen, Crowdsourced data reveal multinational connectivity, population demographics, and possible nursery ground of endangered oceanic manta rays in the Red Sea. *Aquatic Conservation: Marine and Freshwater Ecosystems*. **32**, 1774–1786 (2022).

25. L. Gordon, T. Vierus, First photographic evidence of oceanic manta rays (*Mobula birostris*) at two locations in the Fiji islands. *PeerJ*. **10**, e13883 (2022).

26. M. M. P. Cabral, J. D. Stewart, T. A. Marques, J. T. Ketchum, A. Ayala-Bocos, E. M. Hoyos-Padilla, H. Reyes-Bonilla, The influence of El Niño Southern Oscillation on the population dynamics of oceanic manta rays in the Mexican Pacific. *Hydrobiologia*. **850**, 257–267 (2023).

27. F. Garzon, Z. T. A. Seymour, Z. L. Monteiro, R. T. Graham, Spatial ecology of a newly described oceanic manta ray population in the Atlantic Ocean. *Mar Biol*. **170**, 68 (2023).

28. S. K. Venables, A. D. Marshall, A. J. Armstrong, J. L. Tomkins, W. J. Kennington, Genome-wide SNPs detect no evidence of genetic population structure for reef manta rays (Mobula alfredi) in southern Mozambique. *Heredity*. **126**, 308–319 (2021).

29. H. Lassauce, C. L. Dudgeon, A. J. Armstrong, L. Wantiez, E. L. Carroll, Evidence of fine-scale genetic structure for reef manta rays Mobula alfredi in New Caledonia. *Endangered Species Research*. **47**, 249–264 (2022).

30. J. L. Whitney, R. R. Coleman, M. H. Deakos, Genomic evidence indicates small island-resident populations and sex-biased behavior of Hawaiian Reef Manta Rays. *BMC Ecology and Evolution*. **in press** (in press).

31. J. Hosegood, E. Humble, R. Ogden, M. de Bruyn, S. Creer, G. M. W. Stevens, M. Abudaya, K. Bassos-Hull, R. Bonfil, D. Fernando, A. D. Foote, H. Hipperson, R. W. Jabado, J. Kaden, M. Moazzam, L. R. Peel, S. Pollett, A. Ponzo, M. Poortvliet, J. Salah, H. Senn, J. D. Stewart, S. Wintner, G. Carvalho, Phylogenomics and species delimitation for effective conservation of manta and devil rays. *Molecular Ecology*. **29**, 4783–4796 (2020).

32. J. D. Stewart, C. S. Beale, D. Fernando, A. B. Sianipar, R. S. Burton, B. X. Semmens, O. Aburto-Oropeza, Spatial ecology and conservation of *Manta birostris* in the Indo-Pacific. *Biological Conservation*. **200**, 178–183 (2016).

33. K. E. R. López, J. J. Guadalupe, M. Gordillo-Romero, A. Montero-Oleas, D. A. Pazmiño, M. Guerrero, M. de L. Torres, Exploring the genetic diversity and population structure of *Mobula birostris* in two key aggregation zones in the Eastern Tropical Pacific. *Marine Ecology Progress Series*. **699**, 75–89 (2022).

34. M. O'Malley, K. A. Townsend, P. Hilton, S. Heinrichs, J. D. Stewart, Characterization of the trade in manta and devil ray gill plates in China and South-east Asia through trader surveys. *Aquatic Conservation: Marine and Freshwater Ecosystems*. **27**, 394–413 (2017).

35. C. A. Ward-Paige, B. Davis, B. Worm, Global population trends and human use patterns of manta and mobula rays. *PLOS ONE*. **8**, e74835 (2013).

36. D. A. Croll, H. Dewar, N. K. Dulvy, D. Fernando, M. P. Francis, F. Galván-Magaña, M. Hall, S. Heinrichs, A. Marshall, D. Mccauley, K. M. Newton, G. Notarbartolo di Sciara, M. O'Malley, J. O'Sullivan, M. Poortvliet, M. Roman, G. Stevens, B. R. Tershy, W. T. White, Vulnerabilities and fisheries impacts: the uncertain future of manta and devil rays. *Aquatic Conservation: Marine and Freshwater Ecosystems*. **26**, 562–575 (2016).

37. C. A. Rohner, A. L. Flam, S. J. Pierce, A. D. Marshall, Steep declines in sightings of manta rays and devilrays (Mobulidae) in southern Mozambique. *PeerJ Preprints*. **5** (2017), doi:https://doi.org/10.7287/peerj.preprints.3051v1.

38. M. Moazzam, "Unprecedented decline in the catches of mobulids: an important component of tuna gillnet fisheries of the Northern Arabian Sea," *IOTC - 14th Working Party on Ecosystems and Bycatch* (IOTC-2018-WPEB14-30, Cape Town, South Africa, 2018).

39. M. Carpenter, D. Parker, M. L. Dicken, C. L. Griffiths, Multi-decade catches of manta rays (*Mobula alfredi, M. birostris*) from South Africa reveal significant decline. *Frontiers in Marine Science*. **10** (2023) (available at https://www.frontiersin.org/articles/10.3389/fmars.2023.1128819).

40. J. M. Lawson, S. V. Fordham, M. P. O'Malley, L. N. K. Davidson, R. H. L. Walls, M. R. Heupel, G. Stevens, D. Fernando, A. Budziak, C. A. Simpfendorfer, I. Ender, M. P. Francis, G. N. di Sciara, N. K. Dulvy, Sympathy for the devil: a conservation strategy for devil and manta rays. *PeerJ*. **5**, e3027 (2017).

614   41.   D. Fernando, J. D. Stewart, High bycatch rates of manta and devil rays in the "small-scale"
615   artisanal fisheries of Sri Lanka. *PeerJ*. **9**, e11994 (2021).

616   42.   J. M. Lawson, S. V. Fordham, "Sharks ahead: Realizing the potential of the convention on
617   migratory species to conserve elasmobranchs," *Shark Advocates International, The*
618   *Ocean Foundation* (Report, Shark Advocates International, The Ocean Foundation,
619   Washingston, D.C, 2018), pp. 1–76.

620   43.   J. D. Stewart, F. R. A. Jaine, A. J. Armstrong, A. O. Armstrong, M. B. Bennett, K. B.
621   Burgess, L. I. E. Couturier, D. A. Croll, M. R. Cronin, M. H. Deakos, C. L. Dudgeon, D.
622   Fernando, N. Froman, E. S. Germanov, M. A. Hall, S. Hinojosa-Alvarez, J. E. Hosegood,
623   T. Kashiwagi, B. J. L. Laglbauer, N. Lezama-Ochoa, A. D. Marshall, F. McGregor, G.
624   Notarbartolo di Sciara, M. D. Palacios, L. R. Peel, A. J. Richardson, R. D. Rubin, K. A.
625   Townsend, S. K. Venables, G. M. W. Stevens, Research priorities to support effective
626   manta and devil ray conservation. *Front. Mar. Sci.* **5** (2018),
627   doi:10.3389/fmars.2018.00314.

628   44.   M. Jakobsson, M. D. Edge, N. A. Rosenberg, The relationship between *Fst* and the
629   frequency of the most frequent allele. *Genetics*. **193**, 515–528 (2013).

630   45.   S. M. Mussmann, M. R. Douglas, T. K. Chafin, M. E. Douglas, BA3-SNPs: Contemporary
631   migration reconfigured in BayesAss for next-generation sequence data. *Methods in*
632   *Ecology and Evolution*. **10**, 1808–1813 (2019).

633   46.   J. K. Pritchard, M. Stephens, P. Donnelly, Inference of population structure using multilocus
634   genotype data. *Genetics*. **155**, 945–959 (2000).

635   47.   F. R. A. Jaine, C. A. Rohner, S. J. Weeks, L. I. E. Couturier, M. B. Bennett, K. A.
636   Townsend, A. J. Richardson, Movements and habitat use of reef manta rays off eastern
637   Australia: offshore excursions, deep diving and eddy affinity revealed by satellite
638   telemetry. *Marine Ecology Progress Series*. **510**, 73–86 (2014).

639   48.   A. O. Armstrong, A. J. Armstrong, M. B. Bennett, A. J. Richardson, K. A. Townsend, C. L.
640   Dudgeon, Photographic identification and citizen science combine to reveal long distance
641   movements of individual reef manta rays *Mobula alfredi* along Australia's east coast.
642   *Marine Biodiversity Records*. **12**, 14 (2019).

643   49.   M. Hirschfeld, C. Dudgeon, M. Sheaves, A. Barnett, Barriers in a sea of elasmobranchs:
644   From fishing for populations to testing hypotheses in population genetics. *Global Ecology*
645   *and Biogeography*. **30**, 2147–2163 (2021).

646   50.   M. H. Deakos, J. D. Baker, L. Bejder, Characteristics of a manta ray *Manta alfredi*
647   -population off Maui, Hawaii, and implications for management. *Marine Ecology Progress*
648   *Series*. **429**, 245–260 (2011).

649   51.   C. D. Braun, G. B. Skomal, S. R. Thorrold, M. L. Berumen, Movements of the reef manta
650   ray (*Manta alfredi*) in the Red Sea using satellite and acoustic telemetry. *Mar Biol*. **162**,
651   2351–2362 (2015).

652   52.   E. Setyawan, A. B. Sianipar, M. V. Erdmann, A. M. Fischer, J. A. Haddy, C. S. Beale, S. A.
653   Lewis, R. Mambrasar, Site fidelity and movement patterns of reef manta rays (*Mobula*
654   *alfredi*: Mobulidae) using passive acoustic telemetry in northern Raja Ampat, Indonesia.
655   *Nat. Conserv. Res.* **3** (2018), doi:10.24189/ncr.2018.043.

53.   L. R. Peel, G. M. W. Stevens, R. Daly, C. A. K. Daly, J. S. E. Lea, C. R. Clarke, S. P. Collin,
M. G. Meekan, Movement and residency patterns of reef manta rays *Mobula alfredi* in the
Amirante Islands, Seychelles. *Marine Ecology Progress Series*. **621**, 169–184 (2019).

54.   A. M. Knochel, N. E. Hussey, S. T. Kessel, C. D. Braun, J. E. M. Cochran, G. Hill, R. Klaus,
T. Checkchak, N. M. Elamin El Hassen, M. Younnis, M. L. Berumen, Home sweet home:
spatiotemporal distribution and site fidelity of the reef manta ray (*Mobula alfredi*) in
Dungonab Bay, Sudan. *Movement Ecology*. **10**, 22 (2022).

55.   E. S. Germanov, S. J. Pierce, A. D. Marshall, I. G. Hendrawan, A. Kefi, L. Bejder, N.
Loneragan, Residency, movement patterns, behavior and demographics of reef manta
rays in Komodo National Park. *PeerJ*. **10**, e13302 (2022).

56.   S. T. Kessel, N. A. Elamin, D. J. Yurkowski, T. Chekchak, R. P. Walter, R. Klaus, G. Hill, N.
E. Hussey, Conservation of reef manta rays (*Manta alfredi*) in a UNESCO World Heritage
Site: Large-scale island development or sustainable tourism? *PLOS ONE*. **12**, e0185419
(2017).

57.   E. S. Germanov, A. D. Marshall, Running the gauntlet: regional movement patterns of
Manta alfredi through a complex of parks and fisheries. *PLOS ONE*. **9**, e110071 (2014).

58.   R. Arauz, E. J. Chávez, E. M. Hoyos-Padilla, A. D. Marshall, First record of the reef manta
ray, Mobula alfredi, from the eastern Pacific. *Marine Biodiversity Records*. **12**, 3 (2019).

59.   V. Le Corre, A. Kremer, Cumulative effects of founding events during colonisation on
genetic diversity and differentiation in an island and stepping-stone model. *Journal of
Evolutionary Biology*. **11**, 495–512 (1998).

60.   Y.-T. Lai, C. K. L. Yeung, K. E. Omland, E.-L. Pang, Y. Hao, B.-Y. Liao, H.-F. Cao, B.-W.
Zhang, C.-F. Yeh, C.-M. Hung, H.-Y. Hung, M.-Y. Yang, W. Liang, Y.-C. Hsu, C.-T. Yao,
L. Dong, K. Lin, S.-H. Li, Standing genetic variation as the predominant source for
adaptation of a songbird. *Proceedings of the National Academy of Sciences*. **116**, 2152–
2157 (2019).

61.   T. Bonnet, M. B. Morrissey, P. de Villemereuil, S. C. Alberts, P. Arcese, L. D. Bailey, S.
Boutin, P. Brekke, L. J. N. Brent, G. Camenisch, A. Charmantier, T. H. Clutton-Brock, A.
Cockburn, D. W. Coltman, A. Courtiol, E. Davidian, S. R. Evans, J. G. Ewen, M. Festa-
Bianchet, C. de Franceschi, L. Gustafsson, O. P. Höner, T. M. Houslay, L. F. Keller, M.
Manser, A. G. McAdam, E. McLean, P. Nietlisbach, H. L. Osmond, J. M. Pemberton, E.
Postma, J. M. Reid, A. Rutschmann, A. W. Santure, B. C. Sheldon, J. Slate, C. Teplitsky,
M. E. Visser, B. Wachter, L. E. B. Kruuk, Genetic variance in fitness indicates rapid
contemporary adaptive evolution in wild animals. *Science*. **376**, 1012–1016 (2022).

62.   T. M. Vignaud, J. A. Maynard, R. Leblois, M. G. Meekan, R. Vázquez-Juárez, D. Ramírez-
Macías, S. J. Pierce, D. Rowat, M. L. Berumen, C. Beeravolu, S. Baksay, S. Planes,
Genetic structure of populations of whale sharks among ocean basins and evidence for
their historic rise and recent decline. *Molecular Ecology*. **23**, 2590–2601 (2014).

63.   M. S. Leslie, P. A. Morin, Structure and phylogeography of two tropical predators, spinner
(*Stenella longirostris*) and pantropical spotted (*S. attenuata*) dolphins, from SNP data. *R
Soc Open Sci*. **5**, 171615 (2018).

64.   N. Nikolic, F. Devloo-Delva, D. Bailleul, E. Noskova, C. Rougeux, C. Delord, P. Borsa, C.
Liautard-Haag, M. Hassan, A. D. Marie, P. Feutry, P. Grewe, C. Davies, J. Farley, D.
Fernando, S. Biton-Porsmoguer, F. Poisson, D. Parker, A. Leone, J. Aulich, M. Lansdell,

F. Marsac, S. Arnaud-Haond, Stepping up to genome scan allows stock differentiation in the worldwide distributed blue shark *Prionace glauca*. *Molecular Ecology*. **32**, 1000–1019 (2023).

65. J. L. Younger, G. V. Clucas, D. Kao, A. D. Rogers, K. Gharbi, T. Hart, K. J. Miller, The challenges of detecting subtle population structure and its importance for the conservation of emperor penguins. *Molecular Ecology*. **26**, 3883–3897 (2017).

66. L. I. E. Couturier, C. L. Dudgeon, K. H. Pollock, F. R. A. Jaine, M. B. Bennett, K. A. Townsend, S. J. Weeks, A. J. Richardson, Population dynamics of the reef manta ray *Manta alfredi* in eastern Australia. *Coral Reefs*. **33**, 329–342 (2014).

67. K. Harty, M. Guerrero, A. M. Knochel, G. M. W. Stevens, A. Marshall, K. Burgess, J. D. Stewart, Demographics and dynamics of the world's largest known population of oceanic manta rays *Mobula birostris* in coastal Ecuador. *Marine Ecology Progress Series*. **700**, 145–159 (2022).

68. J. Rambahiniarison, A. Agustines, K. Alexopoulos, G. Araujo, A. O. Armstrong, S. Arnold, A. Barruga, T. Cañete, S. Conales Jr., K. Delijero, N. P. Enolva, A. L. Flam, E. Keane, J. Labaja, C. G. Legaspi, C. Murie, R. Murray, S. P. Oliver, S. J. Pierce, A. Ponzo, C. A. Rohner, R. Schifferer, S. Snow, M. Spakowski, G. M. W. Stevens, T. Tilgel, J. N. C. Wong, A. A. Yaptinchay, Y. Barr, Distribution of the reef manta ray *Mobula alfredi* and the oceanic manta ray *Mobula birostris* in the Philippines: a collaborative effort for conservation. *Journal of Fish Biology*. **102**, 492–503 (2023).

69. C. A. Rohner, S. J. Pierce, A. D. Marshall, S. J. Weeks, M. B. Bennett, A. J. Richardson, Trends in sightings and environmental influences on a coastal aggregation of manta rays and whale sharks. *Marine Ecology Progress Series*. **482**, 153–168 (2013).

70. C. S. Beale, J. D. Stewart, E. Setyawan, A. B. Sianipar, M. V. Erdmann, Population dynamics of oceanic manta rays (*Mobula birostris*) in the Raja Ampat Archipelago, West Papua, Indonesia, and the impacts of the El Niño–Southern Oscillation on their movement ecology. *Diversity and Distributions*. **25**, 1472–1487 (2019).

71. S. Wright, Evolution in Mendelian populations. *Genetics*. **16**, 97–159 (1931).

72. N. M. Phillips, F. Devloo-Delva, C. McCall, T. S. Daly-Engel, Reviewing the genetic evidence for sex-biased dispersal in elasmobranchs. *Rev Fish Biol Fisheries*. **31**, 821–841 (2021).

73. M. A. McClain, N. Hammerschlag, A. J. Gallagher, J. M. Drymon, R. D. Grubbs, T. L. Guttridge, M. J. Smukall, B. S. Frazier, T. S. Daly-Engel, Age-dependent dispersal and relatedness in tiger sharks (*Galeocerdo cuvier*). *Frontiers in Marine Science*. **9** (2022) (available at https://www.frontiersin.org/articles/10.3389/fmars.2022.900107).

74. Y. P. Papastamatiou, C. G. Meyer, F. Carvalho, J. J. Dale, M. R. Hutchinson, K. N. Holland, Telemetry and random-walk models reveal complex patterns of partial migration in a large marine predator. *Ecology*. **94**, 2595–2606 (2013).

75. J. Thorburn, F. Neat, I. Burrett, L.-A. Henry, D. M. Bailey, C. S. Jones, L. R. Noble, Ontogenetic variation in movements and depth use, and evidence of partial migration in a benthopelagic elasmobranch. *Frontiers in Ecology and Evolution*. **7** (2019) (available at https://www.frontiersin.org/articles/10.3389/fevo.2019.00353).

76. R. J. Y. Perryman, J. Mourier, S. K. Venables, R. F. Tapilatu, E. Setyawan, C. Brown, Reef
manta ray social dynamics depend on individual differences in behaviour. *Animal*
Behaviour. **191**, 43–55 (2022).

77. B. L. Taylor, A. E. Dizon, The need to estimate power to link genetics and demography for
conservation. *Conservation Biology*. **10**, 661–664 (1996).

78. J. R. Hoffman, J. R. Willoughby, B. J. Swanson, K. L. Pangle, D. T. Zanatta, Detection of
barriers to dispersal is masked by long lifespans and large population sizes. *Ecology and*
Evolution. **7**, 9613–9623 (2017).

79. N. K. Dulvy, S. A. Pardo, C. A. Simpfendorfer, J. K. Carlson, Diagnosing the dangerous
demography of manta rays using life history theory. *PeerJ*. **2**, e400 (2014).

80. H. Cayuela, Q. Rougemont, J. G. Prunier, J.-S. Moore, J. Clobert, A. Besnard, L.
Bernatchez, Demographic and genetic approaches to study dispersal in wild animal
populations: A methodological review. *Molecular Ecology*. **27**, 3976–4010 (2018).

81. G. Stevens, D. Fernando, M. Dando, G. Notarbartolo di Sciara, *Guide to the Manta and*
Devil Rays of the World (Princeton University Press, 2018).

82. J. D. Stewart, M. Nuttall, E. L. Hickerson, M. A. Johnston, Important juvenile manta ray
habitat at Flower Garden Banks National Marine Sanctuary in the northwestern Gulf of
Mexico. *Mar Biol*. **165**, 111 (2018).

83. J. H. Pate, A. D. Marshall, Urban manta rays: potential manta ray nursery habitat along a
highly developed Florida coastline. *Endangered Species Research*. **43**, 51–64 (2020).

84. S. Hinojosa-Alvarez, R. P. Walter, P. Diaz-Jaimes, F. Galván-Magaña, E. M. Paig-Tran, A
potential third Manta Ray species near the Yucatán Peninsula? Evidence for a recently
diverged and novel genetic Manta group from the Gulf of Mexico. *PeerJ*. **4**, e2586 (2016).

85. B. K. Peterson, J. N. Weber, E. H. Kay, H. S. Fisher, H. E. Hoekstra, Double Digest
RADseq: An inexpensive method for de novo snp discovery and genotyping in model and
non-model species. *PLoS ONE*. **7**, e37135 (2012).

86. C. Palaiokostas, M. Bekaert, M. G. Khan, J. B. Taggart, K. Gharbi, B. J. McAndrew, D. J.
Penman, A novel sex-determining QTL in Nile tilapia (Oreochromis niloticus). *BMC*
Genomics. **16**, 171 (2015).

87. J. Catchen, P. A. Hohenlohe, S. Bassham, A. Amores, W. A. Cresko, Stacks: an analysis
tool set for population genomics. *Molecular Ecology*. **22**, 3124–3140 (2013).

88. N. C. Rochette, J. M. Catchen, Deriving genotypes from RAD-seq short-read data using
Stacks. *Nat Protoc*. **12**, 2640–2659 (2017).

89. T. Korneliussen, I. Moltke, NgsRelate: a software tool for estimating pairwise relatedness
from next-generation sequencing data. *Bioinformatics (Oxford, England)*. **31** (2015), ,
doi:10.1093/bioinformatics/btv509.

90. R. K. Waples, A. Albrechtsen, I. Moltke, Allele frequency-free inference of close familial
relationships from genotypes or low-depth sequencing data. *Molecular Ecology*. **28**, 35–48
(2019).

91. T. Jombart, adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics. **24**, 1403–1405 (2008).

92.   T. Jombart, S. Devillard, F. Balloux, Discriminant analysis of principal components: a new
method for the analysis of genetically structured populations. *BMC Genetics*. **11**, 94
(2010).

93.   J. A. Thia, Guidelines for standardizing the application of discriminant analysis of principal
components to genotype data. *Molecular Ecology Resources*. **23**, 523–538 (2023).

94.   B. S. Weir, C. C. Cockerham, Estimating f-statistics for the analysis of population structure.
Evolution. **38**, 1358–1370 (1984).

95.   B. Gruber, P. J. Unmack, O. F. Berry, A. Georges, dartr: An r package to facilitate analysis
of SNP data generated from reduced representation genome sequencing. *Molecular
Ecology Resources*. **18**, 691–699 (2018).

96.   E. Pante, B. Simon-Bouhet, marmap: A package for importing, plotting and analyzing
bathymetric and topographic data in R. *PLOS ONE*. **8**, e73051 (2013).

97.   S. Dray, A.-B. Dufour, The ade4 package: Implementing the duality diagram for ecologists.
Journal of Statistical Software. **22**, 1–20 (2007).

98.   S. Purcell, B. Neale, K. Todd-Brown, L. Thomas, M. A. R. Ferreira, D. Bender, J. Maller, P.
Sklar, P. I. W. de Bakker, M. J. Daly, P. C. Sham, PLINK: a tool set for whole-genome
association and population-based linkage analyses. *The American Journal of Human
Genetics*. **81**, 559–575 (2007).

99.   M. Milanesi, S. Capomaccio, E. Vajana, L. Bomba, J. F. Garcia, P. Ajmone-Marsan, L. Colli,
BITE: an R package for biodiversity analyses (2017), p. 181610.

100. M. A. Stoffel, M. Esser, M. Kardos, E. Humble, H. Nichols, P. David, J. I. Hoffman,
inbreedR: an R package for the analysis of inbreeding based on genetic markers.
Methods in Ecology and Evolution. **7**, 1331–1339 (2016).

806