Temporally resolved early BMP-driven transcriptional cascade during human amnion specification

AUTHORS:

Nikola Sekulovski[1], Jenna C. Wettstein[1], Amber E. Carleton[1], Linnea E. Taniguchi[1], Xiaolong Ma[2], Sridhar Rao[1,3,4], Jenna K. Schmidt[5], Thaddeus G. Golos[5,6,7], Chien-Wei Lin[2, †] and Kenichiro Taniguchi[1,3,†]

AFFILIATIONS:

[1]Department of Cell Biology, Neurobiology and Anatomy, Medical College of Wisconsin, Milwaukee, WI 53226, USA

[2]Division of Biostatistics, Institute for Health and Equity, Medical College of Wisconsin, Milwaukee, WI 53226, USA

[3]Department of Pediatrics, Medical College of Wisconsin, Milwaukee, WI 53226, USA

[4]Versiti Blood Research Institute, Milwaukee, WI 53226 USA

[5]Wisconsin National Primate Research Center (WNPRC), Madison, WI, USA

[6]Department of Obstetrics and Gynecology, University of Wisconsin - Madison School of Medicine and Public Health, Madison, WI USA

[7]Department of Comparative Biosciences, University of Wisconsin - Madison School of Veterinary Medicine, Madison, WI, USA

[†]Co-corresponding authors

CONTACT:

Kenichiro Taniguchi
8701 Watertown Plank Road
B4245 BSB CB062
Milwaukee, WI 53226
ktaniguchi@mcw.edu

Chien-Wei Lin
8701 Watertown Plank Road
MEB M1406
Milwaukee, WI 53226
chlin@mcw.edu

ABSTRACT

Amniogenesis, a process critical for continuation of healthy pregnancy, is triggered in a collection of pluripotent epiblast cells as the human embryo implants. Previous studies have established that BMP signaling is a major driver of this lineage specifying process, but the downstream BMP-dependent transcriptional networks that lead to successful amniogenesis remain to be identified. This is, in part, due to the current lack of a robust and reproducible model system that enables mechanistic investigations exclusively into amniogenesis. Here, we developed an improved model of early amnion specification, using a human pluripotent stem cell-based platform in which the activation of BMP signaling is controlled and synchronous. Uniform amniogenesis is seen within 48 hours after BMP activation, and the resulting cells share transcriptomic characteristics with amnion cells of a gastrulating human embryo. Using detailed time-course transcriptomic analyses, we established a previously uncharacterized BMP-dependent amniotic transcriptional cascade, and identified markers that represent five distinct stages of amnion fate specification; the expression of selected markers was validated in early post-implantation macaque embryos. Moreover, a cohort of factors that could potentially control specific stages of amniogenesis was identified, including the transcription factor TFAP2A. Functionally, we determined that, once amniogenesis is triggered by the BMP pathway, TFAP2A controls the progression of amniogenesis. This work presents a temporally resolved transcriptomic resource for several previously uncharacterized amniogenesis states and demonstrates a critical intermediate role for TFAP2A during amnion fate specification.

INTRODUCTION

The amniotic epithelium is the innermost layer of the amniochorionic membrane that physically divides the maternal and fetal environments (Miki et al., 2005; Miki and Strom, 2006). Amniogenesis is initiated during implantation in humans (and in non-human primates), eventually leading to the formation of an amniotic sac structure that surrounds and protects the developing embryo (Enders et al., 1983; Enders et al., 1986; Miki et al., 2005; Sasaki et al., 2016; Shahbazi and Zernicka-Goetz, 2018; Taniguchi et al., 2019). At implantation, the embryo (referred to at this time as a blastocyst) contains three functionally distinct cell types: a collection of unpolarized pluripotent epiblast cells (precursors to the embryo proper and amniotic ectoderm), a surrounding layer of polarized trophectoderm (a placental tissue precursor) and an underlying extraembryonic primitive endoderm (a yolk sac precursor). During implantation, the pluripotent epiblast cells undergo apico-basal polarization to form a cyst with a central lumen, the future amniotic cavity (Carleton et al., 2022). This event is followed by the fate transition of pluripotent epiblast cells at the uterine-proximal pole of the cyst to squamous amniotic ectoderm, forming a sharp boundary between amnion and pluripotent epiblast portions of the cyst (Shahbazi and Zernicka-Goetz, 2018; Shao and Fu, 2022; Taniguchi et al., 2019). This structure, the amniotic sac, represents the substrate for the next essential steps of embryonic development. Recent transcriptomic studies of early human and non-human primate embryos have presented benchmark resources for the amnion transcriptome of primates (Bergmann et al., 2022; Nakamura et al., 2016; Nakamura et al., 2017; Sasaki et al., 2016; Tyser et al., 2021; Yang et al., 2021).

The mechanistic dissection of human amnion development presents a challenge,

since it is ethically unacceptable to manipulate human embryos *in vivo* and technically

difficult to carry out such studies *in vitro*. Non-human primate (NHP) embryos such as

rhesus macaque are a suitable alternative to human embryos, due to their close

ancestral similarities (Nakamura et al., 2021), and have already provided crucial insights

into molecular processes that could potentially regulate amniogenesis. Indeed, genetic

manipulation has been demonstrated in NHP systems to study peri-implantation

embryogenesis (Yang et al., 2021), although this is not yet a widely applicable approach

due to resource availability and logistical complexity of such studies, which also carry

their own unique ethical challenges.

Previously, we developed an *in vitro* system of human amnion development

called Gel-3D (Shao et al., 2017a; Shao et al., 2017b), in which human pluripotent stem

cells (hPSC) are plated on a soft gel substrate consisting of extracellular matrix (ECM).

Within 24 hours, aggregates of hPSC form polarized cysts composed of columnar

pluripotent cells. Over the next three to four days, a 3D environment is introduced by

supplementing the culture medium with a diluted ECM overlay. Under these conditions,

the initially pluripotent cysts spontaneously undergo squamous morphogenesis, lose

pluripotency markers, and begin to express amnion markers; in over 90% of such cysts,

the central lumen becomes entirely surrounded by squamous amnion cells (hPSC-

amnion, (Shao et al., 2017a)). Using this system, we showed that mechanically

activated Bone Morphogenetic Protein (BMP) signaling can trigger amniogenesis and

identified several genes that are now used as amnion markers (e.g., GATA3, TFAP2A,

GABRP, (Shao et al., 2017a; Shao et al., 2017b)). These findings were supported by

later studies using early human embryo-like models as well as by single cell

transcriptomic studies in early human and NHP embryos (Chen et al., 2021; Tyser et al.,

2021; Yang et al., 2021; Zheng et al., 2019; Zheng et al., 2022).

Though this Gel-3D system (Shao et al., 2017a) has been valuable in allowing

initial investigations of the mechanisms underlying amniogenic differentiation, it suffers

from the fact that the initiation of amniogenesis by mechanical cues is stochastic and is

not fully penetrant, hindering reproducible examinations into temporally and spatially

sensitive processes. To better implement mechanistic studies, we developed a highly

reproducible hPSC-based amniogenic system, called Glass-3D^{+BMP}, that does not rely

on the mechanically activated spontaneous BMP signaling to initiate amniogenesis.

Rather, in Glass-3D^{+BMP}, BMP4 ligand is added exogenously, and the amniotic

differentiation response is quickly seen in all cells. Interestingly, while BMP signaling

activation is immediately detectable in all cells, squamous morphogenesis (starting

approximately 24 hours after BMP4 addition) is initially focal within a given cyst, but

then spreads laterally, to form fully squamous amnion cysts by 48 hours. Such amnion

cysts (hPSC-amnion) are uniformly composed of mature amniotic cells that share

similar transcriptomes with amnion cells from a Carnegie stage (CS) 7 human embryo

(dataset from (Tyser et al., 2021)).

Given the robust and synchronous nature of this system, we perform detailed

time-course bulk RNA sequencing analyses of developing Glass-3D^{+BMP} hPSC-amnion

to comprehensively characterize changes in transcriptional characteristics during the

initial stages of amnion fate specification. Strikingly, from 0 to 48 hours after BMP

treatment, developing hPSC-amnion displays transcriptomic characteristics of epiblast,

primitive streak-like or amnion cells that are present in the CS7 human embryo,

revealing a potential amnion differentiation trajectory. Moreover, we systematically

characterize dynamically expressed genes, and identify immediate-, early-,

intermediate- and late-responding genes associated with amniotic differentiation, some

of which are validated and characterized in detail in NHP embryos. Interestingly, in

contrast to the previous idea that TFAP2A is a *pan*-amnion marker expressed

throughout amniogenesis, we establish that TFAP2A is an early/intermediate

responding gene, with a temporal pattern of activation that is distinct from immediate

responding genes such as GATA3, another *pan*-amnion marker. Functional analyses

show that, in the absence of TFAP2A, amniogenesis is incomplete, suggesting that

TFAP2A controls amnion fate progression. Lastly, we identify a population of cells at the

boundary of the amnion and the epiblast in the peri-gastrula macaque embryo that

shares striking molecular and cellular characteristics of cells undergoing amnion

specification in Glass-3D^{+BMP}. Together, using the Glass-3D^{+BMP} amnion platform, these

studies present a transcriptomic map of the early stages of amnion differentiation, and

identify a critical transcriptional pathway dependent on TFAP2A, as well as a potential

site of active amniogenesis in the primate peri-gastrula.

RESULTS

Generation of fully squamous hPSC-derived amniotic cysts in a reproducible and

controlled manner using a Glass-3D^{+BMP} culture system

Targeted mechanistic investigations into amnion fate specification has been limited by

the fact that amniogenesis is not fully penetrant in existing *in vitro* embryo-like models

(Chen et al., 2021; Nasr Esfahani et al., 2019; Shao et al., 2017a; Shao et al., 2017b;

Zheng et al., 2019), prompting us to explore alternative strategies. Since our previous

studies have established that BMP signaling is a major driver of amniogenesis (Shao et

al., 2017a; Shao et al., 2017b), we sought to establish a system in which we could

directly control the activation of BMP signaling. We began with a culture condition called

Glass-3D, in which aggregates of hPSC are cultured in a stiff glass substrate condition

with the culture medium supplemented with a 3D ECM overlay (**Fig. 1A**). Under these

conditions, aggregates of hPSC initiate radial organization and form cysts with a central

lumen within 48 hours (Shao et al., 2017a; Taniguchi et al., 2017; Taniguchi et al.,

2015). Importantly, these cysts are composed entirely of polarized pluripotent cells.

When we examined the polarized membrane spatial proteomics of these hPSC-

pluripotent cysts, we found that BMP receptors are enriched at the basolateral

membrane territory (Wang et al., 2021). To test whether BMP signaling could be directly

activated in these cysts by the addition of BMP to the medium, pluripotent cysts (derived

from H9 (WA09) human embryonic stem cells for all experiments, unless otherwise

noted) were treated with exogenously provided BMP4 at d2 (**Fig. 1B**, Glass-3D^{+BMP}).

Nuclear localization of phosphorylated SMAD1/5 (pSMAD1/5), as detected by

immunofluorescence (IF), was used as a readout for BMP signaling activation (**Fig. 1C**).

At d2 in Glass-3D, prior to the addition of BMP, a polarized pluripotent (SOX2$^+$) cyst

structure has formed, and a small central lumen is visible (**Fig. 1C** – 0-min); this

structure is similar to that of the developing epiblast immediately prior to the initiation of

amniogenesis (Shahbazi and Zernicka-Goetz, 2018; Taniguchi et al., 2019). Strikingly,

within 10 minutes after BMP addition, uniform nuclear pSMAD1/5 enrichment is seen in

all treated cells at an equivalent level (**Fig. 1C** – 10-min, fluorescent quantitation in **Fig.**

1D). This is quickly followed by the expression of GATA3, a *pan*-amnion marker and a

well-known immediate downstream BMP target that has been shown to facilitate

pluripotency exit (Gunne-Braden et al., 2020). GATA3 expression is detected in all cells

within 3 hours (**Fig. 1E**), suggesting that BMP signaling is uniformly activated.

Interestingly, TFAP2A, another transcription factor thought to be an early *pan*-amnion

marker, is not detectable in GATA3 positive cells 3 hours after BMP addition (**Fig. 1E**).

After 48-hr of BMP4 treatment, all cysts show uniform squamous morphogenesis, loss

of pluripotency markers (NANOG, SOX2) and activated expression of additional *pan*-

amnion markers, TFAP2A and ISL1 (**Fig. 1F-H**).

To investigate whether exogenous BMP4 ligand is continuously needed

throughout the 48-hour time-course in order to form fully squamous hPSC-amnion, d2

pluripotent cysts were cultured with BMP4 for 24 hours (d2 to d3), then cultured without

BMP4 for the following 24 hours (d3 to d4), and then harvested at d4 (4 days after initial

plating, **Fig. 1B**). Interestingly, these 24-hr BMP treatment samples show fully

squamous amnion cysts that are morphologically and molecularly similar to those

treated over 48-hr. That is, they show uniformly squamous cyst morphogenesis and

prominent expression of GATA3, TFAP2A and ISL1, (**Fig. 1I,J**, quantitation for

fluorescent intensity and nuclear aspect ratios in **Fig. 1K,L**), as well as relatively similar

nuclear pSMAD1/5 signals (**Fig. 1M**). Moreover, in the 24-hr BMP treatment samples

(**Fig. 2A**), time-course analyses show that heterogeneous GATA3 expression is visible

in a few nuclei at 3-hr and 6-hr, becoming more prominent by 12-24-hr (**Fig. 2B-D**).

Early GATA3 expression is followed by initiation of TFAP2A expression by 12-hr (**Fig.**

2B,D). Two other "*pan*-amnion" genes, ISL1 and HAND1 become activated by 24-hr

(**Fig. 2C,E**), revealing a previously unrecognized GATA3 → TFAP2A → ISL1/HAND1

amniotic gene activation order (**Fig. 2B,C**, mRNA expression levels in **Fig. 2D,E**).

Importantly, SOX2 expression is abundant at 6-hr, fades in some nuclei at 12-hr and is

largely diminished by 24-hr (**Fig. 2B,C**, mRNA expression levels in **Fig. 2F**). NANOG

expression follows a similar time course to that of SOX2, revealing a gradual loss of

pluripotency (**Fig. 2F**). Samples treated with BMP4 for 24-hr or 48-hr display similar

levels of amnion and pluripotency markers (**Fig. 2G**), suggesting that BMP4 treatment

may be dispensable after the first 24 hours. To test this possibility, we treated cysts with

BMP4 for 24 hours (d2-d3) and subsequently treated with LDN-193189, a small

molecule inhibitor for the BMP receptor kinase activity (d3-d4). This treatment resulted

in fully squamous cysts that are positive for GATA3 and ISL1/2 (**Fig. 2H**), confirming

that BMP signaling initiates the early amniogenesis cascade, but is not required for its

progression.

Together, these results show that BMP4 treatment of d2 pluripotent cysts in a

Glass-3D culture environment provides a robust amniogenic model that enables

mechanistic investigations. We confirmed this finding in a human induced pluripotent

stem cell line (1196a, **Fig. S1**, (Chen et al., 2014)), further establishing this protocol as

an effective model of amnion fate conversion *in vitro*. Given that BMP4 treatment from

d3 to d4 is dispensable, 24-hour BMP4 treatment between d2 and d3 is used in all

remaining experiments using Glass-3D^{+BMP}, unless otherwise noted.

Transcriptomic characterization of developing Glass-3D^{+BMP} hPSC-amnion

To comprehensively understand the transcriptomic changes during hPSC-amnion

development in the Glass-3D^{+BMP} system, we performed bulk RNA sequencing analyses

for samples taken at 0-, 0.5-, 1-, 3-, 6-, 12-, 24- and 48-hr timepoints (post-BMP4

treatment). Samples were also taken from pluripotent monolayers and d4 pluripotent

cysts (**Fig. 3A** (experimental outline), **3B** (heatmap), **3C** (principal component analysis

(PCA)), see **Table 1A** for normalized and filtered counts). Overall, a total of 12,891

genes were detected among all sequenced samples. Similar to known immediate BMP-

responding genes (e.g., *ID2*, *BAMBI*), *GATA3* shows a significant increase by 1-hr, and

continuously increases overtime (**Fig. 3D**, >100-fold increase by 1-hr). Other markers

also display expected expression patterns (**Fig. 3E-G**). For example, *TFAP2A* and *ISL1*

show significant increases by 3-hr and 24-hr, respectively (**Fig. 3E**), and pluripotency

markers (**Fig. 3G**, *NANOG*, *OCT4/POU5F1*, *SOX2*) show reduction overtime. A

comparison of gene expression between d4 Glass-3D pluripotent cysts and d4 Glass-

3D^{+BMP} hPSC-amnion (48-hr timepoint) shows 1,820 upregulated and 2,092 down-

regulated (cut-off = 2-fold, p<0.05) genes in d4 Glass-3D^{+BMP} hPSC-amnion (**Fig. 3H**,

see **Table 1B,C** for full lists). Highly upregulated markers include known advanced

amnion state markers such as *GABRP*, *IGFBP3* and *EPAS1* (**Fig. 3H**), and highly

downregulated genes include *SOX2* and *NANOG* (**Fig. 3H**, see **Table 1D**,**E** for Gene

Ontology (GO) enrichment analyses (Biological Processes) for all significantly

upregulated and downregulated genes (> 2-fold, FDR < 0.05)).

The transcriptomic characteristics of developing amnion cells from a Carnegie

stage (CS) 7 human embryo were recently described by Tyser *et al.* (**Fig. 4A** shows a

UMAP (Uniform Manifold Approximation and Projection) plot with the original

coordinates and annotations, (Tyser et al., 2021)). In this analysis, the authors state that

yolk sac and connecting stalk tissues were discarded during dissection, but amnion

tissue was intact and included in the single cell preparation that was analyzed (Tyser et

al., 2021). Consistent with this notion, our analysis in which expression was

superimposed on the UMAP plot shows that *pan*-amnion markers such as *GATA3*,

TFAP2A and *ISL1* are broadly expressed in the rod-shaped cell population marked

"Ectoderm" (**Fig. 4B**, areas within the inset "**i**" are shown in **4B-D**; see **Fig. S2A** for

uncropped UMAP expression plots), suggesting that this region contains cells of the

amniotic lineage. We next examined which cells in the CS7 human embryo share

transcriptomic similarities to developing hPSC-amnion, and found that, indeed, the 48-hr

Glass-3D^{+BMP} samples exhibit clear transcriptional similarities to cells marked

"Ectoderm" in the CS7 human embryo (**Fig. 4C**), validating the human amnion-like

characteristics of our hPSC-amnion model. Interestingly, the earlier timepoints also

show overlaps with cell types of the CS7 human embryo (**Fig. 4C**, uncropped plot

shown in **Fig. S2B**). While pluripotent monolayer, d4 Glass-3D pluripotent cyst, as well

as 0-, 0.5-, 1-, 3- and 6-hr samples transcriptomically resemble Tyser "Epiblast" cells

(orange dots in **Fig. 4C**), 12-hr samples show transcriptomic similarities with cells

labeled in the Tyser analysis as "Primitive Streak"; these cells map close to the

Epiblast/Primitive Streak boundary in the Tyser UMAP (green dots in **Fig. 4C**). By 24-hr,

the transcriptomes are most similar to Tyser "Primitive Streak" cells proximal to the

amniotic "Ectoderm" (blue dots in **Fig. 4C**). Despite their "Primitive Streak" annotation,

these 24-hr cells adjacent to the amniotic "Ectoderm" population display relatively low

levels of *TBXT* (a major marker of primitive streak cells, **Fig. 4D**, (Beddington and

Robertson, 1999; Tyser et al., 2021; Zheng et al., 2019)), marking them as distinct from

the "Primitive Streak" cells progressing to "Nascent Mesoderm" (**Fig. 4D**). These

bioinformatic analyses suggest that the Glass-3D^{+BMP} amnion may develop from

pluripotent epiblast-like cells that are transcriptomically equivalent to the CS7 epiblast

cells, and, then traverse $TBXT^{low}$ primitive streak-like intermediate states as the lineage

specification continues before acquiring an amniotic fate by 48-hr (dotted arrow, **Fig.**

4C).

Given that the hPSC-amnion differentiation trajectory in the Glass-3D^{+BMP} model

overlaps with cells in the CS7 human embryo, it appears that active amniogenesis

processes may be present in the early post-implantation embryo. Thus, we applied an

RNA velocity analysis (Bergen et al., 2020) to the Tyser data; in such an analysis,

lineage progression is inferred from comparing spliced and unspliced mRNA. **Fig. 5A**

(inset "**i**") shows that the Tyser velocity data show differentiation trajectories very similar

to the one inferred from the time-course analysis of the Glass-3D^{+BMP} model

(trajectories starting from "Epiblast" → "Primitive Streak" → "Ectoderm").

To further validate this notion in a primate system, we performed detailed IF

expression analyses for GATA3, TFAP2A, ISL1 and SOX2 in an early cynomolgus

macaque embryo staged between CS6 and CS7 (**Fig. 5B-E**). In this CS6/7 embryo, a

clear amniotic sac is seen, with SOX2$^+$ epiblast cells on one side and contiguous ISL1$^+$

squamous amniotic ectoderm cells that are also positive for GATA3 and TFAP2A on the

other (**Fig. 5B-E**, fluorescent intensity quantitation shown in **5C,E**). Surface ectoderm

bridging neural ectoderm and amniotic ectoderm is not seen at this stage; this finding is

consistent with the previous histological studies in human (the Carnegie Collection,

found in the Virtual Human Embryo Project) and NHP pre- and peri-gastrula embryos

(Enders et al., 1983; Enders et al., 1986; Yang et al., 2021). Although amniotic

mesenchyme is not evident, some mesenchymal cells are observed underlying the

epiblast (**Fig. 5F**, green dotted bracket). These mesenchymal cells show abundant

nuclear pSMAD1/5: these cells are likely derived from the primitive streak (**Fig. 5F**).

Strikingly, at the boundary between the epiblast and amnion, a gradual transition in cell

morphology is readily seen: while cells most proximal to the epiblast are columnar, there

is a gradual reduction in height until fully squamous cells are seen (**Fig. 5B,D,F,H**),

represented by the changes in nuclear aspect ratio (**Fig. 5H**). Molecularly, the

expression of GATA3, TFAP2A, ISL1 and SOX2 is correlated with such columnar to

squamous cell shape change (**Fig. 5B-E**). While SOX2 expression shows a reducing

trend, the expression of GATA3, TFAP2A and ISL1 gradually increases in more distal

squamous cells (**Fig. 5B-E**, quantitation in **5C,E**). Furthermore, GATA3, TFAP2A and

ISL1 each display distinct expression patterns within the boundary cell population, as

the expression of GATA3 and TFAP2A is first detected in columnar SOX2$^+$ cells, but in

these cells, ISL1 is not observed (**Fig. 5C**). However, a weak but clear expression of

ISL1 can be found within ~10 distally located cells, from the first detected

GATA3$^+$/TFAP2A$^+$ cell. Similar observations are seen in another early cynomolgus

macaque embryo (**Fig. S3**, staged at CS7). This cell shape change associated with the

ordered transcriptional events (from GATA3, TFAP2A then ISL1) is similar to the

ordered morphological and molecular changes seen in developing Glass-3D^{+BMP} hPSC-

amnion. Thus, the results based on these bioinformatic and expression analyses

provide evidence for the presence of amniotic fate progression in early post-gastrula

embryos at the boundary between amnion and epiblast (a schematic model shown in

Fig. 5I).

Gene expression dynamics in developing Glass-3D^{+BMP} hPSC-amnion

To identify key players potentially involved in amnion fate specification at different

stages, we comprehensively characterized all significantly upregulated genes based on

when significant upregulations are seen: "immediate" (**Table 2A**, up in 0.5- or 1-hr

samples), "early" (**Table 2B**, 3- or 6-hr), "intermediate-1" (**Table 2C**, 12-hr),

"intermediate-2" (**Table 2D**, 24-hr) and "late" (**Table 2E**, 48-hr); genes that were kept

after a more stringent filtering (described in Materials and Methods) are shown in **Fig.**

6), enabling the identification of additional temporal markers.

Previous studies have identified several *pan*-amnion markers that are known

immediate BMP targets (e.g., *GATA3* and others in the list of "immediate" genes, **Table**

2A), which are expressed in other early BMP-driven tissue morphogenesis. Among the

known *pan*-amnion markers, ISL1 is a highly promising marker for early lineage

specified amnion cells (Yang et al., 2021; Zhao et al., 2021; Zheng et al., 2022). In

accord with this, we found that ISL1 expression is limited to squamous amnion cells and

is not seen in the epiblast or in transitioning boundary cells (**Fig. 5B**). Notably, in Glass-

3D^{+BMP}, *ISL1* first shows a significant upregulation at 24-hr (intermediate -2, **Fig. 7A**).

This correlates with the timeframe at which squamous cells are first observed (**Fig. 2**).

This finding suggests that amnion specification is likely initiated prior to 24 hours in

Glass-3D^{+BMP} and that genes that are upregulated between 12-24-hr could be used as

additional markers for amnion specification. Consistent with the studies by Tyser *et al.*,

we identified *DLX5* (716.76-fold increase between 12- and 24-hr, and 2.4-fold increase

from 24- to 48-hr, **Fig. 7B**). Interestingly, however, our expression analysis of the Tyser

et al. dataset reveals a broader *DLX5* expression pattern compared to *ISL1* (the ISL1

expression domain map is similar to that of TFAP2A, **Fig. 4B**). Indeed, our IF analyses

for DLX5 in developing hPSC-amnion as well as in a CS7 cynomolgus macaque

embryo show expression patterns similar to TFAP2A (**Fig. 7C,D**). These results further

validate ISL1 as a marker for specified amnion cells and establish that DLX5 as an

additional marker for cells undergoing amniogenesis as well as mature amnion cells.

Functional tests of an immediate-, GATA3, and an early-, TFAP2A, responding

transcription factors

Immediate- and early-responding transcription factors are functionally relevant

candidates that initiate amnion specification and drive the expression of early amnion

lineage specified markers such as ISL1. Among the genes in the immediate- and early-

responding categories, we screened for genes associated with transcription, and

identified several genes, including immediate – GATA3 and early – TFAP2A. As noted

above, GATA3 expression precedes TFAP2A expression by approximately 12 hours.

Both of these factors show highly enriched expression, and both have previously

established functions as pioneer transcription factors (Rothstein and Simoes-Costa,

2020; Takaku et al., 2016; Tanaka et al., 2020; Van Otterloo et al., 2022; Zaret and

Carroll, 2011). Therefore, we examined the role of GATA3 and TFAP2A by generating

hPSC-amnion in Glass-3D[+BMP] derived from H9 cells lacking GATA3 or TFAP2A (two

clonal lines with unique mutations were used for each analysis, see **Fig. S4** for KO

validation). Interestingly, *GATA3*-KO hPSC grown in Glass-3D^{+BMP} develop into fully

squamous cysts that express amnion markers, possibly indicating a functional

redundancy with other expressed GATA genes (e.g., GATA2, data not shown). In

contrast, while controls show uniformly squamous amniotic cells at 48-hr, *TFAP2A*-KO

cysts contain both squamous (ISL1$^+$) and columnar (NANOG$^+$/SOX2$^+$) cell populations

(**Fig. 8A** (Glass-3D hPSC-pluripotent cysts),**8B,C** (Glass-3D^{+BMP} hPSC-amnion cysts)):

these KO cysts display some resemblance to control 24-hr intermediate cysts (**Fig.**

2B,C). These results suggest that amniogenesis may be halted in the absence of

TFAP2A. To examine this further at transcriptomic levels, we performed time-course

bulk RNA-seq for control and *TFAP2A*-KO hPSC-amnion at 0-, 12-, 24- and 48-hr, and

performed a transcriptome similarity analysis using the Tyser et al. human CS7 embryo

dataset. Indeed, while an expected Epiblast to Ectoderm differentiation trajectory similar

to **Fig. 4C** is found in controls (**Fig. 8D**, left), *TFAP2A*-KO samples do not overlap with

Ectoderm cells, even at 48-hr (**Fig. 8D**, right). Rather, 48-hr *TFAP2A*-KO samples show

transcriptomic characteristics similar to some Primitive Streak cells (**Fig. 8D**, uncropped

plots shown in **Fig. S5**). These results reveal a paused amniogenesis phenotype in the

absence of TFAP2A, providing a previously unrecognized notion that TFAP2A is critical

for maintaining amnion fate progression.

DISCUSSION

In this study, we developed a reproducible and faithful model for robust mechanistic

examinations of human amnion fate specification, and identified a BMP4-dependent

amniogenic transcriptional cascade, as well as a role for TFAP2A in maintaining amnion

fate progression. Moreover, we established DLX5 as an additional marker for lineage

progressing amnion cells, and provided evidence for an active amniogenic

transcriptional cascade at the epiblast/amnion boundary in peri-gastrula human and

non-human primate embryos. These findings and the temporally resolved amniogenic

gene lists provided in this study offer additional avenues for further exploration of

human amniogenesis.

Previous studies established several broad amnion markers (e.g., GATA3,

TFAP2A and ISL1) that are expressed in developing, as well as, differentiated amnion

cells. However, comprehensive studies have not been performed to fully characterize

how each of these markers corresponds to amnion fate specification steps, especially

since BMP signaling is a primary trigger of amniogenesis and a large set of BMP target

genes are activated. In this study, a time-course transcriptomic analysis was employed

to identify genes that display temporal-specific dynamics among genes that show more

static expression patterns: this enabled us to define "stage specific" genes that can be

used to distinguish amniogenic fate progression states. In particular, we found that

GATA3 is an immediate BMP target during amniogenesis (along with other known

targets such as *ID1*, *ID2* and *BAMBI*); a similar BMP4 → GATA3 transcriptional pathway

has been well-established previously in other systems (Abe et al., 2021; Bonilla-Claudio

et al., 2012; Gunne-Braden et al., 2020; Vincentz et al., 2016).

Early activation of GATA3 is followed by the expression of second-tier "early"

genes such as *TFAP2A* by 6-hr (clear TFAP2A protein is detected by 12-hr). Our

functional analyses indicate that loss of TFAP2A results in incomplete amniogenesis,

with failure to downregulate pluripotent genes and loss of some differentiated amnion

markers. Thus, TFAP2A, a previously recognized pioneer factor (Rothstein and Simoes-

Costa, 2020; Van Otterloo et al., 2022), appears to be an important component that

ensures the proper progression of the amniogenesis process.

Several hours following the activation of *TFAP2A*, third-tier "intermediate" genes

such as *ISL1* (ISL1 protein by 24-hr: only in squamous cells) are expressed. It has been

previously established that *ISL1* is a key amnion marker in early primate embryos (Yang

et al., 2021; Zhao et al., 2021; Zheng et al., 2022). Interestingly, our IF analyses of early

peri-gastrulation cynomolgus macaque embryos show that ISL1 expression is restricted

to amnion cells that are already squamous in nature (**Fig. 5**). Additionally, our time-

course transcriptomic analyses in Glass-3D^{+BMP} reveal that *ISL1* expression is seen at

12-24-hr, later than *TFAP2A* activation but much earlier than activation of more

differentiated amnion markers such as *GABRP*, *EPAS1* and *IGFBP3* (48-hr). Together,

these results suggest that amnion cells progress from GATA3$^+$, to GATA3$^+$/TFAP2A$^+$, to

GATA3$^+$/TFAP2A$^+$/ISL1$^+$, to mature amnion cells that express these three markers as

well as the more mature amnion markers such as GABRP, EPAS1 and IGFBP3.

The first amniogenesis event is seen upon implantation, in the epiblast cells

underlying the polar trophectoderm, establishing the amniotic sac, an asymmetric

amnion-epiblast cystic structure (Enders et al., 1983; Enders et al., 1986; Shahbazi and

Zernicka-Goetz, 2018; Taniguchi et al., 2017). In this study, we provide evidence for the

active progression of amniogenic fate conversion at the amnion-epiblast boundary of

the cynomolgus macaque peri-gastrula. What is the difference between amniogenesis

that occurs during implantation and during gastrulation? A previous study by Sasaki *et*

al. (Sasaki et al., 2016) showed that, in a macaque embryo staged at PreG1 (just

initiating implantation), a stage before *BMP4* expression is seen in the amnion at PreG2, *BMP2* is abundantly expressed in the visceral endoderm. However, the visceral endoderm underlies the epiblast population that gives rise to the embryo proper (not amnion), and *CER1* is also abundantly expressed in the visceral endoderm (Sasaki et al., 2016). Thus, BMP2 in the visceral endoderm in PreG1 embryo is thought not to be the trigger for the initial amniogenesis processes that forms the amniotic sac, and it is currently unclear how BMP signaling is activated in the nascent amnion cells during implantation *in vivo*. Using a Gel-3D hPSC-amnion system, we previously showed that amniogenesis is mechanosensitive, and that a soft substrate can induce BMP signaling in a mechanically dependent manner in developing hPSC aggregates/cysts. This suggests that mechanical signaling might play an important role in the initial BMP signaling activation and downstream amniogenesis during implantation *in vivo*. In contrast, since a continuous BMP signaling cascade is established in epiblast cells of the early primate gastrula (e.g., driving primitive streak formation, primordial germ cell formation), we speculate that the amniogenesis process that occurs during and post gastrulation may be independent of a mechanical cue. It is, however, important to note that the pluripotency states are distinct in the epiblast cells during implantation compared to during gastrulation; it has been suggested that the epiblast cells are in the formative state during implantation and in the primed state during gastrulation (Kinoshita and Smith, 2018; Shahbazi et al., 2017; Smith, 2017). Thus, it is possible that the mechanisms regulating responses to BMP ligands, as well as BMP-dependent transcriptional machineries may differ.

Overall, our detailed temporally resolved analyses of Glass-3D^{+BMP} hPSC-

amnion development, combined with validation in early macaque embryos has revealed

several previously unrecognized findings with implications for BMP-dependent

amniogenic transcriptional machineries during amniogenesis, mechanisms continuing

amniogenesis in peri-gastrula macaque embryos, and functional characterization of the

transcriptional machinery required for amnion fate specification.

MATERIALS AND METHODS

hESC lines

Human embryonic stem cell line H9 was used in this study (WA09, P30, P48, WiCell;

NIH registration number: 0062). All protocols for the use of hPSC lines were approved

by the Human Stem Cell Research Oversight Committee at the Medical College of

Wisconsin. All hPSC lines were maintained in a feeder-free system for at least 20

passages and authenticated as karyotypically normal at the indicated passage number.

Karyotype analysis was performed at Cell Line Genetics. All hPSC lines tested negative

for mycoplasma contamination (LookOut Mycoplasma PCR Detection Kit, Sigma-

Aldrich). All transgenic and KO hESC lines in this study used H9 as the parental line. In

summary, hESC were maintained in a feeder-free culture system with mTeSR1

medium, or with 50%/50% mix of mTeSR1 and mTeSR plus (STEMCELL

Technologies). hESC were cultured on 1% (v/v) Geltrex (Thermo Fisher Scientific), or

with Cultrex SCQ (Bio-techne) coated 6 well plates (Nunc). Cells were passaged as

small clumps every 4 to 5 days with Dispase (Gibco). All cells were cultured at 37°C

with 5% CO_2. Media was changed every day. hESC were visually checked every day to

ensure the absence of spontaneously differentiated, mesenchymal-like cells in the

culture. Minor differentiated cells were scratched off the plate under a dissecting scope

once identified. The quality of all hESC lines was periodically examined by

immunostaining for pluripotency markers and successful differentiation to three germ

layer cells.

Cynomolgus macaque

Animals

The female and male cynomolgus macaque were housed and cared for at the Wisconsin

National Primate Research Center (WNPRC). All procedures were performed in

accordance with the NIH Guide for the Care and Use of Laboratory Animals and under

approval of the University of Wisconsin College of Letters and Sciences and Vice

Chancellor Office for Research and Graduate Education Institutional Animal Care and

Use Committee (protocol g005061).

Animal breeding and pregnancy detection

Beginning on day 8 post-onset of menses, the female was housed with a compatible

male and monitored for breeding. Animals were pair-housed until day 16-20 post-onset

of menses. A 2-3 mL blood draw was performed daily from day 8 post-onset of menses

until day 16 to assess the timing of ovulation based on the estradiol peak and rise in

progesterone in serum. Serum samples were analyzed for estradiol (25 μL) and

progesterone (20 μL) using a cobas e411 analyzer equipped with

ElectroChemiLuminescence technology (Roche, Basal, Switzerland) according to

manufacturer instructions. Results were determined via a calibration curve which was

instrument-generated by 2-point calibration using traceable standards and a master

curve provided via the reagent barcode. Inter-assay coefficient of variation (CV) was

determined by assaying aliquots of a pool of rhesus plasma. For estradiol, the limit of

quantitation (LOQ) was 25 pg/mL, the intra-assay CV was 2.02%, and the inter-assay

CV was 5.05%. For progesterone, the LOQ was 0.2 ng/mL, the intra-assay CV was

1.37%, and the inter-assay CV was 4.63%. A transabdominal ultrasound was performed

to detect pregnancy as early as 14 days post-ovulation. The ultrasound measurements

in combination with the timing of ovulation were used to estimate the day of conception

and gestational age of the pregnancy.

Terminal perfusion uterine collection, paraffin embedding, sectioning and staining

The pregnant female was sedated with intramuscular ketamine (>15 mg/kg) followed by

IV sodium pentobarbital (>35 mg/kg) and then perfused with 4% paraformaldehyde (PFA)

via the left ventricle. The entire uterus and cervix were removed. The serosa and

superficial myometrium were scored for better fixative penetration and to denote dorsal

and ventral orientation. Tissues were fixed in 4% PFA with constant stirring and solution

changes were performed every 24 hours for a total of 72 h. The uterus was serially

sectioned from the fundus to the internal cervical os into 4 mm slices using a dissection

box. Cassettes with tissue sections were transferred into 70% ethanol, routinely

processed in an automated tissue processor and embedded in paraffin for histological

analyses (5 μm sections). Fixed uterine tissue were cut in 5 μm thick cross-sections,

mounted on slides, deparaffinized in xylene and rehydrated in an ethanol series. Antigen

retrieval was performed by boiling in citrate buffer. Sections were blocked 4% goat serum

in PBS at RT for at least 3-hr. Subsequent immunolocalization was performed using

commercially available primary antibodies, incubated overnight at 4 °C in 4% serum, at

the dilutions as shown in **Table 3**. Immunofluorescent detection was performed using

secondary antibodies tagged with a fluorescent dye (fluorophores excitation = 488, 555,

and 647nm), and counterstained with DAPI. Negative controls were performed in which

the primary antibody was substituted with the same concentration of normal IgG of the

appropriate isotype. Images were obtained with a Zeiss LSM980 microscope.

Glass-3D^{+BMP} hPSC-amnion formation assays

Glass-3D protocol used to generate hPSC-pluripotent cysts was previously described

(Hamed et al., 2023; Shao et al., 2017a; Taniguchi et al., 2017; Taniguchi et al., 2015).

In brief, hPSC monolayers were dissociated using Accutase (Sigma-Aldrich) at 37 °C for

10 min, centrifuged and resuspended in mTeSR1/mTeSR plus 50:50 mix containing 10

µM Y-27632 (STEMCELL Technologies). hPSC were plated as single cells at 20,000

cells cm^{-2}, unless otherwise specified, onto a 1% (v/v) gel coated wells (Geltrex

(Thermo Fisher Scientific), or with Cultrex SCQ (Bio-techne)). After 24 hours (on day 1),

culture medium was replenished with fresh mTeSR1/mTeSR plus 50:50 media without

Y-27632, supplemented with 2% (v/v) Geltrex or with Cultrex SCQ. Thereafter, media

change was performed daily until day 4 (using mTeSR1/mTeSR plus 50:50 media

supplemented with 1% (v/v) Geltrex or with Cultrex SCQ), unless otherwise noted. This

Glass-3D culture condition results in the formation of hPSC-cysts, cysts with a central

lumen surrounded by pluripotent columnar cells. The Glass-3D^{+BMP} culture condition

was used to generate hPSC-amnion, cysts with a central lumen surrounded by

squamous amnion-like cells. To induce amniogenesis, exogenous BMP ligand

(20ng/mL, Bio-techne) was added to developing pluripotent cysts between day 2 to 3 for

24 hours, or between day 2 to 4 for 48 hours, unless otherwise noted (e.g., for time-

course analyses). Samples were harvested at day 4 (48-hr since starting BMP4

treatment). LDN-193189 (500nM, STEMCELL Technologies) was used to disrupt BMP

signaling.

DNA constructs

piggyBac-CRISPR/Cas9 (pBACON) constructs

A piggyBac-CRISPR/Cas9 (pBACON) vector that contains SpCas9-T2A-puro and hU6-

gRNA expression cassettes flanked by piggyBac transposon terminal repeat elements

(pBACON-puro), which allows subcloning of annealed oligos containing gRNA

sequence at *BbsI* site, has been previously described (Shao et al., 2017b; Townshend

et al., 2020; Wang et al., 2021). gRNA targeting genomic sites and oligo sequences to

generate pBACON-GFP-h*GATA3* (primers: CRISPR_hGATA3_CDS1#1_s and

CRISPR_hGATA3_CDS1#1_as) and -h*TFAP2A* (CRISPR_hTFAP2A_CDS2_s and

CRISPR_hTFAP2A_CDS1_as) are found in **Fig. S4** and **Table 3B**; these were

designed using publicly available tools (http://crispr.mit.edu or

https://www.idtdna.com/site/order/designtool/index/CRISPR_CUSTOM).

piggyBac-based transgenic and genome edited hESC lines

PB constructs (3μg) and pCAG-ePBase (Lacoste et al., 2009) (1μg) were co-transfected

into H9 hESC (70,000 cells cm^{-2}) using GeneJammer transfection reagent (Agilent

Technologies). To enrich for successfully transfected cells, drug selection (puromycin,

2μg/mL for 4 days; neomycin, 250μg/mL for 5 days) was performed 48 to 72 hours after

transfection. hESC stably expressing each construct maintained the expression of

pluripotency markers. For inducible constructs, DOX (500ng/mL) treatment was

performed for 48 hours for neomycin-selected *TFAP2A*-KO rescue pools.

During pBACON-based genome editing, puro-selected cells were cultured at low

density (300 cells cm^{-2}) for clonal selection. Established colonies were manually picked

and expanded for screening indel mutations using PCR amplification of a region

spanning the targeted gRNA region (genomic DNA isolated using DirectPCR Lysis

Reagent (Tail) (VIAGEN), primers: hGATA3, PCR_GATA3_RI_fw#2 and

PCR_GATA3_NI_rv#2; hTFAP2A, PCR_TFAP2A_RI_fw and PCR_TFAP2A_NI_rv),

which were subcloned into pPBCAG-GFP (Chen and LoTurco, 2012) at EcoRI and NotI

sites, and sequenced (Seq-3'TR-pPB-Fw). At least 12 to 15 bacterial colonies were

sequenced to confirm genotypic clonality. Western blots for each protein were also

performed to further validate the KO lines. Control cells are H9 hESC in all loss-of-

function experiments.

Western blot

SDS PAGE gels (10%) and PVDF membranes were used. Membranes were blocked

using Intercept (TBS) Blocking Buffer (Licor), total protein quantification was performed

by using Revert 700 Total Protein Stain (Licor), and primary antibody overnight

incubation was performed at 4˚C, followed by 30-min IRDye (Licor) secondary antibody

incubation at room temperature. Blots were imaged using Licor Odyssey Infared

Imaging system.

Immunostaining

hPSC staining protocols have been previously described (e.g., (Hamed et al., 2023;

Shao et al., 2017a; Wang et al., 2021)). In short, samples (hPSC-cysts, hPSC-amnion)

grown on the coverslip were rinsed with phosphate-buffered saline (PBS; Gibco) twice,

fixed with 4% paraformaldehyde (Sigma-Aldrich) in 1x PBS for 40 to 60 min, then rinsed

with PBS three times, and permeabilized with 0.1% SDS (Sigma-Aldrich) solution for 40

min. The samples were blocked in 4% heat-inactivated goat serum (Gibco) or 4%

normal donkey serum (Gibco) in PBS 1 hour to overnight at 4°C. The samples were

incubated with primary antibody solution prepared in blocking solution at 4°C overnight,

washed three times with PBS (10 min each), and incubated in blocking solution with

goat or donkey raised Alexa Fluor–conjugated

secondary antibodies (Thermo Fisher Scientific) at room temperature for 2 hours. DNA

counterstaining was performed using Hoechst 33342 (nucleus, Thermo Fisher

Scientific), Alexa Fluor–conjugated WGA (membrane, Thermo Fisher Scientific). All

samples were mounted on slides using 90% glycerol (in 1x PBS). When mounting cyst

samples, modeling clays were used as spacers between coverslips and slides to

preserve lumenal cyst morphology.

Confocal microscopy, live imaging and image analysis

Confocal Images of fixed samples were acquired using Zeiss LSM980 microscope.

Measurements of cell nucleus dimensions, cyst epithelial thickness, and cyst orientation

angle were performed manually using the Measurement tool in ImageJ (NIH).

RNA isolation and qRT-PCR analysis

Total RNA extraction was performed using TRIzol (Thermo Fisher). The quantity and

quality of total RNA was determined by spectrometry and denaturing agarose gel

electrophoresis, respectively. The cDNA was synthesized from total RNA ($1\mu g$) using

High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems). Quantitative real-

time RT-PCR (qPCR) analysis of mRNA expression was performed using a CFX96

Real-Time PCR Detection System (BioRad) with Power Up™ SYBR Green Master Mix

(Applied Biosystems). Primers (**Table 3B**) were designed using NCBI primer design tool

to span at least one exon/intron junction. Data were normalized against *GAPDH* and are

shown as the average fold increase ± standard error of the mean (SEM) with relative

expression being calculated using the 2-ΔΔCT method as described previously (Livak

and Schmittgen, 2001). After amplification, the specificity of the PCR was determined by

both melt-curve analysis and gel electrophoresis to verify that only a single product of

the correct size was present.

RNA sequencing and bioinformatics

The quality of total RNA was determined using TapeStation RNA ScreenTape. RNA

was enriched for mRNA using NEBNext Poly(A) mRNA Magnetic Isolation Module

(E7490). cDNA and library preparation were performed using NEBNext® Ultra™ II

Directional RNA Library Prep kit for Illumina (New England Biolabs). Library

concentration was determined using NEBNext® Library Quant Kit for Illumina (New

England Biolabs) or by Qubit (Thermo). Samples were then pooled and were

sequenced using Illumina sequencers: NextSeq550 (Versiti Blood Research Institute) or

using NovaSeq 6000 (S4 flowcell, the DNA services laboratory of the Roy J. Carver

Biotechnology Center at the University of Illinois at Urbana-Champaign). Sequence data

was analyzed using the RNA-Seq workflow in Partek Flow software (Partek Inc., St.

Louis, MO, USA). Raw sequencing reads were mapped to the human genome hg38

using STAR version 2.7.8a with default parameter settings. Aligned reads were then

quantified to Ensembl Transcripts release 104 using Partek's method and genes with

fewer than 10 reads in all samples were excluded from the subsequent analyses. Gene

counts were then filtered based on gene_biotype for protein_coding genes, followed by

excluding features where maximum <= 30.0 reads. Gene expression was then

normalized using DESeq2 (Median Ratio). Differentially expressed genes were selected

using as criteria FDR < 0.05 and absolute fold change > 2.

GO enrichment analysis

GO analysis were performed on significantly upregulated (1,820) or downregulated

(2,092) genes in Glass-3D^{+BMP} amnion-cysts vs Glass-3D pluripotent-cysts (From **Fig.**

3).

Mapping and gene expression analyses of Tyser et al. Carnegie Stage 7 human embryo

single cell RNA sequencing dataset

The original UMAP coordinates and annotations for all 1,195 cells were used

throughout. The processed data were downloaded from http://www.human-gastrula.net,

which was used to generate a UMAP plot (using DimPlot in R package Seurat), as well

as to perform expression analyses (FeaturePlot in Seurat).

RNA velocity (scvelo) analysis of Tyser et al. Carnegie Stage 7 human embryo single

cell RNA sequencing dataset

RNA velocities across all 1,195 single cells were embedded on the original UMAP

coordinates shown in Tyser *et al.* (Tyser et al., 2021). The Tyser et al. Carnegie Stage 7

human embryo raw fastq dataset was obtained from ArrayExpress (accession code: E-

MTAB-9388). From the aligned BAM files, a loom file was generated using the function

run_smartseq2 mode in software velocyto (v0.17) with default parameters to create a

count matrix made of spliced and unspliced read counts (with Human genome

annotation hg19). scVelo (Python package v.0.2.5 (Bergen et al., 2020) was used to

examine inferred RNA velocity and visualized the velocities on the UMAP embedding by

stream plot (with default parameters except smooth was set to 1).

Transcriptomic similarity analysis between bulk RNA sequencing and Tyser et al.

Carnegie Stage 7 human embryo single cell RNA sequencing datasets

Transcriptomic similarities between bulk RNA sequencing samples and the Tyser *et al.*

dataset were examined and visualized using original UMAP coordinates from Tyser *et*

al.. In summary, each bulk RNA-seq sample was first downsampled to 100 pseudo-

single cell data using R package projectLSI (Granja et al., 2019), which was followed by

data integration with Tyser *et al.* dataset in UMAP dimension reduction space. A medoid

from 100 pseudo-single cell data was used as the representation of each bulk RNA-seq

sample, and, for each sample, we identified the most transcriptomically similar cell from

Tyser *et al.* dataset, based on Euclidean distance on UMAP space.

Data availability

The raw data, unfiltered count matrix and processed count matrix will be available to the

database of Genotypes and Phenotypes (dbGaP) upon publication.

Statistical analyses

Graphs were generated using Prism 7 (GraphPad Software) or R,

and statistical analyses were performed using Prism 7. Fig. 1D, 5 independent samples

were counted at each timepoint (total of > 50 cells/sample). For analysis we used one-

way ANOVA with post-hoc Dunnet's multiple comparisons test. In Fig. 1K, 3

independent samples were counted (total of > 75 cells were plotted). For analysis we

used two-tail Student-T test. In Fig. 1J, 3 independent samples were counted (total of >

50 cells were plotted). For analysis we used one-way ANOVA with post-hoc Tukey's

multiple comparison test. In Fig. 2, relative mRNA expression was detected in n=3 for

each timepoint. For analysis we used one-way ANOVA with post-hoc Tukey's multiple

comparison test. *, significant (p-value<0.05); ns, non-significant. All experiments were

repeated at least three times.

FIGURE LEGENDS

Figure 1. Highly penetrant amniogenesis in a Glass-3D^{+BMP} culture system. (**A,B**) hPSC-derived pluripotent cyst (hPSC-pluripotent cyst, Glass-3D in (**A**)) and amnion cyst (Glass-3D^{+BMP} in (**B**)) assays. BMP4 treatment (20ng/mL) leads to uniformly squamous amnion cysts by Day 4. (**C**) Confocal optical sections of d2 cysts harvested prior to (0-min) and 10-, 20-, 30- and 60-min after BMP4 treatment, stained with indicated markers. Staining for phosphorylated SMAD1/5 reveals a prominent nuclear pSMAD1/5 enrichment within 10-min of BMP4 treatment in all cells. (**D**) Quantification of nuclear pSMAD1/5: five independent samples were counted, more than 50 cells were counted per sample. * indicates statistically significant changes (p<0.05) compared to 0-min timepoint. (**E**) Abundant GATA3 expression (in green), but not TFAP2A (red), is seen by 3 hours after BMP4 treatment while SOX2 expression is maintained. (**F-H**) Pluripotency (NANOG, SOX2) and amnion (ISL1/2, GATA3, TFAP2A) marker expression analyses of d4 cysts grown in the Glass-3D (top) or in Glass-3D^{+BMP} (bottom) conditions. (**I,J**) Confocal micrographs of d4 cysts grown in the Glass-3D^{+BMP} system stained for phosphorylated EZRIN, RADIXIN and MOESIN (pERM), an apical membrane marker, as well as with indicated amnion markers (top – cysts were treated with BMP4 for 24 hours between d2 and d3; bottom – treated with BMP4 for 48 hours from d2 to d4). (**K,L**) Quantification of GATA3 and ISL1/2 fluorescent intensities in 24- (blue, n = 115 cells) and 48-hr (red, n = 76 cells) BMP4 treatment samples (**K**), as well as of nuclear aspect ratios of d4 hPSC-cyst (black, n = 63 cells), and 24- (blue, n = 45 cells) and 48-hr (red, n = 52) BMP4-treated cysts (harvested at d4), (**L**). (**M**) pSMAD1/5 staining (green) shows similar nuclear enrichment between 24- and 48-hr BMP4 treatment at d4 (also stained with indicated markers). Scale bars = 50μm.

Figure 2. Time-course analysis of Glass-3D^+BMP hPSC-amnion development. (A) Timeline of the Glass-3D^+BMP hPSC-amnion assay. Twenty four-hr BMP treatment is sufficient to generate uniformly squamous hPSC-amnion cysts. (**B,C**) Optical sections of time-course Glass-3D^+BMP4 hPSC-amnion cysts (harvested at indicated timepoints after BMP4 treatment) stained with indicated markers (inverted black and white fluorescent signals shown for individual channels, to aid visualization). GATA3, an immediate BMP target is seen by 3-hr (yellow arrowheads). hPSC-amnion cysts with prominent nuclear TFAP2A (**B**) and ISL1/2 (**C**) are seen by 12- and 24-hr, respectively (yellow arrowheads), while SOX2 shows gradual decrease. Consistent changes are also observed at mRNA transcript levels using quantitative RT-PCR (**D-F**, also see the expression of additional markers: *HAND1* (amniotic) and *NANOG* (pluripotent); * indicates statistically significant changes (p<0.05) compared to 0-hr timepoint). (**G**) Relative mRNA expression analyses of indicated pluripotency (top) and amniotic (bottom) markers between d4 Glass-3D hPSC-pluripotent cyst (black), and d4 Glass-3D^+BMP hPSC-amnion with 24-hr (blue) or 48-hr (red) BMP treatment; * indicates statistically significant changes (p<0.05). (**H**) BMP inhibitor (LDN-193189) treatment outline as well as confocal micrographs of d4 Glass-3D^+BMP hPSC-amnion (top, control – 24-hr BMP4 treatment between d2 and d3) and hPSC-amnion treated with LDN-193189 between d3 and d4 after BMP4 treatment (between d2-d3). Scale bars = 50μm.

Figure 3. Temporally-resolved transcriptomic characterization of Glass-3D^+BMP hPSC-amnion. (A) Experimental timeline (from d2) of the time-course bulk RNA

sequencing analysis. Colored arrowheads indicate timepoints at which samples were

harvested. (**B**) A heatmap showing a hierarchical clustering analysis of all detected

genes (12,940): bulk transcriptomes from triplicate samples are shown at each

timepoint. A gradient scale indicates Z-score. (**C**) Principal component analysis (PCA)

plot of time-course bulk transcriptomes, using the most variable genes, revealing

progressive transcriptomic changes overtime (indicated by the dotted arrow). Note that

coloring is consistent with the arrowheads in (**A**). (**D-G**) Time-course plots of normalized

expression values for selected known immediate BMP target (**D**), general amnion (**E**),

advanced amnion (**F**) and pluripotency genes (**G**). (**H**) A volcano plot analysis showing

up- (red) and down- (blue) regulated genes in Glass-3D^{+BMP} hPSC-amnion cysts (gene

expression compared to d4 pluripotent hPSC-cysts): black lines represent cutoffs for p-

value <0.05 and 2-fold difference.

Figure 4. Transcriptome comparison of developing Glass-3D^{+BMP} hPSC-amnion

cysts and a Carnegie stage (CS) 7 human embryo. (**A**) A UMAP plot displaying the

original single cell transcriptome coordinates of the CS7 human embryo described in

Tyser *et al.*, shown with the original annotations. The inset (**i**) is used for expression

analyses in (**B**). (**B**) Expression of amnion markers – *GATA3*, *TFAP2A* and *ISL1* –

superimposed onto the UMAP plot in (**A** – **i**). (**C**) Transcriptome similarity analysis

comparing the CS7 human embryo single cell dataset and the Glass-3D^{+BMP} time-

course bulk RNA-seq dataset, showing the human CS7 embryo UMAP plot (inset) with

colored cells that are transcriptomically similar to each timepoints as indicated (orange –

pluripotent cysts/monolayer and 0-, 0.5-, 1-, 3- and 6-hr post-BMP4; green – 12-hr post-

BMP4; blue – 24-hr; magenta – 48-hr). While pluripotent and earlier timepoints (orange

colored cells) overlaps with Epiblast cells, some 12-hr (green) and all 24-hr (blue)

samples are similar to a subpopulation of Primitive Streak cells, and 48-hr (magenta)

samples overlap exclusively with Ectoderm (see Materials and Methods for detailed

bioinformatics pipeline). (**D**) *TBXT* expression transposed onto the CS7 human embryo

UMAP plot, revealing *TBXT*low Primitive Streak sub-population that show transcriptomic

similarities to 24-hr post-BMP samples (indicated by dotted circle). Gradient scales

indicate expression level (orange = high, gray = low). Uncropped plots are found in **Fig.**

S2.

Figure 5. Transcriptional characterizations of amnion-epiblast squamocolumnar

boundary of the CS6/7 cynomolgus monkey amniotic sac. (**A**) RNA velocity

analysis (scvelo) of the *Tyser et al.* CS7 human embryo dataset, showing predicted

lineage progression trajectories from Epiblast to Ectoderm traversing the *TBXT*low

Primitive Streak sub-population. (**B-G**) Optical sections of a CS6/7 cynomolgus

macaque embryo stained for indicated transcription factors (GATA3, ISL1/2, pSMAD1/5

and SOX2 (**B**,**D**,**F**)), as well as quantitation for relative fluorescent intensity in proximal

(closer to epiblast) to distal (closer to amnion) cells of indicated markers at the amnion-

epiblast boundary (**C**, **E** and **G** are quantitation of images in **B**, **D** and **F**, respectively).

Each dot represents a single cell. (**i**), (**ii**) and (**iii**) indicate insets in (**F**). Half of the

amniotic sac shown. Quantitation shows that, while GATA3+/TFAP2A+ cells are also

SOX2+, ISL1+ cells are only seen after SOX2 expression is no longer detected: ISL1

expression is exclusive to squamous amnion cells. Moreover, the expression of

pSMAD1/5 and TFAP2A is reduced in most distal amnion (**ii**, **iii**). Scale bars 200μm. (**H**)

Nuclear aspect ratio quantitation of amnion-epiblast boundary cells in (**B**), (**C**) and (**D**),

mean+SEM shown. Dotted vertical line in each plot indicates amnion-epiblast boundary

(**C,E,G,H**). (**I**) A schematic representation of the CS6/7 cynomolgus macaque amniotic

sac: the confocal micrographs in (**B**) were edited to represent SOX2$^+$ epiblast

(pseudocolored in blue), primitive streak-derived disseminating (green),

GATA3$^+$/TFAP2A$^+$/SOX2^{low} transition boundary (magenta) and GATA3$^+$/TFAP2A$^+$/ISL1$^+$

lineage specified amnion (red) cells.

Figure 6. Characterization of differentially expressed genes identified through the

Glass-3D^{+BMP} hPSC-amnion time-course sequencing analysis. (**A-E**) Combined and

selected genes are shown for each category (A – immediate; B – early; C –

intermediate-1; D – intermediate-2; E – late). Genes for each category are selected

based on significant upregulation (FDR<0.05, fold>2, based on Z-scores) by 1-

(immediate, n = 5), 3- (early, n = 9), 6- to 12- (intermediate-1, n = 30), 24- (intermediate-

2, n = 178) and 48-hr (late, n = 157) post-BMP4 treatment, compared to the previous

timepoint. Vertical lines indicate timepoints at which significant changes are seen.

Figure 7. Establishment of DLX5 as a new marker of amniogenesis similar to

TFAP2A. (**A,B**) Expression analyses for *ISL1* (**A**) and *DLX5* (**B**) in the time-course bulk

RNA sequencing (top) and in the single cell CS7 human embryo (bottom) datasets. (**C**)

Confocal optical sections of Glass-3D hPSC-pluripotent cysts (left) and Glass-3D^{+BMP4}

hPSC-amnion cysts (right, harvested at d4) stained with indicated markers. (**D**)

Histological section of a cynomolgus monkey amniotic sac at CS7 stained with indicated

markers (SOX2$^+$ epiblast; TFAP2A$^+$/DLX5$^+$/SOX2$^-$ amnion). Insets (**i** and **ii**) indicate the

amnion-epiblast boundaries. DLX5 and TFAP2A share a similar expression pattern at

the boundary. Similar to pSMAD1/5 and TFAP2A (shown in **Fig. 5F**), most distal amnion

cells show reduced DLX5 expression. Surface ectoderm bridging neural ectoderm and

amniotic ectoderm is not seen at this stage. Amniotic mesenchyme is observed

underlying the TFAP2A$^+$/DLX5$^+$ amniotic epithelium at this stage. Scale bars = 500μm.

Figure 8. Analyses of Glass-3D^{+BMP} hPSC-amnion in control and *TFAP2A*-KO

background. (**A,B**) Optical sections of control and *TFAP2A*-KO cysts in Glass-3D (**A**,

pluripotent) as well as in Glass-3D^{+BMP} (**B**). While control-like pluripotent cysts are

formed in Glass-3D, several NANOG$^+$ cells are still seen in Glass-3D^{+BMP} in the

absence of *TFAP2A*. (**C**) Confocal micrographs of d4 control and *TFAP2A*-KO Glass-

3D^{+BMP} hPSC-amnion cysts stained with indicated markers, showing a d3-like

intermediate morphology with ISL1$^+$/SOX2$^-$ and ISL1$^-$/SOX2$^+$ cells. (**D**) Transcriptome

similarity analysis comparing the CS7 human embryo single cell dataset with the time-

course (0-, 12-, 24- and 48-hr post-BMP) bulk dataset from control and *TFAP2A*-KO

Glass-3D^{+BMP} hPSC-amnion. Orange, green, blue and magenta colored cells are

transcriptomically similar to 0-, 12-, 24- and 48-hr post-BMP bulk RNA sequencing

samples, respectively. While a clear differentiation trajectory from Epiblast to Ectoderm

is seen in control (left), a halted amnion lineage progression is seen in the *TFAP2A*-KO

background. Scale bars = 50μm.

Table 1. Glass-3D^{+BMP} hPSC-amnion time-course bulk RNA sequencing dataset.

(**A**) normalized and filtered count matrix of the time-course bulk-RNA sequencing. (**B-C**)

Gene expression comparison analysis between 48-hr (d4) Glass-3D^{+BMP} hPSC-amnion

and d4 Glass-3D hPSC-cysts: lists of 1,813 up-regulated (**B**) and 2,088 down- (**C**)

regulated genes. (**D-E**) Gene Ontology analyses (Biological Processes) for top all

significantly up- (**D**) and down- (**E**) regulated genes.

Table 2. Lists of immediate, early, intermediate and late genes. An excel file

containing temporally resolved gene lists: (**A**) immediate (0.5- and 1-hr, 57 genes); (**B**)

early (3- and 6-hr, 135 genes); (**C**) intermediate-1 (12-hr, 81 genes); (**D**) intermediate-2

(24-hr, 642 genes); (**E**) late (48-hr, 909 genes). Genes that showed significant

upregulation compared to the previous timepoint (e.g., 0.5-hr compared to 0-hr, 3-hr

compared to 1-hr) were retained. Duplicated genes found in the list from earlier

timepoint lists are removed. For example, *GATA3* is significantly upregulated throughout

the time course, but is only included in the immediate category. Genes with averaged

normalized counts < 100 were excluded.

Table 3. Lists of reagents used in this study. (**A-C**) Lists of antibodies (**A**), primers

(**B**) and plasmids (**C**).

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. Glass-3D⁺ᴮᴹᴾ hPSC-amnion formation using 1196a. Confocal

micrographs of 1196a human induced pluripotent stem cell line grown in the Glass-

3D⁺ᴮᴹᴾ condition, stained with indicated markers. Scale bars = 50μm.

Figure S2. Original UMAP plots. (**A**) UMAP expression plots of the Tyser *et al.* CS7

human embryo single cell RNA sequencing dataset used in this study (*GATA3*,

TFAP2A, *ISL1, TBXT* and *DLX5*) as well as (**B**) time-course transcriptome similarity

UMAP plot.

Figure S3. Quantitation for relative fluorescent intensity in proximal and distal

cells of indicated markers at the amnion-epiblast boundary in a CS7 cynomolgus

macaque embryo.

Figure S4. Validation of *GATA3*- and *TFAP2A*-KO hPSC lines. (**A,B**) Sequenced

genotyping results of *GATA3*-KO #1 and #2 lines (**A**), representative sequence traces

for each mutations are shown in (**B**). (**C-E**) Genotyping results of *TFAP2A*-KO #1 and

#2 lines (**C**), representative sequence traces for each mutation are shown in (**D**).

Western blot analysis for TFAP2A in control and *TFAP2A*-KO #1 and #2 lysates (**E**, β-

ACTIN is used as a loading control).

Figure S5. Uncropped transcriptional similarity UMAP plots for control and

***TFAP2A*-KO shown in Fig. 8D.**

ACKNOWLEDGEMENTS

We thank A. Brivanlou (Rockefeller University) for ePB transposase. We thank Michaela

Patterson, Matthew Veldman and Deborah Gumucio for their insightful comments to the

manuscript, as well as Versiti Blood Research Institute Molecular Biology Core Lab and

the DNA Services laboratory of the Roy J. Carver Biotechnology Center at the

University of Illinois at Urbana-Champaign for sequence services. We also thank the

Wisconsin National Primate Research Center (WNPRC) Veterinary, Scientific Protocol

Implementation, Pathology and Animal Services Unit staff for providing animal care, and

assisting in procedures including breeding, pregnancy monitoring, and sample

collection. The contents of this manuscript are solely the responsibility of the authors

and do not represent the official views of the NIH.

Funding: This work was supported by NIH grants R01-HD098231 (K.T.) and P51-

OD011106 (to the WNPRC) as well as by MCW CBNA Start-up funds (K.T.).

Competing interests: The authors declare no competing interest. **Data and materials**

availability: All data needed to evaluate the conclusions in the paper are present in the

paper and the Supplementary Materials. Further information and requests for reagents

may be directed to Kenichiro Taniguchi.

896 <u>REFERENCES</u>

897 Abe, M., Cox, T.C., Firulli, A.B., Kanai, S.M., Dahlka, J., Lim, K.C., Engel, J.D., and

898 Clouthier, D.E. (2021). GATA3 is essential for separating patterning domains during

899 facial morphogenesis. Development *148*.

Beddington, R.S., and Robertson, E.J. (1999). Axis development and early asymmetry in mammals. Cell *96*, 195-209.

Bergen, V., Lange, M., Peidli, S., Wolf, F.A., and Theis, F.J. (2020). Generalizing RNA velocity to transient cell states through dynamical modeling. Nature biotechnology *38*, 1408-1414.

Bergmann, S., Penfold, C.A., Slatery, E., Siriwardena, D., Drummer, C., Clark, S., Strawbridge, S.E., Kishimoto, K., Vickers, A., Tewary, M*., et al.* (2022). Spatial profiling of early primate gastrulation in utero. Nature *609*, 136-143.

Bonilla-Claudio, M., Wang, J., Bai, Y., Klysik, E., Selever, J., and Martin, J.F. (2012). Bmp signaling regulates a dose-dependent transcriptional program to control facial skeletal development. Development *139*, 709-719.

Carleton, A.E., Duncan, M.C., and Taniguchi, K. (2022). Human epiblast lumenogenesis: From a cell aggregate to a lumenal cyst. Seminars in cell & developmental biology *131*, 117-123.

Chen, F., and LoTurco, J. (2012). A method for stable transgenesis of radial glia lineage in rat neocortex by piggyBac mediated transposition. Journal of neuroscience methods *207*, 172-180.

Chen, H.M., DeLong, C.J., Bame, M., Rajapakse, I., Herron, T.J., McInnis, M.G., and O'Shea, K.S. (2014). Transcripts involved in calcium signaling and telencephalic neuronal fate are altered in induced pluripotent stem cells from bipolar disorder patients. Translational psychiatry *4*, e375.

Chen, K., Zheng, Y., Xue, X., Liu, Y., Resto Irizarry, A.M., Tang, H., and Fu, J. (2021).

Branching development of early post-implantation human embryonic-like tissues in 3D

stem cell culture. Biomaterials *275*, 120898.

Enders, A.C., Hendrickx, A.G., and Schlafke, S. (1983). Implantation in the rhesus

monkey: initial penetration of endometrium. The American journal of anatomy *167*, 275-

298.

Enders, A.C., Schlafke, S., and Hendrickx, A.G. (1986). Differentiation of the embryonic

disc, amnion, and yolk sac in the rhesus monkey. The American journal of anatomy

177, 161-185.

Granja, J.M., Klemm, S., McGinnis, L.M., Kathiria, A.S., Mezger, A., Corces, M.R.,

Parks, B., Gars, E., Liedtke, M., Zheng, G.X.Y.*, et al.* (2019). Single-cell multiomic

analysis identifies regulatory programs in mixed-phenotype acute leukemia. Nature

biotechnology *37*, 1458-1465.

Gunne-Braden, A., Sullivan, A., Gharibi, B., Sheriff, R.S.M., Maity, A., Wang, Y.F.,

Edwards, A., Jiang, M., Howell, M., Goldstone, R.*, et al.* (2020). GATA3 Mediates a

Fast, Irreversible Commitment to BMP4-Driven Differentiation in Human Embryonic

Stem Cells. Cell stem cell *26*, 693-706 e699.

Hamed, M.M., Taniguchi, K., and Duncan, M.C. (2023). Monitoring Effects of Membrane

Traffic Via Changes in Cell Polarity and Morphogenesis in Three-Dimensional Human

Pluripotent Stem Cell Cysts. Methods in molecular biology *2557*, 83-98.

Kinoshita, M., and Smith, A. (2018). Pluripotency Deconstructed. Development, growth

& differentiation *60*, 44-52.

Lacoste, A., Berenshteyn, F., and Brivanlou, A.H. (2009). An efficient and reversible

transposable system for gene delivery and lineage-specific differentiation in human

embryonic stem cells. Cell stem cell *5*, 332-342.

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data

using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402-

408.

Miki, T., Lehmann, T., Cai, H., Stolz, D.B., and Strom, S.C. (2005). Stem cell

characteristics of amniotic epithelial cells. Stem cells *23*, 1549-1559.

Miki, T., and Strom, S.C. (2006). Amnion-derived pluripotent/multipotent stem cells.

Stem Cell Rev *2*, 133-142.

Nakamura, T., Fujiwara, K., Saitou, M., and Tsukiyama, T. (2021). Non-human primates

as a model for human development. Stem cell reports *16*, 1093-1103.

Nakamura, T., Okamoto, I., Sasaki, K., Yabuta, Y., Iwatani, C., Tsuchiya, H., Seita, Y.,

Nakamura, S., Yamamoto, T., and Saitou, M. (2016). A developmental coordinate of

pluripotency among mice, monkeys and humans. Nature *537*, 57-62.

Nakamura, T., Yabuta, Y., Okamoto, I., Sasaki, K., Iwatani, C., Tsuchiya, H., and

Saitou, M. (2017). Single-cell transcriptome of early embryos and cultured embryonic

stem cells of cynomolgus monkeys. Scientific data *4*, 170067.

Nasr Esfahani, S., Shao, Y., Resto Irizarry, A.M., Li, Z., Xue, X., Gumucio, D.L., and Fu,

J. (2019). Microengineered human amniotic ectoderm tissue array for high-content

developmental phenotyping. Biomaterials *216*, 119244.

Rothstein, M., and Simoes-Costa, M. (2020). Heterodimerization of TFAP2 pioneer

factors drives epigenomic remodeling during neural crest specification. Genome Res

30, 35-48.

Sasaki, K., Nakamura, T., Okamoto, I., Yabuta, Y., Iwatani, C., Tsuchiya, H., Seita, Y.,

Nakamura, S., Shiraki, N., Takakuwa, T*., et al.* (2016). The Germ Cell Fate of

Cynomolgus Monkeys Is Specified in the Nascent Amnion. Developmental cell *39*, 169-

185.

Shahbazi, M.N., Scialdone, A., Skorupska, N., Weberling, A., Recher, G., Zhu, M.,

Jedrusik, A., Devito, L.G., Noli, L., Macaulay, I.C*., et al.* (2017). Pluripotent state

transitions coordinate morphogenesis in mouse and human embryos. Nature *552*, 239-

243.

Shahbazi, M.N., and Zernicka-Goetz, M. (2018). Deconstructing and reconstructing the

mouse and human early embryo. Nature cell biology.

Shao, Y., and Fu, J. (2022). Engineering multiscale structural orders for high-fidelity

embryoids and organoids. Cell stem cell *29*, 722-743.

Shao, Y., Taniguchi, K., Gurdziel, K., Townshend, R.F., Xue, X., Yong, K.M.A., Sang, J.,

Spence, J.R., Gumucio, D.L., and Fu, J. (2017a). Self-organized amniogenesis by

human pluripotent stem cells in a biomimetic implantation-like niche. Nature materials

16, 419-425.

Shao, Y., Taniguchi, K., Townshend, R.F., Miki, T., Gumucio, D.L., and Fu, J. (2017b).

A pluripotent stem cell-based model for post-implantation human amniotic sac

development. Nature communications *8*, 208.

Smith, A. (2017). Formative pluripotency: the executive phase in a developmental

continuum. Development *144*, 365-373.

Takaku, M., Grimm, S.A., Shimbo, T., Perera, L., Menafra, R., Stunnenberg, H.G.,

Archer, T.K., Machida, S., Kurumizaka, H., and Wade, P.A. (2016). GATA3-dependent

cellular reprogramming requires activation-domain dependent recruitment of a

chromatin remodeler. Genome Biol *17*, 36.

Tanaka, H., Takizawa, Y., Takaku, M., Kato, D., Kumagawa, Y., Grimm, S.A., Wade,

P.A., and Kurumizaka, H. (2020). Interaction of the pioneer transcription factor GATA3

with nucleosomes. Nature communications *11*, 4136.

Taniguchi, K., Heemskerk, I., and Gumucio, D.L. (2019). Opening the black box: Stem

cell-based modeling of human post-implantation development. The Journal of cell

biology *218*, 410-421.

Taniguchi, K., Shao, Y., Townshend, R.F., Cortez, C.L., Harris, C.E., Meshinchi, S.,

Kalantry, S., Fu, J., O'Shea, K.S., and Gumucio, D.L. (2017). An apicosome initiates

self-organizing morphogenesis of human pluripotent stem cells. The Journal of cell

biology *216*, 3981-3990.

Taniguchi, K., Shao, Y., Townshend, R.F., Tsai, Y.H., DeLong, C.J., Lopez, S.A.,

Gayen, S., Freddo, A.M., Chue, D.J., Thomas, D.J.*, et al.* (2015). Lumen Formation Is

an Intrinsic Property of Isolated Human Pluripotent Stem Cells. Stem cell reports *5*, 954-

962.

Townshend, R.F., Shao, Y., Wang, S., Cortez, C.L., Esfahani, S.N., Spence, J.R.,

O'Shea, K.S., Fu, J., Gumucio, D.L., and Taniguchi, K. (2020). Effect of Cell Spreading

on Rosette Formation by Human Pluripotent Stem Cell-Derived Neural Progenitor Cells.

Front Cell Dev Biol *8*, 588941.

Tyser, R.C.V., Mahammadov, E., Nakanoh, S., Vallier, L., Scialdone, A., and Srinivas,

S. (2021). Single-cell transcriptomic characterization of a gastrulating human embryo.

Nature *600*, 285-289.

Van Otterloo, E., Milanda, I., Pike, H., Thompson, J.A., Li, H., Jones, K.L., and Williams,

T. (2022). AP-2alpha and AP-2beta cooperatively function in the craniofacial surface

ectoderm to regulate chromatin and gene expression dynamics during facial

development. eLife *11*.

Vincentz, J.W., Casasnovas, J.J., Barnes, R.M., Que, J., Clouthier, D.E., Wang, J., and

Firulli, A.B. (2016). Exclusion of Dlx5/6 expression from the distal-most mandibular

arches enables BMP-mediated specification of the distal cap. Proceedings of the

National Academy of Sciences of the United States of America *113*, 7563-7568.

Wang, S., Lin, C.W., Carleton, A.E., Cortez, C.L., Johnson, C., Taniguchi, L.E.,

Sekulovski, N., Townshend, R.F., Basrur, V., Nesvizhskii, A.I.*, et al.* (2021). Spatially

resolved cell polarity proteomics of a human epiblast model. Sci Adv *7*.

Yang, R., Goedel, A., Kang, Y., Si, C., Chu, C., Zheng, Y., Chen, Z., Gruber, P.J., Xiao,

Y., Zhou, C.*, et al.* (2021). Amnion signals are essential for mesoderm formation in

primates. Nature communications *12*, 5126.

Zaret, K.S., and Carroll, J.S. (2011). Pioneer transcription factors: establishing

competence for gene expression. Genes & development *25*, 2227-2241.

Zhao, C., Reyes, A.P., Schell, J.P., Weltner, J., Ortega, N.M., Zheng, Y., Björklund,

Å.K., Rossant, J., Fu, J., Petropoulos, S.*, et al.* (2021). Reprogrammed blastoids

contain amnion-like cells but not trophectoderm. bioRxiv, 2021.2005.2007.442980.

Zheng, Y., Xue, X., Shao, Y., Wang, S., Esfahani, S.N., Li, Z., Muncie, J.M., Lakins,

J.N., Weaver, V.M., Gumucio, D.L.*, et al.* (2019). Controlled modelling of human

epiblast and amnion development using stem cells. Nature *573*, 421-425.

Zheng, Y., Yan, R.Z., Sun, S., Kobayashi, M., Xiang, L., Yang, R., Goedel, A., Kang, Y.,

Xue, X., Esfahani, S.N.*, et al.* (2022). Single-cell analysis of embryoids reveals lineage

diversification roadmaps of early human development. Cell stem cell *29*, 1402-1419

e1408.

1039

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A ISL1

B DLX5

Tyser *et al.*
CS7 embryo

Tyser *et al.*
CS7 embryo

C Glass-3D Glass-3D +BMP4

DLX5

TFAP2A

SOX2

MERGE

D CS7

DLX5 TFAP2A SOX2

i

ii

Figure 8