Why not both? A case study measuring cortisol and corticosterone in poison frogs

Sarah E. Westrick [a], Ryan T. Paitz [b], and Eva K. Fischer [a]

[a] Department of Evolution, Ecology and Behavior, University of Illinois Urbana-Champaign, 505

S Goodwin Ave Urbana, IL 61801

[b] School of Biological Sciences, Illinois State University, Campus Box 4720 Normal, IL 61790

Corresponding author: Sarah E. Westrick, westse@illinois.edu

Abstract

A general tenet in stress physiology is that the hypothalamic-pituitary-adrenal (HPA) axis

predominantly produces one glucocorticoid (GC) in response to stressors. Researchers typically

measure only the GC believed to be dominant in their species of interest; however, the two main

glucocorticoids produced by vertebrates– cortisol and corticosterone — show variation across

species in absolute levels, relative proportions, and stress responsivity. Therefore, generalizing

assumptions and focusing on a single GC may not tell the whole story. In this study, we

measured both cortisol and corticosterone in water samples collected from five species of

Dendrobatid poison frogs. Understanding the full picture of the physiological stress response in

poison frogs is of special interest due to their susceptibility to environmental stressors and

conservation status. Counter to the dogma that corticosterone is dominant and more abundant in

amphibians, we found cortisol was more abundant than corticosterone in four of the five species.

Dendrobates tinctorius was unique in releasing equal amounts of cortisol and corticosterone.

When we stimulated the HPA axis in *D. tinctorius*, we found corticosterone was responsive to

adrenocorticotropic hormone (ACTH), while cortisol did not change. Comparing plasma and

water samples, we found water-borne cortisol was positively correlated with cortisol in plasma

and the relationship between plasma and water-borne corticosterone was positive but not

statistically significant. These results challenge the broadly accepted assumption that

corticosterone is 'dominant' in amphibians and add to the growing literature highlighting the

importance of measuring both GCs to understand (stress) physiology.

Keywords: ACTH challenge, anurans, corticosterone, cortisol, Dendrobatids, glucocorticoids

1.   Introduction

Researchers across disciplines are interested in understanding stress. The ability to measure

stress physiology non-invasively in the lab and in the wild is key for behavioral ecologist and

conservation biologists seeking to understand animal behavior in a changing world. In studying

stress physiology, we often measure glucocorticoids (GCs), most commonly cortisol and

corticosterone, produced by the adrenal cortex in response to stress (Sapolsky et al., 2000).

Glucocorticoids are of keen interest not only as a readout of stress, but also because they elicit a

variety of physiological, morphological, and behavioral responses (MacDougall-Shackleton et

al., 2019). For example, we know elevated levels of corticosterone can have negative impacts on

tadpoles through reducing growth rate, altering morphology, and shortening time to

metamorphosis which affects growth and survivorship later in life (Glennemeier and Denver,

2002; Kulkarni and Gramapurohit, 2017).

Most studies measure only a single GC, the one considered to be "dominant". The dominant

GC is formally defined as the one released in response to adrenocorticotropin hormone (ACTH),

though dominance is often assumed based on higher abundance. Which GC is dominant varies

across species, with generalizations assumed based on taxon. Yet a key thing that is often

overlooked in stress physiology is the fact that other glucocorticoids are present, often in

substantial amounts, and can be produced independently of the HPA axis being activated during

times of stress (Bornstein et al. 2016). Thus, to fully understand the role glucocorticoids play in

population health, we need to move beyond simply studying the glucocorticoid we assume to be

dominant.

1.1 Cortisol vs corticosterone –

Glucocorticoids are not only involved in the "stress response" (MacDougall-Shackleton et

al., 2019). In fact, glucocorticoids are involved in many physiological processes including, but

not limited to, energy metabolism (Vegiopoulos and Herzig, 2007), immune response (Cruz-

Topete and Cidlowski, 2015), growth and development (Seckl and Meaney, 2004), reproduction

(Whirledge and Cidlowski, 2017), and cardiovascular function (e.g., Cruz-Topete et al., 2016).

Because of their widespread influences, changes in GCs and/or their receptors can be immensely

impactful on survival and fitness. While acutely elevated GC levels often have adaptive

functions, in many species chronically elevated GCs can disrupt normal physiology and decrease

fitness (reviewed in Bonier et al., 2009).

Cortisol and corticosterone are often considered interchangeable in their physiological

function (Sutanto and De Kloet, 1987). Typically, either cortisol or corticosterone is considered

the dominant GC produced by a species because it is more abundant at baseline and increases in

response to stressors or ACTH. ACTH challenges are often used to measure the responsiveness

of the adrenals, to identify the dominant GC, and to validate that the method of hormone

measurement is detecting an increase in HPA axis activity (Palme, 2019; Touma and Palme,

2005). However, the GC that is not responsive or less responsive to ACTH can also play an

important physiological role. For instance, in humans, cortisol is considered the primary effector

hormone of the HPA (Brück, 1983). However, corticosterone has been studied in humans for its

role in the central nervous system (Raubenheimer et al., 2006) and pre-natal responses to

stressors (Wynne-Edwards et al., 2013). In other cortisol-dominant species, the rate of

conversion from corticosterone to aldosterone may vary resulting in more or less circulating

corticosterone in cortisol-dominant species (Koren et al., 2012).

Despite large interspecific variation in GC levels and relative proportions of cortisol and

corticosterone (Koren et al., 2012), we commonly make default assumptions about which GC is

more abundant (and therefore often assumed to be dominant) based on taxa. The consequence of

this generalization is that studies typically measure only the GC that is generally considered

dominant, such as measuring only cortisol in fish and only corticosterone in birds. Yet, these

assumptions and generalizations can be misleading. For example, we find GC dominance varies

across closely and more distantly related mammals. In rodents, corticosterone is the considered

the dominant GC in rats and mice (Crisler et al., 2020), while cortisol is dominant in North

American red squirrels (*Tamiasciurus hudsonicus*) and Arctic ground squirrels (*Spermophilus*

parryii) (Boonstra and Mccoll, 2000). Yet another species in Rodentia, the golden hamster

(*Mesocricetus auratus*), secretes approximately equal amounts of both cortisol and

corticosterone (Albers et al., 1985; Kollack-Walker et al., 1997; Ottenweller et al., 1985).

Interestingly, a recent study of GC concentrations in hair found wide among-individual variation

in the ratio of cortisol to corticosterone across three otariid species (Stellar sea lions [*Eumetopias*

jubatus]; California sea lions [*Zalophus californianus*], and Northern fur seals [*Callorhinus*

ursinus]), though cortisol and corticosterone were correlated within-individuals suggesting either

cortisol or corticosterone could be measured to assess HPA axis activity (Keogh et al., 2020).

Similarly, there are also population differences in the ratio of cortisol to corticosterone in wild

sea otters (*Enhydra lutris*) across the species range with fecal corticosterone concentrations

ranging from around 10-85% of cortisol concentrations (Larson et al., 2009). These highly

variable findings within and across species suggest that we should use caution when following

dogmatic assumptions about which GC to measure based on, order, family, or even population

within the same species (Hancock, 2010).

As for variation in (relative) abundance, more studies are showing the importance of testing

interspecific variation in GC responses even among closely related species. For instance, across

mammals, we see all combinations of which GC responds to acute stress. Some species respond

with an increase in cortisol but not corticosterone (mice [*Mus musculus*] Gong et al., 2015; Javan

rusa deer [*Rusa timorensis*] van Mourik et al., 1985; tuco-tucos [*Ctenomys talarum*] Vera et al.,

2012, 2011), while in others corticosterone responds but not cortisol (clouded leopard [*Neofelis

nebulosa*] Young et al. 2004), and still in others *both* GCs respond (ferrets [*Mustela putorius*]

Rosenthal et al., 1993; red deer [*Cervus elaphus*] Smith and Bubenik, 1990). Sea otters are an

interesting case where some studies have found an increase in cortisol in response to ACTH

(Murray et al., 2020; Wasser et al., 2000), yet others find an acute stressor (catching and

handling) causes an increase in corticosterone suggesting a non-ACTH-driven regulation of

corticosterone (Larson et al., 2009). Across cortisol-dominant mammals, cortisol and

corticosterone are often, but not always, positively correlated with one another (reviewed in

Koren et al., 2012). Baleen whales are a great example of how species vary in their correlation

between the two GCs with a positive correlation in the North Atlantic right (*Eubalaena glacialis*)

and blue (*Balaenoptera musculus*) whales but not in the bowhead whale (*Balaena mysticetus*)

(Hunt et al., 2018). In baleen whales, the two GCs may synchronize when under severe stress or

in an energetically demanding physiological state such as reproduction (Hunt et al., 2018, 2017).

While in some cases there may be financial or logistical limitations preventing the measurement

of both GCs, whenever possible we should explore both to discover potentially interesting and

informative patterns.

1.2 Glucocorticoids in amphibians –

Non-invasively studying the stress physiology of amphibians can be instrumental in

understanding how species respond to multifactor environmental stressors, such as habitat

modification, climate variability, and disease, and can inform conservation strategies for at-risk

species (Kumar and Umapathy, 2019; Narayan, 2013; Palme, 2019; Walls and Gabor, 2019).

Corticosteroids also play critical roles in metamorphosis (Denver, 2009; Kulkarni and Buchholz,

2014). Corticosterone is typically referenced as the dominant and more abundant glucocorticoid

in amphibians (Moore and Jessop, 2003; Norris and Carr, 2013; Wingfield and Romero, 2011),

and occasionally listed as amphibians' only glucocorticoid (Katsu and Iguchi, 2016).

Consequently, the vast majority of studies on stress physiology in amphibians only measure

corticosterone. This dogma prevails despite a history of cortisol being detected in several species

of anurans (frogs and toads) (e.g., Dale, 1962; Krug et al., 1983) and a growing number of

studies demonstrating that the assumption of corticosterone dominance and/or greater abundance

does not hold for all species of amphibians. For example, hellbenders (*Cryptobranchus*

alleganiensis) mainly produce cortisol in response to a stressor (Hopkins et al., 2020).

Additionally, cortisol has been measured in multiple species of amphibians (terrestrial and

aquatic) using dermal swabs (Santymire et al., 2018), and cortisol increases in response to ACTH

in green treefrogs (*Hyla cinerea*), American toads (*Anaxyrus americanus*), and red-spotted newts

(*Notophthalamus viridescens*) (Santymire et al., 2018). Previous studies have suggested aquatic

amphibia rely on cortisol as their primary GC, as many fishes do, whereas semi-terrestrial or

terrestrial amphibia typically use corticosterone (Jungreis et al., 1970), yet the vast majority of

studies on a popular aquatic model species, the clawed frog (*Xenopus* sp.), measure

corticosterone (e.g., Jaudet and Hatey, 1984; Kulkarni and Buchholz, 2012; Shewade et al.,

2020). Measuring GCs in more amphibians will help us discern the role of different GCs across

different life histories and inform our general understanding of amphibian physiology. The

myriad of contradictory studies about the primary GC in amphibians brings into question

whether corticosterone can indeed be assumed to be the primary GC as we expand our study of

endocrinology to more non-traditional species.

1.3 Stress physiology in Neotropical poison frogs –

Many Neotropical poison frog (Dendrobatidae) species are threatened due to risks including

agrochemicals (Angulo et al., 2019), climate change (Whitfield et al., 2007), smuggling (Auliya

et al., 2016), and habitat destruction or fragmentation (Becker et al., 2007; Cushman, 2006). Due

to their susceptibility to the amphibian chytrid fungus *Batrachochytrium dendrobatidis* (*Bd*)

(Courtois et al., 2015) and conservation status (Guillory et al., 2019), the development of non-

invasive measurements of GCs is important for studying poison frog physiology and informing

conservation efforts. Neotropical poison frogs are primarily diurnal and terrestrial, and have been

gaining attention in behavioral ecology and neuroethology due to their interesting parental

behavior (Lötters et al., 2007; Roland and O'Connell, 2015; Westrick et al., 2023). In addition,

poison frogs vary in size but generally are relatively small, with an average snout-vent length

(SVL) of ~17-60 mm. Thus, repeated plasma sampling is not feasible, and we need to further

explore non-invasive methods of studying the endocrinology of poison frogs.

Our goal for this study was to investigate which glucocorticoid is the primary hormone of the

HPA axis in Dendrobatid poison frogs. Both cortisol (Fischer and O'Connell, 2020) and

corticosterone (Nowicki et al., 2022) have been measured in poison frogs, albeit in different

species and sample types (plasma from *Dendrobates tinctorius* and water from *Ranitomeya*

imitator, respectively). We tested interspecific variation in the relationship between cortisol and

corticosterone and focused on the response of both glucocorticoids to a physiological stressor in

one species. We used non-invasive water-borne hormone sampling to allow for repeated

measures from the same individuals (Baugh et al., 2018; Gabor et al., 2013; Rodríguez et al.,

2022). To examine species variation in glucocorticoid dominance among poison frogs, we took

"baseline" water samples from five species of Dendrobatids: the golden poison frog (*Phyllobates*

terribilis), mimic poison frog (*Ranitomeya imitator*), Zimmermann's poison frog (*Ranitomeya*

variabilis), dyeing poison frog (*Dendrobates tinctorius*), and Anthony's poison frog

(*Epipedobates anthonyi*). These species share many behaviors, such as parental care, and are

endemic to tropical regions of Central and South America, but vary in body size and toxicity in

the wild (Wells, 2007). *P. terribilis* and *D. tinctorius* are the largest of the group (~37-60 mm

SVL), *E. anthonyi* are intermediate (~19 to 25 mm SVL), and both *Ranitomeya* are among the

smallest poison frogs (~17-22 mm SVL). To confirm whether the more abundant GC is also

dominant, i.e. elevates in response to an acute stress signal, we additionally performed an ACTH

challenge in our primary focal species (*D. tinctorius*). In *D. tinctorius*, we then compared

glucocorticoid concentrations between water and plasma samples from the same individuals to

assess whether water samples are representative of circulating hormone levels in blood. All

together this study informs our understanding of the 'stress' physiology of this particularly

vulnerable clade of frogs and contributes important data to the puzzle of GC dominance in

amphibians.

2.   Materials and methods

2.1 Animal husbandry –

We housed all frogs in our breeding colony at University of Illinois Urbana-Champaign in

glass terraria (18x18x18", 12x12x18", or 36x18x18" Exo Terra®) with soil, sphagnum moss,

live tropical plants, and coconut shelters. We kept frogs on a 12:12 h light cycle and fed them

flightless *Drosophila* fruit flies (*D. melanogaster* or *D. hydei*) dusted with vitamin supplements

three times weekly. Our automated misting system kept humidity >75% and the room was kept

at 70-76ºF. We housed adult frogs in groups of two to three individuals of the same species,

either one or two males with one female (*E. anthonyi, D. tinctorius, R. variabilis, R. imitator*) or

as a group of two to three juveniles that were too young to be sexed at the time of sampling (*P.*

terribilis). We weighed frogs prior to sampling by placing them in a plastic container on a scale

and subtracting the weight of the container. All procedures were approved by UIUC Institutional

Animal Care and Use Committee (protocol #20147).

2.2 Hormone sampling –

To collect water-borne hormones, we placed individual frogs in glass containers with RO

water based on size. For the smaller poison frogs (*E. anthonyi, R, variabilis*, and *R. imitator*), we

used 4cm x 4cm x 5cm hexagonal glass jars with lids filled with 15 mL of RO water. For the

larger frogs (*D. tinctorius* and *P. terribilis*), we used 6.5cm x 6.5cm x 6cm glass tubs with lids

and 40 mL of RO water. Our goal was to use enough water to cover frogs' highly vascularized

pelvic skin in a small enough container that the frogs would stay in the water and not climb the

walls. We left frogs in the water for one hour before returning them to their home terraria or a

new water tub for additional sampling. We collected all samples 4.5-8 hours into the frog's 12 h

light cycle. We filtered water samples with Whatman® 1 filter papers to remove any large

particulates, such as fecal matter, prior to steroid hormone extraction (see Section 2.4). To

compare across species, we collected n=84 water samples (*D. tinctorius* n = 6, *E. anthonyi* n =

20, *P. terribilis* n = 20, *R. imitator* n = 20, *R. variabilis* n = 18). With n=32 samples, we

subsampled half of the water for additional projects and stored each subsample separately. We

stored all water samples at -20ºC until extraction. For the plasma and water-borne hormone

comparison, we collected water samples as described above for *D. tinctorius* (n = 10)

immediately prior to collecting plasma. For n=5 samples, we split water samples in half for

additional projects. To collect plasma samples, we anesthetized frogs with benzocaine (Orajel™)

and euthanized via rapid decapitation. We used heparinized microhematocrit capillary tubes to

collect trunk blood from the body. We combined all the collected blood from an individual in a

microcentrifuge tube and centrifuged for 5 min at 5,000 rpm to separate the plasma. Using a

micropipette, we collected the plasma supernatant and stored all plasma samples at -20°C.

2.3 ACTH hormone challenge –

For the ACTH hormone challenges, we used a within-individual repeated measures design

with 18 adult *D. tinctorius* (9 male, 9 female). Each frog experienced an injection of 0.5 ug

ACTH per g body weight (Sigma-Aldrich®, product no. A7075) diluted in 0.9% saline and a

vehicle injection of only 0.9% saline with 6 days between injections. The order of injections was

counterbalanced across individuals such that half of the frogs were first injected with vehicle,

and half were first injected with ACTH. Prior to the start of trials, we dissolved the ACTH in

saline at a concentration of 0.125 ug/µL and aliquoted the solution into insulin syringes which

were frozen at -20°C until the day of injection, along with saline vehicle aliquots. On the day of

the trial, we thawed the aliquots of saline and ACTH and calculated the volume needed to inject

each frog with a dosage of 0.5 ug ACTH per g body weight, based on previous work using

ACTH challenges in Neotropical frogs (Baugh et al., 2018). Since the precision of the insulin

syringes was limited, we rounded to the nearest 10 µL to get the volume for injection with both

ACTH and saline. For each trial, we collected a water-borne hormone sample, as described

above, for one hour prior to injection as our "pre-injection" sample. We then used insulin

syringes to inject frogs intraperitoneally. After injecting, we immediately placed the frogs in a

new water bath for one hour to collect the "post-injection" sample. After the post-injection

sample, we returned frogs to their home terraria. We repeated this process the following week

with the alternate treatment (ACTH or saline). We performed all trials between five and seven

hours after the lights turned on. We split each water sample (four samples per 18 frogs = 72

samples total) in half for additional projects and stored subsamples separately.

2.4 Steroid hormone extraction –

For all water samples, we used C18 cartridges (Waters™ Corporation, Milford, MA) for

solid-phase extraction to collect steroid hormones. With cartridges on a vacuum manifold, we

used a vacuum pump to prime the filters with 6 mL 100% methanol. We then rinsed the

cartridges with 6 mL RO water before passing the sample through the cartridge. We eluted the

hormones from the cartridges with 4 mL of 100% methanol into 13x100mm borosilicate glass

vials. We dried samples under a stream of nitrogen gas in a 37°C dry heating block until

completely dry. We stored desiccated samples at -20°C until reconstituted with assay buffer for

enzyme linked immunosorbent assay (ELISA) analysis (see Section 2.6).

To measure glucocorticoids from plasma samples, we used the dissociation reagent included

in our ELISA kits (see Section 2.6). We combined 5 µL of dissociation reagent and 5 µL of

plasma, vortexed, and incubated for a minimum of 5 min at room temperature. We then added

450 µL of assay buffer to make a 1:100 dilution before running samples on ELISA plates.

2.6 Enzyme linked immunosorbent assays –

We used DetectX® ELISA kits from Arbor Assays™ (Ann Arbor, MI) to measure total

cortisol and corticosterone concentration in plasma and water samples. We used each kit

according to the manufacturer's instructions except for using the same assay buffer (catalog no.

X065) for both kits to allow us to split samples across cortisol and corticosterone plates,

following manufacturer recommendations. For the corticosterone kit, we followed instructions

for the 100 µL format because the standard curve for the 100 µL format can measure smaller

concentrations than the 50 µL format. The cortisol ELISA plate required 50 µL of sample for

each well, or 100 µL to run samples in duplicate. Initially, we reconstituted samples with 225 µL

of assay buffer to ensure an even split of the original sample across both cortisol and

corticosterone plates. Starting with the cortisol ELISA plate, we pipet 50 µL of samples and

standards into wells in duplicate (100 µL in total). We followed the rest of the manufacturer's

instructions for adding the conjugate and antibody to the cortisol plate. We then added 105 µL of

assay buffer to further increase the volume of the remaining 125 µL of each sample so we could

use the 100 µL assay format for corticosterone to allow for measurement of lower

concentrations. We pipet 100 µL of each sample into wells in duplicate (200 µL in total) on the

corticosterone ELISA plate and proceeded according to the manufacturer's instructions. For both

cortisol and corticosterone, we measured the optical density at 450 nm. After removing samples

with a coefficient of variation (CV) > 20% (n = 110 out of 352 total measurements), the average

intra-assay CV across all samples was 8.4% for cortisol and 8.2% for corticosterone. The cross

reactivity for the cortisol ELISA plate with corticosterone is reported by the manufacturer as

1.2% and for the corticosterone ELISA plate cross reactivity with cortisol is reported as 0.38%.

We used a four-parameter logistic curve (4PLC) regression to fit the respective standard

curve on each plate using MyAssay software recommended by the manufacturer

(www.MyAssay.com). Based on this curve, we calculated the concentration of cortisol and

corticosterone (pg/mL of assay buffer) for each sample and averaged across duplicates. Due to

low concentrations of both GCs in some species, some samples were too low to fall on the

standard curve (n=30 out of 352 total measurements), though some had concentrations of cortisol

that were too high (n=12). To be conservative, for samples that were off the standard curve in

either direction (too low or too high), we censored the data at the minimum or maximum

detection limits of the assay, respectively. To confirm the censored data was not skewing our

findings, we ran additional analyses with only samples measuring within the assay detection

limits and CV < 20%, and the overall pattern of results remained the same (Supplemental

Materials, Figure S1). To calculate the release rate from the concentration, we multiplied the

concentration by the volume of assay buffer used for initially reconstituting the sample (0.225

mL) to get mass (pg) of the hormone in the sample and divided by collection time (one hour) to

get the release rate. For samples that were split in half prior to extraction, we multiplied the rate

(pg/hr) by two to allow comparison with samples that were not split.

2.7 Statistical analysis –

We used R v4.2.2 (R Core Team, 2022) in RStudio v3.1.446 (Posit Team, 2023) for all

statistical analyses and ran all analyses on log transformed hormone release rates to normalize

residuals. We first tested each assay for parallelism by combining 24 additional water samples

from adult *D. tinctorius* to create a highly concentrated pooled sample and then serial diluting by

1/2 of the reconstituted extracts for a dilution curve with seven samples in total. This serial

dilution sample was split four ways to test parallelism across four different ELISA plates for

steroid hormones, including cortisol and corticosterone. We confirmed parallelism by comparing

the slope of the serially diluted curve to the slope of the standard curve using an interaction term

in a linear model. We found the dilution curves were not significantly different from the standard

curve for their respective hormone (cortisol: ß=-0.34, SE = 1.08, p = 0.76; corticosterone: ß=-

5.09, SE = 3.06, p = 0.12).

Across species, we were primarily interested in whether the release rates of the two GCs

differed within species, rather than absolute GC differences between species. To this end, we fit

a linear mixed model for log transformed hormone release rate with a fixed effect of the

interaction between species and glucocorticoid type (corticosterone vs cortisol) with sample ID

as a random effect. We ran a Type III ANOVA on the linear mixed model to assess the overall

effect of the interaction term. We then ran post-hoc simultaneous pairwise comparisons using the

multcomp package (Hothorn et al., 2008) to specifically look at the difference in release rate

between cortisol and corticosterone within species (e.g., *D. tinctorius* cortisol vs. *D. tinctorius*

corticosterone). We did not include sex as a covariate since not all individuals could be

confidently sexed at time of sampling. However, in a reduced model with only individuals that

we could confidently sex, we found no overall effect of sex on GC production (Table S1). To

increase our sample size for *D. tinctorius* in the species comparison, we included samples on the

first day of trials before the injection during ACTH challenges in the analysis (n=22 out of 29

total measurements of *D. tinctorius* samples). Our final species comparison dataset included 29

measurements from *D. tinctorius* (15 corticosterone, 14 cortisol), 25 measurements from *E.*

anthonyi (12 corticosterone, 13 cortisol), 23 measurements from *P. terribilis* (8 corticosterone,

15 cortisol), 22 measurements from *R. imitator* (8 corticosterone, 14 cortisol), and 19

measurements from *R. variabilis* (3 corticosterone, 16 cortisol).

For the ACTH challenges, we fit a linear mixed effects model to ask if there was an effect of

treatment (ACTH or control) on either cortisol or corticosterone. The model had the following

main effects: GC type (cortisol or corticosterone), time point (pre- or post-injection), and

injection type (ACTH or saline). We also included a three-way interaction between GC type,

time point, and injection type. This interaction indicated if the difference between pre- and post-

injection varied by injection type and whether this relationship varied by GC type. In other

words, is there a change in cortisol from pre- to post-injection based on injection type and is this

change (or lack thereof) the same or different in corticosterone? We included trial ID as a nested

random effect within frog ID to account for repeated measures across trials and individuals. We

used post-hoc pairwise comparisons using the multcomp package to interpret interaction effects.

After filtering for CV, our sample size was 26 pre- and 26 post-injection cortisol measurements

and 27 pre- and 30 post-injection corticosterone measurements. Because our main interest was

how the within-individual response to the injection varied across injection types, all four samples

for each individual were run on the same plate to reduce possible confounding effects of inter-

plate variation within individuals and trials.

We calculated the original plasma sample concentration by multiplying the concentration

measured in the assay by the dilution factor. To examine the relationship between water-borne

glucocorticoids and circulating glucocorticoids in plasma, we fit linear models (R packages lme4

(Bates et al., 2015) and lmerTest (Kuznetsova et al., 2016)) for both cortisol and corticosterone

with plasma concentrations (log transformed pg/mL) as the fixed effect predicting water

hormone release rate (log transformed pg/hr) for the matched plasma-water samples.

3.   Results

3.1 Species variation in relative glucocorticoid abundance –

Overall, we found species differed in which glucocorticoid was more prevalently released in

the water ($F_{9,49.44}=14.49$, $p < 0.0001$; Fig 2). Specifically, pairwise comparisons within species

revealed *D. tinctorius* was unique in releasing almost equal amounts of corticosterone and

cortisol, while the other four species released more cortisol (Table 1; Fig 2). This pattern

remained the same if we exclude all samples that did not fall within the range of the assays

(Table S2; Fig S1).

371   *Table 1.* Results from simultaneous multiple pairwise comparisons of means for hormone release

372   rate across species. P-values < 0.05 are in bold.

	log(release rate, pg/hr): cortisol - corticosterone			
species	*estimates*	*std. error*	*z-value*	*p-value*
D. tinctorius	0.036	0.33	0.11	1.00
E. anthonyi	1.68	0.36	4.73	**< 0.0001**
P. terribilis	1.27	0.39	3.22	**0.006**
R. imitator	2.07	0.40	5.13	**< 0.0001**
R. variabilis	2.89	0.60	4.78	**< 0.0001**

Figure 1. Species variation in relative glucocorticoid abundance. (A) For four out of the five

species tested, the release rate of corticosterone was lower than the release rate of cortisol,

suggesting cortisol is the dominant glucocorticoid in these species. In *D. tinctorius*, we found

cortisol and corticosterone were released at the same rate. Boxplots for corticosterone are teal

(left) and for cortisol are purple (right). Grey dots indicate the mean for each group. Black dots

represent individual samples. (NS = not significant, ** = p < 0.001, *** = p < 0.0001) (B) The

species names correspond to the respective photos beneath the x-axis labels. The tree below the

photos indicates the phylogenetic relationships between species (Guillory et al., 2019).

3.2 ACTH challenges in D. tinctorius –

The effect of an injection (ACTH vs vehicle) varied by GC and also injection type (type of

GC x injection type x time point: ß= -0.97, p = 0.024; Table 2; Fig 2). We found decreased GC

release rates post-injection (ß=-0.59, p = 0.007; Table 2) and this effect was driven by a decrease

in corticosterone in saline injected frogs (pairwise comparison of corticosterone between pre-

and post-injection saline: ß = 0.59, z = 0.21, p = 0.031; Fig 2). After an injection, frogs had

higher corticosterone release rates when injected with ACTH compared to saline alone (pairwise

comparison of corticosterone between ACTH and saline post-injection: ß = 1.36, z = 5.86, p

<0.001; Fig 2). Cortisol release did not vary pre- and post-injection for either type of injection

(pairwise comparisons of cortisol between pre- and post-injection: saline ß = -0.02, z = -0.09, p =

1.00, ACTH ß = -0.03, z = -0.03, p = 1.00; Fig 2).

Table 2. Results from linear model for hormone release rate across injection type and time

points. The reference group is corticosterone, control (saline) injection, and pre-injection time

point. P values < 0.05 are listed in bold.

Predictors	log(release rate, pg/hr)		
	Estimates	CI	p
intercept	4.69	4.34 – 5.04	**<0.001**
glucocorticoid type (cortisol)	0.17	-0.26 – 0.60	0.429
injection type (ACTH)	0.40	-0.08 – 0.89	0.098
time point (post)	-0.59	-1.02 – -0.17	**0.007**
cortisol x ACTH	0.06	-0.54 – 0.66	0.842
cortisol x post-injection	0.61	0.00 – 1.22	**0.048**
ACTH x post-injection	0.95	0.37 – 1.53	**0.002**
cortisol x ACTH x post-injection	-0.97	-1.80 – -0.13	**0.024**
Random Effects			
σ^2	0.29		
τ_{00} trial:frogID	0.12		
τ_{00} frogID	0.02		
ICC	0.32		
N trial	35		
N frogID	18		
Observations	109		
Marginal R^2 / Conditional R^2	0.298 / 0.525		

Figure 2. Glucocorticoid response to an ACTH injection compared to a saline control injection. (A) We found no difference in cortisol release rate between samples taken before and after injection of either saline or ACTH. (B) In contrast, corticosterone release rate was higher after an ACTH injection than after a saline injection. Purple boxplots indicate cortisol data. Boxplots for corticosterone data are in teal. Lighter colors (on the left within injection groups) indicate the control/saline injection for the respective glucocorticoid. Grey dots indicate the mean for each treatment group. Black dots are individual water samples. (* = p < 0.05, *** = p < 0.001)

3.3 Plasma vs. water glucocorticoids –

When we compared water glucocorticoids with plasma glucocorticoids in *D. tinctorius*, the concentration of cortisol in plasma was positively correlated with the release rate of cortisol in water (ß=1.10, CI=0.51 – 1.70, R^2/R^2 adjusted = 0.818/0.782 p = 0.005; Fig 3). We found a slightly positive but statistically non-significant relationship between corticosterone concentration in plasma and corticosterone release rate in water (ß=0.20, CI = -0.31 – 0.72, R^2/R^2 adjusted = 0.108/-0.019, p = 0.387; Fig 3).

Figure 3. Correlations between glucocorticoids in plasma and water. The concentration of cortisol, but not corticosterone, in plasma is positively correlated with the concentration in water samples from the same individuals. Grey shaded area is the confidence interval around the statistically significant linear regression for cortisol.

4. Discussion

Glucocorticoids are involved in many physiological processes that are essential for survival and fitness (MacDougall-Shackleton et al., 2019; Nicolaides et al., 2015); therefore, understanding glucocorticoids and HPA axis dynamics can inform conservation efforts and improve our understanding of how animals are being impacted by environmental changes and threats (Busch and Hayward, 2009; Millspaugh and Washburn, 2004). Often the GC that responds most to an ACTH challenge (the dominant GC) is also the more abundant GC (e.g. Flament et al., 2012). To better understand the stress physiology of Dendrobatids, we measured water-borne glucocorticoids in multiple poison frogs and performed an ACTH hormone challenge and plasma-water correlations in our focal species, *D. tinctorius*. Our findings challenge the common assumption that corticosterone is both more abundant and the dominant GC in frogs and – more broadly – the assumptions that relative abundance predicts dominance.

We demonstrate that focusing only on the ACTH-responsive GC can overlook potentially

important ACTH-independent regulation of other abundant GCs.

4.1 Species variation in glucocorticoid dominance –

Results from all five poison frog species across the *Dendrobatidae* family contradict the

common assumption that corticosterone is both the dominant and most abundant GC in

amphibians (Moore and Jessop, 2003; Norris and Carr, 2013; Wingfield and Romero, 2011), and

in particular in terrestrial amphibians (Jungreis et al., 1970). In water-borne hormone samples

from five poison frog species, we found that most species released more cortisol than

corticosterone. On average, both *Ranitomeya* species released more than twice as much cortisol

as corticosterone (Fig 1). The difference between cortisol and corticosterone was less dramatic

for *P. terribilis* and *E. anthonyi* but cortisol was still released at more than 1.25x the rate of

corticosterone (Fig 1). In contrast, in *D. tinctorius* we detected approximately equal amounts of

both glucocorticoids (Fig 1). In our study, we opportunistically chose these five species of poison

frogs, and it would be interesting to know if this pattern extends to all Dendrobatids.

Most studies comparing cortisol and corticosterone within-species are done in mammals with

few systematic studies in other taxa (Vera et al., 2017). Despite the generally accepted rule that

corticosterone dominant and therefore more abundant in amphibians, the few studies that have

looked at both cortisol and corticosterone in amphibians report a range of empirical results with

no consistent pattern. In some species, corticosterone is the dominant glucocorticoid, as

canonically expected. For example, Rio Grande Leopard tadpoles released more corticosterone

than cortisol overall and increased corticosterone release rates in response to an ACTH

challenge, but decreased cortisol release rates (Forsburg et al., 2019). Similarly, in American

bullfrogs (*Rana catesbeiana*), corticosterone in plasma responded to ACTH whereas cortisol was

undetectable (Johnston et al., 1967). In contrast, other studies in amphibians find cortisol is the

more abundant glucocorticoid, as we find here for multiple species of poison frogs, and

responsive to ACTH. For instance, in fecal samples from two species of Panamanian harlequin

frogs (*Atelopus certus* and *Atelopus glyphus*) both cortisol and corticosterone were responsive to

ACTH, but cortisol metabolites were detected at much higher concentrations overall (Cikanek et

al., 2014). Yet in other taxa, both cortisol and corticosterone are detectable at relatively equal

concentrations, similar to our finding for *D. tinctorius*. In the common water frog (*Rana*

esculenta), cortisol and corticosterone are detectable in plasma and both are elevated in ovulatory

and postovulatory phases compared to the preovulatory phase (Gobbetti and Zerani, 1993).

Further equivocal results about cortisol and corticosterone in amphibians, including

contradictory results in the same species (such as *Xenopus laevis* and *Rana pipens*), are nicely

summarized in Table 2 in Hopkins et al. (2020). With empirical support for dominance of either

GC in amphibians, as well as variation in the relationship between dominance and abundance,

we caution against making a general assumption about which GC is responsive to stress, even

across closely related species.

These inconsistent findings on GC dominance in amphibians and other taxa beg the question

of what could explain this variation. It has been suggested that cortisol and corticosterone may in

fact have some different physiological functions and be differentially regulated within a species.

Focusing only on the ACTH-responsive or "dominant" GC ignores ACTH-independent

regulation of the other GC, which may be just as important (Bornstein et al., 2008; Bornstein and

Chrousos, 1999). The two GCs could be playing different roles across age classes and metabolic

functions. For instance, in a study on zebra finches (*Taeniopygia guttata*), the more abundant GC

varied by tissue type with more corticosterone (the dogmatic primary GC of birds) detected in

plasma, but more cortisol detected in immune tissues (Schmidt and Soma, 2008). Additionally,

corticosterone and cortisol may differentially bind to species-specific corticosteroid-binding

globulins resulting in differing amounts of bound and unbound corticosteroid available in

circulation (Breuner and Orchinik, 2002; Martin and Ozon, 1975; Westphal, 1986).

Pharmacological studies also suggest differential glucocorticoid receptor-mediated gene

regulation based on which type of corticosteroid is available (e.g. Nixon et al., 2016; Pariante et

al., 2003) and differential transport of each corticosteroid across the blood-brain barrier (e.g.

Karssen et al., 2001).

Notably, there is some evidence in frogs that the dominant glucocorticoid relative

glucocorticoid abundance may shift throughout development from tadpole to adult (Dale, 1962;

Jungreis et al., 1970; Krug et al., 1983; McClelland and Woodley, 2021). Previous studies have

found cortisol to be dominant in aquatic larval tadpoles (Dale, 1962), but corticosterone to be

dominant in terrestrial adults of the same species (Jungreis et al., 1970). This developmental shift

is not unique to amphibians. For instance, in rabbits (*Oryctolagus cuniculus*), the ratio of cortisol

to corticosterone shifts from cortisol-dominant to corticosterone-dominant as they develop from

fetus to adult (Mulay et al., 1973). The functional reason for this potential ontogenetic shift in

glucocorticoid dominance across development is unclear and warrants further study across

species and developmental stages. At a mechanistic level, the enzymes necessary for cortisol

production may be regulated by different stimuli than the enzymes for corticosterone production,

and these stimuli may be restricted to specific developmental stages. In particular, researchers

should explore non-ACTH routes of GC release (reviewed in Bornstein et al., 2008; Bornstein

and Chrousos, 1999; Lesouhaitier et al., 1995). Release of glucocorticoids by extrapituitary

mechanisms, including neurotransmitters, neuropeptides, and cytokines, is far less studied and

less understood than activation of the adrenals by ACTH (Bornstein et al., 2008; Bornstein and

Chrousos, 1999; Lesouhaitier et al., 1995). Amphibians are especially promising for studies

across developmental stages given their biphasic life histories that lead to dramatically different

physiological challenges at different life stages.

4.2 Glucocorticoid response to adrenal activation in D. tinctorius –

Our second goal in this study was to identify which glucocorticoid was responsive to

activation of the adrenal glands via injection of ACTH, i.e. the dominant GC by the standard

definition. In *D. tinctorius*, we found the release rate of corticosterone decreased following a

saline injection, indicating frogs were habituating to the sampling process and no longer

mounting a stress response, however an ACTH injection prevented this decrease in

corticosterone, suggesting a stress response relative to controls (Fig 2). Interestingly, cortisol did

not respond to either an injection of ACTH or saline (Fig 2) suggesting cortisol is not responsive

to acute adrenal activation and does not fluctuate across short time periods (i.e., does not show

an increased stress response or habituation). In combination, these results suggest corticosterone

is more responsive to acute adrenal activation than cortisol, despite approximately equal amounts

of cortisol and corticosterone prior to injections. Typically, the GC that responds most to adrenal

activation through an ACTH challenge is also the more abundant GC, but this was not the case in

D. tinctorius. Golden hamsters are another example where the more abundant GC is not the GC

that responds to acute adrenal activation. A previous study found corticosterone was higher than

cortisol at baseline in golden hamsters, yet cortisol but not corticosterone was elevated in

response to chronic stress (Ottenweller et al., 1985). Our finding that the GC that responded to an

ACTH challenge is not the most abundant GC in *D. tinctorius* further muddies our understanding

of which GC plays a primary role in stress physiology in poison frogs.

Our current study shows corticosterone responds to ACTH, however a previous study in *D.*

tinctorius found an increase in cortisol in plasma during an energetically costly behavior, tadpole

transport (Fischer and O'Connell, 2020). Taken together these results suggest corticosterone and

cortisol may serve distinct functions and can be released via distinct mechanisms depending on

context. As discussed above, there are many non-ACTH routes of GC production, including

neurotransmitters and neuropeptides involved in parental care (Bornstein and Chrousos, 1999).

The parental state involves a myriad of physiological changes (Numan, 2020; Rosenblatt and

Snowdon, 1996) and perhaps the unique demands of parenting rely on the production of

glucocorticoids through non-ACTH routes. Though speculative, we suggest that distinct ACTH

and ACTH-independent mechanisms could provide modular regulation of parental care versus

other stressors (McGlothlin and Ketterson, 2008).

In addition, the functionality of cortisol and corticosterone within a species may be more

distinct than we typically assume (Schmidt and Soma, 2008; Vera et al., 2017), leading to

preferential expression of one GC over the other in some contexts and synchronized release in

other contexts. In some species, cortisol is more responsive to acute stress than corticosterone

(Gong et al., 2015; Koren et al., 2012; Venkataseshu and Estergreen, 1970; Vera et al., 2012,

2011), however the two GCs may also simultaneously elevate in more chronic states with high

energetic burdens. For instance, both cortisol and corticosterone increase during pregnancy in

humans (Wintour et al., 1978), little brown bats (*Myotis lucifugus,* Reeder et al., 2004), North

Atlantic right whales (Hunt et al., 2017), and Asian elephants (*Elephas maximus*, Kajaysri and

Nokkaew, 2014). In essence, the ratio of the two GCs in circulation may be dependent on

physiological state (Gong et al., 2015). In the context of increased cortisol during parenting in *D.*

tinctorius (Fischer and O'Connell, 2020), an increase in corticosterone, but not cortisol,

following an ACTH injection suggests non-ACTH routes of GC production may be important in

upregulating cortisol during behavioral states with higher energetic demand. Such modular

regulation could occur through changes in the regulation of the enzyme required to produce

cortisol (17α-hydroxylase). Further studies are needed to explore these intriguing possibilities.

We note our current study and Fischer and O'Connell (2020) measured hormones in different

sample types (plasma vs. water); however, we did find a positive correlation between plasma

concentration and water release rate for cortisol. Additionally, water-borne hormones are an

integrative measure of hormone release rate over time, rather than measuring one moment in

time as in plasma. The collection of water-borne hormones and plasma is inherently separated in

time and therefore, perhaps our null expectation should not be a tight relationship between

plasma and water-borne hormones. Moreover, the relationship between plasma and water-borne

hormones could be life stage-dependent in amphibians, as previously demonstrated in Northern

Leopard frogs (*Lithobates pipiens*; McClelland and Woodley, 2021). Finally, even when plasma

and water hormone levels are correlated, GCs may differ in their release rates such that relative

abundances differ between plasma and water. For example, in túngara frogs (*Physalaemus*

pustulosus), cortisol was present in higher concentrations than corticosterone in water-borne

samples, but corticosterone was higher in plasma samples (Baugh et al., 2018). Thus, while most

studies of water-borne hormones in amphibians find a positive relationship between plasma and

water measurements as we did here, the complexities of this relationship require further study.

4.3 Conclusions

Altogether, these findings highlight the importance of measuring both cortisol and

corticosterone when investigating the complex, multivariate roles of GCs in physiology,

development, and behavior. Even in well-studies species, measuring both GCs could bring to

light new information about the differential regulation and functionality of cortisol and

corticosterone, which may not be as interchangeable as we assume (Hancock, 2010). Our results

suggest poison frogs go against the textbook dogma of corticosterone being dominant and

therefore more abundant in terrestrial amphibians. We found cortisol released at higher rates than

corticosterone in multiple species. *D. tinctorius* were distinctive in releasing nearly equal

amounts of corticosterone and cortisol yet follow the general assumption of corticosterone

dominance by responding to ACTH with an increase in corticosterone. These findings inspire

more questions than they answer. For instance, why does *D. tinctorius* show different relative

GC abundance than other poison frogs? Are cortisol and corticosterone independently regulated

through ACTH and non-ACTH routes? Are poison frogs unique among terrestrial frogs in

producing more cortisol than corticosterone? We clearly have a lot left to learn about the stress

physiology of this unique clade of frogs and amphibians in general. We can say with confidence

that there is interesting, under-appreciated, and unexplored variation in HPA activity and

glucocorticoids, and that simplifying assumptions about the relationship between GC abundance
and dominance do not hold.

Ultimately, we need more studies measuring both GCs whenever possible. We encourage
more researchers to investigate both GCs in their own focal species, regardless of the dogma, as
the "non-dominant" GC may in fact be more abundant. With data across more species, we can
look for broader patterns in GC dominance and abundance to investigate whether there are
adaptive and/or ecological explanations for the diverse patterns emerging across species,
populations, and life stages. We can also test hypotheses about potential unique physiological
roles for cortisol versus corticosterone, which may have previously been obscured due to
measurement of only a single GC. Additionally, measuring both cortisol and corticosterone may
better inform conservation practices through improving our understanding of stress physiology
across species and environments.

Declarations of Competing Interest none

CRediT authorship contribution statement
SEW: conceptualization, methodology, validation, formal analysis, investigation, data curation,
writing – original draft, writing – review & editing, visualization, project administration, funding
acquisition. **RTP:** conceptualization, methodology, writing – review & editing, supervision.
EKF: conceptualization, methodology, writing – review & editing, supervision, funding
acquisition.

Acknowledgements We thank members of the Fischer Lab for their feedback on the manuscript
and for the caretaking of the frogs.

Funding This study was supported by the National Science Foundation (#2010714 to SEW;
#2146058 to EKF) and University of Illinois Urbana-Champaign (start-up funds and RB21025 to
EKF).

Data statement Data and code are available on FigShare (doi: 10.6084/m9.figshare.23537409).

References

Albers, H.E., Yogev, L., Todd, R.B., Goldman, B.D., 1985. Adrenal corticoids in hamsters: role in circadian timing. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology 248, R434–R438. https://doi.org/10.1152/ajpregu.1985.248.4.R434

Angulo, A., von May, R., Icochea, J., 2019. A reassessment of the extinction risk of the Critically Endangered Oxapampa poison frog *Ameerega planipaleae* (Dendrobatidae). Oryx 53, 557–560. https://doi.org/10.1017/S0030605318000571

Auliya, M., García-Moreno, J., Schmidt, B.R., Schmeller, D.S., Hoogmoed, M.S., Fisher, M.C., Pasmans, F., Henle, K., Bickford, D., Martel, A., 2016. The global amphibian trade flows through Europe: the need for enforcing and improving legislation. Biodivers Conserv 25, 2581–2595. https://doi.org/10.1007/s10531-016-1193-8

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting linear mixed-effects models using lme4. Journal of Statistical Software 67, 1–48. https://doi.org/10.18637/jss.v067.i01

Baugh, A.T., Bastien, B., Still, M.B., Stowell, N., 2018. Validation of water-borne steroid hormones in a tropical frog (*Physalaemus pustulosus*). General and Comparative Endocrinology 261, 67–80. https://doi.org/10.1016/j.ygcen.2018.01.025

Becker, C.G., Fonseca, C.R., Haddad, C.F.B., Batista, R.F., Prado, P.I., 2007. Habitat Split and the Global Decline of Amphibians. Science 318, 1775–1777. https://doi.org/10.1126/science.1149374

Bonier, F., Martin, P.R., Moore, I.T., Wingfield, J.C., 2009. Do baseline glucocorticoids predict fitness? Trends in Ecology and Evolution 24, 634–642. https://doi.org/10.1016/j.tree.2009.04.013

Boonstra, R., Mccoll, C.J., 2000. Contrasting Stress Response of Male Arctic Ground Squirrels and Red Squirrels. Journal of Experimental Zoology 286, 390–404.

Bornstein, S.R., Chrousos, G.P., 1999. Adrenocorticotropin (ACTH)- and Non-ACTH-Mediated Regulation of the Adrenal Cortex: Neural and Immune Inputs. The Journal of Clinical Endocrinology & Metabolism 84, 1729–1736. https://doi.org/10.1210/jcem.84.5.5631

Bornstein, S.R., Engeland, W.C., Ehrhart-Bornstein, M., Herman, J.P., 2008. Dissociation of

ACTH and glucocorticoids. Trends in Endocrinology & Metabolism 19, 175–180.

https://doi.org/10.1016/j.tem.2008.01.009

Breuner, C.W., Orchinik, M., 2002. Plasma binding proteins as mediators of corticosteroid action

in vertebrates. Journal of Endocrinology 175, 99–112.

https://doi.org/10.1677/joe.0.1750099

Brück, K., 1983. Functions of the Endocrine System, in: Schmidt, R.F., Thews, G. (Eds.),

Human Physiology. Springer, Berlin, Heidelberg, pp. 658–687.

https://doi.org/10.1007/978-3-642-96714-6_29

Busch, D.S., Hayward, L.S., 2009. Stress in a conservation context: A discussion of

glucocorticoid actions and how levels change with conservation-relevant variables.

Biological Conservation 142, 2844–2853. https://doi.org/10.1016/j.biocon.2009.08.013

Cikanek, S.J., Nockold, S., Brown, J.L., Carpenter, J.W., Estrada, A., Guerrel, J., Hope, K.,

Ibáñez, R., Putman, S.B., Gratwicke, B., 2014. Evaluating Group Housing Strategies for

the Ex-Situ Conservation of Harlequin Frogs (*Atelopus* spp.) Using Behavioral and

Physiological Indicators. PLOS ONE 9, e90218.

https://doi.org/10.1371/journal.pone.0090218

Courtois, E.A., Gaucher, P., Chave, J., Schmeller, D.S., 2015. Widespread Occurrence of Bd in

French Guiana, South America. PLOS ONE 10, e0125128.

https://doi.org/10.1371/journal.pone.0125128

Crisler, R., Johnston, N.A., Sivula, C., Budelsky, C.L., 2020. Chapter 4 - Functional Anatomy

and Physiology, in: Suckow, M.A., Hankenson, F.C., Wilson, R.P., Foley, P.L. (Eds.),

The Laboratory Rat (Third Edition), American College of Laboratory Animal Medicine.

Academic Press, pp. 91–132. https://doi.org/10.1016/B978-0-12-814338-4.00004-0

Cruz-Topete, D., Cidlowski, J.A., 2015. One Hormone, Two Actions: Anti- and Pro-

Inflammatory Effects of Glucocorticoids. NIM 22, 20–32.

https://doi.org/10.1159/000362724

Cruz-Topete, D., Myers, P.H., Foley, J.F., Willis, M.S., Cidlowski, J.A., 2016. Corticosteroids

Are Essential for Maintaining Cardiovascular Function in Male Mice. Endocrinology

157, 2759–2771. https://doi.org/10.1210/en.2015-1604

Cushman, S.A., 2006. Effects of habitat loss and fragmentation on amphibians: A review and

prospectus. Biological Conservation 128, 231–240.

https://doi.org/10.1016/j.biocon.2005.09.031

Dale, E., 1962. Steroid excretion by larval frogs. General and Comparative Endocrinology 2,

171–176. https://doi.org/10.1016/0016-6480(62)90001-1

Denver, R.J., 2009. Stress hormones mediate environment-genotype interactions during

amphibian development. General and Comparative Endocrinology, Karl Lederis Special

Issue 164, 20–31. https://doi.org/10.1016/j.ygcen.2009.04.016

Fischer, E.K., O'Connell, L.A., 2020. Hormonal and neural correlates of care in active versus

observing poison frog parents. Hormones and Behavior 120, 104696.

https://doi.org/10.1016/j.yhbeh.2020.104696

Flament, A., Delleur, V., Poulipoulis, A., Marlier, D., 2012. Corticosterone, cortisol,

triglycerides, aspartate aminotransferase and uric acid plasma concentrations during *foie*

gras production in male mule ducks (*Anas platyrhynchos × Cairina moschata*). British

Poultry Science 53, 408–413. https://doi.org/10.1080/00071668.2012.711468

Forsburg, Z.R., Goff, C.B., Perkins, H.R., Robicheaux, J.A., Almond, G.F., Gabor, C.R., 2019.

Validation of water-borne cortisol and corticosterone in tadpoles: Recovery rate from an

acute stressor, repeatability, and evaluating rearing methods. General and Comparative

Endocrinology 281, 145–152. https://doi.org/10.1016/j.ygcen.2019.06.007

Gabor, C.R., Bosch, J., Fries, J.N., Davis, D.R., 2013. A non-invasive water-borne hormone

assay for amphibians. Amphibia-Reptilia 34, 151–162. https://doi.org/10.1163/15685381-

00002877

Glennemeier, K.A., Denver, R.J., 2002. Small changes in whole-body corticosterone content

affect larval Rana pipiens fitness components. General and Comparative Endocrinology

127, 16–25. https://doi.org/10.1016/S0016-6480(02)00015-1

Gobbetti, A., Zerani, M., 1993. Prostaglandin E2 and prostaglandin F2α involvement in the

corticosterone and cortisol release by the female frog, *Rana esculenta*, during ovulation.

Journal of Experimental Zoology 267, 164–170. https://doi.org/10.1002/jez.1402670209

Gong, S., Miao, Y.-L., Jiao, G.-Z., Sun, M.-J., Li, H., Lin, J., Luo, M.-J., Tan, J.-H., 2015.

Dynamics and Correlation of Serum Cortisol and Corticosterone under Different

Physiological or Stressful Conditions in Mice. PLOS ONE 10, e0117503.

https://doi.org/10.1371/journal.pone.0117503

Guillory, W.X., Muell, M.R., Summers, K., Brown, J.L., 2019. Phylogenomic Reconstruction of

the Neotropical Poison Frogs (Dendrobatidae) and Their Conservation. Diversity 11, 126.

https://doi.org/10.3390/d11080126

Hancock, T.M., 2010. A duality in mammalian glucocorticoid signaling evidence against the

dominant glucocorticoid/interchangeability assumption (M.Sc.). Queen's University

(Canada), Ontario, Canada.

Hopkins, W.A., DuRant, S.E., Beck, M.L., Ray, W.K., Helm, R.F., Romero, L.M., 2020.

Cortisol is the predominant glucocorticoid in the giant paedomorphic hellbender

salamander (*Cryptobranchus alleganiensis*). General and Comparative Endocrinology

285, 113267. https://doi.org/10.1016/j.ygcen.2019.113267

Hothorn, T., Bretz, F., Westfall, P., 2008. Simultaneous inference in general parametric models.

Biometrical Journal 50, 346–363. https://doi.org/10.1002/bimj.200810425

Hunt, K.E., Lysiak, N.S., Moore, M., Rolland, R.M., 2017. Multi-year longitudinal profiles of

cortisol and corticosterone recovered from baleen of North Atlantic right whales

(*Eubalaena glacialis*). General and Comparative Endocrinology 254, 50–59.

https://doi.org/10.1016/j.ygcen.2017.09.009

Hunt, K.E., Lysiak, N.S.J., Matthews, C.J.D., Lowe, C., Fernández Ajó, A., Dillon, D., Willing,

C., Heide-Jørgensen, M.P., Ferguson, S.H., Moore, M.J., Buck, C.L., 2018. Multi-year

patterns in testosterone, cortisol and corticosterone in baleen from adult males of three

whale species. Conservation Physiology 6, coy049.

https://doi.org/10.1093/conphys/coy049

Jaudet, G.J., Hatey, J.L., 1984. Variations in aldosterone and corticosterone plasma levels during

metamorphosis in *Xenopus laevis* tadpoles. General and Comparative Endocrinology 56,

59–65. https://doi.org/10.1016/0016-6480(84)90061-3

Johnston, C., Davis, J., Wright, F., Howards, S., 1967. Effects of renin and ACTH on adrenal

steroid secretion in the American bullfrog. American Journal of Physiology-Legacy

Content 213, 393–399. https://doi.org/10.1152/ajplegacy.1967.213.2.393

Jungreis, A.M., Huibregtse, W.H., Ungar, F., 1970. Corticosteroid identification and

corticosterone concentration in serum of *Rana pipiens* during dehydration in winter and

summer. Comparative Biochemistry and Physiology 34, 683–689.

https://doi.org/10.1016/0010-406X(70)90293-8

Kajaysri, J., Nokkaew, W., 2014. Assessment of Pregnancy Status of Asian Elephants *(Elephas*

maximus) by Measurement of Progestagen and Glucocorticoid and Their Metabolite

Concentrations in Serum and Feces, Using Enzyme Immunoassay (EIA). J. Vet. Med.

Sci. 76, 363–368. https://doi.org/10.1292/jvms.13-0103

Karssen, A.M., Meijer, O.C., van der Sandt, I.C.J., Lucassen, P.J., de Lange, E.C.M., de Boer,

A.G., de Kloet, E.R., 2001. Multidrug Resistance P-Glycoprotein Hampers the Access of

Cortisol But Not of Corticosterone to Mouse and Human Brain. Endocrinology 142,

2686–2694. https://doi.org/10.1210/endo.142.6.8213

Katsu, Y., Iguchi, T., 2016. Corticosterone, in: Handbook of Hormones. Elsevier, pp. 527-e95A-

3. https://doi.org/10.1016/B978-0-12-801028-0.00228-2

Keogh, M.J., Gastaldi, A., Charapata, P., Melin, S., Fadely, B.S., 2020. Stress-related and

reproductive hormones in hair from three north Pacific otariid species: Steller sea lions,

California sea lions and northern fur seals. Conservation Physiology 8, coaa069.

https://doi.org/10.1093/conphys/coaa069

Kollack-Walker, S., Watson, S.J., Akil, H., 1997. Social Stress in Hamsters: Defeat Activates

Specific Neurocircuits within the Brain. J. Neurosci. 17, 8842–8855.

https://doi.org/10.1523/JNEUROSCI.17-22-08842.1997

Koren, L., Whiteside, D., Fahlman, Å., Ruckstuhl, K., Kutz, S., Checkley, S., Dumond, M.,

Wynne-Edwards, K., 2012. Cortisol and corticosterone independence in cortisol-

dominant wildlife. General and Comparative Endocrinology 177, 113–119.

https://doi.org/10.1016/j.ygcen.2012.02.020

Krug, E.C., Honn, V., Battista, J., Nicoll, C.S., 1983. Corticosteroids in Serum of *Rima*

catesbeiana during Development and Metamorphosis. General and Comparative

Endocrinology 52, 232–241. https://doi.org/10.1016/0016-6480(83)90117-X

Kulkarni, P.S., Gramapurohit, N.P., 2017. Effect of corticosterone on larval growth, antipredator

behaviour and metamorphosis of *Hylarana indica*. General and Comparative

Endocrinology, Special issue on International Symposium on Comparative

Endocrinology and Integrative Physiology - CEIP 2015 251, 21–29.

https://doi.org/10.1016/j.ygcen.2016.09.001

Kulkarni, S.S., Buchholz, D.R., 2014. Corticosteroid signaling in frog metamorphosis. General
and Comparative Endocrinology, 2nd Meeting of The North American Society for
Comparative Endocrinology 203, 225–231. https://doi.org/10.1016/j.ygcen.2014.03.036

Kulkarni, S.S., Buchholz, D.R., 2012. Beyond Synergy: Corticosterone and Thyroid Hormone
Have Numerous Interaction Effects on Gene Regulation in *Xenopus tropicalis* Tadpoles.
Endocrinology 153, 5309–5324. https://doi.org/10.1210/en.2012-1432

Kumar, V., Umapathy, G., 2019. Non-invasive monitoring of steroid hormones in wildlife for
conservation and management of endangered species — A review. INDIAN J EXP
BIOL.

Kuznetsova, A., Brockhoff, P., Christensen, R., 2016. lmerTest: Tests in linear mixed effects
models. R package version 3.0.0, https://cran.r-project.org/package=lmerTest.

Larson, S., Monson, D., Ballachey, B., Jameson, R., Wasser, S.K., 2009. Stress-related hormones
and genetic diversity in sea otters (*Enhydra lutris*). Marine Mammal Science 25, 351–
372. https://doi.org/10.1111/j.1748-7692.2008.00260.x

Lesouhaitier, O., Esneu, M., Kodjo, M.K.L., Hamel, C., Contesse, V., Yon, L., Remy-Jouet, I.,
Fasolo, A., Fournier, A., Vandesande, F., Pelletier, G., Conlon, J.M., Roubos, E.W.,
Feuilloley, M., Delarue, C., Leboulenger, F., Vaudry, H., 1995. Neuroendocrine
Communication in the Frog Adrenal Gland. jzoo 12, 255–264.
https://doi.org/10.2108/zsj.12.255

Lötters, S., Jungfer, K.-H., Henkel, F.W., Schmidt, W., 2007. Poison frogs: biology, species and
captive care. Edition Chimaira, Frankfurt am Main.

MacDougall-Shackleton, S.A., Bonier, F., Romero, L.M., Moore, I.T., 2019. Glucocorticoids and
"Stress" Are Not Synonymous. Integrative Organismal Biology 1, obz017.
https://doi.org/10.1093/iob/obz017

Martin, B., Ozon, R., 1975. Steroid-Protein Interactions in Nonmammalian Vertebrates. II.
Steroids Binding Proteins in the Serum of Amphibians; a Physiological Approach.
Biology of Reproduction 13, 371–380. https://doi.org/10.1095/biolreprod13.4.371

McClelland, S.J., Woodley, S.K., 2021. Water-borne corticosterone assay is a valid method in
some but not all life-history stages in Northern Leopard Frogs. General and Comparative
Endocrinology 312, 113858. https://doi.org/10.1016/j.ygcen.2021.113858

McGlothlin, J.W., Ketterson, E.D., 2008. Hormone-mediated suites as adaptations and
evolutionary constraints. Philosophical Transactions of the Royal Society B: Biological
Sciences 363, 1611–1620. https://doi.org/10.1098/rstb.2007.0002

Millspaugh, J.J., Washburn, B.E., 2004. Use of fecal glucocorticoid metabolite measures in
conservation biology research: considerations for application and interpretation. General
and Comparative Endocrinology 138, 189–199.
https://doi.org/10.1016/j.ygcen.2004.07.002

Moore, I.T., Jessop, T.S., 2003. Stress, reproduction, and adrenocortical modulation in
amphibians and reptiles. Hormones and Behavior 43, 39–47.
https://doi.org/10.1016/S0018-506X(02)00038-7

Mulay, S., Giannopoulos, G., Solomon, S., 1973. Corticosteroid Levels in the Mother and Fetus
of the Rabbit During Gestation. Endocrinology 93, 1342–1348.
https://doi.org/10.1210/endo-93-6-1342

Murray, M.J., Young, M.A., Santymire, R.M., 2020. Use of the ACTH challenge test to identify
the predominant glucocorticoid in the southern sea otter (*Enhydra lutris nereis*).
Conservation Physiology 8, coz116. https://doi.org/10.1093/conphys/coz116

Narayan, E.J., 2013. Non-invasive reproductive and stress endocrinology in amphibian
conservation physiology. Conservation Physiology 1, cot011–cot011.
https://doi.org/10.1093/conphys/cot011

Nicolaides, N.C., Kyratzi, E., Lamprokostopoulou, A., Chrousos, G.P., Charmandari, E., 2015.
Stress, the Stress System and the Role of Glucocorticoids. Neuroimmunomodulation 22,
6–19. https://doi.org/10.1159/000362736

Nixon, M., Mackenzie, S.D., Taylor, A.I., Homer, N.Z.M., Livingstone, D.E., Mouras, R.,
Morgan, R.A., Mole, D.J., Stimson, R.H., Reynolds, R.M., Elfick, A.P.D., Andrew, R.,
Walker, B.R., 2016. ABCC1 confers tissue-specific sensitivity to cortisol versus
corticosterone: A rationale for safer glucocorticoid replacement therapy. Science
Translational Medicine 8, 352ra109-352ra109.
https://doi.org/10.1126/scitranslmed.aaf9074

Norris, D.O., Carr, J.A., 2013. Chapter 9 - Comparative Aspects of Vertebrate Adrenals, in:
Norris, D.O., Carr, J.A. (Eds.), Vertebrate Endocrinology (Fifth Edition). Academic
Press, San Diego, pp. 291–315. https://doi.org/10.1016/B978-0-12-394815-1.00009-4

839 Nowicki, J.P., Rodríguez, C., Lee, J.C., Goolsby, B.C., Yang, C., Cleland, T.A., O'Connell,

840   L.A., 2022. Evidence for Empathy in Pair Bonding Poison Frogs.

841   https://doi.org/10.1101/2022.09.25.509360

842 Numan, M., 2020. The parental brain: mechanisms, development, and evolution. Oxford

843   University Press, New York.

844 Ottenweller, J.E., Tapp, W.N., Burke, J.M., Natelson, B.H., 1985. Plasma cortisol and

845   corticosterone concentrations in the golden hamster, (*Mesocricetus auratus*). Life

846   Sciences 37, 1551–1558. https://doi.org/10.1016/0024-3205(85)90188-2

847 Palme, R., 2019. Non-invasive measurement of glucocorticoids: Advances and problems.

848   Physiology & Behavior 199, 229–243. https://doi.org/10.1016/j.physbeh.2018.11.021

849 Pariante, C.M., Kim, R.B., Makoff, A., Kerwin, R.W., 2003. Antidepressant fluoxetine enhances

850   glucocorticoid receptor function in vitro by modulating membrane steroid transporters.

851   British Journal of Pharmacology 139, 1111–1118. https://doi.org/10.1038/sj.bjp.0705357

852 Posit Team, 2023. RStudio: Integrated Development Environment for R.

853 R Core Team, 2022. R: A language and environment for statistical computing.

854 Raubenheimer, P.J., Young, E.A., Andrew, R., Seckl, J.R., 2006. The role of corticosterone in

855   human hypothalamic– pituitary–adrenal axis feedback. Clinical Endocrinology 65, 22–

856   26. https://doi.org/10.1111/j.1365-2265.2006.02540.x

857 Reeder, D.M., Kosteczko, N.S., Kunz, T.H., Widmaier, E.P., 2004. Changes in baseline and

858   stress-induced glucocorticoid levels during the active period in free-ranging male and

859   female little brown myotis, *Myotis lucifugus* (Chiroptera: Vespertilionidae). General and

860   Comparative Endocrinology 136, 260–269. https://doi.org/10.1016/j.ygcen.2003.12.020

861 Rodríguez, C., Fusani, L., Raboisson, G., Hödl, W., Ringler, E., Canoine, V., 2022. Androgen

862   responsiveness to simulated territorial intrusions in *Allobates femoralis* males: Evidence

863   supporting the challenge hypothesis in a territorial frog. General and Comparative

864   Endocrinology 326, 114046. https://doi.org/10.1016/j.ygcen.2022.114046

865 Roland, A.B., O'Connell, L.A., 2015. Poison frogs as a model system for studying the

866   neurobiology of parental care. Current Opinion in Behavioral Sciences 6, 76–81.

867   https://doi.org/10.1016/j.cobeha.2015.10.002

868 Rosenblatt, J.S., Snowdon, C.T. (Eds.), 1996. Parental Care: Evolution, Mechanisms, And

869   Adaptive Significance, Advances in the Study of Behavior. Academic Press.

Rosenthal, K.L., Peterson, M.E., Quesenberry, K.E., Lothrop, C.D., 1993. Evaluation of plasma
cortisol and corticosterone responses to synthetic adrenocorticotropic hormone
administration in ferrets. Am J Vet Res 54, 29–31.

Santymire, R.M., Manjerovic, M.B., Sacerdote-Velat, A., 2018. A novel method for the
measurement of glucocorticoids in dermal secretions of amphibians. Conservation
Physiology 6, coy008. https://doi.org/10.1093/conphys/coy008

Sapolsky, R.M., Romero, L.M., Munck, A.U., 2000. How do glucocorticoids influence stress
responses? Integrating permissive, suppressive, stimulatory, and preparative actions.
Endocrine Reviews 21, 55–89. https://doi.org/10.1210/er.21.1.55

Schmidt, K.L., Soma, K.K., 2008. Cortisol and corticosterone in the songbird immune and
nervous systems: local vs. systemic levels during development. American Journal of
Physiology-Regulatory, Integrative and Comparative Physiology 295, R103–R110.
https://doi.org/10.1152/ajpregu.00002.2008

Seckl, J.R., Meaney, M.J., 2004. Glucocorticoid Programming. Annals of the New York
Academy of Sciences 1032, 63–84. https://doi.org/10.1196/annals.1314.006

Shewade, L.H., Schoephoerster, J.A., Patmann, M.D., Kulkarni, S.S., Buchholz, D.R., 2020.
Corticosterone Is Essential for Survival Through Frog Metamorphosis. Endocrinology
161, bqaa193. https://doi.org/10.1210/endocr/bqaa193

Smith, J.H., Bubenik, G.A., 1990. Plasma concentrations of glucocorticoids in white-tailed deer:
the effect of acute ACTH and dexamethasone administration. Can. J. Zool. 68, 2123–
2129. https://doi.org/10.1139/z90-295

Sutanto, W., De Kloet, E.R., 1987. Species-Specificity of Corticosteroid Receptors in Hamster
and Rat Brains. Endocrinology 121, 1405–1411. https://doi.org/10.1210/endo-121-4-
1405

Touma, C., Palme, R., 2005. Measuring Fecal Glucocorticoid Metabolites in Mammals and
Birds: The Importance of Validation. Annals of the New York Academy of Sciences
1046, 54–74. https://doi.org/10.1196/annals.1343.006

van Mourik, S., Stelmasiak, T., Outch, K.H., 1985. Changes in plasma levels of cortisol and
corticosterone after acute ACTH stimulation in rusa deer (*Cervus rusa timorensis*).
Comparative Biochemistry and Physiology Part A: Physiology 81, 545–549.
https://doi.org/10.1016/0300-9629(85)91023-0

Vegiopoulos, A., Herzig, S., 2007. Glucocorticoids, metabolism and metabolic diseases.

Molecular and Cellular Endocrinology 275, 43–61.

https://doi.org/10.1016/j.mce.2007.05.015

Venkataseshu, G.K., Estergreen, V.L., 1970. Cortisol and Corticosterone in Bovine Plasma and

the Effect of Adrenocorticotropin. Journal of Dairy Science 53, 480–483.

https://doi.org/10.3168/jds.S0022-0302(70)86234-8

Vera, F., Antenucci, C.D., Zenuto, R.R., 2011. Cortisol and corticosterone exhibit different

seasonal variation and responses to acute stress and captivity in tuco-tucos (*Ctenomys*

talarum). General and Comparative Endocrinology 170, 550–557.

https://doi.org/10.1016/j.ygcen.2010.11.012

Vera, F., Zenuto, R., Antenucci, C.D., 2017. Expanding the actions of cortisol and corticosterone

in wild vertebrates: A necessary step to overcome the emerging challenges. General and

Comparative Endocrinology 246, 337–353. https://doi.org/10.1016/j.ygcen.2017.01.010

Vera, F., Zenuto, R.R., Antenucci, C.D., 2012. Differential Responses of Cortisol and

Corticosterone to Adrenocorticotropic Hormone (ACTH) in a Subterranean Rodent

(*Ctenomys talarum*). Journal of Experimental Zoology Part A: Ecological Genetics and

Physiology 317, 173–184. https://doi.org/10.1002/jez.1711

Walls, S.C., Gabor, C.R., 2019. Integrating Behavior and Physiology Into Strategies for

Amphibian Conservation. Frontiers in Ecology and Evolution 7.

Wasser, S.K., Hunt, K.E., Brown, J.L., Cooper, K., Crockett, C.M., Bechert, U., Millspaugh, J.J.,

Larson, S., Monfort, S.L., 2000. A Generalized Fecal Glucocorticoid Assay for Use in a

Diverse Array of Nondomestic Mammalian and Avian Species. General and Comparative

Endocrinology 120, 260–275. https://doi.org/10.1006/gcen.2000.7557

Wells, K., 2007. The Ecology and Behavior of Amphibians. The University of Chicago Press,

Chicago.

Westphal, U., 1986. Steroid-Protein Interactions Revisited, in: Westphal, U. (Ed.), Steroid-

Protein Interactions II, Monographs on Endocrinology. Springer, Berlin, Heidelberg, pp.

1–7. https://doi.org/10.1007/978-3-642-82486-9_1

Westrick, S.E., Moss, J.B., Fischer, E.K., 2023. Who cares? An integrative approach to

understanding the evolution of behavioural plasticity in parental care. Animal Behaviour

200, 225–236. https://doi.org/10.1016/j.anbehav.2022.10.005

Whirledge, S., Cidlowski, J.A., 2017. Glucocorticoids and Reproduction: Traffic Control on the
933  Road to Reproduction. Trends in Endocrinology & Metabolism 28, 399–415.
934  https://doi.org/10.1016/j.tem.2017.02.005
935 Whitfield, S.M., Bell, K.E., Philippi, T., Sasa, M., Bolaños, F., Chaves, G., Savage, J.M.,
936  Donnelly, M.A., 2007. Amphibian and reptile declines over 35 years at La Selva, Costa
937  Rica. Proceedings of the National Academy of Sciences 104, 8352–8356.
938  https://doi.org/10.1073/pnas.0611256104
939 Wingfield, J.C., Romero, L.M., 2011. Adrenocortical Responses to Stress and Their Modulation
940  in Free-Living Vertebrates, in: Comprehensive Physiology. John Wiley & Sons, Ltd, pp.
941  211–234. https://doi.org/10.1002/cphy.cp070411
942 Wintour, E.M., Coghlan, J.P., Oddie, C.J., Scoggins, B.A., Walteis, W. a. W., 1978. A
943  Sequential Study of Adrenocorticosteroid Level in Human Pregnancy. Clinical and
944  Experimental Pharmacology and Physiology 5, 399–403. https://doi.org/10.1111/j.1440-
945  1681.1978.tb00690.x
946 Wynne-Edwards, K.E., Edwards, H.E., Hancock, T.M., 2013. The Human Fetus Preferentially
947  Secretes Corticosterone, Rather than Cortisol, in Response to Intra-Partum Stressors.
948  PLOS ONE 8, e63684. https://doi.org/10.1371/journal.pone.0063684
949
950

A

log(rate of hormone release, pg/hr)

hormone   corticosterone   cortisol

***   **   NS   ***   ***

B

E. anthonyi   *P. terribilis*   *D. tinctorius*   *R. imitator*   *R. variabilis*