Unraveling the Genomic Diversity and Admixture History of Captive Tigers in the United States

Ellie E. Armstrong[1*&], Jazlyn A. Mooney[1,2*&], Katherine A. Solari[1], Bernard Y. Kim[1], Gregory S. Barsh[3,4], Victoria B. Grant[1], Gili Greenbaum[5], Christopher B. Kaelin[4], Katya Panchenko[1], Joseph K. Pickrell[6], Noah Rosenberg[1], Oliver A. Ryder[7], Tsuya Yokoyama[1], Uma Ramakrishnan[8], Dmitri A. Petrov[1,9,10], Elizabeth A. Hadly[1,11,12, 13&]

[1]Department of Biology, Stanford University, Stanford, CA, USA.
[2]Department of Quantitative and Computational Biology, University of Southern California, Los Angeles, CA, USA
[3]HudsonAlpha Institute for Biotechnology, Huntsville, AL 35806, USA
[4]Department of Genetics, School of Medicine, Stanford University, Stanford, CA, USA
[5]Department of Ecology, Evolution & Behavior, The Hebrew University of Jerusalem, Jerusalem, Israel
[6]Gencove, Inc., New York, NY, 11101, USA
[7]San Diego Zoo Wildlife Alliance, Escondido, CA, USA,
[8]National Centre for Biological Sciences, TIFR, Bellary Road, Bangalore, India 560065
[9]Chan Zuckerberg BioHub, San Francisco, CA USA.
[10]Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA, USA.
[11]Department of Earth System Science, Stanford University, Stanford, CA, USA
[12] Woods Institute for the Environment, Stanford University, Stanford, CA, USA
[13]Center for Innovation in Global Health, Stanford University, Stanford, CA, USA

*Contributed equally
&Corresponding authors emails: ellieearmstrong@gmail.com; jazlynmo@usc.edu; hadly@stanford.edu

Abstract

Genomic studies of rare and endangered species have focused broadly on describing diversity patterns and resolving phylogenetic relationships, with the overarching goal of informing conservation efforts. However, few studies have investigated the genomic diversity potentially housed in captive populations. For tigers (*Panthera tigris*) in particular, captive individuals vastly outnumber those in the wild, yet the diversity of the captive population remains largely unexplored. Here, we present the first large-scale genetic study of the private (non-zoo) captive tiger population in the United States (U.S.), also known as 'Generic' tigers. We find that the U.S. Generic tiger population has an admixture fingerprint comprising all six extant wild tiger subspecies (*P. t. altaica*, Amur; *P. t. tigris*, Bengal; *P. t. corbetti*, Indochinese; *P. t. jacksoni*,

Malayan; *P. t. amoyensis*, South China; *P. t. sumatrae*, Sumatran). We show that the Generic

tiger population has a comparable amount of genetic diversity to most wild subspecies,

relatively few private variants, and fewer deleterious mutations. We also observe inbreeding

coefficients that are similar to wild populations, suggesting that inbreeding in captive populations

is not pervasive, although there are some individuals within the Generic population that are

substantially inbred. Our results elucidate the admixture history of the Generic tiger population

in the U.S. Additionally, we develop a reference panel for tigers and show that it can be used

with imputation to accurately distinguish individuals and assign ancestry even with ultra-low

coverage (0.25×) data. The study and reference panel will provide a resource to assist in tiger

conservation efforts.

Main

The Anthropocene has been characterized by population declines, isolation, and

extinction, leading to a shift in the abundance and distribution of thousands of species on Earth

(1). The tiger (*Panthera tigris*) is one of the most iconic and captivating terrestrial species on the

planet and exemplifies the severe and rapid declines in population size and reduced

connectivity that threatens the survival of many species (2). Although some tiger populations

and subspecies have boasted recent recovery efforts, others have gone extinct in the wild (3) or

have been completely extirpated (4) due to human pressures. Today, tigers are composed of six

subspecies (*P. t. altaica*, Amur; *P. t. tigris*, Bengal; *P. t. corbetti*, Indochinese; *P. t. jacksoni*,

Malayan; *P. t. amoyensis*, South China; *P. t. sumatrae*, Sumatran) that have been genetically

and geographically separated for at least the last 10,000 years (5, 6). Their geographic ranges

span the Russian Far East and northeast China (*P. t. altaica*), Bangladesh, Bhutan, India, and

Nepal (*P. t. tigris*), Myanmar, Thailand, and Laos (*P. t. corbetti*), the island of Sumatra (*P. t.

sumatrae*), and the Malay Peninsula (*P. t. jacksoni*), respectively, while the South China tiger (*P.

t. amoyensis*) is extinct in the wild with the last sighting more than 30 years ago (7). All

subspecies have undergone severe population bottlenecks over the last century, primarily due

to human activities such as hunting and the expansion of agriculture, which have directly

reduced tiger numbers, habitat availability, and prey (8).

In contrast to wild populations, the global captive tiger population is now estimated to

include 15,000–20,000 individuals worldwide, a number that is at least five times larger than the

wild population (7). Most populations are largely unregulated, such as farmed and other

privately owned tigers, while others, such as those in accredited zoos, are bred with the

intention of serving as a diversity reservoir for dwindling wild populations. In the United States,

the Association of Zoos and Aquariums (AZA), manages several tiger populations as distinct

subspecies, specifically the Amur (1950s-present), Sumatran (1950s-present), Malayan (1980s-

present), and for a time the "Bengal" (white tigers; 1960s-2011) tiger subspecies.

The idea that captive populations may serve as diversity reservoirs is a persistent theme

and justification of the existence of captive populations (9). For example, the regulated Amur

tiger captive population was established in the early 1950s and has since increased to over

1,400 individuals in captivity worldwide (AZA; 10). Previous work which investigated diversity

levels of 12 microsatellites in the Amur captive population in North America found that the

population may be a reservoir of genetic diversity which is now extinct in the wild (11). This

notion suggests that captive breeding programs with effective management are able to maintain

genetic diversity and avoid inbreeding depression despite the small number of founders (in the

case of the Amur tigers, 29 males and 28 females; (10). Genomic perspectives on whether this

is true for many captive populations are rare and the conclusions are inconsistent across

species (12–17).

Though there are a large number of tigers in captivity, only a small fraction of these

tigers (less than 1%) reside in regulated or accredited facilities, such as accredited zoos (7).

Outside of regulated establishments, individuals and entities breed, own, and sell tigers both

legally and illegally. The formation of the privately-owned, captive population in the U.S. likely

coincided with the establishment of zoos and circuses in the early 1900s, since these entities

were known to exchange individuals (8), but ultimately the ancestry and establishment timing of

the captive, privately-owned population is unknown. Over time, the mission of AZA zoos began

to run counter to circuses and other private ownership practices, but by the 2000s, estimates

put the captive tiger population in the United States at greater than 5,000 individuals (8, 18).

Many of the unregulated facilities do not report basic or reliable population data and have

emerged as a concern globally with the illicit farming of big cats for profit being exposed in

Thailand (19), South Africa (20), and most recently the United States, which was popularized by

the release of Netflix's *Tiger King*. In the United States, up until the recent passing of federal

legislation (21; Big Cat Public Safety Act), there was very little oversight over the breeding,

owning, and selling of tigers.

The pace of breeding in privately owned tiger populations has prompted concerns of

inbreeding, especially since phenotypes (i.e. striped whites, snow whites, and golden tigers) that

rely on rare, recessive mutations (22, 23) are considered to be of high economic value. In fact,

the white tiger ("Bengal") breeding program was discontinued by the AZA when the intentional

selection for the single variant responsible for producing the white phenotype (22, 23) (occurring

naturally, but rarely, in wild Bengal tiger populations) was determined to not be relevant to

conservation initiatives. This population was also intentionally mixed with Amur tigers in an

attempt to prevent inbreeding depression before the program was ultimately discontinued (8). A

previous study documented additional signatures of admixture in captive, privately-owned tigers

from various localities using mitochondrial DNA (mtDNA) and microsatellites, and suggested

that captive tigers contain unique diversity compared to their wild and captive single-ancestry

counterparts (24). A separate study also posited that the captive Amur population may contain

diversity now extinct in the wild (11). However, neither of the previous studies examined the

extent of inbreeding or genetic load in the population using genomic data, which has

significantly more power than typical microsatellite datasets to answer such questions. It

remains unclear how or if this large population of privately-owned, captive tigers could possibly

fit into current management or conservation plans, and whether their genomes hold relic or

unique diversity, or on the other hand, if they show indications of severe inbreeding or high

genetic load.

For the purposes of conservation genomics, the most useful definition of genetic load is

one that describes a reduction in fitness at the population level due to deleterious mutations

(25). The genetic load of a population can affect overall population health and viability, and can

be modified by stochastic processes such as drift, admixture, and inbreeding (26, 27). Recent

genomic-scale work in wild populations has shown purging of deleterious mutations in some

groups (28–30), while other groups (31, 32) appear to have a large fraction of deleterious

mutations remaining. In fact, recent work in wild Bengal tigers has shown purging of deleterious

variation in smaller populations relative to larger populations, alongside continued inbreeding

depression due to high frequency deleterious alleles (33). However, there is a dearth of

literature on genetic load and inbreeding in captive tiger populations, or other tiger subspecies,

at the genomic scale. As genetic rescue, translocation, and captive breeding programs are

being implemented more frequently in endangered species for strategic management (34, 35), a

broader understanding of genetic diversity and genetic load within captive and wild tiger

populations warrants further investigation.

Here, we use genomic data to examine admixture and population structure, quantify

genetic diversity and genetic load, and investigate the extent of inbreeding in the captive,

privately-owned tiger population in the United States (also known as 'Generic' tigers) using

whole-genome sequence data obtained from individuals in accredited sanctuaries. These

sanctuaries do not breed, sell, or buy tigers, but house tigers previously rescued from

unregulated facilities or those that have been forfeited by private owners. We combine our

newly sequenced individuals with previously published data, resulting in a dataset representing

255 unique individuals across all six extant wild subspecies and U.S. Generic tigers (Table 1).

These data allow us to determine how admixture events have shaped the genomic landscape of
the privately-owned captive population. We compare the Generic tigers to their potential source
populations (tigers of single ancestry, i.e., subspecies), and examine how diversity is partitioned
across each group. Then, we test for potential signs of inbreeding and quantify the total amount
of deleterious variation in each population. Lastly, we show that ultra-low coverage (0.25×) data
can be imputed sufficiently using reference haplotypes from single ancestry tigers to determine
ancestry and perform individual identification. Our results demonstrate that low-coverage
sequencing and imputation provide a simple and cost-effective alternative compared to
microsatellites and custom SNP panels to identify the source populations and identity of illegally
traded individuals or wildlife materials from tigers and for long-term monitoring of captive and
wild populations.

Table 1: Number of individuals (duplicates removed) used in each analyses by method (imputed/unimputed) with brief description of filters used. See Methods and Supplementary Methods for additional and specific details.

Group	Total Individuals		Total	PCA	Admixture	Heterozygosity (unimputed, <20% missing)	ROH, IBD, Genetic Load (unrelated, >5X)	SFS (unrelated, >5X, oven sampling)	Local Ancestry (unimputed)
Amur	38	Unimputed	32	32	32	28	22	10 & 6	32
		Imputed	6	6	6				
Bengal	23	Unimputed	23	23	23	23	21	10 & 6	23
		Imputed	0	0	0				
Indochinese	6	Unimputed	6	6	6	6	6	6	6
		Imputed	0	0	0				
Malayan	22	Unimputed	22	22	22	22	20	10 & 6	22
		Imputed	0	0	0				
South China	4	Unimputed	1	1	1	1	1		
		Imputed	3	3	3				
Sumatran	18	Unimputed	18	18	18	18	15	10 & 6	18
		Imputed	0	0	0				
Generic	144	Unimputed	68	68	68	50	18	10 & 6	68
		Imputed	76	76	76				

Results

Population structure and ancestry in Generic tigers

Using both imputed and unimputed individuals, we investigated population structure and
ancestry of the Generic tigers using a combination of principal component analysis (PCA) and
supervised ADMIXTURE. Both the supervised ADMIXTURE analysis (Fig. 1A and
Supplementary Notes) and PCA (Supplementary Fig. S1), show that Generic tigers are admixed

between the six extant tiger subspecies, with individuals varying in the proportion of their

genome derived from each ancestral population (Fig. 1A). Investigation of mitochondrial

haplotypes revealed clustering with all but the Sumatran and South China haplotypes

(Supplementary Fig. S2-4). Local ancestry analyses (unimputed samples only) revealed tracts

from all subspecies except South China, which was not included due to its sample size (N = 1;

Supplementary Fig. S5). Overall, we found that none of the Generic tigers tested have single-

subspecies ancestry. Generic individuals, on average, are primarily admixed between the Amur

and Bengal tiger subspecies (mean: 38.6% and 28.2% of ancestry derived from each

subspecies, respectively) and derive the least ancestry from the Sumatran and Malayan

populations (mean: 11.1% and 5.0%, respectively; Fig. 1A). Our clustering analyses and PCA

results showed that the captive population clusters together by their dominant ancestry in both

PC space and the identity-by-state (IBS) tree (Fig. 1B; Supplementary Fig. S6C). We also

observe some clustering based on putative geographic origin in the United States and Canada

(Fig. 1C), though it is visually less distinguishable than the clustering by ancestry. These results

suggest that structure of the Generic population is primarily caused by the breeding events that

established the captive population in the United States and there is weak evidence for

subsequent geographic structuring, indicating that there is frequent exchange of individuals

between breeders. We see further evidence of historical admixture based on mitochondrial

haplotype results, where Generic individuals cluster with Amur, Indochinese, and Bengal tiger

subspecies. We observed only one Generic sample (SRR836354) grouping with Malayan tigers

in mitochondrial haplotype analyses, and none grouping with Sumatran or South China tigers

(Supplementary Fig. S2-4). Further information can be found in *Supplementary Notes*.

Interestingly, our analyses also identified several individuals previously labeled as

single-ancestry subspecies to be Generic individuals. These individuals were originally labeled

as Amur (SRR7651464-67, SRR7651470) and Bengal (SRR836354) subspecies in public

repositories, but we observe that they all have a mixed ancestry background based on both

PCA (Supplementary Fig. S17) and subsequent admixture analysis (Fig. 1A). These individuals

lack additional metadata but are presumed to have originated in Asia (Supplementary Table 1)

and thus are labeled as 'Asia' in the admixture plot (Fig 1A).

Fig. 1: Ancestral diversity in Generic tigers. **A)** ADMIXTURE analysis showing ancestry of all
individuals, wild and Generic. **B)** PCA of North American Generic tigers where color
corresponds to an individual's major ancestry component. **C)** PCA of Generic tigers where color
corresponds to the region of North America from which the individual originated. For panels B &
C one individual was removed as an outlier; details can be found in *Supplementary Notes*.

Tiger subspecies diversity and population history

Since the Generic tigers form a well-admixed population, we next investigated their

genetic diversity, inbreeding, and mutation load relative to the single-ancestry, wild subspecies.

Primarily, we wanted to determine whether Generic tigers were potential reservoirs of genetic

diversity that is absent in the wild, as has been previously suggested. Given that bottlenecks

have occurred both in wild populations and during the process of founding the various captive

tiger populations, the surviving diversity in the Generic population in comparison to wild

populations is unknown. To address this, we estimated the number of distinct alleles per-locus,

calculated the proportion of sites that are variant and segregating in the Generic tigers, but fixed

in the wild subspecies, and calculated observed heterozygosity for each individual using only

unimputed individuals (Fig. 2 & Supplementary Figs. S7-8). ADZE (36) estimates allelic and

shared diversity through a rarefaction approach. We show that trajectories for allelic diversity

suggested the Indochinese subspecies (Supplementary Fig. S7) contained the most diversity,

followed by the Bengal subspecies. However, due to the small sample size for Indochinese

tigers (N = 6), it is unclear whether their allelic diversity scales with sample size as it does in the

Bengal tiger subspecies (Supplementary Fig. S8).

Next, we asked whether there might be an enrichment of variant sites that are solely

segregating within the Generic population but fixed within any reference wild subspecies (Fig.

2A). We found that most sites that were segregating within the Generic tigers are also

segregating in at least one of the wild subspecies (Fig. 2A). Specifically, less than 10% of sites

are exclusively segregating within the Generic tigers (Fig. 2A). Finally, we examined

heterozygosity. Generic tiger heterozygosity fell well within the range of other wild subspecies

(Fig. 2B). Similar to what we observed with ADZE, Bengal and Indochinese tigers had the

highest heterozygosity. Taken together, our analyses suggest that the Generic population is not

a major reserve of unique genetic diversity, rather, it contains diversity that is present in the wild.

Fig. 2: Measurements of shared and within subspecies genetic diversity across tigers. **A)**
Proportion of segregating sites that are fixed in each wild subspecies compared to Generic
tigers. The mean and standard deviation are computed across 10 replicates of captive
individuals (unimputed individuals). South China is not included since there is only one
individual. The designation of "All" indicates the site is fixed across all wild subspecies but
segregating in the Generic population. **B)** Observed heterozygosity distributions across
unimputed individuals, we have applied a horizontal jitter to each data point (See
Supplementary Methods and Imputation and filtering for details).

Due to captive breeding practices and a small founder population, we examined whether

the Generic tiger population showed signs of inbreeding. Specifically, we estimated inbreeding

using multiple metrics, including runs of homozygosity (ROH), shared identity-by-descent (IBD)

segments, and inbreeding coefficients from both SNPs (F_{SNP}) and ROH (F_{ROH}) using only

unimputed data. We observed that the Amur population had, on average, the most Type C (i.e.,

long runs, informative about recent inbreeding) ROH and highest IBD-sharing (Fig. 3 and

Supplementary Figs. S9-10), suggesting that Amur tigers experienced more severe inbreeding

than the Generic population. Further, Amur tigers also had the largest amount of Type B (i.e.,

intermediate length) ROH, which indicates a long-term, small population size (37). Indeed, wild

Amur tigers were documented as having experienced an extreme bottleneck in the 1930s/1940s

(11, 38), possibly explaining these extreme patterns compared to other subspecies.

When examining two other metrics of inbreeding (F_{SNP} and F_{ROH}) we observed that the

Generic tiger population was once again not an outlier. Instead, most Generic individuals had

similar inbreeding values to individuals in the other subspecies (Fig. 3). However, there are a

few individuals within the Generic population that are quite inbred. The fluctuating levels of

inbreeding matched the previous analysis of heterozygosity, where we observed variable levels

of genome-wide heterozygosity across the Generic population (Fig. 2B). Additionally, we also

found that when examining the site frequency spectra (SFS), the Generic tigers also had the

largest proportion of singletons (Supplementary Fig. S11). The large variance in inbreeding

coefficients (Type C ROH, F_{ROH}, and F_{SNP}) suggest that in the case of Generic tigers, admixture

between distinct subspecies likely increased heterozygosity in some individuals, but inbreeding

within individual facilities eroded it (39). Taken together, our results show that the Generic

population is not any more inbred than wild tiger populations, nor is it more diverse.

Fig. 3: Quantification of different types of ROH across tiger subspecies and measurements of inbreeding. **(Top)** Length of different classes of ROH (A; short, B; intermediate, C; long) for each tiger subspecies and Generic tigers. **(Bottom)** Total proportion of genome in Type C ROH (F_{ROH}) for each individual per population. We have labeled each individual with their corresponding inbreeding coefficient measured as F_{SNP}, the darkest color (black) is the lowest value, and the brightest color (orange) corresponds to the largest value.

Tiger subspecies mutation load

We last sought to characterize the prevalence of putatively deleterious variation in the captive population, relative to each subspecies (unimputed samples). Variants were categorized as putatively deleterious if they were nonsynonymous, derived mutations with a SIFT score less than 0.05 (see *Supplementary Methods* for more detail). We implemented multiple approaches to count deleterious variants in the genome: 1) counting homozygous derived genotypes; 2) counting the total number of homozygous and heterozygous derived genotypes (counting

variants); and 3) counting alleles where we tabulate twice the number of homozygous derived

genotypes plus heterozygous genotypes (counting alleles).

Given that the Generic population primarily derived its ancestry from the Amur and

Bengal subspecies (Fig. 1A), we expected that the distribution of deleterious variation in

Generic tigers would be comparable to the Bengal and Amur subspecies. Indeed, the observed

distribution of both neutral and deleterious variants in Generic tigers was more similar to the

Amur tigers and Bengal tigers than other wild subspecies (Fig. 4 & Supplementary Fig. S12).

Overall, we observed similar levels of deleterious variation across Sumatran, Malayan,

Indochinese, and Bengal subspecies. Though the mean number of derived deleterious variants

was slightly, but consistently, higher in the Sumatran subspecies for all counting methods

(Supplementary Fig. S12). We found that, irrespective of subspecies, individuals with the most

putatively deleterious derived homozygotes also tended to have the largest inbreeding

coefficients, quantified with either F_{SNP} or F_{ROH} (Supplementary Fig. S14). Thus, inbreeding likely

led to an overall decrease in fitness for all tigers, irrespective of their ancestry. Our observation

that the Generic tigers had the lowest average numbers of derived deleterious mutations,

suggests that the Generic tigers may have been able to either purge some of the variants that

are deleterious in homozygous form being carried by the other subspecies, or the admixture

process reduced their frequency as homozygotes.

Next, we examined whether putatively deleterious and nonsynonymous homozygous

variation tended to be enriched within ROH versus outside of ROH. Recall that counting

homozygotes is relevant when we are interested in variants that could be recessive, here we

are asking whether putatively deleterious and nonsynonymous variants tend to be in ROH,

which would allow them to act recessively and expose them to selection. We identified a

significant enrichment of both nonsynonymous and putatively deleterious derived homozygotes

within ROH versus outside of ROH across all subspecies (Supplementary Fig. S15). We saw

that the Generic tigers were a clear standout in terms of carrying nonsynonymous derived

homozygotes within ROH (odds ratio ~2.9). However, the odds of putatively deleterious derived

homozygotes falling within ROH in the Generic tigers are comparable to any of the wild

subspecies. In the case of putatively deleterious derived homozygotes within ROH, the standout

group became the Indochinese tigers, though they also had the largest 95% CIs

(Supplementary Fig. S15). Taken together, these results support that we cannot predict how

admixture will affect the burden or distribution of nonsynonymous and putatively deleterious

derived homozygotes.

300
301
302 **Fig. 4**: Quantifying genetic load in each subspecies by counting derived neutral and deleterious
303 variants. **A)**: Scaled counts of derived neutral alleles in each tiger subspecies and Generic
304 tigers. **B)**: Scaled counts of derived deleterious alleles in each tiger subspecies and Generic
305 tigers. Extended figure (Supplementary Fig. S12) contains the count of homozygotes and count
306 of variants.
307
Creation of a reference panel for future tiger conservation

Given that we collated the largest genomic dataset of tigers to date, we also tested

whether the data were sufficient to create an accurate reference panel for low-coverage

sequencing and imputation. An imputation pipeline can be particularly useful in wildlife forensics

where the ancestry or identity of the individual is unknown, but also for cost-effective

sequencing of many individuals. Imputation is generally used in the context of low-coverage

data to increase the reliability of genotype calls and improve the accuracy of downstream

analyses, and has recently been advocated as an alternative to reduced representation

approaches in non-model species, which often amplify a small and potentially biased portion of

the genome (40). The primary goal of the reference panel was to assess whether we could

accurately impute ultra-low coverage data for the purpose of individual identification and

ancestry determination in tigers. The full data was split into a training set (reference panel)

composed of 106 single-ancestry individuals (5, 6) with coverage between 2×-43×. Then,

imputation was performed using *lowimpute* on 86 low- and ultra-low coverage (0.25×-6×)

samples (Supplementary Fig. S1). We tested accuracy in a set of nine individuals (six

unimputed and three imputed), where four of the individuals were also represented in the

reference panel (See Supplementary Notes for additional details). To test all samples, we

randomly downsampled reads from each individual to five different coverages (0.25×, 0.5×, 1×,

2×, 5×) in addition to no down sampling (using all the reads in a standard variant calling pipeline

without imputation). Downsampled reads were run through the imputation pipeline, and

accuracy of the imputation was tested using variant calls (via non-reference discordance; NDR),

ancestry predictions, and the ability to uniquely identify individuals.

We found that as sequencing depth increased, the accuracy of variant calls, as

measured by NDR, also increased (Supplementary Table 2). NDR ranged from 7.71% at the

highest depth (6×) to 14.49% at the lowest depth (0.25×). We also observed variance in NDR

depending on whether a sample was included in the reference panel, as expected, samples

from the reference panel had the lowest NDR (Supplementary Table 2). Despite the variance in

NDR, ancestry (Supplementary Fig. S16) and the ability to identify individuals (Supplementary

Table 4), even at ultra-low coverage (0.25×), remained accurate. Our results show that low-

coverage sequencing and imputation are a sufficient and low-cost alternative to high-coverage

sequencing or genotyping for the purposes of ancestry and individual identification using this

reference panel for tigers.

340
Discussion

Admixture and variation

We found that all Generic tigers sampled from the U.S. are admixed and contain

ancestry from all six extant tiger subspecies (Fig. 1A). Most Generic tigers derive a majority of

their ancestry from the Amur and Bengal subspecies (Fig. 1A), but individuals varied in the

proportion of ancestry derived from any one subspecies (Fig. 1A, Supplementary Fig. S5). The

length distribution of local ancestry segments (Supplementary Fig. S5) indicated that there are

likely very few individuals in the extant U.S. Generic tiger population that have directly

descended from recently introduced, single-ancestry sources. The timing and sources of these

introductions is unknown. Though we observe ancestry from South China tigers in many of the

captive samples, the low number of available samples for South China generally supports

additional investigations before definitive conclusions can be made about their ancestry

contributions to the Generic population.

Our findings differ from the results obtained from Luo et al. 2008, where 30 microsatellite

loci and mitochondrial (4kb) data were used to investigate samples from captive tigers (N = 105)

obtained between 1982 and 2002 from 14 countries (including the U.S.). By comparing their

data to voucher specimens (39 Amur; 2 South China; 33 Indochinese; 22 Malayan; 17 Bengal;

21 Sumatran), they found that 49 of the captive tigers belonged to single subspecies ancestry,

while 52 were found to be of admixed ancestry (24). A total of 18 tigers did not match the

suspected ancestry (i.e. the tiger was found to be admixed compared to the source reporting

single subspecies ancestry or vice versa). Luo and colleagues (2008) did not describe the origin

of the sampled individuals in detail, but it is possible that individuals of single-subspecies

ancestry were being actively mixed into the privately-owned captive tiger population when those

samples were collected. Alternatively, it is possible that the small number of microsatellites used

in the previous study have comparatively limited power to distinguish the complex ancestry of

the Generic tigers compared to thousands of SNPs, as has been the case in other systems (41–

43). Interestingly, Luo and colleagues (2008) found that captive tigers with mixed subspecies

ancestry primarily contained ancestry from the Indochinese tiger lineage, whereas the tigers

analyzed herein contained mostly Amur and Bengal ancestry. Additional genomes for captive

non-zoo individuals from different countries would provide insight into whether tiger ancestry

proportions are a useful fingerprint distinguishing continental or country-specific tiger

populations that could be leveraged for wildlife investigations.

While there is no doubt that the Generic tiger population carries some unique variation

simply because of the large number of individuals in captivity (i.e., census size), we have shown

in our sampling that the population does not contain more variation or more unique genetic

variation than its single-ancestry counterparts (Fig. 2, Supplementary Figs. S7-8), contrary to

previous studies (24). In other words, Generic tigers have average heterozygosity (Fig. 2B),

possess very few unique segregating sites (Supplementary Fig. S7-8), and when compared to

other subspecies, most sites that are segregating in the Generic population are also segregating

in the wild populations (Fig. 2A). While there may be some beneficial alleles circulating in the

Generic population, predicting which loci are adaptive is a challenging task for even the most

well-studied species, and implementing gene-based management strategies can also bring

about harmful side-effects (44, 45).

Deleterious variation

The Generic population contains less deleterious recessive homozygous variation and

relatively low amounts of Class C ROH relative to wild populations (Fig. 3 and Fig. 4). Given the

degree of admixture in the Generics, the low frequency of deleterious mutations is expected, yet

surprising given the purported inbreeding of 'farmed' tigers (46). However, Generic tigers also

have the largest enrichment of nonsynonymous variation (counting alleles, counting variants,

counting derived homozygotes) within ROH versus outside. In other words, though Generic

tigers have a comparable amount of nonsynonymous and putatively derived homozygotes

compared to their wild ancestors, they tend to carry more of the nonsynonymous variation within

ROH than their ancestors. Thus, the results of admixture on the genomic landscape in Generic

tigers are, in this case, not obvious. Heterozygosity and load are similar across all tigers, likely

due to a composite effect of admixture between distinct lineages increasing heterozygosity,

followed by severe bottlenecks of repeated founding events and deliberate inbreeding. The

observed outcome of admixture in the Generics is the result of chance but is not necessarily

predictive of other metrics of inbreeding, load, or heterozygosity, and also does not negate

possible effects of outbreeding, which we did not examine here. The increase in

nonsynonymous load within versus outside of ROH, and population-level heterozygosity falling

between major ancestry sources has also been observed in human populations, specifically in

admixed Latin American founder populations (39).

Ancestry inference for unknown individuals

The final objective of this study was to compile a reference dataset and imputation panel

capable of identifying the ancestry and identity of individuals. At present, tiger forensic work

relies primarily on mitochondrial and microsatellite data (47). Mitochondrial markers in felid

lineages contain nuclear mitochondrial inserts (numts), making interrogating these regions

difficult and error-prone (48, 49). Further, mitochondrial markers are only indicative of the

maternal lineage of an individual and are not able to identify individuals with mixed ancestry. For

complex admixture scenarios, examining only a handful of microsatellites may cause biased

results or in the case of species or subspecies-level hybridization distort ancestry signals (50–

53). As sequencing costs decline, arrays and panels are set to emerge as the predominant and

cost-effective way to query forensic and low-quality samples, compared to microsatellites (54,

55). Arrays and panels require either a sufficient number of SNPs (56, 57) or if a reduced set is

needed to be cost-effective, a very careful selection of SNPs which are reflective of ancestry or

relatedness. Low-coverage WGS provides an avenue for rapid and simple identification, which

could be performed on any number of sequencing machines from Illumina to the portable

Oxford Nanopore. The reference dataset and imputation pipeline from our study provide a

robust way to identify individuals and their ancestry rapidly and cost-effectively, even from very

low-coverage samples.

Conclusions

There has been ongoing discussion regarding how captive populations might contribute

to conservation, and particularly whether they harbor unique alleles beneficial to the survival of

the species or represent historical variation that is no longer present in the wild. Here, we have

shown that the captive, privately-owned tiger population does not contain significant unique

variation compared to the wild subspecies. Additional investigations into the remaining diversity

in captive zoo populations are unlikely to reveal additional diversity since many of these

lineages are the putative founders of the Generic tiger population. Several notable zoo

populations, such as giraffes (58) and orangutans (59) have sourced animals for captive

breeding programs from multiple, genetically distinct populations, subspecies, or species. If we

continue to consider captive populations as diversity reservoirs for wild species, additional

investigation into the potential outcomes and interplay of such processes (admixture, inbreeding

vs outbreeding depression) is extremely important. We further caution that deleterious variation

analyses, especially in non-model organisms, should be viewed skeptically. Deleterious

variation analyses generally rely on annotations from other organisms and the accurate

inference of an ancestral state, both of which can be imprecise analyses even in the most well-

studied model organisms (60–62). Computational annotations take a holistic approach to

determining the likely consequences of variation, but until functional validation of variants

occurs, genetic load should be integrated into management plans with extreme caution.

Cumulatively, our analyses are concordant with the known history of tiger subspecies in

terms of historic bottlenecks and recent inbreeding, yet we also reveal the unique history of the

large population of Generic tigers in the United States and provide a comprehensive overview of

all extant subspecies and available data. Most Generic tigers contain ancestry from all six wild

tiger subspecies in their genomes. Contrary to previous hypotheses, most Generic tigers do not

show signs of severe, recent inbreeding, nor do they hold unique diversity. Thus, the role they

might play (if any) in terms of conservation of genetic diversity is unclear. Barring outbreeding

concerns, we find no genetic reason why Generic tigers from the U.S. would not be useful in

augmenting tiger populations, since most individuals do not contain an excess of deleterious

mutations or appear to be inbred. Whether it is wise to keep the tiger subspecies separate to

preserve their genetic uniqueness, or whether certain circumstances warrant lineage mixing

remains to be seen for the tiger and many other species. South China tigers have already been

genetically rescued by another subspecies (63), and what many populations may require for

survival is more individuals, but where they are sourced is a critical choice. For example, here

we showed that the Sumatran tiger population in particular may be in need of conservation

action due to an excess of homozygous deleterious alleles (Fig. 4 Supplementary Fig. S12 &

13), but further investigation into the single-ancestry captive populations will further elucidate

the genetic resources available for tiger conservation and simulations could help determine the

best sources for possible translocations (33).

The U.S. Generic population is the product of unique evolutionary admixture events

which may help us better understand the consequences of mixing divergent, wild lineages in the

future. To encourage future study and aid illegal trade and trafficking investigations, we also

present a reference panel that in conjunction with imputation can accurately identify the

ancestry and identity of an individual using even ultra low-coverage sequencing, demonstrating

the validity of this approach. Low-coverage WGS is an attractive alternative to creating marker

panels, which often are affordable and accessible only with caveats (e.g. when running many

markers or many samples), that often exclude conservation budgets and sample needs. Taken

together, our results are not only informative for the conservation management of tigers, but

also provide a cost-effective path for others in the conservation community to follow for

generating a reference panel when using low-coverage sequencing for identification of

individuals and their ancestral origins.

Methods

Sample collection and sequencing

A total of 154 tiger samples were collected opportunistically during routine vet care from

sanctuary facilities by vet and sanctuary staff or from existing biobank collections

(Supplementary Table 1). All samples were extracted using a Qiagen DNeasy kit (Cat. No.

69504) and samples prepared using a modified Nextera library prep protocol (64). 77 of these

samples (listed as 'Unimputed' in Supplementary Data 1) were sequenced between

approximately 2× and 5× depth. The remaining 77 samples ('Imputed' in Supplementary Data 1)

were sequenced at approximately 0.25×. Tigers collected in sanctuaries or with unknown

ancestry are labeled as "Generic" for the purposes of this manuscript. We additionally collected

data on the putative location of birth of each tiger from the sanctuaries in North America. These

locations were then translated into one of six regions: West (Washington, Oregon, California,

Nevada, Montana, Idaho, Wyoming, Colorado, Utah), Southwest (Arizona, New Mexico, Texas,

Oklahoma), Midwest (North Dakota, South Dakota, Nebraska, Kansas, Minnesota, Iowa,

Missouri, Wisconsin, Illinois, Michigan, Indiana, Ohio), Southeast (Arkansas, Louisiana,

Tennessee, Mississippi, Alabama, Georgia, Florida, Kentucky, West Virginia, Virginia, North

Carolina, South Carolina), and Northeast (Maine, New Hampshire, Massachusetts, New York,

Pennsylvania, Maryland, Delaware, Connecticut, Vermont, Rhode Island, New Jersey) and

Canada (Canada was not subdivided since N = 2 and both individuals were from the same

facility).

Variant-calling and reference panel construction

An additional 100, publicly available (as of December 2019) tiger genome samples were

downloaded from NCBI. Reads were mapped to the GenTig1.0 genome (65) using BWA-MEM

v0.7.17 (66) and variant calling was subsequently performed by Gencove using the Genome

Analysis Toolkit (GATK) v4.1.4.1 (67) according to best practices. Initial variant calling was

performed on all samples available at the time, excluding those sequenced at 0.25×. All of these

samples are referred to as 'unimputed' for the purposes of this manuscript. Initial variant calling

yielded a total of 23,579,569 variable sites across the entire genome. We restricted calls to

biallelic sites using BCFtools v1.6 (68), and subsequently filtered for quality, missingness, and

depth. See *Supplementary Methods* for additional details. The final data set contained a total of

7,519,430 sites across unimputed individuals and a callable genome of 2,174,711,735 base

pairs.

In order to select individuals to build the reference panel and accurately split individuals

into groups for kinship estimation, we conducted Principal Component Analysis (PCA) to ensure

that all individuals in the unimputed dataset were clustering according to subspecies using

PLINK v2 (69). Individuals were subsequently split into ancestry groups to form the reference

panel, which included representatives from all six tiger subspecies, but no Generic individuals.

Further, we tested several methods for detecting relatedness using pedigreed individuals in the

dataset, which were subsequently used to identify and remove duplicates. Additional information

can be found in *Supplementary Methods*.

Imputation and filtering
Using only single-subspecies ancestry individuals verified here and by previous studies,

we developed a reference panel to impute variants for an additional 86 individuals (labeled as

'imputed' in Supplementary Data 1) through the *loimpute* pipeline developed by Gencove and

available at www.gencove.com (70). The 86 imputed individuals were composed primarily of

individuals sequenced at ultra-low coverage (N = 75; 0.25×), but also included two individuals

from a Canadian Zoo sequenced at approximately 3×, and an additional 9 samples that became

publicly available after the initial variant calling had been performed (see Supplementary Data 1

for details).

We combined the VCFs for imputed individuals with the unimputed individuals using

BCFtools *merge*. Because imputation emits a call for every site in the reference pipeline, we

restricted the merged sites VCF to retain only the quality sites identified after initial variant

calling and filtering. We further checked for imputation accuracy using concordance measures,

specifically non-reference discordance (71), and examined the accuracy of ancestry and

relatedness measures over a variety of coverages. Additional information can be found in

Supplementary Methods.

Genetic diversity
We used several approaches to investigate genetic diversity across tigers. The first

approach was to create equal sized groups of (N=10) individuals and tabulate the proportion of

sites that were SNPs in the Generic population and fixed in each wild subspecies. We used the

same reference groups from each subspecies and generated 10 replicate samples (with

replacement) of the Generic tigers to check whether these proportions varied across the

Generic population.

Observed heterozygosity was also calculated as the total number of heterozygous sites

divided by the number of callable sites (2,174,711,735) in the genome. Observed homozygous

sites were counted in each subspecies using VCFtools (72) using the '--het' flag and

heterozygous sites were calculated by subtracting the (O)HOM (observed homozygosity)

column from the NSITES (the total sites queried) column. The number of callable sites was

determined as the total number of base pairs minus the sites with mappability scores < 1 for

autosomal scaffolds. We additionally tested to see if heterozygosity was correlated with

missingness. See *Supplementary Methods* for details.

Runs of homozygosity (ROH)
545   GARLIC(73) was used to detect ROH. The error was set at 0.001, the window size at

700, and centromeres were set as 0,0 since no centromere information was available. To

ensure that ROH were not mistakenly called on regions with an excess of missing calls, each

ROH was then intersected with the number of callable sites in each window (see *Heterozygosity*

section for details). ROH larger than 100kb and containing callable sites within one standard

deviation ($\pm$0.066) of the mean coverage (0.655) were retained. ROH were binned into different

size classes A (short), B (intermediate), and C (long) per subspecies. Binning was based on the

use of a Gaussian mixture function that fits a model to the ROH length distribution within the

group. Type A ROH are typically indicative of linkage-disequilibrium blocks. Type B ROH are

informative about long-term small population sizes and cryptic relatedness. The presence of

Type C ROH indicates recent inbreeding in the population. F_{ROH} was computed as the total

fraction of the genome within a type C ROH.

Identity-by-descent segments (IBD)
558   IBD segments were called using TRUFFLEv1.38 with parameters '--segments --missing

1 --maf 0 --nofiltering' (74). Segments greater than 2Mb where the fraction of coverage by

callable sites (count listed above) was within one standard deviation ($\pm$0.032) of the mean

coverage (0.660) were retained. IBD scores were computed using the same approach from

Nakatsuka et al. (75), where we normalize shared IBD by the number of sampled individuals.

Putatively neutral and deleterious variation
We polarized and annotated sites by first filtering the data to scaffolds that corresponded

to main chromosomes from felCat8 (76); GCF_000181335.2). Coordinates were identified using

liftOver (77). The remaining sites, with felCat8 coordinates, were annotated with an impact and

consequence using VEP v92. We removed intergenic sites, splice acceptor, splice donor, splice

region annotations, and selected the most damaging impact for a given transcript. Sites were

coded as nonsynonymous (NS), synonymous (SYN), or loss of function (LOF) and SIFT scores

were added (78). Information from VEP was combined with a SIFT score to find putatively

neutral (SYN with SIFT score greater than 0.05) and putatively deleterious sites (NS or LOF

with SIFT score less than 0.05). See *Supplementary Methods* for details about annotations and

SIFT scores.

Ancestral bases were identified with Progressive CACTUS (79) alignment of the

GenTig1.0 genome with 9 other felid genomes; further details can be found in *Supplementary*

Methods. To assess load in each subspecies, we used only unimputed individuals with at least

5× coverage and less than 5% missing data. Lastly, we scaled the number of sites per individual

by subtracting the total number of variant sites across all individuals from missing sites to get

the total number of called sites. Each count was divided by the number of callable sites for that

individual; the proportion was multiplied by the average number of callable sites across all

subspecies.

Data Availability
Whole genome sequence data associated with this study has been deposited into NCBI under

bioproject number PRJNA976043 and will be released upon acceptance. Raw variant call files

(VCFs) have been uploaded to dryad repository https://doi.org/10.5061/dryad.k0p2ngff1.

Code Availability

All code associated with this project can be found at https://github.com/jaam92/Tigers/.

Acknowledgements

We would like to acknowledge each of the sanctuaries and zoos who contributed samples, knowledge, and time to this study, specifically the Performing Animal Welfare Society, the Exotic Feline Rescue Center, In-Sync Exotics, Turpentine Creek Wildlife Refuge, Carolina Tiger, and others. We especially acknowledge Joe Taft and Rebecca Rizzo of EFRC, Jackie Gai and Ed Stewart of PAWS, and Vicky Keahy of In-Sync, who have spent many hours discussing this project with us. This project would not have been possible without Bill Nimmo and Kizmin Reeves of Tigers in America. In addition, we thank the San Diego Frozen Zoo for access to their collections and their collaborative spirit. We also thank Sara Huston Katsanis for procuring the Carolina Tiger Rescue and Tiger World specimens and supporting the mentoring grant for V. Grant.

Author contributions

EEA, EAH, and UR conceived the study; KP, TY, VBG, and EEA generated the sequence data; EEA, JM, GG, KAS, BYK, CBK, JKP analyzed the data. All authors contributed to editing, guiding analyses, and writing the manuscript.

Declaration of interests

J.K.P. is an employee of Gencove, Inc.

References

1. R. Dirzo, *et al.*, Defaunation in the Anthropocene. *Science* **345**, 401–406 (2014).

2. J. Walston, *et al.*, Bringing the Tiger Back from the Brink—The Six Percent Solution. *PLoS Biol.* **8**, e1000485 (2010).

3. R. Tilson, K. Traylor-Holzer, Q. M. Jiang, The decline and impending extinction of the South China tiger. *Oryx* **31**, 243–252 (1997).

4. H.-R. Xue, *et al.*, Genetic ancestry of the extinct Javan and Bali tigers. *J. Hered.* **106**, 247–
257 (2015).

5. Y.-C. Liu, *et al.*, Genome-wide evolutionary analysis of natural history and adaptation in the
world's tigers. *Curr. Biol.* **28**, 3840–3849 (2018).

6. E. E. Armstrong, *et al.*, Recent Evolutionary History of Tigers Highlights Contrasting Roles
of Genetic Drift and Selection. *Mol. Biol. Evol.* **38**, 2366–2379 (2021).

7. S.-J. Luo, Y.-C. Liu, X. Xu, Tigers of the World: Genomics and Conservation. *Annu Rev*
Anim Biosci **7**, 521–548 (2019).

8. R. Tilson, P. J. Nyhus, *Tigers of the World: The Science, Politics and Conservation of*
Panthera tigris (Academic Press, 2009).

9. D. G. Kleiman, K. V. Thompson, C. K. Baer, *Wild Mammals in Captivity: Principles and*
Techniques for Zoo Management, Second Edition (University of Chicago Press, 2010).

10. K. Traylor-Holzer, AZA North American Regional Amur Tiger Studbook. *Minnesota*
Zoo/IUCN Conservation Breeding Specialist (2007).

11. P. Henry, *et al.*, In situ population structure and ex situ representation of the endangered
Amur tiger. *Mol. Ecol.* **18**, 3173–3184 (2009).

12. K. Ralls, J. D. Ballou, Managing genetic diversity in captive breeding and reintroduction
programs in *Transactions of the 57th North American Wildlife and Natural Resources*
Conference, (1992).

13. D. J. Fraser, How well can captive breeding programs conserve biodiversity? A review of
salmonids. *Evol. Appl.* **1**, 535–586 (2008).

14. J. R. Willoughby, *et al.*, The impacts of inbreeding, drift and selection on genetic diversity in
captive breeding populations. *Mol. Ecol.* **24**, 98–110 (2015).

15. H. Ito, R. Ogden, T. Langenhorst, M. Inoue-Murayama, Contrasting results from molecular
and pedigree-based population diversity measures in captive zebra highlight challenges
facing genetic management of zoo populations. *Zoo Biol.* **36**, 87–94 (2017).

16. R. Ogden, *et al.*, Benefits and pitfalls of captive conservation genetic management:
Evaluating diversity in scimitar-horned oryx to support reintroduction planning. *Biol.*
Conserv. **241**, 108244 (2020).

17. B. R. Wright, C. J. Hogg, E. A. McLennan, K. Belov, C. E. Grueber, Assessing evolutionary
processes over time in a conservation breeding program: a combined approach using
molecular data, simulations and pedigree analysis. *Biodivers. Conserv.* **30**, 1011–1029
(2021).

18. R. Garner, A Census of Federally Regulated Big Cat Populations within the United States
as of December 2020. *Journal of Zoological and Botanical Gardens* **2**, 517–530 (2021).

19. E. Cohen, The fall and reincarnation of Thailand's Tiger Temple. *International Journal of*
Tourism Anthropology **7**, 115–131 (2019).

20. V. L. Williams, A. J. Loveridge, D. J. Newton, D. W. Macdonald, A roaring trade? The legal
651   trade in Panthera leo bones from Africa to East-Southeast Asia. *PLoS One* **12**, e0185996
652   (2017).

653   21. R. M. Grijalva, *et al.*, BIG CAT PUBLIC SAFETY ACT [H. rept. 116-430 for H.r. 1380]
654   (2020) (May 10, 2023).

655   22. X. Xu, *et al.*, The genetic basis of white tigers. *Curr. Biol.* **23**, 1031–1035 (2013).

656   23. X. Xu, *et al.*, The genetics of tiger pelage color variations. *Cell Res.* **27**, 954–957 (2017).

657   24. S.-J. Luo, *et al.*, Subspecies genetic assignments of worldwide captive tigers increase
658   conservation value of captive populations. *Curr. Biol.* **18**, 592–596 (2008).

659   25. G. Bertorelle, *et al.*, Genetic load: genomic estimates and applications in non-model
660   animals. *Nat. Rev. Genet.* **23**, 492–503 (2022).

661   26. D. Charlesworth, J. H. Willis, The genetics of inbreeding depression. *Nat. Rev. Genet.* **10**,
662   783–796 (2009).

663   27. L. F. Keller, D. M. Waller, Inbreeding effects in wild populations. *Trends Ecol. Evol.* **17**,
664   230–241 (2002).

665   28. J. A. Robinson, C. Brown, B. Y. Kim, K. E. Lohmueller, R. K. Wayne, Purging of Strongly
666   Deleterious Mutations Explains Long-Term Persistence and Absence of Inbreeding
667   Depression in Island Foxes. *Curr. Biol.* **28**, 3487–3494.e4 (2018).

668   29. C. Grossen, F. Guillaume, L. F. Keller, D. Croll, Purging of highly deleterious mutations
669   through severe bottlenecks in Alpine ibex. *Nat. Commun.* **11**, 1001 (2020).

670   30. Y. Xue, *et al.*, Mountain gorilla genomes reveal the impact of long-term population decline
671   and inbreeding. *Science* **348**, 242–245 (2015).

672   31. L. Smeds, H. Ellegren, From high masked to high realized genetic load in inbred
673   Scandinavian wolves. *Mol. Ecol.* **32**, 1567–1580 (2023).

674   32. J. A. Mooney, C. D. Marsden, A. Yohannes, R. K. Wayne, K. E. Lohmueller, Long-term
675   Small Population Size, Deleterious Variation, and Altitude Adaptation in the Ethiopian Wolf,
676   a Severely Endangered Canid. *Mol. Biol. Evol.* **40** (2023).

677   33. A. Khan, *et al.*, Genomic evidence for inbreeding depression and purging of deleterious
678   genetic variation in Indian tigers. *Proc. Natl. Acad. Sci. U. S. A.* **118** (2021).

679   34. A. R. Whiteley, S. W. Fitzpatrick, W. C. Funk, D. A. Tallmon, Genetic rescue to the rescue.
680   *Trends Ecol. Evol.* **30**, 42–49 (2015).

681   35. K. A. Witzenberger, A. Hochkirch, Ex situ conservation genetics: a review of molecular
682   studies on the genetic consequences of captive breeding programmes for endangered
683   animal species. *Biodivers. Conserv.* **20**, 1843–1861 (2011).

684   36. Z. A. Szpiech, M. Jakobsson, N. A. Rosenberg, ADZE: a rarefaction approach for counting
685   alleles private to combinations of populations. *Bioinformatics* **24**, 2498–2504 (2008).

37.   T. J. Pemberton, *et al.*, Genomic patterns of homozygosity in worldwide human
populations. *Am. J. Hum. Genet.* **91**, 275–292 (2012).

38.   S. Alasaad, R. C. Soriguer, G. Chelomina, Y. P. Sushitsky, J. Fickel, Siberian tiger's recent
population bottleneck in the Russian Far East revealed by microsatellite markers. *Mamm.*
Biol. **76**, 722–726 (2011).

39.   J. A. Mooney, *et al.*, Understanding the Hidden Complexity of Latin American Population
Isolates. *Am. J. Hum. Genet.* **103**, 707–726 (2018).

40.   M. M. Watowich, *et al.*, Best practices for genotype imputation from low-coverage
sequencing data in natural populations. *Mol. Ecol. Resour.* (2023)
https:/doi.org/10.1111/1755-0998.13854.

41.   S. J. Zimmerman, C. L. Aldridge, S. J. Oyler-McCance, An empirical comparison of
population genetic analyses using microsatellite and SNP data for a species of
conservation concern. *BMC Genomics* **21**, 382 (2020).

42.   N. Liu, L. Chen, S. Wang, C. Oh, H. Zhao, Comparison of single-nucleotide polymorphisms
and microsatellites in inference of population structure. *BMC Genet.* **6 Suppl 1**, S26 (2005).

43.   C. Gärke, *et al.*, Comparison of SNPs and microsatellites for assessing the genetic
structure of chicken populations. *Anim. Genet.* **43**, 419–428 (2012).

44.   M. Kardos, A. B. A. Shafer, The Peril of Gene-Targeted Conservation. *Trends Ecol. Evol.*
33, 827–839 (2018).

45.   P. S. Miller, P. W. Hedrick, MHC Polymorphism and the Design of Captive Breeding
Programs: Simple Solutions Are Not the Answer. *Conserv. Biol.* **5**, 556–558 (1991).

46.   J. Guo, Tigers in trouble: year of the tiger. *Nature* **449**, 16–18 (2007).

47.   D. Karmacharya, *et al.*, Species, sex and geo-location identification of seized tiger
(Panthera tigris tigris) parts in Nepal—A molecular forensic approach. *PLoS One* **13**,
e0201639 (2018).

48.   J.-H. Kim, *et al.*, Evolutionary analysis of a large mtDNA translocation (numt) into the
nuclear genome of the Panthera genus species. *Gene* **366**, 292–302 (2006).

49.   W. Zhang, *et al.*, Highly conserved D-loop-like nuclear mitochondrial sequences (Numts) in
tiger (Panthera tigris). *J. Genet.* **85**, 107–116 (2006).

50.   L. Szatmári, *et al.*, A comparison of microsatellites and genome-wide SNPs for the
detection of admixture brings the first molecular evidence for hybridization between Mustela
eversmanii and M. putorius (Mustelidae, Carnivora). *Evol. Appl.* **14**, 2286–2304 (2021).

51.   A. Miralles, *et al.*, Inconsistent estimates of hybridization frequency in newts revealed by
SNPs and microsatellites. *Conserv. Genet.* **25**, 215–225 (2024).

52.   L. K. Vaughan, *et al.*, The use of plasmodes as a supplement to simulations: A simple
example evaluating individual admixture estimation methodologies. *Comput. Stat. Data*
Anal. **53**, 1755–1766 (2009).

53. B. M. Fitzpatrick, Estimating ancestry and heterozygosity of hybrids using molecular
markers. *BMC Evol. Biol.* **12**, 131 (2012).

54. R. Ogden, Unlocking the potential of genomic technologies for wildlife forensics. *Mol. Ecol.*
Resour. **11 Suppl 1**, 109–116 (2011).

55. M. Kayser, P. de Knijff, Improving human forensics through advances in genetics,
genomics and molecular biology. *Nat. Rev. Genet.* **12**, 179–192 (2011).

56. N. A. Rosenberg, L. M. Li, R. Ward, J. K. Pritchard, Informativeness of genetic markers for
inference of ancestry. *Am. J. Hum. Genet.* **73**, 1402–1422 (2003).

57. N. A. Rosenberg, Algorithms for selecting informative marker panels for population
assignment. *J. Comput. Biol.* **12**, 1183–1201 (2005).

58. L. B. Lackey, Giraffe studbook Giraffa camelopardalis North American regional/global.
Association of Zoos and Aquariums and Disney's.

59. G. L. Banes, B. M. F. Galdikas, L. Vigilant, Reintroduction of confiscated and displaced
mammals risks outbreeding and introgression in natural populations, as evidenced by
orang-utans of divergent subspecies. *Sci. Rep.* **6**, 22026 (2016).

60. V. Pejaver, *et al.*, Calibration of computational tools for missense variant pathogenicity
classification and ClinGen recommendations for PP3/BP4 criteria. *Am. J. Hum. Genet.* **109**,
2163–2177 (2022).

61. L. Liu, *et al.*, Biological relevance of computationally predicted pathogenicity of noncoding
variants. *Nat. Commun.* **10**, 330 (2019).

62. H. Li, *et al.*, Exome variant discrepancies due to reference-genome differences. *Am. J.*
Hum. Genet. **108**, 1239–1250 (2021).

63. W. Zhang, *et al.*, Sorting Out the Genetic Background of the Last Surviving South China
Tigers. *J. Hered.* **110**, 641–650 (2019).

64. M. Baym, *et al.*, Inexpensive multiplexed library preparation for megabase-sized genomes.
PLoS One **10**, e0128036 (2015).

65. E. E. Armstrong, *et al.*, Genome Report: Chromosome-level draft assemblies of the snow
leopard, African leopard, and tiger (Panthera uncia, Panthera pardus pardus, and Panthera
tigris). *bioRxiv*, 2022.04.26.489474 (2022).

66. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv [q-bio.GN] (2013).

67. A. McKenna, *et al.*, The Genome Analysis Toolkit: a MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

68. P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *Gigascience* **10** (2021).

69. C. C. Chang, *et al.*, Second-generation PLINK: rising to the challenge of larger and richer
datasets. *Gigascience* **4**, 7 (2015).

70. K. Wasik, *et al.*, Comparing low-pass sequencing and genotyping for trait mapping in
pharmacogenetics. *BMC Genomics* **22**, 197 (2021).

71. R. Arthur, J. O'Connell, O. Schulz-Trieglaff, A. J. Cox, Rapid genotype refinement for
whole-genome sequencing data using multi-variate normal distributions. *Bioinformatics* **32**,
2306–2312 (2016).

72. P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
(2011).

73. Z. A. Szpiech, A. Blant, T. J. Pemberton, GARLIC: Genomic Autozygosity Regions
Likelihood-based Inference and Classification. *Bioinformatics* **33**, 2059–2062 (2017).

74. A. Dimitromanolakis, A. D. Paterson, L. Sun, Fast and Accurate Shared Segment Detection
and Relatedness Estimation in Un-phased Genetic Data via TRUFFLE. *Am. J. Hum. Genet.*
105, 78–88 (2019).

75. N. Nakatsuka, *et al.*, The promise of discovering population-specific disease-associated
genes in South Asia. *Nat. Genet.* **49**, 1403–1407 (2017).

76. J. U. Pontius, *et al.*, Initial sequence and comparative analysis of the cat genome. *Genome*
Res. **17**, 1675–1689 (2007).

77. M. Haeussler, *et al.*, The UCSC Genome Browser database: 2019 update. *Nucleic Acids*
Res. **47**, D853–D858 (2019).

78. P. C. Ng, S. Henikoff, SIFT: Predicting amino acid changes that affect protein function.
Nucleic Acids Res. **31**, 3812–3814 (2003).

79. J. Armstrong, *et al.*, Progressive Cactus is a multiple-genome aligner for the thousand-
genome era. *Nature* **587**, 246–251 (2020).