GATA1-defective immune-megakaryocytes as possible drivers of idiopathic pulmonary fibrosis

Francesca Gobbo[1,3], Maria Zingariello[2], Paola Verachi[3], Mario Falchi[4], Francesca Arciprete[2], Fabrizio Martelli[5], Angelo Peli[6], Maria Mazzarini[3,7], Jeff Vierstra[7], Carolyn Mead-Harvey[8], Amylou C. Dueck[8], Giuseppe Sarli[1], Stefano Nava[9,10], Giacomo Sgalla[11], Luca Richeldi[11] and Anna Rita Migliaccio[2,7]

[1]Department of Veterinary Medical Sciences, University of Bologna, Ozzano dell'Emilia (Bologna) 40064, Italy; [2]Unit of Microscopic and Ultrastructural Anatomy, Department of Medicine, University Campus Bio-Medico, Rome 00128, Italy; [3]Department of Biomedical and Neuromotor Sciences, University of Bologna, Bologna, Italy; [4] National Center HIV/AIDS Research, Istituto Superiore di Sanità, Rome 00161, Italy; [5]National Center for Preclinical and Clinical Research and Evaluation of Pharmaceutical Drugs, Istituto Superiore di Sanità, Rome 00161, Italy; [6]Department for Life Quality Studies, University of Bologna, Rimini Campus, Rimini 47921, Italy; [7]Altius Institute for Biomedical Sciences, Seattle, WA 98121, USA; [8]Mayo Clinic, Department of Quantitative Health Sciences, Division of Clinical Trials and Biostatistics, Scottsdale, AZ 85259, USA; [9]IRCCS Azienda Ospaliero-Universitaria di Bologna, Respiratory and Critical Care Unit, Bologna 40138, Italy; [10] Department of Medical and Surgical Sciences, University of Bologna, Bologna 40138, Italy; [11]Department of Medical and Surgical Sciences Fondazione Policlinico Universitario "A. Gemelli" IRCCS, Università Cattolica del Sacro Cuore, Roma 00168, Italy.

Corresponding Author: Anna Rita Migliaccio, Altius Institute for Biomedical Sciences, Seattle, WA, USA. amigliaccio@altius.org

Word counts: Title: 10 words, abstract: 148 words, text 9157 words, references: 83, Figures 13 and Tables 2. **10** Supplementary Figures + 3 supplementary Tables (plus 2 links for a total of 12 movies on Zenodo) for a total of 15 supplementary items.

Summary

Idiopathic pulmonary fibrosis (IPF) is a progressive fibrotic lung disorder with limited
therapeutic options. Insufficient understanding of driver mutations and poor fidelity of
currently available animal models has limited the development of effective therapies. Since
GATA1 deficient megakaryocytes sustain myelofibrosis, we hypothesized that they may also
induce fibrosis in lungs. We discovered that lungs from IPF patients and *Gata1^{low}* mice
contain numerous GATA1negative immune-poised megakaryocytes that, in mice, have
defective RNA-seq profiling and increased TGF-β1, CXCL1 and P-selectin content. With
age, *Gata1^{low}* mice develop fibrosis in lungs. Development of lung fibrosis in this model is
prevented by *P-selectin* deletion and rescued by P-selectin, TGF-β1 or CXCL1 inhibition.
Mechanistically, P-selectin inhibition decreases TGF-β1 and CXCL1 content and increases
GATA1positive megakaryocytes while TGF-β1 or CXCL1 inhibition decreased CXCL1
only. In conclusion, *Gata1*^{low} mice are a novel genetic-driven model for IPF and provide a
link between abnormal immune-megakaryocytes and lung fibrosis.

Keywords

Idiopathic pulmonary fibrosis, megakaryocytes, GATA1, TGF-β, P-selectin, CXCL1.

Introduction

Idiopathic pulmonary fibrosis (IPF) is a chronic, progressive form of pneumonia associated with fibrosis of unknown etiology[1]. IPF represents the prototypic form of progressive pulmonary fibrosis (PPF), also associated with other non-idiopathic conditions. IPF occurs primarily in older adults (usually >60 years of age) and leads to progressive deterioration of pulmonary function, and eventual death from chronic or acute respiratory failure[1,2]. Its incidence is 3-9 cases every 100,000 inhabitants and manifests itself prevalently in males with known risk factors such as smoking[3]. Studies of IPF and PPF pathobiology are hampered by limited availability of lung biopsies due to the fact that lung biopsy is no longer needed for diagnosis. While animal models of IPF/PPF have been proposed, the fidelity between the human disease and the currently available models is in general poor. Current animal models employed to study the pathobiology and to identify candidate treatments for IPF are animals (usually mice or rats) forced to breath bleomycin[4], a chemotherapeutic compound that has lung toxicity as its major adverse effect[5]. Animals subjected to bleomycin develop acute inflammation and fibrosis in the lungs which eventually leads to their death within one-two weeks. In spite of its widespread use, bleomycin has several limitations which reduce its clinical relevance. First, the aggressive progression of fibrosis in this model does not resemble that of IPF, where fibrosis develops slowly over many years. Second, IPF usually affects individuals without previous exposures to chronic inflammatory insults, suggesting that inflammation represents a confounding factor in translating the results from these models to humans. Third and most importantly, the lesions developed in this model are associated with acute alveolar damage, including edema, and have a bronchiolo-centric localization, likely the consequence of the intra-tracheal route of the drug administration[6]. By contrast, in IPF fibrosis occurs mostly in the basal lobes of the lung, is predominantly subpleural, and tends to extend to the para-septal parenchyma generating the typical honeycomb pattern revealed by high-resolution computed tomography (HRCT) of the chest[7]. Such different anatomic localization strongly suggests that the cells responsible for fibrosis in the animal models and in IPF patients are different. Nevertheless, studies in bleomycin mice, validated by observations in patients, have implicated abnormally high levels of several pro-fibrotic proteins, such as transforming growth factor-β (TGF-β)[8], interleukin-8 (IL-8)[9], and the adhesion receptor P-selectin (P-SEL)[10], among others, in the etiology of IPF. In the case of TGF- β, the availability of chemical inhibitors[11] and validation experiments demonstrating that mice forced to breath adenoviruses containing TGF-β develop fibrosis in the lung[12,13],

have inspired clinical trials with pirfenidone and nintedanib, two pleiotropic proinflammatory

inhibitors also targeting TGF-β, in IPF[14–17]. However, these two drugs have been shown to be

effective in slowing down the decline of lung function, but not in preventing disease

progression in most IPF patients[14–17]. Therefore, the only cure for IPF is lung transplantation,

an option available to only a limited number of patients[1–3]. The clinical needs of IPF patients

highlight the need for novel, possibly genetically driven, animal models which more closely

resemble the pathological mechanisms of IPF in patients. It is conceivable that these models

may also shed light on the genetic drivers for IPF which are still largely unknown. In fact,

although familial clustering and association studies suggest that the disease has a genetic

basis[3,18], a causative genetic link is yet to be established. The identification that loss-of-

function mutations in the *telomerase* gene[19], or in genes, such as *ZCCH8*, which encode

proteins required for expression of the *telomerase*[20], in patients with familiar IPF raised great

hope that the genetic cause of the disease had been finally discovered[21]. However, induction

of telomerase-deficiency in mice increases the senescence of both hematopoietic and alveolar

stem cells but the mice, although more susceptible to bleomycin-induced lung fibrosis, do not

develop IPF at baseline[22,23]. More recently, loss-of-functions mutations in genes encoding

surfactant proteins, the expression of which is restricted to alveolar type II cells, have been

found in 2-25% of the patients with sporadic and familiar IPF[24]. The pathobiology role of the

SPTFC mutations in the etiology of IPF was validated by the observation that Tamoxifen-

driven conditional knock-in mice harboring one of the *SPTFC* mutations (isoleucine-to-

threonine at codon 73 (I73T)) found in the patients develop an aggressive IPF phenotype and

die within few weeks from the administration of the drug[25]. However, consideration of

possible founder effects suggests that the mutation status (either in telomerase-related or

surfactant protein genes), is known for only 4% of all the acquires and congenital cases of

IPF[26]. The fact that the driver mutations for >90% of the patients is not known, highlights the

need for additional studies to identify other pathobiology pathways that may sustain the

disease.

Recent experiments have demonstrated that activation of *c-Jun*, a gene downstream to TGF-

β, induces fibrosis in multiple organs, including bone marrow (BM) and lung, in mice[27,28]. In

the case of myelofibrosis (MF), the most severe of the Philadelphia-chromosome negative

myeloproliferative neoplasms[29,30], c-jun is activated by increased microenvironment

bioavailability of TGF-β produced by abnormal megakaryocytes (MK)[31,32]. In fact, the MK in

the BM of patients with myelofibrosis remain immature due to a RSP14 ribosomopathy

induced by the driver mutations that reduces the translation of the mRNA for GATA1[33,34], the

transcription factor essential for their proper maturation[35]. The causative role of this abnormal

MK in the etiology of the disease has been validated by experiments demonstrating that mice

expressing either the driver mutations or low levels of *Gata1* (*Gata1^low mice)* only in MK[36]

develop myelofibrosis with age[37]. The abnormalities of human and murine GATA1

hypomorphic MK include a content greater than normal of TGF-β[38], CXCL1, the murine

equivalent of IL-8[39], and P-SEL[40], three of the proinflammatory proteins that have been

implicated also in the development of IPF.

MK are important cellular components of the lungs where they exert the function to release

platelets (Plt) intravascularly[41,42] and to participate in immune reactions against opportunistic

infections[43,44]. In fact, by leveraging single cell RNA-seq and other methods, recent studies

have demonstrated the existence of three distinct types of MK, each one exerting a different

function[44–46]: The BM of adult mice and humans contains the Plt-producing, which

correspond to the cells previously defined as mature MK, and the niche-supportive and

immune MK, both of which have immature morphology. The Plt-producing MK are the most

frequent, have high levels of ploidy, express genes related to thrombopoiesis, including the

transcription factor GATA1, and are identified by the CD42b/ARNL markers. Niche-

supportive MK are mostly >=8N and express high levels of PF4 and TGF-β, and possibly of

other pro- inflammatory cytokines, which regulate growth of hematopoietic stem cells, and

low levels of GATA1. These cells express the CD61 and MYLK4 markers and may be

induced in vitro by thrombopoietin to acquire a mature morphology and to produce Plt.

Immune-poised MK are low ploidy with signatures of inflammatory response and myeloid

leukocyte activation genes. These cells can be identified by the CD41 and LPS1 markers by

confocal microscopy and by the CD41 and CD53 markers by flow cytometry. They have

phagocytic activity and can stimulate T cell expansion and have been identified in lungs

where their numbers increase in response to infectious challenges and are activated by

neutrophil netosis to phagocytize the infective agents[43,44]. Although previous studies had

already implicated MK abnormalities in the development of fibrosis in the lung of

bleomycin-treated mice[47], the pathobiological role of these cells in IPF is far from

established. Based on the driving role exerted by GATA1 hypomorphic MK in establishing

fibrosis in the BM[33,34,37], we hypothesized that IPF may be driven by a specific MK

subpopulation with an abnormally low GATA1 content. Here, we tested this hypothesis by

comparing the MK subclasses and their GATA1 content in the lung from patients with IPF

and the non-diseased tissue of lung cancer patients and assessed whether lungs from mice

carrying the *Gata1*[low] mutation contain abnormal MK and became fibrotic with age.

Results

The lung from patients with IPF contains an excess of GATA1 hypomorphic megakaryocytes biased for the immune lineage

To test the role of MK in the development of IPF, we first compared the frequency and GATA1 content of MK (CD42b[pos] cells) in lungs from IPF patients with those observed in non-diseased sections from patients with lung cancer (three subject per group) by confocal microscopy (**Figures 1, S1A**). Available clinical information on the patients is described in **Table S1**. Hematoxylin-Eosin and reticulin staining of consecutive sections confirmed abnormal architecture and presence of fibrosis in sections from IPF patients while the overall normal architecture of the sections from No-IPF lungs indicated that they did not contain obvious cancerous tissues (**Figure 1A**). CD42[pos] MK were identified with *similar* low frequency in sections from IPF and No-IPF lungs. In No-IPF patients, MK were detected inside blood vessels and most of them (80%) contained robust nuclear levels of GATA1 (**Figures 1B, S1A**). By contrast, in IPF-patients, MK were detected within the lung parenchyma in close association with alveolar epithelial cells and <20% of them contained detectable levels of GATA1 (**Figures 1B,C, S1A**). To be noted that, as expected, none of the lung cells which were not stained by CD42b expressed GATA1.

The identity of the MK subpopulations present in the lung was assessed by confocal microscopy studies with markers specific for the immune-, HSC-supportive and Plt-poised MK (**Figures 2, S1B**). All MK subpopulations were recognized in the lung from both No-IPF and IPF patients. Again, in No-IPF patients, MK were localized prevalently within blood vessels while in IPF patients they were localized near the alveoli. Interestingly, the frequency of immune-MK was significantly greater (by 2-fold, $p<0.05$) in lungs from IPF vs. No-IPF patients while that of HSC-supportive and Plt-poised MK was similar in the two groups.

Megakaryocytes in the lungs from Gata1[low] mice have an immature morphology and became biased for the immune-lineage with age

To assess the effects of reduced GATA1 content on the MK subpopulations present in the lung, we compared frequency, maturation state and lineage fate of MK present in mice

harboring the *Gata1*^low^ mutation and their wild-type (WT) littermates (**Figures 3,S2,S3** and

Table 1). MK (CD42b^pos^ cells) are readily identified in lungs from both *Gata1*^low^ and WT

littermates by both immune-histochemistry and -fluorescent analysis (**Figures 3A,B,**

S2A,S3). As expected, the nucleus of <10% of CD42b^pos^ cells from the lung of *Gata1*^low^ mice

contains GATA1 while this transcription factor is clearly detected in the vast majority

(>90%) of the MK from the WT littermates (**Figures 3B, S3B**). Also in these mouse studies,

expression of GATA1 was restricted to the cells in the lung which express the MK marker

CD42b.

The localization and maturation state of MK in lung sections from *Gata1*^low^ and their WT

littermates was determined by morphological and flow cytometry observations (**Figures 3, S2**

and **Table 1**). MK were observed either intravascularly or within the perivascular interstitium

in the lungs from both WT and *Gata1*^low^ littermates (**Figure S2A**). Most of the CD42b^pos^

cells in the lungs from WT mice displayed a cytoplasm with clearly divided into Plt

territories (Plt-forming MK, PfMKs) and appeared small, probably because they had already

released Plt which surrounded them (**Figures 2A,S3A, Table 1**). When detected in the

alveolar septa, WT MK were arranged into clusters (**Figure S3A**). By contrast, the MK in the

lung from *Gata1*^low^ mice had an immature morphology which included a circular shape and

reduced levels of Plt territories (**Figures 3A, S3A, Table 1**). Only a minority of the *Gata1*^low^

MK were PfMKs surrounded by CD42b positive Plt (**Figure S3A).** These *Gata1*^low^ PfMKs

were found near blood vessels inside the septa, had a reduced cytoplasm which extended

itself between the epithelial cells (**Figure S3A**).

Statistical analyses of the frequency and maturation state of the MK observed in the lungs

from a large cohort of mice indicate that, without controlling for the effects of age or sex,

indicated that the difference in the frequency of total, small (both reduced in *Gata1*^low^ mice)

and large (increased in *Gata1*^low^ mice) MK in the lung from WT and *Gata1*^low^ mice is

statistically significant (total p=0.001, small p<0.001, large p=0.007) (**Table 1**).

The morphological observations on the MK maturation state were confirmed by flow

cytometry studies on the cell ploidy state determined by CD42b and propidium iodide

staining. There was a trend indicating that the lung of *Gata1*^low^ mice contain less total cells

and more CD61^pos^/CD41^pos^ MK by flow cytometry (**Figure 3C,D**). However, these the

interpretation of these data is biased by the extensive manipulation necessary to prepare

single cell lung suspensions and by the low number of mice (3-5) analyzed. By flow

cytometry, we determined that, as expected[46]*,* the ploidy of CD42b^pos^ cells in WT lungs

ranged from 2N up to 64N with approximately 60% of the cells displaying a DNA content

>8N (**Figure 3E**). By contrast, the DNA content of the majority (**~80**%) of CD42b^{pos} cells

from *Gata1^{low}* lungs was 4N with few (<20%) MK having a DNA content >8N, confirming

that the CD42b cells from the lung of *Gata1^{low}* mice are less mature than the WT ones.

In the case of MK, low ploidy cells may represent either Plt-poised MK at the beginning of

their maturation process or MK poised to exert immune- or HSC-supporting functions. To

assess whether enrichment for low ploidy cells observed in the lung of *Gata1^{low}* mice might

underly a switch among subpopulations, confocal microscopy studies with markers specific

for immune-, HSC-supportive and Plt-poised MK subpopulations were performed (**Figures**

4D, S4). These studies included WT mice as control and analyzed age (young and old) as

independent variable. MK with a HSC-supportive or Plt-poised phenotype were recognized in

the lungs from both young and old WT and *Gata1^{low}* mice at a frequency significantly greater

than normal at both ages (**Figures 4,S4**). By contrast, immune-MK were present in robust

numbers in the lung from old *Gata1^{low}* mice but were barely detectable in those from young

Gata1^{low} mice and from young and old WT mice (**Figures 4,S4**).

These results indicate that the lungs from old Gata1^{low} mice contain increased levels of

immune-MK vs. normal.

Megakaryocytes from Gata1^{low} mice express a defective immune-signature

To understand the consequences of reduced GATA1 content on the functions of the MK,

CD41^{pos} cells (>95% CD41^{pos} by re-analyses) purified from the BM and lung of 15-months

old WT and *Gata1^{low}* mice were profiled by RNAseq (**Figures 5, S5**). Principal component

analysis (PCA**)** of the RNAseq data from individual mice indicate a strong separation and

clustering among groups, with the differences between tissue of provenience (BM or lung)

accounting for approximately 79% of the variance in the expression data (**Figures 5A, S5A**).

Overall, the CD41^{pos} cells from the BM and lung of WT mice expressed significantly

different levels of 11,828 genes (3,185 upregulated, 8,643 downregulated) while the number

of genes significantly expressed at different levels by CD41^{pos} cells from BM and lung of

Gata1^{low} mice were 17,819 (4,848 upregulated, 2,919 downregulated). Comparison of WT

and *Gata1^{low}* cells from the same origin revealed that the mutation induced extensive changes

in gene expression in both organs. The genes significantly different in *Gata1^{low}* BM vs WT

were 3,257 (1,861 upregulated, 1,396 downregulated) while those in the lung were 2,126

(818 upregulated and 1,308 downregulated) (**Figure S5B**). Interestingly, although both WT

and *Gata1*[low] MK were purified according to the CD41 marker, the CD41[pos] cells from the

lung of *Gata1*[low] mice expressed levels of *CD41* (*Itga2b*) significantly greater than those of

thee corresponding cells from WT mice (foldchange=2.329, p=0.001), while the expression

of *CD61* (*Itgb3*) and *CD42b* (*GpI2b*) was similar between the two groups (**Figure 5B, Table**

2), confirming the morphological and flow cytometry observations indicating that the

Gata1[low] lung from old mice are enriched for immature cells. Comparison of the expression

of representative genes characterizing the signatures of the three recently described Mk

subpopulations[46] indicates that most of the expression changes are found among genes that

characterize immune-MK (*Clc3*, *Clec4a2*, *Clec4e*, *Clec5a*, *Ctsd*, etc, upregulated and *Cxcl1*,

Ccxl10, *Il1b*, *Ccl3*, *Cxcl5*, *Lcn2*, etc, downregulated) (**Figure 5B** and **Table 2**). Overall, the

abnormalities in the expression signature of MK purified from the BM and lung of *Gata1*[low]

mice were similar (**Table 2**), suggesting that they were mostly driven by insufficient

expression of the transcription factor rather than by pro-inflammatory cytokines present in the

lung microenvironment.

Collectively, these results indicate that the immune-MK from the BM and lung of old

Gata1[low] mice are functionally corrupted.

Both WT and Gata1*[low] *mice are prone to display inflammation in lung with age

Since comparative studies on the baseline pathologies expressed by mice of different strains

indicate that CD1 mice, the background where the *Gata1*[low] mutation is hosted, suffer from

acute inflammatory pathology, including pneumonia[48], we first determined whether the

presence of the *Gata1*[low] mutation would exacerbate the baseline inflammation that is

expected to be present in the lung from CD1 mice as they age (**Figures 6,S2** and **Table 1**).

Histological features consistent with interstitial pneumonia are observed in the lungs from a

significant number (>40%) of both the WT and *Gata1*[low] mice analyzed (**Figures 6, S2B,** and

Table 1). More specifically, interstitial pneumonia (mild, moderate and severe) characterized

by infiltration of lymphocytes, macrophages and neutrophils in the alveolar septae and in the

peri-bronchial-vascular spaces, was observed in 9 out of 12 WT mice analyzed and in 27 out

of 37 of the *Gata1*[low] mice (**Figure 6, Table 1**). In *Gata1*[low] mice, pneumonia extended from

the alveolar septae to the subpleural interstitium (**Figure S2B**). Pneumonia was associated

with the presence of infiltrating neutrophils that were often found surrounding the MK

(**Figure S2C**). Without controlling for the effects of age or sex, there are no statistically

significant differences in severity of lung inflammation between the two genotypes (**Table 1**).

Sensitivity analyses adjusting for age and for sex also failed to find a statistically significant

difference (**Table 1**).

These data indicate that the presence of the *Gata1^{low}* mutation does not exacerbate the

baseline interstitial pneumonia developed by mice of the CD1 strain.

Fibrosis is present only in the lung from old Gata1^{low} mice

Although the fibrosis in the lung from bleomycin-treated animal models is associated with an

inflammatory state induced by the chemical insult, it is debated whether inflammation has a

direct causative role in the development of IPF. Since the lung from *Gata1^{low}* and WT

littermates display similar levels of inflammation, we hypothesized that a comparison

between the level of fibrosis developed in the lungs of the two strains might clarify whether

in IPF inflammation and fibrosis are associated.

In contrast to the lack of difference in inflammation between the two strains, there was an

elevated incidence of fibrosis detected by both semiquantitative and quantitative criteria in

the lungs from *Gata1*^{low} mice while fibrosis was seldomly observed in those from their WT

littermates (**Figures 7,S6,S7A** and **Table 1**). In fact, both by reticulin and Masson's

trichrome staining, fibrosis is not detectable (66.7%) or is overall mild (Grade 1, 33.3%) in

the lungs from WT mice. By contrast, it is absent only in a minority (27.6%) of the lungs

from *Gata1*^{low} mice and is present at mild (Grade 1) and moderate/severe (Grade 2-3) levels

in 41.4% of the mice (**Table 1**) with a penetrance of 85.7% at >22 months of age. Of note,

both by Masson staining and immunofluorescent-microscopy with Col 1A and Col III

antibodies, MK were consistently found embedded within areas of collagen deposition

(**Figure S7**).

Statistical analyses with a large cohort of mice indicate that, without controlling for the

effects of age or sex, there are statistically significant differences between WT and *Gata1*^{low}

mice in the level of lung fibrosis as assessed by semiquantitative (*p*=0.01) or by quantitative

methods (*p*=0.006) (**Table 1**). In multivariable analysis, the formal test of interaction between

genotype and age was statistically significant (*p*=0.02), indicating that age is a statistically

significant modifier on the relationship between genotype and lung fibrosis. By contrast,

there is insufficient evidence to conclude there is a relationship between sex and fibrosis

grade (*p*=0.99) (**Figure 7D,E**).

These data indicate that fibrosis in the lungs is an age-associated trait in mice harboring the

Gata1^{low} genetic lesion.

Lungs from old WT and Gata1^{low} mice contain elevated levels of P-SEL, TGF-β1 and

CXCL1 which only in Gata1^{low} mice are highly expressed by megakaryocytes

Previous results in bleomycin mice have indicated that IPF is associated with greater than

normal expression of several pro-inflammatory factors including P-SEL (high in endothelial

cells); TGF-β (produced by multiple cell types including alveolar macrophages, epithelial and

endothelial cells, interstitial fibroblasts and alveolar Type II cells); and CXCL1, the murine

equivalent of human interleukin-8 (IL-8) in bronchoalveolar lavages[10,49–59]. High levels of

TGF-β are also expressed by the alveolar Type II cells of the Tamoxifen-driven conditional

knock-in *SPTFCI73T* mouse model[25]. However, the underlying inflammatory state present in

the lung of these models hampers assessing which cell type is responsible to produce the pro-

inflammatory cytokines responsible for fibrosis. We hypothesized that this assessment may

be facilitated by investigations in our model that discriminates between inflammatory

(present in both WT and *Gata1^{low}* lungs) and fibrotic (*Gata1^{low}* lungs only) events. To this

aim, we compared the content of P-SEL, TGF-β1 and CXCL1 in the lungs from *Gata1^{low}* and

WT littermates by immunohistochemistry (**Figure 8**). Strikingly, the lungs from WT mice

contained robust levels of both TGF-β1 and CXCL1 while P-SEL was barely detectable. The

levels of both cytokines were 20-fold and 2-fold greater, respectively, and P-SEL was

expressed at robust levels in the lungs from *Gata1^{low}* mice.

The relatively simple histological architecture of the lung allowed us to identify the cell types

in which these pro-inflammatory proteins are expressed (**Figure 9**). TGF-β1 and CXCL1

were expressed at comparable levels by endothelial cells, macrophages and neutrophils from

lungs of both WT and *Gata1^{low}* mice while P-SEL and TGF-β1 were clearly expressed at

levels greater than normal in the alveolar type II cells and in MK from *Gata1^{low}* mice. The

high content of P-SEL and TGF-β1 detected in *Gata1^{low}* MK by immune-histochemistry is

consistent with their high levels (*TGF-β1*: fold-change 1.49, *p*=0.006) and *P-sel*: fold change

1.63, *p*=0.031) mRNA detected by RNAseq of *Gata1^{low}* MK (**Table 2**).

These results suggest that high expression of P-SEL, TGF-β1 and CXCL1 by endothelial

cells, bronchial epithelium and macrophages are drivers for the inflammation observed in the

lung of both WT and *Gata1^{low}* littermates, but that their expression by alveolar Type II cells

and MK is responsible for fibrosis which is developed only by the lungs of the mutant mice.

Fibrosis in the lung from Gata1^{low} mice is prevented by genetic ablation of P-selectin

To assess the role of P-SEL in the pathogenesis of lung fibrosis, we compared the MK

profiling and the histopathology of lungs from *Gata1*^{low} mice with and without the *P-sel* gene

with age (**Figures 3A,7C-E, 8, 10, S3A, S8** and **Tables 1, S2**).

In the lung of mice carrying only the *P-sel*^{null} mutation, used as control, the total numbers of

MK were similar to those observed in WT littermates (**Table S2**) although cells with a large

morphology were more frequent than normal (**Figures 3A, S3A** and **Table S2).** Moreover,

the lungs from *P-sel*^{null} mice manifest levels of inflammation not significantly different from

that expressed by WT mice but do not express fibrosis (**Figure 10, 7C-E** and **Table S2**).

These results indicate that deletion of *P-sel* has modest effects on the baseline inflammatory

phenotype of CD1 mice.

The lungs from *Gata1*^{low} and *P-sel*^{null}*Gata1*^{low} mice contain similar numbers of MK (**Figure**

3A, and **Table 1**). There was a trend toward higher levels of severe (grade 2/3) inflammation

in the *Gata1*^{low} lung compared to *P-sel*^{null}*Gata1*^{low} mice (43.2% vs 27.8%) which is not

statistically significant (unadjusted p=0.27; age-adjusted p=0.08; sex-adjusted p=0.34)

(**Figure 10, Table 1**). However, lung fibrosis is higher among *Gata1*^{low} compared to *P-*

sel^{null}*Gata1*^{low} mice (Mean [±SD]=12.8 [±11.56] vs 5.2 [±5.3], unadjusted p=0.01). After

adjusting for differences in sex or in age (age-adjusted p=0.04; sex-adjusted p=0.002), the

difference in lung fibrosis between the two group is statistically significant (**Figure 7C-E,**

Figure 10, Table 1).

Mechanistically, reduction of fibrosis in mice lacking *P-sel* is associated with barely

detectable levels not only, as expected due to the genetic lesion, of P-SEL, but also of TGF-β

while the levels of CXCL1 remained as high as in WT mice (**Figure 8**). As expected, single

cell analyses indicate that genetic deletion of *P-sel* reduced P-SEL content in endothelial

cells, alveolar Type II cells, bronchial epithelium, macrophages but did not affect the content

of TGF-β and CXCL1 in these cells with respect to WT (compare **Figure S8 with Figure 9**).

By contrast, deletion of *P-sel* in *Gata1*^{low} mice specifically reduced the content of TGF-β in

the MK with respect to that observed in *Gata1*^{low} mice (compare **Figure S8** with **Figure 9**),

providing support for the hypothesis that MK are the driver for the increased

microenvironmental bioavailability associated with lung fibrosis in *Gata1*^{low} mutants.

In conclusion, ablation of *P-sel* has modest effects on inflammation but, after adjusting for

age and sex, significantly reduces fibrosis in the lungs of *Gata1*^{low} mice. Reduction of fibrosis

is associated with reduced frequency of morphologically immature MK and of MK producing
TGF-β1.

Inhibitors of P-selectin, TGF-β or CXCL1 rescues the fibrosis in the lung from Gata1^{low}

Inhibitors of P-selectin, TGF-β or CXCL1 rescues the fibrosis in the lung from Gata1^{low}
mice

To assess the pathobiological role of pro-inflammatory factors in the development of lung
fibrosis in *Gata1*^{low} mice, we tested whether it could be reversed by treating 12-months old
Gata1^{low} mice with the P-SEL neutralizing antibody RB40.34, or the TGF-β1 trap AVID200
or the CXCR1/2, the receptors for CXCL1, inhibitor Reparixin (**Figure 11**).

The P-SEL neutralizing antibody significantly reduced both inflammation and fibrosis in the
lungs. By day 40, 50% of the treated mice had no sign of lung inflammation, and fibrosis
persisted at low level by reticulin staining and was undetectable by Masson staining (**Figure
11A-C**).

The TGF-β1 trap had modest effects on lung inflammation and did not reduce fibrosis by
reticulin staining although it significantly decreased the collagen fibers detected by Masson
by day 78 (**Figure 11D-F**).

CXCR1/2 inhibition transiently reduced lung inflammation by day 20 and significantly
reduced fibrosis by reticulin staining both at day 20 and 37 and by Masson staining at day 20
(**Figure 11G-I**).

In conclusion, the P-SEL, TGF-β1 or CXCL1 inhibitors as single agents are all effective in
reverting fibrosis in the lungs from *Gata1*^{low} mice although a dissociation with the effects on
inflammation and a temporal gradient of efficacy was observed among the three drugs with
inhibition of P-SEL being the most effective.

Mechanistic insights on the effects exerted by the three pro-inflammatory inhibitors on the
fibrosis of the lung were obtained by investigating their effects on the TGF-β1, CXCL1 and
GATA1 content of the MK (**Figures 12,13** and **S9,S10**). Treatment with the P-SEL inhibitor
significantly reduced both the TGF-β1 and CXCL1 content by day 12 and the effects
persisted up to day 40 (**Figure 12**). At the single cell level, treatment with the P-SEL
inhibitor reduced the TGF-β and CXCL1 content in the MK (**Figure S9**). Treatment with this
inhibitor also increased the GATA1 content of the MK (**Figure 13**). By contrast, treatment
with either the TGF-β1 or the CXCR1/2 inhibitor significantly reduced only the CXCL1
content by the end of the treatment without affecting that of TGF-β1 (**Figure 12**). These

reductions also were more pronounced at the level of the MK (**Figure S9**). The TGF-β1 and

the CXCR1/2 inhibitor did not significantly affect the GATA1 content of MK which

remained low (**Figure 13, S10**).

These results provide further support to the hypothesis that in *Gata1^{low}* mice IPF is sustained

by pro-inflammatory proteins, such as TGF-β1 and CXCL1, produced by the abnormal MK.

Discussion

Studies on IPF are challenged by the fact that most of the anatomo-pathology biorepositories

contain lung specimens from a very limited number of IPF patients and from virtually no

healthy individuals. We succeeded obtaining lung sections from three IPF patients and

overcame the unavailability of healthy controls by analyzing sections from the lung of three

patients with lung cancer which did not include evident tumor tissue. Using these specimens,

we determined that although the frequency of MK was similar in the two groups, there were

important differences in terms of nuclear GATA1 content (barely detectable in IPF),

localization (within blood vessels in No-IPF and in the parenchyma near the alveolar

epithelium in IPF) and subpopulations (enriched for immune-MK in IPF) between the MK in

the two groups. These results suggest that immune-MK corrupted by low GATA1 content

exert a pathological role in the etiology of IPF.

We then validated these observations by determining whether the lung from mice harboring

the *Gata1^{low}* mutation which impairs expression of this gene only in cells of the

hematopoietic lineage and more specifically in MK, contain MK biased toward the immune

lineage and develop an IPF-like phenotype with age. Of note Gata1 is not expressed in the

non-hematopoietic cells within the lung. We describe that the lungs from old *Gata1^{low}* mice

contain MK expressing low levels of GATA1, an immature morphology (paucity of Plt

territories and low ploidy), biased toward the immune lineage and localized mostly in the

parenchyma near the alveoli. Furthermore, *Gata1^{low}* mice develop with age fibrosis in the

lung. These results support the hypothesis suggested by the human studies described above

that abnormalities in the immune-MK populations sustained by low GATA1 content

represent a driver for IPF.

Overall, the histological features of the fibrosis observed in the lung from *Gata1^{low}* mice are

more similar to that observed in IPF in patients than that displayed by the currently available

bleomycin model of the disease: i) as IPF patients, *Gata1^{low}* mice progressively develop both

reticulin and collagen fibrosis in the lungs as they age, recapitulating the chronic nature of the

disease, while in the bleomycin model fibrosis is an acute event retained until the insult

persist; ii) although the lung from *Gata1^{low}* mice never acquires the honeycomb morphology

typical of those from IPF patients, in the *Gata1*^{low} lung, as in those from the patients, fibrosis

is present predominantly in the subpleural region of the basal lobes and, occasionally, in the

para-septal parenchyma while in bleomycin-treated animals, fibrosis has a bronco centric

localization that reflects the site where the insult is applied; iii) as in some IPF patients,

inflammation is not the primary cause of fibrosis in the lung from *Gata1^{low}* mice while

fibrosis is a direct consequence of inflammation in bleomycin-treated animals. Our

conclusion that inflammation is not the primary driver of fibrosis in *Gata1^{low}* mice is based

on robust considerations: i) inflammation is observed also in the lung from WT and *P-sel^{null}*

mice which never develop fibrosis; ii) genetic deletion of *P-sel* prevents development of

fibrosis in the lung of mice carrying the *Gata1^{low}* mutation but does not reduce the base-line

inflammation of this organ; iii) pharmacological inhibition of either P-SEL, TGF-β or

CXCR1/2, has modest, if any, effect on inflammation but rescues the fibrosis in the lung of

Gata1^{low} mice. We suggest that inflammation is a confounding factor rather than a cause of

fibrosis in IPF and that *Gata1^{low}* mice represent a model for IPF more stringent than the

bleomycin-treated mice currently used to study the etiology of this disease. Additional studies

comparing the megakaryocyte subpopulations present in the lungs from IPF patients and from

patients with inflammation but not fibrosis, such as idiopathic non-specific interstitial

pneumonia, chronic hypersensitivity pneumonitis or connective tissue disease related

idiopathic lung disease, are necessary to clarify the MK abnormalities that specifically drive

IPF in patients, This study will be performed in the near future under the umbrella of PUMA,

a retrospective observational protocol approved by the Human Subject Committee of Campus

Biomedico and Policlinico Gemelli on March 21^{st}, 2023.

It is very interesting that the anatomical localization, and other features, of the fibrosis in the

lung of *Gata1^{low}* mice are very similar to that of the recently described Tamoxifen-treated *SP-*

CI73T mouse model[25]. Furthermore, in the lung of *Gata1^{low}* mice the corrupted MK are found

closely associated with the alveolar epithelium and are surrounded by fibers of collagen 1A

and III. These observations induce us to speculate that *Gata1^{low}* MK sustain fibrosis by

secreting factors that alter the metabolism of the surfactant proteins expressed by the alveolar

type II cells turning them into pro-fibrotic cells. Additional support for this hypothesis is that

the content of proinflammatory cytokines of the alveolar Type II cells in *Gata1^{low}* lung, as

those of the corresponding cells from Tamoxifen-treated *SP-CI73T* mouse model[25], include

high levels of TGF-β.

In the bleomycin-animal model, inflammation is initially associated with accumulation of

neutrophils within the lung, which are subsequently replaced by lymphocytes[60]. This

accumulation is associated with increases in the levels of systemic and microenvironmental

pro-inflammatory cytokines (TGF-β, CXCL1 and others), which precede the appearance of

the pro-fibrotic markers (fibronectin and pro-collagen 1) and of fibrosis[5]. Abnormal

expression of P-SEL by endothelial cells[10] and of TGF-β by numerous cell types, including

endothelial cells, epithelial cells, alveolar type II cells and macrophages[53,54], has been

implicated in the development of fibrosis in the lungs of bleomycin-treated animals. High

levels of TGF-β produced by the alveolar Type II cells were also associated with the

development of fibrosis in the Tamoxifen-treated *SP-CI73T* mouse model[25]. By contrast,

although CXCL1 is elevated in bronco-alveolar lavages of these mice[61], the cell population

responsible for its production in the lung of mouse models has yet to be identified. Most of

the findings in mice have been validated by studies involving IPF patients[55,57,58,62 50–52]. An

important exception is P-SEL that in the patients is expressed at high levels not by the

endothelial cells but by Plt, cells which originate from the MK[49]. The debate on which cell(s)

is responsible to establish the proinflammatory molecules milieu responsible for fibrosis in

the lung is expected since the studies in all the models available up to now, including the

Tamoxifen-treated *SP-CI73T* mice, are confounded by inflammation. The fact that the strain

hosting the *Gata1^{low}* mutation is well known to express a baseline inflammatory state with

age, allowed us to use a combination of genetic and pharmacological approaches to dissect

the identity of the cells specifically responsible to sustain fibrosis in the lung of our model.

By immune-histochemistry, endothelial and epithelial cells, alveolar macrophages and

interstitial fibroblasts from the lung of WT and *P-sel^{null}* mice, which express inflammation

but no fibrosis, contain robust levels of P-SEL, TGF-β and/or CXCL1, suggesting that the

pro-inflammatory profile of these cells is responsible for the baseline inflammation of the

CD1 strain. By contrast, MK, and to some extent alveolar type II cells, are the only cells that

contain robust levels of these pro-inflammatory mediators specifically in the lung from

Gata1^{low} mice, suggesting that they are responsible to establish the pro-inflammatory milieu

that drives fibrosis in our model. Support for the pro-inflammatory role of the MK is

provided by their RNA-seq profiling indicating that those of the lung from *Gata1^{low}* mice

express upregulated levels of TGF-β (fold-change=1.49, *p*=0.006) and CXCR2, the receptor

for CXCL1[63,64] (fold change=1.95, p=0.007) while the levels of CXCL1 were downregulated

(fold-change=-2.843, P=0.000). The discrepancy between the reduced levels of *CXCL1*

mRNA detected in *Gata1*[low] MK by RNA-seq and their high *CXCL1* content detected by

immunohistochemistry may be explained by the high levels of mRNA for *CXCR2*, a receptor

which internalize its ligand without routing it to the degradation system, expressed by these

cells which allow for greater accumulation of the protein in the cytoplasm. The RNAseq

profiling also displays downregulation of other factors involved in the MK-specific anti-

microbial response in this organ: *CXCL1* fold-change=2.8, *p*=0.000), *CXCL5* (fold-change -

2.691, *p*=0.017, and *CXCL10* (fold-change -1.88, *p*=0.033)[63–66], confirming that the GATA1-

deficient immune MK detected in the lung from old *Gata1*[low] mice are corrupted.

The pathogenetic role of P-SEL, TGF-β and CXCL1 in the fibrosis of the lung from *Gata1*[low]

mice was further dissected by evaluating the effects of the deletion of *P-sel* and

pharmacological inhibition of P-SEL, TGF-β and CXCR1/2 on the lung phenotype of these

mice. Genetic deletion of *P-sel* prevented mice carrying the *Gata1*[low] mutation to develop

fibrosis in the lung with age. This effect was associated with reduced content of both TGF-β

and CXCL1 in the MK from this organ. Furthermore, inhibition of P-SEL reduced fibrosis

(both as reticulin and collagen fibers), the frequency of MK with low GATA1 content and the

content of TGF-β and CXCL1 in the lung for up to day 40 of treatment. By contrast, TGF-β

inhibition reduced collagen fibers and CXCL1 content only by the end of the treatment (day

72) while that of CXCR1/2 reduced collagen fibers transiently (day 20 only), reticulin fibers

both at day 20 and 37 and CXCL1 content only at day 37. The efficacy of the CXCR1/2

inhibitor to consistently reduce only reticulin fibers is consistent with the reported efficacy of

this drug in Bleomycin-treated mice that are a model for early stages of the disease[67,68]. Of

note, both the TGF-β and CXCR1/2 inhibitors did not affect neither the TGF-β nor the

GATA1 content of the MK, which remained, respectively high and low, in the lung of the

treated *Gata1*[low] mice. Although the effects exerted by the potent and specific TGF-β1 trap on

the fibrosis of the lung from *Gata1*[low] mice are inferior to those reported for additional drugs

which target the same pathway tested in bleomycin-treated models, they are consistent with

the poor performance of those clinically tested up to now in IPF[14–17].

Overall, these data suggest that in *Gata1*[low] mice fibrosis in the lung is triggered by an

abnormal population of immune-MK which initiates a sequence of pathological effects

exerted by multiple pro-inflammatory proteins over time. It may be speculated that the first

step is accumulation of immune-MK that became engaged into a process of pathological

emperipolesis mediated by high levels of P-SEL[28]. This pathological emperipolesis leads to

death of the MK by para-apoptosis and release of the content of their cytoplasm into the

microenvironment, increasing the bioavailability of TGF-β and CXCL1, and possibly of other

proinflammatory cytokines not identified as yet, in the microenvironment. The release of

these pro-inflammatory cytokines sustains a mesenchymal cell transition responsible for

fibrosis. Since the *Gata1*^low MK from the lung, although embedded in collagen fibers, do not

contain these proteins in their cytoplasm, we exclude fibrosis is mediated by niche-poised

MK, a population responsible to synthesize the extracellular matrix during organo-genesis[45]

and that, when pathologically reactivated in adulthood, promote scarring of the BM[28]. We

suggest that, as in Tamoxifen-treated *SP-CI73T* mice, the cells undergoing the mesenchymal

transition responsible for fibrosis in the *Gata1*^low lung are the alveolar Type II cells which

became corrupted by the nearby MK.

In addition to the poor fidelity of the No-IPF controls used in the human experiments

discussed above, our study has other limitations: i) the mutation status of the IPF patients

analyzed is not known. However, given the low frequency of the telomerase-related or

surfactant proteins mutations found in IPF patients, it is unlikely that our patients harbored

any of the mutations already known; ii) the *Gata1*^*low* mutation reduces the content of the

protein in all the megakaryocyte subpopulations and not only in those with immune

functions. Unfortunately, conditional knock-out mice with reduced/ablated GATA1

expression only in the immune MK are not available. The available conditional *PF4 Cre*

strain[69] is not suited for our studies because in this strain GATA1 expression is completely

ablated in all the megakaryocyte sub-populations, making impossible to identify

abnormalities induced by reduced GATA1 content in specific MK subpopulations. Since,

compared to the other subpopulations, the development of immune-MK is specifically

sensitive to the expression of *Pu1*, double mutants in which the *Gata1*^*low* mutation is

combined with conditional *PF4 Cre-Pu1* knock-out mice may be more suited to rigorously

assess the role of immune MK in the development of IPF. Unfortunately, these mutants are

not yet available; iii) our assumption that the corrupted *Gata1*^*low* MK and the altered alveolar

Type II cells are casually linked is based only on the anatomical localization of the two cell

types. Further studies, which will rigorously assess whether the *Gata1*^*low* MK alter the

functions of alveolar Type II cells by determining their expression profiling using specific

lineage-tracking genetic mouse models, which again are not currently available, are necessary

to formally assess whether MK and alveolar type II cells are related in the development of

IPF of our model.

In spite of these limitations, we believe that our data are important because they open new

perspectives for the etiology and cure of a disease the genetic lesion of which is currently

unknown for >95% of the patients and that has limited therapeutic options. In this regard,

cells of BM origin had been already associated with development of IPF[70], leading to the

interesting hypothesis that, at least in some patients, IPF may have a hematopoietic origin that

may be treated with BM transplantation[71]. However, the consideration that IPF patients are

old and fragile and unlikely to survive the fully ablating conditioning regimes available in

2004 when these studies were performed has left this hypothesis unexplored. The more

patient friendly regimens under development (i.e. hematopoietic stem cell depletion of the

host by treatment with anti-cKIT antibodies)[72-74] may make now this approach feasible even

in IPF patients. The fact that reduced conditioning in transplanted patients with

myelofibrosis, a population as fragile as that affected by IPF, gave satisfactory results

supports the feasibility of transplants with this regiment in older patients[75]. This hypothesis

deserved to be tested by proof-of-principle experiments in animal models.

In conclusion, *Gata1^{low}* mice are a novel genetically driven IPF animal model and provide

strong evidence that immune-MK alterations play a major role in the development of this

disease. The *Gata1*^{low} model also clarifies that P-SEL dysregulation predates TGF-β and

CXCL1 in the pathobiology of IPF and should be considered for treatment of these patients.

Materials and Methods

Human Subjects. Consecutive lung sections from IPF and non-IPF patients (three different

patients per group) were provided, respectively, by the Institute of Human Pathology,

Policlinico Sant'Orsola, Bologna, Italy and by Policlinico Campus Bio-Medico (Rome).

Details of the source of the subjects included in the study are provided in **Table S1**.

Mice. Male and female WT, *P☐sel*^{null}, *Gata1*^{low} and *P☐sel*^{null}*Gata1*^{low} mice were generated

in the animal facility of Istituto Superiore di Sanità which is the recipient of the NIH animal

welfare assurance number F16-00089, as previously described[40]. All the mutations are in the

CD1 background. The study was approved by the Italian Ministry of Health (protocol n.

419/2015-PR and D9997.121) and conducted according to the European Directive

2010/63/EU for the protection of animals used for scientific purposes.

Treatments. <u>*P-SEL inhibitors:*</u> Twenty 8-11 months-old *Gata1*[low] mice were randomly
divided in two groups which were treated either with vehicle (2% v/v dimethyl sulfoxide in
H₂O) or with biotinylated mAb RB40.34 (BD Pharmigen, San Diego, CA; 30 μg/mouse
every other day by tail vein injection), as described[76,77]. Mice were sacrificed after 12-40 days
for histopathologic determinations. <u>*TGF-β1 inhibitors*</u>: *Gata1*[low] mice (10 - 12 months-old)
were weighted and randomly divided into two groups (four males and four females/group)
and treated with either vehicle (PBS) or with the TGF-β1 trap AVID200 (5mg/Kg twice a
week by intraperitoneal injection), as described[78]. Mice were treated either for 42 (six mice,
three females and three males with AVID200 and 2 females with vehicle) or 72-78 (eight
mice, four females and four males in each group) days. At day 42, 72 or 78 from the
beginning of the treatment, mice were euthanized, and their lung removed for histopathology
observations. <u>*CXCR1/CXCR2 inhibitors*</u>: Sixteen eight-month-old *Gata1*[low] mice were
anesthetized with 2-to-3% isoflurane and implanted subcutaneously with an ALZET®
Osmotic Pump (model 2002) pre-filled with 200μL of vehicle (sterile saline) or the
CXCR1/CXCR2 inhibitor Reparixin (7.5mg/h/Kg in sterile saline) as described[79,80]. Mice
were sacrificed 20 or 37 days after the beginning of the treatment, and their lungs removed
for histopathological determinations.

Histochemistry. Lung and BM from WT, *Gata1*[low], *P☐sel*[null] and *P☐sel*[null]*Gata1*[low] mice
harvested from mice that had previously received intratracheal inoculation of 10% (vol/vol)
phosphate☐buffered formalin and then fixed in 10% (vol/vol) phosphate☐buffered formalin,
paraffin embedded and cut into consecutive 3μm sections, according to standard procedures.
Consecutive sections from the mouse organs, as well as from BM and lungs of the patients,
were stained with Hematoxylin☐Eosin (H&E, cat no. 01HEMH2500 and 01EOY101000,
respectively; Histo-Line Laboratories, Milan, Italy), Reticulin (Bio-Optica, Milan, Italy) or
Masson's trichrome (Bio-Optica) and evaluated with a Nikon Eclipse E600 microscope
(Nikon Instruments Europe BV, Amsterdam, NL) equipped with the Imaging Source "33"
Series USB 3.0 Camera (DFK 33UX264; Bremen, DE). Inflammation was assessed on de-
identified H&E stained sections by an experienced veterinarian pathologist and expressed on
the basis of a semiquantitative scale based on the numbers of inflammatory cells (neutrophils,
macrophages and lymphocytes) that were infiltrating the alveolar septae and/or peribronchial-
vascular spaces: grade 0 (absence of inflammatory cells); grade 1 (rare inflammatory cells in
<10% of the area); grade 2 (moderate numbers of inflammatory cells in 10-30% of the area)
and grade 3 (presence of large numbers inflammatory cells in >30% of the area). <u>Fibrosis</u>.

Semiquantitative and quantitative assessment of fibrosis were performed on sections stained

with Reticulin or Masson's trichrome staining by an experienced veterinary pathologist and

by Image analysis using the software ImageJ (version 1.52t, National Institutes of Health,

Bethesda, MD, USA), respectively. The semiquantitative scoring was the following: grade 0

(presence of collagen fibers only at the bronchovascular level and absence of septal alveolar

fibrosis); grade 1 (presence of focal mature collagen fibers in the alveolar-septal interstitium);

grade 2 (deposition of multifocal to coalescing collagen fibers in the alveolar-septa

interstitium); grade 3 (deposition of collagen bundles that irregularly thicken the alveolar

septa and alterations of the overall pulmonary architecture) (**Figure S6A**). Quantitative image

analysis was performed on five different photomicrographs randomly selected per each lung

section/mouse (area of each photomicrograph=1.49mm^2) (**Figure S6B**). The number of mice

analyzed per experimental group is indicated in the legends. Fibrosis was expressed as

percent of the area stained in blue with respect to the total area of the field censored of the

empty alveolar spaces. Statistical analyses confirmed a good correlation between the

semiquantitative and quantitative determinations of fibrosis (**Figure S6C**).

Immunofluorescence (IF) studies. Mouse and human lung and sections (3μm thick each)

were dewaxed in xylene and treated with EDTA (buffer pH=8), incubated for 20min in a

pressure cooker (110-120°C, high pressure) for antigen retrieval and then with antibodies

against CD42b (cat no. ab183345, Abcam, Cambridge, UK) or/and GATA1 (cat no. sc-265,

Santa Cruz Biotechnology, Santa Cruz, CA, USA), Collagen I (cat no. NB600-408, Novus

biologicals, Centennial, CO, USA), Collagen III (cat no. ab7778, Abcam) or Collagen IV (cat

no. ab6586, Abcam) over night at 4°C. Cell binding of the primary antibodies was visualized

with a secondary antibody goat anti-rat Alexa Fluor 488 (cat no. ab150165, Abcam) or anti-

rabbit Alexa Fluor 555 (cat no. ab150078, Abcam), as appropriate. Sections were

counterstain with DAPI (cat no. D9542-5MG, Sigma Aldrich, Saint Louis, MI, USA) and

mounted with Fluor-shield histology mounting medium (cat no. F6182-10MG, Sigma

Aldrich). The slides were examined using a Nikon Eclipse Ni microscope equipped with the

appropriate filter cubes to distinguish the fluorochromes employed. The images were

recorded with a Nikon DS-Qi1Nc digital camera and NIS Elements software BR 4.20.01 and

selected sections were acquired using FV1000 confocal microscope (Olympus, Tokyo, Japan)

equipped with a confocal spectral imaging system (Olympus Fluoview 1000). The number of

MK was counted in 10 fields per section and expressed as mean value.

Confocal Microscopy. Femurs were fixed in formaldehyde (10% v/v with neutral buffer),
treated for 1h with a decalcifying kit (Osteodec; Bio-Optica, Milan, Italy) and included in
paraffin. Three micron-thick BM sections were dewaxed in xylene and antigens were
retrieved by treatment with EDTA buffer (pH=8) for 20' in a pressure cooker (110-120°C,
high pressure). For each sample almost 3 consecutive sections were obtained. One was
incubated with antibodies against CD41 (rat anti mouse, Santacruz Biotechnology INC.
(MWReg30): sc-19963) and LSP1 (EPR5997), another one with antibodies against CD61 (B-
7, mouse anti mouse, Santacruz Biotechnology INC., sc-46655) conjugated with Alexa 488,
and MYLK (polyclonal rabbit anti-mouse, Proteintech, 24309-1-AP); and the last one
incubated with antibodies against CD42b (mouse monoclonal, Santacruz Biotechnology INC,
sc-271171) and ARNTL (rabbit anti mouse/human NB100-2288, Bio-Techne), for 2 hours at
room temperature. After extensive washing, sections were incubated for 45□min at room
temperature with the following secondary antibodies: CD41 was visualized with the antibody
goat anti rat Alexa Fluor 488 (Catalog #A-11006, Invitrogen); the LSP1, CD42b and MYLK
antibodies were visualized with the antibody donkey anti-rabbit Alexa Fluor 555 (Catalog #
A-31572, Invitrogen) and ARNTL was visualized with the antibody goat anti mouse Alexa
Fluor 488 (Catalog #A32723, Invitrogen), CD61 is directly conjugated with Alexa 488. All
sections were mounted with Mounting Medium With DAPI - Aqueous, Fluoroshield (Catalog
#ab104139, Abcam) and examined using a confocal microscope Zeiss LSM 900 (Carl Zeiss
GmbH, Jena, Germany) in Airyscan mode, using a (Zeiss) planapo objective 60x oil A.N.
1,42. Excitation light was obtained by a Laser Dapi 408 nm (DAPI), an Argon Ion Laser (488
nm) for Alexa 488 and a Diode Laser HeNe (561 nm) for Alexa 555. DAPI emission was
recorded from 415 to 485 nm, Alexa 488 emission was recorded from 495 to 550 nm, Alexa
555 emission was recorded from 583 to 628 nm. Images recorded have an optical thickness of
0.3 μm and were processed with the Zen Blue (3.2) (Carl Zeiss GmbH, Jena,Germany) and
ImageJ (1.53t – National Institutes of Health, USA, http://imagej.nih.gov/ij) softwares. All
images were collected with the same detector value for DAPI, Alexa 488 and Alexa 555,
respectively. The background of each signal was calculated splitting the channels and the
single channel background obtained in areas external to cells. The number of positive cells
were obtained in the whole section(s).

Immunohistochemistry (IHC). Three micron-thick lung sections were dewaxed and
rehydrated. Endogenous peroxidase was blocked by immersion in H_2O_2 in methanol (3%,
v/v) for 30' at room temperature. Antigen retrieval was performed by incubation in citrate

buffer, pH=6.0 for 1min in a microwave oven at 750W, followed by cooling at room

temperature for 20'. After blocking of non-specific antigenic sites by incubation with goat

serum in PBS (10%, v/V) for 30' at room temperature, slides were incubated with CD42b

(cat no. ab183345, Abcam, Cambridge, UK), biotinylated anti-P-SEL (cat no. 553743, BD

Pharmigen), anti-TGF-β1 (cat no. sc-130348, Santa Cruz Biotechnology), anti-CXCL1 (cat

no. ab86436, Abcam), anti-CXCR1 (cat no. GTX100389, Genetex, Irvine, CA, USA) and

anti-CXCR2 (cat no. ab14935, Abcam) antibodies overnight at 4°C. Binding sites were

revealed with the commercial avidin-biotin-peroxidase kit (ABC Kit Elite, Vector,

Burlingame, CA) and the chromogen 3,3'-diaminobenzidine (0.05%, Histo-Line

Laboratories) either directly or previous incubation with suited secondary biotinylated

antibodies. Slides were counterstained with Harris hematoxylin (Histo-Line Laboratories) and

mounted with HistoMount DPX medium (cat no. 01BM500H, Histo-Line Laboratories).

Image analysis was performed using ImageJ program and the content of antibody binding

was calculated as intensity of the brown staining above the threshold in areas ($1.49mm^2$ each)

from 5 different photomicrographs randomly selected for each sample. The intensity of the

staining of individual cells recognized according to standard morphological criteria was

arbitrarily assessed as "–" (below detection), "+" (detectable) and "++" (strongly detectable).

Flow cytometry of lung megakaryocytes. Lungs were incubated in Dulbecco's modified

Eagle Medium (cat no. D8537, Sigma-Aldrich, Saint Lois, M0, USA), supplemented with

10% (v/v) fetal bovine serum (cat no. F7524, Signa Aldrich), penicillin/streptomycin (cat. no.

15140122), vitamins (cat no. 11120052), Glutamax (cat. no. 10566016), non-essential amino

acids (cat no. 11140050) and sodium pyruvate (cat no. 11360070) plus collagenase type II

(1mg/mL, cat no. 17101015) (all from Gibco; Grand Island, NY, USA), as described[46]. After

30min at 37°C, lungs were mashed through a 100-µm cell strainer (BD Pharmingen) and

washed with cold media and centrifuged at 1200 rpm for 5 minutes. ACK Lysing Buffer (cat

no. A1049201, Gibco) was added to the pellet for 5 minutes in ice. The ACK was washed out

with 50 mL of PBS and centrifuged at 1200 rpm for 5min. Cells were then incubated with

phycoerythrin (PE)-CD41 (cat no. 558040) and fluorescein isothiocyanate (FITC)-CD61 (cat.

no. 553346). Dead cells were excluded by Sytox Blue staining (0.002mM, Molecular Probes,

Eugene, OR, USA). For DNA content determinations, cells were then incubated with

phycoerythrin (PE)-CD42b (cat. no. 555473) and FITC-CD61, washed and fixed with 4%

paraformaldehyde (Sigma-Aldrich) for 60min, permeabilized with Triton-X100 (0.2% v/v in

PBS, cat. no. 108643; Merck, Darmstadt, Germany) for 15min and incubated with propidium

iodide (40 µg/mL in PBS) and RNase A (100units/mL, cat. no. 10109142001, Merck) for

30min at 37□. All the antibodies are from BD Pharmingen. Cells were analyzed with Gallios

(Beckman Coulter, Brea, CA, USA) and the flow cytometry data processed with the Kaluza

program (version 2.1, Beckman Coulter).

RNA seq analysis. MK were purified from the BM and lungs of two *Gata1^{low}* and three WT

littermates (11-25 months of age. Total RNA was collected using the Direct-zol^{TM} RNA

MiniPrep (>200,000 sorted MK) or MicroPrep (<200,000 sorted MK) (Zymo Reaserch).

Illumina libraries were constructed using SMARTer Stranded Total RNA-Seq Kit v3 - Pico

Input Mammalian (TaKaRa, Japan). RNA-seq libraries were sequenced at read lengths 2

x 151bp on the Illumina Novaseq 6000 (Illumina). Aligment was performed using RNA-

STAR (2.3.1) using the ENCODE 4 default transcriptome annotation (GENCODE v25).

Gene counts were obtained using featureCounts and FPKM per gene were calculated using

Cufflinks. Differentially expressed genes (DEGs) between the two groups of lungs MK from

Gata1^{low} mice and WT mice were analyzed with R package DESeq2 and the GO Enrichment

Analysis was performed with R package enrichplot, DOSE[81] and clusterProfiler [82,83].

Multiple testing correction was performed using the Benjamini-Hochberg method and

significance was established at an FDR of 5%.

Gene Set Enrichment Analysis. GSEA was performed using GSEA software GSEA 4.3.1

wiTH default parameters. The CURATED gene set (m2.all.v2022.1.Mm.symbols.gmt) used

for   the   analysis   was   downloaded   from   MSigDB

(https://www.broadinstitute.org/gsea/msigdb/collections.jsp). A total of 32,527 expressed

genes were included in the analysis. Genes were pre-ranked in order of their differential

expression between MK from *Gata1^{low}* lungs and from CD1 lungs.

Statistical methods. Categorical variables are presented as counts and percentages by

genotype; Fisher's exact tests were used to test the univariate association between genotype

and outcome. Continuous variables are presented as mean (standard deviation [SD]), median,

and range). Statistical analyses was performed by parametric t-test or one-way ANOVA for

multiple comparisons, as appropriate.   Multiple linear regression and ordinal logistic

regression were used to model relationships between genotype and outcomes, adjusted by age

or by sex. Multiple linear regression was used to test the interaction between age and

genotype on the outcome of percent fibrosis. Where interaction was found, simple linear

regression was used to assess the relationships between fibrosis outcomes and age, stratified

by genotype. Alignment between semiquantitative and quantitative levels of fibrosis was

assessed using box plots. All hypothesis testing was two-sided, with p<0.05 used as the

threshold for statistical significance. Analyses were conducted using SAS software (version

9.4; SAS Institute Inc., Cary, NC, USA), R (version 3.6.3), and Prism 8 (version 8.0.2;

GraphPad Software, La Jolla, CA).

Study approval. This is not a human study subjected to prior IRB approval according to the

Declaration of Helsinki for Studies Involving Human Subjects. All human specimens were

provided as de-identified material by public repositories.

Data availability. Data will be deposited in publicly accessible databases upon acceptance of
the manuscript.

Author contributions

769   FG, MZ, PV, MF, FA, FM and GS performed experiments and analyzed data. MM and JV

770   performed the RNA-seq analysis. CMH and ACD did the statistical analysis. AP, GS, SN,

771   and LR reviewed the data and wrote the manuscript. ARM, designed the experiments,

772   analyzed and reviewed the data and wrote the manuscript. All the authors read the

773   manuscript, concurred with its content and approved the submission.

Acknowledgments

775   This study was supported by grants from the National Cancer (P01-CA108671 ARM) and

776   Associazione Italiana Ricerca Cancro (AIRC IG23525, ARM). Profs. Antonia D'Errico, Ilaria

777   Bassi and Simone Carotti are gratefully acknowledged for providing access to the tissue

778   repositories of St. Orsola Hospital, Bologna and Campus Bio-medico, Rome, Italy.

Declaration of Interest

780   The authors declare that the research was conducted in the absence of any commercial or

781   financial relationships that could be construed as a potential conflict of interest. ARM

782   received funding for research from Dompé Pharmaceutics, Forbius, Novartis for other

783   projects.

References

(1)   Lederer DJ, Martinez FJ. Idiopathic Pulmonary Fibrosis. *NEJM* **2018**;*378*(19),1811–
1823.

(2)　Martinez FJ et al. Idiopathic Pulmonary Fibrosis. *Nat Rev Dis Primers*
2017;*3*(1),17074.

(3)　Richeldi L, Collard HR, Jones MG. Idiopathic Pulmonary Fibrosis. *The Lancet*.
2017;13,389(10082),1941-52.

(4)　Mekhael O et al. Mouse Models of Lung Fibrosis. *Methods Mol Biol* **2021**;*2299*,291–
321.

(5)　Hay J, Shahzeidi S, Laurent G. Mechanisms of Bleomycin-Induced Lung Damage.
Arch Toxicol **1991**,*65*(2),81–94.

(6)　Borzone G, Moreno R, Urrea R, Meneses M, Oyarzún M, Lisboa C. Bleomycin-
Induced Chronic Lung Damage Does Not Resemble Human Idiopathic
Pulmonary Fibrosis. *Am J Respir Crit Care Med* **2001**;*163*(7),1648–53.

(7)　Nicholson AG, Fulford LG, Colby TV, du Bois RM, Hansell D M, Wells AU. The
Relationship between Individual Histologic Features and Disease Progression in
Idiopathic Pulmonary Fibrosis. *Am J Respir Crit Care Med* **2002**;*166*(2),173–7.

(8)　Fernandez IE, Eickelberg O. The Impact of TGF-β on Lung Fibrosis. *Proc Am Thorac*
Soc **2012**;*9*(3),111–6.

(9)　Yang L et al. IL-8 Mediates Idiopathic Pulmonary Fibrosis Mesenchymal Progenitor
Cell Fibrogenicity. *Am J Physiol-Lung Cell Molec Physiol* **2018**;*314*(1), L127–L136.

(10)　Serrano-Mollar, A. P-Selectin Upregulation in Bleomycin Induced Lung Injury in
Rats: Effect of N-Acetyl-L-Cysteine. *Thorax* **2002**;*57*(7),629–34.

(11)　Huang, C-Y, Chung C-L, Hu T-H, Chen J-J, Liu P-F, Chen C-L. Recent Progress in
TGF-β Inhibitors for Cancer Therapy. *Biomedicine & Pharmacotherapy* **2021**;*134*,
111046.

(12)　Sime PJ, Xing Z, Graham FL, Csaky KG, Gauldie J. Adenovector-Mediated Gene
Transfer of Active Transforming Growth Factor-Beta1 Induces Prolonged Severe
Fibrosis in Rat Lung. *JCI* **1997**;*100*(4),768–76.

(13)　Liu, J-YR. Transforming Growth Factor-β₁ Overexpression in Tumor Necrosis Factor-
α Receptor Knockout Mice Induces Fibroproliferative Lung Disease. *Am J Respir Cell*
Mol Biol **2001**;*25*(1),3–7.

(14)   Noble PW et al. Pirfenidone in Patients with Idiopathic Pulmonary Fibrosis

(CAPACITY): Two Randomized Trials. *Lancet* **2011**;*377*(9779),1760–9.

(15)   King TEJ et al. A Phase 3 Trial of Pirfenidone in Patients with Idiopathic Pulmonary

Fibrosis. *N Engl J Med* **2014**;*370*(22),2083–92.

(16)   Richeldi L et al. Design of the INPULSIS^TM Trials: Two Phase 3 Trials of Nintedanib

in Patients with Idiopathic Pulmonary Fibrosis. *Respir Med* **2014**;*108*(7),1023–30.

(17)   Cerri S et al. Real-Life Comparison of Pirfenidone and Nintedanib in Patients with

Idiopathic Pulmonary Fibrosis: A 24-Month Assessment. *Respir Med* **2019**;*159*,

105803.

(18)   Hunninghake GM et al. Interstitial Lung Disease in Relatives of Patients with

Pulmonary Fibrosis. *Am J Respir Crit Care Med* **2020**;*201*(10),1240–8.

(19)   Arias-Salgado EG et al. Genetic Analyses of Aplastic Anemia and Idiopathic

Pulmonary Fibrosis Patients with Short Telomeres, Possible Implication of DNA-

Repair Genes. *Orphanet J Rare Dis* **2019**;*14*(1),82.

(20)   Gable DL et al. ZCCHC8, the Nuclear Exosome Targeting Component, Is Mutated in

Familial Pulmonary Fibrosis and Is Required for Telomerase RNA Maturation. *Genes*

Dev **2019**;*33*(19–20),1381–96.

(21)   Armanios M. Telomerase and Idiopathic Pulmonary Fibrosis. *Mutation*

Research/Fundamental and Molecular Mechanisms of Mutagenesis **2012**;*730*(1–

2),52–8.

(22)   Zhang K et al. Pulmonary Alveolar Stem Cell Senescence, Apoptosis, and

Differentiation by P53-Dependent and -Independent Mechanisms in Telomerase-

Deficient Mice. *Cells* **2021**;*10*(11),2892

(23)   Yasutomo, K. Genetics and Animal Models of Familial Pulmonary Fibrosis. *Int*

Immunol **2021**;*33*(12),653–7

(24)   Mulugeta S, Nureki S-I, Beers MF. Lost after Translation: Insights from Pulmonary

Surfactant for Understanding the Role of Alveolar Epithelial Dysfunction and Cellular

Quality Control in Fibrotic Lung Disease. *Am J Physiol-Lung Cell Mol Physiol*

2015;*309* (6),L507–L525.

(25) Nureki, S-I. et al. Expression of Mutant Sftpc in Murine Alveolar Epithelia Drives
Spontaneous Lung Fibrosis. *J Clin Inv* **2018**;*128*(9),4008–24.

(26) Zhang D, Newton CA. Familial Pulmonary Fibrosis. *Chest* **2021**;*160*(5),1764–73.

(27) Wernig G et al. Unifying Mechanism for Different Fibrotic Diseases. *Proc Nat Acad*
Sci **2017**;*114*(18),4757–62.

(28) Malara A, Abbonante V, Zingariello M, Migliaccio A, Balduini A. Megakaryocyte
Contribution to Bone Marrow Fibrosis: Many Arrows in the Quiver. *Mediterr J*
Hematol Infect Dis **2018**;*10*(1),e2018068–e2018068.

(29) Tefferi A. Primary Myelofibrosis: 2023 Update on Diagnosis, Risk☐stratification, and
Management. *Am J Hematol* **2023**;*98*(5),801–21.

(30) Passamonti F, Mora B. Myelofibrosis. *Blood* 2023;141(16),1954–70.

(31) Wen JQ et al. Targeting Megakaryocytic-Induced Fibrosis in Myeloproliferative
Neoplasms by AURKA Inhibition. *Nat Med* **2015**;*21*(12),1473–80.

(32) Ciurea SO. Pivotal Contributions of Megakaryocytes to the Biology of Idiopathic
Myelofibrosis. *Blood* **2007**;*110*(3),986–93.

(33) Gilles L et al. Downregulation of GATA1 Drives Impaired Hematopoiesis in Primary
Myelofibrosis. *J Clin Invest* **2017**;*127*(4),1316–20.

(34) Vannucchi AM et al. Abnormalities of GATA-1 in Megakaryocytes from Patients with
Idiopathic Myelofibrosis. *Am J Pathol* **2005**;*167*(3),849–58.

(35) Shivdasani RA, Orkin SH. The Transcriptional Control of Hematopoiesis. *Blood* **1996**;
87(10),4025–39.

(36) Shivdasani RA, Fujiwara Y, McDevitt MA, Orkin SH. A Lineage-Selective Knockout
Establishes the Critical Role of Transcription Factor GATA-1 in Megakaryocyte
Growth and Platelet Development. *EMBO Journal* **1997**;*16*(13),3965–73.

(37) Vannucchi AM et al. Development of Myelofibrosis in Mice Genetically Impaired for
GATA-1 Expression (GATA-1(Low) Mice). *Blood* **2002**;*100*(4),1123–32.

(38) Zingariello M et al. Characterization of the TGF-Beta1 Signaling Abnormalities in the
Gata1low Mouse Model of Myelofibrosis. *Blood* **2013**;*121*(17),3345–63.

(39)   Zingariello M et al. Resident Self-Tissue of Proinflammatory Cytokines Rather Than
874   Their Systemic Levels Correlates with Development of Myelofibrosis in Gata1low
875   Mice. *Biomolecules*. **2022**;12(2),234

876   (40)   Spangrude GJ et al. P-Selectin Sustains Extramedullary Hematopoiesis in the
877   Gata1low Model of Myelofibrosis. *Stem Cells* **2016**;*34*(1),67–82.

878   (41)   Lefrançais E et al. The Lung Is a Site of Platelet Biogenesis and a Reservoir for
879   Haematopoietic Progenitors. *Nature* **2017**;*544*(7648),105–9.

880   (42)   Trowbridge EA, Martin JF, Slater DN. Evidence for a Theory of Physical
881   Fragmentation of Megakaryocytes, Implying That All Platelets Are Produced in the
882   Pulmonary Circulation. *Thromb Res* **1982**;*28*(4),461–75.

883   (43)   Pariser DN et al. Lung Megakaryocytes Are Immune Modulatory Cells. *J Clin Invest*
884   **2021**;*131*(1),e137377.

885   (44)   Yeung AK, Villacorta-Martin C, Hon S, Rock JR, Murphy GJ. Lung Megakaryocytes
886   Display Distinct Transcriptional and Phenotypic Properties. *Blood Adv*
887   **2020**;*4*(24),6204–17.

888   (45)   Wang H et al. Decoding Human Megakaryocyte Development. *Cell Stem Cell* **2021**;
889   *28*(3),535-549.e8.

890   (46)   Sun S et al. Single-Cell Analysis of Ploidy and the Transcriptome Reveals Functional
891   and Spatial Divergency in Murine Megakaryopoiesis. *Blood* **2021**;*138*(14),1211–24.

892   (47)   Zhou Y et al. Megakaryocytes Participate in the Occurrence of Bleomycin-Induced
893   Pulmonary Fibrosis. *Cell Death Dis* **2019**;*10*(9),648.

894   (48)   Brayton CF, Treuting PM, Ward JM. Pathobiology of Aging Mice and GEM. *Vet*
895   *Pathol* **2012**;*49*(1),85–105.

896   (49)   Crooks MG, Fahim A, Naseem KM, Morice AH, Hart SP. Increased Platelet
897   Reactivity in Idiopathic Pulmonary Fibrosis Is Mediated by a Plasma Factor. *PLoS*
898   *One* **2014**;*9*(10),e111347.

899   (50)   Kass DJ et al. Comparative Profiling of Serum Protein Biomarkers in Rheumatoid
900   Arthritis–Associated Interstitial Lung Disease and Idiopathic Pulmonary Fibrosis.
901   *Arthritis & Rheumatology* **2020**;*72*(3),409–19.

902   (51)   Guiot J, Moermans C, Henket M, Corhay J-L, Louis R. Blood Biomarkers in
903   Idiopathic Pulmonary Fibrosis. *Lung* **2017**;*195*(3),273–80.

904   (52)   Papiris SA et al. High Levels of IL-6 and IL-8 Characterize Early-on Idiopathic
905   Pulmonary Fibrosis Acute Exacerbations. *Cytokine* **2018**;*102*,168–72.

906   (53)   Khalil N, O'Connor R. The Role of TGF-β in Bleomycin Induced Pulmonary Fibrosis.
907   In: Transforming Growth Factor-β in Cancer Therapy. **2008**;Vol 1:pp581-594 *Humana*
908   *Press*

909   (54)   Chen X et al. Inhibition of Wnt/β-Catenin Signaling Suppresses Bleomycin-Induced
910   Pulmonary Fibrosis by Attenuating the Expression of TGF-B1 and FGF-2. *Exp Mol*
911   *Pathol* **2016**;*101*(1),22–30.

912   (55)   Saito A, Horie M, Micke P, Nagase T. The Role of TGF-β Signaling in Lung Cancer
913   Associated with Idiopathic Pulmonary Fibrosis. *Int J Mol Sci* **2018**;*19*(11),3611.

914   (56)   Khalil N, Corne, S, Whitman C, Yacyshyn H. Plasmin Regulates the Activation of
915   Cell-Associated Latent TGF-Beta 1 Secreted by Rat Alveolar Macrophages after in
916   Vivo Bleomycin Injury. *Am J Respir Cell Mol Biol* **1996**;*15*(2),252–59.

917   (57)   Khalil N et al. Increased Production and Immunohistochemical Localization of
918   Transforming Growth Factor- *α* in Idiopathic Pulmonary Fibrosis. *Am J Respir Cell*
919   *Mol Biol* **1991**;*5*(2),155–62.

920   (58)   Xu YD, Hua J, Mui A, O'Connor R, Grotendorst G, Khalil N. Release of Biologically
921   Active TGF-B1 by Alveolar Epithelial Cells Results in Pulmonary Fibrosis. *Am J*
922   *Physiol-Lung Cell and Mol Physiol* **2003**;*285*(3),L527–L539.

923   (59)   Liang J et al. Single-Cell Transcriptomics Identifies Dysregulated Metabolic Programs
924   of Aging Alveolar Progenitor Cells in Lung Fibrosis. *bioRxiv*
925   **2020**;2020.07.30.227892.

926   (60)   Izbicki G, Segel MJ, Christensen TG, Conner MW, Breuer R. Time Course of
927   Bleomycin-Induced Lung Fibrosis. *Int J Exp Pathol* **2002**;*83*(3),111–9.

928   (61)   Redente EF et al. Age and Sex Dimorphisms Contribute to the Severity of Bleomycin-
929   Induced Lung Injury and Fibrosis. *Am J Physiol-Lung Cell Mol Physiol* **2011**;*301*(4),
930   L510–L518.

(62) Khalil N, O'Connor RN, Flanders KC, Unruh H. TGF-Beta 1, but Not TGF-Beta 2 or
TGF-Beta 3, Is Differentially Present in Epithelial Cells of Advanced Pulmonary
Fibrosis: An Immunohistochemical Study. *Am J Respir Cell Mol Biol* **1996**;*14*(2),131–
138.

(63) Cheng Y, Ma X, Wei Y, Wei X-W. Potential Roles and Targeted Therapy of the
CXCLs/CXCR2 Axis in Cancer and Inflammatory Diseases. *Biochim Biophys Acta*
(BBA) - Reviews on Cancer **2019**,*1871*(2),289–312.

(64) Liu Q et al. The CXCL8-CXCR1/2 Pathways in Cancer. *Cytokine Growth Factor Rev*
2016;*31*,61–71.

(65) Mei J et al. CXCL5 Regulates Chemokine Scavenging and Pulmonary Host Defense to
Bacterial Infection. *Immunity* **2010**;*33*(1),106–17.

(66) Elemam N, Talaat I, Maghazachi A. CXCL10 Chemokine: A Critical Player in RNA
and DNA Viral Infections. *Viruses* **2022**;*14*(11),2445.

(67) Cheng I-Y. et al. Particulate Matter Increases the Severity of Bleomycin-Induced
Pulmonary Fibrosis through KC-Mediated Neutrophil Chemotaxis. *Int J Mol Sci* **2019**;
21(1),227.

(68) Mattos MS et al. CXCR1 and CXCR2 Inhibition by Ladarixin Improves Neutrophil-
Dependent Airway Inflammation in Mice. *Front Immunol* **2020**;*11:566953*.

(69) Bruns I et al. Megakaryocytes regulate hematopoietic stem cell quiescence through
CXCL4 secretion. Nat Med. 2014 Nov;20(11),1315-20.

(70) Hashimoto N, Jin H, Liu T, Chensue SW, Phan SH. Bone Marrow–Derived 947
Progenitor Cells in Pulmonary Fibrosis. *JCI* 2004;113 (2),948,243–52.

(71) Dunsmore SE, Shapiro SD. The bone marrow leaves its scare: New concepts for
Pulmonary fibrosis. JCI **2004**; 113(2),180-2.

(72) Czechowicz A, Kraft D, Weissman IL, Bhattacharya D. Efficient
Transplantationvia Antibody-Based Clearance of Hematopoietic Stem Cell Niches.
Science (1979) 2007;318 (5854),1296–99.

(73) Czechowicz A et al. Selective Hematopoietic Stem Cell Ablation Using CD117-
Antibody-Drug-Conjugates Enables Safe and Effective Transplantation with Immunity
Preservation. Nat Commun 2019;10(1),617-29.

(74)   Chan Y et al. Non-Genotoxic Restoration of the Hematolymphoid System in Fanconi
962   Anemia. Transplant Cell Ther 2023;29(3),164.e1-164.e9.

963   (75)   McLornan D et al. Myeloablative and Reduced-Intensity Conditioned Allogeneic
964   Hematopoietic Stem Cell Transplantation in Myelofibrosis: A Retrospective Study by
965   the Chronic Malignancies Working Party of the European Society for Blood and
966   Marrow Transplantation. *Biol. Blood Marrow Transplant.* 2019;25(11), 2167–71.

967   (76)   Embury SH et al. The Contribution of Endothelial Cell P-Selectin to the Microvascular
968   Flow of Mouse Sickle Erythrocytes in Vivo. *Blood* **2004**;*104*(10),3378–85.

969   (77)   Verachi P et al. Preclinical Studies on the Use of a P-Selectin-Blocking Monoclonal
970   Antibody to Halt Progression of Myelofibrosis in the Gata1 Mouse Model. *Exp*
971   *Hematol* **2022**;117,43-61 .

972   (78)   Varricchio L et al. TGF-B1 Protein Trap AVID200 Beneficially Affects
973   Hematopoiesis and Bone Marrow Fibrosis in Myelofibrosis. *JCI Insight*
974   **2021**;*6*(18),e145651.

975   (79)   Dunbar A et al. CXCL8/CXCR2 Signaling Mediates Bone Marrow Fibrosis and
976   Represents a Therapeutic Target in Myelofibrosis. *Blood* **2023**;141(20),20508-19.

977   (80)   Verachi P et al. The CXCR1/CXCR2 Inhibitor Reparixin Alters the Development of
978   Myelofibrosis in the Gata1low Mice. *Front Oncol* **2022**;*12*,853484.

979   (81)   Yu G, Wang LG, Yan G-R, He Q-Y. DOSE: An R/Bioconductor Package for Disease
980   Ontology Semantic and Enrichment Analysis. *Bioinformatics* **2015**;*31*(4),608–9.

981   (82)   Yu G, Wang LG, Han Y, He Q-Y. ClusterProfiler: An R Package for Comparing
982   Biological Themes Among Gene Clusters. *OMICS* **2012**;*16*(5),284–7.

983   (83)   Wu T et al. ClusterProfiler 4.0: A Universal Enrichment Tool for Interpreting Omics
984   Data. *The Innovation* **2021**;*2*(3),100141.

986
987

Table 1: Comparison of age, sex, and histological parameters between WT vs *Gata1*[low] mice and *Gata1*[low] mice vs P-sel[null]Gata1[low]

	Gata1[low] (n=47)	WT (n=18)	Unadjusted p-value	Age-adjusted p-value	Sex-adjusted p-value	P-sel[null]Gata1[low] (n=25)	Unadjusted p-value	Age-adjusted p-value	Sex-adjusted p-value
Age (months)									
N (Missing)	47 (0)	18 (0)				25 (0)			
Mean (±SD)	12.6 (±5.44)	8.7 (±5.56)	0.01[a]			10.4 ± (4.04)	0.07[a]		
Median	12.0	10.5				11.0			
Range	2.0, 24.0	2.0, 18.0				3.0, 16.0			
Sex, n (%)			0.37[b]				0.62[b]		
Female	5 (31.3%)	5 (31.3.7%)				14 (56.0%)			
Male	11 (68.8%)	11 (68.8%)				11 (44.0%)			
N (Missing)	45 (2)	16 (8)				25 (0)			
Total MK in the lung/144mm^2									
N (Missing)	22 (25)	9 (9)				10 (15)			
Mean (±SD)	1.5 (±0.91)	3.6 (±2.42)	0.001[a]	<0.001[c]	0.004[c]	15 (±4.9)	0.80	0.<80[a]	0.14[c]
Median	1.3	3.0				1.5			
Range	1.1, 8.3	0.0, 3.8				8, 22			
Small MK/144mm^2									
N (Missing)	22 (25)	9 (9)				10 (15)			
Mean (±SD)	0.7 (±0.84)	3.4 (±2.53)	<0.001[a]	<0.001[b]	<0.001[c]	0.8 (±0.39)	0.70[a]	0.48[c]	0.01[c]
Median	0.4	3.0				0.6			
Range	0.0, 3.6	1.0, 8.3				0.2, 1.4			
Large MK/144mm^2							0.37[a]	0.43[c]	0.72[c]
N (Missing)	22 (25)	9 (9)				10 (15)			
Mean (±SD)	0.9 (±0.59)	0.3 (±0.42)	0.007[a]	0.009[c]	0.02[c]	0.7 (±0.36)	0.37[a]	0.43[c]	0.72[c]
Median	0.7	0				0.6			

Table 1: Comparison of age, sex, and histological parameters between WT vs *Gata1*[low] mice and *Gata1*[low] mice vs P-sel[null]Gata1[low]

	Gata1[low] (n=47)	WT (n=18)	Unadjusted p-value	Age-adjusted p-value	Sex-adjusted p-value	P-sel[null]Gata1[low] (n=25)	Unadjusted p-value	Age-adjusted p-value	Sex-adjusted p-value
Range	0.0, 2.3	0.0, 1.2				0.2, 1.2			
Severity of lung inflammation, n (%)			>0.99[b]	0.85[d]	0.86[d]		0.27[b]	0.08[d]	0.34[d]
0	10 (27.0%)	3 (25.0%)				9 (50.0%)			
1	11 (29.7%)	4 (33.3%)				4 (22.2%)			
2 or 3	16 (43.2%)	5 (41.7%)				5 (27.8%)			
N (Missing)	37 (10)	12 (6)				18 (7)			
Semiquantitative fibrosis, n (%)			0.01[b]	0.02[d]	0.001[d]		0.009[b]	0.01[d]	<0.001[d]
0	8 (27.6%)	8 (66.7%)				12 (66.7%)			
1	9 (31.0%)	4 (33.3%)				5 (27.8%)			
2 or 3	12 (41.4%)	0 (0.0%)				1 (5.6%)			
N (Missing)	29 (18)	12 (6)				18 (7)			
Quantitative fibrosis									
N (Missing)	29 (18)	12 (6)				18 (7)			
Mean (±SD)	12.8 (±11.56)	3.0 (±3.10)	0.006[a]	0.01[c]	0.002[c]	5.2 (±5.32)	0.01[a]	0.04[c]	0.002[c]
Median	13.2	1.9				3.1			
Range	0.0, 45.1	0.0, 9.0				0.3, 16.3			

[a]Equal variance two sample t-test p-value; [b]Fisher's exact p-value; [c]Parameter p-value from multiple linear regression with age or sex as covariate; [d]Parameter p-value from ordinal logistic regression with age or sex as covariate.

Table 2: Expression signature for genes which characterize HSC-supportive, immune- and Plt-poised MK populations in CD41pos cells purified from the bone marrow and lung of old *Gata1^low* mice with respect to the corresponding populations from WT littermates.

-Poised	ID	BM *Gata1low* VS BM Wilde Type			Lung *Gata1low* VS Lung Wilde Type		
		log2 FoldChange	pvalue	padj	log2 FoldChange	pvalue	padj
HSC	Cxcr2	4.820	0.000	0.000	1.945	0.007	0.129
HSC	Cxcr4	0.170	0.704	0.913	1.108	0.020	0.235
HSC	Gata2	-2.372	0.000	0.002	1.430	0.016	0.210
HSC	Igf1	1.002	0.305	0.692	-1.515	0.051	0.366
HSC	Itgb3	-0.077	0.845	0.962	0.297	0.463	0.843
HSC	Mylk2	0.451	0.671	0.899	0.361	0.697	0.927
HSC	Mylk4	-0.590	0.658	0.894	1.599	0.197	
HSC	Sox9	-1.096	0.237	0.636	-1.338	0.099	0.491
HSC	Tgfb1	-0.335	0.527	0.832	1.488	0.006	0.115
HSC	Tpm1	-0.687	0.156	0.532	-1.314	0.007	0.135
HSC	Tpm2	-2.037	0.001	0.028	-2.039	0.001	0.029
HSC	Wnt3a	-2.283	0.068	0.359	1.398	0.185	
IMMUNE	Ccl3	-2.660	0.000	0.001	1.686	0.007	0.127
IMMUNE	Clec4a2	3.790	0.000	0.000	2.407	0.002	0.065
IMMUNE	Clec4e	1.451	0.018	0.165	1.470	0.022	0.250
IMMUNE	Clec5a	4.401	0.000	0.000	2.109	0.002	0.061
IMMUNE	Ctsd	1.171	0.075	0.378	1.983	0.003	0.073
IMMUNE	Cxcl1	1.013	0.274	0.669	-2.843	0.000	0.012
IMMUNE	Cxcl10	-2.476	0.004	0.061	-1.881	0.033	0.301
IMMUNE	Cxcl12	-3.969	0.000	0.000	-1.189	0.058	0.395
IMMUNE	Cxcl5	-5.023	0.001		-2.691	0.017	
IMMUNE	Cybb	4.240	0.000	0.009	3.122	0.008	0.144
IMMUNE	Hck	4.154	0.000	0.000	1.278	0.095	0.482
IMMUNE	Il1a	-0.621	0.585	0.859	0.443	0.666	0.918
IMMUNE	Il1b	1.775	0.000	0.006	1.248	0.011	0.171
IMMUNE	Il6	-3.290	0.000	0.002	-2.186	0.007	0.132
IMMUNE	Itga2b	-1.794	0.007	0.085	2.329	0.001	0.032
IMMUNE	Sp1	0.424	0.165	0.546	-0.226	0.475	0.847
IMMUNE	Lcn2	5.421	0.000	0.000	-2.205	0.004	0.099
PLT	Gp1ba	-2.511	0.010	0.110	-0.340	0.726	0.935
PLT	Pf4	-1.230	0.228	0.628	2.035	0.054	0.376
PLT	Ppbp	-2.383	0.112		1.563	0.361	
PLT	Selp	0.108	0.883	0.883	1.628	0.031	0.294
PLT	Tpo	-2.694	0.112		0.416	0.777	
PLT	Vwf	-0.346	0.777	0.940	1.203	0.325	0.770

Figure 1. The megakaryocytes in the lung from patients with IPF are hypomorphic for GATA1. A) Hematoxilin&Eosin (H&E) and reticulin staining of the corresponding consecutive sections confirming that the lung from the IPF patient has an overall abnormal architecture with fibrosis. By contrast the architecture of the lung section from the lung cancer patient is overall normal, confirming that we were analyzing a region of the organ free of cancerous cells. Magnification 400x in the panels on the top and 200x in the middle and bottom panels. (B) Double IF staining with anti-GATA1 (green) and CD42b (red, as a marker of MK) of lung sections from one representative IPF patient and one representative no-IPF patient, as indicated. Nuclei are counterstained with DAPI. The insert shows a representative IPF MK at high power fields to show the absence of green signals in its nucleus. The signal captured in the individual channels in additional representative MK are presented in **Figure S1A**. The white and red dotted lines indicate vessels (containing autofluorescent red blood cells) and alveoli (lined by epithelial cells recognized by the morphology of their nuclei, IPF). C) Frequency of CD42b^{pos} MK per area (0.488mm^2) and percentage of GATA1^{pos} MK (over the total number of CD42b^{pos} cells) observed in the lungs from IPF and no-IPF patients, as indicated. Data are presented as Mean(±SD) and as value in individual subjects (each dot represents a single subject). The data are analyzed by One-way ANOVA and significant p values are indicated within the panels.

Figure 2. The lung from patients with IPF contains numerous megakaryocytes biased for the immune lineage. A) Confocal microscopy analyses for markers of immune (CD41, green/LSP1, red), HSC-supporting (CD61 green/MYLK4 red) and Plt-poised (CD42b red/ARNTL green) MK in the lung from one representative no-IPF and one IPF patients, as indicated. The signal captured in the individual channels in additional representative MKs are presented in **Figure S1B**. The white and red dotted lines indicate vessels (containing autofluorescent red blood cells) and bronchial epithelium (lined by epithelial cells recognized as cuboidal or columnar cells, IPF). To be noted that the CD41^{pos} cell in the No-IPF section is LSP1^{neg}. Original magnification 400x. B-D) Frequency of total CD41^{pos} and CD41^{pos}/LPS1^{pos} (immune-MK). D) CD61^{pos} and CD61^{pos}/MYLK^{pos} (HSC-supporting MK, E) and total CD42b^{pos} and CD42b^{pos}/ARNTL^{pos} (Plt-MK, F) MK detected in the lungs from three No-IPF and three IPF patients, as indicated. Data are presented as Mean(±SD) and as value in individual subjects (each dot represents a single subject). The data are analyzed by One-way ANOVA and significant p values are indicated within the panels.

Figure 3. The megakaryocytes in the lungs from Gata1low mice have the morphology of immature cells (reduced platelet territories and low ploidy levels) and are hypomorphic for GATA1. A) IHC with CD42b of lung sections from WT (wild-type) and *Gata1low* littermates, as well as from *P-sel^null* and *P-sel^null Gata1low* mice. The panels present representative MK with immature or mature (presence Plt-forming MK, PfMK) or absence of platelet territories. Scale bar 50μm. Magnification 1000x (See **Figure S2A** for further detail). B) Confocal microscopy with CD42b (red) and GATA1 (green) of MK in the lung from representative WT and *Gata1low* littermates. White and red dotted lines indicate vessels (containing autofluorescent red blood cells) and bronchial epithelium (lined by epithelial cells recognized by their canonical morphology). Original Magnification 400x. Frequency of CD42b^pos cells/area (0.488mm^2) and percentage of GATA1^pos CD42b^pos (over total number of CD42b^pos cells) cells observed in lungs from multiple mice are presented. C) Total cell number in lungs of old WT and *Gata1low* mice, as indicated. D) Representative flow cytometry analyses with the gate used to determine the frequency of CD41^pos/CD61^pos cells in lungs of WT and *Gata1low* mice, as indicated. The frequency observed in multiple experiments is presented on the right. E) Flow cytometry histograms with the level of propidium iodide (PI) incorporated by cells in the CD42b^neg (control) and CD42b^pos (MK) gate of lung cells from WT and *Gata1low* mice. CD42b^neg cells were analyzed to determine the staining of cells with a DNA content of 2N. The panel on the right shows the frequency of CD42b^pos cells with DNA content 2N, 4N and >8N in lungs from *Gata1low* mice and WT littermates (both 15-months old). In B) and E) data are presented as Mean (±SD) and as value in individual mice (each symbol a single mouse) and were analyzed by ANOVA. Statistically significant p values are indicated within the panels.

Figure 4. A) Confocal microscopy analyses with markers for the immune (CD41, green/LSP1, red), HSC-supporting (CD61 green/MYLK4 red) and Plt-poised (CD42b red/ARNTL green) MK in the lung from representative WT and *Gata1low* mice at 2-3 (young) and 15-months (old) of age, as indicated. MK are indicated by arrows. White dotted lines indicate the contour of blood vessels (recognized by the autofluorescent red blood cells), red dotted lines indicate the profile of the bronchial epithelium and blue dotted lines indicate alveolar spaces contoured by the alveolar epithelium. Original Magnification 400x. The signal from representative MK captured in the individual channels are presented in **Figure S4**. To be noted that Ptl-poised MK are localized within vessels both in the WT and *Gata1low* mice. B-D) Frequency of CD41pos/LPS1pos (immune-MK), CD61pos/MYLKpos (HSC-supporting MK) and CD42bpos/ARNTLpos (Plt-poised) MK detected in the lungs from multiple young and old WT and *Gata1low* mice. Data are presented as Mean(±SD) and as value in individual subjects (each dot represents a single mouse). The data are analyzed by One-way ANOVA and significant p values are indicated within the panels.

Figure 5. The megakaryocytes from Gata1low mice express a defective immune-signature. A) Principal component analysis (PCA) of the RNAseq data obtained with MK purified from the BM and lungs of individual 15-months old WT and *Gata1low* littermates (each dot an individual mouse) as indicated. B) Heatmaps for the level of expression of representative genes which characterize the signature of immune (blue), HSC-supporting (green) and Plt-poised (red) MK described by 43. Statistically significant (p<0.05) differences are indicated by red dots. The significant upregulation of CD41 in the lung of *Gata1low* with respect to WT confirms the confocal microscopy data indicating that *Gata1low* contain more immune-MK than WT mice. Although immune-MK are higher in number, the overall immune-gene signature of the *Gata1low* MK is defective.

Figure 6. Inflammation is present in lungs from wild-type and Gata1low littermates alike. (A) Stack images of the lungs from representative WT and *Gata1low* littermates (both 20 months old) stained by H&E or Masson's trichrome, as indicated. Original magnification 40x. Abbreviations: cr=cranial: cd=caudal; d=dorsal; l=lateral and m=medial. B) H&E staining of sections of the lungs from representative WT and *Gata1low* littermates expressing interstitial pneumonia at grade 0, 1, 2 and 3, as indicated. Red and green arrows indicate lymphocytes and neutrophils respectively. Magnification 200x. The frequency of mice expressing the different grades of inflammation in the two groups is indicated on the right. (see **Table 1** for further detail).

Figure 7. Fibrosis is present only in lungs from old *Gata1^low* mice. (A) Stack images of the lungs from representative WT and *Gata1^low* littermates (both 20 months old) stained by Masson's trichrome, as indicated (consecutive sections of the images presented in Figure 6). Original magnification 40x. Abbreviations: cr=cranial: cd=caudal; d=dorsal; l=lateral and m=medial. B) H&E, Reticulin and Masson's trichrome staining of consecutive sections from the lungs of representative WT and *Gata1^low* littermates at 3- and 24-months of age, as indicated. Magnification 100x. C-E) Quantification of fibrosis by reticulin staining (C) and association of fibro *Gata1^low*sis in the lung evaluated by Masson's trichrome staining with sex (D) and age (E), stratified by WT, *Gata1^low*, *P-sel^null* and *Gata1^low P-sel^null* littermates, as indicated. P-values from One-way ANOVA in C and logistic (D) and simple linear (E) regression models, stratified by genotype are overlayed, along with p-value for the interaction term from a multivariable model including sex and genotype (D) or age and genotype (E) as covariates (see **Table 1** for further detail).

Figure 8. Lungs from old *Gata1^{low}* mice contain levels greater than normal of P-SEL, TGF-β1 and CXCL1. A) Immunohistochemical staining for P-SEL, TGF-β1 and CXCL1 of lung sections from representative old WT, *Gata1^{low}*, *P-sel^{null}* and *Gata1^{low}P-sel^{null}* mice, as indicated. Magnification 200x. B) Quantification by computer assisted imaging of the P-SEL, TGF-β1 and CXCL1 content in the lungs from the four experimental groups. Data are presented as Mean(±SD) and were analyzed by Tukey's multiple comparisons test. Values observed in individual mice are presented as dots. Statistically significant p values (<0.05) among groups are indicated withing the panels.

Figure 9. While P-SEL, TGF-β1 and CXCL1 are contained at high levels in multiple cell types in the lung from WT and *Gata1^low* mice, they are contained at high levels in megakaryocytes only in that from *Gata1^low* mice. P-SEL, TGF-β1 and CXCL1 content in endothelial cells, alveolar type II cells, bronchial epithelium, macrophages, neutrophils and MK from representative WT and *Gata1^low* littermates (both 13-months old) as indicated. Cells were identified according to the following morphological criteria: Endothelial cells were recognized as flat cells delimiting the vessels lumina; alveolar type II cells were identified as cuboidal cells along the alveolar walls; bronchial epithelium was defined as pseudostratified to mono-stratified cuboidal cells along the airways of the bronchial area; macrophages were identified on the basis of their round nucleus and abundant cytoplasm; neutrophils on the basis of their small size and kidney shaped nuclei and MK on the basis of their size (10 times greater than that of any other cell type) and the polylobate morphology of their nuclei. Magnification 400x. Semi-quantitative estimates of protein content are indicated as undetectable (-), present (+) and strongly present (++).

Figure 10. Genetic ablation of P-selectin prevents the development of fibrosis in the lung from *Gata1low* mice. A) H&E, Reticulin and Masson's trichrome staining of consecutive sections of lungs from representative WT, *Gata1low*, *P-selnull* and *P-selnullGata1low* mice (all 12-15 months old), as indicated. Magnification 40x. Results observed in multiple mice per experimental group are presented and analyzed in detail in **Figure 7** and **Table 1**.

Figure 11. Treatments with inhibitors of either P-selectin, TGF-β or CXCR1/2 all prevent the development of fibrosis in the lung from old *Gata1^low* mice. A, D and G) H&E, Reticulin and Masson's trichrome staining of consecutive sections from lungs of representative *Gata1^low* mice treated either with vehicle or the P-SEL antibody (12 or 40 days, B), or the TGF-β inhibitor (42 and 78 days, E) or the CXCR1/2 inhibitor (20 or 37 days). H), as indicated. Magnification 200x. C, F and I) Number of *Gata1^low* mice (in percent of the total) expressing Grade 0, 1, 2 or 3 of inflammation when treated with either vehicle or the P-SEL (C), TGF-β (F) or CXCR1/2 (I) inhibitor. Results of the vehicle groups at different time points were pooled, because similar. D, G, J) Level of fibrosis detected by reticulin or Masson staining in consecutive lung sections from mice treated either with vehicle or with the P-SEL (D), the TGF-β (G), or the CXCR1/2 (I) inhibitor. Data are presented as Mean (±SD) and as data per individual mouse (each dot a single mouse). Statistical analyses were performed by one-way ANOVA and p values statistically significant between groups are indicated within the panels.

Figure 12. In the lung from *Gata1^low* mice, treatment with the P-SEL inhibitor reduces the content of both TGF-β1 and CXCL1 while those with the TGF-β1 or the CXCR1/2 inhibitor reduce the levels of CXCL1 only. A) Immunohistochemical staining for TGF-β1 and CXCL1 of consecutive lung sections from representative *Gata1^low* mice treated either with vehicle or with the P-SEL antibody (12 or 40 days), or the TGF-β (42 and 78 days) or the CXCR1/2 (20 or 37 days) inhibitor. Magnification 400x. B,C) Quantification of the TGF-β1 (B) and CXCL1 (C) content of lungs from *Gata1^low* mice treated either with vehicle or inhibitors for P-SEL, TGF-β1 or CXCR1/2, as indicated. Data are presented as Mean (±SD) and as values per individual mice (each symbol a different mouse) and were analyzed by one-way Anova. P values statistically significant are indicated within the panels.

Figure 13. Only the treatment with the P-SEL inhibitor increases the GATA1 content of the mega-karyocytes in the lung from *Gata1^low* mice. Double immune-fluorescence staining with anti-GATA1 (green) and CD42b of lung sections from representative *Gata1^low* mice treated either with vehicle or with the P-SEL (day 12 and 40), TGF-β1 (42 and 78 days) or CXCLR1/2 (20 or 37 days) inhibitors, as indicated. Nuclei were highlighted with DAPI (blue). Magnification 400x. The signal captured in the individual chan-nels are presented in Figure S10). Total number of MK (CD42bpos cells) and of MK expressing detectable levels of GATA1 in the lungs from *Gata1^low* mice treated either with vehicle or the P-SEL, TGF-β1 or CXCR1/2 inhibitor, as indicated. Data are presented as Mean (±SD) and as values per individual mice (each symbol a different mouse) and were analyzed by one-way Anova. P values statistically significant are indicated within the panels.