Prothrombin Knockdown Protects Podocytes and Reduces Proteinuria in Glomerular Disease

Running Headline: Prothrombin Modulates Podocyte Health

Amanda P. Waller,[1] Katelyn J. Wolfgang,[1] Eman Abdelghani,[1,2] Zachary S. Stevenson,[1] Kaushik Muralidharan,[1] Tasha K. Wilkie,[1] William E. Smoyer,[1,3] Marvin T. Nieman,[4] and Bryce A. Kerlin[1,2]

[1]Center for Clinical & Translational Research, The Abigail Wexner Research Institute at Nationwide Children's, Columbus, OH

[2]Division of Hematology/Oncology/Blood & Marrow Transplantation, Department of Pediatrics, The Ohio State University College of Medicine, Columbus, OH

[3]Division of Nephrology and Hypertension, Department of Pediatrics, The Ohio State University College of Medicine, Columbus, OH

[4]Department of Pharmacology, Case Western Reserve University School of Medicine, Cleveland, OH

Support: This project was supported by grants K08DK103982, R03DK118315, and R01DK124549 (to BAK) from the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Additional support was provided by the George & Elizabeth Kelly Foundation (Lewis Center, OH; to BAK). This study was also supported (in part) by research funding from the CSL Behring – Professor Heimburger Award to BAK.

Corresponding Author: Bryce A. Kerlin, MD
Center for Clinical & Translational Research
The Abigail Wexner Research Institute at Nationwide Children's
700 Children's Dr., W325
Columbus, OH 43205
P: 614-355-2729
F: 614-722-3330
bryce.kerlin@nationwidechildrens.org

Abstract Word Count: 221
Text Word Count: 3714
Figures: 7
Tables: 0
Supplemental Figures: 10
Supplemental Tables: 2
References: 98

Keywords: Prothrombin, Proteinuria, Podocyte, Nephrotic Syndrome, Glomerular Disease

ABSTRACT

Chronic kidney disease (CKD) is a leading cause of death, and its progression is driven by glomerular podocyte injury and loss, manifesting as proteinuria. Proteinuria includes urinary loss of coagulation zymogens, cofactors, and inhibitors. Importantly, both CKD and proteinuria significantly increase the risk of thromboembolic disease. Prior studies demonstrated that anticoagulants reduced proteinuria in rats and that thrombin injured cultured podocytes. Herein we aimed to directly determine the influence of circulating prothrombin on glomerular pathobiology. We hypothesized that (pro)thrombin drives podocytopathy, podocytopenia, and proteinuria. Glomerular proteinuria was induced with puromycin aminonucleoside (PAN) in Wistar rats. Circulating prothrombin was either knocked down using a rat-specific antisense oligonucleotide or elevated by serial intravenous infusions of prothrombin protein, which are previously established methods to model hypo- (LoPT) and hyper-prothrombinemia (HiPT), respectively. After 10 days (peak proteinuria in this model) plasma prothrombin levels were determined, kidneys were examined for (pro)thrombin co-localization to podocytes and podocytopathy, podocytopenia, proteinuria, and plasma albumin were measured. LoPT significantly reduced podocyte exposure to prothrombin, podocytopathy, and proteinuria with improved plasma albumin. In contrast, HiPT significantly increased podocytopathy and proteinuria. Podocytopenia was significantly improved in LoPT vs. HiPT rats. In summary, prothrombin knockdown ameliorated podocytopathy, podocytopenia, and proteinuria in rats with PAN-induced glomerular proteinuria. Thus, (pro)thrombin antagonism may be a viable strategy to simultaneously provide thromboprophylaxis and prevent podocytopathy-mediated CKD progression.

INTRODUCTION

Kidney disease is the 8[th] leading cause of death in the U.S. and cardiovascular disease is the leading cause of mortality in patients with chronic kidney disease (CKD).[1-9] Glomerular diseases are a leading cause of CKD and end-stage kidney disease (ESKD).[6, 10] The glomerular filtration barrier is composed of three layers: (1) fenestrated endothelial cells, (2) collagen basement membrane, and (3) podocytes which interdigitate to form the slit diaphragm (a specialized adherence junction).[11] Proteinuria, glomerular podocyte injury (podocytopathy) leading to podocyte loss (podocytopenia), and glomerulosclerosis are hallmarks of glomerular disease-mediated CKD.[1, 4, 12-16] Podocytes are terminally differentiated epithelial cells that maintain glomerular endothelial cell phenotype via paracrine signaling, synthesize basement membrane components, and mechanically resist pulsatile intracapillary blood pressure.[11, 17, 18] Podocyte function is thus critical to glomerular filtration barrier maintenance and podocyte injury and/or death leads to urinary loss of plasma proteins (proteinuria).[18-21] While proteinuria due to acute podocyte dysfunction is potentially reversible, it becomes irreversible with ≥20% podocyte depletion and glomerulosclerotic lesions form on the extraluminal surface of glomerular capillaries in the areas of podocyte loss.[1, 12, 22] Proteinuria, in turn, is a key driver of CKD progression, leading to renal tubular inflammation and interstitial fibrosis.[1] Thus, podocyte preservation and proteinuria constraint is crucial to delay or prevent both CKD progression and CKD-related cardiovascular disease.

Nephrotic syndrome (NS) is comprised of a group of primary glomerular diseases that collectively are a leading cause of ESKD, the most severe form of CKD.[6] NS is clinically characterized by massive proteinuria, hypoalbuminemia, edema, and dyslipidemia.[23, 24] The monogenic, familial forms of NS are predominantly due to mutations in podocyte-specific genes and idiopathic NS is typified by podocytopathy.[1, 25] Rodent models of podocytopathy-mediated proteinuria have thus become important tools for the investigation of NS pathogenesis and CKD progression.[10, 12] The massive proteinuria of NS includes urinary loss of albumin and other plasma proteins, including coagulation system zymogens, cofactors, and inhibitors with variable synthetic compensation that manifests as an acquired hypercoagulopathy.[26, 27] Thus, NS is itself a profoundly prothrombotic condition with increased risk for both venous and arterial thrombosis, even before CKD onset.[8, 26, 28-33]

We and others have recently shown that prothrombin (*F2*) may directly injure podocytes during NS.[34-36] Thrombin injured conditionally immortalized rat podocytes via protease-activated receptor (PAR) 1 in a PAR4-dependent manner and thrombin inhibition reduced proteinuria in two mechanistically distinct rat NS models.[34, 36-38] Moreover, prothrombin colocalization to podocytes *in vivo* was proportional to proteinuria, strongly suggesting that prothrombin, originating from the plasma compartment, interacts with podocytes to drive podocytopathy, podocytopenia, and proteinuria during NS. The objective of this study was to test this hypothesis in the well-established puromycin aminonucleoside (PAN)-induced rat NS model.[10, 34, 36, 39]

Transgenic rat *F2* hypo- and hyper-expressing models are not yet available to facilitate examination of this phenomenon.[40-43] However, antisense oligonucleotides (ASOs) have been used to successfully knock-down mouse prothrombin expression to levels comparable to those seen in $F2^{lox/-}$ knockdown mice.[44-47] Thus, ASOs provide an alternative means to achieve *in vivo* prothrombin knockdown. Intravenous administration of human prothrombin protein has been used to model hyperprothrombinemia in mice.[48] We thus utilized ASO-mediated prothrombin knockdown and serial prothrombin infusions to mimic hypo- and hyper-prothrombinemia (LoPT and HiPT), respectively. Using this strategy, we demonstrated that plasma prothrombin levels modulate both podocyte health and proteinuria in experimental rat NS.

METHODS

Animals

All experimental rat protocols were approved by the Institutional Animal Care and Use Committee at the Abigail Wexner Research Institute, in accordance with the NIH Guide for the Care and Use of Laboratory Animals. Wistar rats (body weight ~150 g, age ~45-50 d) were used for all experiments (**Table S1**). Subcutaneous ASO or intravenous prothrombin was administered at the indicated doses and frequencies to achieve LoPT and HiPT conditions, respectively (**Figure S1)**. Proteinuria was induced with a single tail vein injection of PAN (MilliporeSigma, Burlington, MA; 50 mg/kg diluted in saline) on Day 0.[10, 34, 39] Anti-*F2* ASO (401027) treatment (60 mg/kg subcutaneous twice weekly) was begun on Day -17.5 to induce LoPT prior to onset of podocyte injury and continued through Day +9. Prothrombin infusions were begun on Day 0, immediately following PAN (or saline) administration. Control rats received twice weekly scrambled ASO (141923) on Days -17.5 to +9, saline tail vein injections on Days 0, 3, 6, and 9 (instead of PAN and/or prothrombin). Sham rats were treated identically to control rats but did receive PAN on Day 0. Morning spot urine samples were collected on Days 0 (before PAN or saline; **Figure S2**) and 10 to determine urinary protein-to-creatinine ratio. Following Day 10 urine collection, the rats were anesthetized with 3% isoflurane and blood was collected from the inferior vena cava through a 23-G needle into final concentration 0.32% NaCitrate/1.45 µM Corn Trypsin Inhibitor (CTI; Prolytix, Essex Junction, VT) and processed to Platelet Poor Plasma (PPP), as previously described.[39] Urine and plasma were stored at -80ºC until analyzed. Following exsanguination, both kidneys were removed, decapsulated, and placed in ice cold PBS. One quarter kidney was set aside for immunohistology and glomeruli were isolated from the remainder of both kidneys.

Glomerular Isolation and Podocyte Flow Cytometry

Glomeruli were isolated from renal cortex using a standard sequential sieving method, washed and resuspended in ice cold PBS, then manually counted in a hemocytometer, as previously described.[49-51] An aliquot of the glomerular suspension (~2,500 glomeruli) was dissociated to a single cell suspension in Digestion Buffer at 37ºC on a 1400/min shaker with intermittent shearing through an 18-G needle for 26 minutes. Tissue debris was

removed with a 40-micron filter, the cells were washed with 10 mL HBSS, centrifuged at 1500 rpm for 5 min, then resuspended in 1 mL HBSS supplemented with 0.1% BSA and 4',6-diamidino-2-phenylindole (DAPI; 1 µg/mL). The cells were then prepared for flow cytometry analysis by staining with Live/Dead near IR dye and FITC-conjugated synaptopodin antibody (Fitzgerald, North Acton, MA; 1:5) in Perm/Wash buffer. The cells were analyzed on a LSR II flow cytometer (BD Biosciences) and synaptopodin expression was gated only on cells with a DAPI-positive intact nucleus. Isotype controls, unstained cells, and an open channel were used to identify and calibrate for autofluoresence. Synaptopodin-positive podocytes were normalized to bead standards to estimate podocyte counts which were then divided by the starting number of glomeruli to enumerate podocytes/glomerulus. Reassuringly, we observed a direct linear relationship between podocyte counts determined by flow cytometry and a histologic method (see Supplemental Methods and **Figure S3**).[52] A second aliquot of the glomerular cell suspension was stained with FITC-TUNEL (1:5) and APC-conjugated synaptopodin antibody (1:4) to determine % TUNEL-positive podocytes. Flow cytometry data points for each individual rat represents an analysis of 100,000 events.

Immunofluorescence Histology

Paraffin-embedded rat kidney sections were deparaffinized and processed for synaptopodin and thrombin co-immunofluorescence, as previously described.[34] However, the sections were photobleached for 72 hours prior to deparaffinization using a custom-built lightbox to reduce background autofluorescence.[53] Lightbox construction was adapted from previous descriptions using mirrored acrylic panels and a broad-spectrum LED panel.[54-56] Images were captured with a Leica DMI 4000B inverted fluorescence microscope (Deerfield, IL) and analyzed with ZEN Black software (Zeiss USA, Thornwood, NY). Colocalization analysis was performed on a pixel-by-pixel basis, wherein every pixel within the defined region of interest is plotted in a scattergram based on fluorescence intensity of each channel. The software calculates the colocalization coefficient as the proportion of colocalized pixels (positive on both channels) within the total population of a single channel, green (synaptopodin-positive) for the purposes of this study.[57]

Coagulation Parameters

Plasma prothrombin concentration was measured with rat- and human-specific ELISA (MyBioSource Inc, San Diego, California). Prothrombin activity was determined chromogenically using a commercially available assay (Rox Prothrombin; DiaPharma, West Chester, OH), as previously described,[58] and are expressed as a percentage of rat pooled normal plasma (rPNP). Endogenous thrombin potential (ETP) was determined on a 1:1 dilution of PPP with the Technothrombin TGA kit (Technoclone, Vienna, Austria) using TGA RC low reagent, and read on a Spectramax M2 fluorescent plate reader (Molecular Devices, Sunnyvale, California), as previously described.[39, 58]

Statistical Analyses

One- or two-way ANOVA (analysis of variance) for multiple group comparisons, using SigmaStat software (Systat, San Jose, CA). When a significant difference was identified by ANOVA, post hoc tests were performed using the Student-Newman-Keuls technique. Statistical significance was defined as $P<0.05$. Figures were prepared using GraphPad Prism (Boston, MA). Data are presented as mean ± SE.

See Supplemental Methods for Additional Detail.

RESULTS

Antisense Oligonucleotide-Mediated Hypoprothrombinemia in Healthy Rats

ASO 401027, which is 100% complementary to a 3' segment of rat *F2* mRNA, effectively abolished *F2* mRNA expression in primary rat hepatocytes vs. control ASO (0.00±0.00 vs. 1.0±0.01 relative fold change; *P*<0.001; **Figure S4**), whereas ASO 401025 (75% complementary to rat *F2* mRNA) less effectively suppressed *F2* expression (0.63±0.14 relative fold change; *P*=0.024). ASO 401027 also reduced prothrombin activity in primary rat hepatocyte protein lysates to 1.41±0.16 vs. 9.13±0.13 mIU/mL in control ASO lysates (*P*<0.001), whereas ASO 401025 less efficiently reduced activity to 2.31±0.10 mIU/mL (*P*<0.001). In dose finding studies, ASO 401027 was administered subcutaneously to healthy rats twice weekly for 17.5 days (**Figure 1**). These data suggested that maximal plasma prothrombin activity reduction (~32% of rPNP) was achieved with 60 mg/kg/dose. To mimic the planned full-length experiment, this dose (60 mg/kg twice weekly) was then continued for 27.5 days after which plasma prothrombin activity was reduced to 31.6±3.52% (vs. 100.3±1.45% in rPNP; *P*<0.001) and endogenous thrombin potential (ETP) was reduced to 514.3±58.7 nM*min (vs. 2451.0±140.5 nM*min; *P*<0.001). Thus, whereas prothrombin activity was reduced by ~68%, ETP was reduced by ~71% suggesting that circulating prothrombin reduction impaired thrombin-dependent coagulation cascade amplification.[59] Hepatic synthesis is the major source of circulating prothrombin and hepatic *F2* mRNA expression was essentially abolished by this 27.5 day ASO treatment regimen (*P*<0.01).[60, 61] Thus, ASO 401027 effectively induced sustained LoPT in otherwise healthy rats.

Serial Prothrombin Infusion-Mediated Hyperprothrombinemia in Healthy Rats

The hemostatic system is highly conserved amongst vertebrate species, including mammals.[62, 63] Thus, rPNP thrombin generation was augmented by addition of human prothrombin as expected (**Figure S5**). Moreover, rat thrombomodulin effectively regulated the spiked-in human prothrombin. These observations are similar to those previously observed using human prothrombin added to mouse PNP and indicate that human prothrombin is compatible with rat pro- and anti-coagulant systems.[48] In preliminary experiments, intravenously administered

human prothrombin (31.25 mg/kg) generated peak plasma prothrombin activity of ~207% 1-hour post-dose with an apparent half-life of ~61 hours (**Figure 2**), which is similar to that previously reported in humans.[64] To mimic the planned full-length experiment, 31.25 mg/kg prothrombin was administered as a loading dose on day 0 with maintenance doses of 16.67 mg/kg on days 3, 6, and 9. On day 10, 24 hours after the day 9 dose, prothrombin activity was 186.8±10.4% (vs. 100.3±1.45% in rPNP; $P<0.001$). Similarly, day 10 ETP was elevated to 5766.0±207.0 (vs. 2451.0±140.5 nM*min; $P<0.001$). Thus, whereas prothrombin activity was enhanced by ~86%, ETP was enhanced by ~135% suggesting that elevated circulating prothrombin enhanced thrombin-dependent coagulation cascade amplification. These data demonstrate that human prothrombin induces hyperprothrombinemia (HiPT) in otherwise healthy rats for up to 10 days.

Prothrombin Modulation during Puromycin Aminonucleoside-Induced Rat Nephrotic Syndrome

Peak NS disease activity occurs at day 8-11 following PAN administration, we therefore evaluated prothrombin levels at day 10 to ensure that the manipulations observed above in otherwise healthy rats were sustained during PAN-mediated NS (PAN-NS).[10, 34, 39] Similar to the primary rat hepatocyte and *in vivo* dose finding studies described above, ASO 401027 treated rats (LoPT) had significantly reduced hepatic *F2* expression ($P<0.05$ vs Control) whereas *F2* expression was not significantly affected by PAN (Sham; $P=0.35$ vs Control) or prothrombin infusions (HiPT; $P=0.28$ and $P=0.98$ vs. Control and Sham, respectively; **Figure 3**). PAN-NS rats treated with ASO 401027 (LoPT) exhibited significantly reduced day 10 plasma prothrombin concentrations (0.92±0.10 μM) vs. Control (4.07±0.17 μM) and Sham (4.20±0.15 μM) whereas prothrombin infusions (HiPT) resulted in significantly elevated concentrations (8.16±0.38 μM; **Figures S6 and 3**). The LoPT and HiPT changes in prothrombin concentrations translated into significant changes in chromogenic prothrombin activity: LoPT 29.5±3.9% and HiPT 222.3±12.8% vs. Control (105.3±1.2%) and Sham (123.6±7.7%; **Figure 3**). Congruently, ETP was reduced to 306.7±62.9 nM*min in LoPT (vs. 2893.0±122.0 and 3417.0±235.8 nM*min in Control and Sham rats respectively; $P<0.001$) whereas HiPT rats had significantly elevated ETP (4406.0±407.2 nM*min; $P<0.05$). These data demonstrate successful manipulation of plasma prothrombin levels during rat PAN-NS.

Plasma Prothrombin Levels Dictated Thrombin-Podocyte Interactions during Puromycin Aminonucleoside-Induced Rat Nephrotic Syndrome

We previously demonstrated that thrombin colocalized to podocytes during rat PAN-NS.[34] We thus predicted that thrombin-podocyte interactions would fluctuate with plasma prothrombin levels. As expected, thrombin-podocyte colocalization was significantly reduced in LoPT rats (7.8±1.2% vs. 18.9±1.8% in Sham; P<0.01) to levels similar to Control (5.2±1.5%; P=0.30; **Figure 4**). Meanwhile, thrombin-podocyte colocalization was significantly higher than Control in both Sham (P<0.01) and HiPT (18.1±3.4; P<0.05) rat PAN-NS. The latter data suggest that thrombin interactions with podocyte-expressed protease-activated receptors may be saturated at physiologic prothrombin concentrations.[34] Nonetheless, these data strongly suggest that plasma prothrombin levels dictate podocyte exposure to thrombin during glomerular proteinuria.

Hypoprothrombinemia Diminished *in situ* Podocyte Injury

We and others previously demonstrated that *in vitro* thrombin-exposure leads to DNA nicking that is consistent with terminal podocyte injury.[34, 35] Importantly, podocytes may be terminally injured prior to detachment from the glomerular capillary surface.[19] Thus, to evaluate the health status of *in situ* podocytes we performed TUNEL assays on dissociated cells from isolated glomeruli. Importantly, neither LoPT nor HiPT altered podocyte DNA-nicking in otherwise healthy rats (**Figure S7**). However, HiPT significantly increased TUNEL-positive podocytes (17.4±0.5%) in PAN-NS (11.7±1.8% vs Sham; P<0.05; **Figure 5**). In contrast, LoPT reduced TUNEL-positivity (6.9±1.5%; P<0.05 vs. Sham) to levels not significantly different from Control (6.5±1.5%; P=0.90). Podocyte DNA-nicking was correlated with prothrombin levels, and thrombin-colocalization during PAN-NS (**Figure S8**).

Hyperprothrombinemia Reduced *in situ* Podocyte Survival

Glomerular filtration barrier function and CKD progression are highly dependent on glomerular podocyte loss. We thus evaluated *in situ* podocyte survival by counting remaining podocytes in dissociated glomerular cells. Whereas LoPT and HiPT did not alter podocyte counts in otherwise healthy rats (**Figure S7**), HiPT PAN-NS rats

had significantly lower podocyte counts (134.2±21.5 podocytes/glomerulus) vs. either LoPT (350.9±65.3; *P*<0.01) or Control (389.0±41.8; *P*<0.001; **Figure 6**). However, the LoPT and HiPT conditions did not translate into significantly improved or diminished podocyte counts vs. Sham. Meanwhile, podocyte counts were correlated with prothrombin levels, thrombin-colocalization, and podocyte injury in PAN-NS rats (**Figure S9**). These data suggest that hyperprothrombinemia may accelerate podocyte loss during PAN-NS.

Prothrombin Regulated Proteinuria and Plasma Albumin during Puromycin Aminonucleoside-Induced Rat Nephrotic Syndrome

Podocyte injury and loss leads to glomerular filtration barrier dysfunction resulting in proteinuria and hypoalbuminemia.[10, 16, 18, 20] Consistent with preserved podocyte function, LoPT PAN-NS rats had significantly lower day 10 proteinuria values in comparison to Sham (6.6±0.7 vs. 15.0±3.7 mg/mg creatinine; *P*<0.05) that was not significantly different than Control (1.0±0.1 mg/mg; **Figure 7**). Meanwhile, HiPT rats had significantly worse proteinuria (34.9±9.3 mg/mg; *P*<0.05). Consistent with improved overall protein homeostasis, plasma albumin was significantly improved in LoPT vs. Sham (3.32±0.03 vs. 3.07±0.08 g/dL; *P*<0.05) to values that were not significantly different from Control (3.41±0.02 g/dL). However, HiPT did not significantly worsen hypoalbuminemia (3.05±0.1 g/dL). Importantly, neither LoPT nor HiPT altered proteinuria or plasma albumin in otherwise healthy rats (**Figure S7**). In PAN-NS proteinuria was significantly correlated with prothrombin levels and podocyte injury (**Figure 7**). Similarly, plasma albumin was correlated with prothrombin levels, podocyte injury, podocyte survival, and proteinuria (**Figure S10**).

DISCUSSION

Proteinuria, podocytopathy, and podocytopenia all contribute to glomerular dysfunction and progression of chronic kidney disease.[8, 16, 22, 65-71] The data presented in this study demonstrate that circulating levels of prothrombin (the zymogen precursor of thrombin) modulated *in vivo* podocyte thrombin exposure, injury, survival, and function in the setting of proteinuric glomerular disease. We and others have previously shown that thrombin injures cultured podocytes *in vitro* and that *in vivo* thrombin inhibition preserves glomerular function as determined by proteinuria reduction.[34-38] In contrast to the previous studies which utilized anticoagulant agents which may have off-target effects, this study directly manipulated prothrombin levels *in vivo*, providing direct evidence that (pro)thrombin is involved in glomerular disease pathogenesis. We previously demonstrated that thrombin injures *in vitro* podocytes via its cognate protease-activated receptors.[34] Collectively, these data suggest that thrombin originates as circulating prothrombin that is pathologically filtered during proteinuria leading to intraglomerular thrombin generation and direct podocytopathic thrombin signaling. Thus, interruption of this signaling pathway may represent a novel therapeutic target to slow or halt glomerular disease progression toward CKD/ESKD.

Mounting indirect evidence from *in vitro* studies and *in vivo* pharmacologic manipulation strongly suggest that thrombin is a pathological driver of podocyte injury in the setting of proteinuria.[34-38] The present study provides direct evidence supporting this hypothesis. HiPT rats exhibiting prothrombin levels ~195% of normal developed worse proteinuria along with increased podocyte injury. The human thrombophilic single nucleotide polymorphism, *F2* G20210A (rs1799963), is associated with prothrombin levels ~115-170% of normal.[48] To our knowledge, rs1799963 has not been associated with enhanced CKD progression or incidence, but most studies were not directly looking for such a signal, were likely underpowered, or were investigating CKD-related thrombotic risk.[72-74] It is also possible that since the reference interval for prothrombin extends to 150% that the mildly increased levels associated with this polymorphism do not lead to meaningfully increased CKD risk.[75] However, the present data suggest that rs1799963 should be more carefully analyzed as a potential driver of proteinuria-mediated CKD progression. Protein overload, achieved with ≥3 g/kg cumulative albumin dose, is another commonly employed method to induce transient proteinuria in animal models.[10, 76] However, it is unlikely

that HiPT mimics protein overload since the cumulative prothrombin protein dose (81.26 mg/kg) used in these studies is only 2.7% of the minimum albumin dose required to induce proteinuria and did not induce proteinuria in the absence of PAN (**Figure S7**).

In contrast, highly specific ASO-mediated prothrombin knockdown (to ~24% of normal) resulted in diminished podocyte thrombin exposure, podocytopathy, podocytopenia, proteinuria, and improved plasma albumin. Prothrombin is predominantly synthesized in the liver but several other organs, including the kidney, also produce prothrombin.[61] Moreover, ASO therapy has been demonstrated to reduce target mRNA in many tissues, including both liver and kidney.[77] Thus, while it is clear from these experiments that ASO-mediated prothrombin knockdown improves podocyte function and survival, the origin of podocytopathic prothrombin remains ill-defined. However, manipulation of circulating prothrombin did not induce proteinuria in healthy animals, strongly suggesting that proteinuria is a prerequisite for (pro)thrombin-mediated podocyte injury and that podocytopathic prothrombin thus likely originates from the plasma compartment. Alternatively, podocyte injury or proteinuria may stimulate glomerular cells or adjacent tissues to upregulate local prothrombin synthesis. A previous study identified prothrombin protein in tubular epithelial cells, but not glomeruli, suggesting that juxtaglomerular tubules could be a relevant local source of prothrombin synthesis.[61]

We previously demonstrated that *in vitro* thrombin-mediated podocytopathy is dependent upon activation of podocyte-expressed protease-activated receptors.[34] Activation of these receptors is dependent upon enzymatic cleavage and exposure of a cryptic tethered ligand by thrombin or other enzymes.[78] Once generated, thrombin has a <1 minute half-life *in vivo*,[75, 79-81] suggesting that prothrombin is most likely converted to thrombin locally in the glomerulus, perhaps even on the podocyte surface.[79, 82] However, the mechanism by which prothrombin is converted to intraglomerular thrombin during proteinuria remains undefined. The canonical hemostatic mechanism is cleavage of prothrombin by the factor Xa/Va prothrombinase complex. While prothrombin and cofactor V have been found in the urine of nephrotic patients, factor X has not.[83, 84] Additionally, a proteomic study has demonstrated that neither factor X nor V protein are expressed by healthy murine podocytes.[85] Thus, while podocytes do express tissue factor, the complete complement of canonical prothrombinase components

do not appear to be available.[82, 86, 87] Alternatively, adjacent kidney cells have been reported to synthesize both factor X and V.[88] Moreover, these components may originate from the plasma during proteinuria. It is thus possible that the canonical prothrombinase mechanism is at work in the diseased glomerulus. Alternatively, one or more putative tissue prothrombinases may explain glomerular thrombin activity.[89] Amongst the latter, fibrinogen-like protein 2 (*Fgl2*) is expressed by podocytes and is thus a leading candidate.[85, 90-92] This possibility is particularly intriguing as it would provide a potential target to ameliorate (pro)thrombin-mediated podocytopathy without increasing the risk of CKD-related bleeding.[93]

There are several limitations to consider when interpreting these experiments. Although the mammalian coagulation system is highly conserved, it is possible that the use of human prothrombin in these studies could have resulted in subtle species-dependent differences in prothrombin activation efficiency or protease-activated receptor signaling responses in the rat glomerulus. The persistent hyperprothrombinemia after multiple human prothrombin injections argues against xenoimmunity in this model for up to 10 days. Nonetheless, the use of rat prothrombin or creation of a gene duplication-mediated hyperprothrombinemia model would provide more specific evidence. Longer term experiments should be conducted to determine if these short-term improvements translate into slowed or halted CKD progression such as improvements in glomerular filtration rate loss. Although PAN is a relatively podocyte-specific toxin, it is possible that there are off-target PAN effects on the coagulation system. However, PAN reportedly does not alter protein synthesis in isolated rat hepatocytes, making it unlikely that PAN directly altered coagulation protein synthesis.[94] Finally, these data, generated in a rat NS model, may not be generalizable to other forms of glomerular disease that lead to CKD such as diabetic nephropathy, lupus nephritis, or hypertensive nephropathy and thus, should be examined in other animal models of glomerular disease.

In summary, prothrombin is able to modulate both podocyte function and survival in the PAN model of nephrotic syndrome. To determine the importance of prothrombin-mediated podocyte injury and its potential as a novel therapeutic target to slow CKD progression will require additional studies. Important next steps include defining the relevance of this mechanism in CKD models, determining the prothrombinase mechanism driving thrombin

formation in the glomerulus, and evaluating its suitability as a druggable target using thrombin inhibitors, especially those that are already approved by relevant regulatory agencies for use in humans with kidney disease and may thus be suitable for rapid clinical translation.[95] Because proteinuria imparts increased thrombotic risk,[29, 31, 96-98] renally excreted anticoagulant agents that may inhibit glomerular thrombin should be investigated as a novel means to simultaneously reduce thrombotic complications and slow proteinuria-mediated CKD progression.

***Acknowledgments*:**

This project was supported by grants K08DK103982, R03DK118315, and R01DK124549 (to BAK) from the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Additional support was provided by the George & Elizabeth Kelly Foundation (Lewis Center, OH; to BAK). This study was also supported (in part) by research funding from the CSL Behring – Professor Heimburger Award to BAK. The authors are indebted to Dr. Brett P. Monia, Ionis Pharmaceuticals, for providing the antisense oligonucleotides, Matthew Curtis, Carl Zeiss Microscopy, for developing the Zen Blue software algorithm for histological podocyte counting, and Dave Dunaway, Abigail Wexner Research Center, for assistance with flow cytometry protocol refinement.

***Authorship Contributions*:**
Experiments were conducted by APW, KJW, EA, ZSS, KM, TKW, and BAK. APW and BAK analyzed data; APW and BAK prepared the figures and wrote the paper; all authors read, edited, and approved the final version; WES and MTN provided important intellectual contributions; BAK was responsible for funding, oversight, and coordination of the project.

***Conflict of Interest Disclosures*:** The authors declare no competing financial interests.

REFERENCES

1.   Breyer MD, Susztak K. The next generation of therapeutics for chronic kidney disease. *Nat Rev Drug Discov* 2016; **15:** 568-588.

2.   Gansevoort RT, Correa-Rotter R, Hemmelgarn BR*, et al.* Chronic kidney disease and cardiovascular risk: epidemiology, mechanisms, and prevention. *Lancet* 2013; **382:** 339-352.

3.   Said S, Hernandez GT. The link between chronic kidney disease and cardiovascular disease. *J Nephropathol* 2014; **3:** 99-104.

4.   Webster AC, Nagler EV, Morton RL*, et al.* Chronic Kidney Disease. *Lancet* 2017; **389:** 1238-1252.

5.   Weiner DE, Carpenter MA, Levey AS*, et al.* Kidney function and risk of cardiovascular disease and mortality in kidney transplant recipients: the FAVORIT trial. *Am J Transplant* 2012; **12:** 2437-2445.

6.   Saran R, Robinson B, Abbott KC*, et al.* US Renal Data System 2017 Annual Data Report: Epidemiology of Kidney Disease in the United States. *Am J Kidney Dis* 2018; **71:** A7.

7.   Virani SS, Alonso A, Aparicio HJ*, et al.* Heart Disease and Stroke Statistics-2021 Update: A Report From the American Heart Association. *Circulation* 2021; **143:** e254-e743.

8.   Go AS, Tan TC, Chertow GM*, et al.* Primary Nephrotic Syndrome and Risks of ESKD, Cardiovascular Events, and Death: The Kaiser Permanente Nephrotic Syndrome Study. *J Am Soc Nephrol* 2021; **32:** 2303-2314.

9.   Kochanek KD, Xu J, Arias E. Mortality in the United States, 2019. *NCHS Data Brief* 2020**:** 1-8.

10.   Pippin JW, Brinkkoetter PT, Cormack-Aboud FC*, et al.* Inducible rodent models of acquired podocyte diseases. *Am J Physiol Renal Physiol* 2009; **296:** F213-229.

11.   Butt L, Unnersjo-Jess D, Hohne M*, et al.* A molecular mechanism explaining albuminuria in kidney disease. *Nat Metab* 2020; **2:** 461-474.

12.   Lemley KV. Glomerular pathology and the progression of chronic kidney disease. *Am J Physiol Renal Physiol* 2016; **310:** F1385-1388.

13.   Durvasula RV, Shankland SJ. Podocyte injury and targeting therapy: an update. *Current opinion in nephrology and hypertension* 2006; **15:** 1-7.

14.   Rask-Madsen C, King GL. Diabetes: Podocytes lose their footing. *Nature* 2010; **468:** 42-44.

15.   Yu D, Petermann A, Kunter U*, et al.* Urinary podocyte loss is a more specific marker of ongoing glomerular damage than proteinuria. *J Am Soc Nephrol* 2005; **16:** 1733-1741.

16.   Wickman L, Afshinnia F, Wang SQ*, et al.* Urine podocyte mRNAs, proteinuria, and progression in human glomerular diseases. *J Am Soc Nephrol* 2013; **24:** 2081-2095.

17.   Garg P. A Review of Podocyte Biology. *Am J Nephrol* 2018; **47 Suppl 1:** 3-13.

18.   Nagata M. Podocyte injury and its consequences. *Kidney Int* 2016; **89:** 1221-1230.

19.   Mundel P, Shankland SJ. Podocyte biology and response to injury. *J Am Soc Nephrol* 2002; **13:** 3005-3015.

20.   Shankland SJ. The podocyte's response to injury: role in proteinuria and glomerulosclerosis. *Kidney Int* 2006; **69:** 2131-2147.

21.   Kopp JB, Anders HJ, Susztak K*, et al.* Podocytopathies. *Nat Rev Dis Primers* 2020; **6:** 68.

22.   Wharram BL, Goyal M, Wiggins JE*, et al.* Podocyte depletion causes glomerulosclerosis: diphtheria toxin-induced podocyte depletion in rats expressing human diphtheria toxin receptor transgene. *J Am Soc Nephrol* 2005; **16:** 2941-2952.

23.   Eddy AA, Symons JM. Nephrotic syndrome in childhood. *Lancet* 2003; **362:** 629-639.

24.   Orth SR, Ritz E. The nephrotic syndrome. *N Engl J Med* 1998; **338:** 1202-1211.

25.   Lovric S, Ashraf S, Tan W*, et al.* Genetic testing in steroid-resistant nephrotic syndrome: when and how? *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 2016; **31:** 1802-1813.

26.   Schlegel N. Thromboembolic risks and complications in nephrotic children. *Semin Thromb Hemost* 1997; **23:** 271-280.

27.   Andersen RF, Palmfeldt J, Jespersen B*, et al.* Plasma and urine proteomic profiles in childhood idiopathic nephrotic syndrome. *Proteomics Clin Appl* 2012; **6:** 382-393.

28.   Kerlin BA, Ayoob R, Smoyer WE. Epidemiology and pathophysiology of nephrotic syndrome-associated thromboembolic disease. *Clin J Am Soc Nephrol* 2012; **7:** 513-520.

29.   Kerlin BA, Blatt NB, Fuh B*, et al.* Epidemiology and risk factors for thromboembolic complications of childhood nephrotic syndrome: a Midwest Pediatric Nephrology Consortium (MWPNC) study. *J Pediatr* 2009; **155:** 105-110, 110 e101.

30.   Loscalzo J. Venous thrombosis in the nephrotic syndrome. *N Engl J Med* 2013; **368:** 956-958.

31.   Mahmoodi BK, ten Kate MK, Waanders F*, et al.* High absolute risks and predictors of venous and arterial thromboembolic events in patients with nephrotic syndrome: results from a large retrospective cohort study. *Circulation* 2008; **117:** 224-230.

32.   Singhal R, Brimble KS. Thromboembolic complications in the nephrotic syndrome: pathophysiology and clinical management. *Thromb Res* 2006; **118:** 397-407.

33.   Waller AP, Troost JP, Parikh SV*, et al.* Nephrotic syndrome disease activity is proportional to its associated hypercoagulopathy. *Thromb Res* 2021; **201:** 50-59.

34.   Sharma R, Waller AP, Agrawal S*, et al.* Thrombin-Induced Podocyte Injury Is Protease-Activated Receptor Dependent. *J Am Soc Nephrol* 2017; **28:** 2618-2630.

35.   Wang H, Madhusudhan T, He T*, et al.* Low but sustained coagulation activation ameliorates glucose-induced podocyte apoptosis: protective effect of factor V Leiden in diabetic nephropathy. *Blood* 2011; **117:** 5231-5242.

36.   Yamashita J, Nakajima K, Ohno Y*, et al.* Protective effects of antithrombin on puromycin aminonucleoside nephrosis in rats. *Eur J Pharmacol* 2008; **589:** 239-244.

37.   Baroni EA, Costa RS, da Silva CG, *et al.* Heparin treatment reduces glomerular injury in rats with adriamycin-induced nephropathy but does not modify tubulointerstitial damage or the renal production of transforming growth factor-beta. *Nephron* 2000; **84:** 248-257.

38.   Benchetrit S, Golan E, Podjarny E, *et al.* Low molecular weight heparin reduces proteinuria and modulates glomerular TNF-alpha production in the early phase of adriamycin nephropathy. *Nephron* 2001; **87:** 155-160.

39.   Kerlin BA, Waller AP, Sharma R, *et al.* Disease Severity Correlates with Thrombotic Capacity in Experimental Nephrotic Syndrome. *J Am Soc Nephrol* 2015; **26:** 3009-3019.

40.   Sun WY, Witte DP, Degen JL, *et al.* Prothrombin deficiency results in embryonic and neonatal lethality in mice. *Proc Natl Acad Sci U S A* 1998; **95:** 7597-7602.

41.   Xue J, Wu Q, Westfield LA, *et al.* Incomplete embryonic lethality and fatal neonatal hemorrhage caused by prothrombin deficiency in mice. *Proc Natl Acad Sci U S A* 1998; **95:** 7603-7607.

42.   Mullins ES, Kombrinck KW, Talmage KE, *et al.* Genetic elimination of prothrombin in adult mice is not compatible with survival and results in spontaneous hemorrhagic events in both heart and brain. *Blood* 2009; **113:** 696-704.

43.   Sun WY, Coleman MJ, Witte DP, *et al.* Rescue of prothrombin-deficiency by transgene expression in mice. *Thromb Haemost* 2002; **88:** 984-991.

44.   Crosby JR, Zhao C, Zhang H, *et al.* Reversing Antisense Oligonucleotide Activity with a Sense Oligonucleotide Antidote: Proof of Concept Targeting Prothrombin. *Nucleic Acid Ther* 2015; **25:** 297-305.

45.   Arumugam PI, Mullins ES, Shanmukhappa SK, *et al.* Genetic diminution of circulating prothrombin ameliorates multiorgan pathologies in sickle cell disease mice. *Blood* 2015; **126:** 1844-1855.

46.   Turpin B, Miller W, Rosenfeldt L, *et al.* Thrombin drives tumorigenesis in colitis-associated colon cancer. *Cancer Res* 2014; **74:** 3020-3030.

47.   Horowitz NA, Blevins EA, Miller WM, *et al.* Thrombomodulin is a determinant of metastasis through a mechanism linked to the thrombin binding domain but not the lectin-like domain. *Blood* 2011; **118:** 2889-2895.

48.   Aleman MM, Walton BL, Byrnes JR, *et al.* Elevated prothrombin promotes venous, but not arterial, thrombosis in mice. *Arterioscler Thromb Vasc Biol* 2013; **33:** 1829-1836.

49.   Agrawal S, Guess AJ, Chanley MA, *et al.* Albumin-induced podocyte injury and protection are associated with regulation of COX-2. *Kidney Int* 2014; **86:** 1150-1160.

50.   Smoyer WE, Gupta A, Mundel P, *et al.* Altered expression of glomerular heat shock protein 27 in experimental nephrotic syndrome. *J Clin Invest* 1996; **97:** 2697-2704.

51.   Guess A, Agrawal S, Wei CC, *et al.* Dose- and time-dependent glucocorticoid receptor signaling in podocytes. *Am J Physiol Renal Physiol* 2010; **299:** F845-853.

52.   Lemley KV, Bertram JF, Nicholas SB, *et al.* Estimation of glomerular podocyte number: a selection of valid methods. *J Am Soc Nephrol* 2013; **24:** 1193-1202.

53.   Viegas MS, Martins TC, Seco F, *et al.* An improved and cost-effective methodology for the reduction of autofluorescence in direct immunofluorescence studies on formalin-fixed paraffin-embedded tissues. *Eur J Histochem* 2007; **51:** 59-66.

54.   Sun Y, Ip P, Chakrabartty A. Simple Elimination of Background Fluorescence in Formalin-Fixed Human Brain Tissue for Immunofluorescence Microscopy. *J Vis Exp* 2017.

55.   Duong H, Han M. A multispectral LED array for the reduction of background autofluorescence in brain tissue. *J Neurosci Methods* 2013; **220:** 46-54.

56.   Neumann M, Gabel D. Simple method for reduction of autofluorescence in fluorescence microscopy. *J Histochem Cytochem* 2002; **50:** 437-439.

57.   Manders EMM, Verbeek FJ, Aten JA. Measurement of co-localization of objects in dual-colour confocal images. *J Microsc* 1993; **169:** 375-382.

58.   Waller AP, Agrawal S, Wolfgang KJ*, et al.* Nephrotic syndrome-associated hypercoagulopathy is alleviated by both pioglitazone and glucocorticoid which target two different nuclear receptors. *Physiol Rep* 2020; **8:** e14515.

59.   Wolberg AS, Campbell RA. Thrombin generation, fibrin clot formation and hemostasis. *Transfus Apher Sci* 2008; **38:** 15-23.

60.   Jamison CS, Degen SJ. Prenatal and postnatal expression of mRNA coding for rat prothrombin. *Biochim Biophys Acta* 1991; **1088:** 208-216.

61.   Stenberg LM, Brown MA, Nilsson E*, et al.* A functional prothrombin gene product is synthesized by human kidney cells. *Biochem Biophys Res Commun* 2001; **280:** 1036-1041.

62.   Kaye S, Stokol T. Hemostatic Testing in Companion Exotic Mammals. *Vet Clin North Am Exot Anim Pract* 2022; **25:** 613-630.

63.   Weyand AC, Shavit JA. Zebrafish as a model system for the study of hemostasis and thrombosis. *Curr Opin Hematol* 2014; **21:** 418-422.

64.   Monroe DM, Hoffman M, Roberts HR. Molecular Biology and Biochemistry of the Coagulation Factors and Pathways of Hemostasis. In: K. K, Lichtman MA, Beutler E, Kipps TJ*, et al.* (eds). *Williams Hematology*. McGraw-Hill Education: NY, NY, 2010.

65.   Fukuda A, Chowdhury MA, Venkatareddy MP*, et al.* Growth-dependent podocyte failure causes glomerulosclerosis. *J Am Soc Nephrol* 2012; **23:** 1351-1363.

66.   Fukuda A, Wickman LT, Venkatareddy MP*, et al.* Urine podocin:nephrin mRNA ratio (PNR) as a podocyte stress biomarker. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 2012; **27:** 4079-4087.

67.   Kikuchi M, Wickman L, Hodgin JB*, et al.* Podometrics as a Potential Clinical Tool for Glomerular Disease Management. *Seminars in nephrology* 2015; **35:** 245-255.

68.   Nishizono R, Kikuchi M, Wang SQ*, et al.* FSGS as an Adaptive Response to Growth-Induced Podocyte Stress. *J Am Soc Nephrol* 2017; **28:** 2931-2945.

69.   Sato Y, Wharram BL, Lee SK*, et al.* Urine podocyte mRNAs mark progression of renal disease. *J Am Soc Nephrol* 2009; **20:** 1041-1052.

70. Wanner N, Hartleben B, Herbach N, *et al.* Unraveling the role of podocyte turnover in glomerular aging and injury. *J Am Soc Nephrol* 2014; **25:** 707-716.

71. Zoja C, Abbate M, Remuzzi G. Progression of chronic kidney disease: insights from animal models. *Current opinion in nephrology and hypertension* 2006; **15:** 250-257.

72. Bauer A, Limperger V, Nowak-Gottl U. End-stage renal disease and thrombophilia. *Hamostaseologie* 2016; **36:** 103-107.

73. Lutz J, Menke J, Sollinger D, *et al.* Haemostasis in chronic kidney disease. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 2014; **29:** 29-40.

74. Tripathi G, Sankhwar SN, Sharma RK, *et al.* Role of thrombotic risk factors in end-stage renal disease. *Clin Appl Thromb Hemost* 2010; **16:** 132-140.

75. Mann KG, Brummel K, Butenas S. What is all that thrombin for? *J Thromb Haemost* 2003; **1:** 1504-1514.

76. Wagner MC, Campos-Bilderback SB, Chowdhury M, *et al.* Proximal Tubules Have the Capacity to Regulate Uptake of Albumin. *J Am Soc Nephrol* 2016; **27:** 482-494.

77. Hung G, Xiao X, Peralta R, *et al.* Characterization of target mRNA reduction through in situ RNA hybridization in multiple organ systems following systemic antisense treatment in animals. *Nucleic Acid Ther* 2013; **23:** 369-378.

78. Han X, Nieman MT, Kerlin BA. Protease-activated receptors: An illustrated review. *Res Pract Thromb Haemost* 2021; **5:** 17-26.

79. Lane DA, Philippou H, Huntington JA. Directing thrombin. *Blood* 2005; **106:** 2605-2612.

80. Jesty J. The kinetics of inhibition of alpha-thrombin in human plasma. *J Biol Chem* 1986; **261:** 10313-10318.

81. Ruhl H, Muller J, Harbrecht U, *et al.* Thrombin inhibition profiles in healthy individuals and thrombophilic patients. *Thromb Haemost* 2012; **107:** 848-853.

82. Drake TA, Morrissey JH, Edgington TS. Selective cellular expression of tissue factor in human tissues. Implications for disorders of hemostasis and thrombosis. *Am J Pathol* 1989; **134:** 1087-1097.

83. Pang L, Li Q, Li Y, *et al.* Urine proteomics of primary membranous nephropathy using nanoscale liquid chromatography tandem mass spectrometry analysis. *Clin Proteomics* 2018; **15:** 5.

84. Wang Y, Zheng C, Wang X, *et al.* Proteomic profile-based screening of potential protein biomarkers in the urine of patients with nephrotic syndrome. *Mol Med Rep* 2017; **16:** 6276-6284.

85. Rinschen MM, Godel M, Grahammer F, *et al.* A Multi-layered Quantitative In Vivo Expression Atlas of the Podocyte Unravels Kidney Disease Candidate Genes. *Cell Rep* 2018; **23:** 2495-2508.

86. Apostolopoulos J, Moussa L, Tipping PG. The cytoplasmic domain of tissue factor restricts physiological albuminuria and pathological proteinuria associated with glomerulonephritis in mice. *Nephron Exp Nephrol* 2010; **116:** e72-83.

87. Yamabe H, Yoshikawa S, Ohsawa H, *et al.* Tissue factor production by cultured rat glomerular epithelial cells. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* 1993; **8:** 519-523.

88. Liu N, Ono T, Suyama K, *et al.* Mesangial factor V expression colocalized with fibrin deposition in IgA nephropathy. *Kidney Int* 2000; **58:** 598-606.

89. de Ridder GG, Lundblad RL, Pizzo SV. Actions of thrombin in the interstitium. *J Thromb Haemost* 2016; **14:** 40-47.

90. Chan CW, Chan MW, Liu M, *et al.* Kinetic analysis of a unique direct prothrombinase, fgl2, and identification of a serine residue critical for the prothrombinase activity. *J Immunol* 2002; **168:** 5170-5177.

91. Prunotto M, Farina A, Lane L, *et al.* Proteomic analysis of podocyte exosome-enriched fraction from normal human urine. *J Proteomics* 2013; **82:** 193-229.

92. Boerries M, Grahammer F, Eiselein S, *et al.* Molecular fingerprinting of the podocyte reveals novel gene and protein regulatory networks. *Kidney Int* 2013; **83:** 1052-1064.

93. Pavord S, Myers B. Bleeding and thrombotic complications of kidney disease. *Blood Rev* 2011; **25:** 271-278.

94. Kovacs AL, Seglen PO. Inhibition of hepatocytic protein degradation by methylaminopurines and inhibitors of protein synthesis. *Biochim Biophys Acta* 1981; **676:** 213-220.

95. Derebail VK, Rheault MN, Kerlin BA. Role of direct oral anticoagulants in patients with kidney disease. *Kidney Int* 2020; **97:** 664-675.

96. Htike N, Superdock K, Teehan G, *et al.* Evaluating proteinuria and nephrotic syndrome in patients with venous thromboembolism. *Am J Med Sci* 2012; **343:** 124-126.

97. Kato S, Chernyavsky S, Tokita JE, *et al.* Relationship between proteinuria and venous thromboembolism. *J Thromb Thrombolysis* 2010; **30:** 281-285.

98. Mahmoodi BK, Gansevoort RT, Veeger NJ, *et al.* Microalbuminuria and risk of venous thromboembolism. *JAMA* 2009; **301:** 1790-1797.

Figure 1: Antisense Oligonucleotide-Mediated Hypoprothrombinemia in Healthy Rats. (**A**)Twice weekly treatment with ASO 401027 for 17.5 days reduced plasma prothrombin enzymatic activity in a dose-dependent manner that appeared to plateau at ~60 mg/kg/dose (relative to rat pooled normal plasma prothrombin activity; *n*=2 per dose level). Twice weekly ASO 401027 at 60 mg/kg/dose significantly reduced plasma prothrombin enzymatic activity (**B**) and endogenous thrombin potential (ETP, **C**, **D**) measured at day 27.5 (*n*=3). (**E**) RT-qPCR demonstrated that twice weekly ASO 401027 at 60 mg/kg/dose significantly reduced day 27.5 hepatic prothrombin (*F2*) gene expression (relative to β-actin (*Actb*)) (*n*=3). ***$P<0.001$

Figure 2: Serial Prothrombin Infusion-Mediated Hyperprothrombinemia in Healthy Rats. (**A**) A single dose of intravenous prothrombin (31.25 mg/kg) increased plasma prothrombin enzymatic activity to a peak of ~207% at 1-hour post-dose with an apparent half-life of ~61 hours (relative to rat pooled normal plasma prothrombin activity; *n*=2 per time point). This 31.25 mg/kg loading dose was then followed by maintenance doses of 16.67 mg/kg on days 3, 6, and 9 resulting in significantly increased plasma prothrombin enzymatic activity (**B**) and endogenous thrombin potential (ETP, **C**, **D**) on day 10 (*n*=3). ***P<0.001

Figure 3: Prothrombin Modulation during Rat PAN-NS. On day 10, (**A**) RT-qPCR revealed that ASO 401027 significantly reduced hepatic prothrombin (*F2*) transcript levels (relative to β-actin (*Actb*)) whereas *F2* expression was not significantly altered by PAN-induced proteinuria or prothrombin infusions (*n*=3-12). Plasma prothrombin protein quantity (**B**), enzymatic activity (**C**), and endogenous thrombin potential (ETP, **D**, **E**) were significantly reduced in LoPT rats and significantly increased in HiPT rats in comparison to both control rats and PAN-NS (Sham) rats. *n*=3-12; *P<0.05, **P<0.01, ***P<0.001; ○: Control; ▲: LoPT (ASO-mediated hypoprothrombinemia); ●: Sham; ■: HiPT (prothrombin infusion-mediated hyperprothrombinemia)

Figure 4: Plasma Prothrombin Levels Dictated Thrombin-Podocyte Interactions during Rat PAN-NS. (**A**) Representative immunofluorescence histology of glomeruli on day 10 (original magnification, x60). (**B**) Relative proportion of synaptopodin-positive pixels with colocalized thrombin calculated using the colocalization coefficient feature of ZEN Black software (Zeiss USA). n=3-12 rats per group; each point represents averaged data taken from 20 random glomeruli per rat, *$P<0.05$, **$P<0.01$; ○: Control; ▲: LoPT (ASO-mediated hypoprothrombinemia); ●: Sham; ■: HiPT (prothrombin infusion-mediated hyperprothrombinemia)

Figure 5: Hypoprothrombinemia Diminished *in situ* Podocyte Injury. (**A**) Representative dot plot and subsequent gating strategy used to identify the proportion of synaptopodin-positive podocyte nuclei that were TUNEL-positive, indicating DNA nicking (a marker of terminal podocyte injury). (**B**) Representative histograms of TUNEL-positive podocyte nuclei by treatment group. (**C**) On day 10, LoPT PAN-NS rats had significantly reduced TUNEL-positive podocyte nuclei, in contrast HiPT PAN-NS rats had significantly increased TUNEL-positive podocyte nuclei. *n*=3-12 rats per group; each point represents data derived from 100,000 events per rat, *$P<0.05$, **$P<0.01$, ***$P<0.001$; SSC-A: side scatter-area; FSC-A: forward scatter-area; DAPI-UV: 4′,6-diamidino-2-phenylindole; Synaptopodin-APC: allophycocyanin-labeled anti-synaptopodin antibody; TUNEL-FITC: Fluorescein isothiocyanate-terminal deoxynucleotidyl transferase dUTP nick end labeling; ○: Control; ▲: LoPT (ASO-mediated hypoprothrombinemia); ●: Sham; ■: HiPT (prothrombin infusion-mediated hyperprothrombinemia)

Figure 6: Hyperprothrombinemia Reduced *in situ* Podocyte Survival. (**A**) Representative dot plot and subsequent gating strategy used to count synaptopodin-positive podocytes. (**B**) Representative histograms of podocytes by treatment group. (**C**) On day 10, HiPT rats had significantly reduced podocyte counts in comparison to control and LoPT rats. n=3-12 rats per group; each point represents data derived from 100,000 events per rat; *P<0.05; SSC-A: side scatter-area; FSC-A: forward scatter-area; DAPI-UV: 4′,6-diamidino-2-phenylindole; Synaptopodin-FITC: Fluorescein isothiocyanate-labeled synaptopodin; ○: Control; ▲: LoPT (ASO-mediated hypoprothrombinemia); ●: Sham; ■: HiPT (prothrombin infusion-mediated hyperprothrombinemia)

Figure 7: Prothrombin Regulated Proteinuria and Plasma Albumin during Rat PAN-NS. (A) Day 10 proteinuria was significantly reduced in LoPT rats whereas HiPT rats had significantly more proteinuria than any other group. (**B**) Plasma albumin was significantly improved in LoPT rats. Proteinuria was significantly correlated with plasma prothrombin protein quantity (**C**), enzymatic activity (**D**), and endogenous thrombin potential (ETP, **E**). Proteinuria correlation with podocyte-prothrombin colocalization, TUNEL-positive podocytes, and podocyte counts per glomerulus are shown in panels **F-H**, respectively. *n*=3-12 rats per group; *P<0.05, **P<0.01; ○: Control; ▲: LoPT (ASO-mediated hypoprothrombinemia); ●: Sham; ■: HiPT (prothrombin infusion-mediated hyperprothrombinemia)