Sperm production and allocation in response to risks of sperm competition in the black soldier fly *Hermetia illucens*

Frédéric Manas*, Carole Labrousse, Christophe Bressac

Institut de Recherche sur la Biologie de l'Insecte (IRBI), UMR 7261 CNRS-University of Tours, 37200 Tours, France

*Corresponding author

Correspondence: frederic.manas@univ-tours.fr

Abstract

In polyandrous species, sperm competition between males for the offspring paternity goes on after copulation via the competition of their ejaculates for the fertilisation of female's oocytes. Males allocating more spermatozoa are favored. Different models of sperm competition predict adaptative plasticity of males in the production and allocation of their spermatozoa. These predictions were tested experimentally in the black soldier fly (BSF) *Hermetia illucens*. In this new farmed insect, the adult's biology is little known despite the economic interest of larvae for bioconversion and as an animal feedstuff. Two sets of experiments were carried out to modify the risks of sperm competition perceived by the males. The first consisted of placing adult males in different social contexts (alone or in groups of 10 – i.e. mean risks of sperm competition) and then measuring their sperm production. The second took place at the beginning of the copulation; pairs with males from the two mean risks of sperm competition treatments were transferred to different contexts of immediate risks of sperm competition (empty cages, cages containing 10 males or cages containing 10 females), then the spermathecae of the females were collected in order to count the number of spermatozoa allocated by the males. Virgin males reared in groups of 10 showed more spermatozoa in their seminal vesicles than males alone. Regarding sperm allocation, spermathecae of females in groups of 10 males, as well as those in groups of 10 females, had more spermatozoa than those maintained alone. Copulation duration was not affected by these treatments. Males respond to the treatments by a greater investment in sperm production and transfer. A control of sperm storage by the female is also not excluded. These manipulations of sperm competition risks showed that sperm production and allocation are dependent on social context in BSF. The existence of these mechanisms evidences a plasticity of reproduction under socio-sexual situations. It underlines the importance of studying finely the control of fertility in this farmed insect.

Keywords: Sexual selection, Reproduction, Farming insect, Social context, Copulation, *Spermatheca*

Introduction

The struggle for reproduction is an important selective pressure leading to many evolutionary adaptations which is particularly typified by the competition between males of polyandrous species (Andersson, 1994). Fifty years ago, Parker (1970) theorized that intrasexual competition between males could be expressed both before and after copulation, as it could continue within female reproductive organs, in the form of sperm competition - i.e. 'the competition within a single female between the sperm from two or more males over the fertilization of the ova'.

Many physiological (Pizzari & Parker, 2009; Godwin et al., 2017), morphological (Córdoba-Aguilar et al., 2003) and behavioral (Alcock, 1994; Cueva del Castillo, 2003; Barbosa, 2012) traits have been interpreted in light of this paradigm shift (Parker et al., 1998; Wigby and Chapman, 2004). For example, longer spermatozoa swimming faster, bigger testes producing more spermatozoa or mate-guarding strategies are selected by sperm competition as they maximize male's fertilization success in the competition (Alcock, 1994; LaMunyon & Samuel, 1999; Godwin et al., 2017; Lüpold et al., 2020). Among these traits, plasticity in sperm production (i.e. spermatogenesis), as well as the sperm allocated to each copulation have been the subject of many predictions (Parker, 1970; Parker et al., 1997; Parker & Pizzari, 2010). Based on the costs to males of spermatozoa and seminal fluid contents (Dewsbury, 1982; Olsson et al., 1997), theoretical models predict fitness benefits when males strategically adjust ejaculate expenditure depending on the mating context (Parker & Pizzari, 2010). These models are basically divided into two categories : those based on the "risks" of sperm competition (Parker, 1990; Parker et al., 1997) and those based on the "intensity" of sperm competition (Ball & Parker, 1996, 1997). The risks of sperm competition are defined as the probability that the sperm of a male will compete with the sperm of other males for fertilization of a defined set of ova (Parker, 1990; Parker et al., 1997) whereas the intensity of sperm competition, usually applied to external fertiliser, is the actual number of ejaculates competing for a given set of eggs (Ball & Parker, 1996, 1997).

The predictions of "risks" models have been successfully tested in many organisms, including rodents, fish, and many insects (delBarco-Trillo, 2011). For example, in *Drosophila melanogaster*, sperm production increases when males are housed with other males for a long period of time (Moatt et al., 2014) – increasing what has been named "mean risks of sperm competition" (Engqvist & Reinhold, 2005). Moreover the arrival of rival males during copulation – increasing the "immediate risks of sperm competition" (Engqvist & Reinhold, 2005) - induces focal males to transfer more spermatozoa to the female (Garbaczewska et al., 2013). On the contrary, "intensity" models predictions are more challenging to test and little evidence is available as to the veracity of their predictions (Kelly & Jennions, 2011 but see Sloan et al., 2018).

The quantity of sperm produced or allocated is not the only component of copulation modified according to the context of sperm competition. For instance, the duration of copulation is particularly studied as it can be considered as a proxy for the amount of sperm allocated (Bretman et al., 2009; Barbosa, 2011), although it is not always true (see Weggelaar et al., 2019). Regardless of the sperm allocation, copulation duration is also predicted to vary with sperm competition risks (Alcock, 1994). By copulating longer, males undertake mate guarding, thus preventing the female from remating (Alcock, 1994), which is a widespread behavior in insects (Lorch et al., 1993; Cueva del Castillo, 2003; Barbosa, 2011).

In this study, we aimed to test the predictions of risks of sperm competition models in the black soldier fly *Hermetia illucens* (BSF), a species of great interest for mass-rearing and organic waste bioconversion (Tomberlin & van Huis, 2020). More precisely, we tested wether males adjust their spermatozoa production and allocation in response to the mean and immediate risks of sperm competition Despite its economic interest, studies on adults BSF and their reproductive biology are scarce. Giunti et al., (2018) reported a high prevalence of same-sex sexual behaviors in adults BSF, which can be associated to a high degree of polygyny in other species (MacFarlane et al., 2010). Multiple matings have been reported (Permana et al., 2020; Hoffmann et al., 2021) and morphological traits including complex spermathecae, long and numerous spermatozoa, large testes (Munsch-Masset et al., 2023) strongly suggest post-copulatory sexual selection pressures in this species. Here we experimentally manipulated the risks of sperm competition to examine the phenotypic plasticity in ejaculate expenditure. First, we tested whether

87 long-term exposure to other males could affect sperm production (mean risks of sperm competition) in
88 males' seminal vesicles. Secondly, we assessed if the sudden appearance or disappearance of rivals
89 (immediate risks of sperm competition) coupled with different mean sperm competition treatments could
90 affect the duration of copulation and the amount of sperm stored in the spermathecae of females.

Materials and methods

Rearing conditions

Males and females tested in this study came from a mass rearing maintained in lab conditions since
2018. The strain used is a mix from 2 french commercial lines (Innovafeed and Biomimetic) and one lab line
from Wageningen. Black soldier flies were reared under controlled conditions. Adults – ca 200 to 300 -
were hosted in 50x50x50 cm cages at 24°C and were provided with a cotton ball saturated with water to
maintain moisture. They were exposed to a 12 hours day/night regimen with Philips TLD 36W-84
fluorescent tubes positioned at 10 cm from the cages and providing 2000 to 6000 lux. After collection from
the rearing cage, eggs and larvae were maintained at 27°C, the developing substrate was the Gainesville
diet (Tomberlin & Sheppard, 2002), no additional moisture was added during development. Pupae were
collected and maintained in groups at 24°C with sawdust until emergence. Emerging flies were collected
and sexed daily for experiments. Females were isolated in 15x15x15cm cages in groups of 20 females per
cage. As for males, they were isolated differently depending on the treatment (see below). Both sexes had
access to a cotton ball saturated with water before the experiment. When flies were not used for
experiments, they were transferred to the rearing cage.

Production of spermatozoa

Males of same age (n = 19) were maintained under low risk of sperm competition situation. These
individuals were singled, placed in individual 120 mL plastic containers preventing any visual or physical
contact with other males and limiting olfactory cues. The second treatment consisted in placing ten males
of the same age in a 960 mL plastic container allowing physical, visual, and chemosensory contacts, to
simulate a high risks of sperm competition (n = 24 individuals dissected). Males from both treatments were
kept in these conditions from 5 to 8 days with access to a cotton ball saturated with water.

Allocation of spermatozoa

As BSF will not initiate copulations when only one male and one female are placed in a cage (Personal
observations), the first step for the experiment on the immediate risks of sperm competition involved
transferring 20 virgin males from both treatments (10 singled males and 10 grouped males) to one
15x15x15 cm cage containing 20 virgin females. Males from the singled treatments and from the grouped
treatment were differentiated by marking a different coloured spot on their thorax.
Individuals remained in contact for 5 hours and fourteen observations sessions were performed. The
cage was constantly watched during the contact time to ensure that only first matings for each individual
were recorded.
Once copulations began, each mating pair was gently placed on the lid of a petri dish and transferred
in a cage of similar size containing either no individuals to simulate low immediate risks of sperm
competition (n = 38 individuals composed by n = 23 grouped males and n = 15 singled males), 10 males to
simulate high immediate risks of sperm competition (n = 38 individuals composed by n = 24 grouped males
and n = 14 singled males), or 10 females to test the ability of males to recognise genuine competitors (n =
43 individuals composed by n = 27 grouped males and n = 16 singled males). Time was recorded once
copulations were completed to record duration, and 'high risks' pairs – i.e. pairs that mated in presence of
conspecifics - were removed from the cage immediately after separation to ensure that female did not re-
mate. Pairs were kept together in petri dishes within which there is not enough space for extra copulations
to occur (see Munsch-Masset et al., 2023), until dissection of the female reproductive tract.

Figure 1 - Procedure for the mating experiment to measure the allocation of spermatozoa. Males from two mean risks of sperm competition treatments were transferred to a mating cage. After the begining of mating, the pair was transferred to a different cage according to the immediate risks of sperm competition treatment.

Dissections and collection of data

Since age can affect the number of spermatozoa in seminal vesicles (Munsch-Masset et al., 2023), we dissected males of similar ages (n = 5 males of 5 days, n = 28 males of 6 days and n = 10 males of 8 days after emergence, and we controlled the age in the statistic models, see below). Dissections were performed under a Nikon SMZ745T stereomicroscope (x3.35 magnification) (Nikon, Japan) in PBS saline buffer using fine forceps. For all males, the abdomen was opened after decapitation to collect seminal vesicles which

were then placed on a slide and gently uncoiled with fine forceps. Males were photographed under a Nikon
SMZ745T stereomicroscope (x3.35 magnification) (Nikon, Japan) with a Leica IC 80 HD camera (Leica,
Germany) to measure the head width using ImageJ. This measure can be considered as a reliable proxy of
the size of the individuals (Jones & Tomberlin, 2021). In the same way, the seminal vesicles were
photographed and their whole length was measured with ImageJ. A drop of DAPI was then applied to the
preparation to label the nuclei of the spermatozoa for blind-counting in a section of one of the two seminal
vesicles using a fluorescence microscope (Olympus CX40, Japan) with a x20 objective as in Munsch-Masset
et al., (2023). The length of this section was also measured to obtain the ratio between the sperm-counted-
section and the whole seminal vesicles. Then, this ratio was multiplied to the number of sperm counted
within the portion to obtain the total number of spermatozoa in the seminal vesicles. Finally, this was
doubled for the total number of sperm of one male.
The dissection of females took place the day after copulation, knowing that females lay eggs two/three
days after mating (Munsch-Masset et al., 2023). The two individuals of a pair were photographed under a
Nikon SMZ745T stereomicroscope (x3.35 magnification) (Nikon, Japan) with a Leica IC 80 HD camera (Leica,
Germany) to measure the head width using ImageJ like the males frome the mean risks experiment these
measures and were taken into account in the models for both the sperm production and allocation as
covariates. For all individuals, the abdomen was opened to collect the three spermathecae which were
then placed on a microscope slide. Before crushing them with a microscope coverslip to release the
spermatozoa, a drop of DAPI was applied to the spermatheca to mark the nucleus of the spermatozoa
which were blind-counted under a fluorescence microscope(Olympus CX40, Japan) under a x20 objective.

Statistical analyses
To test our hypotheses, linear mixed models (LMM) were used with the "lmer" function in the "lme4"
package in R (Bates et al., 2015). We included the head size, the size of the seminal vesicles and the
age of the male as covariates and the mean risks of sperm competition treatment as a fixed effect to
explain the number of spermatozoa in the seminal vesicles. For the mating experiment, we included
the head size of the female, the head size of the male, the age of the pair, copulation duration as
covariates and the two risks of sperm competition treatments – mean risks and immediate risks – as
well as their interaction as fixed effects to explain the number of spermatozoa in the spermathecae.
For the latter, we also added the number of individuals in the cage prior to the begining of mating as
it could influence sperm allocation.To study the copulation duration, we used cox proportional hazard
model with the "coxph" function in the "survival package" in R (Therneau, 2019). In the same way,
age, sizes of the male and the female and both sperm competition risks treatments were included as
fixed effects in the model.

The day of sampling was included as a random effect to account for variability inherent to each series
in the sperm count in the models. The fixed effects in our models were tested using the "lmerTest" package
(Kuznetsova et al., 2017), with type III ANOVA F statistics using Satterthwaite approximations for the linear
mix models and with type III ANOVA Chi statistics for the survival model. The assumptions of the linear
mixed model, including normality of residuals, constant variance, and absence of multicollinearity among
the independent variables were checked graphically. As heteroscedasticity in the models was deteted, a
logarithmic transformation was applied on the response variables. We also assessed the proportional
hazards assumption of the cox model using Schoenfeld residuals and found no significant violations of this
assumption.
All statistical analyses were performed using R version 4.0.2 (R Core Team, 2020). The significance level
was set at alpha = 0.05 for all tests. Quantitative data are presented as means ± standard errors (SE) and
hazard ratios (HR) are reported for cox models.

Results

190   On average, 5 hours of contact at this population density – 20 females and 20 males - allowed 7 ± 0.72
191 mating to occur, meaning that 35 % actually mated. Through the different observations, grouped males

192 copulated significantly more than singled males (Fisher's exact test: P = 0.03). In total n = 45 copulations
193 from singled males and n = 74 copulations from grouped males and were observed.

Production of spermatozoa

The number of spermatozoa found in the seminal vesicles of the males was neither related to their size
($F_{1,38}$ = 0.61 ; P = 0.44, β ± SE = -0.21 ± 0.28) nor with their age ($F_{1,38}$ = 0.62 ; P = 0.44, β ± SE = 0.06 ± 0.08)
nor with the length of their seminal vesicles ($F_{1,38}$ = 1.81 ; P = 0.19, β ± SE = 0.07 ± 0.05). However, the
treatment of mean risks of sperm competition showed a significant effect on the number of spermatozoa
in the seminal vesicles of males ($F_{1,38}$ = 7.96 ; P < 0.01 ; full model R^2 = 0.20) (Fig.1). Males kept in groups
had a mean 43 % increase in the number of spermatozoa (mean ± SE : 15578 ± 1105, n = 24) in their seminal
vesicles compared to singled males (mean ± SE : 10920 ± 1200, n = 19).

Figure 2 - The number of spermatozoa in the seminal vesicles of males according to the mean sperm
competition treatment (either the male single or the male within a group of 10 males) Box plots show
median (horizontal bars), upper, and lower quartiles (borders of the box). Whiskers extend from the
10th to the 90th percentiles. Different letters indicate a significant difference at α = 0.05.

Allocation of spermatozoa

The number of spermatozoa found in the female's spermathecae was neither related to the size of the
male ($F_{1,107.7}$ = 0.61 ; P = 0.44, β ± SE = 0.15 ± 0.19), nor with their age ($F_{3, 14.1}$ = 0.07; P = 0.78; β ± SE = -0.02
± 0.09), nor to the copulation duration ($F_{1,104.02}$ = 0.01 ; P = 0.96; β ± SE = 0.01 ± 0.01) (Fig.3). The number
of spermatozoa in the female's spermathecae was marginally and positively correlated with the number of
individuals in the cage before the beginning of mating ($F_{1,104.07}$ = 3.81 ; P = 0.05; β ± SE = 0.01 ± 0.01). Timing
of mating was not different between singled and grouped males (W = 1465, P = 0.27, singled males mean
± SE = 33.6 ± 0.79, grouped males ± SE = 32.32 ± 0.66).
Females that mated with males from the two mean sperm competition risks treatments stored the
same amount of spermatozoa ($F_{1,105.30}$= 0.23 ; P = 0.63, Low treatment males : 4614± 273, High treatment
males : 4709 ± 270) and the interaction bewteen the mean and the immediate risks treatments was not
significant ($F_{2,103.11}$ = 0.08; P = 0.92).
However, the number of spermatozoa found in the female's spermathecae was related to immediate
sperm competition risks treatment ($F_{2,99.56}$ = 6.8; P < 0.01; full model R^2 = 0.43) (Fig.4). Females that mated
in empty cages stored less spermatozoa than the ones that mated in presence of males (β ± SE = -0.38 ±
0.13; t = -2.84; P < 0.01) and the ones that mated in presence of females (β ± SE = -0.43 ± 0.13; t = -3.29; P
< 0.01).There was no significative difference (t = -0.37, P = 0.71) between the content of spermathecae of
females mated with males in the 10 males treatment (mean ± SE : 4943 ± 376, n = 38) and those in the 10
females treatment (mean ± SE : 5554 ± 284, n = 43). Females mated with males in presence of conspecifics
– either males or females - had a mean 60 % increase in the number of spermatozoa (mean ± SE : 5268 ±

233, n = 81 with n = 38 with pairs 10 males and n = 43 pairs with 10 females) compared to the pairs mating
alone (mean ± SE : 3406 ± 268, n = 38).

Figure 3 - The number of spermatozoa in the spermathecae of females according to the copulation
duration. Each point is an individual female, n = 119. The dashed line represents a non significant
relationship between these two variables, linear regression: $R^2 < 0.01$).

Figure 4 - The number of spermatozoa in the spermathecae of females according to the immediate
risk of sperm competition (either the pair mating alone, with 10 males or with 10 females). The colors
represent the mean sperm competition risks treatment (singled males in green and grouped-males
in orange).Box plots show median (horizontal bars), upper, and lower quartiles (borders of the box).
Whiskers extend from the 10th to the 90th percentiles. Different letters indicate a significant
difference at $\alpha = 0.05$.

Copulation duration

241 The mean time of copulation for all pairs was mean ± SE = 35.16 ± 1.19. Immediate sperm competition
242 risks ($\chi^2 = 0.88$; P = 0.64), mean sperm competition risks treatments ($\chi^2 = 0.09$; P = 0.76) and their
243 interaction ($\chi^2 = 3.43$; P = 0.18) showed no effects on copulation duration. In the same way, the size of the
244 female ($\chi^2 = 0.31$; P = 0.58), the size of the male ($\chi^2 = 0.60$; P = 0.44) and age of the pair ($\chi^2 = 1.32$; P =
245 0.25) were not related to copulation duration.

Discussion

BSF Males had more spermatozoa in their seminal vesicles when they were grouped, and females of
pairs that mated in presence of conspecific stored more spermatozoa in their spermathecae. In line with
the predictions of the sperm competition theory, the males of the BSF respond, on one hand, to mean risks
of sperm competition (long-term exposure to rivals) by producing more spermatozoa in their seminal
vesicles and on the other hand, to the immediate risks of sperm competition (sudden exposure to rivals)
by allocating more spermatozoa in a copulation. In contrast, copulation duration was neither related to
sperm competition risks treatments, nor to the number of transferred spermatozoa.
Regardless of sperm competition risks, it has been shown that ejaculate expenditure could be condition
dependent (Perry & Rowe, 2010; Kaldun & Otti, 2016; Wylde et al., 2020), or sometimes associated with
secondary sexual signals (Mautz et al., 2013; Polak et al., 2021). It seems not to be the case in the BSF
where it has already been shown that male size does not affect sperm production (Jones & Tomberlin,
2021; Munsch-Masset et al., 2023). In the same way, we show here that it does not affect sperm allocation.
Interestingly, it appears that males producing more spermatozoa (reared under high risk of sperm
competition) do not transfer more spermatozoa to females. Although it is not the case, one would expect
that the amount of spermatozoa available to males might be partly determinant of the amount allocated
to a copulation (Engqvist & Reinhold, 2005) and finally stored by females. For the latter, it is impossible to
disentangle the role of the male from the role of the female on the storage of spermatozoa.
We cannot exclude the male mating rate hypothesis (Vahed & Parker, 2012) - i.e. the possibility that
the increase in sperm production in group-reared males is due to a higher estimate of the number of mating
opportunities based on density. Such an increase in sperm production could be explained by the risk of
sperm competition or by the number of opportunities for successive mating, although these two
explanations are not mutually exclusive. Furthermore, while allocating the same quantity of spermatozoa
to a unique mating (see results), we observed that grouped males had higher success in mating than singled
males, suggesting that density induce a greater number of copulation attempts. To investigate this
hypothesis further, a different experimental set-up would have to be set up to allow the possibility of
multiple copulations per treatment.
In contrast to some other species like the yellow dung fly *Scatophaga stercoraria* (Simmons et al., 1999)
or the scorpionfly *Panorpa cognata* (Engqvist & Sauer, 2003), copulation duration is not related to the
amount of sperm transferred by the male in BSF. Sperm transfer dynamics that do not follow a linear
relationship with time are not rare (Weggelaar et al., 2019) and it seems to be the case in the black soldier
fly. Here, durations of copulations were not different between the three treatments. Apart from sperm
transfer dynamics, plasticity in copulation duration when males are exposed to rivals can be associated
with active mate guarding (Lorch et al., 1993; Alcock, 1994; Cueva del Castillo, 2003), a behavior that BSF
males do not appear to exhibit (Giunti et al., 2018), as confirmed here. Also, it would appear that once
copulation has begun, surrounding males lose interest in the pair, unlike during courtship when challenger
males may pounce on the pair attempting copulation (Julita et al., 2020).
Numerous cues can be used by males to assess the risk of sperm competition. For example, another
Diptera, *Drosophila melanogaster* uses combinations of cues as diverse as visual, contacts, chemosensory,
and sounds to detect rivals (Bretman et al., 2011). It has been suggested that BSF uses acoustic signals to
identify conspecifics without differentiating females from potential rivals (Giunti et al., 2018), leading
frequent same-sex sexual behaviors. These behaviors are observed with males displaying aedaegus
eversion during courtship (Personal observations, Giunti et al., 2018), which may indicate that males of the
BSF attempt to copulate indifferently with males and females. Same-sex sexual behaviors are common in
species with high density of individuals where females are hard to distinguish (Scharf & Martin, 2013) as in
BSF rearings, but wether these are current in the wild is not known. Interestingly, we found that BSF males
appeared to adjust the number of spermatozoa allocated in a copulation when they were with conspecifics,
regardless of whether these were males or females. Although this effect may be due to an effect of
population density, this sperm adjustment is in line with a potential absence of sex recognition in BSF
males.
Like many aspects of BSF biology, pre-copulatory sexual selection processes in this species are not
precisely known. Sexual dimorphism is reduced and preliminary results indicate that male size does not
play a role in female's mates selection (Unpublished data). BSF was described as using leks to mate

(Tomberlin & Sheppard, 2001). Those structures are defined as aggregated males display sites that females attend primarily for the purpose of fertilization (Höglund & Alatalo, 1995). Supposedly aggressive intrasexual interactions were also observed but females were said to be 'similarly greeted' than males in the supposed lek sites, except that these interactions ended in a mating (Tomberlin & Sheppard, 2001). We did not notice any aggregating area akin to a lek in our rearing conditions (Benelli et al., 2014), furthermore the possible lack of sex recognition may question the hypothesis of the BSF actually being a lekking species.

Previous studies have demonstrated the occurrence of multiple mating in BSF (Permana et al., 2020; Hoffmann et al., 2021). Consistently with sperm competition theory, our findings suggest that males invest more in sperm production and allocation as a strategy to overcome rivals in this competitive reproductive environment. However, a bet hedging strategy is not evidenced here because males copulating in the presence of virgin females do not spare their sperm reserves in the perspective of the insemination of a maximum number of mates. Besides sperm competition, the complexity of female spermathecae in this species (Munsch-Masset et al., 2023) strongly suggests that post-copulatory intersexual selection mechanisms are at work, such as cryptic female choice or sperm precedence (Pascini & Martins, 2017).

BSF is a species of great economic interest in animal production for its potential as a feed source (Tomberlin & van Huis, 2020). Rearing conditions are certainly very different from natural conditions which are still little known. Not all individuals in a breeding mate, and only a proportion of them mate several times (Unpublished data). When it comes to controlling reproduction for example to make genetic selection, the understanding of sexual selection processes is a key factor that need to be integrated in the rearings – e.g. by determining if different contexts of competition are good for productivity.

Acknowledgments

We thank Elisabeth Herniou the proofreading of the paper. We thank Hélène Girotvergne for technical assistance.

Funding

FM was funded by the Doctoral School 'Santé, Sciences Biologiques et Chimie du Vivant'. This work is part of the BioSexFly program funded by the Centre Val de Loire region.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

Analyses reported in this article can be reproduced using the data and script provided by Frédéric Manas (2023) (https://zenodo.org/records/10078561). https://doi.org/10.5281/zenodo.10078561

References

Alcock J (1994) Postinsemination Associations Between Males and Females in Insects: The Mate-Guarding Hypothesis. *Annual Review of Entomology*, **39**, 1–21. https://doi.org/10.1146/annurev.en.39.010194.000245

Andersson M (1994) *Sexual selection*. Princeton University Press.

Ball MA, Parker GA (1996) Sperm Competition Games: External Fertilization and "Adapative" Infertility. *Journal of Theoretical Biology*, **180**, 141–150. https://doi.org/10.1006/jtbi.1996.0090

Ball MA, Parker GA (1997) Sperm Competition Games: Inter- and Intra-species Results of a Continuous External Fertilization Model. *Journal of Theoretical Biology*, **186**, 459–466. https://doi.org/10.1006/jtbi.1997.0406

Barbosa F (2011) Copulation duration in the soldier fly: the roles of cryptic male choice and sperm
competition risk. *Behavioral Ecology*, **22**, 1332–1336. https://doi.org/10.1093/beheco/arr137
Barbosa F (2012) Males responding to sperm competition cues have higher fertilization success in a soldier
fly. *Behavioral Ecology*, **23**, 815–819. https://doi.org/10.1093/beheco/ars035
Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models using lme4. *Journal of*
Statistical Software, **67**, 1–48. https:// doi.org/10.18637/jss.v067.i01
Benelli G, Daane KM, Canale A, Niu CY, Messing RH, Vargas RI (2014) Sexual communication and related
behaviours in Tephritidae: current knowledge and potential applications for Integrated Pest
Management. *Journal of Pest Science*, **87**, 385–405. https://doi.org/10.1007/s10340-014-0577-3
Bretman A, Fricke C, Chapman T (2009) Plastic responses of male Drosophila melanogaster to the level of
sperm competition increase male reproductive fitness. *Proceedings of the Royal Society B: Biological*
Sciences, **276**, 1705–1711. https://doi.org/10.1098/rspb.2008.1878
Bretman A, Westmancoat JD, Gage MJG, Chapman T (2011) Males use multiple, redundant cues to detect
mating rivals. *Current Biology*, **21**, 617–622. https://doi.org/10.1016/j.cub.2011.03.008
Córdoba-Aguilar A, Uhía E, Rivera AC (2003) Sperm competition in Odonata (Insecta): the evolution of
female sperm storage and rivals' sperm displacement. *Journal of Zoology*, **261**, 381–398.
https://doi.org/10.1017/S0952836903004357
Cueva del Castillo R (2003) Body size and multiple copulations in a neotropical grasshopper with an
extraordinary mate-guarding duration. *Journal of Insect Behavior*, **16**, 503–522.
https://doi.org/10.1023/A:1027303323242
Dewsbury DA (1982) Ejaculate cost and male choice. *The American Naturalist*, **119**, 601–610.
https://doi.org/10.1086/283938
Engqvist L, Reinhold K (2005) Pitfalls in experiments testing predictions from sperm competition theory.
Journal of Evolutionary Biology, **18**, 116–123. https://doi.org/10.1111/j.1420-9101.2004.00792.x
Engqvist L, Sauer KP (2003) Determinants of sperm transfer in the scorpionfly Panorpa cognata: male
variation, female condition and copulation duration. *Journal of Evolutionary Biology*, **16**, 1196–1204.
https://doi.org/10.1046/j.1420-9101.2003.00613.x
Garbaczewska M, Billeter JC, Levine JD (2013) Drosophila melanogaster males increase the number of
sperm in their ejaculate when perceiving rival males. *Journal of Insect Physiology*, **59**, 306–310.
https://doi.org/10.1016/j.jinsphys.2012.08.016
Giunti G, Campolo O, Laudani F, Palmeri V (2018) Male courtship behaviour and potential for female mate
choice in the black soldier fly Hermetia illucens L. (Diptera: Stratiomyidae). *Entomologia Generalis*, **38**,
29–46. https://doi.org/10.1127/entomologia/2018/0657
Godwin JL, Vasudeva R, Michalczyk Ł, Martin OY, Lumley AJ, Chapman T, Gage MJG (2017) Experimental
evolution reveals that sperm competition intensity selects for longer, more costly sperm. *Evolution*
Letters, **1**, 102–113. https://doi.org/10.1002/evl3.13
Hoffmann L, Hull KL, Bierman A, Badenhorst R, Bester-van der Merwe AE, Rhode C (2021) Patterns of
genetic diversity and mating systems in a mass-reared black soldier fly colony. *Insects*, **12**, 480.
https://doi.org/10.3390/insects12060480
Höglund J, Alatalo RV (1995) *Leks:* Princeton University Press.
Jones BM, Tomberlin JK (2021) Effects of adult body size on mating success of the black soldier fly, Hermetia
illucens (L.) (Diptera: Stratiomyidae). *Journal of Insects as Food and Feed*, **7**, 5–20.
https://doi.org/10.3920/JIFF2020.0001
Julita U, Lusianti Fitri L, Eka Putra R, Dana Permana A (2020) Mating success and reproductive behavior of
black soldier fly Hermetia illucens l. (Diptera, Stratiomyidae) in tropics. *Journal of Entomology*, **17**, 117–
127. https://doi.org/10.3923/je.2020.117.127
Kaldun B, Otti O (2016) Condition-dependent ejaculate production affects male mating behavior in the
common bedbug *Cimex lectularius*. *Ecology and Evolution*, **6**, 2548–2558.
https://doi.org/10.1002/ece3.2073
Kelly CD, Jennions MD (2011) Sexual selection and sperm quantity: meta-analyses of strategic ejaculation.
Biological Reviews, **86**, 863–884. https://doi.org/10.1111/j.1469-185X.2011.00175.x
Kuznetsova A, Brockhoff PB, Christensen RHB (2017) **lmerTest** Package: Tests in Linear Mixed Effects
Models. *Journal of Statistical Software*, **82**. https://doi.org/10.18637/jss.v082.i13

LaMunyon CW, Samuel W (1999) Evolution of sperm size in nematodes: sperm competition favours larger
sperm. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, **266**, 263–267.
https://doi.org/10.1098/rspb.1999.0631

Lorch Patrick D, Wilkinson GeraldS , Reillo Paul R (1993) Copulation duration and sperm precedence in the
stalk-eyed fly Cyrtodiopsis whitei (Diptera : Diopsidae). *Behavioral Ecology and Sociobiology*, **32**.
https://doi.org/10.1007/BF00183785

Lüpold S, de Boer RA, Evans JP, Tomkins JL, Fitzpatrick JL (2020) How sperm competition shapes the
evolution of testes and sperm: a meta-analysis. *Philosophical Transactions of the Royal Society B:*
Biological Sciences, **375**, 20200064. https://doi.org/10.1098/rstb.2020.0064

MacFarlane GR, Blomberg SP, Vasey PL (2010) Homosexual behaviour in birds: frequency of expression is
related to parental care disparity between the sexes. *Animal Behaviour*, **80**, 375–390.
https://doi.org/10.1016/j.anbehav.2010.05.009

Mautz BS, Møller AP, Jennions MD (2013) Do male secondary sexual characters signal ejaculate quality? A
meta-analysis. *Biological Reviews*, **88**, 669–682. https://doi.org/10.1111/brv.12022

Moatt JP, Dytham C, Thom MD (2014) Sperm production responds to perceived sperm competition risk in
male Drosophila melanogaster. *Physiology & Behavior*, **131**, 111–114.
https://doi.org/10.1016/j.physbeh.2014.04.027

Munsch-Masset P, Labrousse C, Beaugeard L, Bressac C (2023) The reproductive tract of the black soldier
fly (Hermetia illucens) is highly differentiated and suggests adaptations to sexual selection. *Entomologia*
Experimentalis et Applicata. https://doi.org/10.1111/eea.13358

Olsson M, Madsen T, Shine R (1997) Is sperm really so cheap? Costs of reproduction in male adders, *Vipera*
berus. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, **264**, 455–459.
https://doi.org/10.1098/rspb.1997.0065

Parker GA (1970) Sperm competition and its evolutionary consequences in the insects. *Biological Reviews*,
45, 525–567. https://doi.org/10.1111/j.1469-185X.1970.tb01176.x

Parker GA (1990) Sperm Competition Games: Raffles and Roles. *Proceedings: Biological Sciences*, **242**, 120–
126. https://doi.org/10.1098/rspb.1990.0114

Parker GA, Ball MA, Stockley P, Gage MJG (1997) Sperm competition games: a prospective analysis of risk
assessment. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, **264**, 1793–1802.
https://doi.org/10.1098/rspb.1997.0249

Parker GA, Birkhead TR, Møller AP (1998) Sperm competition and sexual selection. *Sperm competition and*
the evolution of ejaculates: towards a theory base, 3–54.

Parker GA, Pizzari T (2010) Sperm competition and ejaculate economics. *Biological Reviews*, **85**, 897–934.
https://doi.org/10.1111/j.1469-185X.2010.00140.x

Pascini TV, Martins GF (2017) The insect spermatheca: an overview. *Zoology*, **121**, 56–71.
https://doi.org/10.1016/j.zool.2016.12.001

Permana AD, Fitri LL, Julita U (2020) Influence of mates virginity on black soldier fly, Hermetia illucens L.
(Diptera: stratiomyidae) mating performance. *Jurnal Biodjati*, **5**, 174–181.
https://doi.org/10.15575/biodjati.v5i2.9049

Perry JC, Rowe L (2010) Condition-dependent ejaculate size and composition in a ladybird beetle.
Proceedings of the Royal Society B: Biological Sciences, **277**, 3639–3647.
https://doi.org/10.1098/rspb.2010.0810

Pizzari T, Parker GA (2009) Sperm competition and sperm phenotype. In: *Sperm Biology* , pp. 207–245.
Elsevier. https://doi.org/10.1016/B978-0-12-372568-4.00006-9

Polak M, Hurtado-Gonzales JL, Benoit JB, Hooker KJ, Tyler F (2021) Positive genetic covariance between
male sexual ornamentation and fertilizing capacity. *Current Biology*, **31**, 1547-1554.e5.
https://doi.org/10.1016/j.cub.2021.01.046

Scharf I, Martin OY (2013) Same-sex sexual behavior in insects and arachnids: prevalence, causes, and
consequences. *Behavioral Ecology and Sociobiology*, **67**, 1719–1730. https://doi.org/10.1007/s00265-
013-1610-x

Simmons LW, Parker GA, Stockley P (1999) Sperm Displacement in the Yellow Dung Fly, Scatophaga
stercoraria: An Investigation of Male and Female Processes. *The American Naturalist*, **153**, 302–314.
https://doi.org/10.1086/303171

Sloan NS, Lovegrove M, Simmons LW (2018) Social manipulation of sperm competition intensity reduces
seminal fluid gene expression. *Biology Letters*, **14**, 20170659. https://doi.org/10.1098/rsbl.2017.0659
Therneau T (2019) The survival package. https://github.com/therneau/survival
Tomberlin JK, Van Huis A (2020) Black soldier fly from pest to 'crown jewel' of the insects as feed industry:
an historical perspective. *Journal of Insects as Food and Feed*, **6**, 1–4.
https://doi.org/10.3920/JIFF2020.0003
Tomberlin JK, Sheppard DC (2001) Lekking behavior of the black soldier fly (Diptera: Stratiomyidae). *The
Florida Entomologist*, **84**, 729. https://doi.org/10.2307/3496413
Tomberlin JK, Sheppard DC (2002) Factors influencing mating and oviposition of black soldier flies (Diptera:
Stratiomyidae) in a colony. *Journal of Entomological Science*, **37**, 345–352.
https://doi.org/10.18474/0749-8004-37.4.345
delBarco-Trillo J (2011) Adjustment of sperm allocation under high risk of sperm competition across taxa:
a meta-analysis. *Journal of Evolutionary Biology*, **24**, 1706–1714. https://doi.org/10.1111/j.1420-
9101.2011.02293.x
Vahed K, Parker DJ (2012) The evolution of large testes: sperm competition or male mating rate? *Ethology*,
118, 107–117. https://doi.org/10.1111/j.1439-0310.2011.01991.x
Weggelaar TA, Commandeur D, Koene JM (2019) Increased copulation duration does not necessarily reflect
a proportional increase in the number of transferred spermatozoa. *Animal Biology*, **69**, 95–115.
https://doi.org/10.1163/15707563-00001078
Wigby S, Chapman T (2004) Sperm competition. *Current Biology*, **14**, R100–R103.
https://doi.org/10.1016/j.ympev.2004.01.013
Wylde Z, Crean A, Bonduriansky R (2020) Effects of condition and sperm competition risk on sperm
allocation and storage in neriid flies. *Behavioral Ecology*, **31**, 202–212.
https://doi.org/10.1093/beheco/arz178
471

Mean risks of sperm competition

Plasticity in production of spermatozoa

Mating cage

Immediate risks of sperm competition

Plasticity in allocation of spermatozoa