1

_N_-6-methyladenosine (m6A) Promotes the Nuclear Retention of mRNAs with Intact 5′

Splice Site Motifs

Eliza S. Lee[1,6], Harrison. W. Smith[1], Yifan E. Wang[1], Sean S. J. Ihn[1], Leticia Scalize de Olivera[1],

Nevraj S. Kejiou[1], Yijing L. Liang[2], Syed Nabeel-Shah[3,4], Robert Y. Jomphe[1,5], Shuye Pu[4], Jack

F. Greenblatt[3,4], Alexander F. Palazzo[1,7]

1- University of Toronto, Department of Biochemistry, Canada

2- Centre for Computational Medicine, Hospital for Sick Children, Toronto, Canada

3- University of Toronto, Department of Molecular Genetics, Canada

4- Terrence Donnelly Centre for Cellular & Biomolecular Research, Toronto, Canada

5- Cell Biology Program, Hospital for Sick Children, Toronto, Canada

6- Present Address: Department of Biochemistry, University of Colorado Boulder, CO

7- Corresponding author

Abstract

In humans, misprocessed mRNAs containing intact 5′ Splice Site (5′SS) motifs are nuclear retained and targeted for decay by ZFC3H1, a component of the Poly(A) Exosome Targeting complex, and U1-70K, a component of the U1 snRNP. In *S. pombe*, the ZFC3H1 homolog, Red1, binds to the YTH-domain containing protein Mmi1 to target certain RNA transcripts to nuclear foci for nuclear retention and decay. Here we show that YTHDC1 and YTHDC2, two YTH domain-containing proteins that bind to *N*-6-methyladenosine (m6A) modified RNAs, interact with ZFC3H1 and U1-70K, and are required for the nuclear retention of mRNAs with intact 5′SS motifs. Disruption of m6A deposition inhibits both the nuclear retention of these transcripts and their accumulation in YTHDC1-enriched foci that are adjacent to nuclear speckles. Endogenous RNAs with intact 5′SS motifs, such as intronic polyadenylated transcripts, tend to be m6A-modified at low levels. Thus, m6A modification acts in a conserved quality control mechanism that targets misprocessed mRNAs for nuclear retention and decay.

Introduction

Eukaryotic cells are divided into two compartments: the nucleus where pre-mRNAs are synthesized and processed, and the cytoplasm where mature mRNAs are translated into proteins. This subcellular division allows for extensive quality control at the level of both RNA synthesis and RNA processing to promote the nuclear retention and degradation of spurious transcripts and misprocessed mRNAs before they access the translation machinery in the cytoplasm (Palazzo and Lee 2018; Garland and Jensen 2020; Palazzo et al. 2024). Misprocessed mRNAs typically contain intact RNA processing *cis*-elements which promote retention and decay (Legrain and Rosbash 1989; Abad et al. 2008; Lee et al. 2015).

One class of misprocessed mRNAs are intronic polyadenylated (IPA) transcripts (Tian et al. 2007). These are generated by cryptic 3′cleavage/polyadenylation signals found in introns. Although most of these cryptic 3′cleavage/polyadenylation signals are suppressed (Kaida et al. 2010), they can be used at low levels to generate IPA transcripts that tend to encode protein fragments (Ogami et al. 2017), which are often toxic as they can compete with endogenous proteins for their binding partners (Veitia 2007). IPA transcripts contain intact 5' splice site (5′SS) motifs, which are present at the boundary between the upstream exon and the intron where the 3′ cleavage occurs. We and others have shown that the presence of an intact 5′SS motif in an RNA promotes its nuclear retention and decay (Legrain and Rosbash 1989; Abad et al. 2008; Lee et al. 2015). These transcripts are recognized by the U1 snRNP, and the U1 component U1-70K is required for their nuclear retention and decay (Fortes et al. 2003; Abad et al. 2008; Goraczniak et al. 2009; Lee et al. 2022). We previously found that U1-70K associates with ZFC3H1 (Lee et al. 2022), a component of the Poly(A) Exosome Targeting (PAXT) complex. Moreover, we and others have shown that ZFC3H1 is required for the nuclear retention and decay of RNAs with

intact 5′SS motifs including IPA transcripts (Ogami et al. 2017; Lee et al. 2022). PAXT also

contains the nuclear poly(A) binding protein, PABPN1, and the RNA helicase MTR4, which is

known to feed RNAs into the nuclear exosome, the major RNase complex in human cells (Meola

et al. 2016; Ogami et al. 2017; Ogami and Manley 2017; Silla et al. 2018, 2020). Although

MTR4 is not required for nuclear retention, PABPN1 likely plays a role (Lee et al. 2022).

The PAXT complex is conserved between humans and *S. pombe*, where it is known as the

MTREC (<u>Mt</u>l1-<u>Re</u>d1 <u>C</u>ore) complex, and is composed of Mtl1, the paralogue of Mtr4, Red1, the

paralogue of ZFC3H1 and Pab2, a poly(A) binding protein (Sugiyama and Sugioka-Sugiyama

2011; Zhou et al. 2015; Shichino et al. 2020). MTREC promotes the nuclear retention and decay

of meiotic mRNAs that are inappropriately synthesized in interphase cells. MTREC is recruited

to these RNAs through a complex of three proteins consisting of Erh1, Pir1 and Mmi1

(Sugiyama and Sugioka-Sugiyama 2011; Sugiyama et al. 2016; Wei et al. 2021; Shichino et al.

2020). Note that Pir1 has a Serine/Proline-rich region that shows a limited homology with the N-

terminus of ZFC3H1 (Wei et al. 2021) and that Mmi1 contains a YTH domain, which in other

systems recognizes *N*-6-methyladenosine (m6A) modified RNAs. In *S. pombe*, no m6A

modification is detectable in mRNAs (Ishigami et al. 2021) and instead Mmi1 binds to specific

RNA motifs (Chen et al. 2011; Sugiyama and Sugioka-Sugiyama 2011; Wang et al. 2016).

MTREC and Mmi1 are also involved in the nuclear retention and decay of non-coding RNAs

(ncRNAs) and transposable element-derived RNAs (Sugiyama et al. 2016; Vo et al. 2019; Zhou

et al. 2015). When RNA decay is inhibited, MTREC-targeted RNAs accumulate in a

Mmi1/Red1-enriched nuclear foci (Sugiyama and Sugioka-Sugiyama 2011; Shichino et al. 2018,

2020).

In humans, YTH domains recognize m6A modified RNAs (Xu et al. 2014). This modification is enriched in several regions of mRNAs, including large exons, regions surrounding the stop codon, and 3′UTRs (Dominissini et al. 2012; Meyer et al. 2012), and is depleted around exon-exon junctions (Yang et al. 2022; Uzonyi et al. 2023; He et al. 2023; Luo et al. 2023). The m6A modification is catalyzed by the METTL3 methyltransferase, which forms a complex with METTL14 and other factors (Liu et al. 2014). This modification can be removed by the demethylases ALKBH5, FTO and perhaps ALKBH3 (Jia et al. 2011; Zheng et al. 2013; Ueda et al. 2017). m6A is predominantly recognized by five YTH domain containing proteins in humans, including YTHDC1 and YTHDC2, which are localized to the nucleoplasm (Xu et al. 2021).

Initial studies indicated that the m6A modification, and m6A-binding proteins promote mRNA export (Roundtree et al. 2017; Zheng et al. 2013; Lesbirel et al. 2018; Lesbirel and Wilson 2019); however, more recent work has found that the levels of m6A modification correlate with nuclear retention (Tang et al. 2024). Moreover, m6A and YTHDC1 are required for the suppression of RNAs generated from transposable elements, enhancer RNAs and other ncRNAs (Liu et al. 2020), reminiscent of the function of MTREC and Mmi1 in *S. pombe*. Like Mmi1 and Red1, YTHDC1 and ZFC3H1 are found in nuclear foci that contain mRNA (Cheng et al. 2021; Nayler et al. 2000; Wang et al. 2021). These foci partially overlap nuclear speckles, which are required for the nuclear retention of mRNA with intact 5′SS motifs (Lee et al. 2022).

Here we show that m6A modification is required for the efficient nuclear retention of mRNAs that contain intact 5′SS motifs, which are found in IPA transcripts, and retains these in nuclear foci present adjacent to nuclear speckles.

Results

YTHDC1 and YTHDC2 interact with both ZFC3H1 and U1-70K

In *S. pombe*, the MTREC associates with Mmi1, a nuclear YTH domain-containing protein (Figure 1A, left) (Sugiyama and Sugioka-Sugiyama 2011; Chen et al. 2011; Zhou et al. 2015; Shichino et al. 2020). To determine whether components of the equivalent human complex (known as PAXT, Figure 1A right) interact with nuclear YTH domain-containing proteins, we expressed either FLAG-HA-YTHDC1 or FLAG-ZFC3H1-HA in HEK293 cells and assessed their binding partners by co-immunoprecipitation in the absence or presence of RNase, which removed all detectable RNA from the lysate (Figure S1A). In FLAG-HA-YTHDC1 immunoprecipitates, we detected both endogenous ZFC3H1 and U1-70K, even in the presence of RNase (Figure 1B, S1B). FLAG-HA-YTHDC1 immunoprecipitates also contained YTHDC2 but not any detectable ALKBH5 (Figure 1B, S1B). In FLAG-ZFC3H1-HA immunoprecipitates, we detected endogenous YTHDC1, YTHDC2 and U1-70K, even in the presence of RNase, but not eIF4AIII or G3BP2 (Figure 1C-D, S1C). In all cases, the plasmid-expressed proteins were present at levels that were less than 10 fold the endogenous protein level and had only minor effects on the levels of their interacting partners (Figure S1D-J). When we immunoprecipitated a smaller fragment of FLAG-ZFC3H1 (FLAG-ZFC3H1 1-1233), which lacks the C-terminal TPR repeats but contains all known protein interaction-domains (Wang et al. 2021), we again detected YTHDC1, YTHDC2 and U170K in the precipitates, but not ALKBH5, eIF4AIII, METTL3 or ALY (Figure 1E-F).

These results indicate that nuclear YTH proteins associate with ZFC3H1 and U1-70K, two factors required for the nuclear retention of RNAs that contain intact 5′SS motifs.

DRACH motifs are required for the efficient nuclear retention of mRNAs with an intact 5′SS motif

The m6A methyltransferase METTL3 preferentially modifies RNAs at consensus DRACH motifs (G/A/U, A/G, A, C, A/U/C) (Meyer et al. 2012; Dominissini et al. 2012). To test whether m6A was required for nuclear retention, we mutated the DRACH motifs in the *fushi-tarazu* (*ftz*) reporter which we previously used to study the effect of the 5′SS motif on nuclear retention (Lee et al. 2015). For the parental intronless *ftz* reporter (*ftz-Δi*) we identified ten putative DRACH motifs and for the 5′SS motif containing version (*ftz-Δi-5′SS*) we found eleven motifs (Figure 2A). We mutated all but of one of the DRACH sites to generate two other reporters, *no-m6A-ftz-Δi* and *no-m6A-ftz-Δi-5′SS* respectively. Note that one DRACH motif, present near the consensus 5′SS motif, was not altered. When the *ftz-Δi-5′SS* reporter was expressed in human osteosarcoma cell lines (U2OS), elimination of most DRACH motifs drastically reduced m6A deposition (Figure 2B). When the distribution of the mRNAs produced from these various reporters was assessed by RNA fluorescent in situ (FISH) hybridization, we found that although the presence of an intact 5′SS motif promoted nuclear retention, elimination of most DRACH motifs inhibited this retention (Figure 2C-D). Elimination of the DRACH motifs in *ftz-Δi* had no effect on its export. When we expressed versions of *ftz-Δi* and *ftz-Δi-5′SS* mRNA with optimized DRACH motifs (AGACT) to enhance methylation (creating *e-m6A-ftz-Δi* and *e-m6A-ftz-Δi-5′SS*; Figure 2A), we did not observe a significant change in their distribution (Figure 2D).

The m6A methyltransferase, METTL3, is required for the nuclear retention of mRNAs with an intact 5′SS motif

Next, we tested whether the small molecule STM2457, which sterically blocks the S-adenosylmethionine-binding site of METTL3 and inhibits its RNA methylation activity (Yankova et al. 2021), also inhibited nuclear retention of mRNA with intact 5′SS motifs. When U2OS cells were treated with 100 μM STM2457 for 18 hours, *ftz-Δi-5′SS* mRNA was no longer nuclear retained when compared to DMSO-treated cells (Figure 2E-F). Note that STM2457-treatment had no effect on the nuclear export of *ftz-Δi* mRNA (Figure 2E-F) and minimal effects on the levels of ZFC3H1, YTHDC1, YTHDC2 or U1-70K (Figure S2A-B).

To further confirm this result, we depleted METTL3 using a combination of four lentiviral-delivered shRNAs (Figure 3G). We found that in METTL3-depleted cells, *ftz-Δi-5′SS* mRNA was more cytoplasmic than in cells treated with control shRNA (Figure 3H-I). Again, the nuclear export of *ftz-Δi* mRNA was not affected.

In aggregate, these results demonstrate that the m6A modification is required for the nuclear retention of reporter mRNAs containing intact 5′SS motifs.

YTHDC1 and YTHDC2 are required for the nuclear retention of 5′SS motif containing reporter mRNAs

To test whether YTHDC1 and YTHDC2 are required for the nuclear retention of mRNAs with intact 5′SS motifs, we co-depleted these two proteins (Figure 3A) using two lentivirus-delivered shRNAs for each gene. In parallel, we depleted ZFC3H1, which is required for the

retention of mRNAs with intact 5′SS motifs (Lee et al. 2022), and ALKBH5, an m6A

demethylase (results described in the next section; Figure 3A).

We observed that in YTHDC1/2-co-depleted cells, *ftz-Δi-5′SS* mRNA was no longer

efficiently retained in the nucleus when compared to control shRNA-treated cells (Figure 3B-C).

The degree to which nuclear retention was inhibited was similar to ZFC3H1-depleted cells

(Figure 3B-C). Both YTHDC1/2-co-depletion and ZFC3H1-depletion had no effect on the

nuclear export of our reporter mRNAs that lacked either the 5′SS motif (*ftz-Δi*) or most DRACH

motifs (*no-m6A-ftz-Δi-5′SS*; Figure 3C). Note that YTHDC1/2-depletion had some effect on the

levels of ZFC3H1, and vice versa (Figure S3A-B).

Next, to validate our results with more natural IPA-generating genes, we monitored IPA

transcripts generated from two mini genes, *PCF11-IPA* and *CCDC71-IPA* (Figure 3D-F, S3C-E),

that we had previously characterized (Lee et al. 2022). Both reporters are based on endogenous

IPA transcripts and require both ZFC3H1 and U1-70K for their efficient nuclear retention. These

IPA transcripts can then be specifically monitored by FISH using fluorescent probes that

hybridize to the intronic region, just upstream of the 3′ cleavage/polyadenylation site (Figure 3D,

S3C). In agreement with our previous results, the two reporters were more cytoplasmic in cells

co-depleted of YTHDC1 and YTHDC2 when compared to cells treated with control shRNA

(Figure 3E-F, S3D-E).

To examine the effects on endogenous IPA transcripts, we fractionated U2OS cells

(Figure S3F) that were treated with various shRNAs and assessed the levels of endogenously

generated *PCF11* IPA transcripts by qRT-PCR. Depletion of either ZFC3H1 or YTHDC1/2

caused an increase in the total levels and in the cytoplasmic/nuclear ratios of endogenous *PCF11*

IPA transcripts (Figure 3G-H), although depletion of the former had a greater effect than

depletion of the latter.

Collectively, these results suggest that YTHDC1 and YTHDC2 are required for the

efficient nuclear retention of mRNAs with intact 5′SS motifs.

Depletion of the m6A demethylase ALKBH5 inhibits nuclear retention of mRNAs with

intact 5′SS motifs

We reasoned that depletion of enzymes that remove the m6A modification should

enhance the nuclear retention of transcripts with intact 5′SS motifs. To test this, we depleted

ALKBH5 in U2OS cells using lentiviral delivered shRNAs (Figure 3A). Note that due to the

nature of our surprising results (see below) we used two different shRNAs to ensure that our

findings were not due to off target effects.

Unexpectedly, we found that depletion of ALKBH5 inhibited the nuclear retention of *ftz-*

Δi-5′SS mRNA (Figure 3B-C), *PCF11-IPA* (Figure 3E-F) and to a lesser extent *CCDC71-IPA*

(Figure S3D-E). Depletion of ALKBH5 had no effect on the nuclear export of our reporter

mRNAs that lacked either the 5′SS motif (*ftz-Δi*) or DRACH motifs (*no-m6A-ftz-Δi-5′SS*) (Figure

3B-C). Consistent with previous reports (Zheng et al. 2013), we found that ALKBH5 was present

in the nuclear fraction, although it was also present in the cytoplasm (Figure S3G).

From these results, we concluded that ALKBH5 was required for the efficient nuclear

retention of most mRNAs that contained intact 5′SS motifs. Although this result seems counter to

our other findings, it may indicate that ALKBH5 may play additional roles beyond removing

m6A modifications (see discussion).

m6A accumulates in IPA transcripts

We next asked whether m6A methylation marks accumulate within natural IPA transcripts. We analyzed CLIP-seq experiments which used an m6A-specific antibody to enrich for modified RNAs isolated from HEK293 cells. We compared the distribution of these m6A methylation marks in intronic regions that have been previously shown to generate IPA transcripts which are sensitive to ZFC3H1-depletion (Lee et al. 2022). As a comparison, each IPA transcript was randomly matched with an exon that was both length- and expression-matched as determined by high throughput sequencing. This matching process was repeated five times across two separate transcriptome sequencing experiments. Although we observed that the average number of m6A modifications per RNA was similar between the IPA transcripts and the length/expression-matched exons (Figure 4A), significantly fewer IPA transcripts lacked detectable m6A modifications (Figure 4B) and significantly more IPA transcripts had a low number of detectable m6A modifications (1-6 sites) when compared to length/expression-matched exons. In contrast, significantly fewer IPA transcripts had a high number of m6A modifications (>12 sites) when compared to length/expression-matched exons (Figure 4B).

When the density of m6A modifications was averaged along the length of the intronic portion of IPA transcripts, we found that these m6A marks were slightly enriched near the start of the IPA (i.e., near the 5′SS motif) and near the 3′cleavage/polyadenylation site, when compared to the density of m6A averaged along all exons, where m6A was depleted at both ends near exon-exon boundaries (Figure 4C) as seen by other groups (Yang et al. 2022; Uzonyi et al. 2023; He et al. 2023; Luo et al. 2023). When we looked more closely at three example genes that generate IPA transcripts, *PCF11*, *CSNK1E*, and *NCOA5,* we found that m6A marks were present

throughout the intronic region that generates the IPA transcript (Figure 4D-F). In some cases, as

in the *NCOA5* IPA transcript, there were only a few m6A marks, but these tended to be near one

of the ends of the IPA transcript and also present in the upstream exon. Note that the level of all

these IPA transcripts increase in ZFC3H1-depleted U2OS cells (Figure 4D-F) as previously

reported (Lee et al. 2022).

mRNAs with intact 5′SS motifs accumulate in nuclear foci containing YTHDC1 and this

requires DRACH motifs

Previously, we found that 5′SS motif containing mRNAs are targeted and then retained in

nuclear foci and that their nuclear retention requires nuclear speckles (Lee et al. 2015, 2022). We

also found that ZFC3H1 and U1-70K are dispensable for the targeting of these mRNAs to

nuclear speckles but instead prevent these RNAs from exiting nuclear foci (Lee et al. 2022).

With this in mind, we asked if m6A modifications affect the trafficking of mRNAs in and out of

nuclear foci.

To determine the localization of mRNAs at early time points post-synthesis, plasmids

containing *ftz-Δi-5′SS* and *no-m6A-ftz-Δi-5′SS* reporter genes were microinjected into the nuclei

of U2OS cells to allow for rapid synchronous expression. After allowing for one hour of mRNA

expression, a time point when nuclear speckle targeting is highest (Akef et al. 2013), cells were

fixed and immunostained for SC35, a nuclear speckle marker (Spector et al. 1991) and probed

for *ftz* mRNA by FISH. We observed robust nuclear speckle targeting of both constructs (Figure

5A-B). This result indicated that within the context of *ftz-Δi-5′SS*, m6A was not required for

nuclear speckle targeting.

Next, we assessed whether m6A affected the trafficking of mRNAs with intact 5′SS

motifs out of nuclear speckles. We thus monitored the mRNA distribution 24h after transfection

of the plasmids with reporter genes. At this time point, the distribution of the reporter mRNA

reaches steady state and localization in foci is dictated by the rates of entry and egress (Akef et

al. 2013). In these cells, *ftz-Δi-5′SS* was mostly localized in foci that contained YTHDC1 (Figure

5C, see arrows in the first row). These foci were adjacent to nuclear speckles. When *no-m6A-ftz-*

Δi-5′SS mRNA was visualized, we observed a slight but consistent colocalization with the

nuclear speckle marker SC35, despite the fact that this mRNA was well exported (Figure 5C).

There was a lower level of colocalization between this mRNA and YTHDC1. These trends were

confirmed by Pearson correlation analysis (Figure 5D). These results indicate that m6A promotes

the transfer of mRNAs from nuclear speckles to nearby YTHDC1-containing nuclear foci

(Figure 5E).

Discussion

Here we show that m6A modifications and the m6A binding proteins, YTHDC1 and

YTHDC2, are required for the nuclear retention of mRNAs containing intact 5′SS motifs.

Moreover, we find that the m6A-binding proteins YTHDC1 and YTHDC2 associate with

ZFC3H1 and U1-70K. It is thus likely that these factors form a complex that targets these

misprocessed mRNAs for nuclear retention and possibly degradation by the nuclear exosome,

which is known to receive substrate RNAs from the PAXT complex. Recent work has

demonstrated that m6A modifications are excluded from regions surrounding exon-exon splice

site junctions by the exon-junction complex (EJC) (Yang et al. 2022; Uzonyi et al. 2023; He et

al. 2023; Luo et al. 2023). Thus, m6A modifications accumulate on unprocessed or misprocessed

RNAs, and in conjunction with intact 5′SS motifs, recruit ZFC3H1 to promote RNA nuclear

retention and decay (Figure 5E). Furthermore, our findings suggest that mRNAs with 5′SS are

first targeted to nuclear speckles, and then transferred to YTHDC1-enriched foci, a nuclear

structure that has been likely documented by other groups (Silla et al. 2018; Wang et al. 2021;

Cheng et al. 2021).

Our new results suggest that m6A modifications act as a layer of quality control to

suppress the expression of unwanted transcripts. Previous reports have implicated m6A in

promoting mRNA export, and this is partially based on the analysis of ALKBH5-depletion,

which we believe has indirect effects on mRNA metabolism. More recent publications have

found that m6A acts as a repressive mark both in terms of nuclear export (Smalec et al. 2022;

Tang et al. 2024), RNA stability (Liu et al. 2020) and even feedback to silence chromatin regions

that produce transposable element-derived RNA (Chen et al. 2021; Liu et al. 2021). One of the

reports that linked m6A to mRNA nuclear export also observed that upon the co-depletion of

METTL3 and METTL14 there was a large increase in the cytoplasmic accumulation of m6A-

modified mRNAs, suggesting that m6A actually represses nuclear mRNA export (Roundtree et

al. 2017). Other studies have found that when m6A modified mRNAs are in the cytoplasm, they

are targeted for decay by UPF1 and the CCR4-NOT deadenylase complex, and this may

represent a fail-safe mechanism to ensure the silencing of any m6A-enriched RNA that leaks out

of the nucleus (Wang et al. 2014; Du et al. 2016; Boo et al. 2022).

One unresolved question is why the depletion of ALKBH5 promoted the export of IPA

transcripts. One possible explanation is that ALKBH5-depletion disrupted nuclear speckles, as

previously reported (Zheng et al. 2013), which are required for the retention of reporter mRNAs

with intact 5′SS motifs (Lee et al. 2022). When we stain ALKBH5-depleted U2OS cells for the

nuclear speckle marker SC35, we do not see any obvious changes in nuclear speckles (data not

shown). Nevertheless, it is possible that ALKBH5-depletion alters some property of speckles that

is not obvious by gross inspection of SC35 immunofluorescence. Another possible explanation is

that there is some compensatory effect in ALKBH5-depleted cells by the other demethylases

(FTO and perhaps ALKBH3), but this needs to be tested.

Finally, our results strongly indicate that links between the m6A recognition proteins (i.e.

YTH domain containing proteins) and the PAXT complex are evolutionarily conserved. In both

humans and *S. pombe*, YTH domain containing proteins appear to be involved in suppressing

transposable element-derived RNAs, suggesting additional conserved features. In *S. pombe*, both

the MTREC complex and Mmi1 act together to degrade meiotic mRNAs that are inappropriately

expressed in interphase cells. MTREC and Mmi1 are bridged by Erh1, which has a human

homolog, ERH, that has been recently shown to regulate the stability of transposable element-

derived RNAs and meiotic mRNAs (McCarthy et al. 2021). Thus, it is likely that many aspects

of the MTREC-Mmi1 pathway in *S. pombe* are conserved in humans, and our latest results

further confirm this idea. Future experiments need to be performed to determine whether other

aspects of MTREC and Mmi1 function, including their regulation by TOR signalling and their

ability to stimulate heterochromatin formation (Vo et al. 2019; Wei et al. 2021) are conserved in

human cells.

Material and Methods

Plasmids constructs and antibodies

All plasmids for reporter constructs in pcDNA3.0, pcDNA3.1 (V5-His) or pcDNA5 are as previously described (Akef et al. 2013; Lee et al. 2015, 2020, 2022; Palazzo et al. 2007). For full length FLAG-ZFC3H1-HA, an insert within the previously published plasmid (FLAG-ZFC3H1 (1-1233)) was deleted by PCR. The *no-m6A-ftz* and *e-m6A-ftz* reporters were synthesized and inserted in pUC57 plasmids (Genscript), subcloned using HindIII and SmaI sites and inserted into pcDNA3.0 or pcDNA3.1 (V5-His) respectively. The FLAG-HA-tagged YTHDC1 plasmid was purchased from Addgene (plasmid # 85167).

Antibodies used in this study include rabbit polyclonals against FLAG (Sigma, F7425), METTL3 (ABclonal, A8370), YTHDC1 (Abcam, ab122340), YTHDC2 (Abcam, ab176846), ALKBH5 (Atlas Antibodies, HPA007196), ZFC3H1 (Bethyl laboratories, A301-457A), U1-70K (Abcam, ab83306), ALY (Zhou et al. 2000), G3BP2 (Proteintech, 16276-1-AP), HA (Sigma, SAB4300), Lamin B1 (Abcam, ab16048), eIF4AIII (generous gift from Dr. Lykke-Anderson's lab), TRAP-α (Görlich et al. 1990), and mouse monoclonals against SC35 (Sigma, Clone 35), FG-nucleoporins (Sigma, mAb414) and α-tubulin (Sigma, DM1A). Immunofluorescence was performed with Alexa647-conjugated secondaries (ThermoFisher) and HRP-conjugated secondaries (Cell Signaling). All antibodies were diluted 1:1000 to 1:2000 for western blotting.

Cell culture, DNA transfection, cell fractionation, and METTL3 inhibition by STM2457

U2OS and HEK293T cells were grown in DMEM media (Wisent) supplemented with 10% fetal bovine serum (FBS) (Wisent) and 5% penicillin/streptomycin (Wisent). DNA

transfection experiments were performed as previously described (Akef et al. 2013; Lee et al.

2015, 2020, 2022; Palazzo et al. 2007). U2OS and HEK cells were transfected with the

appropriate amount of DNA plasmid according to the manufacturer's protocol using GenJet

U2OS or Lipo293T DNA *in vitro* transfection reagent (SignaGen Laboratories) for 18 to 24 hrs.

Cell fractionation was performed as previously described (Lee et al. 2020). Equivalent

amounts of nuclear and cytoplasmic lysates were analysed.

For METTL3 inhibition by STM2457, U2OS cells were transfected with the appropriate

plasmids as described above. 6h post-transfection, cells were treated with 100 µM STM2457

(Sigma-Aldrich) or in DMSO for 18h. Cells were fixed and fluorescent in situ hybridization was

performed as described below.

Fluorescent in situ hybridization (FISH) staining, immunostaining and analysis of nuclear
356

speckle targeting

Fluorescent in situ hybridization and immunostaining were performed as previously

described (Gueroussov et al. 2010; Lee et al. 2015, 2020, 2022).

Analysis of nuclear speckle targeting (Figure 5B) was performed as previously described

(Akef et al. 2013). Each analysed region of interest consists of a rectangular region drawn to

cover a single SC35-positive speckle and its surroundings (1–4 µm^2). Note that individual

nuclear speckles were categorized using Pearson correlation intervals of 0.1, from strongest

(Pearson correlation of 1 to 0.9) to weakest (Pearson correlation of 0.6 to 0.5). Any Pearson

correlation < 0.5 was scored as "not localized". Analysis of RNA foci (Figure 5D) was similarly

performed except that each analysed region of interest consists of a rectangular region drawn to

cover an RNA foci and its surroundings (1–4 μm^2) and Pearson correlation between the RNA

and either SC35 or YTHDC1 was tabulated.

Lentiviral delivered shRNA protein depletion

The lentiviral delivered shRNA protein depletion was performed as previously described

(Akef et al. 2013; Lee et al. 2015, 2020, 2022; Palazzo et al. 2007). HEK293T were plated at

50% confluency on 60 mm dishes and transiently transfected with the gene specific shRNA

pLKO.1 plasmid (Sigma), packaging plasmid (Δ8.9) and envelope (VSVG) vectors using

Lipo293T DNA *in vitro* transfection reagent (SignaGen Laboratories) according to the

manufacture's protocol. Viruses were harvested from the media 48 h post-transfection and added

to U2OS cells pre-treated with 8 μg/ml hexadimethrine bromide. Cells were selected with 2

μg/ml puromycin media for at least 4 to 6 days. Western blotting was used to determine the

efficiency of YTHDC1/YTHDC2, METTL3, ALKBH5 and ZFC3H1 depletion. The shRNAs

constructs (Sigma) used in this study are as follows: **ZFC3H1-2** "TRCN0000432333" 5'-

CCGGGACTGA TGACATCGCT AATTTCTCGA GAAATTAGCG ATGTCATCAG

TCTTTTTTG-3', **YTHDC1-1** "TRCN0000243987" 5'- CCGGTGGATT TGCAGGCGTG

AATTACTCGA GTAATTCACG CCTGCAAATC CATTTTTG -3', **YTHDC1-2**

"TRCN0000243988" 5'-CCGGCACCAG AGACCAGGGT ATTTACTCGA GTAAATACCC

TGGTCTCTGG TGTTTTTG -3', **YTHDC2-1** "TRCN0000157638" 5'-CCGGCGGAAG

CTAAATCGAG CCTTTCTCGA GAAAGGCTCG ATTTAGCTTC CGTTTTTTG-3',

YTHDC2-2 "TRCN0000232759" 5'-CCGGCAGAAG TGGCATAGCT TATTTCTCGA

GAAATAAGCT ATGCCACTTC TGTTTTTG -3', **ALKBH5-1** "TRCN0000064783" 5'-

CCGGG AAAGG CTGTTGGCAT CAATACTCGA GTATTGATGC CAACAGCCTT

TCTTTTTG-3', **ALKBH5-2** "TRCN0000064787" 5'-CCGGCCTCAG GAAGACAAGA

TTAGACTCGA   GTCTAATCTT   GTCTTCCTGA   GGTTTTTG-3',   **METTL3-1**

"TRCN0000034714" 5'-CCGGGCCTTA ACATTGCCCA CTGATCTCGA GATCAGTGGG

CAATGTTAAG GCTTTTTG-3', **METTL3-2** "TRCN0000034717" 5'- CCGGGCCAAG

GAACAATCCA   TTGTTCTCGA   GAACAATGGA   TTGTTCCTTG   GCTTTTTG-3',

METTL3-3 "TRCN0000289743" 5'-CCGGGCAAGT ATGTTCACTA TGAAACTCGA

GTTTCATAGT GAACATACTT GCTTTTTG-3', **METTL3-4** "TRCN0000289812" 5'-

CCGGCGTCAG   TATCTTGGGC   AAGTTCTCGA   GAACTTGCCC   AAGATACTGA

CGTTTTTG-3'.

FLAG co-immunoprecipitation

FLAG co-immunoprecipitation was performed as previously described (Lee et al. 2022),

except with these modifications. 1X 150 mm dish of HEK293 cells were transfected with either

FLAG-HA-YTHDC1, FLAG-ZFC3H1-HA or FLAG-ZFC3H1 (1-1233) using Lipo293T DNA

in vitro transfection reagent according to manufacturer's instructions (SignaGen Laboratories).

18 to 24h post transfection the cell pellets were trypsinized, collected and washed with 3 times

with in ice-cold 1X PBS and flash frozen in liquid nitrogen. In parallel, another 150 mm dish of

untransfected HEK293 cells of similar confluency was collected. The pellets were lysed in 2 ml

lysis buffer (20 mM Tris-HCl, pH 8, 137 mM NaCl, 1% NP-40, 2 mM EDTA and *cOmplete*

mini-protease inhibitor (Roche)) for 10 min rocking on a nutator at 4 °C. To ensure that the

nuclear proteins were released from chromatin, the lysed sample was passed through a 25¾″

needle 5X times to shear the chromatin before incubation. The lysate was cleared by spinning at

16,100*g* for 10 min. ~950 µl of the lysate was added to ~50 to 60 µl FLAG M2 beads (Sigma-

Aldrich, A2220), previously preincubated overnight with 5% BSA and 667 µg/ml Ultra-Pure

salmon sperm DNA (Invitrogen), and incubated for 3h or overnight at 4 °C. The remaining ~50

µl of the lysate was reserved as "INPUT", and 5X sample buffer was added and the sample was

boiled at 95°C for 5 min. Following incubation, the FLAG M2 beads was washed 2 to 5 times

with 300 mM wash buffer (20 mM Tris-HCl, pH 8, 300 mM NaCl, 0.05% NP-40, 2 mM EDTA

and *cOmplete* mini-protease inhibitor (Roche)), one time with 137 mM wash buffer (same as for

300 mM wash buffer, except with 137 mM NaCl) and transferred to a new Eppendorf tube.

Subsequently, for RNase digested samples, ~400µl of 1:1000 RNase A/T1 (Thermo Scientific,

EN055) diluted in 137 mM wash was added to the FLAG M2 beads and incubated for 20 min at

4°C. Samples that were not RNase digested were treated in the same manner, except the RNase

A/T1 was omitted in the incubation mixture. The samples were washed twice with 137 mM wash

buffer, transferred to a new Eppendorf tube and 50 to 100 µl 2.5X sample buffer was added to

sample and boiled a 95°C for 5 min. Samples were separated on an SDS-PAGE gel and

transferred onto a blot for immunoblotting.

RNaseA/T1 treatment

The RNase efficacy in our co-immunoprecipitation experiments was validated by

showing the loss of RNA integrity upon RNase treatment using a denaturing agarose gel. The co-

immunoprecipitate lysate was treated with or without RNaseA/T1 followed by RNA extraction

by Trizol extraction, followed by isopropanol precipitation. RNA samples were denatured using

formamide and formaldehyde in MOPS buffer and incubated at 65°C for 30 min. Denatured

RNA was run on an agarose gel supplemented with formaldehyde and stained with SYBR Gold.

Integrity was assessed by visualizing the presence of intact 28S and 18S rRNA bands in the gel.

m6A-miCLIP-seq

miCLIP-seq was performed as previously described (Linder et al. 2015), with modifications as detailed below. Briefly, cells were grown in independent batches to represent biological replicates. We extracted total RNA from HEK293 cells using TRIzol according to manufacturer's instructions. After RNA was treated with DNase I, 12 μg of total RNA was used for miCLIP-seq. 2 μl of RNA Fragmentation Reagents (Invitrogen AM8740) was added to the RNA suspended in 20 μl of sterile water. The mixture was incubated in a heat block at 75°C for exactly 12 minutes. Reaction was stopped by adding 2 μl of stop solution (Invitrogen AM8740). The fragmented RNA was incubated with 10 μg of m6A antibody (MBL RN131P) for 2 hours at 4°C. RNA-antibody complexes were UV crosslinked twice with 0.15 J/cm^2 at 254 nm in a Stratalinker 1800. 100μl of Protein G Dynabeads were added, and the samples were rotated for 2 hours at 4°C. Samples were washed three times using iCLIP high salt buffer with 5 min rotation in the cold room for each wash. RNA dephosphorylation was performed using FastAP and T4 polynucleotide kinase. Pre-adenylated L3 adaptors were ligated to the 3′ends of RNAs using the enhanced CLIP ligation method, as detailed previously (Nabeel-Shah and Greenblatt 2023). RNA was 5'-end-labeled with ^{32}P using T4 polynucleotide kinase (New England Biolabs catalogue number M0201L). Samples were separated using 4–12% BisTris-PAGE and transferred to a nitrocellulose membrane (Protran). RNA was recovered from the membrane by digesting proteins with proteinase K (Thermo Fisher catalogue number 25530049). Recovered RNA was reverse transcribed into cDNA using barcoded iCLIP primers. The cDNA was size-selected (low: 70 to 85 nt, middle: 85 to 110 nt, and high: 110 to 180 nt) using 6% TBE-Urea gels. cDNA was circularized using CircLigase™ II ssDNA ligase to add the adaptor to the 5'-end. We added betain at a final concentration of 1 M in the reaction and mixture was incubated

for 2 hours at 60°C, as detailed previously (Nabeel-Shah and Greenblatt 2023). Circularized

cDNA was digested at the internal BamHI site for linearization. Linearized cDNA was PCR

amplified using Phusion High-Fidelity PCR Master Mix (NEB, M0531S).

The final PCR libraries were agarose gel purified on purification columns (QIAGEN).

The eluted DNA was mixed at a ratio of 1:5:5 from the low, middle, and high fractions and

submitted for sequencing on an Illumina NextSeq 500 platform to obtain single-end 51

nucleotide reads with 40 million read depth per sample. miCLIP-seq data is deposited at Gene

Expression Omnibus (GEO, https://www.ncbi.nlm.nih.gov/geo/) with the accession code

GSE230846 for the series, which contains several other datasets submitted as a part of Nabeel-

Shah et al. (under review).

miCLIP-seq analysis

miCLIP-seq analyses were performed as detailed in our previous reports for iCLIP-seq

(Song et al. 2022). Briefly, 51-nt miCLIP-seq raw reads that consist of 3 random positions, a 4-nt

multiplexing barcode, and another 2 random positions, followed by the cDNA sequence were de-

duplicated based on the first 45 nt, and random positions, barcodes, and any 3′-bases matching

Illumina adaptors were removed. Reads shorter than 25 nt were discarded, and the remaining

reads were trimmed to 35nt using Trimmomatic (Bolger et al. 2014). Reads were mapped to the

human genome/transcriptome (Ensembl annotation hg19) using Tophat (Trapnell et al. 2009)

with default settings where reads with a mapping quality < 3 were removed from further

analysis, which also removes multi-mapping reads.

Crosslinked induced truncations (CITS)

The precise m6A site was inferred by identifying m6A antibody crosslink induced

truncation sites, due to the fact that reverse transcriptase does not efficiently transcribe across

crosslinked regions. CLIP Tool Kit (CTK) was used to identify RNA-binding sites at single-

nucleotide resolution (Shah et al. 2017). As miCLIP-seq aims to capture the crosslink-induced

truncation sites (CITS) during cDNA preparation, we therefore called CITS, rather than CIMS

(crosslink-induced mutations), on individual replicates. CITS with FDR ≤ 0.01 were considered

as significant. CITS from 2 replicates were merged at a maximum gap of 1 bp. Each peak was

then extended 2 bp both upstream and downstream of the identified CITS. Subsequently, only

those CITS with a DRACH motif were selected and the coordinates of the base 'A' in the motif

were designated as m6A sites.

Bioinformatic analysis of m6A density and distribution in IPA transcripts

A list of introns that do not overlap with exons on the opposite strand was compiled from

Gencode v.19. To identify ZFC3H1-regulated IPA transcripts, intronic reads from shZFC3H1

U2OS Frac-Seq data (Lee et al. 2022) were analysed using rtracklayer (Lawrence et al. 2009) to

identify introns with enrichment in the first 20% over the last 20%. Next, introns with more

overall reads in shZFC3H1 compared to shControl conditions were selected. IPA 5' splice sites

were inferred from the PolyA feature annotation file from Gencode.

To estimate the levels of IPA transcripts compared to exons in HEK293 cells, previously

published RNAseq data (GEO: GSE165772) was aligned to the Ensemble GRCh38 genome by

STAR (Dobin et al. 2013). Samtools (Danecek et al. 2021) coverage command was used to

obtain the RNAseq coverage (meandepth column) at all the unique IPA transcripts and exon

sites. To compare relative m6A levels each IPA transcripts was randomly matched with an exon

that was both size matched (in nucleotide length, allowing for +/- 10% difference) and

expression matched (mean read depth, allowing for +/- 10% difference). Average m6A reads per

IPA transcripts or exon were compiled. The IPA transcript-exon matching procedure was

repeated five times, to determine whether differences were statistically significant. This entire

procedure was then repeated using two separate RNAseq runs (GSM5049715 and

GSM5049716). It should be noted that many IPAs are present at extremely low levels and this

likely impeded our ability to detect m6A sites; however, this is equally true for the expression-

matched exon group.

To determine the distribution of m6A reads from the miCLIP analysis along IPA

transcripts and exonic regions, m6A peaks were aligned to IPAs or exons using their genomic

coordinates and normalized to the length of the IPA or exon. The distribution was plotted using

the ggplot2 density geometry (Wickham 2009). Most bioinformatical analyses were done using

R Statistical Software v4.2.2 (R Core Team 2022).

Research Contributions

E.S.L., H.W.S., Y.E.W, S.S.J.I., L.S.O. and N.S.K., performed the experiments under the

guidance of A.F.P.. R.E.J. and Y.L.L. performed the bioinformatics analysis of m6A distribution

under the guidance of A.F.P.. S.N.S performed the m6A-miCLIP experiments and S.P. performed

the bioinformatics analysis under the guidance of J.F.G.. E.S.L. and A.F.P. co-wrote the paper,

with input from all other authors.

Funding

This work was supported by a grant to AFP from Natural Sciences and Engineering Research

Council of Canada (FN 492860).

Conflict of Interest

The authors have declared that no conflict of interest exists.

Acknowledgements

We thank members of the Donnelly Centre RNA sequencing facility for performing the m6A-

miCLIP RNA sequencing. We thank G. L. Burke and N. Ahmed, for assistance with the m6A-

miCLIP library preparation, J. Gao for her helpful discussions for the bioinformatics analysis and

W.S. Tse for his technical expertise for our computational needs. Finally, we thank members of

the Palazzo lab for their helpful discussions and insights.

References

Abad X, Vera M, Jung SP, Oswald E, Romero I, Amin V, Fortes P, Gunderson SI. 2008. Requirements for
gene silencing mediated by U1 snRNA binding to a target sequence. *Nucleic Acids Res* **36**: 2338–
2352.

Akef A, Zhang H, Masuda S, Palazzo AF. 2013. Trafficking of mRNAs containing ALREX-promoting
elements through nuclear speckles. *Nucleus* **4**: 326–340.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics **30**: 2114–2120.

Boo SH, Ha H, Lee Y, Shin M-K, Lee S, Kim YK. 2022. UPF1 promotes rapid degradation of m6A-containing
RNAs. *Cell Reports* **39**: 110861.

Chen C, Liu W, Guo J, Liu Y, Liu X, Liu J, Dou X, Le R, Huang Y, Li C, et al. 2021. Nuclear m6A reader YTHDC1
regulates the scaffold function of LINE1 RNA in mouse ESCs and early embryos. *Protein Cell* **12**:
455–474.

Chen H-M, Futcher B, Leatherwood J. 2011. The fission yeast RNA binding protein Mmi1 regulates
meiotic genes by controlling intron specific splicing and polyadenylation coupled RNA turnover.
PLoS One **6**: e26804.

Cheng Y, Xie W, Pickering BF, Chu KL, Savino AM, Yang X, Luo H, Nguyen DTT, Mo S, Barin E, et al. 2021.
N6-Methyladenosine on mRNA facilitates a phase-separated nuclear body that suppresses
myeloid leukemic differentiation. *Cancer Cell* **39**: 958-972.e8.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA,
Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools. *GigaScience* **10**: giab008.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013.
STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

Dominissini D, Moshitch-Moshkovitz S, Schwartz S, Salmon-Divon M, Ungar L, Osenberg S, Cesarkas K,
Jacob-Hirsch J, Amariglio N, Kupiec M, et al. 2012. Topology of the human and mouse m6A RNA
methylomes revealed by m6A-seq. *Nature* **485**: 201–206.

Du H, Zhao Y, He J, Zhang Y, Xi H, Liu M, Ma J, Wu L. 2016. YTHDF2 destabilizes m(6)A-containing RNA
through direct recruitment of the CCR4-NOT deadenylase complex. *Nat Commun* **7**: 12626.

Fortes P, Cuevas Y, Guan F, Liu P, Pentlicky S, Jung SP, Martínez-Chantar ML, Prieto J, Rowe D, Gunderson
SI. 2003. Inhibiting expression of specific genes in mammalian cells with 5' end-mutated U1
small nuclear RNAs targeted to terminal exons of pre-mRNA. *Proc Natl Acad Sci USA* **100**: 8264–
8269.

Garland W, Jensen TH. 2020. Nuclear sorting of RNA. *WIREs RNA* **11**: e1572.

Goraczniak R, Behlke MA, Gunderson SI. 2009. Gene silencing by synthetic U1 adaptors. *Nat Biotechnol*
27: 257–263.

Görlich D, Prehn S, Hartmann E, Herz J, Otto A, Kraft R, Wiedmann M, Knespel S, Dobberstein B,
Rapoport TA. 1990. The signal sequence receptor has a second subunit and is part of a
translocation complex in the endoplasmic reticulum as probed by bifunctional reagents. *J Cell
Biol* **111**: 2283–2294.

Gueroussov S, Tarnawsky SP, Cui XA, Mahadevan K, Palazzo AF. 2010. Analysis of mRNA Nuclear Export
Kinetics in Mammalian Cells by Microinjection. *J Vis Exp* **46**: 2387.

He PC, Wei J, Dou X, Harada BT, Zhang Z, Ge R, Liu C, Zhang L-S, Yu X, Wang S, et al. 2023. Exon
architecture controls mRNA m6A suppression and gene expression. *Science* **379**: 677–682.

Ishigami Y, Ohira T, Isokawa Y, Suzuki Y, Suzuki T. 2021. A single m6A modification in U6 snRNA diversifies
exon sequence at the 5' splice site. *Nat Commun* **12**: 3244.

Jia G, Fu Y, Zhao X, Dai Q, Zheng G, Yang Y, Yi C, Lindahl T, Pan T, Yang Y-G, et al. 2011. N6-
methyladenosine in nuclear RNA is a major substrate of the obesity-associated FTO. *Nat Chem*
Biol **7**: 885–887.

Kaida D, Berg MG, Younis I, Kasim M, Singh LN, Wan L, Dreyfuss G. 2010. U1 snRNP protects pre-mRNAs
from premature cleavage and polyadenylation. *Nature* **468**: 664–668.

Lawrence M, Gentleman R, Carey V. 2009. rtracklayer: an R package for interfacing with genome
browsers. *Bioinformatics* **25**: 1841–1842.

Lee ES, Akef A, Mahadevan K, Palazzo AF. 2015. The consensus 5' splice site motif inhibits mRNA nuclear
export. *PLoS One* **10**: e0122743.

Lee ES, Smith HW, Wolf EJ, Guvenek A, Wang YE, Emili A, Tian B, Palazzo AF. 2022. ZFC3H1 and U1-70K
promote the nuclear retention of mRNAs with 5' splice site motifs within nuclear speckles. *RNA*
28: 878–894.

Lee ES, Wolf EJ, Ihn SSJ, Smith HW, Emili A, Palazzo AF. 2020. TPR is required for the efficient nuclear
export of mRNAs and lncRNAs from short and intron-poor genes. *Nucleic Acids Res* **48**: 11645–
11663.

Legrain P, Rosbash M. 1989. Some cis- and trans-acting mutants for splicing target pre-mRNA to the
cytoplasm. *Cell* **57**: 573–583.

Lesbirel S, Viphakone N, Parker M, Parker J, Heath C, Sudbery I, Wilson SA. 2018. The m6A-methylase
complex recruits TREX and regulates mRNA export. *Sci Rep* **8**: 13827.

Lesbirel S, Wilson SA. 2019. The m6A-methylase complex and mRNA export. *Biochimica et Biophysica*
Acta (BBA) - Gene Regulatory Mechanisms **1862**: 319–328.

Linder B, Grozhik AV, Olarerin-George AO, Meydan C, Mason CE, Jaffrey SR. 2015. Single-nucleotide-
resolution mapping of m6A and m6Am throughout the transcriptome. *Nat Methods* **12**: 767–
772.

Liu J, Dou X, Chen C, Chen C, Liu C, Xu MM, Zhao S, Shen B, Gao Y, Han D, et al. 2020. N6-
methyladenosine of chromosome-associated regulatory RNA regulates chromatin state and
transcription. *Science* **367**: 580–586.

Liu J, Gao M, He J, Wu K, Lin S, Jin L, Chen Y, Liu H, Shi J, Wang X, et al. 2021. The RNA m6A reader
YTHDC1 silences retrotransposons and guards ES cell identity. *Nature* **591**: 322–326.

Liu J, Yue Y, Han D, Wang X, Fu Y, Zhang L, Jia G, Yu M, Lu Z, Deng X, et al. 2014. A METTL3-METTL14
complex mediates mammalian nuclear RNA N6-adenosine methylation. *Nat Chem Biol* **10**: 93–
95.

Luo Z, Ma Q, Sun S, Li N, Wang H, Ying Z, Ke S. 2023. Exon-intron boundary inhibits m6A deposition,
enabling m6A distribution hallmark, longer mRNA half-life and flexible protein coding. *Nat*
Commun **14**: 4172.

McCarthy RL, Kaeding KE, Keller SH, Zhong Y, Xu L, Hsieh A, Hou Y, Donahue G, Becker JS, Alberto O, et al.
2021. Diverse heterochromatin-associated proteins repress distinct classes of genes and
repetitive elements. *Nat Cell Biol* **23**: 905–914.

Meola N, Domanski M, Karadoulama E, Chen Y, Gentil C, Pultz D, Vitting-Seerup K, Lykke-Andersen S,
Andersen JS, Sandelin A, et al. 2016. Identification of a Nuclear Exosome Decay Pathway for
Processed Transcripts. *Mol Cell* **64**: 520–533.

Meyer KD, Saletore Y, Zumbo P, Elemento O, Mason CE, Jaffrey SR. 2012. Comprehensive analysis of
mRNA methylation reveals enrichment in 3' UTRs and near stop codons. *Cell* **149**: 1635–1646.

Nabeel-Shah S, Greenblatt J. 2023. Revised iCLIP-seq Protocol for Profiling RNA–protein Interaction Sites
at Individual Nucleotide Resolution in Living Cells. *BIO-PROTOCOL* **13**: e4688.

Nayler O, Hartmann AM, Stamm S. 2000. The ER Repeat Protein Yt521-B Localizes to a Novel Subnuclear
Compartment. *J Cell Biol* **150**: 949–962.

Ogami K, Manley JL. 2017. Mtr4/ZFC3H1 protects polysomes through nuclear RNA surveillance. *Cell Cycle*
16: 1999–2000.

Ogami K, Richard P, Chen Y, Hoque M, Li W, Moresco JJ, Yates JR, Tian B, Manley JL. 2017. An
Mtr4/ZFC3H1 complex facilitates turnover of unstable nuclear RNAs to prevent their cytoplasmic
transport and global translational repression. *Genes Dev* **31**: 1257–1271.

Palazzo AF, Lee ES. 2018. Sequence Determinants for Nuclear Retention and Cytoplasmic Export of
mRNAs and lncRNAs. *Front Genet* **9**: 440.

Palazzo AF, Qiu Y, Kang YM. 2024. mRNA nuclear export: how mRNA identity features distinguish
functional RNAs from junk transcripts. *RNA Biol* **21**: 1–12.

Palazzo AF, Springer M, Shibata Y, Lee C-S, Dias AP, Rapoport TA. 2007. The signal sequence coding region
promotes nuclear export of mRNA. *PLoS Biol* **5**: e322.

R Core Team. 2022. R: A language and environment for statistical computing. https://www.R-
project.org/.

Roundtree IA, Luo G-Z, Zhang Z, Wang X, Zhou T, Cui Y, Sha J, Huang X, Guerrero L, Xie P, et al. 2017.
YTHDC1 mediates nuclear export of N6-methyladenosine methylated mRNAs. *Elife* **6**: e31311.

Shah A, Qian Y, Weyn-Vanhentenryck SM, Zhang C. 2017. CLIP Tool Kit (CTK): a flexible and robust
pipeline to analyze CLIP sequencing data. *Bioinformatics* **33**: 566–567.

Shichino Y, Otsubo Y, Kimori Y, Yamamoto M, Yamashita A. 2018. YTH-RNA-binding protein prevents
deleterious expression of meiotic proteins by tethering their mRNAs to nuclear foci ed. T.H.
Jensen. *eLife* **7**: e32155.

Shichino Y, Otsubo Y, Yamamoto M, Yamashita A. 2020. Meiotic gene silencing complex MTREC/NURS
recruits the nuclear exosome to YTH-RNA-binding protein Mmi1. *PLOS Genetics* **16**: e1008598.

Silla T, Karadoulama E, Mąkosa D, Lubas M, Jensen TH. 2018. The RNA Exosome Adaptor ZFC3H1
Functionally Competes with Nuclear Export Activity to Retain Target Transcripts. *Cell Rep* **23**:
2199–2210.

Silla T, Schmid M, Dou Y, Garland W, Milek M, Imami K, Johnsen D, Polak P, Andersen JS, Selbach M, et al.
2020. The human ZC3H3 and RBM26/27 proteins are critical for PAXT-mediated nuclear RNA
decay. *Nucleic Acids Res* **48**: 2518–2530.

Smalec BM, Ietswaart R, Choquet K, McShane E, West ER, Churchman LS. 2022. Genome-wide
quantification of RNA flow across subcellular compartments reveals determinants of the
mammalian transcript life cycle. 2022.08.21.504696.
https://www.biorxiv.org/content/10.1101/2022.08.21.504696v1 (Accessed January 5, 2023).

Song J, Nabeel-Shah S, Pu S, Lee H, Braunschweig U, Ni Z, Ahmed N, Marcon E, Zhong G, Ray D, et al.
2022. Regulation of alternative polyadenylation by the C2H2-zinc-finger protein Sp1. *Mol Cell* **82**:
3135-3150.e9.

Spector DL, Fu XD, Maniatis T. 1991. Associations between distinct pre-mRNA splicing components and
the cell nucleus. *EMBO J* **10**: 3467–3481.

Sugiyama T, Sugioka-Sugiyama R. 2011. Red1 promotes the elimination of meiosis-specific mRNAs in
vegetatively growing fission yeast. *EMBO J* **30**: 1027–1039.

Sugiyama T, Thillainadesan G, Chalamcharla VR, Meng Z, Balachandran V, Dhakshnamoorthy J, Zhou M,
Grewal SIS. 2016. Enhancer of Rudimentary Cooperates with Conserved RNA-Processing Factors
to Promote Meiotic mRNA Decay and Facultative Heterochromatin Assembly. *Mol Cell* **61**: 747–
759.

Tang P, Yang J, Chen Z, Du C, Yang Y, Zhao H, Huang L, Li G, Liu F, Dong B, et al. 2024. Nuclear retention
coupled with sequential polyadenylation dictates post-transcriptional m6A modification in the
nucleus. *Mol Cell* **84**: 3758-3774.e10.

Tian B, Pan Z, Lee JY. 2007. Widespread mRNA polyadenylation events in introns indicate dynamic
interplay between polyadenylation and splicing. *Genome Res* **17**: 156–165.

Trapnell C, Pachter L, Salzberg SL. 2009. TopHat: discovering splice junctions with RNA-Seq.
Bioinformatics **25**: 1105–1111.

Ueda Y, Ooshio I, Fusamae Y, Kitae K, Kawaguchi M, Jingushi K, Hase H, Harada K, Hirata K, Tsujikawa K.
2017. AlkB homolog 3-mediated tRNA demethylation promotes protein synthesis in cancer cells.
Sci Rep **7**: 42271.

Uzonyi A, Dierks D, Nir R, Kwon OS, Toth U, Barbosa I, Burel C, Brandis A, Rossmanith W, Le Hir H, et al.
2023. Exclusion of m6A from splice-site proximal regions by the exon junction complex dictates
m6A topologies and mRNA stability. *Mol Cell* **83**: 237-251.e7.

Veitia RA. 2007. Exploring the Molecular Etiology of Dominant-Negative Mutations. *Plant Cell* **19**: 3843–
3851.

Vo TV, Dhakshnamoorthy J, Larkin M, Zofall M, Thillainadesan G, Balachandran V, Holla S, Wheeler D,
Grewal SIS. 2019. CPF Recruitment to Non-canonical Transcription Termination Sites Triggers
Heterochromatin Assembly and Gene Silencing. *Cell Rep* **28**: 267-281.e5.

Wang C, Zhu Y, Bao H, Jiang Y, Xu C, Wu J, Shi Y. 2016. A novel RNA-binding mode of the YTH domain
reveals the mechanism for recognition of determinant of selective removal by Mmi1. *Nucleic*
Acids Res **44**: 969–982.

Wang X, Lu Z, Gomez A, Hon GC, Yue Y, Han D, Fu Y, Parisien M, Dai Q, Jia G, et al. 2014. m6A-dependent
regulation of messenger RNA stability. *Nature* **505**: 117–120.

Wang Y, Fan J, Wang J, Zhu Y, Xu L, Tong D, Cheng H. 2021. ZFC3H1 prevents RNA trafficking into nuclear
speckles through condensation. *Nucleic Acids Research* **49**: 10630–10643.

Wei Y, Lee NN, Pan L, Dhakshnamoorthy J, Sun L-L, Zofall M, Wheeler D, Grewal SIS. 2021. TOR targets an
RNA processing network to regulate facultative heterochromatin, developmental gene
expression and cell proliferation. *Nat Cell Biol* **23**: 243–256.

Wickham H. 2009. *Ggplot2: elegant graphics for data analysis*. Springer, New York.

Xu C, Wang X, Liu K, Roundtree IA, Tempel W, Li Y, Lu Z, He C, Min J. 2014. Structural basis for selective
binding of m6A RNA by the YTHDC1 YTH domain. *Nat Chem Biol* **10**: 927–929.

Xu Y, Zhang W, Shen F, Yang X, Liu H, Dai S, Sun X, Huang J, Guo Q. 2021. YTH Domain Proteins: A Family
of m6A Readers in Cancer Progression. *Frontiers in Oncology* **11**: 629560.

Yang X, Triboulet R, Liu Q, Sendinc E, Gregory RI. 2022. Exon junction complex shapes the m6A
epitranscriptome. *Nat Commun* **13**: 7904.

Yankova E, Blackaby W, Albertella M, Rak J, De Braekeleer E, Tsagkogeorga G, Pilka ES, Aspris D, Leggate
D, Hendrick AG, et al. 2021. Small-molecule inhibition of METTL3 as a strategy against myeloid
leukaemia. *Nature* **593**: 597–601.

Zheng G, Dahl JA, Niu Y, Fedorcsak P, Huang C-M, Li CJ, Vågbø CB, Shi Y, Wang W-L, Song S-H, et al. 2013.
ALKBH5 Is a Mammalian RNA Demethylase that Impacts RNA Metabolism and Mouse Fertility.
Mol Cell **49**: 18–29.

Zhou Y, Zhu J, Schermann G, Ohle C, Bendrin K, Sugioka-Sugiyama R, Sugiyama T, Fischer T. 2015. The
fission yeast MTREC complex targets CUTs and unspliced pre-mRNAs to the nuclear exosome.
Nat Commun **6**: 7050.

Zhou Z, Luo MJ, Straesser K, Katahira J, Hurt E, Reed R. 2000. The protein Aly links pre-messenger-RNA
splicing to nuclear export in metazoans. *Nature* **407**: 401–405.

 Figure Legends

Figure 1. YTHDC1 and YTHDC2 interact with ZFC3H1 and U1-70K

(A) The main components of the MTREC complex (outlined by a dotted line) and associated

factors in *S. pombe* (left), and their human homologs, including the PAXT complex (outlined by

a dotted line) and YTH proteins (right).

(B) HEK293 cells were either transfected with FLAG-tagged YTHDC1 or mock transfected.

After 18-24 hours, cell lysates were collected and subjected to immunoprecipitation using FLAG

M2 beads. Precipitates were then treated with either buffer or a combination of RNase A and

RNase T1 and reisolated to remove RNA-dependent interacting proteins. Samples were

separated by SDS-PAGE and immunoblotted for the indicated proteins. Note that the inputs

represent 5% of the samples used for immunoprecipitation.

(C-D) Similar to (B), except that cells were transfected with FLAG/HA-tagged ZFC3H1 (FLAG-

ZFC3H1-HA).

(E-F) Similar to (B), except that cells were transfected with a FLAG-tagged fragment of

ZFC3H1 containing amino acids 1-1233.

Figure 2. The m6A modification is required for nuclear retention of 5′SS motif containing

mRNAs

(A) Schematic of the intronless fushi-tarazu (*ftz-Δi*) reporters used in this study, with and without

the 5′SS motif in the 3′UTR. Note that the consensus 5′SS motif promotes nuclear retention. The

positions of either consensus DRACH motifs, or their mutated forms that either eliminate the

motifs (AGGCT), or enhance the motifs (AGACT) are indicated.

(B) U2OS cells were transfected with the indicated *ftz* reporters. After 18-24 hours, RNA was

isolated and immunoprecipitated with anti-m6A antibodies or IgG controls. Levels of *ftz* in the

precipitates were assessed by qRT-PCR. Each bar represents the average and standard error of at

three independent experiments. Student t-test was performed, * $p < 0.05$ ** $p < 0.01$ *** $p <$

0.001.

(C-D) U2OS cells were transfected with the indicated *ftz* reporters. 18-24 hours later the cells

were fixed and the mRNA was visualized by fluorescent in situ hybridization (FISH) using

probes against *ftz*. Representative images are shown in (C), scale bar = 20 μm, and quantification

is shown in (D). Each bar represents the average and standard error of at least three independent

experiments, each experiment consisting of at least 30 to 60 cells. Student t-test was performed,

*** $p < 0.001$.

(E-F) U2OS cells were transfected with the indicated *ftz* reporters. 6h post transfection, cells

were treated with 100 μM STM2457 or DMSO. 18 hours later, the cells were fixed and the

mRNA was visualized by FISH. Representative images are shown in (E), scale bar = 20 μm, and

quantification shown in (F). Each bar represents the average and standard error of at least three

independent experiments, each experiment consisting of at least 30 to 60 cells. Student t-test was

performed, * $p < 0.05$.

(G-I) U2OS cells were treated with lentivirus containing four different shRNAs against METTL3

or scrambled control shRNA. (G) Lysates were collected after 96 hours, separated by SDS-

PAGE, and immunoprobed for METTL3 or tubulin. (H-I) Control- or METTL3-depleted cells

were transfected with the indicated *ftz* reporters. 18-24 hours later the cells were fixed and the

mRNA was visualized by FISH. Representative images are shown in (H), scale bar = 20 μm, and

quantification is shown in (I). Each bar represents the average and standard error of three

independent experiments, each experiment consisting of at least 30 to 60 cells. Student t-test was

performed, ** $p < 0.01$.

Figure 3. YTHDC1/YTHDC2 and ALKBH5 are required for the nuclear retention of 5′SS

motif containing mRNAs

(A) U2OS cells were treated with different lentivirus shRNAs against YTHDC1, YTHDC2,

ALKBH5, ZFC3H1 or scrambled control. Note that "ZFC3H1-2" denotes shRNA #2 against

ZFC3H1 as described previously (Lee et al. 2022). Lysates were collected after 96 hours,

separated by SDS-PAGE and immunoprobed for YTHDC1, YTHDC2, ALKBH5, ZFC3H1,

tubulin or mAb414, which recognizes FG-Nucleoporin proteins. To effectively deplete YTHDC1

and YTHDC2, cells were treated with lentivirus containing two shRNAs against YTHDC1 and

another two against YTHDC2.

(B-C) Control- or YTHDC1/2-, ALKBH5-, ZFC3H1-depleted U2OS cells were transfected with

the intronless *ftz* reporter plasmid (+/- 5′SS) and 5′SS motif containing reporter that lacks m6A

methylation sites (*no-m6A-ftz-Δi-5′SS*). 18-24 hours later the cells were fixed and the mRNA was

visualized by FISH. Representative images are shown in (B), scale bar = 20 μm, and

quantification is shown in (C). Each bar represents the average and standard error of three

independent experiments, each experiment consisting of at least 30 to 60 cells. Student t-test was

performed, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

(D) Schematic of *PCF11-IPA* RNA reporter. The position of the FISH probe used to visualize the

IPA transcript is marked in gray and the position of the 3′ cleavage site in the intron is as

indicated.

(E-F) Similar to (B-C), except that Control-, YTHDC1/YTHDC2- or ALKBH5-depleted cells

were transfected with the *PCF11-IPA* RNA reporter.

(G-H) Total RNA was isolated from whole cell lysates (G) or nuclear and cytoplasmic fractions

(H) from control- or YTHDC1/2-, ZFC3H1-depleted U2OS cells. *PCF11* IPA was assessed by

qRT-PCR using oligo-dT to generate cDNA and intronic-specific primers for the amplicon.

Levels were normalized to the control depletion. Note that no signal was detected in reactions

lacking reverse transcriptase (not shown). Each bar represents the average and standard error of

three (G) or two (H) independent experiments.

Figure 4. m6A methylation marks accumulate in IPA transcripts

(A-B) m6A deposition was assessed by m6A-miCLIP RNA-seq from HEK293 cells. For each

intronic region that generates an IPA, an exon that was length (+/-10%) and expression (+/-10%)

matched was randomly selected. This procedure was repeated 5 times to generate 6 sets (IPAs

and exons Rep 1-5). The six sets were then compared for the average number and standard

deviation of m6A sites (A) or by stratifying them based on number of sites (B). To ensure

reproducibility, the relative expression of intronic regions that generate IPAs and exons were

determined based on two independent RNAseq experiments ("Sequencing 1" and "Sequencing

2").

(C) Meta-analysis of m6A density (as determined by m6A-miCLIP RNA-seq) along intronic

portions of IPA transcripts (dark blue) and all exons (cyan). The position of the 5′SS motif and

poly(A) site is indicated. In IPA transcripts, m6A methylation marks are enriched near the 5′SS

motif and poly(A) sites. In exons, m6A methylation marks are depleted from the 5′ and 3′ ends

(i.e. near exon-exon junctions), as documented by others.

(D-F) Three examples of IPA transcripts that accumulate m6A marks. IGV browser tracks are

shown for miCLIP density. Intronic cleavage/polyadenylation sites are indicated by the red

arrows. Total RNA sequencing levels in U2OS cells from previously published work (Lee et al.

2022) is shown. Note that the levels of IPA transcripts are upregulated following ZFC3H1-

depletion (shZFC).

Figure 5. 5′SS motif containing mRNAs are transferred from nuclear speckles to YTHDC1-

containing foci in an m6A-dependent manner

(A-B) Plasmid DNA containing the *ftz-Δi-5′SS* or *no-m6A-ftz-Δi-5′SS* reporter genes was

microinjected into U2OS nuclei. One hour post-injection, cells were fixed and stained for *ftz*

mRNA by FISH and SC35 by immunofluorescence. Representative cells are shown in (B), each

row represents a single field of view. The merged image shows mRNA in red and SC35 in green.

Speckles containing mRNA are indicated by arrows and the scale bar = 10μm. Individual nuclear

speckles (as detected by SC35) were analysed for colocalization with *ftz* mRNA by Pearson's

Correlation Analysis (B) as previously described (Akef et al. 2013). Each bar is the average and

standard error of the mean of three experiments, each experiment consist of 100 nuclear speckles

analysed from 10 cells.

(C-D) Plasmid DNA containing the *ftz-Δi-5′SS* or *no-m6A-ftz-Δi-5′SS* reporter genes was

transfected into U2OS. 24 hours post-transfection, cells were fixed and stained for *ftz* mRNA by

FISH and for YTHDC1 and SC35 by immunofluorescence. Representative cells are shown in

(C), each row represents a single field of view. The merged image shows mRNA in red,

YTHDC1 in green and SC35 in blue. *Ftz* mRNA foci are indicated by arrows and the scale bar =

10μm. Individual *ftz* mRNA foci were analysed for colocalization with either SC35 (y-axis) or

YTHDC1 (x-axis) by Pearson's Correlation Analysis (D). Each dot represents a single mRNA

foci (each cell containing 5-10 foci, cells were obtained from two independent experiments).

(E) Model of how m6A modifications are required for the nuclear surveillance of 5′SS motif

containing mRNAs (see text for more details).

Supplemental Figures

Supplemental Figure 1. (Related to Figure 1)

(A) Co-immunoprecipitation HEK cell lysates were treated with or without RNaseA/T1,

separated on a denaturing agarose gel and stained overnight with SYBR Gold. 28S and 18S

rRNA bands are visible in the mock treated samples but not present in the RNaseA/T1 treated

samples.

(B-C) Densitometry analysis of immunoprecipitations where the fraction of protein in the input

that is present precipitate is quantified (Figure 1B-D).

(D-F) HEK cells were transfected with FLAG-HA-YTHDC1 and lysed after 24 hours. The

samples were separated by SDS-PAGE and immunoblotted with antibodies against HA (which

detects the tagged YTHDC1), YTHDC1 (which detects the tagged and endogenous YTHDC1)

and ZFC3H1. (E) Quantification of tagged and untagged YTHDC1 levels using densitometry

analysis. Levels of each protein were normalized to tubulin and to untransfected samples. (F)

Quantification of endogenous ZFC3H1 levels, normalized similar to (E).

(G-I) Similar to (D-F), except that FLAG-ZFC3H1-HA was transfected in HEK cells.

Supplemental Figure 2. (Related to Figure 2E-F)

852   (A-B) U2OS cells were treated with 100 µM STM2547 or DMSO for 24 hours. Samples were

853   separated by SDS-PAGE and immunoblotted for ZFC3H1, YTHDC1, YTHDC2 and U1-70K (A)

854   which were then quantified by densitometry analysis (B) where protein levels are normalized to

855   tubulin and DMSO treated cells.

Supplemental Figure 3. (Related to Figure 3)

(A-B) Similar to Figure 3A, U2OS cells were treated with different lentivirus shRNAs against

YTHDC1, YTHDC2, ZFC3H1 or scrambled control (shCtrl). Lysates were collected after 96

hours, separated by SDS-PAGE and immunoprobed for YTHDC1, YTHDC2 and ZFC3H1 (A)

and then quantified by densitometry analysis (B) where protein levels are normalized to tubulin

and shCtrl samples.

(C) Schematic of *CCDC71-IPA* RNA reporter. The position of the FISH probe used to visualize

the IPA RNA is marked in gray and the position of the 3′ cleavage site in the intron is as

indicated.

(D-E) Control- or YTHDC1/2-, ALKBH5-, ZFC3H1-depleted U2OS cells were transfected with

the *CCDC71-IPA* RNA reporter, which was analysed similar to Figure 3E-F.

(F) U2OS cells were fractionated into nuclear and cytoplasmic fractions which were separated by

SDS-PAGE and immunoprobed with various protein markers to determine the purity of each

fraction; Lamin B1 (nucleus), tubulin (cytoplasm) and TRAP- α (endoplasmic reticulum).

(G) Similar to (F), except immunoprobed for ALKBH5. To determine the purity of each fraction

ALY (nuclear marker) and tubulin (cytoplasmic marker) were probed.

Lee_Fig2

Lee_Fig3