Acute targeted induction of gut-microbial metabolism disrupts host circadian rhythm

Giorgia Greter[1], Stefanie Oswald[1], Daria Künzli[1], Jiayi Lan[2], Emma Slack[1], Markus Arnoldini[1*]

[1]Department of Health Sciences and Technology, ETH Zurich, 8093, Zurich, Switzerland

[2]Department of Chemistry and Applied Biosciences, ETH Zurich, 8093, Zurich, Switzerland

1. Summary

The gut microbiota and host circadian rhythm mutually influence each other, and microbiota metabolism has been shown to play a role in regulating host circadian function via secretion of fermentation products (FP). Microbial metabolism is dependent on the availability of nutrients for the microbiota, typically through the host's food intake, making it challenging to disentangle the effect of host and microbiota metabolism. In this study, we acutely induced gut microbial metabolic activity without inducing host metabolism in mice. We found that increasing microbial metabolism in the gut altered circadian clock gene expression locally. Actuating microbiota metabolism also reduced host food intake beyond the calories provided by the microbiota, suggesting a systemic signaling effect of microbial metabolism on the host.

Keywords: Gut microbiota, circadian rhythm, microbial metabolism, fermentation products

2. Introduction

From regulating our daily sleep-wake cycles to influencing metabolic processes, the circadian rhythm dominates the rhythm of most life on Earth [1, 2]. The mammalian circadian clock is a fundamental biological process which functions through a tightly regulated network of transcription factors oscillating in positive and negative feedback loops [3, 4, 5, 6, 7]. Light works as a trigger to the hypothalamus, the master pacemaker, which transmits time-of-day signals to the rest of the body to affect fluctuations in physiology and behavior [8, 9]. One of the key behavioral rhythms that is regulated by the circadian rhythm is food intake [10]. In anticipation of the changing food availability with a change in light, the master pacemaker activates orexigenic pathways and hormone production to increase feeding and activity, which in turn triggers anorexigenic pathways through satiety signals to prepare for fasting and rest [11]. Being synchronous with the 24-hour light-dark cycle increases fitness, and conversely, being asynchronous with the external environment is detrimental to health [12, 13, 14].

Recent research has revealed crucial links between the biological clock, feeding behavior, and the intricate ecosystem of the gut microbiota [15, 16]. The circadian oscillations in food intake in the host result in oscillations in gut microbial metabolism and composition in a circadian pattern [17, 18]. Oscillating immune function in the host also plays a role in shaping the gut microbial ecosystem. For example, immunoglobulin (Ig) A, a key antibody that regulates commensal bacteria community composition, is secreted with a circadian rhythm, and host responses to the invasion of pathogenic bacteria are dependent on the time of day [19, 20, 21].

There is a growing interest in the interplay between the gut microbiota and the host, as it is now thought that not only the host circadian rhythm regulates the microbiota, but also vice versa: that the microbiota influences the host circadian rhythm. The microbiota plays a significant role in the amplitude of host clock gene expression oscillations, and is associated

with weight gain and reduced longevity in a dysregulated circadian rhythm [22, 23, 24, 25]. In
line with this, germ-free (GF) mice, which do not have a microbiota, are less sensitive to the
dysregulation of the circadian clock induced by high fat diets [26]. A major way of signaling
between the gut microbiota and the host are microbial fermentation products (FP). FP are
metabolic byproducts of the microbiota which are used by the host for many metabolic and
endocrine functions in the gut-brain axis, as well as for food intake regulation [27, 28, 29]. FP
production by the microbiota triggers satiety signaling in the host through various pathways
and is thought to play a role in regulating the host circadian rhythm due to the circadian nature
of food intake [30, 17]. However, FP concentrations and microbial metabolism are variable
due to the higher order interactions that take place in the complex microbial communities in
the gut, making it difficult to evaluate mechanistic interactions between the microbiota and
the host circadian rhythm [31].

It has previously been challenging to disentangle the host circadian rhythm and eating patterns
from gut microbial metabolism, as metabolic oscillations in the gut microbiota arise from host
food intake. In this study, we use an established intervention to acutely increase microbial
metabolism in the gut without directly providing calories to the host. This disrupted the
normal, host-dependent microbial rhythm, allowing us to evaluate how microbiota activity
regulates host circadian clock expression and subsequent eating patterns. We show that
microbial metabolism regulates host circadian clock gene expression and that this regulation
influences host circadian behavior.

3. Results

The host circadian rhythm influences microbial cyclical fluctuations

The complexity of the gut microbiota and the higher order interactions in the gut can make it
challenging to identify mechanisms by which the microbiota interacts with the host circadian
rhythm [32]. To overcome this challenge, we colonized germ-free (GF) wild-type C57BL/6
mice with a defined minimal microbiota of three species and bred them for several
generations throughout the study. The Easily Accessible Microbiota (EAM) is composed of
Bacteroides thetaiotaomicron (*B. theta*)*, Eubacterium rectale* (*E. rectale*)*,* and *Escherichia*
coli (*E. coli*), and can be used to study bacterial interactions in a well-defined system of
relevant microbiota members. *B. theta* is the most abundant member of the EAM microbiota,
but all three species reach high numbers in the gut of colonized mice (Figure 1A). *B. theta, E.*
rectale, and *E. coli* are members of the phyla Bacteroidetes, Firmicutes, and Proteobacteria
(recently renamed to Bacteroidota, Bacillota, and Pseudomonadota), respectively*,* the three
most abundant phyla in the human gut [15, 33].

To investigate the rhythmic fluctuations in the EAM population, we measured the bacterial
load in EAM mouse feces over two days. Fecal samples were collected every 6 hours and the
bacterial load per gram of feces over time was estimated, as well as the relative abundance of
each species. All three EAM species vary in total population over time, with the population
decreasing at the end of each dark and light cycle (Figure 1B). *E. coli* fluctuates less in
relative abundance compared to *E. rectale* and *B. theta,* which both fluctuate throughout the
dark and light cycles (Figure 1C). Food intake was measured for each mouse using an
isolator-housed TSE Phenomaster system. As a readout for microbial metabolism, hydrogen
gas excretion was also measured over time. Hydrogen is a byproduct of butyrate production,
and *E. rectale* is the only butyrate producer in the EAM, and therefore the dominant hydrogen
producer (Figure S1B) [34, 35]. Hydrogen levels fluctuated in a 24-hour period in the EAM
mouse, closely following mouse food intake, which provides the non-host-available fibers that
the microbiota can metabolize in the gut (Figure 1D). Germ-free mice did not produce
measurable levels of hydrogen despite having the same food intake rhythm (Figure 1E).

Lactulose disrupts the normal rhythm of microbial metabolism

To acutely disrupt the normal rhythm of microbial metabolism in the gut without directly
providing nutrients to the host, we administered lactulose to EAM mice. Lactulose is a
disaccharide that cannot be metabolized by mammals, is known to be metabolized by the gut
microbiota, and is associated with an increase in hydrogen production and reduced colon pH
due to microbial metabolic activity [36, 37, 38]. We administered the lactulose between
Zeitgeber times 3 and 4, several hours after the mice had stopped eating and when hydrogen
production was decreasing (Figure 2A). To account for the effect of the oral gavage treatment
on the microbiota, PBS was administered as a negative control, and all experiments were
replicated in GF mice to control for host effects independently of the microbiota. Mice that
were treated with lactulose increased hydrogen production two hours after treatment
compared to PBS-treated mice, which continued in the normal rhythm of hydrogen production
(Figure 2B). There was a significant difference in hydrogen between the mice in the two
groups at the endpoint of the experiment, indicating that the microbiota had metabolized the
administered lactulose.

Five hours after treatment, after the microbiota had metabolized the lactulose, the mice were
euthanized, and bacterial population sizes in cecum content were measured to evaluate the
effect of lactulose on the EAM microbiota. There was a significant increase in the cecal *E.*
rectale population, but no significant change in the numbers of *B. theta* or *E. coli* (Figure 2C).
Having observed increased hydrogen production in lactulose-treated mice, we further
evaluated the effect of lactulose on microbial metabolism. Microbial metabolism results in an
increase in FP production, which should lead to a decrease in pH in the cecum. There was a
significant drop in cecum content pH when EAM mice were treated with lactulose, which was
associated with an increase in FP concentration in the cecum (Figure S2A and 2D). Lactulose-
treated mice had significantly higher concentrations of acetate, butyrate, and succinate,
indicating increased metabolic activity of all EAM species, despite the lack of significant
increase in *B. theta* and *E. coli* population densities [39, 40]. GF mice that were treated with
lactulose showed a small but significant drop in pH in cecum content, likely due to osmotic
effects (Figure S2A). In GF mice, FP concentrations did not increase after lactulose treatment
(Figure 2E).

In order to disentangle the osmotic from the metabolic effects of lactulose treatment, we
measured the water fraction in feces and cecum content after treatment in EAM mice and GF
mice in all experiments. The ratio of the dry fecal weight to the wet fecal weight of GF mice
decreased with lactulose treatment, indicating an increase in water content in the feces (Figure
S2B). However, the dry to wet feces ratio in EAM mice increased with lactulose treatment,
possibly due to the increased microbial biomass from lactulose metabolism. We further
controlled for the osmotic effect of lactulose by collecting all the feces that were produced by
the mice between the treatment and endpoint of the experiment and normalized the dry weight
to the dry/wet weight ratio (Figure S2C). The osmotic effect of lactulose was weaker than the
metabolic effect in EAM mice feces, indicated by the increase in dry weight in feces. When
we measured the water content in cecum content, we found no significant difference between
treatments in either EAM of GF mice, indicating that the effect of lactulose largely affected
fecal water content rather than cecal water content (Figure S2D). Therefore, we concluded
that the osmotic effect of lactulose was minimal in the cecum in this study and osmotic effects
therefore are not major contributors to the observed changes in bacterial population sizes and
metabolism.

Inducing microbial metabolism alters circadian clock gene expression in GI tissue

To test whether the induction of microbial metabolism changes gene expression driving the
host circadian rhythm, we studied the transcription of several core circadian clock genes in the
mouse gastrointestinal tract (Figure 3A). The circadian clock regulation begins with the core
clock genes (*CLOCK* and *ARNTL),* which activate transcription of the *Period* and

Cryptochrome repressor genes [41, 42]. *CLOCK* and *ARNTL* also activate transcription of the
Reverb and *ROR* genes in a second regulatory loop. Finally, *CLOCK* and *ARNTL* regulate the
transcription of *NFIL3* and *DBP* in a third regulatory loop. These genes all function in a
transcriptional autoregulatory feedback loop throughout the body. As described, EAM mice
were treated with lactulose or PBS between zeitgeber time 3-4 and euthanized roughly five
hours later. Tissue was harvested from the ileum, cecum, and colon for RNA extraction and
RT-PCR. The gene expression of core clock genes was compared between the two treatment
groups. When EAM mice were treated with lactulose, there was a significant increase in
several core clock genes in ileum and colon tissues, but no significant difference in gene
expression in cecum tissue, where it is presumed that most of the microbial metabolism
occurred (Figure 3B and S3A). These differences in expression reflected the fluctuations seen
in peripheral tissues in the normal mouse circadian rhythm [43]. To control for the direct
effect that lactulose may have on the host, this experiment was also conducted in GF mice.
When GF mice were treated with lactulose, there was no significant increase in core clock
genes in any of the studied tissues, indicating that microbial metabolism induced changes in
the peripheral circadian clock (Figure 3C).

Inducing microbial metabolism decreases host food intake in following active cycle

We then tested whether changing host circadian rhythm through specific alteration of gut
microbial metabolism has further effects on circadian behavior of the host. We measured the
rate and total amount of EAM mouse food intake in the dark cycles before and after lactulose
treatment. The dark cycle preceding treatment was used to ensure normal feeding behavior for
each mouse and provided an undisturbed baseline measurement. Lactulose or PBS treatments
were then administered as before at zeitgeber 3-4, and food intake was measured again in the
following dark cycle and compared to the baseline food intake in the previous dark cycle
(Figure 4A). EAM mice treated with lactulose reduced both their food intake rate and total
food intake in the following active cycle, while there was no significant reduction in food
intake in EAM mice that were treated with PBS (Figure 4B and S4A) or in GF mice treated
with lactulose (Figure 4C and S4B). The non-significant trend towards lower food intake in
lactulose-treated GF mice can potentially be explained by the viscosity of the lactulose
solution; viscous solutions given to mice via oral gavage have been shown to decrease food
intake [44]. However, the decrease in food intake is small in GF mice compared to EAM
mice, indicating that actuation of microbiota metabolism is the dominant cause for this of this
effect.

Disruption of circadian food intake patterns is due to systemic effects of microbial
metabolic products

Microbial metabolism of lactulose in the mouse takes place largely in the cecum and colon
[45]. However, core clock gene expression was most significantly altered in the ileum when
mice were administered lactulose and they then consumed less food following treatment,
implying that the effect of microbial metabolism on the host is systemic and not localized to
the point of metabolic processes (Figure 3B and 4B). To test the systemic effect of microbial
metabolism on host food intake, EAM mice were administered a FP mix of succinate, acetate,
and butyrate, as these were the FP that were significantly produced by the EAM in the cecum
after lactulose treatment (Figure 5A). Orally administered FP were presumed to be taken up
by the host in the small intestine, before reaching the cecum [46]. EAM mice reduced their
food intake rate and total food intake following the treatment of FP, indicating that the effect
of the microbiota on host circadian behavior is systemic (Figure 5D).

One potential mechanism by which the microbiota regulates host food intake systemically is
by indirect feeding of the host through fermentation products, which can be used for energy
harvesting by the host [47]. To evaluate whether the systemic effect of the microbiota on host
food intake was solely due to the supplemental energy provided by the FP, we estimated the

total number of kilocalories that were taken up by the host before and after the treatment of
FP, assuming that all FP that were given to the host were used as an energy source (0.19
kilocalories). After accounting for the energy provided by the FP treatment, EAM mice still
reduced their food intake, meaning that they reduced their caloric intake by more than 0.19
kilocalories (Figure 5C).

To further evaluate whether this systemic effect of the FP could be observed when they were
provided to the host indirectly, we estimated the total maximum number of calories that could
be provided to the host by the microbiota after lactulose treatment (Figure 4B). We estimated
that the lactulose treatment provided each mouse with a maximum of 0.52 kilocalories (Fig.
5C). This maximum value assumes that the energy available to the host from lactulose feeding
is equal to the combustion enthalpy of the fed lactulose. This is an overestimation because the
host cannot directly metabolize lactulose and gets energy indirectly after the microbiota has
extracted significant amounts of energy already. After accounting for the supplementary
feeding of the host through FP, we found that EAM mice reduced their food intake following
lactulose treatment by even more than 0.52 kilocalories. This finding indicates that the
microbiota has a systemic effect on circadian behavior through a mechanism other than
energy homeostasis of the host.

210   4. Discussion

The gut microbiome and host circadian rhythm are intricately linked, with evidence
suggesting that the host circadian rhythm can influence the gut microbiota and vice versa [48,
49]. In this study, we show that the microbiota regulates host circadian rhythm and behavior
through metabolic rhythmicity. We first demonstrated that the EAM gut microbiota oscillates
in population and in metabolism in a periodic manner which is closely associated with the
periodic food intake of the host. Similar rhythmic oscillations in bacterial metabolism have
been found independently of microbiota complexities [50, 51, 18]. We disrupted the metabolic
rhythmicity of the EAM by providing a nutrient that is available only to the microbiota during
the host's fasting period to evaluate the effect that microbial metabolism has on host circadian
rhythm. We observed a significant increase of microbial metabolism using one treatment of
lactulose that was comparable to normal production during the host's active state, simulating
the host's active state for the microbiota [17]. Consequently, we observed an increase of
expression of several core clock genes in the ileum and colon tissue [43].

Most microbial metabolism takes place in the cecum, and interestingly, we found no change
in cecum tissue gene expression after treating EAM mice with lactulose. This may indicate a
systemic effect of microbial metabolism on the host, rather than a localized one. In fact, a
systemic relationship between the gut microbiota and small intestinal epithelial rhythmicity
has previously been described as it was found that altering the microbiota through diet causes
changes in small intestinal immune function [52]. The microbiota has a systemic effect on the
host immune system that extends even further than GI tissue – FP induce regulatory T-cell
production and differentiation in the host peripheral tissues [53]. The systemic effect of the
microbiota on the host circadian rhythm influenced host food intake following the lactulose
treatment. We observed an increase in FP concentration in the mouse cecum after
administering lactulose to EAM mice, and a subsequent decrease in food intake in the
following active cycle. We observed a similar decrease in food intake when we administered a
mix of FP directly to the host, indicating that microbial metabolism is in part regulating host
circadian feeding behavior. One possible mechanism by which this may occur is through
microbial feeding of the host [47]. The microbiota regulates host appetite through energy
harvesting of inaccessible fibers: GF mice eat more food in a day than colonized mice to
compensate for the reduced energy extraction in the large intestine that is otherwise provided
by the microbiota [50]. By treating EAM mice with lactulose, we assumed that the host would
receive energy indirectly from microbial fermentation, and that the host would reduce its food

intake in the following dark cycle by the same number of kilocalories. However, after
accounting for the energy that the microbiota was providing the host through lactulose
metabolism, we still observed a reduction in total energy intake in mice. We observed similar
results when we provided the host directly with FP. This finding indicates that providing
energy is not the only mechanism by which FP reduce host feeding, and that there is a further
mechanism that regulates the host's decrease in total energy harvest.

FP may have a systemic effect on the host through various mechanisms, and there is an
association between FP intake and a reduction in food intake [54, 55, 56, 57]. Butyrate dietary
supplementation in mice decreases food intake and promotes fat oxidation, and acetate
supplementation can decrease appetite directly through the blood-brain-barrier [58, 59]. It has
also been found that dietary supplements of fermentable carbohydrates which can be broken
down by the microbiota result in an upregulation of GLP-1 and PYY, as well as a
downregulation in the orexigenic hormone ghrelin [60, 61]. These hormones involved with
satiety signaling are produced in the body in a rhythmic manner [62, 63]. By disrupting the
rhythmic fluctuations of microbial metabolism and increasing FP production in the cecum, the
microbiota may disrupt the circadian satiety signaling hormone production and therefore
disrupt the circadian clock, and subsequently, feeding behavior. It has previously been found
that mice with gene deletions in the circadian clock lose rhythmicity in food intake and FP
concentration in the cecum, and that restoring rhythmic food intake through restricted feeding
restores rhythmicity in FP production, indicating the intricate association between the host
circadian rhythm and the microbiota [64, 65, 66].

In conclusion, our work demonstrates that the gut microbiota regulates host circadian rhythm.
The microbiota breaks down dietary fibers ingested by the host and the metabolic products of
microbial metabolism regulate host circadian gene expression and eating behavior. We
speculate that this regulation extends to other factors associated with host circadian rhythm
such as endocrine signaling, as the effect of the microbiota on the host circadian rhythm was
systemic, and that the microbiota plays a significant role in maintaining regular circadian
behavior through its metabolic byproducts.

271 5. Acknowledgements

We would like to thank Daniel Hoces for providing the base R script which was adapted for
analyzing the Phenomaster data. We would further like to thank Sven Nowok and Dominik
Bacovcin and the animal caretakers at the EPIC facility at ETHZ for maintaining the mouse
lines. We would like to thank Sandra Kaiser, Bidong Nguyen, and Wolf Dietrich-Hardt for
providing the Lactulose dosage for this study. This work was supported by the Swiss National
Science Foundation (No. 1851228) and was supported as a part of NCCR Microbiomes, a
National Centre of Competence in Research, funded by the Swiss National Science
Foundation (No. 180575).

280 6. Author contributions

281 G.G., E.S.,and M.A. designed the research. G.G. performed the experiments and data
282 evaluation, and S.O. and D.K. assisted in the experiments. J.L. performed the mass
283 spectrometry. G.G. and M.A. wrote the manuscript, in cooperation with all authors.

284 7. Declaration of Interest

285 The authors declare no competing interests.

286 8. Figure titles and legends

Figure 1: The host circadian rhythm influences microbial cyclical fluctuations. A: Cell
counts per grams of feces in the EAM mouse (n=8), calculated by qPCR. **B:** Daily

fluctuations of EAM strains in mouse feces. Feces were sampled over 2 days (n=8). Shaded
regions represent the dark cycle in the animal facility, from zt=12 (6PM) to zt=24 (6AM). **C:**
Daily fluctuations in relative abundance of EAM strains in mouse feces. Relative abundance
was calculated as the proportion of total bacterial population. **D:** Circadian food intake and
hydrogen production of the EAM mouse over two days of measurement, where measurements
are taken every 24 minutes (n=16). Curve fit was calculated using formula y~s(x; bs="cs). SE
is illustrated with grey shading surrounding line. **E:** Circadian food intake and hydrogen
production of the GF mouse (n=8).

Figure 2: Lactulose disrupts the normal rhythm of microbial metabolism. A: Schematic
illustrating the experimental procedure. **B:** Hydrogen production increases after lactulose
gavage, but not after PBS gavage (n=16, 8 in each group). Treatment is indicated by a dotted
line at zt=3. An unpaired T-test was performed on final hydrogen measurements (p<0.0001).
C: Bacterial population after gavage (n=22, lactulose=12, PBS=10). *E. rectale* population
increases in lactulose treated mice compared to PBS treated mice (p<0.05). **D:** SCFA
concentration in EAM (n=17, lactulose=10, PBS=7) mice after treatment. Unpaired T-test
revealed an increase in acetate, butyrate, and succinate concentration after multiple testing
correction ("**" = p<0.01, "***" = p<0.001, "****" = p<0.0001). **E:** SCFA concentration in
GF (n=8, 4 in each group) mice after treatment. There is no difference in SCFA concentration
after treatment.

**Figure 3: Inducing microbial metabolism alters circadian clock gene expression in GI
tissue. A:** Schematic of the clock gene network in mice. **B:** Clock gene expression in gut
tissue after EAM mouse treatment (n=17, lactulose=10, PBS=7) measured via RT-PCR.
Unpaired T-test revealed increases in gene expression in the colon and ileum after multiple
testing correction ("*" = p<0.05, "**" = p<0.01). **C:** Clock gene expression in gut tissue after
GF mouse treatment (n=16, 8 in each group). No significant differences were found in gene
expression.

**Figure 4: Inducing microbial metabolism decreases host food intake in following active
cycle. A:** Schematic illustrating the experimental procedure. **B:** EAM mouse (n=15,
lactulose=8, PBS=7) food intake rate in dark phase prior to treatment (control) and following
treatment (treatment). Paired T-test displays a reduction in food intake rate after lactulose
treatment (p<0.01). **C:** GF mouse (n=14, 7 in each group) food intake rate. Paired T-test
displays no difference in food intake rate after either treatment.

**Figure 5: Disruption of circadian food intake patterns is due to systemic effects of
microbial metabolic products. A:** Schematic illustrating the experimental procedure. **B:**
EAM mouse (n=5) food intake rate in dark phase prior to FP treatment (control) and
following treatment (treatment). Paired T-test displays a reduction in food intake rate after FP
treatment (p<0.01). **C:** EAM mouse total caloric intake after accounting for calories provided
by the FP treatment. Paired T-test displays a significant reduction in caloric intake after
accounting for calories provided directly by the FP (p<0.05). **D:** EAM mouse (data calculated
from S4A) total caloric intake after accounting for calories provided by the microbiota after
lactulose metabolism. Paired T-test displays a significant reduction in caloric intake after
accounting for feeding by the microbiota (p<0.05).

9. Methods
5.1 Lead contact

Further information and requests for resources and reagents should be directed to and will be
fulfilled by the lead contact, Markus Arnoldini (Markus.Arnoldini@hest.ethz.ch).

5.2 Materials availability

All materials are available upon request to the corresponding author.

5.3   Mice

All strains and resources are listed in the key resources table.

Wild type C57BL/6 mice were maintained on a standard diet at the ETH Phenomics center. Mice were re-derived Germ-free or bred with the EAM microbiota and kept under strict hygienic conditions in breeding isolators. The EAM microbiota was derived using strains *Bacteroides thetaiotaomicron* [67], *Eubacterium rectale* ATCC, and *Escherichia coli* HS. All experiments began with 9-15 weeks of age with mixed-sex groups. Mice were fed *ad libitum* for the duration of all experiments and were kept with a dark and light period of 12 hours each. All experiments were approved by the Swiss Kantonal authorities (License ZH120/19 and ZH016/21) according to the legal and ethical requirements.

5.4   Metabolic cages

At the start of the experiment, mice were transferred from the breeding isolators to the isolator-housed TSE PhenoMaster® (TSE Systems, Germany) system and were single housed for the duration of the experiment. Mice were acclimated for 4 days after transfer to the metabolic cages. The TSE PhenoMaster® system allows measurements of oxygen, carbon dioxide, and food and water consumption for 160 seconds in 24-minute intervals. Eight metabolic cages were housed in two sterile isolators, and each cage was connected to the system through a HEPA filter. Hydrogen in each cage was measured through a hydrogen sensor (SGP30, Sensirion, Switzerland) which was attached to the TSE Phenomaster® system. Air flow in each cage was set to 0.4 L/min. A two-point calibration of all gases using reference gases was performed within 24 hours of each experiment. The TSE Phenomaster® data and hydrogen data were combined in the data analysis post-experiment.

5.5   Mouse experiments

On the fifth day after mouse transfer to the TSE Phenomaster®, mice were treated by oral gavage with either 132mg Lactulose in 100µL PBS, or 100µL PBS. The oral gavage was administered between Zeitgeber 3-4. To measure circadian rhythm changes in the host and changes in the gut, mice were euthanized 5 hours after the gavage. To measure the effect of lactulose on host food intake, mice were euthanized the day following the treatment. To measure the effect of FP on host food intake, mice were treated between Zeitgeber 3-4 with 66mg of FP in 100µL of PBS, with a 4:3:3 ratio of sodium succinate, sodium acetate, and sodium butyrate, respectively. Mice which did not eat anything in the 24 hours following oral gavage were excluded from the study.

5.6   Bacterial quantification

Mice were euthanized when a peak in hydrogen production was observed following gavage. Feces or cecum content were collected from mice and weighed for bacterial quantification. Bacterial DNA was extracted using the DNeasy PowerSoil Pro Kit (Qiagen, Germany). qPCR was performed using the FastStart Universal SYBR Green Master Mix (Roche, Switzerland). Primers for each tested strain were diluted to a final concentration of 1µM. DNA was amplified using a QuantStudio 7 Flex instrument (Applied Biosystems, MA, USA) or a StepOnePlus Real-Time PCR System instrument (Applied Biosystems, MA, USA), with an initial denaturation step of 95ºC for 10 minutes, followed by 40 cycles of 95ºC for 15 seconds and 60ºC for 60 seconds. Bacteria were quantified in each run using a standard curve of known bacterial DNA concentration. Germ-free cecum content and water samples were used to generate a detection limit. For each experiment, a contamination control for *Blautia pseudococcoides* was included in the qPCR, and mice with a C_T lower than 26 in cecum samples were excluded from the study.

5.7   Cecal pH measurement

Cecum content was collected in a 2mL Eppendorf tube (Sarstedt, Switzerland) and snap
frozen in liquid nitrogen. At the time of measurement, cecum content was thawed and spun
down at 1000rpm for 30 seconds to accumulate all content at the bottom of the tube. A pH
sensor (Sevencompact S220, Mettler Toledo, Ohio, USA) was completely inserted into the
cecum content and pH was measured.

5.8   Short chain fatty acid quantification

Cecum content was collected in a 2mL Eppendorf tube, snap frozen in liquid nitrogen, and
stored at -80°C until further analysis. For UPLC/MS, cecum content was thawed on ice, then
homogenized in 70% isopropanol and centrifuged. The supernatant was used for FP
quantification using a protocol previously described [68]. A 7-point calibration curve was
generated with FP in known concentrations. Calibrators and samples were spiked with a
mixture of isotope-labeled internal standards, derivatized to 3-nitrophenylhydrazones, and
then quenched with 0.1% formic acid.

An ACQUITY UPLC system (I-Class, Waters, MA, USA) coupled with an Orbitrap Q-
Exactive Plus mass spectrometer (Thermo Scientific, San Jose, CA) were used for UPLC/MS
analysis. For the FP quantification a Kinetex 2.6 µm XB-C18, 50×2.1 mm column
(Phenomenex, Torrance, CA, USA) and a flow rate of 250 µL/min was used with a binary
mixture of solvent A (water with 0.1% formic acid) and solvent B (acetonitrile with 0.1%
formic acid). The gradient starts from 90% of A, then gradually decreases to 75% of A within
6 min and then to 0% of A within 1 min. Then a 0% of A is kept for 1 min and a 90% of A is
kept for 1 min to restore the initial solvent ratio. The column was kept at 40°C and the
autosampler at 5°C.
Acquired raw data were imported to Skyline (V21.1) and MS2 peaks were integrated. A
calibration curve for each compound was built based on the ratio between peak area
of unlabeled and labeled form using a home-built R code. Limit of blank was calculated as
mean of the concentration detected in technical blank samples +- 1.645 standard deviation.

5.9   Dry to wet weight ratio calculation¨

Feces and cecum content were collected at the end of the experiment and the wet weight was
measured. Samples were then lyophilized (Alpha 2-4 LD plus, Martin Christ GmbH,
Germany) for 48 hours and dry weight was measured. The ratio of dry weight to wet weight
was calculated to estimate the water content in the feces and cecum content after the gavage.

For measuring total fecal output after gavage, the bedding in each cage was replaced at the
time of the oral gavage. At the end of the experiment, all fecal pellets were collected and
lyophilized. The dry weight of the feces was measured and normalized to the dry to wet
weight ratio of each mouse.

5.10   RNA extraction

Mouse ileum, cecum, and colon tissues were collected in RNAlater (Thermo Scientific, CA,
USA) and snap frozen in liquid nitrogen. Samples were stored at -80°C until further analysis.
For RNA extractions, samples were thawed on ice and weighed. RNA was extracted using a
modified protocol of the Direct-zol RNA kit (Zymo). Samples were mixed with Trizol
(Thermo Scientific, CA, USA) and incubated for 5 minutes at room temperature. Samples
were then homogenized twice with 0.1mm beads (BioSpec Products, Oklahoma, USA) and
1.0mm silicon carbide beads (BioSpec Products, Oklahoma, USA), for 90 seconds at 30Hz.
Further Trizol was mixed with the samples, and centrifuged. Supernatant was mixed with
chloroform (Sigma-Aldrich, Switzerland) and centrifuged. Supernatant was then washed in
100% ethanol and centrifuged in collection tubes. RNA was washed in Pre-wash buffer and
incubated for 15 minutes at room temperature with 60 units of DNAse. Finally, RNA was
washed in Wash buffer and eluted in Nuclease-free water (Invitrogen, Massachusetts, USA).

RNA concentration and quality were measured using a nanodrop machine (ND-1000
Spectrophotometer, Thermo Scientific, CA, USA) and RNA was diluted to a final
concentration of 20ng/µL in nuclease-free water. Reverse transcription was done using the
High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems). cDNA was diluted
1:25 in nuclease-free water for qPCR.

5.11   Circadian clock gene expression

Host circadian clock gene primers were used to estimate expression using RT-PCR. A
housekeeping gene (TATA binding protein) was used for normalization. The RT-PCR was
carried out as previously described, and sample extracted RNA was included in a reaction
with the housekeeping gene to control for DNA contamination. The delta delta C_T method
was used to estimate relative gene expression for each experiment [69].

5.12   Data analysis

Phenomaster quality control

For all measurements, a variable of zeitgeber time (h) was created where 0-12 represents the
mouse light phase and 12-24 represents the mouse dark phase. The day number of the
experiment was calculated based on the zeitgeber time of the first measurement. Each new
day began at zeitgeber time 0, at the start of the light cycle. To account for faulty
measurements due to disruptive events and for measurement noise, some datapoints were
excluded from the raw dataset. Food intake above 1g and water intake above 1mL in one 24-
minute interval for one mouse was considered to have been caused by human disruption and
was excluded. Food intake of 0.01g as well as negative food and drink values were considered
measurement noise and were excluded. For each mouse, the interquartile range of food and
water intake measurements was calculated. Values greater than the 75^{th} percentile + 1.5 times
the interquartile range were considered outliers and excluded from the dataset. Potential
sources of outlier measurements include food and water loss during human disruption, as well
as leaky water bottles. After the data cleanup, mice which had a total daily food intake
difference of more than 1g in the two days prior to treatment were considered not acclimated
to the metabolic cages and were excluded from further analysis.

Hydrogen sensor data

The hydrogen sensors attached to the TSE Phenomaster® system measured hydrogen every
second, and hydrogen measurements were recorded in a separate file. The median and
maximum hydrogen values for each experimental box and reference box measurement were
calculated. Maximum hydrogen values were used for further analysis. If the reference box
measurements remained visually constant for the duration of the experiment, the hydrogen
data was accepted and merged with the Phenomaster data for further analysis.

Metabolic cage statistical analysis

For estimating circadian rhythm of mouse food intake and hydrogen production, data was
extracted from the raw dataset in the two days prior to mouse treatment. For each mouse, the
average hydrogen value and food intake measurement per timepoint was calculated. A general
additive model was used to fit a curve (using R function mgcv::gam) to these points using the
formula y~s(x; bs="cs).

For estimating food intake per day, data was extracted from the beginning of the day prior to
mouse treatment to the end of the first dark cycle following treatment, resulting in data from
two dark cycles. Cumulative food intake and total food intake were calculated per mouse for
each dark cycle, and food intake rate was calculated as the slope of the linear regression of
cumulative food intake. A paired T-test was used to compare food intake rate and total food

intake for each mouse in the dark cycle prior to treatment and in the dark cycle following
treatment.

<u>Energy harvesting estimations</u>

For estimating the indirect caloric intake of the EAM mouse in the lactulose treatment, it was
assumed that the EAM metabolized 100% of the lactulose, and that the entirety of the
fermentation products of the microbiota were absorbed by the host for caloric feeding. The
combustion enthalpy of lactulose was estimated from the combustion enthalpy of lactose
(1346 kcal/mol), and the kilocalories from 132mg of lactulose were calculated to be 0.52 per
mouse [70]. For estimating the caloric intake of the EAM mouse in the FP treatment, it was
assumed that 100% of FP given to the host were used for energy harvesting. The combustion
enthalpies and masses provided to the host of acetate (0.21 kcal/mmol, 19.8 mg), butyrate
(0.52 kcal/mmol, 19.8 mg), and succinate (0.36 kcal/mmol, 26.4 mg) were used to calculate
the maximum possible calories provided to the host (0.19 kcal).

It was assumed that there are 3.94 kilocalories/gram of standard mouse chow [50]. The total
caloric intake was calculated per mouse for the dark cycle preceding and following lactulose
treatment based on the energy from food intake and indirect feeding from the microbiota and
FP treatment. A paired T-test was used to compare caloric intake for each mouse in the dark
cycle prior to treatment and in the dark cycle following treatment.

For estimating clock gene expression differences in GI tissue and calculating FP
concentration in cecum content an unpaired T-test was carried out to compare both groups. To
account for false positives in the data analysis, the Benjamini and Hochberg multiple testing
correction method was used on both these datasets. For all further comparisons between
treatment groups, an unpaired T-test was used to measure differences. All data analysis was
carried out using R V4.2.0.

10.   Supplemental titles and legends

Figure S1: *E. rectale* is the principal hydrogen producer in the EAM. A:Circadian food
intake and hydrogen production of the EAM mouse over two days of measurement, where
measurements are taken every 24 minutes (n=16). Curve fit was calculated using formula
y~s(x; bs="cs). SE is illustrated with grey shading surrounding line. Shaded regions represent
the dark cycle in the animal facility, from zt=12 (6PM) to zt=24 (6AM). **B:** Circadian food
intake and hydrogen production of the incomplete EAM (colonized with *B. theta* and *E. coli*)
mouse (n=4)

Figure S2: 132mg of lactulose does not have a strong osmotic effect on the host. A:
Cecum content pH of EAM (n=17, lactulose=10, PBS=7) and GF (n=16, 8 in each group)
mice after treatment. Unpaired T-test revealed a decrease in pH in EAM mice (p<0.0001) and
GF mice (p<0.05).**B:** Dry to wet feces ratio in EAM (n=15, lactulose=8, PBS=7) and GF
(n=8, 4 in each group) mice. Unpaired T-test shows an increase (p<0.01) in the ratio in EAM
mice after lactulose treatment. **C:** Total wet feces output after normalizing dry feces weight to
mean dry/wet feces ratio in EAM (n=8, 4 in each group) and GF (n=8, 4 in each group) mice.
Unpaired T-test shows an increase (p<0.05) in the total wet feces output in EAM mice after
lactulose treatment. **D:** Dry to wet cecum content ratio in EAM (n=17, lactulose=10, PBS=7).

Figure S3: Inducing microbial metabolism alters circadian clock gene expression in GI
tissue . A: Clock gene expression in gut tissue after EAM mouse treatment (n=17,
lactulose=10, PBS=7) measured via RT-PCR. Unpaired T-test revealed increases in gene
expression in the colon and ileum after multiple testing correction ("*" = p<0.05, "**" =

p<0.01). **B:** Clock gene expression in gut tissue after GF mouse treatment (n=16, 8 in each
group). No significant differences were found in gene expression.

Figure S4: Inducing microbial metabolism decreases host food intake in following active
cycle. A: EAM mouse (n=15, lactulose=8, PBS=7) total food intake in dark phase prior to
treatment (control) and following treatment (treatment). Paired T-test displays a reduction in
food intake rate after lactulose treatment ("**" = p<0.01). **B:** GF mouse (n=14, 7 in each
group) total food intake. Paired T-test displays no difference in food intake rate after either
treatment.

545 ## 11.   References

[1] J. C. Dunlap, "Molecular Bases for Circadian Clocks Review review word for the late 80s and early 90s would have," 1999.

[2] I. Robinson and A. B. Reddy, "Molecular mechanisms of the circadian clockwork in mammals," *FEBS Letters,* vol. 588, no. 15, pp. 2477-2483, 8 2014.

[3] C. Dibner, U. Schibler and U. Albrecht, "The mammalian circadian timing system: Organization and coordination of central and peripheral clocks," *Annual Review of Physiology,* vol. 72, pp. 517-549, 3 2009.

[4] N. Gekakis, D. Stakins, H. B. Nguyen, F. C. Davis, L. D. Wilsbacher, D. P. King, J. S. Takahashi and C. J. Weitz, "Role of the CLOCK Protein in the Mammalian Circadian Mechanism," *Science,* 1998.

[5] M. H. Hastings, A. B. Reddy and E. S. Maywood, "A clockwork web: Circadian timing in brain and periphery, in health and disease," *Nature Reviews Neuroscience,* vol. 4, no. 8, pp. 649-661, 2003.

[6] K. Z. M. J. S. S. S. L. P. W. D. R. J. X. M. E. S. H. M. H. R. S. M. Kume, "mCRY1 and mCRY2 Are Essential Components of the Negative Limb of the Circadian Clock Feedback Loop to coordinated circadian outputs from the nucleus, ulti-mately regulating rhythms in physiology and behavior (Welsh et al Circadian clocks also appear to ex," *Cell,* 1999.

[7] L. P. Shearman, S. Sriram, D. R. Weaver, E. S. Maywood, I. I. Chaves, B. Zheng, K. Kume, C. C. Lee, G. T. J. Van Der Horst, M. H. Hastings and S. M. Reppert, "Interacting Molecular Loops in the Mammalian Circadian Clock," *Science,* 2000.

[8] A. Patke, M. W. Young and S. Axelrod, "Molecular mechanisms and physiological importance of circadian rhythms," *Nature Reviews Molecular Cell Biology,* vol. 21, no. 2, pp. 67-84, 2 2020.

[9] M. R. Ralph, R. G. Foster, F. C. Davis and M. Menaker, "Transplanted suprachiasmatic nucleus determines circadian period," *Science,* 1990.

[10] E. Challet, "The circadian regulation of food intake," *Nature Reviews Endocrinology,* vol. 15, no. 7, pp. 393-405, 7 2019.

[11] J. H. Strubbe and G. Van Dijk, "The temporal organization of ingestive behaviour and its interaction with regulation of energy balance," *Neuroscience and Biobehavioral Reviews,* vol. 26, no. 4, pp. 485-498, 6 2002.

[12] L. M. Beaver, B. O. Gvakharia, T. S. Vollintine, D. M. Hege, R. Stanewsky and J. M. Giebultowicz, "Loss of circadian clock function decreases reproductive fitness in males of Drosophila melanogaster," *Proceedings of the National Academy of Sciences of the United States of America,* vol. 99, no. 4, pp. 2134-2139, 2002.

[13] A. B. Reddy and J. S. O'Neill, "Healthy clocks, healthy body, healthy mind," *Trends in Cell Biology,* vol. 20, no. 1, pp. 36-44, 1 2010.

[14] G. C. Yuan, Y. J. Liu, M. F. Dion, M. D. Slack, L. F. Wu, S. J. Altschuler and O. J. Rando, "Molecular biology: Genome-scale identification of nucleosome positions in S. cerevisiae," *Science,* vol. 309, no. 5734, pp. 626-630, 7 2005.

[15] M. A. Mahowald, F. E. Rey, H. Seedorf, P. J. Turnbaugh, R. S. Fulton, A. Wollam, N. Shah, C. Wang, V. Magrini, R. K. Wilson, B. L. Cantarel, P. M. Coutinho, B. Henrissat, L. W. Crock, A. Russell, N. C. Verberkmoes, R. L. Hettich and J. I. Gordon, "Characterizing a model human gut microbiota composed of members of its two dominant bacterial phyla," *PNAS,* 2009.

[16] R. M. Voigt, C. B. Forsyth, S. J. Green, P. A. Engen and A. Keshavarzian, "Circadian Rhythm and the Gut Microbiome," *International Review of Neurobiology,* vol. 131, pp. 193-205, 2016.

[17] Y. Tahara, M. Yamazaki, H. Sukigara, H. Motohashi, H. Sasaki, H. Miyakawa, A. Haraguchi, Y. Ikeda, S. Fukuda and S. Shibata, "Gut Microbiota-Derived Short Chain Fatty Acids Induce Circadian Clock Entrainment in Mouse Peripheral Tissue," *Scientific Reports,* vol. 8, no. 1, 12 2018.

[18] C. A. Thaiss, D. Zeevi, M. Levy, G. Zilberman-Schapira, J. Suez, A. C. Tengeler, L. Abramson, M. N. Katz, T. Korem, N. Zmora, Y. Kuperman, I. Biton, S. Gilad, A. Harmelin, H. Shapiro, Z. Halpern, E. Segal and E. Elinav, "Transkingdom control of microbiota diurnal oscillations promotes metabolic homeostasis," *Cell,* vol. 159, no. 3, pp. 514-529, 10 2014.

[19] M. M. Bellet, E. Deriu, J. Z. Liu, B. Grimaldi, C. Blaschitz, M. Zeller, R. A. Edwards, S. Sahar, S. Dandekar, P. Baldi, M. D. George, M. Raffatellu and P. Sassone-Corsia, "Circadian clock regulates the host response to Salmonella," *Proceedings of the National Academy of Sciences of the United States of America,* vol. 110, no. 24, pp. 9897-9902, 6 2013.

[20] O. Pabst and E. Slack, "IgA and the intestinal microbiota: the importance of being specific," *Mucosal Immunology,* vol. 13, no. 1, pp. 12-21, 1 2020.

[21] H. A. Penny, R. G. Domingues, M. Z. Krauss, F. Melo-Gonzalez, M. A. Lawson, S. Dickson, J. Parkinson, M. Hurry, C. Purse, E. Jegham, C. Godinho-Silva, M. Rendas, H. Veiga-Fernandes, D. A. Bechtold, R. K. Grencis, K. M. Toellner, A. Waisman, J. R. Swann, J. E. Gibbs and M. R. Hepworth, "Rhythmicity of intestinal IgA responses confers oscillatory commensal microbiota mutualism," *Science Immunology,* vol. 7, no. 75, 9 2022.

[22] V. Acosta-Rodríguez, F. Rijo-Ferreira, M. Izumo, P. Xu, M. Wight-Carter, C. B. Green and J. S. Takahashi, "Circadian alignment of early onset caloric restriction promotes longevity in male C57BL/6J mice," *Science,* 2022.

[23] F. Bishehsari, P. A. Engen, D. Adnan, S. Sarrafi, S. Wilber, M. Shaikh, S. J. Green, A. Naqib, L. B. Giron, M. Abdel-Mohsen and A. Keshavarzian, "Abnormal food timing and predisposition to weight gain: Role of barrier dysfunction and microbiota," *Translational Research,* vol. 231, pp. 113-123, 5 2021.

[24] C. A. Thaiss, M. Levy, T. Korem, L. Dohnalová, H. Shapiro, D. A. Jaitin, E. David, D. R. Winter, M. Gury-BenAri, E. Tatirovsky, T. Tuganbaev, S. Federici, N. Zmora, D. Zeevi, M. Dori-Bachash, M. Pevsner-Fischer, E. Kartvelishvily, A. Brandis, A. Harmelin, O. Shibolet, Z. Halpern, K. Honda, I. Amit, E. Segal and E. Elinav, "Microbiota Diurnal Rhythmicity Programs Host Transcriptome Oscillations," *Cell,* vol. 167, no. 6, pp. 1495-1510.e12, 12 2016.

[25] Y. Wang, Z. Kuang, X. Yu, K. A. Ruhn, M. Kubo and L. V. Hooper, "The intestinal microbiota regulates body composition through NFIL3 and the circadian clock," *Science,* pp. 912-916, 2017.

[26] V. Leone, S. M. Gibbons, K. Martinez, A. L. Hutchison, E. Y. Huang, C. M. Cham, J. F. Pierre, A. F. Heneghan, A. Nadimpalli, N. Hubert, E. Zale, Y. Wang, Y. Huang, B. Theriault, A. R. Dinner, M. W. Musch, K. A. Kudsk, B. J. Prendergast, J. A. Gilbert and E. B. Chang, "Effects of diurnal variation of gut microbes and high-fat feeding on host circadian clock function and metabolism," *Cell Host and Microbe,* vol. 17, no. 5, pp. 681-689, 5 2015.

[27] E. E. Canfora, J. W. Jocken and E. E. Blaak, "Short-chain fatty acids in control of body weight and insulin sensitivity," *Nature Reviews Endocrinology,* vol. 11, no. 10, pp. 577-591, 10 2015.

[28] H. Han, B. Yi, R. Zhong, M. Wang, S. Zhang, J. Ma, Y. Yin, J. Yin, L. Chen and H. Zhang, "From gut microbiota to host appetite: gut microbiota-derived metabolites as key regulators," *Microbiome,* vol. 9, no. 1, 12 2021.

[29] A. Burokas, S. Arboleya, R. D. Moloney, V. L. Peterson, K. Murphy, G. Clarke, C. Stanton, T. G. Dinan and J. F. Cryan, "Targeting the Microbiota-Gut-Brain Axis: Prebiotics Have Anxiolytic and Antidepressant-like Effects and Reverse the Impact of Chronic Stress in Mice," *Biological Psychiatry,* vol. 82, no. 7, pp. 472-487, 10 2017.

[30] S. O. Fetissov, "Role of the gut microbiota in host appetite control: Bacterial growth to animal feeding behaviour," *Nature Reviews Endocrinology,* vol. 13, no. 1, pp. 11-25, 1 2017.

[31] E. Bairey, E. D. Kelsic and R. Kishony, "High-order species interactions shape ecosystem diversity," *Nature Communications,* vol. 7, 8 2016.

[32] A. L. Gould, V. Zhang, L. Lamberti, E. W. Jones, B. Obadia, N. Korasidis, A. Gavryushkin, J. M. Carlson, N. Beerenwinkel and W. B. Ludington, "Microbiome interactions shape host fitness," *PNAS,* 2018.

[33] D. Mariat, O. Firmesse, F. Levenez, V. D. Guimarăes, H. Sokol, J. Doré, G. Corthier and J. P. Furet, "The firmicutes/bacteroidetes ratio of the human microbiota changes with age," *BMC Microbiology,* vol. 9, 2009.

[34] P. Louis and H. J. Flint, "Diversity, metabolism and microbial ecology of butyrate-producing bacteria from the human large intestine," *FEMS Microbiology Letters,* vol. 294, no. 1, pp. 1-8, 5 2009.

[35] P. Louis and H. J. Flint, "Formation of propionate and butyrate by the human colonic microbiota," *Environmental Microbiology,* vol. 19, no. 1, pp. 29-41, 1 2017.

[36] R. L. Bown, J. A. Gibson, G. E. Sladen, B. Hicks and A. M. Dawson, "Effects of lactulose and other laxatives on ileal and colonic pH as measured by a radiotelemetry device," 1974.

[37] S. Y. Wotzka, M. Kreuzer, L. Maier, M. Zünd, M. Schlumberger, B. Nguyen, M. Fox, D. Pohl, H. Heinrich, G. Rogler, L. Biedermann, M. Scharl, S. Sunagawa, W. D. Hardt and B. Misselwitz, "Microbiota stability in healthy individuals after single-dose lactulose challenge—A randomized controlled study," *PLoS ONE,* vol. 13, no. 10, 10 2018.

[38] Z. Zhang, X. Chen, J. Zhao, C. Tian, X. Wei, H. Li, W. Lin, A. Jiang, R. Feng, J. Yuan and X. Zhao, "Effects of a lactulose-rich diet on fecal microbiome and metabolome in pregnant mice," *Journal of Agricultural and Food Chemistry,* vol. 67, no. 27, pp. 7674-7683, 7 2019.

[39] B. Enjalbert, P. Millard, M. Dinclaux, J. C. Portais and F. Létisse, "Acetate fluxes in Escherichia coli are determined by the thermodynamic control of the Pta-AckA pathway," *Scientific Reports,* vol. 7, 2 2017.

[40] A. A. Salyers, M. O'brien and S. F. Kotarski, "Utilization of Chondroitin Sulfate by Bacteroides thetaiotaomicron Growing in Carbohydrate-Limited Continuous Culture," 1982.

[41] K. H. Cox and J. S. Takahashi, "Circadian clock genes and the transcriptional architecture of the clock mechanism," *Journal of Molecular Endocrinology,* vol. 63, no. 4, pp. R93-R102, 11 2019.

[42] J. S. Takahashi, "Transcriptional architecture of the mammalian circadian clock," *Nature Reviews Genetics,* vol. 18, no. 3, pp. 164-179, 3 2017.

[43] S. Wang, Y. Lin, X. Yuan, F. Li, L. Guo and B. Wu, "REV-ERBα integrates colon clock with experimental colitis through regulation of NF-κB/NLRP3 axis," *Nature Communications,* vol. 9, no. 1, 12 2018.

[44] R. Rasoamanana, P. C. Even, N. Darcel, D. Tomé and G. Fromentin, "Dietary fibers reduce food intake by satiation without conditioned taste aversion in mice," *Physiology and Behavior,* Vols. 110-111, pp. 13-19, 2 2013.

[45] G. Rechkemmer, K. Rönnau and W. V. Engelhardt, "Fermentation of polysaccharides and absorption of short chain fatty acids in the mammalian hindgut," *Comparative Biochemistry and Physiology,* vol. 9, no. 4, pp. 563-568, 1988.

[46] M. G. Schmitt, K. H. Soergel, C. M. Wood and J. J. Steff, "Absorption of Short-Chain Fatty Acids from the Human Ileum," *Digestive Diseases,* 1977.

[47] G. den Besten, K. Lange, R. Havinga, T. H. van Dijk, A. Gerding, K. van Eunen, M. Müller, A. K. Groen, G. J. Hooiveld, B. M. Bakker, D.-J. Reijngoud, d. G. Besten, v. T. Dijk and v. K. Eunen, "Gut-derived short-chain fatty acids are vividly assimilated into host carbohydrates and lipids," *Am J Physiol Gastrointest Liver Physiol,* vol. 305, pp. 900-910, 2013.

[48] L. Litichevskiy and C. A. Thaiss, "The Oscillating Gut Microbiome and Its Effects on Host Circadian Biology," *Annual review of Nutrition,* 2022.

[49] A. Zarrinpar, A. Chaix, S. Yooseph and S. Panda, "Diet and feeding pattern affect the diurnal dynamics of the gut microbiome," *Cell Metabolism,* vol. 20, no. 6, pp. 1006-1017, 12 2014.

[50] D. Hoces, J. Lan, W. Sun, T. Geiser, M. L. Stäubli, E. C. Barazzone, M. Arnoldini, T. D. Challa, M. Klug, A. Kellenberger, S. Nowok, E. Faccin, A. J. Macpherson, B. Stecher, S. Sunagawa, R.

Zenobi, W. D. Hardt, C. Wolfrum and E. Slack, "Metabolic reconstitution of germ-free mice by a gnotobiotic microbiota varies over the circadian cycle," *PLoS Biology,* vol. 20, no. 9, 9 2022.

[51] X. Liang, F. D. Bushman and G. A. FitzGerald, "Rhythmicity of the intestinal microbiota is regulated by gender and the host circadian clock," *Proceedings of the National Academy of Sciences of the United States of America,* vol. 112, no. 33, pp. 10479-10484, 8 2015.

[52] T. Tuganbaev, U. Mor, S. Bashiardes, T. Liwinski, S. P. Nobs, A. Leshem, M. Dori-Bachash, C. A. Thaiss, E. Y. Pinker, K. Ratiner, L. Adlung, S. Federici, C. Kleimeyer, C. Moresi, T. Yamada, Y. Cohen, X. Zhang, H. Massalha, E. Massasa, Y. Kuperman, P. A. Koni, A. Harmelin, N. Gao, S. Itzkovitz, K. Honda, H. Shapiro and E. Elinav, "Diet Diurnally Regulates Small Intestinal Microbiome-Epithelial-Immune Homeostasis and Enteritis," *Cell,* vol. 182, no. 6, pp. 1441-1459.e21, 9 2020.

[53] N. Arpaia, C. Campbell, X. Fan, S. Dikiy, J. Van Der Veeken, P. Deroos, H. Liu, J. R. Cross, K. Pfeffer, P. J. Coffer and A. Y. Rudensky, "Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation," *Nature,* vol. 504, no. 7480, pp. 451-455, 2013.

[54] C. L. Adam, S. W. Gratz, D. I. Peinado, L. M. Thomson, K. E. Garden, P. A. Williams, A. J. Richardson and A. W. Ross, "Effects of dietary fibre (pectin) and/or increased protein (casein or pea) on satiety, body weight, adiposity and caecal fermentation in high fat diet-induced obese rats," *PLoS ONE,* vol. 11, no. 5, 5 2016.

[55] E. S. Chambers, A. Viardot, A. Psichas, D. J. Morrison, K. G. Murphy, S. E. Zac-Varghese, K. MacDougall, T. Preston, C. Tedford, G. S. Finlayson, J. E. Blundell, J. D. Bell, E. L. Thomas, S. Mt-Isa, D. Ashby, G. R. Gibson, S. Kolida, W. S. Dhillo, S. R. Bloom, W. Morley, S. Clegg and G. Frost, "Effects of targeted delivery of propionate to the human colon on appetite regulation, body weight maintenance and adiposity in overweight adults," *Gut,* vol. 64, no. 11, pp. 1744-1754, 11 2015.

[56] J. E. Drew, N. Reichardt, L. M. Williams, C. D. Mayer, A. W. Walker, A. J. Farquharson, S. Kastora, F. Farquharson, G. Milligan, D. J. Morrison, T. Preston, H. J. Flint and P. Louis, "Dietary fibers inhibit obesity in mice, but host responses in the cecum and liver appear unrelated to fiber-specific changes in cecal bacterial taxonomic composition," *Scientific Reports,* vol. 8, no. 1, 12 2018.

[57] J. A. Parnell and R. A. Reimer, "Weight loss during oligofructose supplementation is associated with decreased ghrelin and increased peptide YY in overweight and obese adults," *American Journal of Clinical Nutrition,* vol. 89, no. 6, pp. 1751-1759, 1 2009.

[58] G. Frost, M. L. Sleeth, M. Sahuri-Arisoylu, B. Lizarbe, S. Cerdan, L. Brody, J. Anastasovska, S. Ghourab, M. Hankir, S. Zhang, D. Carling, J. R. Swann, G. Gibson, A. Viardot, D. Morrison, E. L. Thomas and J. D. Bell, "The short-chain fatty acid acetate reduces appetite via a central homeostatic mechanism," *Nature Communications,* vol. 5, 4 2014.

[59] Z. Li, C. X. Yi, S. Katiraei, S. Kooijman, E. Zhou, C. K. Chung, Y. Gao, J. K. Van Den Heuvel, O. C. Meijer, J. F. Berbée, M. Heijink, M. Giera, K. W. Van Dijk, A. K. Groen, P. C. Rensen and Y. Wang, "Butyrate reduces appetite and activates brown adipose tissue via the gut-brain neural circuit," *Gut,* vol. 67, no. 7, pp. 1269-1279, 2018.

[60] P. D. Cani, C. Dewever and N. M. Delzenne, "Inulin-type fructans modulate gastrointestinal peptides involved in appetite regulation (glucagon-like peptide-1 and ghrelin) in rats," *British Journal of Nutrition,* vol. 92, no. 3, pp. 521-526, 9 2004.

[61] J. Zhou, R. J. Martin, R. T. Tulley, A. M. Raggio, K. L. Mccutcheon, L. Shen, S. C. Danna, S. Tripathy, M. Hegsted, M. J. Keenan and J. Zhou, "Dietary resistant starch upregulates total GLP-1 and PYY in a sustained day-long manner through fermentation in rodents," *Am J Physiol Endocrinol Metab,* vol. 295, pp. 1160-1166, 2008.

[62] M. Gil-Lozano, E. L. Mingomataj, W. K. Wu, S. A. Ridout and P. L. Brubaker, "Circadian secretion of the intestinal hormone GLP-1 by the rodent L cell," *Diabetes,* vol. 63, no. 11, pp. 3674-3685, 11 2014.

[63] J. Qian, C. J. Morris, R. Caputo, M. Garaulet and F. A. Scheer, "Ghrelin is impacted by the endogenous circadian system and by circadian misalignment in humans," *International Journal of Obesity,* vol. 43, no. 8, pp. 1644-1649, 8 2019.

[64] A. Segers, L. Desmet, S. Sun, K. Verbeke, J. Tack and I. Depoortere, "Night-time feeding of Bmal1−/− mice restores SCFA rhythms and their effect on ghrelin," *Journal of Endocrinology,* vol. 245, no. 1, pp. 155-164, 2020.

[65] S. Sen, S. Dumont, D. Sage-Ciocca, S. Reibel, P. de Goede, A. Kalsbeek and E. Challet, "Expression of the clock gene Rev-erbα in the brain controls the circadian organisation of food intake and locomotor activity, but not daily variations of energy metabolism," *Journal of Neuroendocrinology,* vol. 30, no. 1, 1 2018.

[66] S. Yang, A. Liu, A. Weidenhammer, R. C. Cooksey, D. McClain, M. K. Kim, G. Aguilera, E. D. Abel and J. H. Chung, "The role of mPer2 clock gene in glucocorticoid and feeding rhythms," *Endocrinology,* vol. 150, no. 5, pp. 2153-2160, 5 2009.

[67] E. C. Martens, N. M. Koropatkin, T. J. Smith and J. I. Gordon, "Complex glycan catabolism by the human gut microbiota: The bacteroidetes sus-like paradigm," *Journal of Biological Chemistry,* vol. 284, no. 37, pp. 24673-24677, 9 2009.

[68] G. Liebisch, J. Ecker, S. Roth, S. Schweizer, V. Öttl, H.-F. Schött, H. Yoon, D. Haller, E. Holler, R. Burkhardt and S. Matysik, "Quantification of Fecal Short Chain Fatty Acids by Liquid Chromatography Tandem Mass Spectrometry—Investigation of Pre-Analytic Stability," *Biomolecules,* vol. 9, no. 4, p. 121, 3 2019.

[69] K. J. Livak and T. D. Schmittgen, "Analysis of relative gene expression data using real-time quantitative PCR and the 2-ΔΔCT method," *Methods,* vol. 25, no. 4, pp. 402-408, 2001.

[70] T. H. Clarke and G. Stegeman, "Heats of Combustion of Some Mono-and Disaccharides," 1939.

[71] A. Steimle, A. De Sciscio, M. Neumann, E. T. Grant, G. V. Pereira, H. Ohno, E. C. Martens and M. S. Desai, "Constructing a gnotobiotic mouse model with a synthetic human gut microbiome to study host–microbe cross talk," *STAR Protocols,* vol. 2, no. 2, 6 2021.

[72] S. Brugiroux, M. Beutler, C. Pfann, D. Garzetti, H. J. Ruscheweyh, D. Ring, M. Diehl, S. Herp, Y. Lötscher, S. Hussain, B. Bunk, R. Pukall, D. H. Huson, P. C. Münch, A. C. McHardy, K. D. McCoy, A. J. MacPherson, A. Loy, T. Clavel, D. Berry and B. Stecher, "Genome-guided design of a defined mouse microbiota that confers colonization resistance against Salmonella enterica serovar Typhimurium," *Nature Microbiology,* vol. 2, 11 2016.

Figure 1: The host circadian rhythm influences microbial cyclical fluctuations. A: Cell counts per grams of feces in the EAM mouse (n=8), calculated by qPCR. **B:** Daily fluctuations of EAM strains in mouse feces. Feces were sampled over 2 days (n=8). Shaded regions represent the dark cycle in the animal facility, from zt=12 (6PM) to zt=24 (6AM). **C:** Daily fluctuations in relative abundance of EAM strains in mouse feces. Relative abundance was calculated as the proportion of total bacterial population. **D:** Circadian food intake and hydrogen production of the EAM mouse over two days of measurement, where measurements are taken every 24 minutes (n=16). Curve fit was calculated using formula y~s(x; bs="cs). SE is illustrated with grey shading surrounding line. **E:** Circadian food intake and hydrogen production of the GF mouse (n=8).

Figure 2: Lactulose disrupts the normal rhythm of microbial metabolism. A: Schematic illustrating the experimental procedure. **B:** Hydrogen production increases after lactulose gavage, but not after PBS gavage (n=16, 8 in each group). Treatment is indicated by a dotted line at zt=3. An unpaired T-test was performed on final hydrogen measurements (p<0.0001). **C:** Bacterial population after gavage (n=22, lactulose=12, PBS=10). *E. rectale* population increases in lactulose treated mice compared to PBS treated mice (p<0.05). **D:** SCFA concentration in EAM (n=17, lactulose=10, PBS=7) mice after treatment. Unpaired T-test revealed an increase in acetate, butyrate, and succinate concentration after multiple testing correction ("**" = p<0.01, "***" = p<0.001, "****" = p<0.0001). **E:** SCFA concentration in GF (n=8, 4 in each group) mice after treatment. There is no difference in SCFA concentration after treatment.

Figure 3: Inducing microbial metabolism alters circadian clock gene expression in GI tissue. A: Schematic of the clock gene network in mice. **B:** Clock gene expression in gut tissue after EAM mouse treatment (n=17, lactulose=10, PBS=7) measured via RT-PCR. Unpaired T-test revealed increases in gene expression in the colon and ileum after multiple testing correction ("*" = p<0.05, "**" = p<0.01). **C:** Clock gene expression in gut tissue after GF mouse treatment (n=16, 8 in each group). No significant differences were found in gene expression.

Figure 4: Inducing microbial metabolism decreases host food intake in following active cycle. A: Schematic illustrating the experimental procedure. **B:** EAM mouse (n=15, lactulose=8, PBS=7) food intake rate in dark phase prior to treatment (control) and following treatment (treatment). Paired T-test displays a reduction in food intake rate after lactulose treatment (p<0.01). **C:** GF mouse (n=14, 7 in each group) food intake rate. Paired T-test displays no difference in food intake rate after either treatment.

Figure 5: Disruption of circadian food intake patterns is due to systemic effects of microbial metabolic products. A: Schematic illustrating the experimental procedure. **B:** EAM mouse (n=5) food intake rate in dark phase prior to FP treatment (control) and following treatment (treatment). Paired T-test displays a reduction in food intake rate after FP treatment (p<0.01). **C:** EAM mouse total caloric intake after accounting for calories provided by the FP treatment. Paired T-test displays a significant reduction in caloric intake after accounting for calories provided directly by the FP (p<0.05). **D:** EAM mouse (data calculated from S4A) total caloric intake after accounting for calories provided by the microbiota after lactulose metabolism. Paired T-test displays a significant reduction in caloric intake after accounting for feeding by the microbiota (p<0.05).