**Body size shapes song in honeyeaters**

Eleanor M. Hay[1]*, Matthew D. McGee[1], Craig R. White[1,2], Steven L. Chown[1]

[1] School of Biological Sciences, Monash University, Melbourne, VIC 3800, Australia

[2] Centre for Geometric Biology, School of Biological Sciences, Monash University,

Melbourne, VIC 3800, Australia

*Corresponding author; email: eleanor.m.hay@outlook.com

**ABSTRACT**

Birdsongs are among the most distinctive animal signals. Their evolution is thought to be

shaped simultaneously by habitat structure and by the constraints of morphology. Habitat

structure affects song transmission and detectability, thus influencing song (the acoustic

adaptation hypothesis), while body size and beak size and shape necessarily constrain song

characteristics (the morphological constraint hypothesis). Yet, support for the acoustic

adaptation and morphological constraint hypotheses remains equivocal, and their

simultaneous examination is infrequent. Using a phenotypically diverse Australasian bird

clade, the honeyeaters (Aves: Meliphagidae), we compile a dataset consisting of song,

environmental, and morphological variables for 163 species and jointly examine predictions

of these two hypotheses. Overall, we find that body size constrains song frequency and pace

in honeyeaters. Although habitat type and environmental temperature influence aspects of

song, that influence is indirect, likely via effects of environmental variation on body size,

with some evidence that elevation constrains the evolution of song peak frequency. Our

results demonstrate that morphology has an overwhelming influence on birdsong, in support

of the morphological constraint hypothesis, with the environment playing a secondary role

generally via body size rather than habitat structure. These results suggest that changing body

size, a consequence of both global effects such as climate change, and local effects such as

habitat transformation, will substantially influence the nature of birdsong.

*Keywords:* Acoustic adaptation; morphological constraint; song evolution; spatial covariance;

evolutionary constraint; allometry

## INTRODUCTION

Birdsongs are acoustic signals influencing survival, reproduction, and reproductive isolation [1, 2], thus shaping the evolution of avian diversity [3-5]. Song is a complex trait and influenced by a range of factors but is commonly thought to be mediated by two major factors, one external, one internal. First, signal detectability, which is influenced by the physical properties of the signalling environment [6, 7] is thought to be a major external factor. Consequently, habitat characteristics are proposed to influence the acoustic properties of bird song. The acoustic adaptation hypothesis (AAH) [6] specifically makes the prediction that because low frequency sounds transmit further than high frequency sounds, birds (and other animals) residing in dense habitats, such as forests, will produce lower frequency sounds than their counterparts dwelling in more open environments [8, 9]. Additionally, it is also predicted that birds will sing at slower paces to limit reverberation and further aid signal transmission [9].

Second, birdsong is also subject to constraints imposed by morphology. Body size is one of the most well understood acoustic constraints [10-12] through its effect on the size of the vocal tract and sound producing structures, which places a limit on the lowest frequency an animal can effectively produce [13]. Allometric constraints of acoustic signals are clear: generally, larger animals produce lower frequency signals [1, 10]. In birds, beak morphology influences acoustic signals too. The avian bill is used in coordination with tongue and vocal tract movements to modify sounds [14] and is thought to act as a final modulator of song, influencing both its frequency and temporal components [15-19]. These effects are complicated, however, by the allometric relationship between body size and beak size [19, 20], and by interactions among the biotic and abiotic factors that are thought to influence acoustic signalling. Specifically, beak morphology and body size also affect thermoregulation

and are influenced by environmental temperatures [21-24]. In consequence, what appears to

be a straightforward morphological constraint hypothesis [10-12], is more complicated.

Such complexity, caused by the effects of ecological selection on signal production,

characterises both the acoustic adaptation and morphological constraint hypotheses. For

example, environmental conditions not only influence body size and beak shape, but they

also affect habitat productivity, complexity, and food availability, which, because of dietary

preferences, influence beak shape [14, 25] and, presumably, song characteristics [14, 26, 27].

This may account for mixed outcomes when studies focus on just one of the hypotheses, with

particularly contradictory results for the AAH [28, 29]. Where morphology and habitat have

been considered simultaneously, habitat was found to have little effect on song, body size

influences frequency, and beak size influences the temporal and performance components of

song [30], but understanding of the direct and indirect influences of body size, especially in

relation to the environment, remains incomplete. Therefore, although birds are one of the

most diverse vertebrate clades in which acoustic signalling plays such a conspicuous role in

sexual signalling and speciation [4, 5], the manner in which multiple and sometimes non-

independent factors influence their signals remains opaque [18, 30], and the avian songbook

thus not fully interpretable.

Here, we set out to uncover the effects of both habitat and morphology on bird song at

a macroevolutionary scale, while considering underlying interactions and correlations

between variables. We address this complexity explicitly in a strong inference framework

[31] and use a suite of phylogenetic comparative methods [32] to explore it with

simultaneous investigation of the predictions of the major hypotheses. We focus on a large

and diverse yet monophyletic group of birds to do so, recognising the power that such focus

on a geographically restricted, diverse single clade can bring [33]. The honeyeaters (Aves:

Meliphagidae) are a clade (~190 species) of passerine birds which have been historically

confined to Australasia [34]. They are oscine passerines that learn their songs and are

therefore influenced by cultural transmission [35]. Learned songs are more subject to change

over generations than innate songs and are also thought to be impacted substantially by the

environment because they have greater plasticity [18, 36]. Honeyeaters exhibit exceptional

diversity in characters thought to influence song: body sizes range from <10 g to 350 g [37],

beaks vary from short for feeding on insects to long and decurved for feeding on nectar [38,

39], and species inhabit a wide variety of biomes from arid deserts to the wet tropics [34]

(Fig. 1). Consequently, song in honeyeaters varies greatly among species, with some species

exhibiting complex and variable songs such as the tūī [40], while others have simpler and

more stereotyped songs, such as the noisy miner [41]. Like most bird species, intraspecific

variation does exist in song for some species of honeyeater [e.g. 40, 41], however the

extensive diversity that exists between species is far greater than any potential within-species

variation (Fig. 2). Ultimately, this immense diversity in song, body size, beak shape, and

environments are ideal for disentangling the factors that shape birdsong (Fig. 3).

Specifically, we compile a dataset of song variables (peak frequency, song pace, and

frequency bandwidth), morphological traits (body size and beak shape), and environmental

variables (habitat type, temperature, elevation, and precipitation) for 163 species of

honeyeater (Table S1). We first use a comparative phylogenetic approach consisting of

phylogenetic multiple regression analyses to assess the factors influencing song traits. Models

are also repeated in a Bayesian framework, and the influence of both phylogenetic and spatial

covariance are considered. We next adopt an evolutionary approach, where we first examine

the influence of open and closed habitats on song evolution. This is followed by phylogenetic

Bayesian regression analysis in which we model the effect of continuous morphological and

environmental variables on the phylogenetic variance of song, analogous to the Brownian

rate parameter [42], to obtain further insight of constraints on rates of song evolution. Based

on expectations of the morphological constraint hypothesis, we expect body size to constrain

song frequency and beak morphology to impact the pace of which honeyeaters sing. For the

acoustic adaptation hypothesis, we expect honeyeaters restricted to closed environments to

sing at lower frequencies and slower paces to favour signal transmission.

**METHODS**

**Honeyeater song data**

Song recordings were sourced from xeno-canto (https://xeno-canto.org/), the Macaulay

Library (Cornell Laboratory of Ornithology, https://www.macaulaylibrary.org/), the

Australian National Wildlife Collection sound archive (ANWC), and the Avian Vocalization

Center (AVoCet; https://avocet.integrativebiology.natsci.msu.edu/). We focussed exclusively

on male song and did not consider recordings of female and juvenile song, or other types of

vocalisations such as calls. Where available, expert classifications on recording type available

in databases were followed (e.g. "song" or "call"). Following past studies which have

explored song in honeyeaters, songs were defined as vocalisations including tonal elements,

separated by gaps greater than 1 s [39]. After downloading, recordings were converted from

.mp3 files to .wav files for analysis in Raven Pro v.1.6 (Cornell Laboratory of Ornithology).

This was done in R [43] using the 'warbleR' package [44] and the '*mp32wav*' function with a

sampling rate of 44.1 kHz. Here it is important to note that compression (i.e. MP3 format) of

song recordings has the potential to influence precision of sound frequency measurements by

up to 1 kHz in worst case scenarios [45]. In saying that, compression does not bias

measurements [45], and this precision is likely to have minimal influence in

macroevolutionary studies in which songs and frequency varies largely across species (Table

S1; Fig. 2). To reduce any potential errors or bias cause by poor-quality recordings, we

149 prioritised using the best quality recordings available for each species (recordings scored as

150 "A" or "B" for xeno-canto, recordings ranked four stars or more for Macaulay Library).

151  To maximize the independence of our samples, only one bout of song was analysed

152 from each recording. Maps were generated to visualise the number and location of song

153 recordings in relation to each species breeding range using range maps from BirdLife

154 International [46] (File S1). Recordings made at different geographic locations were assumed

155 to be of different individual birds, and recordings that did not include a clear song example

156 were discarded. Spectrograms were generated for 564 individuals from 163 honeyeater

157 species (1-20 recordings per species; ~85% of honeyeaters sampled; Table S2, File S1) using

158 Raven Pro v.1.6 (Cornell Laboratory of Ornithology). Within Raven Pro, we used a Hann

159 spectrogram with a window size of 256 samples. Following previous studies [e.g. 30, 39, 47],

160 for each song up to 30 notes were manually selected and the following five song variables

161 were extracted: Peak frequency (the frequency at maximum amplitude across the entire song

162 recording, Hz), maximum frequency (upper frequency bound of the highest pitched note in

163 the song, Hz), minimum frequency (lower frequency bound of the lowest pitched note in the

164 song, Hz), song bandwidth (maximum frequency – minimum frequency), and song pace

165 (number of notes/song duration). Peak frequency of each song was estimated using the peak

166 frequency function available in the measurements of Raven Pro (Cornell Laboratory of

167 Ornithology) and amplitude plots were used to measure values of maximum and minimum

168 frequency. The mean value for each song variable was then calculated for each species. All

169 song variables were then $\log_{10}$-transformed prior to statistical analysis to meet parametric

170 assumptions of normality and homogeneity of variance, but also because logarithmic scales

171 of sound frequency correspond to how birds perceive and modulate sound [48].

All five song variables are correlated to some extent, particularly the multiple

measures of frequency (minimum frequency, maximum frequency, and peak frequency; Fig.

S1). Thus, we chose to focus on one of the three frequency variables for simplicity. Out of the

three frequency variables, peak frequency has the most evidence in support of the acoustic

adaptation hypothesis [49]. Additionally peak frequency is thought to be less sensitive to

background noise and recording conditions than minimum and maximum frequencies [50],

and past studies considering only one frequency measurement have typically used peak

frequency [e.g. 29]. For these reasons, we focus on peak frequency over maximum frequency

and minimum frequency and assess the effect of predictor variables on peak frequency, pace,

and bandwidth (Fig. 3). Because we do still include bandwidth, which is calculated from

maximum and minimum frequency, caution needs to be taken with interpreting this variable

due to the potential influence of compression and background noise on the accuracy of this

measure as discussed previously.

**Trait and environmental data**

To consider the influence of morphology on song, measures of body size [37, 51] and beak

width, depth, and length [37, 51] were included (Table S1). As we are only using male song

data, we only considered male beak measurements. Raw measurements of male beak length,

width, and depth were extracted from AVONET [51] and Marki et al. [38] and the mean

value was calculated for each species. All morphological variables were $\log_{10}$-transformed.

From here, beak length, width, and depth were then subjected to principal component

analysis (PCA) using the '*prcomp*' function in R [43]. The first principal component (PC1)

accounted for 92.53% of the cumulative variation and describes bill size. Whereas PC2

accounted for 5.46% of the cumulative variation and describes a contrast between short and

thick bills vs. long and narrow bills (visualised in Fig. 3; Table S3). A phylogenetic

regression of beak PC1 on body size demonstrated that PC1 is strongly correlated with body

size (PGLS regression $\lambda = 0.66$, $\beta = 4.13 \pm 0.22$, $P > 0.0001$, Table S4; Fig. S2). Thus, PC2

was used to represent beak shape in analysis.

To investigate the AAH, habitat type was classified using a 2-point scale, and was

classified as dense or open/semi-open habitats. We used the habitat density variable form the

AVONET database which classifies birds globally into dense, semi-open, and open habitat

types [51]. This defines open habitats as species primarily occurring in desert, grassland, low

shrubs, rocky habitats, cities, and also applies to species living on top of forest canopy; semi-

open as species primarily inhabiting open shrub, scattered bushes, parkland, dry or deciduous

forest, and thorn forest; and closed habitats as species primarily living in tall evergreen forest

with a closed canopy, or in the lower vegetation strata of dense thickets, shrubland or

marshland [51, 52]. Because we are interested in comparing the differences between dense

and open habitats, we grouped honeyeater species with open ($N_{open} = 3$) and semi-open ($N_{semi-open} = 79$)

habitat types together ($N_{open+semi-open} = 82$) and compared them to species from

dense habitats ($N_{dense} = 81$).

Body size and beak morphology are influenced by the environment [21-24, 39],

therefore appropriate bioclimatic variables were extracted from the geographic range of each

species and included in analysis. Because this study is conducted on a macroevolutionary

scale, we focused on classifying these variables at the species-level, and honeyeater breeding

range maps from BirdLife International [46] were used to sample values of mean annual

temperature, annual precipitation, and elevation from the BioClim database of present-day

climatic conditions [53]. These three bioclimatic variables were selected because they are all

commonly expected to influence size and thermoregulation: mean annual temperature was

included because temperature is expected to directly influence thermoregulation [24]; annual

precipitation was selected because this is also strongly liked with thermoregulation and is

especially important for areas with large variation in precipitation such as Australasia [54];

and elevation was included because lower elevations typically have warmer temperatures

than higher elevations, and vary in other ways that have significant impacts on bird size [55,

56]. We used a resolution of 1 km$^2$, and calculated the mean value for each bioclimatic

variable across each species geographic range using the 'raster' package [57] in R [43].

**Phylogeny**

Trait values of closely related species are expected to be more similar than those of distantly

related species. Therefore, to comparatively analyse relationships between habitat,

morphology, and song while accounting for phylogenetic history, we used a recently

reconstructed honeyeater phylogeny from Hay et al. [58]. This phylogeny was constructed

using a combination of nuclear genes, mitochondrial genes, and topological constraints from

previous phylogenomic studies. The phylogeny was trimmed, using the '*drop.tip*' function

from the 'ape' package [59], to the 163 species for which we had song data and was used as a

phylogenetic effect in all analysis.

To visualise correlations between the morphological constraint hypothesis and the

acoustic adaptation hypothesis, song variables, morphological traits, and habitat type were

plotted on the phylogeny (Fig. 2) using the using the functions '*plotTree.wBars*' and

'*contMap*' from the R package 'phytools' [60].

**Analyses**

We used two approaches to assess the acoustic adaptation hypothesis and the morphological

constraint hypothesis. The first is a typical comparative approach using regression analysis to

identify relationships, whereas in the second an evolutionary constraint method is adopted.

When investigating hypotheses in a strong inference context, several biases might

arise. To account for these potential biases a directed acyclic graph [DAG; 61] was created to

visualise causal relationships and the backdoor criterion was applied to determine the

covariates required to assess both hypotheses while considering interactions between the two

hypotheses [62] (Fig. 3). In this case, two confounding biases exist in which a predictor

variable affects both another predictor variable and the response variable of interest [62]. The

first bias exists between body size and beak shape, which are two strongly correlated traits

that have similar predictions for how they could each influence song frequency [19].

Statistically it can be difficult to disentangle effects such interrelated traits might have and

there is a possibility that any effect of beak shape on song might not actually be attributed to

that variable but could in fact be a statistical artefact [19, 20, 63]. Thus, to overcome this, the

effect of body size needs to be controlled and body size must also be included as a predictor

in analysis. The second bias exists between the interrelated effects of phenotypic traits and

the environment. Environment has the potential to affect beak shape and body size, as well as

song, and therefore this needs to be controlled by including environmental predictor variables

in the model as well. Thus, in this scenario, to address both the acoustic adaptation hypothesis

and the morphological constraint hypothesis body size, beak shape, habitat type, temperature,

precipitation, and elevation must all be included as predictor variables in analysis (Fig. 3;

DAG model available at dagitty.net/mz3jqZU). To further consider the direct effects of

environmental variables on body size and beak shape we also use models where we explicitly

test the effect of precipitation, elevation, and temperature on both body size and beak shape.

We first approach the acoustic adaptation hypothesis and the morphological constraint

hypothesis from a typical comparative approach to determine the drivers of song.

Phylogenetic generalised least squares regression (PGLS; [64, 65] was used to determine the

influence of predictor variables on song. PGLS regression tests were run using the 'ape'

[v5.3; 59] and 'nlme' [66] packages in R [43]. An estimated lambda parameter was used to

control for the amount of phylogenetic effect in the model residuals [67].

Spatiophylogenetic modelling [68] was then used to estimate the spatial and

phylogenetic structure in honeyeater song, and to examine the contributions of body size,

beak shape, and environment to variation in song. To model the spatial distributions of each

species, we used honeyeater point occurrence records from Hay et al. [58] which were

originally downloaded from eBird via GBIF [69, 70]. This dataset was reduced to match the

number of species included in our study, and our analysis included a total of 253,742

honeyeater point occurrence records. The spatiophylogenetic model was implemented in the

R package 'INLA' which uses an integrated nested Laplace approximation to estimate the

joint posterior distribution of model parameters [71, 72]. INLA is an approach for latent

Gaussian Markov random field models, providing a significantly faster alternative to Markov

chain Monte Carlo [72]. Prior to analysis, all predictor variables were standardised by

subtracting the mean and dividing by the standard deviation. For the phylogenetic effect,

INLA requires a phylogenetic precision matrix which is the inverse of a phylogenetic

covariance matrix. Prior to inverting, the phylogenetic covariance matrix was standardised by

dividing by its determinant raised to the power of $1/N_{species}$ [68]. To model the spatial effect

across the entire landscape, we followed a previous study that has used this approach to

model speciation patterns in honeyeaters [58], and used a spatial mesh consisting of 859

vertices, which was constructed over honeyeater point occurrence records and averaged

across the spatial random fields of each occurrence record for each species (see

supplementary R script). This approach integrates a spatial random field across each species'

distribution, so each species contributes a single datapoint to the likelihood. This approach

also reduces any bias which could arise from biased data in occurrence records due to

differential sampling effort, and to some extent accounts for variation between wide ranging

and narrowly distributed species. Further information on these models is available in Dinnage

et al. [68]. Default priors were used for both the phylogenetic and spatial effects. The

significance of effects in the models were based on whether the 95% credible intervals of the

effect size overlapped with zero.

We then approach the acoustic adaptation hypothesis and the morphological

constraint hypothesis from an evolutionary perspective to assess the influence of potential

constraints by examining rates of trait evolution using Brownian motion and Ornstein-

Uhlenbeck models [42, 73-75]. We fit multivariate Brownian motion (BM) and Ornstein-

Uhlenbeck (OU) models in R [43] using mvMORPH [76] to assess whether open and closed

habitat have constrained song evolution. We constructed multivariate Ornstein-Uhlenbeck

models and Brownian motion models with either one rate (no evolutionary relationship

between open and closed habitats) or two rates of evolution (open and closed habitats have

evolved independently) and tested the effect on song pace, bandwidth, peak frequency. We

performed stochastic character mapping of open and closed habitats on the phylogeny using

the *make.simmap* function from the 'phytools' package [60] and generated 100 stochastic

character histories using the tip states on the tree. Model fit was compared with AIC. Despite

being limited to testing the influence of discrete predictors, the mvMORPH approach is

beneficial here as a multivariate framework enables us to determine the influence of open and

closed habitats on peak frequency, bandwidth, and pace altogether.

Finally, we apply phylogenetic Bayesian regression [77, 78] to model the effect of

continuous morphological and environmental variables on the phylogenetic variance of song,

analogous to the Brownian rate parameter, to obtain further insight on constraints from an

evolutionary perspective. This approach enables us to examine how continuous predictor

variables such as body size and beak shape may constrain rates of song evolution. Models

were implemented in R with the package 'brms' [79], for Bayesian models using the

probabilistic programming language STAN [77, 78], which is a Markov chain Monte Carlo

(MCMC) sampler that uses techniques based on Hamiltonian Monte Carlo [80]. Here, we

estimate the effect of predictor variables on both the mean to confirm earlier PGLS and INLA

results, as well as the variance to examine constraints on rates of trait evolution. All

continuous variables were transformed to have a mean of 0 and a standard deviation of 0.5.

Phylogenetic relatedness of species in our regression analysis was accounted for with a

phylogenetic covariance matrix calculated using the *vcv* function from the R package 'ape'

[59]. Models were run for at least 2,000 MCMC iterations with 4 chains and convergence

was assessed using the estimated potential scale reduction statistic Rhat [78]. Effects were

judged as significant if the 95% credible intervals of the effect size did not overlap with zero.

**RESULTS**

Multiple PGLS regression models testing the effect of morphological and environmental

variables on song revealed several relationships. Peak frequency ($\beta = -0.27 \pm 0.041$, $P <$

$0.0001$; model $\lambda = 0.44$, $N = 163$) and pace ($\beta = -0.31 \pm 0.087$, $P < 0.001$; model $\lambda = 0.51$, $N$

$= 163$) are negatively related to body size, indicating that overall, smaller birds produce

higher frequency sounds and faster paced songs (Fig. 4a; Table S4). Beak shape and habitat

type had no impact on any of the song variables (Fig. 4a; Table S4). By contrast, bandwidth

($\beta = -0.020 \pm 0.0071$, $P = 0.005$; model $\lambda = 0.44$, $N = 163$) is negatively related to

temperature, in which cooler mean annual temperatures are associated with broader

bandwidths (Fig. 4a; Table S4). Body size in honeyeaters is positively related to precipitation

suggesting that honeyeaters are larger in wetter areas ($\beta = 0.0001 \pm 0.000017$, $P = 0.0021$;

model $\lambda = 0.99$, $N = 163$; Table S4). A bivariate PGLS model of habitat type on body size

identified honeyeaters to be larger in closed than open environments ($\beta = 0.056 \pm 0.024$, $P =$

$0.021$; model $\lambda = 1.00$, $N = 163$; Table S4).

Spatiophylogenetic analysis revealed considerable spatial and phylogenetic structure

to the variation in honeyeater songs and show that song traits are variable across the

distribution of honeyeaters (Fig. 4b-d). Consistent with PGLS models, frequency and pace

are negatively related to body size (Fig. 4b; Fig. 4d; Table S5). Models further confirm the

effects of temperature on frequency bandwidth, indicating that species in warmer areas sing

with narrower bandwidths (Fig. 4c). Congruent with PGLS results, a positive relationship

exists between precipitation and body size (Table S5).

Model comparisons of multivariate Ornstein-Uhlenbeck models and Brownian motion

models indicate that open and closed habitat types have influenced the evolution of song in

honeyeaters. The best model based on AIC was an Ornstein-Uhlenbeck model with two rates

of evolution (Fig. 5a; Table S6), implying that song has evolved toward different

evolutionary optima in open and closed habitats. Estimated theta values extracted from these

models show that values tend to be lower for closed habitats than for open habitats,

suggesting that lower values of frequency, bandwidth, and pace are selected for in closed

habitats.

Bayesian phylogenetic regressions confirm all results of PGLS models (Table S7).

When modelling the influence of predictor variables on the phylogenetic variance of song, we

find peak frequency to be constrained by elevation. This indicates that honeyeaters with at

lower elevations have higher rates of evolution, whereas higher elevations constrain the rate

of peak frequency evolution (Fig. 5b; Table S7).

**DISCUSSION**

In this study we used honeyeaters to test the predictions of two key hypotheses thought to influence acoustic signals: the morphological constraint hypothesis and the acoustic adaptation hypothesis. We found evidence to support the predictions of the morphological constraint hypothesis: body size constrains frequency and pace. We found no influence of beak shape on song, however. The predictions of the acoustic adaptation hypothesis were not fully supported, but temperature was found to influence bandwidth, and we found some evidence habitat type has influenced the evolution of song. However, these findings are likely mediated by environmental effects on body size. Nonetheless, our evolutionary approach showed that elevation constrains the evolution of peak frequency. Overall, we found that the main constraint of song is body size, and this is ultimately modulated by underlying environmental effects of precipitation on body size, which has consequences for song evolution.

**Morphological constraints on song**

In the case of the morphological constraint hypothesis, we found overwhelming evidence that body size constrains peak frequency: smaller-sized honeyeaters produce higher frequency songs. This outcome is congruent with previous findings for other bird groups [18, 28-30, 47, 81], and likely the outcome of a proximal mechanitic relationship where larger birds have larger sound-producing organs, such as the syrinx [10-12]. In turn this raises the question of what factors influence body size. In African tinkerbirds, temperature and latitude account for variation in body size, ultimately having consequences on song frequency [82].

Among honeyeater species, no evidence exists for latitudinal variation in body size (Bergmann's rule) [58], but the current analyses show that that honeyeater body size is negatively related to annual precipitation, indicating that species from wetter areas tend to be

larger than those from drier areas. This could be in part driven by some form of tropical

temperature effect but is also likely the outcome of aridity gradients and honeyeater

responses to them across Australasia. Body size is fundamental to an organism's ability to

maintain homeostasis [23, 24], and precipitation is a key factor in thermoregulation [54, 83].

Australia has a particularly strong precipitation gradient from the coast to the interior [84].

The influence of such precipitation gradients on body size in honeyeaters is apparent in arid

zone honeyeaters which comprise chats (genus *Epithanura*) and the gibberbird (*Ashbyia*

*lovensis*). These honeyeater species are relatively small, weighing 9-13.5 g (*Epithanura* spp.)

and 18 g (*Ashbyia lovensis*), respectively [37] (Table S1), and occupy the driest areas across

the study region. Furthermore, this interplay between body size, song frequency, and the

environment could suggest that indirect effects via ecological selection, in this case

precipitation, on body size in honeyeaters can have consequences for signal production.

The morphological constraint from beak shape is not as strong as that for body size

and no evidence was found for an influence of beak shape on song. This is contrary to

expectations as a previous study including 88 species of honeyeater found both beak size and

shape to negatively influence pace and minimum frequency, respectively [39]. However, that

study did not consider influences of habitat in these models, which was acknowledged as a

limitation because of the complex relationships between beak size and shape, foraging

ecology, and the environment [39]. Beak morphology in general is understood to act as a

mechanical constraint on song and influence movement [14, 16], the extent of which is

underpinned by environmental conditions and beak specialisation for dietary preferences. For

example, mechanics behind small and slender beaks of insectivores prioritise closing velocity

for catching insects, whereas curved and narrow beaks of nectarivores prioritise nectar

transfer [14], and these factors can influence song. Ultimately, this complex relationship

between beak morphology, the environment, and song highlights the importance of exploring

effects of habitat and morphology altogether.

Numerous past studies have identified beak attributes, prominently beak size, to

impact song frequency [15, 16, 18, 39], but we find no evidence for an impact of beak shape

on peak frequency. Disentangling the effects of beak size and shape on song is complicated

by allometric effects of body size on beak size which could confound the relationship

between beak size and many song features. In the present study we accounted for this

allometry by using the second principal component of beak variables, representing shape, and

controlled for body size in all analysis. The discrepancy between the present study and

previous work is not overly surprising given that we focus on beak shape rather than size.

However, only a few past studies have explicitly acknowledged this statistical concern [19,

30] and caution is required when interpreting findings which might not have fully accounted

for body size because any effect of beak shape on song variables could be mistakenly

attributed [19, 20, 63]. Beyond statistical concerns, fully understanding the influence of beak

shape on song is not entirely straightforward and requires more attention because behaviours

such as adjusting gape and coordinated movements of the beak with the tongue also alter

amplification and resonance of song [14].

Body size also constrains song pace: larger honeyeaters sing at slower paces. This

negative effect of body size on song pace is consistent with previous studies of birds [28, 30],

however, it has generally been attributed to beak size, and little discussion of the relationship

between body size and pace exists throughout the literature. Certainly, beak shape does

constrain pace in honeyeaters, but there is clearly an effect of body size too. One potential

explanation is that body size influences air intake and output which could determine song

pace. Larger birds have larger lungs and vocal producing organs, but also have greater

oxygen consumption, and take larger and fewer breaths than smaller birds [85] which could

explain differences in pace. Morphological structures such as the glottis which acts as a valve

between the lungs and mouth, and the trachea which connects the syrinx in birds to the lungs,

determine air movement. In most birds the trachea narrows towards the lungs [86],

potentially limiting air movement, and could cause larger lungs take longer to fill and empty,

perhaps placing a constraint on the pace at which birds can sing.

**Environmental constraints on song**

Empirical evidence for the acoustic adaptation hypothesis has remained mixed and its

predictions are generally not supported [9, 29, 36, 49]. In the present study, no direct

evidence is uncovered for an effect of habitat type on song at a macroevolutionary scale in

honeyeaters. The hypothesised selection imposed by closed habitats is likely not strong

enough to have a macroevolutionary influence, or perhaps open habitats are also subject to

their own constraints on the transmission and reception of signals. Signals produced in open

habitats can experience degradation from wind and environmental attenuation through the

atmosphere, ground, and vegetation [7]. Conversely, any constraint of closed habitats could

possibly be overcome through behaviours such as movement. Honeyeaters in general have

high dispersal ability [38, 58] and smaller sized birds could implement behaviours such as

increased movement to compensate if acoustic signals are not being detected [87].

Our evolutionary approach did, however, indicate that habitat type influences the

evolution of song. Here, model comparisons of multivariate Ornstein-Uhlenbeck models and

Brownian motion models indicate song in open and closed habitats has evolved toward

different optima. This was generally found to support the predictions of the acoustic

adaptation hypothesis: lower values for peak frequency, bandwidth, and pace tended to be

preferred in closed habitats. However, analysis revealed that honeyeaters tend to be larger in

closed than open environments. Such underlying relationships suggest that any effect we

uncover for habitat type influencing song is more likely to be driven by underlying variation

in body size, and ultimately provides further support for body size being the primary

constraint of song.

Although little support is uncovered for the AAH, we find some evidence that other

environmental variables influence song in honeyeaters. In particular, bandwidth is

consistently found to be negatively related to mean annual temperature, indicating that

honeyeater species in warmer areas sing with narrower bandwidths or less frequency

variation. Although the impacts of temperature on bandwidth could be related to habitat type,

body size is more likely the main influence. As we have shown, body size in honeyeaters is

structured by precipitation, however body size in birds is also commonly structured by

temperature [24, 55] and there is likely an underlying relationship between temperature and

body size in honeyeaters. Indeed, both temperature and precipitation jointly influence

thermoregulation and desiccation, and can affect body size [54]. Body size is understood to

place a limit on the lowest pitched sound a bird can produce, and larger species use wider

bandwidths than smaller species [13].

Lastly, our evolutionary approach also found that elevation constrains the rate of peak

frequency evolution, and that species occurring at lower elevations tended to have faster rates

of peak frequency evolution than those at higher elevations. One of the most notable areas for

elevation in honeyeaters is New Guinea, and interestingly our spatial maps (Fig. 4b-c)

indicate that New Guinea is associated with higher frequencies, greater bandwidths, and

faster paces. New Guinea also contains the greatest proportion of species richness for

honeyeaters, where up to 41 species co-occur [38]. Past studies have found that niche

partitioning along elevational zones in New Guinea has enabled many divergent species to

co-exist here [88], and perhaps the constraints of elevation on peak frequency evolution in

this area has contributed to this co-existence of species. Nevertheless, it is also likely that

niche partitioning has resulted in divergent body sizes which has then influenced peak

frequency, and the result of elevation constraining peak frequency evolution could simply be

an artefact of this.

**Conclusion**

This study provides support for the morphological constraint hypothesis and firmly

establishes body size as the main constraint of song in honeyeaters. Effects of habitat type

and any apparent environmental drivers on song have stronger associations with body size

than with habitat structure *per se*.

500           These findings have implications for the future of birdsong in a rapidly changing

world. Changing climates are resulting in a reduction in avian body size [23, 89], and habitat

loss and transformation include declines of larger sized birds as part of overall changes to

local assemblage composition [90, 91]. Our results therefore forecast an increase in higher

pitched songs as body sizes decline. Overall, we demonstrate that morphology and habitat are

deeply intertwined and can jointly influence song. Thus, climate change and habitat loss will

not only have consequences for acoustic signalling and song evolution, but will also

transform the avian songbook.

**Acknowledgements**

We thank Leo Joseph and Margaret Cawsey at ANWC for assistance with accessing recordings and acknowledge receipt of media from The Macaulay Library at the Cornell Lab of Ornithology. Members of the Chown Lab, Melodie McGeoch, and members of the McGeoch lab provided feedback on aspects of this work. Two anonymous reviewers provided useful comments on a previous version of the manuscript.

**Funding**

EMH was supported by an Australian Government Research Training Program (RTP) Scholarship, a Monash Graduate Excellence Scholarship (MGES), and a Faculty of Science Dean's Postgraduate Research Scholarship. SLC was funded by Australian Research Council Discovery Project DP170101046. MDM was funded by Australian Research Council Discovery Early Career Research Award DE180101558.

**Authors' contributions**

EMH and SLC conceptualised the study. Trait and environmental variables were gathered by EMH. Processing of song recordings and song measurements were completed by EMH. Analysis was completed by EMH with supervision and input from MDM, CRW, and SLC. EMH wrote the manuscript with input from SLC, MDM, and CRW. All authors read and approved the final manuscript.

**REFERENCES**

1.    Bradbury J.W., Vehrencamp S.L. 1998 *Principles of Animal Communication*.

Sunderland, Massachusetts, USA, Sinauer.

2.    Catchpole C.K., Slater P.J.B. 2003 *Bird Song: Biological Themes and Variations*.

New York, Cambridge University Press.

3.    Coyne J.A., Orr H.A. 2004 *Speciation*. Sunderland, Massachusetts USA, Sinauer

Associates.

4.    Tobias J.A., Ottenburghs J., Pigot A.L. 2020 Avian diversity: speciation,

macroevolution, and ecological function. *Annu Rev Ecol Evol Syst* **51**, 533-560.

(doi:10.1146/annurev-ecolsys-110218-025023).

5.    Price T. 2008 *Speciation in Birds*. Colorado, Roberts and Co.

6.    Morton E.S. 1975 Ecological sources of selection on avian sounds. *Am Nat* **109**, 17-

34. (doi:doi:10.1086/282971).

7.    Wiley R.H., Richards D.G. 1978 Physical constraints on acoustic communication in

the atmosphere: implications for the evolution of animal vocalizations. *Behav Ecol*

*Sociobiol* **3**, 69-94. (doi:10.1007/BF00300047).

8.    Brumm H., Naguib M. 2009 Environmental acoustics and the evolution of bird song.

*Adv Study Behav* **40**, 1-33.

9.    Ey E., Fischer J. 2009 The "Acoustic Adaptation Hypothesis"—a review of the

evidence from birds, anurans and mammals. *Bioacoustics* **19**, 21-48.

(doi:10.1080/09524622.2009.9753613).

10.   Ryan M.J., Brenowitz E.A. 1985 The role of body size, phylogeny, and ambient noise

in the evolution of bird song. *Am Nat* **126**, 87-100. (doi:10.1086/284398).

11.   Bertelli S., Tubaro P.L. 2002 Body mass and habitat correlates of song structure in a

primitive group of birds. *Biol J Linn Soc* **77**, 423-430. (doi:10.1046/j.1095-

8312.2002.00112.x).

12.   Ey E., Pfefferle D., Fischer J. 2007 Do age- and sex-related variations reliably reflect

body size in non-human primate vocalizations? A review. *Primates* **48**, 253-267.

(doi:10.1007/s10329-006-0033-y).

13.   Friis J.I., Sabino J., Santos P., Dabelsteen T., Cardoso G.C. 2021 The allometry of

sound frequency bandwidth in songbirds. *Am Nat* **197**, 607-614.

(doi:10.1086/713708).

14.   Krishnan A. 2023 Biomechanics illuminates form-function relationships in bird bills.

*J Exp Biol* **226**, 1-13. (doi:10.1242/jeb.245171).

15.   Palacios M.G., Tubaro P.L. 2000 Does beak size affect acoustic frequencies in

woodcreepers? *Condor* **102**, 553-560. (doi:10.1093/condor/102.3.553).

16.   Huber S.K., Podos J. 2006 Beak morphology and song features covary in a population

of Darwin's finches (*Geospiza fortis*). *Biol J Linn Soc* **88**, 489-498.

(doi:10.1111/j.1095-8312.2006.00638.x).

17.   Demery A.-J.C., Burns K.J., Mason N.A. 2021 Bill size, bill shape, and body size

constrain bird song evolution on a macroevolutionary scale. *J Ornithol* **138**, 1-11.

(doi:10.1093/ornithology/ukab011).

18.   Mason N.A., Burns K.J., Tobias J.A., Claramunt S., Seddon N., Derryberry E.P. 2017

Song evolution, speciation, and vocal learning in passerine birds. *Evolution* **71**, 786-

796. (doi:10.1111/evo.13159).

19.   Friis J.I., Sabino J., Santos P., Dabelsteen T., Cardoso G.C., Jennions M.D. 2022

Ecological adaptation and birdsong: how body and bill sizes affect passerine sound

frequencies. *Behav Ecol* **33**, 798-806. (doi:10.1093/beheco/arac042).

20. Freckleton R.P. 2010 Dealing with collinearity in behavioural and ecological data: model averaging and the problems of measurement error. *Behav Ecol Sociobiol* **65**, 91-101. (doi:10.1007/s00265-010-1045-6).

21. Tattersall G.J., Arnaout B., Symonds M.R.E. 2017 The evolution of the avian bill as a thermoregulatory organ. *Biol Rev* **92**, 1630-1656. (doi:10.1111/brv.12299).

22. Greenberg R., Cadena V., Danner R.M., Tattersall G.J. 2012 Heat loss may explain bill size differences between birds occupying different habitats. *PLoS One* **7**, e40933. (doi:10.1371/journal.pone.0040933).

23. Dubiner S., Meiri S. 2022 Widespread recent changes in morphology of Old World birds, global warming the immediate suspect. *Glob Ecol Biogeogr* **31**, 791-801. (doi:10.1111/geb.13474).

24. McQueen A., Klaassen M., Tattersall G.J., Atkinson R., Jessop R., Hassell C.J., Christie M., Victorian Wader Study Group, Australasian Wader Studies Group, Symonds M.R.E. 2022 Thermal adaptation best explains Bergmann's and Allen's Rules across ecologically diverse shorebirds. *Nat Commun* **13**, 4727. (doi:10.1038/s41467-022-32108-3).

25. Grant P.R., Grant B.R. 2006 Evolution of character displacement in Darwin's finches. *Science* **313**, 224-226. (doi:10.1126/science.1128374).

26. Podos J. 2001 Correlated evolution of morphology and vocal signal structure in Darwin's fnches. *Nature* **409**, 185-188. (doi:10.1038/35051570).

27. Podos J., Nowicki S. 2004 Beaks, adaptation, and vocal evolution in Darwin's finches. *Bioscience* **54**, 501-510. (doi:10.1641/0006-3568(2004)054[0501:BAAVEI]2.0.CO;2).

28.  Mason N.A., Burns K.J. 2015 The effect of habitat and body size on the evolution of

vocal displays in Thraupidae (tanagers), the largest family of songbirds. *Biol J Linn*

*Soc* **114**, 538-551. (doi:10.1111/bij.12455).

29.  Mikula P., Valcu M., Brumm H., Bulla M., Forstmeier W., Petruskova T.,

Kempenaers B., Albrecht T. 2021 A global analysis of song frequency in passerines

provides no support for the acoustic adaptation hypothesis but suggests a role for

sexual selection. *Ecol Lett* **24**, 477-486. (doi:10.1111/ele.13662).

30.  Derryberry E.P., Seddon N., Derryberry G.E., Claramunt S., Seeholzer G.F.,

Brumfield R.T., Tobias J.A. 2018 Ecological drivers of song evolution in birds:

disentangling the effects of habitat and morphology. *Ecol Evol* **8**, 1890-1905.

(doi:10.1002/ece3.3760).

31.  Platt J.R. 1964 Strong inference: certain systematic methods of scientific thinking

may produce much more rapid progress than others. *Science* **146**, 347-353.

(doi:10.1126/science.146.3642.347).

32.  Revell L.J., Harmon L.J. 2022 *Phylogenetic Comparative Methods in R*. Princeton

University Press.

33.  Marquet P.A., Fernández M., Navarette A.A., Valdovinos C. 2004 Diversity

emerging: towards a deconstruction of biodiversity patterns. In *Frontiers of*

*Biogeography: New Directions in the Geography of Nature* (eds. Lomolino M.V.,

Heaney L.R.), pp. 191-209. Sunderland, Sinauer Associates.

34.  Higgins P.J., Ford H.A., Christidis L. 2008 Family Meliphagidae. In *Handbook of the*

*Birds of the World: Penduline-tits to Shrikes* (eds. del Hoyo J., Elliott A., Christie

D.A.), pp. 498–691. Barcelona, Spain, Lynx Edicions.

35.  Slater P. 1986 The cultural transmission of bird song. *Trends Ecol Evol* **1**, 94-97.

36.    Hardt B., Benedict L. 2020 Can you hear me now? A review of signal transmission

634            and experimental evidence for the acoustic adaptation hypothesis. *Bioacoustics* **30**,

635            716-742. (doi:10.1080/09524622.2020.1858448).

636    37.    Dunning J.B. 2008 *CRC Handbook of Avian Body Masses*. Taylor & Francis Group,

637            LLC.

638    38.    Marki P.Z., Kennedy J.D., Cooney C.R., Rahbek C., Fjeldsa J. 2019 Adaptive

639            radiation and the evolution of nectarivory in a large songbird clade. *Evolution* **73**,

640            1226-1240. (doi:10.1111/evo.13734).

641    39.    Friedman N.R., Miller E.T., Ball J.R., Kasuga H., Remes V., Economo E.P. 2019

642            Evolution of a multifunctional trait: shared effects of foraging ecology and

643            thermoregulation on beak morphology, with consequences for song evolution. *Proc*

644            *Royal Soc B* **286**, 20192474. (doi:10.1098/rspb.2019.2474).

645    40.    Hill S.D., Amiot C., Ludbrook M.R., Ji W. 2015 Seasonal variation in the song

646            structure of tui (*Prosthemadera novaeseelandiae*). *N Z J Ecol* **39**, 110-115.

647    41.    Jurisevic M.A., Sanderson K.J. 2016 The vocal repertoires of six honeyeater

648            (Meliphagidae) species from Adelaide, South Australia. *Emu* **94**, 141-148.

649            (doi:10.1071/mu9940141).

650    42.    O'Meara B.C., Ané C., Sanderson M.J., Wainwright P.C. 2006 Testing for different

651            rates of continuous trait evolution using likelihood. *Evolution* **60**, 922-933.

652            (doi:10.1111/j.0014-3820.2006.tb01171.x).

653    43.    R Core Team. 2013 R: a language and environment for statistical computing.

654    44.    Araya-Salas M., Smith-Vidaurre G. 2017 warbleR: an R package to streamline

655            analysis of animal acoustic signals. *Methods Ecol Evol* **8**, 184-191.

656            (doi:10.1111/2041-210x.12624).

45.    Araya-Salas M., Smith-Vidaurre G., Webster M. 2017 Assessing the effect of sound

file compression and background noise on measures of acoustic signal structure.

*Bioacoustics* **28**, 1-17. (doi:10.1080/09524622.2017.1396498).

46.    BirdLife International, HBW. 2018 Bird species distribution maps of the world.

(doi:http://datazone.birdlife.org/species/requestdis.).

47.    Pearse W.D., Morales-Castilla I., James L.S., Farrell M., Boivin F., Davies T.J. 2018

Global macroevolution and macroecology of passerine song. *Evolution* **72**, 944-960.

(doi:10.1111/evo.13450).

48.    Cardoso G.C. 2013 Using frequency ratios to study vocal communication. *Anim*

*Behav* **85**, 1529-1532. (doi:10.1016/j.anbehav.2013.03.044).

49.    Boncoraglio G., Saino N. 2007 Habitat structure and the evolution of bird song: a

meta-analysis of the evidence for the acoustic adaptation hypothesis. *Funct Ecol* **21**,

134-142. (doi:10.1111/j.1365-2435.2006.01207.x).

50.    Zollinger S.A., Podos J., Nemeth E., Goller F., Brumm H. 2012 On the relationship

between, and measurement of, amplitude and frequency in birdsong. *Anim Behav* **84**,

e1-e9. (doi:10.1016/j.anbehav.2012.04.026).

51.    Tobias J.A., Sheard C., Pigot A.L., Devenish A.J.M., Yang J., Sayol F., Neate-Clegg

M.H.C., Alioravainen N., Weeks T.L., Barber R.A., et al. 2022 AVONET:

morphological, ecological and geographical data for all birds. *Ecol Lett* **25**, 581-597.

(doi:10.1111/ele.13898).

52.    Tobias J.A., Sheard C., Seddon N., Meade A., Cotton A.J., Nakagawa S. 2016

Territoriality, social bonds, and the evolution of communal signaling in birds. *Front*

*Ecol Evol* **4**, 1-15. (doi:10.3389/fevo.2016.00074).

53.    Fick S.E., Hijmans R.J. 2017 WorldClim 2: new 1-km spatial resolution climate

surfaces for global land areas. *Int J Climatol* **37**, 4302-4315. (doi:10.1002/joc.5086).

54.    James F.C. 1970 Geographic size variation in birds and its relationship to climate.

683           *Ecol* **51**, 365-390. (doi:10.2307/1935374).

684    55.    Olson V.A., Davies R.G., Orme C.D.L., Thomas G.H., Meiri S., Blackburn T.M.,

685           Gaston K.J., Owens I.P.F., Bennett P.M. 2009 Global biogeography and ecology of

686           body size in birds. *Ecol Lett* **12**, 249-259. (doi:10.1111/j.1461-0248.2009.01281.x).

687    56.    Blackburn T.M., Ruggiero A. 2001 Latitude, elevation and body mass variation in

688           Andean passerine birds. *Glob Ecol Biogeogr* **10**, 245-259. (doi:10.1046/j.1466-

689           822X.2001.00237.x).

690    57.    Hijmans R.J. 2015 Raster: geographic data analysis and modeling. *R package*.

691           (doi:https://cran.r-project.org/package=raster).

692    58.    Hay E.M., McGee M.D., Chown S.L. 2022 Geographic range size and speciation in

693           honeyeaters. *BMC Ecol Evol* **22**, 1-14. (doi:10.1186/s12862-022-02041-6).

694    59.    Paradis E., Claude J., Strimmer K. 2004 APE: analyses of phylogenetics and

695           evolution in R language. *J Bioinform* **20**, 289-290.

696           (doi:10.1093/bioinformatics/btg412).

697    60.    Revell L.J. 2012 phytools: an R package for phylogenetic comparative biology (and

698           other things). *Methods Ecol Evol* **3**, 217-223. (doi:10.1111/j.2041-

699           210X.2011.00169.x).

700    61.    Textor J., van der Zander B., Gilthorpe M.S., Liskiewicz M., Ellison G.T. 2016

701           Robust causal inference using directed acyclic graphs: the R package 'dagitty'. *Int J*

702           *Epidemiol* **45**, 1887-1894. (doi:10.1093/ije/dyw341).

703    62.    Arif S., MacNeil M.A. 2022 Predictive models aren't for causal inference. *Ecol Lett*

704           **25**, 1741-1745. (doi:10.1111/ele.14033).

705    63.    Goodhue D.L., Lewis W., Thompson R. 2017 A multicollinearity and measurement

706           error statistical blind spot: correcting for excessive false positives in regression and

707           PLS. *MIS Quarterly* **41**, 667-684. (doi:https://www.jstor.org/stable/26635007).

708    64.    Grafen A. 1989 The phylogenetic regression. *Philos Trans R Soc Lond B Biol Sci* **326**,

709           119-157.

710    65.    Martins E.P., Hansen T.F. 1997 Phylogenies and the comparative method: a general

711           approach to incorporating phylogenetic information into the analysis of interspecific

712           data. *Am Nat* **149**, 646-667.

713    66.    Pinheiro J., Bates D., DebRoy S., Sarkar D., Team R.C. 2013 nlme: linear and

714           nonlinear mixed effects models.

715    67.    Revell L.J. 2010 Phylogenetic signal and linear regression on species data. *Methods*

716           *Ecol Evol* **1**, 319-329.

717    68.    Dinnage R., Skeels A., Cardillo M. 2020 Spatiophylogenetic modelling of extinction

718           risk reveals evolutionary distinctiveness and brief flowering period as threats in a

719           hotspot plant genus. *Proc Royal Soc B* **287**, 20192817. (doi:10.1098/rspb.2019.2817).

720    69.    GBIF. 2020 GBIF occurrence download (13 August 2020).

721           https://doi.org/10.15468/dl.ee29jh.

722    70.    eBird. 2020 eBird basic dataset. Version: EBD_relSep-2020. Ithaca: Cornell Lab of

723           Ornithology.

724    71.    Martins T.G., Simpson D., Lindgren F., Rue H. 2013 Bayesian computing with

725           INLA: new features. *Comput Stat Data Anal* **67**, 68-83.

726           (doi:10.1016/j.csda.2013.04.014).

727    72.    Rue H., Martino S. 2007 Approximate Bayesian inference for hierarchical Gaussian

728           Markov random field models. *J Stat Plan Inference* **137**, 3177-3192.

729           (doi:10.1016/j.jspi.2006.07.016).

73.   Felsenstein J. 1985 Phylogenies and the comparative method. *Am Nat* **125**, 1-15.

(doi:10.1086/284325).

74.   Hansen T.F. 1997 Stabilising selection and the comparative analysis of adaptation.

*Evolution* **51**, 1341-1351. (doi:10.2307/2411186).

75.   Butler M.A., King A.A. 2004 Phylogenetic comparative analysis: a modeling

approach for adaptive evolution. *Am Nat* **164**, 683-695. (doi:10.1086/426002).

76.   Clavel J., Escarguel G., Merceron G. 2015 mvMORPH: an R package for fitting

multivariate evolutionary models to morphometric data. *Methods Ecol Evol* **6**, 1311-

1319. (doi:10.1111/2041-210x.12420).

77.   Gelman A., Carlin J.B., Stern H.S., Dunson D.B., Vehtari A., Rubin D.B. 2013

*Bayesian Data Analysis*, CRC press.

78.   Carpenter B., Gelman A., Hoffman M.D., Lee D., Goodrich B., Betancourt M.,

Brubaker M., Guo J., Li P., Riddell A. 2017 Stan: a probabilistic programming

language. *J Stat Softw* **76**, 1-32. (doi:10.18637/jss.v076.i01).

79.   Bürkner P.-C. 2017 brms: an R package for Bayesian multilevel models using Stan. *J*

*Stat Softw* **80**, 1-28. (doi:10.18637/jss.v080.i01).

80.   Hoffman M.D., Gelman A. 2014 The No-U-Turn Sampler: adaptively setting path

lengths in Hamiltonian Monte Carlo. *J Mach Learn Res* **15**, 1593-1623.

(doi:10.5555/2627435.2638586).

81.   Derryberry E.P. 2009 Ecology shapes birdsong evolution: variation in morphology

and habitat explains variation in white-crowned sparrow song. *Am Nat* **174**, 24-33.

(doi:10.1086/599298).

82.   Sebastianelli M., Lukhele S.M., Nwankwo E.C., Hadjioannou L., Kirschel A.N.G.

2022 Continent-wide patterns of song variation predicted by classical rules of

biogeography. *Ecol Lett* **25**, 2448–2462. (doi:10.1111/ele.14102).

83.   Albright T.P., Mutiibwa D., Gerson A.R., Smith E.K., Talbot W.A., O'Neill J.J.,

McKechnie A.E., Wolf B.O. 2017 Mapping evaporative water loss in desert

passerines reveals an expanding threat of lethal dehydration. *Proc Natl Acad Sci USA*

**114**, 2283-2288. (doi:10.1073/pnas.1613625114).

84.   Keith D.A. 2017 *Australian Vegetation*. Cambridge University Press.

85.   Frappell P.B., Hinds D.S., Boggs D.F. 2001 Scaling of respiratory variables and the

breathing pattern in birds: an allometric and phylogenetic approach. *Physiol Biochem*

*Zool* **74**, 75-89. (doi:10.1086/319300).

86.   Raftery A. 2013 Avian anaesthesia. *In Practice* **35**, 272-278. (doi:10.1136/inp.f2861).

87.   Dial K.P., Greene E., Irschick D.J. 2008 Allometry of behavior. *Trends Ecol Evol* **23**,

394-401. (doi:10.1016/j.tree.2008.03.005).

88.   Norman J.A., Rheindt F.E., Rowe D.L., Christidis L. 2007 Speciation dynamics in the

Australo-Papuan *Meliphaga* honeyeaters. *Mol Phylogenet Evol* **42**, 80-91.

(doi:10.1016/j.ympev.2006.05.032).

89.   Gardner J.L., Peters A., Kearney M.R., Joseph L., Heinsohn R. 2011 Declining body

size: a third universal response to warming? *Trends Ecol Evol* **26**, 285-291.

(doi:10.1016/j.tree.2011.03.005).

90.   Lindenmayer D.B., Blanchard W., Westgate M.J., Foster C., Banks S.C., Barton P.,

Crane M., Ikin K., Scheele B.C. 2019 Novel bird responses to successive, large-scale,

landscape transformations. *Ecol Monogr* **89**, e01362. (doi:10.1002/ecm.1362).

91.   Bovo A.A.A., Ferraz K.M.P.M.B., Magioli M., Alexandrino E.R., Hasui É., Ribeiro

M.C., Tobias J.A. 2018 Habitat fragmentation narrows the distribution of avian

functional traits associated with seed dispersal in tropical forest. *Perspect Ecol*

*Conserv* **16**, 90-96. (doi:10.1016/j.pecon.2018.03.004).

 **FIGURE CAPTIONS**

**Figure 1.** Morphological diversity of honeyeater body sizes and beak shapes. Images are of

the: crow honeyeater[1] (a; *Gymnomyza aubryana*), barred honeyeater[1] (b; *Gliciphila*

*undulata*), New Holland honeyeater[2] (c; *Phylidonyris novaehollandiae*), tūī[1] (d;

*Prosthemadera novaeseelandiae*), white-eared honeyeater[3] (e; *Nesoptilotis leucotis*), yellow-

faced honeyeater[1] (f; *Caligavis chrysops*), crescent honeyeater[1] (g; *Phylidonyris*

*pyrrhopterus*), noisy miner[2] (h; *Manorina melanocephala*), eastern spinebill[1] (i;

*Acanthorhynchus tenuirostris*), New Caledonian myzomela[1] (j; *Myzomela caledonica*),

white-plumed honeyeater[1] (k; *Ptilotula penicillata*), New Caledonian friarbird[1] (l; *Philemon*

*diemenensis*), dusky myzomela[2] (m; *Myzomela obscura*), dark brown honeyeater[1] (n;

*Lichmera incana*), yellow wattlebird[1] (o; *Anthochaera paradoxa*), yellow honeyeater[2] (p;

*Stomiopera flava*), bell miner[3] (q; *Manorina melanophrys*), white-naped honeyeater[3] (r;

*Melithreptus lunatus*), white-fronted chat[2] (s; *Epthianura albifrons*)*,* spiny-cheeked

honeyeater[3] (t; *Acanthagenys rufogularis*). Image size for each species reflects relative size of

the species. Image credits: [1]Matthias Dehling, [2]Eleanor Hay, [3]Steven Chown.

**Figure 2.** Phylogeny containing the 163 honeyeater species for which we obtained song

variables for and used in analysis. Branches are coloured according to peak frequency. Size

of bars at the tips of the phylogeny represent body size (mass in grams) and are coloured

according to open (white) and closed (black) habitat types. Four examples of honeyeater song

spectrograms are included: blue-faced honeyeater (*Entomyzon cyanotis*; xeno-canto accession

number = XC439288; body size = 105.26 g), singing honeyeater (*Gavicalis virescens*; xeno-

canto accession number = XC334251; body size = 23.33 g), noisy friarbird (*Philemon*

*corniculatus*; xeno-canto accession number = XC287056; body size = 100.71 g), and eastern

spinebill (*Acanthorhynchus tenuirostris;* xeno-canto accession number = XC490800; body

size = 10.44 g). Birds are illustrations by Eleanor Hay.

**Figure 3.** Conceptual diagram explaining the underlying interactions between environmental

and phenotypic variables and their influence on song. Song spectrogram displays frequency

(kHz) over time (s) and is an example from the New Zealand bellbird (*Anthornis melanura*;

xeno-canto accession number = XC153005). The three song variables considered in this

study are indicated with annotations; peak frequency represents the frequency of maximum

amplitude across the entire song, bandwidth is the difference between maximum and

minimum frequency across the song, and pace is the number of notes over song duration.

Beak shape shows the beak variation in honeyeaters, the top is the species with the lowest

value for beak PC2 (Arfak honeyeater; *Melipotes gymnops*) and the bottom is the species

with the highest value for PC2 (Eastern spinebill; *Acanthorhynchus tenuirostris*). Illustrations

by Eleanor Hay.

**Figure 4.** Results from phylogenetic generalised least squares regression models (a) and the

spatiophylogenetic models (c-d), testing the influence of environmental and morphological

traits on song variables. Separate models were run to test the influence on peak frequency,

bandwidth, and pace, and are indicated in the legends. For the PGLS results (a), the predictor

variables have been scaled so the effect sizes are comparable, full model results are available

in Table S4. Maps in panels b-c display the mean spatial random effect for each song variable

which was extracted from the final spatiophylogenetic model. Density plots show the

marginals of the fixed effects in the model, with the mean and 95% credible intervals.

Predictor variables which were identified to have significant effects are highlighted in bold,

and full model results are available in Table S5.

**Figure 5.** Distribution of AIC values from mvMORPH models (a) when testing the effect of

open and closed habitat types on peak frequency, bandwidth, and pace. Note that models

were conducted over 100 stochastic character mapping of open and closed habitats on the

phylogeny. Models compared were single rate Brownian motion (BM1, n = 100), two rate

Brownian motion (BMM, n = 100), single rate Ornstein-Uhlenbeck (OU1, n = 100) and two

rate Ornstein Uhlenbeck (OUM, n = 100). Full model results are available in Table S6. Panel

b displays outcomes of Bayesian phylogenetic regression models (a), testing the influence of

environmental and morphological traits on the variance of peak frequency, bandwidth, and

pace. Models are indicated with the legend and full model results are available in Table S7.

855 **Figure 1.**

Blue-faced honeyeater (*Entomyzon cyanotis*)

Singing honeyeater (*Gavicalis virescens*)

Peak frequency (Hz)

947.5      7752

□ Open habitat    ■ Closed habitat

Noisy friarbird (*Philemon corniculatus*)

Eastern spinebill (*Acanthorhynchus tenuirostris*)

862 **Figure 2.**

869 **Figure 3.**

879 **Figure 4.**

**Figure 5.**

899