Body size shapes song in honeyeaters

Eleanor M. Hay[1]*, Matthew D. McGee[1], Craig R. White[1,2], Steven L. Chown[1]

[1] School of Biological Sciences, Monash University, Melbourne, VIC 3800, Australia

[2] Centre for Geometric Biology, School of Biological Sciences, Monash University,

Melbourne, VIC 3800, Australia

*Corresponding author; email: eleanor.m.hay@outlook.com

ABSTRACT

Birdsongs are among the most distinctive animal signals. Their evolution is thought to be

shaped simultaneously by habitat structure and by the constraints of morphology. Habitat

structure affects song transmission and detectability, thus influencing song (the acoustic

adaptation hypothesis), while body size and beak size and shape necessarily constrain song

characteristics (the morphological constraint hypothesis). Yet, support for the acoustic

adaptation and morphological constraint hypotheses remains equivocal, and their

simultaneous examination is infrequent. Using a phenotypically diverse Australasian bird

clade, the honeyeaters (Aves: Meliphagidae), we jointly examine predictions of these two

hypotheses. We find that body size constrains song frequency and pace in honeyeaters, while

beak shape constrains the rate of song pace evolution. Although habitat type and

environmental temperature influence aspects of song, that influence is indirect, via effects of

environmental variation on body size. Our results demonstrate that morphology has an

overwhelming influence on birdsong, in support of the morphological constraint hypothesis,

with the environment playing a secondary role generally via body size rather than habitat

structure. These results suggest that changing body size, a consequence of both global effects

such as climate change, and local effects such as habitat transformation, will substantially

influence the nature of birdsong.

Keywords: Acoustic adaptation; morphological constraint; song evolution; spatial covariance;

evolutionary constraint; allometry

INTRODUCTION

Birdsongs are acoustic signals influencing survival, reproduction, and reproductive isolation [1,2], thus shaping the evolution of avian diversity [3-5]. Song is thought to be mediated by two major factors, one external, one internal. First, signal detectability, which is influenced by the physical properties of the signalling environment [6,7] is thought to be a major external factor. Consequently, habitat characteristics are proposed to influence the acoustic properties of bird song. The acoustic adaptation hypothesis (AAH) [6] specifically makes the prediction that because low frequency sounds transmit further than high frequency sounds, birds (and other animals) residing in dense habitats, such as forests, will produce lower frequency sounds than their counterparts dwelling in more open environments [8,9].

Second, birdsong is also subject to constraints imposed by morphology. Body size is one of the most well understood acoustic constraints [10-12] through its effect on the size of the vocal tract and sound producing structures, which places a limit on the lowest frequency an animal can effectively produce [13]. Allometric constraints of acoustic signals are unambiguous: larger animals produce lower frequency signals [1,10]. In birds, beak shape influences acoustic signals too. The avian bill is used in coordination with tongue and vocal tract movements to modify sounds [14] and is thought to act as a final modulator of song, influencing both its frequency and temporal components [15-19]. These affects are complicated, however, by the allometric relationship between body size and beak size [19-21], and by interactions among the biotic and abiotic factors that are thought to influence acoustic signalling. Specifically, beak shape and body size also affect thermoregulation and are influenced by environmental temperatures [22-25]. In consequence, what appears to be a straightforward morphological constraint hypothesis [10-12], is more complicated.

Such complexity, caused by the effects of ecological selection on signal production,

characterises both the acoustic adaptation and morphological constraint hypotheses. For

example, environmental conditions not only influence body size and beak shape, but they

also affect habitat productivity, complexity, and food availability, which, because of dietary

preferences, influence beak shape [14,26] and, presumably, song characteristics [14,27,28].

This may account for mixed outcomes when studies focus on just one of the hypotheses, with

particularly contradictory results for the AAH [29,30]. Where morphology and habitat have

been considered simultaneously, habitat was found to have little effect on song, body size

influences frequency, and beak shape influences the temporal and performance components

of song [31], but understanding of the direct and indirect influences of body size, especially

in relation to the environment, remains incomplete. Therefore, although birds are one of the

most diverse vertebrate clades in which acoustic signalling plays such a conspicuous role in

sexual signalling and speciation [4,5], the manner in which multiple and sometimes non-

independent factors influence their signals remains opaque [18,31], and the avian songbook

thus not fully interpretable.

Here, we therefore set out to uncover the effects of both habitat and morphology on

bird song, while considering underlying interactions and correlations between variables. We

address this complexity explicitly in a strong inference framework [32] and use a suite of

phylogenetic comparative methods [33] to explore it with simultaneous investigation of the

predictions of the major hypotheses. We focus on a large and diverse yet monophyletic group

of birds to do so, recognising the power that such focus on a geographically restricted,

diverse single clade can bring [34]. The honeyeaters (Aves: Meliphagidae) are a clade (~190

species) of passerine birds which have been historically confined to Australasia [35]. They

are oscine passerines that learn their songs and are therefore influenced by cultural

transmission. Learned songs are more subject to change over generations than innate songs

and are also thought to be impacted substantially by the environment because they have

greater plasticity [18,36]. Honeyeaters exhibit exceptional diversity in characters thought to

influence song: body sizes range from <10 g to 350 g [37], beaks vary from short for feeding

on insects to long and decurved for feeding on nectar [38,39], and species inhabit a wide

variety of biomes from arid deserts to the wet tropics [35] (Fig. 1). This extensive diversity in

body size, beak shape, and environments are ideal for disentangling the factors that shape

birdsong (Fig. 2).

Specifically, we compile a dataset of song variables (peak frequency, song pace, and

frequency bandwidth), morphological traits (body size and beak shape), and environmental

variables (habitat type, temperature, elevation, and precipitation) for 163 species of

honeyeater (Table S1). We first use a comparative phylogenetic approach consisting of

phylogenetic multiple regression analyses to assess the factors influencing song traits.

Models are also repeated in a Bayesian framework, and the influence of both phylogenetic

and spatial covariance are considered. We next adopt an evolutionary approach, where we

first examine the influence of open and closed habitats on song evolution. This is followed by

phylogenetic Bayesian regression analysis in which we model the effect of continuous

morphological and environmental variables on the phylogenetic variance of song, analogous

to the Brownian rate parameter [40], to obtain further insight of constraints on rates of song

evolution.

METHODS

Honeyeater song data

Song recordings were sourced from xeno-canto (https://xeno-canto.org/), the Macaulay

Library (Cornell Laboratory of Ornithology, https://www.macaulaylibrary.org/), the

Australian National Wildlife Collection sound archive (ANWC), and the Avian Vocalization

Center (AVoCet; https://avocet.integrativebiology.natsci.msu.edu/). We focussed exclusively

on male song and did not consider recordings of female and juvenile song, or other types of

vocalisations such as calls. Where available, expert classifications on recording type available

in databases were followed (e.g. "song" or "call") and the best quality recordings available

for each species were prioritised (recordings scored as "A" or "B" for xeno-canto, recordings

ranked four stars or more for Macaulay Library). However, lower quality recordings were

considered when no other recordings were available.

131   After downloading, recordings were converted from .mp3 files to .wav files for

analysis in R [41] using an online converter (www.online-audio-converter.com) with a

sampling rate of 44.1 kHz. To maximize the independence of our samples, only one example

of song was analysed from each recording. Recordings made at different geographic locations

were assumed to be of different individual birds, and recordings that did not include a clear

song example were discarded. Spectrograms were generated for 866 individuals from 163

honeyeater species (1-36 recordings per species; ~85% of honeyeaters sampled; Table S2)

using Raven Pro v.1.6 (Cornell Laboratory of Ornithology) with a window size of 256

samples. Following previous studies [e.g. 31,39,42], for each song up to 30 notes were

selected and the following five song variables were extracted: Peak frequency (the frequency

at maximum amplitude across the entire song recording, Hz), maximum frequency (upper

frequency bound of the highest pitched note in the song, Hz), minimum frequency (lower

frequency bound of the lowest pitched note in the song, Hz), song bandwidth (maximum

frequency – minimum frequency), and song pace (number of notes/song duration). Peak

frequency of each song was estimated using the peak frequency function available in the

measurements of Raven Pro (Cornell Laboratory of Ornithology). The mean value for each

song variable was then calculated for each species. All song variables were then $\log_{10}$-

transformed prior to statistical analysis to meet parametric assumptions of normality and

homogeneity of variance, but also because logarithmic scales of sound frequency correspond

to how birds perceive and modulate sound [43].

All five song variables are correlated to some extent, particularly the multiple

measures of frequency (minimum frequency, maximum frequency, and peak frequency; Fig.

S1). Out of the three frequency variables, peak frequency is the most essential for signal

transmission [8], and has the most evidence in support of the acoustic adaptation hypothesis

[44]. Additionally peak frequency is thought to be less sensitive to background noise and

recording conditions than minimum and maximum frequencies [45]. For these reasons, we

focus on peak frequency over maximum frequency and minimum frequency and assess the

effect of predictor variables on peak frequency, pace, and bandwidth (Fig. 2).

Trait and environmental data

To consider the influence of morphology on song measures of body size [37,46] and beak

width, depth, and length [46] were included (Table S1). As we are only using male song data,

we only considered male beak measurements. Raw measurements of beak length, width, and

depth were extracted from AVONET [46] and the mean value was obtained for each species.

All morphological variables were log transformed. From here, beak length, width, and depth

were then subjected to principal component analysis (PCA) using the '*prcomp*' function in R

[41]. The first principal component (PC1) accounted for 92.53% of the cumulative variation

and describes bill size. Whereas PC2 accounted for 5.46% of the cumulative variation and

describes a contrast between short and thick bills vs. long and narrow bills (visualised in Fig.

2; Table S3). A phylogenetic regression of beak PC1 on body size demonstrated that PC1 is

strongly correlated with body size (PGLS regression $\lambda = 0.66$, $\beta = 4.13 \pm 0.22$, $P > 0.0001$,

Table S4). Thus, PC2 was used to represent beak shape in analysis.

To investigate the AAH, habitat density was first classified using a 2-point scale, and

habitat type was classified as dense or open/semi-open habitats. We followed previous

studies which have classified birds globally into dense, semi-open, and open habitat types

[46]. As we are interested in comparing the differences between dense and open habitats, we

grouped honeyeater species with open (N_{open} = 3) and semi-open ($N_{semi-open}$ = 79) habitat

types together ($N_{open+semi-open}$ = 82) and compared them to species from dense habitats (N_{dense}

= 81). Because body size and beak shape are influenced by the environment [22-25,39],

appropriate bioclimatic variables were extracted from the geographic range of each species

and included in analysis. Honeyeater breeding range maps from BirdLife International [47]

were used to sample values of mean annual temperature, annual precipitation, and elevation

from the BioClim database of present-day climatic conditions at 1 km^2 resolution [48], using

the 'raster' package [49] in R [41]. The mean value of each variable was then calculated for

each species.

Phylogeny

Trait values of closely related species are expected to be more similar than that of distantly

related species. Therefore, to comparatively analyse relationships between habitat,

morphology, and song while accounting for phylogenetic history, we used a recently

reconstructed honeyeater phylogeny from Hay et al. [50]. This phylogeny was constructed

using a combination of nuclear genes, mitochondrial genes, and topological constraints from

previous phylogenomic studies. The phylogeny was trimmed, using the '*drop.tip*' function

from the 'ape' package [51], to the 163 species for which we had song data and was used as a

phylogenetic effect in all analysis.

To visualise correlations between the morphological constraint hypothesis and the

acoustic adaptation hypothesis, song variables, morphological traits, and habitat type were

plotted on the phylogeny (Fig. 3) using the using the functions '*plotTree.wBars*' and

'*contMap*' from the R package 'phytools' [52].

Analyses

We used two approaches to assess the acoustic adaptation hypothesis and the morphological

constraint hypothesis. The first is a typical comparative approach using regression analysis to

identify relationships, whereas in the second an evolutionary constraint method is adopted.

When investigating hypotheses in a strong inference contest, several biases might

arise. To account for these potential biases a directed acyclic graph [DAG; 53] was created to

visualise causal relationships and the backdoor criterion was applied to determine the

covariates required to assess both hypotheses while considering interactions between the two

hypotheses [54] (Fig. 2). In this case, two confounding biases exist in which a predictor

variable affects both another predictor variable and the response variable of interest [54]. The

first bias exists between body size and beak shape, which are two strongly correlated traits

that have similar predictions for how they could each influence song frequency [19].

Statistically it can be difficult to disentangle effects such interrelated traits might have and

there is a possibility that any effect of beak shape on song might not actually be attributed to

that variable but could in fact be a statistical artefact [19-21]. Thus, to overcome this, the

effect of body size needs to be controlled and body size must also be included as a predictor

in analysis. The second bias exists between the interrelated effects of phenotypic traits and

the environment. Environment has the potential to affect beak shape and body size, as well as

song, and therefore this needs to be controlled by including environmental predictor variables

in the model as well. Thus, in this scenario, to address both the acoustic adaptation

hypothesis and the morphological constraint hypothesis body size, beak shape, habitat type,

temperature, precipitation, and elevation must all be included as predictor variables in

analysis (Fig. 2; DAG model available at dagitty.net/mz3jqZU). To further consider the direct

effects of environmental variables on body size and beak shape we also use models where we

explicitly test the effect of precipitation, elevation, and temperature on both body size and

beak shape.

We first approach the acoustic adaptation hypothesis and the morphological constraint

hypothesis from a typical comparative approach to determine the drivers of song.

Phylogenetic generalised least squares regression (PGLS; [55,56] was used to determine the

influence of predictor variables on song. PGLS regression tests were run using the 'ape'

[v5.3; 51] and 'nlme' [57] packages in R [41]. An estimated lambda parameter was used to

control for the amount of phylogenetic effect in the model residuals [58].

Spatiophylogenetic modelling [59] was then used to estimate the spatial and

phylogenetic structure in honeyeater song, and to examine the contributions of body size,

beak shape, and environment to variation in song. To model the spatial distributions of each

species, honeyeater point occurrence records from Hay et al. [50] were used. Our analysis

included a total of 253,742 honeyeater point occurrence records. The spatiophylogenetic

model was implemented in the R package 'INLA' which uses an integrated nested Laplace

approximation to estimate the joint posterior distribution of model parameters [60,61]. INLA

is an approach for latent Gaussian Markov random field models, providing a significantly

faster alternative to Markov chain Monte Carlo [61]. Prior to analysis, all predictor variables

were standardised by subtracting the mean and dividing by the standard deviation. For the

phylogenetic effect, INLA requires a phylogenetic precision matrix which is the inverse of a

phylogenetic covariance matrix. Prior to inverting, the phylogenetic covariance matrix was

standardised by dividing by its determinant raised to the power of $1/N_{species}$ [59]. To model

the spatial effect across the entire landscape, we followed a previous study that has used this

approach to model speciation patterns in honeyeaters [50], and used a spatial mesh consisting

of 859 vertices, which was constructed over honeyeater point occurrence records and

averaged across the spatial random fields of each occurrence record for each species. This

approach integrates a spatial random field across each species' distribution. Default priors

were used for both the phylogenetic and spatial effects. The significance of effects in the

models were based on whether the 95% credible intervals of the effect size overlapped with

zero.

We then approach the acoustic adaptation hypothesis and the morphological

constraint hypothesis from an evolutionary perspective to assess the influence of potential

constraints by examining rates of trait evolution using Brownian motion and Ornstein-

Uhlenbeck models [40,62-64]. We fit multivariate Brownian motion (BM) and Ornstein-

Uhlenbeck (OU) models in R [41] using mvMORPH [65] to assess whether open and closed

habitat have constrained song evolution. We constructed multivariate Ornstein-Uhlenbeck

models and Brownian motion models with either one rate (no evolutionary relationship

between open and closed habitats) or two rates of evolution (open and closed habitats have

evolved independently) and tested the effect on song pace, bandwidth, peak frequency. We

performed stochastic character mapping of open and closed habitats on the phylogeny using

the *make.simmap* function from the 'phytools' package [52] and generated 100 stochastic

character histories using the tip states on the tree. Model fit was compared with AIC. Despite

being limited to testing the influence of discrete predictors, the mvMORPH approach is

beneficial here as a multivariate framework enables us to determine the influence of open and

closed habitats on peak frequency, bandwidth, and pace altogether.

Finally, we apply phylogenetic Bayesian regression [66,67] to model the effect of

continuous morphological and environmental variables on the phylogenetic variance of song,

analogous to the Brownian rate parameter, to obtain further insight on constraints from an

evolutionary perspective. This approach enables us to examine how continuous predictor

variables such as body size and beak shape may constrain rates of song evolution. Models

were implemented in R with the package 'brms' [68], for Bayesian models using the

probabilistic programming language STAN [66,67], which is a Markov chain Monte Carlo

(MCMC) sampler that uses techniques based on Hamiltonian Monte Carlo [69]. Here, we

estimate the effect of predictor variables on both the mean to confirm earlier PGLS and INLA

results, as well as the variance to examine constraints on rates of trait evolution. All

continuous variables were transformed to have a mean of 0 and a standard deviation of 0.5.

Phylogenetic relatedness of species in our regression analysis was accounted for with a

phylogenetic covariance matrix calculated using the *vcv* function from the R package 'ape'

[51]. Models were run for at least 2,000 MCMC iterations with 4 chains and convergence

was assessed by ensuring the estimated potential scale reduction statistic Rhat reached 1 [67].

Effects were judged as significant if the 95% credible intervals of the effect size did not

overlap with zero.

RESULTS

Multiple PGLS regression models testing the effect of morphological and environmental

variables on song revealed several relationships. Peak frequency ($\beta = -0.25 \pm 0.043$, $P <$

0.0001; model $\lambda = 0.48$, $N = 163$) and pace ($\beta = -0.31 \pm 0.089$, $P < 0.001$; model $\lambda = 0.55$, N

$= 163$) are negatively related to body size, indicating that overall, smaller birds produce

higher frequency sounds and faster paced songs (Fig. 4a; Table S4). Beak shape and habitat

density had no impact on any of the song variables (Fig. 4a; Table S4). By contrast, both

bandwidth ($\beta = -0.021 \pm 0.0065$, $P = 0.002$; model $\lambda = 0.43$, $N = 163$) and pace ($\beta = -0.013 \pm$

0.0066, $P = 0.042$; model $\lambda = 0.55$, $N = 163$) are negatively related to temperature, in which

cooler mean annual temperatures are associated with broader bandwidths (more vocal

variation) and faster paced songs (Fig. 4a; Table S4). Bandwidth is also positively related to

precipitation, but the effect is weak ($\beta = 0.000062 \pm 0.000029$, $P = 0.032$; model $\lambda = 0.43$, N

$= 163$; Fig. 4a; Table S4). Body size in honeyeaters is positively related to precipitation

suggesting that honeyeaters are larger in wetter areas ($\beta = 0.000062 \pm 0.000019$, $P < 0.001$;

model $\lambda = 0.99$, $N = 163$; Table S4). Bivariate PGLS models of habitat type on body size

identified honeyeaters to be larger in closed than open environments ($\beta = 0.052 \pm 0.024$, $P =$

0.034; model $\lambda = 0.99$, $N = 163$; Table S4).

Spatiophylogenetic analysis revealed considerable spatial and phylogenetic structure

to the variation in honeyeater songs and show that song traits are variable across the

distribution of honeyeaters (Fig. 4b-d). Consistent with PGLS models, frequency and pace

are negatively related to body size (Fig. 4b; Fig. 4d; Table S5). Models further confirm the

effects of precipitation and temperature on frequency bandwidth, indicating that species in

warmer and drier areas sing with narrower bandwidths (Fig. 4c). However, when spatial

covariance is considered alongside phylogenetic covariance, the effect of temperature and

pace is reduced and no longer significant (Fig. 4d; Table S5). The effect of precipitation on

body size remains (Table S5).

Model comparisons of multivariate Ornstein-Uhlenbeck models and Brownian motion

models indicate that open and closed habitat types have influenced the evolution of song in

honeyeaters. The best model based on AIC was an Ornstein-Uhlenbeck model with two rates

of evolution (Fig. 5a; Table S6), implying that song has evolved toward different

evolutionary optima in open and closed habitats. Estimated theta values extracted from these

models show that values tend to be lower for closed habitats than for open habitats,

suggesting that lower values of frequency, bandwidth, and pace are selected for in closed

habitats.

Bayesian phylogenetic regressions confirm all results of PGLS models (Table S7).

When modelling the influence of predictor variables on the phylogenetic variance of song,

we find song pace to be constrained by beak shape. This indicates that honeyeaters with

shorter and wider beaks have higher rates of evolution, whereas longer and narrower beaks

constrain the rate of song pace evolution (Fig. 5b; Table S7).

DISCUSSION

In this study we used honeyeaters to test the predictions of two key hypotheses thought to

influence acoustic signals: the morphological constraint hypothesis and the environmental

adaptation hypothesis. We found evidence to support the predictions of the morphological

constraint hypothesis: body size constrains frequency and pace, and beak shape constrains

rate of song pace evolution. The predictions of the environmental adaptation hypothesis were

not fully supported, but temperature and precipitation were found to influence bandwidth,

and we did find some evidence habitat type has influenced the evolution of song. However,

these findings are likely mediated by environmental effects on body size. Overall, we found

that the main constraint of song is body size, and this is ultimately modulated by underlying

environmental effects of precipitation on body size, which has consequences for song

evolution.

Morphological constraints on song

In the case of the morphological constraint hypothesis, we found overwhelming evidence that

body size constrains peak frequency: smaller-sized honeyeaters produce higher frequency

songs. This outcome is congruent with previous findings for other bird groups [18,29-

31,42,70], and likely the outcome of a proximal mechanitic relationship where larger birds

have larger sound-producing organs, such as the syrinx [10-12]. In turn this raises the

question of what factors influence body size. In African tinkerbirds, temperature and latitude

account for variation in body size, ultimately having consequences on song frequency [71].

Among honeyeater species, no evidence exists for latitudinal variation in body size

(Bergmann's rule) [50], but the current analyses show that that honeyeater body size is

negatively related to annual precipitation, indicating that species from wetter areas tend to be

larger than those from drier areas. This could be in part driven by some form of tropical

temperature effect but is also likely the outcome of aridity gradients and honeyeater

responses to them across Australasia. Body size is fundamental to an organism's ability to

maintain homeostasis [24,25], and precipitation is a key factor in thermoregulation [72,73].

Australia has a particularly strong precipitation gradient from the coast to the interior [74].

The influence of such precipitation gradients on body size in honeyeaters is apparent in arid

zone honeyeaters which comprise chats (genus *Epithanura*) and the gibberbird (*Ashbyia*

lovensis). These honeyeater species are relatively small, weighing 9-13.5 g (*Epithanura* spp.)

and 18 g (*Ashbyia lovensis*), respectively [37] (Table S1), and occupy the driest areas across

the study region. Furthermore, this interplay between body size, song frequency, and the

environment could suggest that indirect effects via ecological selection, in this case

precipitation, on body size in honeyeaters can have consequences for signal production.

The morphological constraint from beak shape is not as strong as that for body size

and no direct evidence is found for an influence of beak shape on song. Longer and narrower

beak shapes do, however, constrain the rate of song pace evolution in honeyeaters. Beak size

has been negatively associated with pace in Australian honeyeaters previously [39], and a

wide range of other bird clades including medium ground finches [16], white-crowned

sparrows [70], ovenbirds [31], and tanagers [17]. Beak shape is understood to act as a

mechanical constraint on song and hinder movement [16], the extent of which is underpinned

by beak specialisation for dietary preferences. For example, mechanics behind small and

slender beaks of insectivores prioritise closing velocity for catching insects, whereas curved

and narrow beaks of nectarivores prioritise nectar transfer [14]. Our finding that longer and

narrower beaks constrain rate of song pace evolution in honeyeaters is congruent with these

biomechanical constraints of beak shape.

Numerous past studies have identified beak attributes, prominently beak size, to

impact song frequency [15,16,18,39], but we find no evidence for an impact of beak shape on

peak frequency. Disentangling the effects of beak size and shape on song is complicated by

allometric effects of body size on beak size which could confound the relationship between

beak size and many song features. In the present study we accounted for this allometry by

using the second principal component of beak variables, representing shape, and controlled

for body size in all analysis. The discrepancy between the present study and previous work is

not overly surprising given that we focus on beak shape rather than size. However, only a few

past studies have explicitly acknowledged this statistical concern [19,31] and caution is

required when interpreting findings which might not have fully accounted for body size

because any effect of beak shape on song variables could be mistakenly attributed [19-21].

Beyond statistical concerns, fully understanding the influence of beak shape on song is not

entirely straightforward and requires more attention because behaviours such as adjusting

gape and coordinated movements of the beak with the tongue also alter amplification and

resonance of song [14].

Body size also constrains song pace: larger honeyeaters sing at slower. This negative

effect of body size on song pace is consistent with previous studies of birds [29,31], however,

it has generally been attributed to beak size, and little discussion of the relationship between

body size and pace exists throughout the literature. Certainly, beak shape does constrain pace

in honeyeaters, but there is clearly an effect of body size too. One potential explanation is

that body size influences air intake and output which could determine song pace. Larger birds

have larger lungs and vocal producing organs, but also have greater oxygen consumption, and

take larger and fewer breaths than smaller birds [75] which could explain differences in pace.

Morphological structures such as the glottis which acts as a valve between the lungs and

mouth, and the trachea which connects the syrinx in birds to the lungs, determine air

movement. In most birds the trachea narrows towards the lungs [76], potentially limiting air

movement, and could cause larger lungs take longer to fill and empty, perhaps placing a

constraint on the pace at which birds can sing.

Environmental constraints on song

Empirical evidence for the acoustic adaptation hypothesis has remained mixed and its

predictions are generally not supported [9,30,36,44]. In the present study, no direct evidence

is uncovered for an effect of habitat type on song at a macroevolutionary scale in

honeyeaters. The hypothesised selection imposed by closed habitats is likely not strong

enough to have a macroevolutionary influence, or perhaps open habitats are also subject to

their own constraints on the transmission and reception of signals. Signals produced in open

habitats can experience degradation from wind and environmental attenuation through the

atmosphere, ground, and vegetation [7]. Conversely, any constraint of closed habitats could

possibly be overcome through behaviours such as movement. Honeyeaters in general have

high dispersal ability [38,50] and smaller sized birds could implement behaviours such as

increased movement to compensate if acoustic signals are not being detected [77].

Our evolutionary approach did, however, indicate that habitat type influences the

evolution of song. Here, model comparisons of multivariate Ornstein-Uhlenbeck models and

Brownian motion models indicate song in open and closed habitats has evolved toward

different optima. This was generally found to support the predictions of the acoustic

adaptation hypothesis: lower values for peak frequency, bandwidth, and pace tended to be

preferred in closed habitats. However, analysis revealed that honeyeaters tend to be larger in

closed than open environments. Such underlying relationships suggest that any effect we

uncover for habitat type influencing song is more likely to be driven by underlying variation

in body size, and ultimately provides further support for body size being the primary

constraint of song.

Although little support is uncovered for the AAH, we find some evidence that other

environmental variables influence song in honeyeaters. Initially, song pace is found to be

weakly negatively related to mean annual temperature, however this effect is reduced and is

no longer significant when spatial covariance is considered in analysis, highlighting the

importance of considering spatial covariance alongside phylogenetic covariance in analysis.

Nevertheless, bandwidth is consistently found to be negatively related to mean annual

temperature and positively related to annual precipitation. These findings indicate that

honeyeater species in warmer and drier areas sing with narrower bandwidths or less

frequency variation. Although the impacts of temperature and precipitation on bandwidth

could be related to habitat type, body size is more likely the main influence. As we have

shown, body size in honeyeaters is structured by precipitation, it is understood to place a limit

on the lowest pitched sound a bird can produce, and larger species use wider bandwidths than

smaller species [13].

Conclusion

This study provides support for the morphological constraint hypothesis and firmly

establishes body size as the main constraint of song in honeyeaters, with beak shape

additionally constraining the rate of song pace evolution. Effects of habitat type and any

apparent environmental drivers on song have stronger associations with body size than with

habitat structure *per se*.

These findings have implications for the future of birdsong in a rapidly changing

world. Changing climates are resulting in a reduction in avian body size [24,78], and habitat

loss and transformation include declines of larger sized birds as part of overall changes to

local assemblage composition [79,80]. Our results therefore forecast an increase in higher

pitched songs as body sizes decline. Moreover, beak shapes are also expected to be

influenced by climate change [81], and habitat loss and fragmentation can cause local

extinction of certain beak shapes [79,80], all of which has further implications for song

evolution. Because beak shape influences rates of song pace evolution, ecosystem loss of

certain beak shapes will cause loss of unique song repertoires, and climate-induced changes

in beak shape will likely alter rates of song evolution. Certainly, we demonstrate that

morphology and habitat are deeply intertwined and can jointly influence song. Thus, climate

change and habitat loss will not only have consequences for acoustic signalling and song

evolution, but will also transform the avian songbook.

Acknowledgements

We thank Leo Joseph and Margaret Cawsey at ANWC for assistance with accessing

recordings and acknowledge receipt of media from The Macaulay Library at the Cornell Lab

of Ornithology. Members of the Chown Lab, Melodie McGeoch, and members of the

McGeoch lab provided feedback on aspects of this work.

Funding

EMH was supported by an Australian Government Research Training Program (RTP)

Scholarship, a Monash Graduate Excellence Scholarship (MGES), and a Faculty of Science

Dean's Postgraduate Research Scholarship. SLC was funded by Australian Research Council

Discovery Project DP170101046. MDM was funded by Australian Research Council

Discovery Early Career Research Award DE180101558.

Authors' contributions

EMH and SLC conceptualised the study. Trait and environmental variables were gathered by

EMH. Processing of song recordings and song measurements were completed by EMH.

Analysis was completed by EMH with supervision and input from MDM, CRW, and SLC.

EMH wrote the manuscript with input from SLC, MDM, and CRW. All authors read and

approved the final manuscript.

REFERENCES

1.   Bradbury JW, Vehrencamp SL. 1998 *Principles of Animal Communication*.

Sunderland, Massachusetts, USA: Sinauer.

2.   Catchpole CK, Slater PJB. 2003 *Bird Song: Biological Themes and Variations*. New

York: Cambridge University Press.

3.   Coyne JA, Orr HA. 2004 *Speciation*. Sunderland, Massachusetts USA: Sinauer

Associates.

4.   Tobias JA, Ottenburghs J, Pigot AL. 2020 Avian diversity: speciation,

macroevolution, and ecological function. *Annu Rev Ecol Evol Syst* **51**, 533-60.

(10.1146/annurev-ecolsys-110218-025023)

5.   Price T. 2008 *Speciation in birds*. Colorado: Roberts and Co.

6.   Morton ES. 1975 Ecological sources of selection on avian sounds. *Am Nat* **109**, 17-

34. (doi:10.1086/282971)

7.   Wiley RH, Richards DG. 1978 Physical constraints on acoustic communication in the

atmosphere: implications for the evolution of animal vocalizations. *Behav Ecol*

Sociobiol **3**, 69-94. (10.1007/BF00300047)

8.   Brumm H, Naguib M. 2009 Environmental acoustics and the evolution of bird song.

Adv Study Behav **40**, 1-33.

9.   Ey E, Fischer J. 2009 The "Acoustic Adaptation Hypothesis"—a review of the

evidence from birds, anurans and mammals. *Bioacoustics* **19**, 21-48.

(10.1080/09524622.2009.9753613)

10.   Ryan MJ, Brenowitz EA. 1985 The role of body size, phylogeny, and ambient noise

in the evolution of bird song. *Am Nat* **126**, 87-100. (10.1086/284398)

11.   Bertelli S, Tubaro PL. 2002 Body mass and habitat correlates of song structure in a

primitive group of birds. *Biol J Linn Soc* **77**, 423-30. (10.1046/j.1095-

8312.2002.00112.x)

12.   Ey E, Pfefferle D, Fischer J. 2007 Do age- and sex-related variations reliably reflect

body size in non-human primate vocalizations? A review. *Primates* **48**, 253-67.

(10.1007/s10329-006-0033-y)

13.   Friis JI, Sabino J, Santos P, Dabelsteen T, Cardoso GC. 2021 The allometry of sound

frequency bandwidth in songbirds. *Am Nat* **197**, 607-14. (10.1086/713708)

14.   Krishnan A. 2023 Biomechanics illuminates form-function relationships in bird bills.

J Exp Biol **226**, 1-13. (10.1242/jeb.245171)

15.   Palacios MG, Tubaro PL. 2000 Does beak size affect acoustic frequencies in

woodcreepers? *The Condor* **102**, 553-60. (10.1093/condor/102.3.553)

16.   Huber SK, Podos J. 2006 Beak morphology and song features covary in a population

of Darwin's finches (*Geospiza fortis*). *Biol J Linn Soc* **88**, 489-98. (10.1111/j.1095-

8312.2006.00638.x)

17.   Demery A-JC, Burns KJ, Mason NA. 2021 Bill size, bill shape, and body size

constrain bird song evolution on a macroevolutionary scale. *J Ornithol* **138**, 1-11.

(10.1093/ornithology/ukab011)

18.   Mason NA, Burns KJ, Tobias JA, Claramunt S, Seddon N, Derryberry EP. 2017 Song

evolution, speciation, and vocal learning in passerine birds. *Evolution* **71**, 786-96.

(10.1111/evo.13159)

19.   Friis JI, Sabino J, Santos P, Dabelsteen T, Cardoso GC, Jennions MD. 2022

Ecological adaptation and birdsong: how body and bill sizes affect passerine sound

frequencies. *Behav Ecol* **33**, 798-806. (10.1093/beheco/arac042)

20.   Freckleton RP. 2010 Dealing with collinearity in behavioural and ecological data:

539   model averaging and the problems of measurement error. *Behav Ecol Sociobiol* **65**,

540   91-101. (10.1007/s00265-010-1045-6)

541   21.   Goodhue DL, Lewis W, Thompson R. 2017 A multicollinearity and measurement

542   error statistical blind spot: correcting for excessive false positives in regression and

543   PLS. *MIS Quarterly* **41**, 667-84. (https://www.jstor.org/stable/26635007)

544   22.   Tattersall GJ, Arnaout B, Symonds MRE. 2017 The evolution of the avian bill as a

545   thermoregulatory organ. *Biol Rev* **92**, 1630-56. (10.1111/brv.12299)

546   23.   Greenberg R, Cadena V, Danner RM, Tattersall GJ. 2012 Heat loss may explain bill

547   size differences between birds occupying different habitats. *PLoS One* **7**, e40933.

548   (10.1371/journal.pone.0040933)

549   24.   Dubiner S, Meiri S. 2022 Widespread recent changes in morphology of Old World

550   birds, global warming the immediate suspect. *Glob Ecol Biogeogr* **31**, 791-801.

551   (10.1111/geb.13474)

552   25.   McQueen A, Klaassen M, Tattersall GJ, Atkinson R, Jessop R, Hassell CJ, et al. 2022

553   Thermal adaptation best explains Bergmann's and Allen's Rules across ecologically

554   diverse shorebirds. *Nat Commun* **13**, 4727. (10.1038/s41467-022-32108-3)

555   26.   Grant PR, Grant BR. 2006 Evolution of character displacement in Darwin's finches.

556   *Science* **313**, 224-6. (10.1126/science.1128374)

557   27.   Podos J. 2001 Correlated evolution of morphology and vocal signal structure in

558   Darwin's fnches. *Nature* **409**, 185-8. (10.1038/35051570)

559   28.   Podos J, Nowicki S. 2004 Beaks, adaptation, and vocal evolution in Darwin's finches.

560   *Bioscience* **54**, 501-10. (10.1641/0006-3568(2004)054[0501:BAAVEI]2.0.CO;2)

29. Mason NA, Burns KJ. 2015 The effect of habitat and body size on the evolution of

vocal displays in Thraupidae (tanagers), the largest family of songbirds. *Biol J Linn*

Soc **114**, 538-51. (10.1111/bij.12455)

30. Mikula P, Valcu M, Brumm H, Bulla M, Forstmeier W, Petruskova T, et al. 2021 A

global analysis of song frequency in passerines provides no support for the acoustic

adaptation hypothesis but suggests a role for sexual selection. *Ecol Lett* **24**, 477-86.

(10.1111/ele.13662)

31. Derryberry EP, Seddon N, Derryberry GE, Claramunt S, Seeholzer GF, Brumfield

RT, et al. 2018 Ecological drivers of song evolution in birds: disentangling the effects

of habitat and morphology. *Ecol Evol* **8**, 1890-905. (10.1002/ece3.3760)

32. Platt JR. 1964 Strong inference: certain systematic methods of scientific thinking may

produce much more rapid progress than others. *Science* **146**, 347-53.

(10.1126/science.146.3642.347)

33. Revell LJ, Harmon LJ. 2022 *Phylogenetic comparative methods in R*. Princeton

University Press.

34. Marquet PA, Fernández M, Navarette AA, Valdovinos C. 2004 Diversity emerging:

towards a deconstruction of biodiversity patterns. In: *Frontiers of biogeography: New*

directions in the geography of nature. p. 191-209. Sunderland: Sinauer Associates.

35. Higgins PJ, Ford HA, Christidis L. 2008 Family Meliphagidae. In: *Handbook of the*

Birds of the World: Penduline-tits to shrikes. p. 498–691. Barcelona, Spain: Lynx

Edicions.

36. Hardt B, Benedict L. 2020 Can you hear me now? A review of signal transmission

and experimental evidence for the acoustic adaptation hypothesis. *Bioacoustics* **30**,

716-42. (10.1080/09524622.2020.1858448)

37.   Dunning JB. 2008 *CRC Handbook of Avian Body Masses*. Taylor & Francis Group,

LLC.

38.   Marki PZ, Kennedy JD, Cooney CR, Rahbek C, Fjeldsa J. 2019 Adaptive radiation

and the evolution of nectarivory in a large songbird clade. *Evolution* **73**, 1226-40.

(10.1111/evo.13734)

39.   Friedman NR, Miller ET, Ball JR, Kasuga H, Remes V, Economo EP. 2019 Evolution

of a multifunctional trait: shared effects of foraging ecology and thermoregulation on

beak morphology, with consequences for song evolution. *Proc Royal Soc B* **286**,

20192474. (10.1098/rspb.2019.2474)

40.   O'Meara BC, Ané C, Sanderson MJ, Wainwright PC. 2006 Testing for different rates

of continuous trait evolution using likelihood. *Evolution* **60**, 922-33. (10.1111/j.0014-

3820.2006.tb01171.x)

41.   R Core Team. 2013 R: a language and environment for statistical computing.

42.   Pearse WD, Morales-Castilla I, James LS, Farrell M, Boivin F, Davies TJ. 2018

Global macroevolution and macroecology of passerine song. *Evolution* **72**, 944-60.

(10.1111/evo.13450)

43.   Cardoso GC. 2013 Using frequency ratios to study vocal communication. *Anim Behav*

85, 1529-32. (10.1016/j.anbehav.2013.03.044)

44.   Boncoraglio G, Saino N. 2007 Habitat structure and the evolution of bird song: a

meta-analysis of the evidence for the acoustic adaptation hypothesis. *Funct Ecol* **21**,

134-42. (10.1111/j.1365-2435.2006.01207.x)

45.   Zollinger SA, Podos J, Nemeth E, Goller F, Brumm H. 2012 On the relationship

between, and measurement of, amplitude and frequency in birdsong. *Anim Behav* **84**,

e1-e9. (10.1016/j.anbehav.2012.04.026)

46. Tobias JA, Sheard C, Pigot AL, Devenish AJM, Yang J, Sayol F, et al. 2022

AVONET: morphological, ecological and geographical data for all birds. *Ecol Lett*

25, 581-97. (10.1111/ele.13898)

47. Bird species distribution maps of the world. Version 2018.1 [Internet]. 2018.

Available from: http://datazone.birdlife.org/species/requestdis.

48. Fick SE, Hijmans RJ. 2017 WorldClim 2: new 1-km spatial resolution climate

surfaces for global land areas. *Int J Climatol* **37**, 4302-15. (10.1002/joc.5086)

49. Hijmans RJ. 2015 Raster: geographic data analysis and modeling. *R package*.

(https://cran.r-project.org/package=raster)

50. Hay EM, McGee MD, Chown SL. 2022 Geographic range size and speciation in

honeyeaters. *BMC Ecol Evol* **22**, 1-14. (10.1186/s12862-022-02041-6)

51. Paradis E, Claude J, Strimmer K. 2004 APE: analyses of phylogenetics and evolution

in R language. *J Bioinform* **20**, 289-90. (10.1093/bioinformatics/btg412)

52. Revell LJ. 2012 phytools: an R package for phylogenetic comparative biology (and

other things). *Methods Ecol Evol* **3**, 217-23. (10.1111/j.2041-210X.2011.00169.x)

53. Textor J, van der Zander B, Gilthorpe MS, Liskiewicz M, Ellison GT. 2016 Robust

causal inference using directed acyclic graphs: the R package 'dagitty'. *Int J*

Epidemiol **45**, 1887-94. (10.1093/ije/dyw341)

54. Arif S, MacNeil MA. 2022 Predictive models aren't for causal inference. *Ecol Lett* **25**,

1741-5. (10.1111/ele.14033)

55. Grafen A. 1989 The phylogenetic regression. *Philos Trans R Soc Lond B Biol Sci* **326**,

119-57.

56. Martins EP, Hansen TF. 1997 Phylogenies and the comparative method: a general

approach to incorporating phylogenetic information into the analysis of interspecific

data. *Am Nat* **149**, 646-67.

57.   Pinheiro J, Bates D, DebRoy S, Sarkar D, Team RC. 2013 nlme: Linear and nonlinear
mixed effects models.

58.   Revell LJ. 2010 Phylogenetic signal and linear regression on species data. *Methods*
Ecol Evol **1**, 319-29.

59.   Dinnage R, Skeels A, Cardillo M. 2020 Spatiophylogenetic modelling of extinction
risk reveals evolutionary distinctiveness and brief flowering period as threats in a
hotspot plant genus. *Proc Royal Soc B* **287**, 20192817. (10.1098/rspb.2019.2817)

60.   Martins TG, Simpson D, Lindgren F, Rue H. 2013 Bayesian computing with INLA:
new features. *Comput Stat Data Anal* **67**, 68-83. (10.1016/j.csda.2013.04.014)

61.   Rue H, Martino S. 2007 Approximate Bayesian inference for hierarchical Gaussian
Markov random field models. *J Stat Plan Inference* **137**, 3177-92.
(10.1016/j.jspi.2006.07.016)

62.   Felsenstein J. 1985 Phylogenies and the comparative method. *Am Nat* **125**, 1-15.
(10.1086/284325)

63.   Hansen TF. 1997 Stabilising selection and the comparative analysis of adaptation.
Evolution **51**, 1341-51. (10.2307/2411186)

64.   Butler MA, King AA. 2004 Phylogenetic comparative analysis: a modeling approach
for adaptive evolution. *Am Nat* **164**, 683-95. (10.1086/426002)

65.   Clavel J, Escarguel G, Merceron G. 2015 mvMORPH: an R package for fitting
multivariate evolutionary models to morphometric data. *Methods Ecol Evol* **6**, 1311-9.
(10.1111/2041-210x.12420)

66.   Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. 2013 *Bayesian*
Data Analysis. CRC press.

67. Carpenter B, Gelman A, Hoffman MD, Lee D, Goodrich B, Betancourt M, et al. 2017

Stan: a probabilistic programming language. *J Stat Softw* **76**, 1-32.

(10.18637/jss.v076.i01)

68. Bürkner P-C. 2017 brms: an R package for Bayesian multilevel models using Stan. *J*

Stat Softw **80**, 1-28. (10.18637/jss.v080.i01)

69. Hoffman MD, Gelman A. 2014 The No-U-Turn Sampler: adaptively setting path

lengths in Hamiltonian Monte Carlo. *J Mach Learn Res* **15**, 1593-623.

(10.5555/2627435.2638586)

70. Derryberry EP. 2009 Ecology shapes birdsong evolution: variation in morphology and

habitat explains variation in white-crowned sparrow song. *Am Nat* **174**, 24-33.

(10.1086/599298)

71. Sebastianelli M, Lukhele SM, Nwankwo EC, Hadjioannou L, Kirschel ANG. 2022

Continent-wide patterns of song variation predicted by classical rules of

biogeography. *Ecol Lett* **25**, 2448–62. (10.1111/ele.14102)

72. James FC. 1970 Geographic size variation in birds and its relationship to climate.

Ecology **51**, 365-90. (10.2307/1935374)

73. Albright TP, Mutiibwa D, Gerson AR, Smith EK, Talbot WA, O'Neill JJ, et al. 2017

Mapping evaporative water loss in desert passerines reveals an expanding threat of

lethal dehydration. *Proc Natl Acad Sci USA* **114**, 2283-8. (10.1073/pnas.1613625114)

74. Keith DA. 2017 *Australian vegetation*. Cambridge University Press.

75. Frappell PB, Hinds DS, Boggs DF. 2001 Scaling of respiratory variables and the

breathing pattern in birds: an allometric and phylogenetic approach. *Physiol Biochem*

Zool **74**, 75-89. (10.1086/319300)

76. Raftery A. 2013 Avian anaesthesia. *In Practice* **35**, 272-8. (10.1136/inp.f2861)

77.   Dial KP, Greene E, Irschick DJ. 2008 Allometry of behavior. *Trends Ecol Evol* **23**,

682   394-401. (10.1016/j.tree.2008.03.005)

683   78.   Gardner JL, Peters A, Kearney MR, Joseph L, Heinsohn R. 2011 Declining body size:

684   a third universal response to warming? *Trends Ecol Evol* **26**, 285-91.

685   (10.1016/j.tree.2011.03.005)

686   79.   Lindenmayer DB, Blanchard W, Westgate MJ, Foster C, Banks SC, Barton P, et al.

687   2019 Novel bird responses to successive, large-scale, landscape transformations. *Ecol*

688   *Monogr* **89**, e01362. (10.1002/ecm.1362)

689   80.   Bovo AAA, Ferraz KMPMB, Magioli M, Alexandrino ER, Hasui É, Ribeiro MC, et

690   al. 2018 Habitat fragmentation narrows the distribution of avian functional traits

691   associated with seed dispersal in tropical forest. *Perspect Ecol Conserv* **16**, 90-6.

692   (10.1016/j.pecon.2018.03.004)

693   81.   Ryding S, Klaassen M, Tattersall GJ, Gardner JL, Symonds MRE. 2021 Shape-

694   shifting: changing animal morphologies as a response to climatic warming. *Trends*

695   *Ecol Evol* **36**, 1036-48. (10.1016/j.tree.2021.07.006)

FIGURE CAPTIONS

Figure 1. Morphological diversity of honeyeater body sizes and beak shapes. Images are of

the: crow honeyeater[1] (a; *Gymnomyza aubryana*), barred honeyeater[1] (b; *Gliciphila*

undulata), New Holland honeyeater[2] (c; *Phylidonyris novaehollandiae*), tūī[1] (d;

Prosthemadera novaeseelandiae), white-eared honeyeater[3] (e; *Nesoptilotis leucotis*), yellow-

faced honeyeater[1] (f; *Caligavis chrysops*), crescent honeyeater[1] (g; *Phylidonyris*

pyrrhopterus), noisy miner[2] (h; *Manorina melanocephala*), eastern spinebill[1] (i;

Acanthorhynchus tenuirostris), New Caledonian myzomela[1] (j; *Myzomela caledonica*),

white-plumed honeyeater[1] (k; *Ptilotula penicillata*), New Caledonian friarbird[1] (l; *Philemon*

diemenensis), dusky myzomela[2] (m; *Myzomela obscura*), dark brown honeyeater[1] (n;

Lichmera incana), yellow wattlebird[1] (o; *Anthochaera paradoxa*), yellow honeyeater[2] (p;

Stomiopera flava), bell miner[3] (q; *Manorina melanophrys*), white-naped honeyeater[3] (r;

Melithreptus lunatus), white-fronted chat[2] (s; *Epthianura albifrons),* spiny-cheeked

honeyeater[3] (t; *Acanthagenys rufogularis*). Image size for each species reflects relative size of

the species. Image credits: [1]Matthias Dehling, [2]Eleanor Hay, [3]Steven Chown.

Figure 2. Conceptual diagram explaining the underlying interactions between environmental

and phenotypic variables and their influence on song. Song spectrogram displays frequency

(kHz) over time (s) and is an example from the New Zealand bellbird (*Anthornis melanura*;

XC153005). The three song variables considered in this study are indicated with annotations;

peak frequency represents the frequency of maximum amplitude across the entire song,

bandwidth is the difference between maximum and minimum frequency across the song, and

pace is the number of notes over song duration. Beak shape shows the beak variation in

honeyeaters, the top is the species with the lowest value for beak PC2 (Arfak honeyeater;

Melipotes gymnops) and the bottom is the species with the highest value for PC2 (Eastern

spinebill; *Acanthorhynchus tenuirostris*). Illustrations by Eleanor Hay.

Figure 3. Phylogeny containing the 163 honeyeater species for which we obtained song

variables for and used in analysis. Branches are coloured according to peak frequency. Bars

at the tips of the phylogeny represent body size and are coloured according to open (white)

and closed (black) habitat types. Birds are illustrations by Eleanor Hay.

Figure 4. Results from phylogenetic generalised least squares regression models (a) and the

spatiophylogenetic models (c-d), testing the influence of environmental and morphological

traits on song variables. Separate models were run to test the influence on peak frequency,

bandwidth, and pace, and are indicated in the legends. For the PGLS results (a), the predictor

variables have been scaled so the effect sizes are comparable, full model results are available

in Table S4. Maps in panels b-c display the mean spatial random effect for each song variable

which was extracted from the final spatiophylogenetic model. Density plots show the

marginals of the fixed effects in the model, with the mean and 95% credible intervals.

Predictor variables which were identified to have significant effects are highlighted in bold,

and full model results are available in Table S5.

Figure 5. Distribution of AIC values from mvMORPH models (a) when testing the effect of

open and closed habitat types on peak frequency, bandwidth, and pace. Note that models

were conducted over 100 stochastic character mapping of open and closed habitats on the

phylogeny. Models compared were single rate Brownian motion (BM1, n = 100), two rate

Brownian motion (BMM, n = 100), single rate Ornstein-Uhlenbeck (OU1, n = 100) and two

rate Ornstein Uhlenbeck (OUM, n = 100). Full model results are available in Table S6. Panel

b displays outcomes of Bayesian phylogenetic regression models (a), testing the influence of

environmental and morphological traits on the variance of peak frequency, bandwidth, and

pace. Models are indicated with the legend and full model results are available in Table S7.

Figure 1.

791 **Figure 2.**

Figure 3.

816 **Figure 4.**

826 **Figure 5.**

843