Ryanodine Receptor Stabilization Therapy Suppresses Ca^{2+}-Based Arrhythmias in a Novel Model of Metabolic HFpEF

Aaron D. Kaplan[1,2], Liron Boyman[1,3,4], Christopher W. Ward[1,5,6], W. Jonathan Lederer[1,3,4], Maura Greiser[1,3,6]*

[1] Center for Biomedical Engineering and Technology, University of Maryland School of Medicine, Baltimore, Maryland, USA.

[2] Division of Cardiovascular Medicine, Department of Medicine, University of Maryland School of Medicine, Baltimore, Maryland, USA

[3] Department of Physiology, University of Maryland School of Medicine, Baltimore, Maryland, USA.

[4] Greenebaum Comprehensive Cancer Center, University of Maryland School of Medicine, Baltimore, Maryland, USA

[5] Department of Orthopedics, University of Maryland School of Medicine, Baltimore, Maryland, USA.

[6] Claude D. Pepper Older Americans Independence Center, University of Maryland School of Medicine, Baltimore, Maryland, USA.

* Corresponding author: Maura Greiser.

email: mgreiser@som.umaryland.edu

Keywords: Heart Failure, Arrhythmia, Calcium Handling, Metabolic Stress,

Abstract

Heart Failure with preserved ejection fraction (HFpEF) is the most prevalent form of heart failure worldwide and its significant mortality is associated with a high rate of sudden cardiac death (SCD; 30% - 40%). Chronic metabolic stress is an important driver of HFpEF, and clinical data show metabolic stress as a significant risk factor for ventricular arrhythmias in HFpEF patients. The mechanisms of SCD and ventricular arrhythmia in HFpEF remain critically understudied and empirical treatment is ineffective. To address this important knowledge gap, we developed a novel preclinical model of metabolic-stress induced HFpEF using Western diet (High fructose and fat) and hypertension induced by nitric oxide synthase inhibition (with L-NAME) in wildtype C57BL6/J mice. After 5 months, mice display all clinical characteristics of HFpEF and present with stress-induced sustained ventricular tachycardia (VT). Mechanistically, we found a novel pattern of arrhythmogenic intracellular Ca^{2+} handling that is distinct from the well-characterized changes pathognomonic for heart failure with reduced ejection fraction. In addition, we show that the transverse tubular system remains intact in HFpEF and that arrhythmogenic, intracellular Ca^{2+} mobilization becomes hyper-sensitive to ß-adrenergic activation. Finally, in proof-of-concept experiments we show *in vivo* that the clinically used intracellular calcium stabilizer dantrolene, which acts on the Ca^{2+} release channels of the sarcoplasmic reticulum (SR), the ryanodine receptors, acutely prevents stress-induced VT in HFpEF mice. Therapeutic control of SR Ca^{2+} leak may present a novel mechanistic treatment approach in metabolic HFpEF.

Main Text

Heart failure (HF) with preserved ejection fraction (HFpEF), the most prevalent form of HF worldwide, is associated with significant morbidity and high mortality[1]. Initially considered a disorder of diastolic dysfunction related to hypertensive left-ventricular remodeling, HFpEF is now understood as a heterogeneous disease widely seen in patients with significant metabolic stress driven by underlying comorbidities of obesity, dyslipidemia, hypertension and decreased insulin sensitivity [1, 2]. A high rate of sudden cardiac death (SCD; ~30-40 %) and ventricular arrhythmias contribute to HFpEF morbidity and mortality, and evidence that empiric anti-arrhythmic treatment is ineffective in HFpEF [3, 4] suggests that arrhythmias in metabolic stress-related HFpEF may arise via a distinct and as yet unidentified mechanism[3].

To investigate arrhythmia mechanisms pertaining to the large clinical cohort of metabolic stress related HFpEF we developed a novel preclinical model in wildtype mice (C57BL6/J). The model is based solely on mimicking the stressors (hypercaloric Western diet, hypertension, obesity) found in patients with metabolic stress related HFpEF, without relying on genetically altered animals. To achieve this, we added high dietary fructose (20% of dietary calories), an important driver of metabolic stress, to the well-established murine HFpEF model produced by high fat diet and hypertension-inducing nitric oxide synthase inhibition (with L-NAME)[5]. After 5 months, this **f**ructose-**m**etabolic HFpEF model (FM-HFpEF) demonstrated, *in vivo,* key aspects of the targeted clinical human HFpEF phenotype, including obesity, sarcopenia, decreased insulin sensitivity, high serum levels of brain natriuretic peptide (BNP), decreased global longitudinal strain (GLS) with normal ejection fraction (EF) and, importantly, stress-induced ventricular tachycardia (VT; Fig.1A-E).

Ca^{2+} "leaking" from its stores in the sarcoplasmic reticulum (SR) is the key mechanism underlying Ca^{2+} based arrhythmogenesis in heart failure with reduced

ejection fraction (HFrEF)(6, 7). Here we show a massive arrhythmogenic intracellular
Ca^{2+} leak in FM-HFpEF ventricular myocytes, much larger than in HFrEF, evidenced by
a ~ 10-fold increase in spontaneous Ca^{2+} spark frequency and ~ 9-fold increase in spark
mediated SR Ca^{2+} leak (Fig.1F). Consistent with this massive SR Ca^{2+} leak was a
dramatic increase in spontaneous Ca^{2+} wave activity (~7-fold higher in FM-HFpEF
myocytes compared to control myocytes, Fig.1G), the cellular trigger for Ca^{2+} based
arrhythmias(6, 7).

We show that ß-adrenergic challenge *in vivo* elicits VT in FM-HFpEF mice
(Fig.1E). We also find that *in vitro* ß-adrenergic challenge (isoproterenol 50 nmol/L) in
single FM-HFpEF cardiomyocytes elicited a much higher frequency of arrhythmogenic
Ca^{2+} waves than in control cells, thus further supporting the Ca^{2+} basis of arrhythmias in
metabolic HFpEF (Fig.1G). Importantly, this high degree of intracellular Ca^{2+} mobilization
in vitro, together with the inducibility of sustained VTs *in vivo*, demonstrate an intact and
highly sensitive ß-adrenergic response in FM-HFpEF.

A crucial underlying mechanism of Ca^{2+} based arrhythmogenesis in
HFrEF is a higher diastolic Ca^{2+} concentration in the cytosol ($[Ca^{2+}]_i$), which sensitizes
the Ca^{2+} release channels of the SR, the ryanodine receptors type 2 (RyR2) to release
Ca^{2+} (8). Surprisingly, calibrated measurements of $[Ca^{2+}]_i$ in single cardiomyocytes
electrically paced at physiological frequencies showed that diastolic $[Ca^{2+}]_i$ was not
increased in FM-HFpEF myocytes (2 Hz, Fig.1Hi,ii) despite an increase in the magnitude
of their systolic $[Ca^{2+}]_i$ (Fig.1Hi-iii). In addition, Ca^{2+} transients from FM-HFpEF mice
were shorter or unchanged from controls at physiological stimulation rates of 1-4 Hz,
another striking difference from HFrEF where Ca^{2+} transients are prolonged (Fig.1I). To
define the underlying mechanisms of this FM-HFpEF Ca^{2+} phenotype, we first examined
the ultrastructure of the transverse and axial tubular system (TATS) of living, single
cardiomyocytes. Sub-diffraction Airy scan imaging revealed no changes in TATS
density, organization or orientation in FM-HFpEF (Fig.2A). These results are contrary to
the severe reduction and disorganization of TATS pathognomonic for HFrEF, where the
structural disruption of the cardiac dyad, the close apposition (~20 nm) of L-type Ca^{2+}
channels ($Ca_v1.2$) with RyR2, causes arrhythmogenic SR Ca^{2+} release(9).

We next quantified the expression and post-translational modification (PTM) of
dyadic and SR Ca^{2+} handling proteins. In ventricles from FM-HFpEF mice $Ca_v1.2$
expression was unchanged. Surprisingly, RyR2 expression and its phosphorylation
(Serine2808) were significantly reduced (Fig.2B), another important distinction from
changes seen in HFrEF. In addition, calsequestrin 2 (CSQ2), the major intra-SR Ca^{2+}
buffering protein, was significantly reduced, as was the expression and phosphorylation
of phospholamban (PLB), the regulatory protein of the SR Ca^{2+} ATPase (SERCA)
(Fig.2B). Taken together, these results suggest a significant remodeling of the cardiac
dyad due to an altered stoichiometry of the $Ca_v1.2$-RyR2 nanodomain. In addition, the
expression and PTM of key SR Ca^{2+} handling proteins are reduced (schematic shown in
Fig. 2D, E). While future work is necessary to decode the ensemble contribution of these
changes to the distinct HFpEF Ca^{2+} signaling, reduced CSQ2 is known to cause
ventricular arrhythmias via SR Ca^{2+} leak in cardiomyocytes due to reduced intra SR Ca^{2+}
buffering (10).

To test a novel mechanistic approach for anti-arrhythmic therapy in HFpEF we
performed proof-of-concept experiments with dantrolene, a ryanodine receptor
stabilizing agent. Dantrolene is used clinically for the treatment of malignant
hyperthermia(11) and has recently been shown to suppress Ca^{2+} based arrhythmias in

human hearts *ex vivo*(12). Here we show that acute, mechanistic Ca^{2+} stabilizing
treatment with dantrolene of FM-HFpEF mice *in vivo* (40 mg/kg i.p) significantly reduced
arrhythmia burden and completely abolished stress-induced VTs (Fig.2C).

In summary, we report a novel preclinical model of metabolic HFpEF that
phenocopies the complex cardiometabolic features and stress-induced ventricular
tachycardias seen in clinical HFpEF related to metabolic stress (Fig.1A-E)(1, 3). We
show here for the first time that metabolic stress induced HFpEF produces Ca^{2+} based
arrhythmogenesis distinct from the "Ca^{2+} phenotype" pathognomonic for HFrEF (Fig.2D,
E). Specifically, a massive intracellular Ca^{2+} leak, unchanged diastolic $[Ca^{2+}]_i$, intact
TATS and reduced expression of key SR Ca^{2+} handling proteins underlie
arrhythmogenic Ca^{2+} signaling in HFpEF. Importantly, these changes are also distinct
from a form of HFpEF secondary to genetic hypertension (Dahl salt-sensitive rats) that
progresses to HFrEF, where the ß-adrenergic responsiveness of intracellular Ca^{2+}
mobilization was blunted(13). Another novel and distinct (from HFrEF) finding in FM-
HFpEF was a decrease in CSQ2 expression. Previous work established that even small
reductions of CSQ2 expression produce significant Ca^{2+} dependent arrhythmias, due to
both the reduced Ca^{2+} binding capacity of the SR and increased RyR2 activity(10).

Finally, we demonstrate in proof-of-concept experiments that dantrolene, a
clinically relevant ryanodine receptor stabilizer, acutely abolishes stress-induced VTs in
FM-HFpEF mice. These results represent the first mechanistic, antiarrhythmic treatment
concept (RyR2 stabilization) for VTs in metabolic HFpEF. Importantly, dantrolene, which
is contraindicated in HFrEF due to its negative inotropy, could be used clinically in
HFpEF patients. Future work will further evaluate the long-term effects and clinical
treatment potential of ryanodine receptor stabilization therapy in metabolic stress related
HFpEF.

121
122   **Materials and Methods**
123   Male C57BL6/J mice were used in this study. Cardiac myocyte isolation, Ca^{2+} imaging
124   and sub-diffraction microscopy were performed as previously described(14, 15). Detailed
125   descriptions are provided in the Supporting Information.
126

Acknowledgments
Sources of Funding: University of Maryland Claude D. Pepper Center Grant P30
AG028747 (MG).7U19 AI090959 (WJL &LB); Frontiers in Anesthesia Research Award
from International Anesthesia Research Society (WJL & LB); R01 HL142290 (WJL
&CWW); 5R35GM140822 (WJL); U01 HL116321 (WJL); Special BioMET/UMB Funds
(WJL). R01 AR071618 (CWW).

References

1. B. A. Borlaug, K. Sharma, S. J. Shah, J. E. Ho, Heart Failure With Preserved Ejection Fraction. *Journal of the American College of Cardiology* **81**, 1810–1834 (2023).

2. B. A. Borlaug, *et al.*, Obesity and heart failure with preserved ejection fraction: new insights and pathophysiological targets. *Cardiovascular research* **118**, 3434–3450 (2023).

3. M. F. Yuyun, S. Kinlay, J. P. Singh, J. Joseph, Are arrhythmias the drivers of sudden cardiac death in heart failure with preserved ejection fraction? A review. *ESC Heart Fail* **10**, 1555–1569 (2022).

4. A. Gutierrez, *et al.*, Nonsustained ventricular tachycardia in heart failure with preserved ejection fraction. *Pacing Clin Electrophysiol* **43**, 1126–1131 (2020).

5. G. G. Schiattarella, *et al.*, Nitrosative stress drives heart failure with preserved ejection fraction. *Nature* **568**, 351–356 (2019).

6. S. M. Pogwizd, D. M. Bers, Cellular basis of triggered arrhythmias in heart failure. *Trends Cardiovasc Med* **14**, 61–66 (2004).

7. A. M. Gomez, *et al.*, Defective excitation-contraction coupling in experimental cardiac hypertrophy and heart failure. *Science* **276**, 800–6 (1997).

8. D. A. Eisner, J. L. Caldwell, A. W. Trafford, D. C. Hutchings, The Control of Diastolic Calcium in the Heart. *Circulation Research* **126**, 395–412 (2020).

9. L. S. Song, *et al.*, Orphaned ryanodine receptors in the failing heart. *Proc Natl Acad Sci U S A* **103**, 4305–10 (2006).

10. N. Chopra, *et al.*, Modest reductions of cardiac calsequestrin increase sarcoplasmic reticulum Ca2+ leak independent of luminal Ca2+ and trigger ventricular arrhythmias in mice. *Circ Res* **101**, 617–26 (2007).

11. G. G. Harrison, Malignant hyperthermia. Dantrolene--dynamics and kinetics. *Br J Anaesth* **60**, 279–286 (1988).

12. S. A. George, *et al.*, Ryanodine receptor inhibition with acute dantrolene treatment reduces arrhythmia susceptibility in human hearts. *Am J Physiol Heart Circ Physiol* **325**, H720–H728 (2023).

13. P. J. Kilfoil, *et al.*, Distinct features of calcium handling and β-adrenergic sensitivity in heart failure with preserved *versus* reduced ejection fraction. *J Physiol* **598**, 5091–5108 (2020).

14. M. Greiser, *et al.*, Calcium and bicarbonate signaling pathways have pivotal, resonating roles in matching ATP production to demand. *Elife* **12** (2023).

15. M. Greiser, *et al.*, Tachycardia-induced silencing of subcellular Ca2+ signaling in atrial myocytes. *The Journal of clinical investigation* **124**, 4759–72 (2014).

173 **Figures**

Figure 1

Figure 1. Arrhythmogenic Ca²⁺ signaling in metabolic HFpEF. Metabolic HFpEF model in
mice. **A**. Body composition: **i**. Fat, **ii**. Bodyweight and **iii**. Lean Mass. **B**. Insulin tolerance test
(ITT) **C**. Transthoracic echocardiography **i**. Ejection fraction (EF), **ii.** Left ventricular mass (LVM)
iii. Global longitudinal strain (GLS), ******: $p<0.01$. **D**. Serum levels of B-type natriuretic peptide
(BNP). **E.** EKGs and arrhythmia score. ******: $p<0.01$. Calcium imaging. **F.** Ca²⁺ sparks in HFpEF. **i.**
Exemplars of Ca²⁺ sparks. **ii.** Ca²⁺ spark frequency, **iii.** Ca²⁺ spark-mediated SR Ca²⁺ leak, **iv.**
Ca²⁺ spark amplitude, **v.** Ca²⁺ spark duration as full duration half maximum (FDHM). ********:
$p<0.0001$; *******: $p<0.001$; ******: $p<0.01$; *****: $p<0.05$. **G.** Ca²⁺ waves. **i.** Exemplars of spontaneous Ca²⁺
waves. **ii.** Exemplars of spontaneous Ca²⁺ waves after isoproterenol treatment. *denotes
stimulated beat. **iii.** Ca²⁺ wave frequency **iv.** Ca²⁺ wave free survival **v.** Ca²⁺ wave frequency (+
isoproterenol). *******: $p<0.001$; ******: $p<0.01$. **H.** [Ca²⁺]ᵢ in HFpEF myocytes. **i.** Stimulated Ca²⁺
transients (0.5-2 Hz). Summary data of **ii.** diastolic [Ca²⁺]ᵢ, **iii.** Systolic [Ca²⁺]ᵢ. **I. i, ii.** Stimulated
Ca²⁺ transients (1-4 Hz). **iii.** Ca²⁺ transient amplitude **iv.** Ca²⁺ transient decay.

Figure 2

Figure 2. A. Transverse-Axial Tubule system (TATS). AiryScan images of living ventricular myocytes stained with the membrane dye di-8-ANEPPS (5 µmol/L) to label the TATS. **ii.** Total length of the TATS. **iii.** Number of junctions that connect intersecting perpendicular tubules. **iv.** The length of the TATS at each direction within the cells. **B.** Protein expression levels of key Ca^{2+} handling proteins. $Ca_V1.2$: α1c subunit of the L-type Ca^{2+} channel, SERCA: SR Ca^{2+} ATPase, PLB: phospholamban, PLB-P: phosphorylated-PLM (Serine16), CSQ2: calsequestrin 2, RyR2: Ryanodine receptor type 2, RyR2-P: phosphorylated RyR2 (Serine2808). **C.** *in vivo* dantrolene treatment of FM-HFpEF mice. **i.** Summary data showing arrhythmia index in FM-HFpEF mice (n=6) pretreated with i.p. dantrolene (40 mg/kg i.p; 30 min) compared to untreated FM-HFpEF mice (n=6). **L.** Table showing changes in intracellular Ca^{2+} handling in HFrEF compared to the FM-HFpEF model. **M.** Schematic depicting organization of the dyadic Ca^{2+} release-apparatus in normal, HFrEF and FM-HFpEF.

7