A case study of *Whirly1* (*WHY1*) evolution in the angiosperms: altered sequence, splicing, and expression in a clade of early-transitional mycoheterotrophic orchids

Rachel M. Muti[1], Craig F. Barrett[2], Brandon T. Sinn[1,3]*

[1] Otterbein University, Department of Biology and Earth Science, 1 South Grove Street, Westerville,
Ohio, USA

[2] West Virginia University, Department of Biology, Morgantown, West Virginia, USA

[3] University of Latvia, Faculty of Biology, 1 Jelgavas iela, Riga, Latvia

* Correspondence:
Brandon T. Sinn
sinn1@otterbein.edu

Keywords: Mycoheterotrophy; *Corallorhiza*; Orchid; Genomic Stability; Intron Retention;
Transcription Factor; Plastome Evolution; *Whirly1*

Abstract

The plastid-targeted transcription factor *Whirly1* (*WHY1*) has been implicated in chloroplast
biogenesis, plastid genome stability, and fungal defense response, which together represent
characteristics of interest for the study of autotrophic losses across the angiosperms. While gene loss
in the plastid and nuclear genomes has been well studied in mycoheterotrophic plants, the evolution
of the molecular mechanisms impacting genome stability are completely unknown. Here we
characterize the evolution of *WHY1* in four early-transitional mycoheterotrophic orchid species in the
genus *Corallorhiza* by synthesizing the results of phylogenetic, transcriptomic, and comparative
genomic analyses with *WHY1* genomic sequences sampled from 21 orders of angiosperms. We found
an increased number of non-canonical *WHY1* isoforms assembled from all but the greenest
Corallorhiza species, including intron retention in some isoforms. Within *Corallorhiza*,
phylotranscriptomic analyses revealed the presence of tissue-specific differential expression of
WHY1 in only the most photosynthetically capable species and a coincident increase in the number of
non-canonical *WHY1* isoforms assembled from fully mycoheterotrophic species. Gene- and codon-
level tests of *WHY1* selective regimes did not infer significant signal of either relaxed selection or
episodic diversifying selection in *Corallorhiza*, but did so for relaxed selection in the late-stage full
mycoheterotrophic orchids *Epipogium aphyllum* and *Gastrodia elata*. Taken together, this study
provides the most comprehensive view of *WHY1* evolution in angiosperms to date. Our analyses
reveal that splicing alteration and decreased expression of *WHY1* are coincident with deceased
plastome stability in a group of early-transitional mycoheterotrophic orchids and that these changes
may precede the selective shifts observed in late-stage mycoheterotrophic species.

1   Introduction

The ability to photosynthesize has been lost dozens of times across the angiosperm Tree of Life, and at least 30 independent losses have occurred in the Orchidaceae (Merckx and Freudenstein, 2010; Barrett et al., 2014; Barrett et al., 2019). Mycoheterotrophy, the derivation of carbon nutrition from fungi (Leake, 1994), is common to all orchids during early development and is a nutritional requirement due to the lack of endosperm in their seeds (Rasmussen, 1995; 2002). In lieu of stored nutrition, orchid seeds have evolved a complex symbiotic relationship with fungi, where orchid seeds germinate only in the presence of an appropriate fungal partner and the nutrition required for embryo development is derived exclusively via the degradation of fungal hyphae which penetrate the orchid cells (Merckx, 2013; Zeng et al., 2016; Yuan et al., 2018). The duration of reliance upon their fungal partner for nutrition has been extended in some orchid species, which have evolved to parasitize fungi for the entirety of their lives. Independent shifts to a mycoheterotrophic condition throughout the angiosperms have independently led to plastid genome (plastome) degradation (Barrett et al., 2014; Wickett et al., 2016; Graham et al., 2017; Timilsena, et al., 2023), elevated rates of nucleotide substitution (Lemaire et al., 2010; Wicke et al., 2016), and oftentimes the loss of morphological structures such as leaves and roots (Leake, 1994).

Corallorhiza is a North American, temperate genus comprising 12 species of morphologically-reduced, mycoheterotrophic orchids for which varying states of plastid genome (plastome) degradation and inferred photosynthetic ability have been characterized (Zimmer et al., 2008; Cameron et al., 2009; Barrett and Freudenstein, 2008; Barrett and Davis, 2012; Barrett et al., 2014; Barrett et al., 2018). The presence of relatively intact plastomes containing the expected repertoire of housekeeping genes in *Corallorhiza* species evidences the clade as a group of early-transitional mycoheterotrophs, in contrast with late-stage mycoheterotrophic species which have highly degraded plastomes and lack many or all plastid housekeeping genes, such as *Epipogium* and *Gastrodia* species (*sensu* Barrett et al., 2012; see also Barrett et al., 2014). *Corallorhiza* species parasitize Basidiomycete fungi that are engaged in mycorrhizal relationships with nearby autotrophic plants, predominantly in the families Russulaceae and Thelephoraceae (Taylor and Bruns, 1997; Taylor et al., 2004; Barrett et al., 2010; Freudenstein and Barrett, 2014). Besides their relationship with fungi, a conspicuous characteristic of *Corallorhiza* species is the complete loss of both leaf laminae and roots. A recurrent theme of morphological reduction has been documented across parasitic and mycoheterotrophic plant lineages (see Leake, 1994), and work has recently focused on genomic content and gene expression in mycoheterotrophic species in order to improve our understanding of the genomic precursors and consequences of this trophic transition (Barrett et al., 2014; Wickett et al., 2016; Graham et al., 2017; Zhang et al., 2017; Yuan et al., 2018; Cai, 2023; Timilsena et al., 2023).

The integrity of the plastome of parasitic and mycoheterotrophic plants is of particular interest as reduction in gene content, increased number of pseudogenes, structural variation, and a reduction in overall genome length have been found to correlate with the degree of external carbon reliance among parasitic angiosperms. Generally, disruptions to the genome such as double-strand DNA breaks are harmful to the organism, and many mechanisms that help to protect against such occurrences have evolved throughout the Tree of Life (Waterworth et al., 2011). One such genome stabilizing mechanism that is increasingly recognized for its involvement in processes such as plastome double-strand break repair is the activity of the Whirly family of transcription factors (*WHY*; Desveaux et al., 2005).

Transcription factors are regulatory gene products that function by binding to DNA
(Latchman, 1993). The Whirly family comprises *WHY1*, *WHY2*, and *WHY3*, which are three plant-
specific, nuclear-encoded genes with DNA binding domains that are named for their whirligig-like
structural conformation (Desveaux et al., 2002; Cappadocia et al., 2010). Crystal structures of the
Whirly transcription factors have been determined as tetramers that have a single-stranded DNA
binding domain that spans two subunits (Cappadocia et al., 2013). Of particular interest is *WHY1*,
the product of which has been implicated to play roles in myriad processes including mediation of
abiotic stressors (Zhuang et al., 2019, 2020; Ruan et al., 2022), induction of double strand DNA
break repair (Cappadocia et al., 2010), and plastid biogenesis (Prikryl et al., 2008). Transcription
factors are crucial to myriad regulatory and developmental processes, which is reflected in the
massive expansions of many TF gene families in plants (Lehti-Shiu et al., 2017). Our present
understanding of TF evolution has largely been informed through the investigations focused on
understanding their roles in morphological or ecological diversification (see de Mendoza et al.,
2013; Lai et al., 2020), rather than how they change in systems which have undergone coincident
extreme loss of morphological and genomic features.

WHY1 has been shown to dually localize to both plastids and the nucleus (Krause et al.,
2005; Grabowski et al., 2008; Prikryl et al., 2008; Isemer et al., 2012; Ren et al., 2017). In
chloroplasts, *WHY1* localizes to the boundary between the thylakoid and nucleoid membrane in
chloroplasts and has been implicated in retrograde signaling regulating H_2O_2 homeostasis and as a
coordinator of photosynthetic gene expression (Lepage et al., 2013; Foyer, et al., 2014; Lin et al.,
2019). Species that have undergone the transition to heterotrophy experience elevated levels of
oxidative stress compared to autotrophic relatives (Suetsugu et al., 2017; Lallemand et al., 2019).
Additionally, *WHY1* proteins stabilize plastid genomes by non-specific binding to the genome which
protects against microhomology-mediated DNA rearrangements, including deletions and
duplications of sequences (Maréchal et al., 2009; Lepage et al., 2013; Zampini et al., 2015).

WHY1 is involved in complex roles in both plant defense responses and genomic
stabilization. It has long been known that mutations which reduce the binding affinity of *WHY1*
correlate with increased infection by some pathogens (Desveaux et al., 2004). *WHY1* has been
shown to bind to specific DNA promoter regions that can induce transcription (Zhuang et al., 2020),
aid in defense response signaling and accumulation of disease resistance (Desveaux et al., 2004;
Desveaux et al., 2005), and also maintain telomere length in the nuclear genome (Desveaux et al.,
2005). Recently, *WHY1* has even been shown capable of negatively regulating the RNA interference
response to two geminiviruses (Sun et al., 2023). Taken together, the literature suggests that
modulation of *WHY1* expression can result in tradeoffs between plant defense and genomic stability.

The roles *WHY1* plays in stabilizing both the nuclear and plastid genomes (Yoo et al., 2007;
Zampini et al., 2015), defense response (Desveaux et al., 2005), and chloroplast development (Qiu
et al., 2022) are central to our choice to study the evolution of this transcription factor. In particular,
experimental work demonstrating plastome destabilization (Lepage et al., 2013; Zampini et al.,
2015), albinism, and variegation in *WHY1* mutants (Prikryl et al., 2008; Ren et al., 2017; Qiu et al.,
2022) makes the gene a compelling target given the reduction in plastome content and length
observed across parasitic angiosperms (Barrett et al., 2014; Wicke et al., 2016). To date, work
characterizing the sequence and expression diversity of *WHY1* has been restricted to model or
agricultural systems (Cappadocia et al., 2013; Ruan et al., 2022; Taylor et al., 2022) and no phylo-
comparative investigations of sequence evolution and selective regime have been conducted. The
implication of *WHY1* in processes associated with the mycoheterotrophic condition frames a

phylogenetically-informed investigation of *WHY1* along a trophic gradient as an important step in
improving our understanding of the evolution of plastid-targeted TFs in heterotrophic plant lineages.

We focus on four species, *C. trifida*, *C. striata*, *C. wisteriana*, and *C. maculata*, which
together comprise an early-transitional trophic gradient to full mycoheterotrophy with sister
relationships among partial and full mycoheterotrophs (Figure 1). The well-characterized
phylogenetic relationships between these four *Corallorhiza* species (Barrett et al., 2018) provide a
powerful framework upon which to investigate *WHY1* evolution during the early stages of transition
to full mycoheterotrophy while accounting for phylogenetic non-independence (*sensu* Felsenstein,
1987). We consider *C. trifida* and *C. wisteriana* to be partial mycoheterotrophs, as their tissues
contain measurable chlorophyll content and their plastomes are the most genetically-intact plastid
genomes in the genus, although photosynthesis has only been directly observed in *C. trifida* (Zimmer
et al., 2008; Cameron et al., 2009; Barrett et al., 2014; Barrett et al., 2018). Conversely, we consider
C. maculata and *C. striata* to be fully mycoheterotrophic, evidenced by their highly reduced
chlorophyll content and degradation of many photosynthesis-related genes (Barrett et al., 2014;
Barrett et al., 2018).

Here we characterize the evolution of *WHY1* across a mycoheterotrophic gradient, framed by
phylogenetic context provided by the broadest taxonomic sampling of the gene to date, including
late-stage mycoheterotrophic orchid species from the genera *Epipogium* and *Gastrodia*. Our
integrative work leverages a combination of novel and publicly available data generated from DNA
sequencing, RNAseq, and Oxford Nanopore sequencing from 110 angiosperm species representing
21 orders. Taken together, the results of our analyses of *WHY1* sequence, expression, splicing, and
selective regime across both a trophic gradient and the angiosperms more broadly suggest that the
gene may play a critical role in maintaining plastome stability after the transition to
mycoheterotrophy.

2   Materials and Methods

Publicly-available Sequences

Annotated *WHY1* sequences were obtained from the Nucleotide and RefSeq NCBI databases
and Orchidstra, an orchid-specific database (Chao et al., 2017). Only *WHY1* sequences containing
canonical *WHY1* ORFs (open reading frames) were retained. Additionally, sequences were excluded
if they did not contain the ssDNA binding region (KGKAAL; *A. thaliana* Q9M9S3) as reported by
Cappadocia et al. (2013; PDB 4KOO). Identical sequences were excluded for taxa with multiple
database accessions. Stop codons were trimmed from sequences, with the exception of premature
stop codons in sequences from known mycoheterotrophs, which were changed to gap characters (-)
for compatibility with downstream methods (e.g. HyPhy, see below). In total, *WHY1* sequences from
110 species were included in the angiosperm-wide alignment (see Alignment section below; Table
S1).

RNAseq and *De Novo* Assembly of Transcripts

Corallorhiza tissues used for RNAseq are those referred to in Sinn and Barrett (2020). In
brief, tissues were categorized as either aboveground (combined stem, flower, and ovary tissues) or
belowground (rhizome tissue, including fungal tissue), and three biological replicates of each tissue
type were sequenced for all four species, with the exception of aboveground *C. maculata* (two

samples) and belowground *C. striata* (two samples) for which library preparation of limited material
was not successful. *De novo* assembly of transcripts from each *Corallorhiza* species was conducted
using Trinity (version v2.13.2, Grabherr et al., 2011). Reads were trimmed using Trimmomatic
(version 0.36, Bolger et al., 2014) with default settings to remove sequencing adapters and low-
quality bases from the read ends. Trinity was provided with a samples file with biological replicate
relationships, and SS library type was set to RF. Trimmed reads were mapped to each assembled
transcript using the splice-aware read mapper BBMAP (version 38.96; Bushnell, 2022) with default
settings, with mapped reads output to SAM format and converted to sorted BAM-formatted files
using SAMtools (version 1.15; Li et al., 2009).

Identification of *WHY1* Transcripts

We used the HMMER suite (version 3.3.2; Eddy, 2011) to create a *WHY1* hidden markov
model (HMM) profile. The HMM profile was built with hmmbuild, which used the complete
angiosperm-wide *WHY1* nucleotide alignment. All assembled *Corallorhiza* transcripts were then
searched against the HMM model using nhmmer (Wheeler and Eddy, 2013). Default parameters were
used for both programs. All assembled isoforms of a transcript identified with the highest E-value
were considered as potential splicing variants of *WHY1*.

PCR Amplification

Primers for amplification of the entire *WHY1* sequence (Table S2) were designed using the
Geneious Prime (version 2020.2.4, https://www.geneious.com) plugin for Primer3 (version 2.3.7;
Untergasser et al., 2012). Oligos were synthesized by Integrated DNA Technologies (Coralville,
Iowa, USA). PCR amplification of *WHY1* for each *Corallorhiza* species was performed using the Taq
RED PCR Kit from Bio-Rad Industries (Hercules, California, USA). PCR was conducted in a
BioRad T100 Thermocycler using the following program: initial template denaturation at 95°C for
three minutes, followed by 30 cycles of denaturation at 95°C for 30 seconds, primer annealing at
52°C for 30 seconds, and template extension at 72°C for 30 seconds. The program ended with a final
extension at 72°C for five minutes and was held at 4°C until retrieval. PCR clean-up was performed
with Agencourt AMPure XP PCR Purification beads (Beckman Coulter Life Sciences; Indianapolis,
Indiana, USA). Purified DNA samples were quantified using a NanoDrop One^C spectrophotometer
(ThermoFisher Scientific; Waltham, Massachusetts, USA).

Nanopore Sequencing

Purified DNA samples were diluted to equimolar concentrations and libraries for long-read
sequencing were prepared according to the Oxford Nanopore Technologies (ONT; Oxford, United
Kingdom) End-Prep protocol (SQK-LSK109). The library of each *Corallorhiza* species received a
unique barcode for Nanopore sequencing using the Native Barcoding Expansion 1-12, PCR-free kit
(EXP-NBD104). The MinION SpotON flow cell (R9.4.1 FLO-MIN 106; ONT) was used for
sequencing. Basecalling was performed using the high-accuracy base calling algorithm as
implemented in the GPU version of Guppy (version 6.2.1+6588110a6; ONT) on an NVIDIA
GeForce RTX 2060 graphics card. Nanopore reads were mapped to the *Dendrobium catenatum*
WHY1 genomic sequence in Geneious Prime using Minimap2 (version 2.17, Li, 2018) with a K-Mer
length set to 15.

Alignment

We aligned all *Corallorhiza WHY1* isoforms using MAFFT (version 7.3.10; Katoh and
Standley, 2013), and the E-INS-i algorithm (Katoh et al., 2002) and the 1PAM scoring matrix, as a
Geneious Prime plugin. Visualization and structural annotation against the canonical *WHY1* sequence

of *Arabidopsis thaliana* (NCBI Q9M9S3) was also conducted using Geneious Prime. Consensus was
determined as majority consensus with a 0% threshold, meaning no minimum frequency was
required for a consensus character if the character was shared by most sequences. All *Corallorhiza*
WHY1 isoforms were translated in Geneious Prime to amino acid sequence and manually trimmed to
the correct ORF. ORF-trimmed *WHY1* translations were aligned with the ORF-trimmed *WHY1*
sequence of *Arabidopsis thaliana* (NCBI NM101308) to verify the presence of the expected
canonical reading frame.

A translation-aware alignment of the canonical form of *WHY1* representing lineages across
the angiosperms was generated using a two-step process. Nucleotides were first aligned and
translated using the frameshift-aware aligner MASCE2 (version 2.0.6; Ranwez et al., 2011) with
default parameters, which inserts gap characters necessary to maintain codon-based statements of
homology across the alignment. This approach was necessary due to the presence of frameshift
mutations in sequences from *Gastrodia elata*, *Epipogium aphyllum*, and *C. striata*. The MACSE2-
processed nucleotide and amino acid alignments were then refined using MAFFT and the E-INS-i
algorithm with a BLOSUM 80 substitution matrix.

An alignment including the complete genomic sequences of *WHY1* from *Phalaenopsis*
equestris (ASM126359v1) and *Dendrobium catenatum* (ASM160598v2) was also generated.
Phalaenopsis equestris (Cai et al., 2015) and *D. catenatum* (Zhang et al., 2016) are the closest
relatives of *Corallorhiza* with sequenced genomes. This DNA alignment was generated to identify
the introns of *WHY1* and to evaluate the exonic content of assembled transcripts. All *Corallorhiza*
isoforms and *Corallorhiza* Nanopore consensus sequences were aligned with the sequences of *P.*
equestris and *D. catenatum* using MAFFT (version 7.3.10; Katoh and Standley, 2013), as a Geneious
Prime plugin, and the E-INS-i algorithm (Katoh et al., 2002).

Visual Screening for Amino Acid Substitutions

Nonsynonymous substitutions within the angiosperm *WHY1* alignment were surveyed
visually in our amino acid alignments via Geneious Prime. Substitutions of interest included those
present exclusively in *Corallorhiza* species, mycoheterotrophs, and the Orchidaceae. The codons of
A. thaliana (Q9M9S3) and *P. equestris* that corresponded to the sites of nonsynonymous
substitutions in *WHY1* of interest were cross-referenced with the annotated *A. thaliana WHY1*
sequence for structure and the *P. equestris WHY1* genomic sequence as included in the DNA
alignment for exon location.

Phylogenetic Inference

IQ-TREE (version 1.6.12; Nguyen et al., 2015) was used to infer phylogenetic relationships
amongst the recovered *WHY1* sequences using automated model choice (Kalyaanamoorthy et al.,
2017), optimal partitioning assessment, and nearest neighbor interchange search was enabled. Node
support was estimated using 1,000 ultrafast bootstrap approximation replicates (Hoang et al., 2018).
Two partitions were defined, which comprise the signal peptide and the highly variable 5' portion of
the chain (positions 1 – 516) and the highly conserved chain region (517 – 1,122), identified using
functional annotations on *WHY1* sequence per *A. thaliana* (Q9M9S3).

Differential Expression

251 *In silico* analysis of differential expression was conducted using scripts provided as
252 components of the Trinity RNAseq pipeline (Haas et al., 2013). Transcript abundance was estimated
253 using the alignment-free estimation method as implemented in Salmon (version 1.2.0, Patro et al.,
254 2017). The strand-specific library type was set to RF, for reverse-forward. Salmon output files

included transcript abundance estimates at both the transcript and gene level. Matrices were built using the abundance_estimates_to_matrix.pl script for both transcript counts and gene expression. Differential expression analysis was run using the R (R Core Team, 2019) package edgeR (Bioconductor version 3.10, Robinson et al., 2010) via the run_DE_analysis.pl script, which identified differential expression using biological replicates of tissues across the replicate conditions aboveground and belowground. TPM (transcripts per million) values were normalized using the TMM (trimmed mean of M-values; Robinson and Oshlack, 2010) approach in order to normalize expression values while maintaining comparability of expression between samples.

Selection Analyses

The *WHY1* nucleotide alignment was tested for statistically significant changes in selection regime using four methods implemented in the command-line, multithreaded version of the Hypothesis Testing using Phylogenies suite (HyPhy; version 2.5.39; Pond and Muse, 2005). We tested for significant change of selective regime using five test branch sets against null reference branch sets comprising all other species in the phylogram: 1) *Corallorhiza* species; 2) *C. maculata + C. striata*; 3) *Corallorhiza + Epipogium aphyllum + Gastrodia elata*; 4) *C. maculata + C. striata + E. aphyllum + G. elata*; 5) *E. aphyllum + G. elata*. The same *Amborella trichopoda*-rooted maximum likelihood *WHY1* topology inferred with IQ-tree was used for all analyses.

RELAX (Wertheim et al., 2015) was used to test for evidence of relaxed selection. RELAX breaks each codon into its three component sites, each with an assigned omega class. Values for omega are calculated as Dn/Ds ratios, using the calculation for the reference branches as the null hypothesis. The value k is the selection intensity parameter and is an exponent value on omega. The alternative model fits a value for k that changes the rate to fit with the test branches. Evidence for intensified selection strength along test branches is indicated by a significant result where the value of k is greater than 1 ($k > 1$, $P < 0.05$). Evidence for relaxed selection along test branches is indicated by a significant value of k less than 1 ($k < 1$, $P < 0.05$). Strength of selection was assessed simultaneously for all species using Fast Unconstrained Bayesian Approximation (FUBAR, Murrell et al., 2013). FUBAR can detect weak, yet pervasive, purifying or diversifying selection at the codon level (posterior probability > 0.90) without the use of test and reference branch sets.

The Branch-Site Unrestricted Statistical Test for Episodic Diversification (BUSTED) was used to test for evidence of gene-wide positive selection (Murrell et al., 2015). BUSTED uses three omega classes defined as $\omega1 \leq \omega2 \leq 1 \leq \omega3$. The $\omega1$ class is the proportion of sites with a very low Dn/Ds ratio. The $\omega2$ class is the proportion of sites just below 1, and $\omega3$ are sites above 1. A value of 1 suggests selective neutrality, and therefore defines the null, or Constrained, model. BUSTED then calculates the log-likelihood of the data for each of the null and alternative models. These ratios are calculated for each site and are called evidence ratios (ERs). They are used as a threshold (χ^2 distribution, $P < 0.01$) but are not a valid test for site-specific likelihood. The null model is rejected if at least one site on a test branch experienced positive selection. Evidence for positive, or diversifying, selection in a gene of a test branch is indicated by a rejection of the null model.

The Adaptive branch-site random effects likelihood (aBSREL) test was used to test for signal of episodic positive selection. aBSREL infers ω values at both the level of sites and branches and can account for rate heterogeneity inherent to complex evolutionary scenarios by partitioning these values into multiple rate classes per branch. aBSREL was conducted in exploratory mode, where all branches were tested and p-values were Holm-Bonferroni corrected, and in the more sensitive *a priori* mode to test for episodic positive selection in each test branch set.

3   Results

De novo Assembly and Identification of *WHY1*

The hmmer suite revealed a single, Trinity-identified gene model and its isoforms as representing *WHY1* from *de novo* assembled transcripts for each *Corallorhiza* species. Trinity assembled a single isoform representing the expected canonical CDS (coding DNA sequence) of *WHY1* for all but *C. striata*, the latest-stage fully-mycoheterotrophic species of *Corallorhiza* sampled. However, mapping reads to each isoform with BBMAP revealed that a five nucleotide indel, the absence of which results in a premature stop codon, was differentially present or absent in the reads of each *Corallorhiza* species (NCBI BioProject PRJNA984634). Trinity differentially incorporated that indel, hereafter referred to as the GTGAA indel, into the isoform pool of each *Corallorhiza* species. The absence of the GTGAA indel in Trinity-assembled isoforms of *C. striata* precluded the recovery of the canonical ORF, but read mapping supports the presence of the five nucleotides necessary to recover the expected ORF in *C. striata*, at a rate of 67.2% of reads in isoform 2 and 76.1% of reads in isoform 4. Those data support that the canonical variant of *WHY1* is also expressed. We analyzed the *C. striata* isoforms as assembled, rather than manually modifying the *C. striata* transcripts to conform to a hypothesized canonical sequence. The inclusion of those five nucleotides would not alter any results presented here, aside from whether a canonical *WHY1* isoform is transcribed in *C. striata*.

The canonical *WHY1* ORF in *C. trifida*, *C. wisteriana*, and *C. maculata* is 795 nucleotides in length and comprises 265 amino acids, two fewer amino acids than that of *A. thaliana*. Non-canonical splicing variants were found in the isoform pools of each *Corallorhiza* species. A splicing variant containing a 79-nucleotide-long sequence of 96.6% mean pairwise identity was recovered from *C. wisteriana*, *maculata*, and *striata*, at nucleotide position 232. The second major variant is the GTGAA indel discussed earlier, which is variously present at nucleotide position 514 in all four species. A third variant identified from the isoform pool of *C. striata* represents a modification of the 3' end of the ORF, with sequence of no obvious homology to that of *WHY1* immediately downstream of the previously discussed five-nucleotide indel that was variously present in our read pools throughout *Corallorhiza*.

Alignment

MAFFT alignment of all raw Trinity-assembled *Corallorhiza WHY1* isoforms resulted in a matrix of 1,302 positions. Pairwise percent identity across isoforms was 81.4% and 88.2% and gaps comprised 11.7% and 8.1% of character states for all isoforms and with the exclusion of the early terminating isoform 1 of *C. striata*, respectively. Alignment of ORF-trimmed transcripts resulted in a matrix of 874 positions (Figure 2). Percent pairwise identity amongst isoforms within a species was highest for *C. trifida* (99.5%) and lowest for *C. striata* (70.6%). A 79-nucleotide-long indel of 96.6% pairwise identity was identified in at least one isoform assembled from all *Corallorhiza* species except *C. trifida*. Translation-aware alignment of ORF-trimmed canonical *Corallorhiza WHY1* sequences, with the differentially present GTGAA indel manually inserted into the otherwise canonical sequence of *C. striata* resulted in an alignment of 795 characters of 98.0% pairwise identity and no gaps. The highest pairwise percent identity was 99.37% between *C. wisteriana* and *maculata* and the lowest was 96.98% between *C. trifida* and *striata*.

The *WHY1* genomic DNA alignment of *Corallorhiza* isoforms, *Corallorhiza* Nanopore sequences, and *P. equestris* and *D. catenatum* genomic sequences had a total length of 12,000 nucleotides (Figure 3), in which the assembled canonical isoforms from each *Corallorhiza* species

contain the expected exons of *WHY1*. One non-canonical isoform of *C. wisteriana*, *C. maculata*, and
C. striata each contained intron 1, for which pairwise percent identity was 96.6% across those
species. Percent pairwise identity of intron 1 between that of *D. catenatum* and *P. equestris* was
82.3%, and similarity between retained introns and that of *P. equestris* ranged from 78.5% to 82.3%
for *C. maculata* and *striata*, respectively. The Nanopore-sequenced and Trinity-assembled *WHY1*
intronic sequences of *C. striata* were identical, while those of *C. maculata* were 94.9% similar, due to
the relative lack of two thiamine nucleotides in the Nanopore sequence.

The angiosperm-scale *WHY1* nucleotide alignment contained *WHY1* sequences from 110
species, including 22 orchid species. The total length of the alignment was 1,122 positions, of which
150 were of identical states across all species. Two partitions roughly corresponding to the transit
peptide and chain regions as annotated in *A. thaliana* (Q9M9S3) were conspicuously visible in the
consensus sequence of the alignment. The first was a highly variable, lineage-specific portion of
sequence ranging from positions 1 – 516 in the alignment. The second was a highly conserved
portion spanning positions 517 – 1,122. The mean pairwise percent identity of the transit region of
Orchidaceae was 51.3%, while that of *Apostasia shenzhenica* compared to either *G. elata* or *E.*
aphyllum was 21.5% and 31.1%, respectively. The mean pairwise percent identity of the transit
region of *Corallorhiza* was 94.4%. The mean pairwise percent identity of the four *Corallorhiza*
species in the alignment was 97.7%, and 81.0% across the Orchidaceae. The grasses had the lowest
mean percent pairwise similarity of any clade, which was 49.36% when each sequence was compared
to each non-grass species in the alignment.

The angiosperm-wide *WHY1* amino acid alignment comprised 376 positions, of which 48
were of identical states across all species. The two partitions in the nucleotide alignment
corresponding to functional annotation in *A. thaliana* (Q9M9S3) were more conspicuous in the amino
acid alignment. Positions 1 – 115 and 116 – 376 corresponded to the transit and chain regions of *A.*
thaliana (Q9M9S3), respectively. The pairwise percent identity of the four *Corallorhiza* species in
the alignment was 97.4% *WHY1*, and across the Orchidaceae it was 77.8%.

Nanopore Sequencing

Nanopore sequencing of genomic *WHY1* sequence from three of the four *Corallorhiza* species
confirmed that the unique sequence in the non-canonical transcripts of *C. wisteriana*, *C. striata*, and
C. maculata represented retention of a *WHY1* intron 1. Sequencing of *WHY1* amplicons generated
read pools ranging from 25,546 to 74,261 reads in *C. trifida* and *C. maculata*, respectively. Mapping
of Nanopore reads against the *D. catenatum* genomic *WHY1* sequence resulted in a mean coverage
depth for the exon 1-2 region ranging from 4,034.9 to 10,617.6 in *C. trifida* and *C. striata*,
respectively (Table S2). Library preparation for Nanopore sequencing of *C. wisteriana WHY1*
amplicons was deemed unsuccessful since the resulting sequence pool contained presumably off
target reads that precluded confident assembly of a *WHY1* consensus sequence from that sample.
Alignment of the Nanopore-sequenced *WHY1* amplicon consensus reads with the *Corallorhiza*
transcript isoforms and the full *WHY1* genomic sequences of *P. equestris* and *D. catenatum* provided
evidence that the non-canonical isoforms of *WHY1* in *Corallorhiza* were a result of alternative
splicing (Figure 4). At least one non-canonical isoform from all but *C. trifida* contains intron 1 of
WHY1.

The sequences obtained from *Corallorhiza* species via Nanopore and RNAseq shared a high
degree of similarity with each other and the *WHY1* sequence in the previously published *P. equestris*
and *D. catenatum* genomes. Intron 1 of *WHY1* was found to have a pairwise percent similarity of

82.3% between *P. equestris* and *D. catenatum*. *Corallorhiza WHY1* intron 1 sequences obtained via
Nanopore sequencing and RNAseq had a mean pairwise percent similarity of 96.9%, while that of *C.*
striata obtained via both RNAseq and Nanopore sequencing had a pairwise percent similarity of
82.3% with *P. equestris*.

Select Amino Acid Substitutions of Interest

Substitutions exclusive to mycoheterotrophic species were evident and in some cases were
exclusive to fully-mycoheterotrophic species. The transit regions of both *E. aphyllum* and *G. elata*
contained some substitutions that shared little homology with the remainder of Orchidaceae. A
phenylalanine-glycine residue at positions 76-77 was exclusive to *C. trifida*, while a leucine-arginine
residue was present for the remainder of Orchidaceae at those positions. While pairwise percent
identity of positions 76-77 was only 13.3% throughout angiosperms, the leucine-arginine residue in
the Orchidaceae had a pairwise percent identity of 90.9% when including the residue from *C. trifida*
and was identical that taxon was excluded. Substitutions of single amino acids comprising the transit
region not found in other orchids were identified in both *C. wisteriana* (positions 46, 95) and *C.*
trifida (position 113), but identical amino acids were identified in some non-orchid autotrophs.

At least one substitution that could impact *WHY1* conformation was identified in
Corallorhiza. The terminal codon of an alpha helix annotated in the structure of *A. thaliana* (9M9S3)
WHY1 (position 336) has been substituted to an isoleucine in *C. trifida* and to a phenylalanine in *C.*
wisteriana, *maculata*, and *striata* (Figure 5). We infer that the ancestral state of position 336 is
leucine, on the basis of that codon state for 89.1% of sampled taxa, including *Amborella trichopoda*.
The only other substitutions at that position were to isoleucine, which was identified in eight
autotrophic angiosperms outside the Orchidaceae, and in *C. trifida*. Additionally, a substitution from
alanine to valine at position 359 was exclusive to *C. striata*, which represents the only occurrence of
that state for this position for 109 other angiosperm species, and the only subgeneric polymorphism
observed in that position.

A five-nucleotide indel resulting in a frameshift (amino acid alignment positions 269 & 270)
was found in *C. striata*, but the pairwise percent identity of downstream sequence with that of *C.*
maculata was high (97.8%) and reads containing the corresponding five nucleotides were identified
in the RNAseq read pool. In fact, RNAseq reads containing the indel were found in the read pools of
all *Corallorhiza* species, suggesting that isoform diversity was conservatively interpreted by our
methods.

We found that the chain region of *E. aphyllum* contained a glycine to arginine substitution in
the characteristic *WHY1* ssDNA binding motif (positions 188-193), a site that was otherwise
conserved throughout the remainder of samples. Additionally, we found that the *E. aphyllum*
sequence contained a residue comprising seven amino acids (positions 294 – 300), the last of which
was a premature stop codon. MAFFT resolved those seven amino acids as an insertion with no
homology to other angiosperm sequences. Six substitutions downstream of that premature stop codon
are exclusive to *E. aphyllum*.

Phylogenetic Inference
Both the Eudicots and Monocots were recovered as monophyletic (BS = 100%). Orchidaceae
was recovered as a monophyletic group (BS = 94%), with *D. catenatum*, *G. elata* and *E. aphyllum* as
an early-diverging paraphyletic grade within the Epidendroids (BS = 98%). *Corallorhiza* species
were recovered as a monophyletic group (BS = 100%) with *C. trifida* sister to *C. striata* + (*C.*

wisteriana, *C. maculata*), rather than the expected relationships recovered in previous work as
depicted in Figure 6.

Differential Expression

Canonical isoforms of *WHY1* were most highly expressed in all four species (see Table XX),
but statistically significant elevated expression in aboveground tissues relative to those belowground
was only detected in *C. trifida*. Gene-level expression of *WHY1* across both tissues was highest for *C.*
wisteriana (244.5) and *trifida* (175.4) and lowest for *C. maculata* (131.9) and *striata* (95.3).
Expression of the canonical isoform of *WHY1* in *C. trifida* ranged from 4.15 to 9.64 TMM in
belowground tissues (median = 7.17 TMM) and from 31.353 to 56.389 TMM in the aboveground
tissues (median = 43.273 TMM). The log fold-change value was 2.56 between belowground and
aboveground tissues for the canonical isoform of *C. trifida*, with a P value of 0.00018 and a false
discovery rate (FDR) of 0.00374. Expression of the *WHY1* nearly canonical isoform in *C. striata*
ranged from 5.547 to 16.859 TMM in belowground tissues (median = 12.31 TMM) and 17.28 and
30.538 TMM in aboveground tissues (median = 23.909 TMM). The *C. striata* canonical *WHY1*
isoform had a log fold-change value of 0.396 between belowground and aboveground tissues, with a
P value of 0.69 and FDR of 0.91. Expression of the canonical isoform of *WHY1* in *C. wisteriana*
ranged from 9.256 to 23.023 TMM in belowground tissues (median = 15.663 TMM) and 34.792 to
74.955 TMM in aboveground tissues (median = 60.80 TMM). The *C. wisteriana* canonical isoform
had a log fold-change value of 1.96 between belowground and aboveground tissues, with P value of
0.02 and FDR of 0.28. The expression of the canonical isoform in *C. maculata* had values of 11.064
and 13.947 TMM in belowground tissues (median = 12.51 TMM) and ranged from 26.179 to 45.999
TMM in aboveground tissues (median = 33.01 TMM). The *C. maculata* canonical isoform had a log
fold-change value of 1.17 between belowground and aboveground tissues, with P value of 0.16 and
FDR of 0.62.

The expression of non-canonical isoforms assembled from the read pools of all four species
varied across tissues and samples, with some tissues or individuals not expressing splicing variants,
and were not statistically different between the two tissue types in any of the four species. The *C.*
trifida non-canonical isoform was expressed at relatively low levels across all aboveground and
belowground tissues, the TMM of which ranged from 1.02 belowground to 4.374 aboveground.
Contrastingly, the TMM of the *C. wisteriana* non-canonical isoform ranged from 0 to 2.017 and was
not detected in two of the belowground replicates for this species. The noncanonical *C. maculata* and
C. striata isoforms were sporadically expressed across tissue types, with the TMM values of one
isoform of *C. maculata* ranging from 0 to 3.564 and those of *C. striata* ranging from 0 to 4.471. Even
in the case of the *C. striata* isoform for which the TMM value was 4.471 in one belowground sample,
the value was either zero or less than 1 in the remainder of samples.

Selection Analyses

The RELAX test inferred statistically significant signal for relaxation of selection pressure for
the following test branch sets: *Corallorhiza* + *E. aphyllum* + *G. elata* (p = 0.001); *C. maculata* + *C.*
striata + *E. aphyllum* + *G. elata* (p = 0.002); and *E. aphyllum* + *G. elata* (p = 0.001). However, runs
of RELAX analyzing test branch sets comprising only *Corallorhiza* (p = 0.30) or *C. maculata* + *C.*
striata (p = 0.62) did not infer significant signal for selection relaxation.

The FUBAR analysis inferred that 254 of 373 codons were under pervasive purifying
selection (posterior probability threshold ≥ 0.9) and that no codons were under pervasive diversifying
selection. Likewise, the BUSTED analyses did not infer statistically significant signal of gene-wide

episodic diversifying selection for any test branch set, where p-values ranged from p = 0.15 for *E.*
aphyllum + *G. elata* to p = 0.50 for *Corallorhiza*.

The aBSREL analysis recovered signal of episodic diversifying selection in seven of 217
branches in the *WHY1* tree (Figure 6) but did not infer statistically significant signal in any test
branch set during analyses conducted in *a priori* mode. The terminal node leading to *Sorghum*
bicolor (p = 2.76 x 10^{-6}) was the only branch in the monocots that showed significant signal of
episodic diversifying selection. Within the eudicots, the branches leading to *Prunus avium* (p = 2.5
x10^{-5}), the *Populus* clade (p = 1.17 x 10^{-7}), the *Quercus* clade (2.35 x 10^{-3}), the *Nicotiana* clade (p =
5.42 x 10^{-3}), the *Coffea* clade (6.96 x 10^{-5}), and the asterid clade (p = 3.7 x 10-4) were inferred to
contain signal of episodic diversifying selection.

4 Discussion

Our work represents the largest-scale investigation of *WHY1* evolution to date and reveals
strong phylogenetic signal for this gene across multiple taxonomic levels in the angiosperms.
Phylogenetic methods have emerged as the gold-standard for the inference of gene orthology
(Münster et al., 1997; Emms and Kelly, 2015; Emms and Kelly, 2019) by which hundreds of genes
have been determined to not only be highly conserved across plant lineages, but to exist as low- or
single-copy (Wu et al., 2006; Duarte et al., 2010; De Smet et al., 2013). The inferred evolutionary
relationships of *WHY1* (Figure 6) were largely congruent with our contemporary understanding of
angiosperm phylogenetic relationships and recent inferences based on genomic-scale datasets
(Angiosperm Phylogeny Group, 2016; Guo et al., 2021; Zhao et al., 2021). Likewise, the inferred
relationships within the Orchidaceae were largely consistent with those inferred in other studies (e.g.,
Givnish et al., 2016; Pérez-Escobar et al., 2021; Serna-Sánchez et al., 2021), apart from *D.*
catenatum, *G. elata*, and *E. aphyllum* forming an early-diverging, paraphyletic grade within the
Epidendroids. However, our recovery of *C. trifida* as sister to the remainder of the species sampled
from *Corallorhiza*, rather than *C. striata*, was not congruent with relationships inferred previously on
the basis of other loci and even genomic-scale datasets (Barrett and Freudenstein, 2008; Barrett et al.,
2014; Barrett et al., 2018). We are not surprised to infer discordance between the evolutionary history
of *WHY1* and that of the phylogenetic history of *Corallorhiza*, since it is accepted that the
evolutionary history of any gene can differ from that of the genome within which it is found (Pamilo
and Nei, 1988; Maddison, 1997). That said, the recovery of the *Corallorhiza* species with the most
intact plastome as sister to those with more degraded plastomes may indicate that the gene-species
tree discordance we infer is due to functional convergence in *WHY1* sequence. For example, the
phenylalanine substitution at position 336 shared by all *Corallorhiza* species aside from *C. trifida*
may affect protein conformation and therefore *WHY1* function. Our findings of general congruence
between the evolutionary history of *WHY1* across angiosperms supports the gene as single- or low-
copy across the 110 angiosperm taxa sampled.

Differences in the degree of nucleotide conservation were evident amongst *WHY1* exons, with
the transit peptide region consistently the most divergent throughout the lineages sampled. The
diversity and evolution of transit peptides, and the apparent discrepancy between our perception of
their functional importance and high sequence divergence, have long been of interest (Bruce, 2001;
Patron and Waller, 2007; Christian et al., 2020). While the functional nature of a transit peptide
might lead to an expectation of conservatism, low sequence similarity and patterns of mutation that
we describe in *WHY1* are emerging as generalizable properties of plant transit peptides. For example,
Christian et al. (2020) found that the transit peptides in the genomes of 15 genera sampled throughout
the angiosperms had a mean pairwise percent identity of just 37.9% and that random indels drive

transit peptide evolution. Our results provide evidence that the transit peptide of *WHY1* evolves
similarly, with a pairwise percent identity of 32.5% across the angiosperms, and with evident
substitutions and indels downstream of a homologous start codon being the most likely drivers of
divergence in the gene region. In contrast, the portion of the chain encoding the ssDNA binding motif
was the most conserved with a mean percent pairwise similarity of 72.1%. The only sizable indel
observed in the gene region was a seven-codon insertion in *E. aphyllum*, a late-stage
mycoheterotrophic orchid. It is likely that *E. aphyllum WHY1* results in a non-functional product,
given that it encodes a premature stop codon that would result in a protein 70 amino acids shorter
than that of any other sampled species. Interestingly, *WHY1* was more conserved overall at the
nucleotide level amongst *Corallorhiza* species than were either expression patterns across tissues or
exon inclusion in sequenced mRNAs across the trophic gradient. Amongst autotrophic species,
nucleotide divergence of *WHY1* is particularly pronounced in the grasses, with the mean pairwise
percent identity amongst all members of that clade versus the remainder of samples being more than
5% lower than that of the same comparison made for either of the late-stage mycoheterotrophic
species sampled. The high substitution rate we inferred for *WHY1* in the grasses is interesting
because the plastomes of many species in the clade are known to contain inversions and structural
heteroplasmy within individual plants, of which the latter has only recently been described from their
relatives the Cyperaceae (Doyle et al., 1992; Lee et al., 2020). The contrasting levels of nucleotide
conservation in the transit and chain portions of *WHY1* across lineages suggest that lineage-specific
functions of the transit peptide sequence could be a fruitful line of investigation, especially given that
Christian et al. (2020) identified bias in amino acid usage between the plastid transit peptide
sequences of monocot and eudicot lineages.

Alignment of *WHY1* revealed lineage-specific substitutions at sites inferred to be involved in
structural conformation in both poorly and highly conserved gene regions. For example, the transit
regions of both late-stage mycoheterotrophic species contained indels with little homology with the
remainder of samples, and the sequence of *E. aphyllum* contained a premature stop codon in the
typically highly conserved chain region followed downstream by four autapomorphic amino
substitutions in an eight amino acid span. We identified substitutions that could underlie functional
change in each *Corallorhiza* species, the most interesting of which was a substitution involving an
alpha helix inferred in previous work conducted in *A. thaliana* (Cappadocia et al., 2013). The
phylogenetic framework provided by our analysis suggests that the plesiomorphic codon state of
position 336 was a leucine, which was substituted to an isoleucine in *C. trifida* and to a phenylalanine
in the remainder of *Corallorhiza* sampled. We infer that the phenylalanine substitution in
Corallorhiza species aside from *C. trifida* would result in the introduction of a benzene ring in which
there were previously only aliphatic hydrocarbons. While our work revealed many nonsynonymous
substitutions throughout *Corallorhiza* sequences, we did not identify nucleotide substitutions that
obviously result in loss of function in these transitional mycoheterotrophic species.

Statistically significant shifts of selective regime were detected for late-stage
mycoheterotrophic and some non-orchid autotrophic lineages. Our inference that 68.1% of *WHY1*
codons are under pervasive purifying selection and none are under pervasive positive selection,
despite the relatively low levels of nucleotide conservatism we documented, are congruent with the
critical function of the gene across the angiosperms as has been documented in model and
agricultural species in previous work. While we inferred disproportionately high substitution rates for
WHY1 in the grasses, we did not find evidence that divergence of those sequences was associated
with diversifying selection. Despite the lack of pervasive positive selection in *WHY1* across the
angiosperms, our identification of significant episodic diversifying selection in seven of 100
autotrophic lineages sampled suggests that a lineage-specific adaptive role of *WHY1* may be

relatively common, though similar signal was not identified in sampled mycoheterotrophs. The lack
of diversifying signal in any of our mycoheterotrophic species does not support neofunctionalization
of the gene, which we find interesting given the myriad functions of *WHY1* and the diversifying
signal identified in lineages across the angiosperms. Our findings together suggest that relaxation of
WHY1 selective constraint occurs after the transition to fully mycoheterotrophy, as significant signal
of relaxed selection was only detected in branch sets containing *E. aphyllum* and *G. elata*, two late-
stage fully mycoheterotrophic orchids.

Our analyses reveal that the expression of *WHY1* in *Corallorhiza* can differ by both tissue
type and across the mycoheterotrophic gradient. Multiple studies have induced and characterized the
effects of differential expression of *WHY1* by exposure to biotic and abiotic stimuli, implicating roles
for the gene ranging from mediating drought stress (Ruan et al., 2022) to pathogen response
(Desveaux et al., 2000; Sun et al., 2023). A minimum level of *WHY1* expression could be expected
due to roles of *WHY1* that are not involved in photosynthesis, such as the maintenance of telomere
length of nuclear chromosomes (Yoo et al., 2007). Our analyses are the first to characterize *WHY1*
expression in non-model or non-cultivated plant species, and therefore baseline expectations for
tissue-specific expression levels for wild species have not yet been established. However, the values
observed in *Corallorhiza* are within the expressive ranges reported in studies of *A. thaliana* (4-76
TPM; Liu, et al., 2012; Mergner et al., 2020) and *Solanum tuberosum* (7-50 TPM; The Potato
Genome Sequencing Consortium, 2011). Additionally, the Klepikova *Arabidopsis* Atlas (Klepikova
et al., 2016) and 1,122 tissue-specific samples available via the *Arabidopsis* RNA-seq Database
(http://ipf.sustech.edu.cn/pub/athrdb/; accessed 8 March 2023) evidence that *WHY1* expression
should be expected to be lower in roots or rhizomes than in leaves, which is congruent with
expression patterns in *C. trifida*. Previous studies reporting *WHY1* expression have been conducted in
species with larger individuals with typical, non-reduced morphologies allowing for finer-scale
investigations of tissue-specific expression than can be conducted in *Corallorhiza*, due to a lack of
leaf laminae and roots in the latter. However, the patterns of expression we found across
Corallorhiza tissues are congruent with those known for *WHY1*. While gene-level expression across
tissues was highest for *C. wisteriana* and *trifida*, tissue-specific expression was only statistically
significant in *C. trifida*, after correcting for repeated testing. Our findings characterizing similar
WHY1 expression amongst the belowground tissues of *Corallorhiza* provide for a hypothesized
minimum expression level of canonical *WHY1*, which is similar to aboveground levels of expression
in *Corallorhiza* species aside from *C. trifida.* Taken together, our results provide evidence for a
trajectory beginning with differential expression of *WHY1* between aboveground and belowground
tissues of the most photosynthetically capable *Corallorhiza* species to similar expression levels
between above and belowground tissues of the latest-stage mycoheterotrophic members of the genus.
These results suggest that downregulation of *WHY1* in aboveground tissues is correlated with
increased mycoheterotrophy, but that it is unlikely to underlie the transition to mycoheterotrophy
since the tissue-level expression patterns of the gene in *C. trifida* remain like those described from
autotrophic plants.

Our work is the first to characterize alternative splicing of, and intron retention in, *WHY1*.
Approximately 70% of plant genes with multiple exons can be expected to be alternatively spliced
(Chamala et al., 2015), and intron retention is a common form of alternative splicing in plants (Ner-
Gaon et al., 2004). Our finding of intron retention in one isoform from each *Corallorhiza* species
aside from *C. trifida* is the first such event described for the gene. It has long been recognized that
intron retention is most common in transcripts of genes like *WHY1* which serve roles related to
photosynthesis and stress response (Ner-Gaon et al., 2004), a finding also supported by work
investigating the effects of plant stressors on levels of alternative splicing (Filichkin et al., 2018;

Jabre et al., 2019). In fact, tissue-specific differential intron retention has been shown to be an inducible stress response in *Populus trichocarpa* (Filichkin et al., 2018). Modification of *WHY1*, including inserted sequence, has long been used to study the effects of mutations on the function of the gene and to induce knockouts (Desveaux et al., 2004; Yoo et al., 2007; Maréchal et al., 2009), work which helped identify the many pathways that *WHY1* is involved in. For example, Yoo et al., (2007) and Maréchal et al. (2009) leveraged knockouts caused by T-DNA insertions into *WHY1* to reveal the critical role the gene plays in maintaining telomere length and plastome stability in *Arabidopsis thaliana*, respectively. Similarly, Prikryl et al. (2008) and Qui et al., (2022) found that double knockout *WHY1* mutants were characterized by a lethal albino phenotype after the development of a few leaves, with Qui et al. (2022) also characterizing divergent splicing and mRNA editing of plastid genes in *WHY1* mutants. Our discovery of intron-retaining transcripts from all *Corallorhiza* species aside from *C. trifida*, and more transcriptional isoforms in *Corallorhiza* species in later stages of mycoheterotrophy, reveals a negative correlation between increased splicing variation and both plastome stability and chlorophyll concentration in tissues (Barrett et al., 2014). We find it unlikely that the intron-containing isoforms result in functional products, since a premature stop codon results from intron retention. The recovery of non-canonical and intron-retaining *WHY1* isoforms across individuals and tissues of *Corallorhiza* could be signal of idiosyncratic spliceosome regulation in mycoheterotrophic species, epitranscriptomic differences (Jabre et al., 2019), differential responses to stress in sampled tissues (Filichkin et al., 2018), or the expression of multiple, divergent copies of the gene. However, we propose that reduced fidelity in the spliceosome of a mycoheterotrophic plant is the most likely cause of the observed splicing variation, given the phylogenetic and sequencing data at hand. We predict that changes in the expression and splicing of *WHY1* across *Corallorhiza* are likely due to the alteration of one or more pathways involved in gene regulation, since we did find nucleotide-level changes that could be responsible.

Our work is the first to characterize the evolution of a transcription factor that could impact the genetic and phenotypic changes that occur along the path to full mycoheterotrophy. The roles that *WHY1* plays in plastome stability (Parent et al., 2011; Lepage, 2013), defense responses (Lin et al., 2020), and leaf senescence (Lin et al., 2019), together position the gene as a worthwhile target for the study of the molecular underpinnings of the transition to mycoheterotrophy. Heterotrophy in plants is associated with genomic restructuring, where a trend of plastome contraction and nuclear genome expansion via rampant repetitive element accumulation has commonly been observed (Barrett et al., 2014; Lyko and Wicke, 2021). Dramatic reductions of plastome length and gene content of mycoheterotrophic plants have been documented, ranging from minimal degradation in early-transitional orchids such as *Corallorhiza* (Barrett and Davis, 2012) to pronounced degradation in late-transitional orchids such as *Epipogium* and *Pogoniopsis* (Schelkunov et al., 2015; Klimpert et al., 2022). Our finding of putatively functional but increasingly alternatively spliced *WHY1* across the *Corallorhiza* trophic gradient is not necessarily surprising, given the minimally destabilized plastomes of the group. For example, Barrett and Davis (2012) found that the plastome of *C. striata*, the most destabilized of the *Corallorhiza* species included here, is only about 6% reduced relative to that of a leafy, autotrophic relative. Despite their relatively intact states, *Corallorhiza* plastomes are in various stages of degradation (Barrett et al., 2014), and our work here reveals a negative correlation between both increased putatively aberrant splicing and nucleotide-level divergence of *WHY1* with plastome stability across the sampled species. Likewise, the plastomes of *Gastrodia elata* and *Epipogium aphyllum*, both late-stage mycoheterotrophs for which our analyses show that *WHY1* contains premature stop codons and significant signal of relaxed selection, are both extremely reduced and syntenically disrupted (Yuan et al., 2018; Chen et al., 2020; Xu et al., 2021). Taken

together, our findings provide evidence of a negative correlation between increased divergence in sequence, splicing of *WHY1*, and plastome stability in early- to late-stage mycoheterotrophic orchids.

5 Conclusions

Our work showcases the rich opportunities afforded by mycoheterotrophic plants not just for the study of the evolution of *WHY1* but for any gene of which homozygous knockouts can result in a fatal phenotype in autotrophic plants. Continued investigation of non-autotrophic plant lineages promises to fill gaps in our understanding of the precursors and consequences of genomic instability, and even the minimum gene space of land plants. Here we presented findings of non-synonymous nucleotide substitutions in functionally annotated regions in *Corallorhiza WHY1* sequence, and a high degree of divergence in *WHY1* in late-stage fully mycoheterotrophic orchids. However, our results together suggest that changes to the expression and splicing of *WHY1* occur prior to the establishment of obviously deleterious genomic substitutions that would render the TF non-functional in late stage mycoheterotrophic orchids. In sum, our work supports a correlation between decreased expression and increased alternative splicing of *WHY1* with plastome degradation in a group of early-transitional mycoheterotrophic orchids but does not implicate divergent *WHY1* function as a primary factor in the transition from partial to full mycoheterotrophy.

6 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

7 Author Contributions

BS and CB contributed to the conception and design of the study. CB collected *Corallorhiza* samples for RNAseq. RM and BS collected all publicly available samples, conducted wet-lab work, Nanopore sequencing, and bioinformatic analyses. RM wrote the first manuscript draft. All authors contributed to manuscript revision and have read and approved of the final version.

8 Funding

Funding was provided by the Undergraduate Student Research Award and the Bert and Jane Horn Endowed Student Research Award from Otterbein University to RM; US National Science Foundation (DEB#0830020), the California State University Program for Research and Education in Biotechnology, and the West Virginia University Program to Stimulate Competitive Research to CB, and a Faculty Scholarship and Development Fund Award from Otterbein University to BS.

9 Acknowledgments

We would like to acknowledge John V. Freudenstein for supplying DNAs used for Nanopore sequencing, the West Virginia University Genomics Core Facility for their services and technical expertise. We also thank the USDA Forest Service and California State Parks for permission to collect material.

10 References

Barrett, C. F. and J. V. Freudenstein. 2008. Molecular evolution of *rbcL* in the mycoheterotrophic

coralroot orchids (*Corallorhiza* Gagnebin, Orchidaceae). *Molecular Phylogenetics and*
Evolution 47: 665-679.

Barrett, C. F., J. V. Freudenstein, D. Lee Taylor, and U. Kõljalg. 2010. Rangewide analysis of
fungal associations in the fully mycoheterotrophic *Corallorhiza striata* complex
(Orchidaceae) reveals extreme specificity on ectomycorrhizal *Tomentella* (Thelephoraceae)
across North America. *American Journal of Botany* 97: 628-643.

Barrett, C. F. and J. I. Davis. 2012. The plastid genome of the mycoheterotrophic *Corallorhiza*
striata (Orchidaceae) is in the relatively early stages of degradation. *American Journal of*
Botany 99(9): 1513-1523.

Barrett, C. F., J. V. Freudenstein, J. Li, D. R. Mayfield-Jones, L. Perez, J. C. Pires, and C.
Santos. 2014. Investigating the path of plastid genome degradation in an early transitional
clade of heterotrophic orchids, and implications for heterotrophic angiosperms. *Molecular*
Biology and Evolution 31(12): 3095-3112.

Barrett, C. F., S. Wicke, and C. Sass. 2018. Dense infraspecific sampling reveals rapid and
independent trajectories of plastome degradation in a heterotrophic orchid complex. *New*
Phytologist 218(3): 1192-1204.

Barrett, C. F., B. T. Sinn, and A. H. Kennedy. 2019. Unprecedented parallel photosynthetic losses
in a heterotrophic orchid genus. *Molecular Biology and Evolution* 36(9): 1884-1902.

Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* 30(15): 2114-2120.

Bruce, B. D. 2001. The paradox of plastid transit peptides: conservation of function despite
divergence in primary structure. *Biochimica et Biophysica Acta* 1541: 2-21.

Bushnell, B. B. 2022. bbtools. Retrieved from https://sourceforge.net/projects/bbmap/files

Cai, J., X. Liu, K. Vanneste, S. Proost, W.-C. Tsai, K.-W. Liu, L.-J. Chen, et al. 2015. The
genome sequence of the orchid *Phalaenopsis equestris*. *Nature Genetics* 47(1): 65-72.

Cai, L. 2023. Rethinking convergence in plant parasitism through the lens of molecular and

population genetic processes. *American Journal of Botany* (early access):10.1002/ajb2.16174
735

736   Cameron, D. D., K. Preiss, G. Gebauer, and D. J. Read. 2009. The chlorophyll-containing
737   orchid *Corallorhiza trifida* derives little carbon through photosynthesis. *New Phytologist*
738   183:358–364.
739

740   Cappadocia, L., A. Maréchal, J-S. Parent, and É. Lepage. 2010. Crystal structures of DNA
741   Whirly complexes and their role in *Arabidopsis* organelle genome repair. *The Plant Cell* 22:
742   1849–1867.
743

744   Cappadocia, L., J.-S. Parent, J. Sygusch, and N. Brisson. 2013. A family portrait: structural
745   comparison of the Whirly proteins from *Arabidopsis thaliana* and *Solanum tuberosum*. *Acta*
746   *Crystallographica Section* F69: 1207-1211.
747

748   Cecchin, M., S. Benfatto, F. Griggio, A. Mori, S. Cazzaniga, N. Vitulo, M. Delledonne, and M.
749   Ballattari. 2018. Molecular basis of autotrophic vs mixotrophic growth in *Chlorella*
750   *sorokiniana*. *Scientific Reports* 8:6465.
751

752   Chamala, S., G. Feng, C. Chavarro, and W. B. Barbazuk. 2015. Genome-wide identification of
753   evolutionarily conserved alternative splicing events in flowering plants. *Frontiers in*
754   *Bioengineering and Biotechnology* 3:33.
755

756   Chao, Y.-T., S.-H. Yen, J.-H. Yeh, W.-C. Chen, and M.-C. Shih. 2017. Orchidstra 2.0 - a
757   transcriptomics resource for the orchid family. *Plant and Cell Physiology* 58(1): e9.
758

759   Chen, S., X. Wang, Y. Wang, G. Zhang, W. Song, X. Dong, M. L. Arnold, W. Wang, J. Miao, W.
760   Chen, and Y. Dong. 2020. Improved *de novo* assembly of the achlorophyllous orchid
761   *Gastrodia elata*. *Frontiers in Genetics* 11: 580568.
762

763   Chernomor, O., A. von Haeseler, and B.Q. Minh. 2016. Terrace aware data structure for
764   phylogenomic inference from supermatrices. *Systematic Biology* 65(6): 997–1008.
765

766   Christian R. W., S. L Hewitt, G. Nelson, E. H. Roalson, and A. Dhingra. 2020. Plastid transit
767   peptides—where do they come from and where do they all belong? Multi-genome and pan-
768   genomic assessment of chloroplast transit peptide evolution. *PeerJ* 8:e9772.
769

de Mendoza, A., A. Sebé-Pedrós, M. S. Ŝestak, M. Matejčić, G. Torruella, T. Domazet-Lošo, and
I. Ruiz-Trillo. 2013. Transcription factor evolution in eukaryotes and the assembly of the
regulatory toolkit in multicellular lineages. *Proceedings of the National Academy of Sciences*
of the United States of America 110(50): E4858-E4866.

De Smet, R., K. L. Adams, K. Vandepoele, M. C. E. Van Montagu, S. Maere, and Y. Van de Peer.
2013. Convergent gene loss following gene and genome duplications creates single-copy
families in flowering plants. *Proceedings of the Academy of Sciences of the United States of*
America 110(8): 2898-2903.

Desveaux, D., C. Després, A. Joyeux, R. Subramaniam, and N. Brisson. 2000. PBF-2 is a novel
single-stranded DNA binding factor implicated in *PR-10a* gene activation in potato. *The*
Plant Cell 12: 1477-1489.

Desveaux, D., J. Allard, N. Brisson, and J. Sygusch. 2002. A new family of plant transcription
factors displays a novel ssDNA-binding surface. *Nature Structural & Molecular Biology* 9(7):
512-517.

Desveaux, D., R. Subramaniam, C. Després, J.-N. Mess, C. Lévesque, P. R. Fobert, J. L.
Dangl, and N. Brisson. 2004. A "whirly" transcription factor is required for salicylic acid
dependent disease resistance in *Arabidopsis*. *Developmental Cell* 6: 229-240.

Desveaux, D., A. Maréchal, and N. Brisson. 2005. Whirly transcription favors: defense gene
regulation and beyond. *TRENDS in Plant Science* 10(2): 1360-1385.

Doyle, J. J., J. I. Davis, R. J. Soreng, D. Garvin, and M. J. Anderson. 1992. Chloroplast DNA
inversions and the origin of the grass family. *Proceedings of the National Academy of*
Sciences of the United States of America 89: 7722-7726.

Duarte, J. M., P. K. Wall, P. P. Edger, L. L. Landherr, H. Ma, J. C. Pires, J. Leebens-Mack, and C.
W. dePamphilis. 2010. Identification of shared single copy nuclear genes in *Arabidopsis*,
Populus, *Vitis* and *Oryza* and their phylogenetic utility across various taxonomic levels. *BMC*
Evolutionary Biology 10: 61.

Eddy, S. R. 2011. Accelerated profile HMM searches. *PLoS Computational Biology* 7:
e1002195.

Emms, D. M. and S. Kelly. 2015. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome Biology* 16: 157.

Emms, D. M. and S. Kelly. 2019. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biology* 20: 238.

Felsenstein, J. 1985. Phylogenies and the comparative method. *The American Naturalist*
125(1):1-15.

Filichkin, S. A., M. Hamilton, P. D. Dharmawardhana, S. K. Singh, C. Sullivan, A. Ben-Hur, A.
S. N. Reddy, and P. Jaiswal. 2018. Abiotic stresses modulate landscape of *Popular*
transcriptome via alternative splicing, differential intron retention, and isoform ratio
switching. *Frontiers in Plant Science* 9:5.

Foyer, C. H., B. Karpinska, and K. Krupinska. 2014. The functions of WHIRLY1 and
REDOXRESPONSIVE TRANSCRIPTION FACTOR 1 in cross tolerance responses in
plants: a hypothesis. *Philosophical Transactions of the Royal Society B* 369: 20130226

Givnish, T. J., D. Spalink, M. Ames, S. P. Lyon, S. J. Hunter, A. Zuluaga, A. Doucette, et al.
2016. Orchid historical biogeography, diversification, Antarctica and the paradox of orchid
dispersal. *Journal of Biogeography* 43: 1905-1916.

Grabherr, M. G., B. J. Haas, M. Yassour, J. Z. Levin, D. A. Thompson, I. Amit, X. Adiconis, et
al. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference
genome. *Nature Biotechnology* 29(7): 644–652.

Grabowski, E., Y. Miao, M. Mulisch, and K. Krupinska. 2008. Single-stranded DNA-binding
protein Whirly1 in barley leaves is located in plastids and the nucleus of the same cell. *Plant*
Physiology 147: 1800-1804.

Graham, S. W., V. K. Y. Lam, and V. S. F. T. Merckx. Plastomes on the edge: the evolutionary
breakdown of mycoheterotroph plastid genomes. *New Phytologist* 214(1):48-55.

Guo, X., D. Fang, S. K. Sahu, S. Yang, X. Guang, R. Folk, S. A. Smith, A. S. Chanderbali, S.
Chen, M. Liu, T. Yang, S. Zhang, X. Liu, X. Xu, P. S. Soltis, D. E. Soltis, and H. Liu. 2021.
Chloranthus genome provides insights into the early diversification of angiosperms. *Nature*
Communications 12:6930.

Haas, B. J., A. Papanicolaou, M. Yassour, M. Grabherr, P. D. Blood, J. Bowden, M. B. Couger,
et al. 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity
platform for reference generation and analysis. *Nature Protocols* 8(8): 1494-1512.

Hoang, D. T., O. Chernomor, A. von Haeseler, B. Q. Minh, and L. S. Vinh. 2018. UFBoot2:
improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* 35:518-
522.

Isemer, R., M. Mulisch, A. Schäfer, S. Kirchner, H.-U. Koop, and K. Krupinksa. 2012.
Recombinant Whirly1 translocates from transplastomic chloroplasts to the nucleus.
Federation of European Biochemical Societies Letters 586:85-88.

Jabre, I., A. S. N. Reddy, M. Kalyna, S. Chaudhary, W. Khokhar, L. J. Byrne, C. M. Wilson, and
N. H. Syed. 2019. Does co-transcriptional regulation of alternative splicing mediate plant
stress responses? *Nucleic Acids Research* 47(6): 2716-2726.

Kalyaanamoorthy, S., B. Q. Minh, T. K. F. Wong, A. von Haeseler, and L. S. Jermiin. 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* 14:
587-589.

Katoh, K., K. Misawa, K. Kuma, and T. Miyata. 2002. MAFFT: a novel method for rapid
multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research*,
30(14): 3059–3066.

Katoh, K. and D. M. Standley. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772-
780.

Klepikova, A. V., A. S. Kasianov, E. S. Gerasimov, M. D. Logacheva, and A. A. Penin. 2016. A
high resolution map of the *Arabidopsis thaliana* developmental transcriptome based on RNA-
seq profiling. *The Plant Journal* 88(6):1058-1070.

Klimpert, N. J., J. L. S. Mayer, D. S. Sarzi, F. Prosdocimi, F. Pinheiro, and S. W. Graham. 2022.
Phylogenomics and plastome evolution of a Brazilian mycoheterotrophic orchid, *Pogoniopsis*
schenckii. *American Journal of Botany* 109(12): 2030-2050.

Krause, K., I. Kilbienski, M. Mulisch, A. Rödiger, A. Schäfer, and K. Krupinksa. 2005.
DNA-binding proteins of the Whirly family in *Arabidiopsis thaliana* are targeted to the
organelles. *Federation of European Biochemical Societies Letters* 579:3707-3712.

Lai, X., H. Chahtane, R. Martin-Arevalillo, C. Zubita, and F. Parcy. 2020. Contrasted
evolutionary trajectories of plant transcription factors. *Current Opinion in Plant Biology*.
54:101-107.

Latchman, D. S. 1993. Transcription factors: an overview. *International Journal of*
Experimental Pathology 74(5): 417–422.

Lallemand, F., M.-L. Martin-Magniette, F. Gilard, B. Gakière, A. Launay-Avon, É. Delannoy,
and M.-A. Selosse. 2019. *In situ* transcriptomic and metabolomic study of the loss of
photosynthesis in the leaves of mixotrophic plants exploiting fungi. *The Plant Journal*
98(5):826-841.

Leake, J. R. 1994. The biology of myco-heterotrophic ("saprophytic") plants. *New Phytologist*
127(2):171-216.

Lee, C., T. Ruhlman, and R. K. Jansen. 2020. Unprecedented intraindividual structural
heteroplasmy in *Eleocharis* (Cyperaceae, Poales) Plastomes. *Genome Biology and Evolution*
12(5): 641-655.

Lehti-Shiu, M. D., N. Panchy, P. Wang, S. Uygun, and S.-H. Shiu. 2017. Diversity, expansion,
and evolutionary novelty of plant DNA-binding transcription factor families. *Biochimica et*
Biophysica Acta 1860(1):3-20.

Lemaire, B., S. Huysmans, E. Smets, and V. S. F. T. Merckx. 2010. Rate accelerations in nuclear
18S rDNA of mycoheterotrophic and parasitic angiosperms. *Journal of Plant Research*
124:561-576.

Lepage, É., É. Zampini, and N. Brisson. 2013. Plastid genome instability leads to reactive
oxygen species production and plastid-to-nucleus retrograde signaling in *Arabidopsis*. *Plant*
Physiology 163:867-881.

Li, H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34(18):
3094-3100.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R.
Durbin, and the 1000 Genome Project Data Processing Subgroup. 2009. The sequence
alignment/map (SAM) format and SAMtools. *Bioinformatics* 25(16): 2078-2079.

Li, Y. -X., Z. -H. Li, A. Schuiteman, M. W. Chase, J. -W. Li, W. -C. Huang, A. Hidayat, et al.
2019. Phylogenomics of Orchidaceae based on plastid and mitochondrial genomes.
Molecular Phylogenetics and Evolution 139(106540).

Lin, W., D. Huang, X. Shi, B. Deng, Y. Ren, W. Lin, and Y. Miao. 2019. H_2O_2 as a feedback
signal on dual-located WHIRLY1 associates with leaf senescence in *Arabidopsis*. *Cells* 8:
1585.

Lin, W., H. Zhang, D. Huang, D. Schenke, D. Cai, B. Wu, and Y. Miao. 2020. Dual-localized
WHIRLY1 affects alicyclic acid biosynthesis via coordination of ISOCHORISMATE
SYNTHASE1, PHENYLALANINE AMMONIA LYASE1, and S-ADENOSYL-L-
METHIONINE-DEPENDENT METHYLTRANSFERASE1. *Plant Physiology* 184: 1884-
1899.

Liu, J. C. Jung, J. Xu, H. Wang, S. Deng, L. Bernad, C. Arenas-Huertero, and N.-H. Chua. 2012.
Genome-wide analysis uncovers regulation of long intergenic noncoding RNAs in
Arabidopsis. *The Plant Cell* 24: 4333-4345.

Lyko, P. and S. Wicke. 2021. Genomic reconfiguration in parasitic plants involves considerable
gene losses alongside global genome size inflation and gene births. *Plant Physiology* 186:
1412-1423.

Maddison, W. P. 1997. Gene trees in species trees. *Systematic Biology* 46(3): 523-536.

Maréchal, A., J. S. Parent, F. Véronneau-Lafortune, A. Joyeux, B. F. Lang, and N. Brisson.
2009. Whirly proteins maintain plastid genome stability in Arabidopsis. *Proceedings of the*
National Academy of Sciences of the United States of America 106(34): 14693– 14698.

Merckx, V. S. F. T. and J. V. Freudenstein. 2010. Evolution of mycoheterotrophy in plants: a
phylogenetic perspective. *New Phytologist* 185(3):605-609.

Merckx, V. S. F. T. 2013. Mycoheterotrophy: an introduction. In: Mycoheterotrophy: the
biology of plants living on fungi. Ed: V. S. F. T. Merckx. 1st ed. Springer: NY. DOI
10.1007/978-1-4614-5209-6.

Mergner, J., M. Fresno, M. List, M. Papacek, X. Chen, A. Chaudhary, P. Samaras, S. Richter, H.
Shikata, M. Messerer, D. Lang, S. Altmann, P. Cyprus, D. P. Zolg, T. Matheson, M.
Bantscheff, R. R. Hazarika, T. Schmidt, C. Dawld, A. Dunkel., T. Hofmann, S. Sprunck, P.
Falter-Braun, F. Johannes, K. F. X. Mayer, G. Jürgens, M. Wilhelm, J. Baumbach, E. Grill,
K. Schneltz, C. Schwechhelmer, and B. Kuster. 2020. Mass-spectrometry-based draft of the
Arabidopsis proteome. *Nature* 579: 409-414.

Münster, T., J. Pahnke, A. Di Rosa, J. T. Kim, W. Martin, H. Saedler, and G. Theissen. 1997.
Floral homeotic genes were recruited from homologous MADS-box genes preexisting in the
common ancestor of ferns and seed plants. *Proceedings of the National Academy of the*
United States of America 94: 3415-2420.

Murrell, B., S. Moola, A. Mabona, T. Weighill, D. Sheward, S. L. K. Pond, and K. Scheffler.
2013. FUBAR: A Fast, Unconstrained Bayesian AppRoximation for inferring selection.
Molecular Biology and Evolution (30): 1196–1205.

Murrell, B., S. Weaver, M. D. Smith, J. O. Wertheim, S. Murrell, A. Aylward, K. Eren, T.
Pollner, D. P Martin, D. M. Smith, K. Scheffler, and S. L. K. Pond. 2015. Gene-wide
identification of episodic selection. *Molecular Biology and Evolution* 32(5): 1365-1371.

Ner-Gaon, H., R. Halachmi, S. Savaldi-Goldstein, E. Rubin, R. Ophir, and R. Fluhr. 2004. Intron
retention is a major phenomenon in alternative splicing in *Arabidopsis*. *The Plant Journal* 39:
877-885.

Nguyen L.-T., H. A. Schmidt, A. von Haeseler, and B. Q. Minh. 2015. IQ-TREE: A fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular*
Biology and Evolution 32(1): 268–274.

Pamilo, P. and M. Nei. 1988. Relationships between gene trees and species trees. *Molecular*
Biology and Evolution 5(5): 568-583.

Parent, J.-S., E. Lepage, and N. Brisson. 2011. Divergent roles for the two polI-like organelle
DNA polymerases of *Arabidopsis*. *Plant Physiology* 156(1): 254-262.

Patro, R., G. Duggal, M. I. Love, R. A. Irizarry, and C. Kingsford. 2017. Salmon provides fast
and bias-aware quantification of transcript expression. *Nature Methods* 14: 417–419.

Patron, N. J. and R. F. Waller. 2007. Transit peptide diversity and divergence: a global analysis of
plastid targeting signals. *BioEssays* 29: 1048-1058.

Pérez-Escobar, O. A., S. Dodsworth, D. Bogarín, S. Bellot, J. A. Balbuena, R. J. Schely, I. A.
Kikuchi, S. K. Morris, N. Epitawalage, R. Cowan, O. Maurin, A. Zuntini, T. Arias, A. Serna-
Sánchez, B. Gravendeel, M. F. Torres Jimenez, K. Nargar, G. Chomicki, M. W. Chase, I. J.
Leitch, F. Forest, and W. J. Baker. 2021. Hundreds of nuclear and plastid loci yield novel
insights into orchid relationships. *American Journal of Botany* 108(7): 1166-1180.

1003

1004   Pond, S. L. K. and S. V. Muse. 2005. HyPhy: hypothesis testing using phylogenies. In:
1005   Statistical Methods in Molecular Evolution. Ed: R. Nielsen. 1st ed. Springer: NY.

1006

1007   Prikryl, J., K. P. Watkins, G. Friso, K. J. van Wijk, and A. Barkan. 2008. A member of the
1008   Whirly family is a multifunctional RNA- and DNA-binding protein that is essential for
1009   chloroplast biogenesis. *Nucleic Acids Research* 36(16): 5152–5165.

1010

1011   Qiu, Z., D. Chen, L. Teng, P. Guan., G. Yu, P. Zhang, J. Song, Q. Zeng, and L. Zhu. 2022.
1012   OsWHY1 interacts with OsTRXz and is essential for early chloroplast development in rice.
1013   *Rice* 15:50.

1014

1015   R Core Team. 2019. R: A language and environment for statistical computing, Vienna, Austria.
1016   Available at: https://www.R-project.org/.

1017

1018   Ranwez, V., S. Harispe, F. Delsuc, and E. J. P. Douzery. 2011. MACSE: multiple alignment of
1019   coding sequences accounting for frameshifts and stop codons. *PLoS ONE* 6(9):e22594.

1020

1021   Rasmussen, H. N. 1995. Terrestrial orchids: from seed to mycotrophic plant. Cambridge:
1022   Cambridge University Press.

1023

1024   Rasmussen, H. N. 2002. Recent developments in the study of orchid mycorrhiza. *Plant and Soil*
1025   244:149-163.

1026

1027   Reddy, A. S., Y. Marquez, M. Kalyna, and A. Barta. 2013. Complexity of the alternative

1028   splicing landscape in plants. *Plant Cell* 25(10): 3657-3683.

1029

1030   Ren, Y., Y. Li, Y. Jiang, B. Wu, and Y. Miao. 2017. Phosphorylation of WHIRLY1 by CIPK14
1031   shifts its localization and dual functions in *Arabidopsis*. *Molecular Plant* 10:749-763.

1032

1033   Robinson, M. D., D. J. McCarthy, and G. K. Smyth. 2010. edgeR: a Bioconductor package for
1034   differential expression analysis of digital gene expression data. *Bioinformatics* 26(1): 139-
1035   140.

1036

1037   Robinson, M. D. and A. Oshlack. 2010. A scaling normalization method for differential
1038   expression analysis of RNA-seq data. *Genome Biology* 11:R25.

1039

1040   Ruan, Q., Y. Wang, H. Xu, B. Wang, X. Zhu, B. Wei, and X. Wei. 2022. Genome-wide
1041   identification, phylogenetic, and expression analysis under abiotic stress conditions of Whirly
1042   (WHY) gene family in *Medicago sativa* L. *Scientific Reports* 12:18676.

1043

1044   Schelkunov, M. I., V. Y. Shtratnikova, M. S. Nuraliev, M.-A. Selosse, A. A. Penin, and M. D.
1045   Logacheva. 2015. Exploring the limits for reduction of plastid genomes: a case study of the
1046   mycoheterotrophic orchids *Epipogium aphyllum* and *Epipogium roseum*. *Genome Biology*
1047   *and Evolution* 7(4):1179-1191.

1048

1049   Serna-Sánchez, M.A., O. A. Pérez-Escobar, D. Bogarín, F. F. Torres-Jimenez, A. C.
1050   Alvarez-Yela, J. E. Arcila-Galvis, C. F. Hall., F. de Barros, F. Pinheiro, S. Dodsworth, M. W.
1051   Chase, A. Antonelli, and T. Arias. 2021. Plastid phylogenomics resolves ambiguous
1052   relationships within the orchid family and provides a solid timeframe for biogeography and
1053   macroevolution. *Scientific Reports* 11:6858

1054

1055   Sinn, B. T. and C. F. Barrett. 2020. Ancient mitochondrial gene transfer between fungi and the
1056   orchids. *Molecular Biology and Evolution* 37(1): 44-57.

1057

1058   Suetsugu, K., M. Yamato, C. Miura, K. Yamaguchi, K. Takahashi, Y. Ida, S. Shigenobu, and I.
1059   Kaminaka. 2017. Comparison of green and albino individuals of the partially
1060   mycoheterotrophic orchid *Epipactis hellebroine* on molecular identities of mycorrhizal fungi,
1061   nutritional modes and gene expression in mycorrhizal roots. *Molecular Ecology* 26:1652-
1062   1669.

1063

1064   Sun, S., S. Li, X. Zhou, and X. Yang. 2023. WRKY1 represses the WHIRLY1 transcription

1065   factor to positively regulate plant defense against geminivirus infection. *PLoS Pathogens*
1066   19(4): e1011319.

1067

1068 Taylor, R. E., C. E. West, and C. H. Foyer. 2022. WHIRLY protein functions in plants. *Food*
1069   *and Energy Security* 00, e379. https://doi.org/10.1002/fes3.379

1070

1071 The Potato Genome Sequencing Consortium. 2011. Genome sequence and analysis of the tuber
1072   crop potato. *Nature* 475: 189-195.

1073

Timilsena, P. R., C. F. Barrett, A. Piñeyro-Nelson, E. K. Wafula, S. Ayyampalayam, J. R.
McNeal, T. Yukawa, T. J. Givnish, S. W. Graham, J. C. Pires, J. I. Davis, C. Ané, D. W.
Stevenson, J. Leebens-Mack, E. Martínez-Salas, E. R. Álvarez-Buylla, and C. W.
dePamphilis. 2023. Phylotranscriptomic analyses of mycoheterotrophic monocots show a
continuum of convergent evolutionary changes in expressed nuclear genes from three
independent nonphotosynthetic lineages. *Genome Biology and Evolution* 15(1):
https://doi.org/10.1093/gbe/evac183

1081

1082 Untergasser, A., I. Cutcutache, T. Koressaar, J. Ye, B. C. Faircloth, M. Remm, and S. G. Rozen.
1083   2012. Primer3-new capabilities and interfaces. *Nucleic Acids Research* 40(15):e115.

1084

1085 Waterworth, W. M., G. E. Drury, C. M. Bray, and C. E. West, C.E. 2011. Repairing breaks in
1086   the plant genome: the importance of keeping it together. *New Phytologist* 192: 805-822.

1087

1088 Wertheim, J. O., B. Murrell, M. D. Smith, S. L. K. Pond, and K. Scheffler. 2015. RELAX:
1089   detecting relaxed selection in a phylogenetic framework. *Molecular Biology and Evolution*
1090   (32): 820–832.

1091

1092 Wicke, S., K. F. Müller, C. W. dePamphilis, D. Quandt, S. Bellot, and G. M. Schneeweiss. 2016.
1093   Mechanistic model of evolutionary rate variation en route to a nonphotosynthetic lifestyle in
1094   plants. *Proceedings of the National Academy of Sciences of the United States of America*
1095   113(32):9045-9050.

1096

1097 Wu, F. L. A. Mueller, D. Crouzillat, V. Pétiard, and S. D. Tanksley. 2006. Combining
1098   bioinformatics and phylogenetics to identify large sets of single-copy orthologous genes
1099   (COSII) for comparative, evolutionary, and systematic studies: a test case in the euasterid
1100   plant clade. *Genetics* 174: 1407-1420.

1101

Xu, Y., Y. Lei, Z. Su, M. Zhao, J. Zhang, G. Shen, L. Wang, J. Li, J. Qi, and J. Wu. 2021. A
    chromosome-scale *Gastrodia elata* genome and large-scale comparative genomic analysis
    indicate convergent evolution by gene loss in mycoheterotrophic and parasitic plants. *The
    Plant Journal* 108: 1609-1623.

Yoo, H. H., C. Kwon, M. M. Lee, and I. K. Chung. 2007. Single-stranded DNA binding factor
    AtWHY1 modulates telomere length homeostasis in *Arabidopsis*. *The Plant Journal* 49:442-
    451.

Yuan, Y., X. Jin, J. Liu, X. Zhao, J. Zhou, X. Wang, D. Wang, C. Lai, W. Xu, J. Huang, L. Zha,
    D. Liu, X. Ma, L. Wang., M. Zhou, Z. Jiang, H. Meng, H. Peng, Y. Liang, R. Li, C. Jiang, Y.
    Zhao, T. Nan, T. Nan, Y. Jin, Z. Zhan, J. Yang, W. Jiang, and L. Huang. 2018. The *Gastrodia
    elata* genome provides insights into plant adaptation to heterotrophy. *Nature Communications*
    9:1615.

Zeng, X., Y. Li, H. Ling, S. Liu, M. Liu, J. Chen, and S. Guo. 2017. Transcriptomic analyses
    reveal clathrin-mediated endocytosis involved in symbiotic seed germination of *Gastrodia
    elata*. *Botanical Studies* 58:31.

Zhang, G.-Q., Q. Xu, C. Bian, W.-C. Tsai, C.-M. Yeh, K.-W. Liu, K. Yoshida, et al. 2016. The
    Dendrobium catenatum Lindl. genome sequence provides insights into polysaccharide
    synthase, floral development and adaptive evolution. *Scientific Reports* 6: 19029.

Zhang, G.-Q., K.-W. Liu, R. Lohaus, Y.-Y. Hsiao, S.-C. Niu, J.-Y. Wang, Y.-C. Lin, Q. Xu, L.-J.
    Chen, K. Yoshida, S. Fujiwara, Z.-W. Wang, Y.-Q. Zhang, N. Mitsuda, M. Wang, G.-H. Liu,
    L. Pecoraro, H.-X. Huang, X.-J. Xiao, M. Lin, X.-Y. Wu, W.-L. Wu, Y.-Y. Chen, S.-B.
    Chang, S. Sakamoto, M. Ohme-Takagi., M. Yagi, S.-J. Zeng, C.-Y. Shen, C.-M. Yeh, Y.-B.
    Luo, W.-C. Tsai, Y. Van de Peer, and Z.J. Liu. 2017. The *Apostasia* genome and the
    evolution of orchids. *Nature* 549:397-383.

Zhuang, K., F. Kong, S. Zhang, C. Meng, M. Yang, Z. Liu, Y. Wang, N. Ma, and Q. Meng. 2019.
    Whirly1 enhances tolerance to chilling stress in tomato via protection of photosystem II and
    regulation of starch degradation. *New Phytologist* 221: 1998-2012.

Zhuang, K., J. Wang, B. Jiao, C. Chen, J. Zhang, N. Ma, and Q. Meng. 2020. WHIRLY1
    maintains leaf photosynthetic capacity in tomato by regulating the expression of *RbcS1* under
    chilling stress. *Journal of Experimental Botany* 12(22): 3653-3663.

1140 Zhao, S. Y., G. D. Wang, W. Y. Zhao, S. Zhang, F. Y. Kong, X. C. Dong, and Q. W. Meng.
1141   2018. Overexpression of tomato WHIRLY protein enhances tolerance to drought stress and
1142   resistance to *Pseudomonas solanacearum* in transgenic tobacco. *Biologia Plantarum* 62(1):
1143   55-68.

1144

1145 Zhao, T., A. Zwaenepoel, J.-Y. Xue, S.-M. Kao, Z. Li, M. E. Schranz, and Y. Van de Peer. 2021.
1146   Whole-genome microsynteny-based phylogeny of angiosperms. *Nature Communications*
1147   12:3498.

1148

1149 Zampini, É., É. Lepage, S. Tremblay-Belzile, S. Truche, and N. Brisson. 2015. Organelle DNA
1150   rearrangement mapping reveals U-turn-like inversions as a major source of genomic
1151   instability in *Arabidopsis* and humans. *Genome Research* 25:645-654.

1152

1153 Zimmer, K., C. Meyer, and G. Gebauer. 2008. The ectomycorrhizal specialist orchid
1154   Corallorhiza trifida is a partial myco-heterotroph. *New Phytologist* 178: 395–400.

1155

11 Data Availability Statement

Reads resulting from RNAseq and Nanopore sequencing are available via NCBI (BioProject
PRJNA984634; released upon acceptance) as components of BAM formatted files where reads are
mapped to the assembled transcript or contig. All sequence alignments analyzed are available in
FASTA format via a public GitHub repository (https://github.com/btsinn/whirly1_evolution.git;
archived via Zenodo upon acceptance).

Tables

Table 1. *In-silico* expression of *WHY1* across species, tissues, and biological replicates. The first row
for a species contains values for gene-level expression, while subsequent rows contain values for a
specific isoform. Trinity isoform and biological replicate identifiers are provided in Table S3.
Expression values are trimmed-mean-of-means-transformed (TMM) transcripts per million (TPM) in
order to maintain comparability amongst biological replicates. An [*] denotes a canonical isoform
while [ψ] denotes an isoform with a retained intron. Values shown have been truncated, TMM to the
tenth, LogFC to the hundredth, and P- and Q-values to the thousandth.

Species Gene or Isoform	Belowground Tissue (Biological Replicate #)			Aboveground Tissue (Biological Replicate #)			LogFC	P-value (Q-value)
C. trifida	1	2	3	1	2	3		
Gene	12.7	11.6	4.7	31.4	71.0	43.7	2.12	0.004 (0.057)
Isoform 1	5.0	1.8	1.0	1.4	2.9	4.3	0.12	0.915 (0.998)
Isoform 2[*]	7.7	9.6	4.1	31.3	56.3	42.0	2.56	0.0001 (0.003)
C. wisteriana	1	2	3	1	2	3		
Gene	14.2	23.0	8.3	84.3	76.7	37.8	1.91	0.019 (0.176)
Isoform 1[ψ]	0	0.8	0	1.0	1	2.0	2.36	0.313 (1)
Isoform 2[*]	14.7	23.0	9.2	74.9	72.6	34.7	1.96	0.021 (0.283)
C. maculata	1	2	NA	1	2	3		
Gene	16.6	12.7		47.0	25.3	30.0	1.17	0.162 (0.624)
Isoform 1[ψ]	2.0	1.8		0.5	0	3.5	-0.51	0.746 (0.970)
Isoform 2[*]	13.9	11.0		45.9	26.1	26.8	1.33	0.119 (0.387)
C. striata	1	2	3	1	2	NA		
Gene	21.3	16.4	7.8	32.1	17.3		0.62	0.525 (0.804)
Isoform 1	0.9	0	2.3	0	0		NA	NA
Isoform 2	16.8	14.5	5.5	30.5	17.2		0.39	0.695 (0.918)
Isoform 3[ψ]	4.4	0.7	0.5	0.8	0		-2.88	0.254 (0.536)

1170

Figure 1. Overview of *Corallorhiza* species included in this study, showing plastid and nuclear
phylogenetic relationships, inflorescence, trophic status (fully vs partially mycoheterotrophic),
plastome size (bp), number of putatively functional plastid genes, and chlorophyll content (mean and
standard deviation in nanograms of total chlorophylls per milligram of plant material). Phylogenetic,
plastome and chlorophyll content data are from Barrett et al. (2014).

Figure 2. Wrapped view of open reading frame-trimmed *Corallorhiza WHY1* isoform alignment.
Disagreements with the consensus sequence are highlighted. An [*] denotes a canonical isoform while
[Ψ] denotes an isoform with a retained intron.

Figure 3. DNA alignment of canonical isoforms and Nanopore-sequenced genomic sequence from
each *Corallorhiza species* with annotated *WHY1* sequence from the *Phalaenopsis equestris* and
Dendrobium catenatum reference genomes. *WHY1* exons 1 and 2 are annotated in yellow and the
locus ID tags for each reference genome are depicted in both the gene (green) and CDS (yellow)
annotations. Disagreements with the consensus sequence are shown in black. Percent identity of
aligned sites is depicted as a histogram. An [*] denotes a canonical isoform while [Ψ] denotes an isoform
with a retained intron.

Figure 4. DNA alignment of *Corallorhiza* isoforms assembled from each *Corallorhiza species* and
Nanopore sequences for *C. trifida*, *maculata*, and *striata* with annotated *WHY1* sequence from the
Phalaenopsis equestris and *Dendrobium catenatum* reference genomes. *WHY1* exons 1 and 2 are
annotated in yellow and the locus ID tags for each reference genome are depicted in both the gene
(green) and CDS (yellow) annotations. Disagreements with the consensus sequence are highlighted.
Percent identity of aligned sites is depicted as a histogram. An [*] denotes a canonical isoform while [Ψ]
denotes an isoform with a retained intron.

Figure 5. Detailed view of amino acid alignment of *WHY1* sequences of orchid species. Note the
phenylalanine substitutions unique to *Corallorhiza wisteriana*, *maculata*, and *striata*, relative to the
other species sampled. The amino acid substitutions of interest are located at the 3' portion of an
region inferred to conform into an alpha helix structure (pink annotation) in *Arabidopsis thaliana*
WHY1 (alignment row two).

Figure 6. Maximum likelihood inferred topology of *WHY1* evolution with support values below 95
derived from 1,000 non-parametric bootstrap replicates, shown. Lineages highlighted in red represent
those for which statistically-significant signal of episodic diversifying selection of *WHY1* was
detected by the adaptive branch-site random effects likelihood test.

Corallorhiza striata
Fully mycoheterotrophic
Plastome size = 137,505 bp
82 functional plastid genes
Chlorophyll content = 1.6 ± 1.7 ng/mg

Corallorhiza trifida
Partially mycoheterotrophic
Plastome size = 149,384 bp
107 functional plastid genes
Chlorophyll content = 31.6 ± 14.0 ng/mg

Corallorhiza wisteriana
Partially mycoheterotrophic
Plastome size = 146,437 bp
103 functional plastid genes
Chlorophyll content = 21.5 ± 14.2 ng/mg

Corallorhiza maculata
Fully mycoheterotrophic
Plastome size = 146,595 bp
88 functional plastid genes
Chlorophyll content = 0.9 ± 1.2 ng/mg

Consensus													
Identity													

Dendrobium catenatum

Phalaenopsis equestris

C. trifida isoform 1
C. trifida isoform 2*
C. trifida 2227 genomic
C. wisteriana isoform 2*
C. wisteriana isoform 1ᵠ
C. maculata isoform 2*
C. maculata isoform 1ᵠ
C. maculata 2123 genomic
C. striata isoform 1
C. striata isoform 2
C. striata isoform 3ᵠ
C. striata 2084 genomic

Multiple sequence alignment showing conservation across Consensus, Identity, *Amborella trichopoda*, *Arabidopsis thaliana*, and orchid taxa including *Corallorhiza trifida* isoform 2, *Corallorhiza wisteriana* isoform 2, *Corallorhiza maculata* isoform 2, *Corallorhiza striata* isoform 2, *Gastrodia elata*, *Epipogium aphyllum*, *Apostasia shenzhenica*, *Apostasia wallichii*, *Neuwiedia zollingeri*, *Vanilla planifolia*, *Dendrobium catenatum*, *Cypripedium formosanum*, *Cymbidium ensifolium*, *Oncidium gower*, *Orchis italica*, *Ophrys sphegodes*, *Phalaenopsis bellina*, *Phalaenopsis modesta*, *Phalaenopsis aphrodite*, *Phalaenopsis schilleriana*, *Phalaenopsis lueddemanniana*, *Phalaenopsis equestris*.