Pesticide-induced resurgence in brown planthopper is mediated by action on a suite of genes that promote juvenile hormone biosynthesis and female fecundity

Yang Gao[a,1], Shao-Cong Su[a,1], Ji-Yang Xing[a], Zhao-Yu Liu[a], Dick R. Nässel[b], Chris Bass[c], Cong-Fen Gao[a], Shun-Fan Wu[a,*]

[a] College of Plant Protection, Nanjing Agricultural University, Nanjing 210095, China.

[b] Department of Zoology, Stockholm University, Stockholm, Sweden

[c] College of Life and Environmental Sciences, Biosciences, University of Exeter, Penryn Campus, Penryn, UK

[1] These authors contributed equally to this work.

*Corresponding author. E-mail address: wusf@njau.edu.cn (S.-F. Wu)

Abstract

Pesticide-induced resurgence, increases in pest insect populations following pesticide application, is a serious threat to the sustainable control of many highly damaging crop pests. Resurgence can result from pesticide-enhanced pest reproduction, however, the molecular mechanisms mediating this process remain unresolved. Here we show that brown planthopper (BPH) resurgence in rice crops following exposure to the pesticide emamectin benzoate (EB) results from the coordinated action of a suite of genes that regulate juvenile hormone (JH) levels, resulting in increased JH titer in adult females and enhanced fecundity. We demonstrate that EB treatment at sublethal levels results in profound changes in female BPH fitness including increased egg maturation and oviposition. This enhanced reproductive fitness results from the EB-mediated upregulation of key genes involved in the regulation of JH, including *JHAMT and Kr-h1* and the downregulation of allatostatin (*AstA*) and allatostatin receptor (*AstAR*) expression. AstA signaling is known to inhibit the production of JH in the corpora allata and hence EB exposure diminishes this inhibitory action. We find that the changes in gene expression following EB exposure are caused by the action of this insecticide on its molecular target, the glutamate-gated chloride channel (GluCl). Collectively, these results provide mechanistic insights into the regulation of negative pesticide-induced responses in insects and reveal the key actors involved in the JH-signaling pathway that underpin pesticide resurgence.

Keywords: resurgence; reproduction; pesticide; allatostatin; brown planthopper

Significance

Pesticides remain a key means of controlling many of the world's insect pests. However, in some cases, pesticide applications can result in resurgence of pest populations due to pesticide-induced increases in fecundity. In the current study, we show that pesticide resurgence in the brown planthopper (BPH) following exposure to the insecticide emamectin benzoate (EB) results from the transcriptional reprogramming of a diverse suite of positive and negative regulators of juvenile hormone (JH), a critical regulator of insect development and reproduction. This in turn leads to profound increases in female BPH reproductive fitness and enhanced fecundity. Our findings unravel the molecular mechanisms mediating pesticide-induced pest resurgence and inform the development of novel strategies to control highly damaging crop pests.

Introduction

Chemical pesticides remain the primary means of controlling many of the world's most damaging arthropod crop pests (Janssen and van Rijn, 2021; Wu et al., 2018a). However, pesticide applications can result in pest resurgence, increases in pest insect populations that exceed natural, untreated population sizes, following an initial reduction of the pest population (Gandara et al., 2024; Guedes et al., 2016; Hardin et al., 1995; Wu et al., 2020). Two mechanisms have been implicated in pest resurgence - the loss of beneficial insects including natural enemies, and pesticide-enhanced reproduction in the pest insect (Wu et al., 2020). In the case of the latter, several pesticides, such as the insecticides triazophos, deltamethrin and the fungicide jinggangmycin, have been reported to stimulate pest reproduction (Ge et al., 2010; Wu et al., 2018b; Zhang et al., 2014). Pesticide-enhanced pest reproduction has been linked to changes in physiology and biochemistry of pest organisms after exposure to pesticides (Guedes et al., 2016; Wu et al., 2020). However, the molecular mechanisms underlying enhanced reproduction associated with pest resurgence remain poorly resolved.

The brown planthopper (BPH), *Nilaparvata lugens* (Stål), is a notorious pest of rice crops throughout Asia causing annual losses of ~300 million dollars across major rice producing countries (Wu et al., 2020; Wu et al., 2018a). BPHs inhibit the growth of rice plants by feeding, and also transmits highly damaging plant viruses including rice grassy stunt virus and rice ragged stunt virus (Sōgawa, 1982). Currently, chemical insecticides play an indispensable role in the control of BPH due to their efficiency,

rapid effect, and low cost. However, due to the widespread and intensive use of

chemical insecticides, BPH has developed resistance to the majority of compounds

used for control (Wu et al., 2018a; Zeng et al., 2023).

Emamectin benzoate (EB) and abamectin are avermectin pesticides, and act as

allosteric modulators of insect glutamate gated chloride channels (GluCls), inhibiting

muscle contractions that lead to the cessation of insect feeding and subsequent death

(Ishaaya et al., 2002). These insecticides exhibit particularly strong activity against

Lepidoptera such as the rice leaffolder, *Cnaphalocrocis medinalis* Guénee, an

important foliage-feeding insect which attacks rice during the vegetative stage

(Chintalapati et al., 2016). Both BPH and the rice leaffolder are migratory pests with

overlapping migratory paths, however, their occurrence period in the field differs by

approximately one month, with leaffolders appearing earlier than BPH. Therefore, the

use of EB to control rice leaffolder has the potential to impact BPHs arriving later, via

exposure to sublethal concentrations of this compound. In this regard, we have

observed that when farmers use EB and abamectin to control leaffolders on rice crops

in China, BPH outbreaks frequently occur in the same field. While sublethal doses of

certain pesticides have been shown to enhance fecundity in BPHs, including the

insecticides triazophos and deltamethrin (Ge et al., 2013; Ge et al., 2010; Zhang et al., 2014;

Zhang et al., 2022b) and the fungicides carbendazim and jinggangmycin (Wu et al.,

2018b), whether avermectins trigger resurgence in BPH via insecticide-enhanced

reproduction remains unclear.

Reproduction in insects is influenced by external factors such as light (Wang et al.,

2021), temperature (Meiselman et al., 2022), humidity (Roy et al., 2015) and nutrition

(Smykal and Raikhel, 2015), and endogenous factors such as the juvenile hormone (JH)

(Santos et al., 2019), ecdysone (Hun et al., 2022), insulin (Ling and Raikhel, 2018) and target

of rapamycin (TOR) pathways (Ahmed et al., 2020; Du et al., 2022; Lu et al., 2016). Of

these, JH has been particularly implicated in insecticide-induced enhanced fecundity,

with triazophos and deltamethrin treatments leading to increased circulating JH III

titers in BPH females (Wu et al., 2020). JH is synthesized and secreted by the corpora

allata in insects (Luo et al., 2021), and can promote reproduction by regulating the

synthesis and secretion of vitellogenin (Vg) in the female fat body, and stimulating the

absorption of Vg by the developing oocyte (Santos et al., 2019). However, the regulation

of JH is complex (Santos et al., 2019) and the key actors involved in JH-mediated

pesticide-enhanced reproduction remains an open question. Previous finding reported

that the GluCl receptor is involved in JH biosynthesis in cockroach (Chiang et al., 2002a;

Chiang et al., 2002b; Liu et al., 2005). However, it is still unknown whether insecticides that

target the GluCl can increase reproduction in pest insects and how the GluCl receptor

regulates JH production.

In this study, we used a range of diverse approaches to investigate the impact of

sublethal doses of avermectins on BPH fecundity, and unravel the molecular

mechanisms mediating enhanced reproduction following exposure to this insecticide

class. We show that avermectin exposure results in profound changes in the

expression of a key suite of genes that in combination regulate JH, resulting in

increased JH titer in adult females, which promotes fecundity. Interestingly, we found

that GluCl channel modulators can inhibit AstA/AstAR expression and thereby

promote JH production.

RESULTS

GluCl allosteric modulators (emamectin benzoate and abamectin) stimulate

fecundity of female *N. lugens*

To investigate whether GluCl modulators affect fecundity in BPH, we first

determined the sub-lethal and median lethal doses of emamectin benzoate (EB) on

4^{th} instar nymphs, as well as on newly emerged males and females of BPH (Table S1).

For this we employed two different bioassay methods, the rice seedling dip bioassay

method and topical application bioassay method (Wu et al., 2018a; Yao et al., 2017), in

order to assess both the systemic and contact toxicity of these insecticides (Table S1).

We then systemically treated 4^{th} instar nymphs of BPH, newly emerged males and

females with the estimated LC_{15} or LC_{50} concentrations of EB and examined the

fecundity of BPH after these individuals mated with treated or untreated individuals.

We use the term "t" to represent individuals treated with EB and "ck" to indicate control

individuals that were treated with insecticide diluent minus insecticide. First, we tested

whether treatment of BPH with EB in nymphal stage impairs the fecundity of BPHs at

the adult stage. For 4^{th} instar nymphs of BPHs, after treatment with the LC_{15} and LC_{50}

concentrations of EB, the number of eggs laid per female of BPHs in ♀t × ♂t crosses

increased by 1.48 and 1.40 times compared with control ♀ck × ♂ck crosses (Figure

1A); the number of eggs laid per female of BPH in ♀t × ♂ck crosses increased by 1.53

and 2.07 times compared with control crosses (Figure 1B); However, the number of

eggs laid by per female of BPH in ♀ck × ♂t crosses did not increase significantly

compared to control ♀ck × ♂ck crosses (Figure 1C).

Table S1. Determination of the toxicity of emamectin benzoate on BPH in systemic and topical application bioassays.

Treatment method	Developmental stages	Slope ± SE	LC_{15} (95%F.L.) (mg/L) or LD_{15} (95%F.L.) (pg/insect)	LC_{50} (95%F.L.) (mg/L) or LD_{50} (95%F.L.) (pg/insect)	χ^2(df)	P value
Systemic route	4[th] nymph	2.56 ± 0.487	0.87 (0.39-1.32)	2.21 (1.53-2.82)	2.68 (4)	0.61
	Adult Male	2.42 ± 0.35	0.80 (0.40-1.20)	2.13 (1.48-2.78)	2.12 (4)	0.72
	Adult Female	2.47 ± 0.41	2.33 (1.09-3.56)	6.12 (4.18-8.00)	3.56 (4)	0.50
Contact route	4[th] nymph	2.52 ± 0.35	15.54 (9.46-21.42)	39.90 (30.66-49.56)	2.60 (4)	0.63
	Adult Male	2.77 ± 0.57	6.72 (2.52-10.92)	15.96 (9.66-21.84)	1.23 (3)	0.75
	Adult Female	3.31 ± 0.76	18.48 (6.72-28.14)	37.80 (23.10-50.40)	0.79 (3)	0.85

Exposure of 4th instar nymphs to the LC_{15} and LC_{50} concentrations of EB in

contact bioassays also significantly stimulated fecundity. After treatment with the LC_{15}

and LC_{50} concentrations of EB, the number of eggs laid by per female of BPH in ♀t ×

♂t crosses increased by 1.18 and 1.26 times compared with the control (♀ck × ♂ck)

(Figure 1D); The number of eggs laid per female of BPH in ♀t × ♂ck crosses

increased by 1.27 and 1.56 times compared with the control crosses (Figure 1E);

However, there was no significant difference in number of eggs laid between ♀ck × ♂t

crosses and controls (♀ck × ♂ck) (Figure 1F). These results reveal that EB stimulates

the fecundity of females following both the systemic and contact routes of exposure.

Figure 1

Figure 1. Fecundity of BPH following exposure to sub-lethal (LC_{15}) and median lethal (LC_{50})

concentrations of emamectin benzoate following system application bioassays (A: ♀t ×♂t; B:

♀t ×♂ck; C: ♀ck ×♂t) and topical application bioassays (D: ♀t ×♂t; E: ♀t ×♂ck; F: ♀ck ×♂t),

respectively. The letter "t" represents treatment with insecticide, while "ck" indicates controls

that was not treated with insecticide. These were exposed as 4^{th} instar nymphs. All data are

presented as the mean ± s.e.m. Different lower-case letters above the bars indicate significant

differences (One-way ANOVA with Tukey's Multiple Range Test, $p < 0.05$).

Next, we examined whether EB treatment of adult BPHs also stimulates

reproduction. Indeed, treating newly emerged adults with the LD_{15} and LD_{50}

concentrations of EB significantly stimulated the number of eggs laid per female

(Figure 1-figure supplement 1A). Furthermore, sub-lethal exposure of 4^{th} instar BPH

nymphs to another GluCl allosteric modulator, abamectin (LC_{15} and LC_{50}

concentrations) was also found to significantly enhance reproduction (Figure 1-figure

supplement 1B).

Figure 1-figure supplement 1

Figure 1-figure supplement 1. (A) Fecundity of BPH when newly emerged adults were

treated with sub-lethal (LD_{15}) and median lethal (LD_{50}) concentrations of emamectin benzoate

via topical application. (B) Fecundity of BPH when 4^{th} instar nymphs were treated with

sub-lethal (LC_{15}) and median lethal (LC_{50}) concentrations of abamectin via systemic exposure.

All data are presented as the mean ± s.e.m. Different lower-case letters above the bars

indicate significant differences (One-way ANOVA with Tukey's Multiple Range Test, $p < 0.05$).

To examine if EB also stimulates egg-laying in other insect species we conducted

bioassays on the small brown planthopper, *Laodelphax striatellus*, the white backed

planthopper, *Sogatella furcifera* and fruit flies *Drosophila melanogaster*. In contrast to

our findings in BPHs, we found that sub-lethal doses (LC_{15} and LC_{50}) of EB inhibits

fecundity of female *L. striatellus*, (Figure1-figure supplement 2A-C) and has no impact

on the fecundity of *S. furcifera*, (Figure1-figure supplement 2D-F). In addition, we

found that sublethal doses (LC_{15}, LC_{30} or LC_{50}) of EB also inhibit fecundity in *D.*

melanogaster (Figure1-figure supplement 2G and H). These results indicate that the

stimulation of reproduction by EB in BPH is species-specific and does not even

extend to related insect species.

Figure 1-figure supplement 2

Figure 1-figure supplement 2. Fecundity of small brown planthopper, *Laodelphax striatellus*,

(A-C) white backed planthopper, *Sogatella furcifera* (D-F) and fruit fly, *Drosophila*

melanogaster (G and H) when larvae and newly emerged adults were treated with sub-lethal

concentrations of emamectin benzoate. All data are presented as the mean ± s.e.m. Different

lower-case letters above the bars indicate significant differences (One-way ANOVA with

Tukey's Multiple Range Test, $p < 0.05$).

The impact of EB treatment on BPH reproductive fitness

To better understand the effects of EB on the reproductive fitness of BPH, the

duration of the preoviposition period, emergence rate, female ratio, female longevity

and female weight were evaluated following exposure using the systemic bioassay.

The preoviposition period of females treated with the LC_{50} of EB decreased

significantly compared with the control (Figure 1-figure supplement 3A). In contrast no

significant effects of EB on emergence rate and female ratio were observed (Figure

1-figure supplement 3B and C). Female survival analysis showed that exposure of 4th

instar nymphs to the LC_{50} of EB has no impact on female longevity (Figure 1-figure

supplement 3D). Interestingly, occurrence of brachypterism (short-wing phenotype) in

females and female weight were significantly increased after EB exposure (Figure

1-figure supplement 3E and F). It was reported that short-winged females usually

display higher fecundity than long-winged females in BPHs and two different insulin

receptors determine the alternative wing morphs (Xu et al., 2015). However, whether

insulin signaling is involved in the fecundity increase of BPHs in our experiments

needs further investigation.

Figure 1-figure supplement 3

Figure 1-figure supplement 3. The impact of emamectin benzoate on the reproductive

fitness of BPH. Fourth instar nymphs were treated with the LC_{50} concentration of emamectin

benzoate in systemic bioassays. (A) Preoviposition period: Preoviposition refers to the period

in an insect's life cycle between the time it becomes an adult and the time it starts laying eggs.

(B) Emergence rate: the rate of emergence of adults; (C) Female ratio: the ratio of female (to

male) insects in the total of emerged adults; (D) Female adult longevity. (E) Brachypterism

female ratio: the ratio of short-winged to long-winged adults; (F) The weight per female. All

data are presented as the mean ± s.e.m. Different lower-case letters above the bars indicate

significant differences (Student's *t* test, $p < 0.05$).

EB promotes ovarian maturation in BPH

To investigate the cause of increased egg-laying following EB exposure we

examined if EB influences ovarian maturation in BPH. We dissected and compared

the ovaries of females treated with the LC$_{50}$ of EB at 1, 3, 5 and 7 days after eclosion

(DAE) with control females. At 3, 5 and 7 DAE, the number of eggs in the ovary of

BPH in the EB treated group were significantly higher than that of controls (Figure 2A

and B). We also observed that EB treatment of female adults also increases the

number of mature eggs in the ovary (Figure 2-figure supplement 1). Hence, we found

that EB does not affect normal egg developmental stages. However, our results

suggest that EB treatment can promote oogenesis and as a result the insects both

produce more eggs in the ovary and more eggs were laid.

Unlike the model insect *Drosophila melanogaster*, studies on oogenesis in the

BPH are relatively limited. Therefore, we used immunohistochemistry to stain eggs at

different stages of BPH development. We captured images with a laser confocal

microscope, to document each stage of oogenesis in this species. We found that

oogenesis in the BPH differs from that in *D. melanogaster* and belongs to the

telotrophic meroistic type, where the oocyte is connected to nutrient-providing cells in

the anterior trophic region of the ovariole through a nutritive cord. Oogenesis in the

BPHs occurs in three main compartments (Figure 2-figure supplement 2):

1. Terminal Filament (TF): This region, composed of terminal filament cells, is

similar to the terminal filament structure found in Drosophila.

2. Germarium: This region is where germline stem cells (GSCs) differentiate into

oocytes and follicle cells. It typically consists of cap cells adjacent to the terminal

filament, germline stem cells, escort cells, and cystoblasts.

3. Egg Chamber: The egg chamber in the BPH primarily contains eggs at six

developmental stages. The distinct layering of follicle cells, visible through

cytoskeletal staining, is the main feature distinguishing the six different developmental

stages listed in the figure legend (see also Figure 2-figure supplement 2).

We next explored whether EB treatment could enhance or impair oogenesis in

BPH. However, dissection of various developmental stages revealed that emamectin

benzoate treatment has no significant negative effects on developmental stages and

morphology of oogenesis in BPH (Figure 2C and D). Considering the number of eggs

laid by EB treated females was larger than in control females (Figure 1 and Figure

1-figure supplement 1), our data indicated that EB treatment of BPHs can both

promote oogenesis and fecundity (egg-laying) in BPHs.

Figure 2

Figure 2. The impact of emamectin benzoate on ovarian maturation in BPH. Fourth instar

nymphs were treated with the LC_{50} concentration of emamectin benzoate in systemic

bioassays. (A) Ovarian development in EB treated BPH at 1, 3, 5 and 7 days after eclosion

(DAE) compared to untreated controls. Scale bar = 1,000 μm. (B) Number of mature eggs in

the ovaries of EB treated 4^{th} instar BPH nymphs measured at 1, 3, 5 and 7 DAE compared to

controls. All data are presented as the mean ± s.e.m. Asterisks indicate values significantly

different from the control using student t test (ns, no significant; *$p < 0.05$ and **$p < 0.01$). (C)

Different developmental stages of BPH eggs. (D) No impairment of emamectin benzoate on

oogenesis of BPH. Scale bar = 100 μm.

Figure 2-figure supplement 1

Figure 2-figure supplement 1. Number of mature eggs in the ovaries of EB treated BPH

female adults compared to controls. The data are presented as the mean ± s.e.m. Asterisks

indicate values significantly different from the control using student t test (*$p < 0.05$).

Figure 2-figure supplement 2

Figure 2-figure supplement 2. The oogenesis of BPHs. See the main text for detailed

information.

Stage 1: Connected to the germarium, the egg is trapezoidal. Stage 2: The egg is larger

than in Stage 1 and is square to rectangular. Stage 3: The egg begins to elongate toward both

poles, taking on an oval shape. Stage 4: The egg continues to enlarge and elongate,

becoming spindle-shaped. Due to rapid growth, follicle cells on the egg surface start dividing,

indicating nuclear division. Stage 5: The egg further enlarges, and most follicle cells display

binucleation, the most notable feature of this stage. Small amounts of lipids start entering the

egg. Stage 5 is the final stage before maturation, after which the egg enters the pedicel, a

structure primarily formed by muscle fibers. The expanded mature egg stretches the muscle

fibers in the pedicel, increasing the spacing between them. Stage 6: This is the mature egg

stage, which occurs after the egg passes through the pedicel. During this stage, the egg

undergoes rapid growth, continuously absorbing vitellogenin and lipids (Nile red staining

confirmed that Stage 6 has the highest lipid content). This absorption causes the egg's surface

color to transition from the transparency seen in Stage 5 to mature eggs' dark, opaque

appearance. Once the egg enters the pedicel and becomes completely opaque, it reaches full

maturity. The mature egg then moves into the lateral oviduct for oviposition, completing the

egg-laying process.

EB exposure increases circulating sugars and the storage of macromolecules

in BPHs

Nutritional status is an important indicator of reproductive fitness. Thus, to

investigate whether EB affects intermediary metabolism and energy storage of BPHs,

glycogen, triacylglyceride (TAG), total protein content, cholesterol and four circulating

carbohydrates were quantified in 4^{th} instar BPH nymphs following exposure to the

LC_{50} of EB.

We found that EB exposure has no impact on glycogen levels (Figure 2-figure

supplement 2A). The amount of TAG in EB-treated BPHs was 27% higher ($p < 0.05$)

than those in controls, but only in BPH of the late fifth instar (5L) stage, with no

significant differences observed in subsequent developmental stages (Figure 2-figure

supplement 2B). The amount of total protein content in EB-treated BPH was higher

than the control groups in the case of all developmental stages from 5L nymph to

7DAE (Figure 2-figure supplement 2C). EB exposure also increased cholesterol levels

at 4 and 5 DAE (Figure 2-figure supplement 2D). Compared with the solvent control,

EB treatment caused significant increases ($p < 0.05$) in the levels of sucrose, glucose,

fructose, and trehalose (Figure 2-figure supplement 2E-H). Thus, collectively, these

data provide evidence that EB exposure leads to energy mobilization and the

metabolism of carbohydrates and lipids in BPHs.

Figure 2-figure supplement 2

Figure 2-figure supplement 2. Amounts of Glycogen (A), TAG (B), total protein content (C),

cholesterol (D) and four circulating sugars including sucrose, glucose, fructose and trehalose

(E-H) after BPH exposure to EB. All data are presented as the mean ± s.e.m. The differences

between the EB-treated and solvent-treated BPH were analyzed using unpaired student t-test

(*, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$).

EB stimulates egg-laying that is mediated by the JH signaling pathway

Given the important role of juvenile hormone (JH) in vitellogenesis and egg

development in insects (Jing et al., 2021; Ling and Raikhel, 2021; Luo et al., 2021; Santos et al.,

2019; Zheng et al., 2022), we asked whether EB-treatment could influence the titer of JH

in BPH. As measured by ELISA, the juvenile hormone titer of BPH nymphs treated

with the LC_{50} concentration of EB was significantly lower than that of controls in

systemic bioassays during the middle and late stages of the 4[th] instar (Figure 3A).

However, at 2, 3 and 4 DAE, the JH titer in the EB treated group was significantly

higher than that of the control (Figure 3A). Interestingly, the titer of another important

insect hormone, the steroid ecdysone, was not significantly different between

EB-treated BPH and solvent-treated controls (Figure 3-figure supplement 1). To

independently validate the results of ELISA assays, we employed HPLC-MS/MS to

measure JH titer in BPH following EB exposure (Bownes and Rembold, 1987; Guo et al.,

2020; Jing et al., 2021). The results showed that the JH III titer significantly decreased

after EB-treatment at the late 4[th] instar nymph stage (Figure 3A and B), but

significantly increased at the third day after eclosion (3 DAE) (Figure 3A and C). We

also investigated the effects of EB treatment on the JH titer of female adults. The data

indicates that the JH titer was also significantly increased in the EB-treated female

adults compared with controls (Figure 3-figure supplement 3A). However, the steroid

ecdysone, was also not significantly different between EB-treated BPH and controls

(Figure 3-figure supplement 3B).

To further investigate the role of JH in EB-enhanced fecundity in BPHs, we treated

BPHs with methoprene and pyriproxyfen, JH analogues or biologically active JH

mimic respectively, to determine whether they can stimulate fecundity in BPH. Both

compounds significantly increased egg-laying in BPHs (Figure 3D). Taken together

these results reveal that EB stimulates an increase in JH titer and that JH signaling

enhances fecundity in BPHs.

Since we found that EB could induce JH synthesis in the BPH, we asked whether

EB could influence the expression of genes that are involved in JH biosynthesis or

degradation. For this we treated 4[th] instar nymphs or newly emerged female adults of

BPHs with the LC_{50} concentration of EB using systemic bioassay, and then collected

early (5E), middle (5M) and late (5L) stage of 5[th] instar nymph and 1-7 DAE female

adults for analysis. Quantitative PCR was then used to examine the expression of key

genes previously implicated in the regulation of JH (Gospocic et al., 2021; Nouzova et al.,

2018).

Farnesoic acid O-methyltransferase (FAMeT) (Liu et al., 2010), is known as an

important enzyme in the JH biosynthetic pathway, catalyzing methylation of farnesoic

acid (FA) to methyl farnesoate (MF) (Nouzova et al., 2018). We found that this gene was

significantly upregulated in 5M instar nymph to 4 DAE after EB- treatment (1.0-fold to

3.0-fold) (Figure 3E). Juvenile hormone acid methyltransferase (JHAMT) (Nouzova et

al., 2018; Shinoda and Itoyama, 2003), which is involved in the biosynthesis of JH, was

also upregulated in EB-treated BPH at 1 to 5 DAE compared to controls (1.5-fold to

3.0-fold) (Figure 3F). We furthermore found that EB treatment in female adults

stimulates JHAMT expression (Figure 3-figure supplement 3C).

Methoprene-tolerant (Met), belongs to the basic helix–loop–helix Per/Arnt/Sim

(bHLH-PAS) family of transcription factors and is the intracellular (nuclear) receptor of

JH (Riddiford, 2020; Zhu et al., 2019). The levels of *met* mRNA increased in EB-treated

BPH at the 5M and 5L instar nymph and 1 to 5 DAE adult stages compared to controls

(1.7-fold to 2.9-fold) (Figure 3G).

Krüppel homolog 1 (*Kr-h1*), a transcriptional target of JH signaling, is reported to

be sensitive to levels of JH and its expression levels are directly correlated with JH

titers (Gospocic et al., 2021; Jindra et al., 2013; Zhang et al., 2022a). We found that *Kr-h1* was

significantly upregulated in the adults of EB-treated BPH at the 5M, 5L nymph and 4 to

5 DAE stages (4.7-fold to 27.2-fold) when 4th instar nymph and female adults were

treated with EB (Figure 3H and Figure 3-figure supplement 3D).

Similarly, the expression levels of vitellogenin (Vg), a key downstream component

of JH signaling triggering ovarian development in insects including BPH (Shen et al.,

2019), was markedly increased in females at 2–5 DAE by EB (1.7-fold to 5.5-fold)

(Figure 3I). During the maturation process, eggs require uptake of vg, and an increase

in vg content can accelerate egg maturation, producing more mature eggs. Our

molecular data suggest that EB treatment leads to an upregulation of *vg* expression.

Based on these findings, we conclude that the increase in egg-laying caused by EB

treatment is due to the upregulation of *vg* (Figure 3I), which raises Vg content,

promoting the uptake of Vg by maturing eggs and resulting in the production of more

mature eggs.

Juvenile hormone esterase (JHE) is the primary JH-specific degradation enzyme

that plays a key role in regulating JH titers (Kamita et al., 2003). Interestingly, we

observed a significant upregulation of JHE mRNA levels in the early and middle 5th

instar nymph stage followed by downregulation in 5L instar nymphs of EB-treated

BPH (1.3-fold to 2.6-fold) (Figure 3J).

In combination these results reveal that EB has a profound impact on the

expression of key genes involved in the biosynthesis of JH or in downstream signaling

pathway genes that might promote egg development and female fecundity.

To further understand whether these JH pathway-related genes were involved in

egg-laying behavior in BPH, we performed RNAi experiments to downregulate the

expression of *Kr-h1* (Figure 3-figure supplement 2A). We found that silencing of *Kr-h1*

diminished *vg* gene expression (Figure 3-figure supplement 2B). Importantly,

diminishment of *JHAMT*, *Met* and *Kr-h1* genes in female BPHs was also found to

suppress egg-laying (Figure 3K-M). However, this phenotype was rescued by EB

treatment, which restored egg-laying to normal levels in BPH injected with *JHAMT*,

Met and *Kr-h1* dsRNA (Figure 3K-M). Note that this rescue is possible since

knockdown of the genes is incomplete when using dsRNA injection (and residual

gene expression allows for EB action). Together these results provide a mechanistic

understanding of how EB enhances fecundity in BPH by modulating the expression of

key genes involved in JH synthesis (Figure 3N).

Figure 3. EB induced reproduction in BPHs is mediated by components of the JH

signaling pathway. (A) The titer of JH III (as measured by ELISA assay) at different

developmental stages of BPH when 4[th] instar nymphs were treated with the median lethal

concentrations of EB. (B and C) The titer of JH III (as measured by HPLC-MS/MS) in BPH

females at 4L and 3 DAE when treated with median lethal concentrations of EB. (D)

Oviposition rate of BPH when 4th instar nymphs were treated with 4 ppm methoprene or 10

ppm pyriproxyfen. (E-J) Expression of selected JH-related genes (*FAMeT*, *JHAMT*, *Met*, *Kr-h1*,

Vg, and *JHE*) in EB-treated BPH. (K) Egg production following silencing of *JHAMT* with or

without EB application. (L) Egg production following silencing of *met* with or without EB

application. (M) Egg production after silencing *Kr-h1* with or without EB application. (N)

Schematic illustrating the proposed impact of EB on the JH signaling pathway leading to

enhanced reproduction. The question mark indicates one or more potential additional signals.

All data are presented as means ± s.e.m. Student's *t* test was used to compare the two

samples. One-way ANOVA with Tukey's multiple comparisons test was used to compare more

than two samples. ns, no significant difference; Asterisks indicate values significantly different

from the control (ns, no significant; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ and ****$p < 0.0001$).

Different lower-case letters above the bars indicate significant differences ($p < 0.05$).

Figure 3-figure supplement 1

Figure 3-figure supplement 1. Ecdysone titer at different developmental stages of BPH when

4th instar nymphs were treated with median lethal concentrations of EB.

Figure 3-figure supplement 2

Figure 3-figure supplement 2. (A) Expression of *Kr-h1* following RNAi knockdown. (B)

Expression of *Vg* when *Kr-h1* was silenced in BPH. All data are presented as means ± s.e.m.

*p < 0.05; Mann–Whitney test.

Figure 3-figure supplement 3

Figure 3-figure supplement 3. (A) The titer of JH III (as measured by HPLC-MS/MS) in BPH

females at adult stages when treated with median lethal concentrations of EB. (B) The titer of

ecdysone titer of BPH when female adults were treated with median lethal concentrations of

EB. (C and D) Expression of *JHAMT* and *Kr-h1* in EB-treated BPH female adults. All data are

presented as means ± s.e.m. Student's *t* test was used to compare the two samples. Asterisks

indicate values significantly different from the control (ns, no significant; *$p < 0.05$).

EB induces JH biosynthesis through the peptidergic AstA/AstAR signaling pathway

The timing and level of JH biosynthesis can be precisely regulated by neuropeptides, stimulatory allatotropins (ATs) and inhibitory allatostatins (Asts), in many insects (Bellés et al., 1999; Kataoka et al., 1989; Kramer et al., 1991; Lorenz et al., 1995; Stay and Tobe, 2007; Verlinden et al., 2015; Wegener and Chen, 2022; Woodhead et al., 1989). Insects can, in a species-specific manner, produce one type of AT and three types of Asts: FGL-amide Ast (AstA) (Woodhead et al., 1989; Yin et al., 2006); myoinhibitory peptide (MIP or AstB) (Stay and Tobe, 2007) and PISCF Ast (AstC) (Wang et al., 2012). In some species, there also exists two paralogue genes of AstCs which are named AstCC and AstCCC (Veenstra, 2009, 2016). Interestingly, the allatostatic activity of these three types of Ast peptides varies between insect species so that in each species only one type of Ast (for example AstA) controls JH production (Nässel, 2002; Stay and Tobe, 2007; Wang et al., 2012; Wegener and Chen, 2022).

Analysis of the BPH neural transcriptome sequence data has revealed the presence of one AT, four types of Asts and four corresponding receptors, allatotropin receptor (A16, ATR), AstA receptor (A2, AstAR), AstB (MIP) receptor (A10, AstBR or MIPR) and AstC receptor (A1, AstCR) (Tanaka et al., 2014). We cloned the five neuropeptide genes (AT, AstA, AstB/MIP, AstCC and AstCCC) and confirmed the sequence obtained from transcriptome data (Figure 4-figure supplement 1) (Tanaka et al., 2014). Interestingly, we found that AstC is missing in the genome of BPH and only AstCC and AstCCC are present (Figure 4-figure supplement 1). Next, we also cloned their corresponding receptors (Veenstra, 2009) including ATR (A16), AstAR (A2), AstBR

(A10) and AstCR (A1) which might be activated by AstCC and/or AstCCC (Audsley et

al., 2013; Veenstra, 2009; Zhang et al., 2022a). Sequence alignments and phylogenetic

analysis are shown in Figure 4-figure supplement 2.

Quantitative PCR was then used to examine if EB treatment could influence the

expression of the genes encoding these neuropeptides and their receptors. Treating

4^{th} instar or female adults BPH with the LC_{50} concentration of EB significantly

increased the expression of *AT*, *ATR* and *AstCCC* while resulting in the

downregulation of *AstA, AstB/mip, AstCC*, AstAR and *AstBR/mipr* at the adult stage

(Figure 4A and B, Figure 4-figure supplement 3A-G). Among these, *AstA* and *AstAR*

were the most downregulated genes after EB treatment (Figure 4A and B, Figure

4-figure supplement 3H) and thus the *AstA/AstAR* signaling system was selected for

subsequent functional analysis. Silencing of *AstAR* in female BPH using RNAi (Figure

4C), resulted in an increased number of eggs laid per female compared with

dsgfp-injected controls (Figure 4D). Interestingly, silencing *AstAR* also resulted in the

upregulation of *JHAMT, Met* and *Kr-h1* which are involved in the JH

biosynthesis/signaling (Figure 4E-H). However, *JHE* was not influenced by AstAR

silencing (Figure 4I). We next investigated whether silencing the *AstAR* gene would

influence the JH titer in BPHs. A significantly increased JH titer was observed in

AstAR silenced BPH compared with controls (Figure 4J). Thus, our data strongly

suggest that AstA is a key inhibitor of JH production in BPH.

Finally, we investigated whether injection of mature Ast and AT peptides could

influence the number of eggs laid and JH titer in BPH. We synthesized one AT, six

AstAs (AstA1 to AstA6), one AstCC and one AstCCC peptide according to their

determined sequences (Figure 4-figure supplement 1). Indeed, we found that injection

of AstA1 and AstA6 in female adults reduced the number of eggs laid per female over

48 h (Figure 4K). We also observed that AstA1 injection decreased the JH titer as

measured 16h and 48h after injection (Figure 4L), and AT injection increased the JH

titer after 2h, but levels returned to normal 4h after injection (Figure 4M). Collectively,

these results provide compelling evidence that EB induces reproduction in BPH

through the AstA/AstAR and JH signaling pathways (Figure 4N) and further supports

the role of AstA and AT in regulation of JH titer in this species.

Figure 4

Figure 4. EB induced reproduction in BPHs is mediated by the AstA/AstAR and JH

signaling pathway.

(A and B) Expression of *AstA* and *AstAR* in different stages of BPH following EB treatment. (C)

Downregulation of *AstAR* using RNAi leads to a reduction in mRNA expression level. (D) Egg

production in female BPH following silencing of *AstAR* gene. (E-I) Expression of selected JH

signaling pathway related genes (*JHAMT*, *Met*, *Kr-h1* and *JHE*) in *AstAR* silenced BPH. (J)

JHIII titer of BPH females after *AstAR* gene silencing determined by HPLC-MS/MS. (K)

Number of eggs laid per female in 48h following injection of the six mature AstA1-AstA6

peptides and one mature AT peptide. Fifty nanoliter of PBS (as control) and seven different

peptides (20 pmol/insect) were injected into female BPHs three days after eclosion. (L and M)

The JH III titer of BPH females at different time points following AstA or AT injection. (N)

Schematic of the proposed role of AstA/AstAR in the regulation of JH following EB

exposure. The question mark indicates one or more possible additional signals. All data are

presented as means ± s.e.m. Student's *t* test was used to compare the two samples. One-way

ANOVA with Tukey's multiple comparisons test was used to compare more than two samples.

ns, no significant difference; Asterisks indicate values significantly different from the control (ns,

no significant; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ and ****$p < 0.0001$). Different lower-case

letters above the bars indicate significant differences ($p < 0.05$).

Figure 4-figure supplement 1

Figure 4-figure supplement 1. Alignments of the amino acid sequences of: (A) AT, (B) AstA,

(C) AstB/MIP, (D) AstCC and (E) AstCCC peptides from select species. AT, AstA, AstB/MIP

and AstCCC are predicted to have a C-terminal amide. The mature peptides belonging to the

same species have been highlighted with the same color. Species names are as follows: Nillu

(*Nilaparvata lugens*), Locmi (*Locusta migratoria*), Scham (*Schistocerca americana*), Homvi

(*Homalodisca vitripennis*), Manse (*Manduca sexta*), Spofr (*Spodoptera frugiperda*), Drome

(*Drosophila melanogaster*), Spoex (*Spodoptera exigua*), Nasvi (*Nasonia vitripennis*), Grybi

(*Gryllus bimaculatus*), Bommi (*Bombyx mori*); Mesma (*Mesobuthus martensii*), Stear

544   (*Stegodyphus araneomorph*), Limpo (*Limulus polyphemus*), Carma (*Carcinus maenas*), Strma

545   (*Strigamia maritima*), Athro (*Athalia rosae*), Apime (*Apis mellifera*), Dapma (*Daphnia magna*).

546   Black lines under the sequences indicate the locations of the disulfide bridges in the mature

547   peptides. The accession numbers of the sequences are listed in Figure 4-figure supplement 1

548   source data.

Figure 4-figure supplement 2

Figure 4-figure supplement 2. Phylogenetic tree of the predicted BPH (*) allatotropin

receptor (A16, ATR), allatostatins A receptor (A2, AstAR), AstB (MIP) receptor (A10, AstBR or

MIPR) and allatostatins C receptor (A1, AstCR) with other insect species. The tree was

generated using the maximum likelihood method. *Drosophila melanogaster* metabotropic

glutamate receptor was included as an outgroup. The accession numbers of the sequences

used for this phylogenetic tree are listed in Figure 4-figure supplement 2 source data.

Figure 4-figure supplement 3

Figure 4-figure supplement 3. EB induced changes in the expression of *AT*, *AstB*, *AstCC*,

***AstCCC*, *ATR*, *AstBR* and *AstCR* in BPH.** All data are presented as means ± s.e.m.

Student's t test was used to compare the two samples. ns, no significant difference; Asterisks

indicate values significantly different from the control (**p* < 0.05, ***p* < 0.01, and ****p* < 0.001).

EB-enhanced fecundity in BPHs is dependent on its molecular target protein,

the GluCl channel

EB and abamectin, are allosteric modulators, which target glutamate-gated

chloride channels (GluCls) (Sparks et al., 2020; Sparks et al., 2021; Wu et al., 2017). Hence,

we examined whether EB-stimulated fecundity in BPH is mediated by its molecular

target GluCl. The full length *GluCl* coding sequence from BPH was cloned and

sequenced (Figure 5-figure supplement 1) and the impact of EB on *GluCl* gene

expression examined using quantitative PCR. Treatment of BPH with the LC_{50}

concentration of EB significantly downregulated *GluCl* gene expression at the 5E and

5M nymph stages, while upregulating *GluCl* gene expression at 2 DAE and 5 DAE in

the adult stage (Figure 5A). We also found that EB treatment in female adults also

increases the *GluCl* gene expression (Figure 5-figure supplement 3). To examine the

role of *GluCl* gene in BPH fecundity, RNAi was used to knockdown expression of this

gene in female adult BPH (Figure 5B). A significant decrease in the number of eggs

laid by per female was observed in *dsGluCl*-injected insects compared with

dsgfp-injected insects (Figure 5C). However, treatment with EB was found to rescue

the decreased egg-laying phenotype induced by *dsGluCl* injection (Figure 5C). Note

that this rescue is possible since knockdown of the genes is incomplete when using

dsRNA injection (and residual gene expression allows for EB action). To investigate

the mechanism by which *GluCl* expression modulates fecundity we examined if

silencing *GluCl* influences JH titer and JH-related gene expression. Indeed, we

observed that RNAi knockdown of *GluCl* leads to a decrease in JH titer (Figure 5D)

and down-regulation of genes including *JHAMT* which is responsible for JH synthesis,

and the JH signaling downstream genes *Met and Kr-h1* (Figure 5E-G). In contrast,

expression of *JHE* was not changed in the *GluCl* knockdown insects (Figure 5H and I).

We also examined whether silencing *GluCl* impacts the *AstA*/*AstAR* signaling

pathway in female adults. Silencing *GluCl* in female adults was found to have no

impact on the expression of *AT*, *AstA*, *AstB*, *AstCC*, *AstAR*, and *AstBR*. However, the

expression of *AstCCC* and *AstCR* was significantly upregulated in *dsGluCl*-injected

insects (Figure 5-figure supplement 2A-H). These results suggest that EB activates

GluCl which induces JH biosynthesis and release, which in turn stimulates

reproduction in BPH (Figure 5J).

Figure 5

Figure 5. EB induced reproduction in brown planthoppers through its molecular target

protein GluCl.

(A) Expression of *GluCl* in EB-treated and untreated BPH. (B) Expression of *GluCl* following

injection of *dsGluCl* in BPH. (C) Egg production after *GluCl* gene knockdown in EB-treated and

untreated BPH. (D) The JH III titer of BPH females after *GluCl* gene silencing as quantified

using the ELISA method. (E-I) Expression patterns of selected JH-related genes (*JHAMT*, *Met*,

Kr-h1 and *JHE*) in *GluCl* silenced BPH. (J) Schematic of the proposed role of GluCl as a

molecular target of EB and EB-enhanced reproduction in BPH. The question mark indicates

one or more possible additional signals. All data are presented as means ± s.e.m. Student's t

test was used to compare the two samples. One-way ANOVA with Tukey's multiple

comparisons test was used to compare more than two samples. ns, no significant difference;

Asterisks indicate values significantly different from the control (ns, no significant; *$p < 0.05$

and **$p < 0.01$). Different lower-case letters above the bars indicate significant differences ($p <$

0.05).

Figure 5-figure supplement 1

Figure 5-figure supplement 1. Phylogenetic analysis of glutamate-gated chloride channels in

different species. The numbers at the nodes of the branches represent the percentage

bootstrap support (1000 replications) for each branch. The *Sogatella furcifera* GABA-gated

chloride channel and *Nilaparvata lugens* nAchR were used as outgroup. Alignment was

performed with amino acid sequences from TM1-7. The receptor names are listed in the tree.

The accession numbers of the sequences used for this phylogenetic tree are listed in Figure

5-figure supplement 1 source data.

Figure 5-figure supplement 2

Figure 5-figure supplement 2. The expression of *AT*, *AstA*, *AstB*, *AstCC*, *AstCCC*, *AstAR*,

AstBR and *AstCR* in BPH injected with *dsGluCl or dsgfp*. All data are presented as means ±

s.e.m. Student's *t* test was used to compare the two samples. ns, no significant. Different

lower-case letters above the bars indicate significant differences ($p < 0.05$).

Figure 5-figure supplement 3

Figure 5-figure supplement 3. The expression of *GluCl* after BPH female adult treated with

EB. All data are presented as means ± s.e.m. Student's *t* test was used to compare the two

samples, *, $p < 0.05$.

Discussion

Pesticide-induced resurgence of pest insects is a serious problem in rice and several other crops (Wu et al., 2020). However, the mechanisms underpinning pesticide-enhanced reproduction in insects remain poorly understood. Here we reveal that a suite of molecular actors underlie this trait that, in combination, mediate profound physiological changes in the reproductive fitness of female BPHs. Our data provide fundamental insights into the molecular mechanisms by which xenobiotics modify insect reproduction and have applied implications for the control of a highly damaging crop pest. We discuss these topics below.

Sublethal doses of the GluCl modulators, EB and abamectin, stimulates fecundity in BPH

We show that in both contact and systemic assays the avermectins EB and abamectin stimulate reproduction in BPHs. Thus, insecticide-enhanced reproduction is likely a key factor in the BPH resurgence observed when farmers use EB and abamectin to control leaffolders in rice crops in China (Yao et al., 2017). Although this is the first report of sublethal doses of avermectins enhancing insect fecundity, our findings are consistent with previous studies, which have shown that certain insecticides, herbicides and fungicides stimulate BPH reproduction (Cheng et al., 2014; Ge et al., 2011; Ge et al., 2013; Wan et al., 2013; Wang et al., 2010; Wu et al., 2020; Yang et al., 2019; Zhang et al., 2014; Zhao et al., 2011). Intriguingly, we show that EB only induces fecundity in female adults and is specific to BPH, with EB exposure failing to enhance

reproduction in two related species, the small brown planthopper, *L. striatellus* and the

white backed planthopper, *S. furcifera*, or the model insect *D. melanogaster*. Thus, the

mechanisms underpinning this trait appear to act exclusively on female BPHs and

may be specific to this species. Nevertheless, our findings are of general interest

since they link the molecular EB target, the GluCl channel and juvenile hormone

signaling to fecundity.

Pesticides may stimulate insect reproduction through a variety of physiological

and molecular mechanisms. Our data reveal that exposure to sub-lethal

concentrations of EB results in profound changes to female BPH fitness, leading to

increases in female weight, contents of total protein, cholesterol, and sugar, as well as

egg production and decreases in duration of the preoviposition period. Some of these

findings exhibit parallels with previous studies, which demonstrated that treating

third-instar BPH nymphs with either deltamethrin, triazophos, or imidacloprid led to

increased soluble sugar levels in the corresponding adults (Yin et al., 2014). Such

metabolites provide the energy that drives BPH reproduction and hence resurgence

(Wu et al., 2020). Thus, together with prior work, our results suggest that pesticides

associated with resurgence stimulate nutritional intake in BPH to fuel enhanced

energy-intensive reproduction.

The JH signaling pathway plays an essential role in EB-induced fecundity in

BPH

JH is a pleiotropic hormone which plays important roles in development and

reproduction in insects (Riddiford, 2008; Santos et al., 2019). Circulating JH titers are

regulated by factors that control JH production in the corpora allata including

biosynthetic enzymes and catabolic enzymes that regulate JH levels. Our results

show that EB increases circulating JH III titers in BPH females over 2–4 days after

eclosion (DAE) and promotes ovarian development. Previous studies have reported

that triazophos and deltamethrin treatments also lead to increased circulating JH III

titers in BPH females over 1–3 days post emergence. Similarly, jinggangmycin

treatments were found to lead to increased JH titers (by approximately 45–50%) in

BPH females over two days post emergence (Xu et al., 2016). Thus, our findings, in

combination with these previous studies, demonstrate that insecticide treatments can

have dramatic effects on the regulation of key insect hormones involved in pest

reproduction, which can in turn drive pesticide resurgence.

Although increased JH titers following pesticide exposure have been correlated

with reduced levels of active JH esterase during the first three days PE (Ge et al., 2010),

the type and number of mechanisms mediating the observed increase in hormone

titer has remained an open question. Our data reveal that elevated JH titer in

EB-exposed BPH is associated with the upregulation of genes that encode

biosynthetic enzymes for JH (*JHAMT*) and downstream signaling genes that can

induce *vg* gene expression (*kr-h1*). Using RNAi we provide functional evidence of the

role of these genes in the regulation of JH III and fecundity of female BPHs, and

demonstrate that EB can restore the reduction in egg production resulting from the

knockdown of *JHAMT*, *met* and *kr-h1* expression.

JHMAT is an enzyme that catalyzes the conversion of inactive precursors of JH to

active JH in the final stages of JH biosynthesis (Nouzova et al., 2018; Shinoda and Itoyama,

2003). Interestingly, while it has not been previously implicated in pesticide resurgence,

treatment of the stored product pest *Sitotroga cerealella* with diallyl trisulfide, an

insecticidal compound in garlic essential oil, was found to increase JHAMT mRNA

levels (Shah et al., 2022). Because JHMAT is the key rate-limiting enzyme in regulation

of JH titer our results suggest that its enhanced expression is a key molecular

mechanism of pesticide resurgence in BPH.

Met is a ligand-activated member of the basic helix–loop–helix Per/Arnt/Sim

(bHLH-PAS) transcription factors and is the intracellular receptor for JH (Riddiford, 2020;

Zhu et al., 2019). Kr-h1 is a zinc finger protein that acts downstream of Met and is

expressed in response to JH signaling. Although the genes encoding these proteins

have not been previously linked to pesticide resurgence, our finding that they are

upregulated following EB exposure, and demonstration of their role of in promoting

fecundity, is consistent with previous studies (Santos et al., 2019). Specifically, treatment

of BPH with JH III or the insecticidal analogs methoprene or pyriproxifen was found to

induce the expression of *Kr-h1* (Jin et al., 2014). Furthermore, knockdown of *Met* and

Kr-h1 in BPH brachypterous females was found to result in delayed ovariole

maturation and this was significantly more pronounced than the response observed in

BPH treated separately with ds*NlMet* or ds*NlKr-h1* (Lin et al., 2015). This finding

provides evidence of a possible interaction between *Met* and *Kr-h1* and, in

combination with our data, suggests that *Met* and *Kr-h1* may act in concert to mediate

EB-enhanced fecundity. However, our results showed that EB treatment can mildly

increase (about 2-fold) expression of the *Met* gene in brown planthoppers (Figure 3G).

Our data furthermore indicated that *Met* and *FAMeT* expression levels were not drastically

influenced by EB compared with *kr-h1* and *vg* (Figure 3H and I). It should be mentioned

that JH action does not directly result in the increase of *Met*. However, we cannot rule out

the possibility of other factors (indirect effects), induced by EB treatment, increases the

mRNA expression level of *Met*. One recent paper reported that downregulation of

transcription factor *CncC* will increase *met* expression in beetles (see Figure 6A in the

reference) (Jiang et al., 2023). And many studies have reported that insecticide treatment

activates the *CncC* gene signaling pathway, which regulates detoxification gene

expression (Amezian et al., 2023; Fu et al., 2024; Hu et al., 2021). Hence, it is possible that EB

might influence the *CncC* gene pathway, which in turn induces *met* expression. This EB

effect on *met* upregulation may be similar to the upregulation of *GluCl* and some other

secondary effects (see below).

It must be mentioned that although we exposed the BPH to EB at 4th instar, we

make the insect feed on the EB-treated rice plants for four days. After that, the insect

will develop into 5th instar, the final nymphal stage of brown planthopper. Since BPH

do not have a pupal stage, this might cause the EB presented to the insects last a

longer time even in the adult stage. Besides this, we found that EB treatment will

increase the weight of adult females, which indicates that EB might increase food

intake in BPHs that might produce more insulin peptide. Hence, insulin might increase

the JH synthesis at the adult stage.

EB-induced fecundity in BPHs is dependent on the allatostatin A signaling

pathway

In addition to regulatory proteins that promote JH production, insects utilize

peptides that inhibit JH biosynthesis. These include the allatostatins: FGLamides

(FGLa; AstA), the W(X)6Wamides (AstB), and the PISCFs (AstC) (Nässel, 2002; Stay et

al., 1996; Stay and Tobe, 2007; Tanaka et al., 2014; Wegener and Chen, 2022; Zhang et al., 2021).

Interestingly, our results showed that EB exposure results in a marked downregulation

of the expression of the genes encoding allatostatin *AstA* and its receptor *AstAR*. We

provide evidence for the functional impact of this on JH synthesis and BPH fecundity

by: i) demonstrating that RNAi knockdown of *AstAR* expression results in increased

JH titer and enhanced egg production, and, ii) showing that injection of female BPH

with synthetic AstA peptide reduces JH titer and decreases egg production. Thus, our

data provide unequivocal evidence that AstA is a key inhibitor of JH production in BPH.

This finding is consistent with previous work which has shown that FGLa/ASTs (AstA)

inhibit JH biosynthesis in cockroaches, and termites (Woodhead et al., 1989; Yagi et al.,

2005). To our knowledge, our study is the first report of insecticides down-regulating

the expression of the neuropeptide receptor, *AstAR*, and linking this to increases in JH

titer and enhanced reproduction in insects.

Interestingly knockdown of *AstAR* resulted in significant increases in the

expression of genes involved in JH synthesis/signaling including *JHAMT*, *Met* and

Kr-h1. Related to this finding, previous work has shown that knockdown of the AstA

receptor gene, *Dar-2*, in *D. melanogaster* results in changes in the expression of

genes encoding *Drosophila* insulin-like peptides (DILPs) and adipokinetic hormone

(AKH) (Hentze et al., 2015). Together with our findings, this demonstrates that AstA

receptors may modulate the expression of numerous downstream genes involves in

metabolism, energy store and reproduction. In the case of pesticide resurgence our

results imply significant cross-talk in the expression of genes that inhibit JH production

and those that promote it.

The GluCl plays an essential role in EB-induced fecundity in BPH

EB and abamectin are allosteric modulators of GluCls (Sparks et al., 2020; Sparks et

al., 2021; Wu et al., 2017). Our data revealed that EB exposure increases the expression

of the GluCl in adult stage of BPH, and knockdown of GluCl expression resulted in a

reduction in both JH levels and egg production and egg laying. There are several

studies showing that insects treated with insecticides display increases in the

expression of target genes. For example, the relative expression level of the

ryanodine receptor gene of the rice stem borer, *Chilo suppressalis* was increased

10-fold after treatment with chlorantraniliprole, an insecticide which targets the

ryanodine receptor (Peng et al., 2017). Besides this, in *Drosophila*, starvation (and low

insulin) elevates the transcription level of the short neuropeptide F and tachykinin

receptors (Ko et al., 2015; Root et al., 2011). In BPHs, reduction in mRNA and protein

expression of a nicotinic acetylcholine receptor α8 subunit is associated with

resistance to imidacloprid (Zhang et al., 2015). RNA interference knockdown of the α8

gene decreased the sensitivity of *N. lugens* to imidacloprid (Zhang et al., 2015). Hence,

expression of receptor genes may be regulated by diverse factors, including

insecticide treatment. In our case, we found that EB upregulates its target gene *GluCl*.

However, we do not suggest that EB functions as transcriptional activator of *GluCl*

and we have no data to suggest a mechanism for how EB treatment changes the

expression of *GluCl* in the BPH. Considering that EB exposure in our experiments is

lasting several days, it might be an indirect (or secondary) effect caused by other

factors downstream GluCl that affect channel expression Possibly the allosteric

interaction with GluCl by EB renders the channel dysfunctional and the cellular

response is to upregulate the channel/receptor to compensate.

Interestingly, the GluCl has been reported to inhibit the biosynthesis of JH in the

cockroach, *Diploptera punctata* (Liu et al., 2005). Recent work has also reported that

modulation of glutamatergic signals may contribute to the photoperiodic control of

reproduction in bean bug, *Riptortus pedestris* (Hasebe and Shiga, 2022)*.* Furthermore,

work on *D. punctata* has revealed that application of the GluCl channel agonist

ivermectin, which like EB belongs to the avermectin family, caused a decline in JH

synthesis in the corpus allatum (Liu et al., 2005). While the inhibitory effect of ivermectin

observed in this previous study differs from the activating effect of EB we observed in

our study, it is consistent with our finding of a role for the GluCl channel in the

regulation of JH regulation. Interestingly, we found that knockdown of the *GluCl* gene

expression results in the down-regulation of *JHAMT*, *Met and Kr-h1*, further revealing

significant convergent relationships between genes underpinning pesticide

resurgence.

Conclusion

Our study has revealed that exposure of BPHs to EB affects a diverse suite of genes that act in combination to enhance JH titer and thus increase female fecundity. A schematic of how these factors promote ovarian maturation in the adult stage of *N. lugens* through the AstA and JH signaling pathway is provided in Figure 6. Our findings provide the foundation for further work to understand how these genes interact and the mechanisms by which their expression is activated or repressed by EB. Furthermore, our findings provide fundamental insights into the molecular response in insects to xenobiotic stress and illustrate that pesticides can have unexpected and negative impacts on pest populations. In this regard our findings also have applied implications for the control of a highly damaging crop pest. Previous studies have reported that avermectins such as abamectin are toxic to the wolf spider *Pardosa pseudoannulata*, which is the main predator of BPH in rice crops (Chen et al., 2017; Sogawa, 2015). Thus, these insecticides both stimulate reproduction in BPH while killing their natural enemies providing a 'perfect storm' for inducing damaging BPH outbreaks. Based on these findings, to avoid BPH resurgence, we suggest that the agents, EB and abamectin, should not be (or rarely be) applied to rice plants at growth stages when BPHs are present. On a more optimistic note, our findings have identified numerous genes that play key roles in BPH reproduction and thus represent promising targets for the development of novel controls against this important pest.

Figure 6. Schematic of the proposed regulatory pathway of EB-enhanced fecundity in BPH. Emamectin benzoate (EB) exposure results in the upregulation of genes that promote

JH signaling pathway (*JHAMT and Kr-h1*) and the downregulation of genes that inhibit it

(allatostatin, Ast*A* and allatostatin A receptor, *AstAR*). This transcriptome reprograming is

dependent on the action of EB on its molecular target the glutamate-gated chloride channel

(GluCl) receptor. The resulting increased JH titer promotes vg synthesis and increased

fecundity in EB exposed insects. We observe significant cross-talk in the expression of genes

that inhibit JH production and those that promote it, with *AstAR* inhibiting the expression of

JHAMT, Met and *Kr-h1* and *GluCl* activating the expression of *JHAMT* which is responsible for

JH synthesis, and the JH signalling downstream genes *Met and Kr-h1.*

Materials and methods

Insects

BPH was initially collected from Wanan, JiangXi Province in 2020, reared on 'Taichung Native 1' (TN1) rice seedlings in the laboratory without exposure to any insecticides. The strain was maintained in a climatic chamber at 27 ± 1°C, with relative humidity of 70 ± 10% and a light: dark = 16 h: 8 h photoperiod.

Chemicals

Emamectin benzoate (95.2%) was obtained from Hebei Weiyuan Co., Ltd. (Hebei, China). Abamectin (96.8%) was obtained from Hebei Weiyuan Co., Ltd. (Hebei, China). Pyriproxyfen (98%) was obtained from Shanghai Shengnong Co., Ltd. (Shanghai, China). Methoprene (S)-(+) (mx7457-100mg) was purchased from Shanghai MaoKang Biotechnology Co., Ltd., (Shanghai, China). Juvenile hormone standard sample (J912305-10mg) was purchased from Shanghai Macklin Biotechnology Co., Ltd., (Shanghai, China).

Bioassay

Different life stages of insects were used to perform bioassay to investigate the effects of insecticide on nymphs and adults. To test whether treatment of the nymph stage of insects would promote reproduction in female, we used 4th instar nymphs of BPH or 3rd instar nymph of *Laodelphax striatellus* and *Sogatella furcifera* to perform bioassays. To test whether treatment of the adult stage of insects would promote reproduction in female, we used newly emerged male and female BPH.

Systemic route: The rice-seeding dipping bioassay method was used to

evaluate the susceptibility of BPH, *L. striatellus* and *S. furcifera* to EB.

Technical-grade insecticides were dissolved in acetone as stock solution then diluted

in a series of six concentrations with water containing 0.1% Triton. Selected rice

seedlings at the 6-8 cm growth stage were dipped in insecticide solutions for 30 s and

then air-dried at room temperature. The roots of the rice seedlings were wrapped with

cotton strips and placed seedlings placed in a plastic cup 5 cm in diameter. Fifteen

insects were introduced into each plastic cup for each replicate. The top of the cup

was then selaed with gauze to prevent escape. All experiments comprised at least

three biological replicates. Control rice seedings were treated with 0.1% Triton X-100

water solution only. All treatments were maintained under standard conditions of 27 ±

1 °C and 70–80% relative humidity with a 16 h light/8 h dark photoperiod. Mortality

was assessed after 4 d for *N. lugens* or 2 d for *L. striatellus* and *S. furcifera* after

treatment with insecticides. The insects were considered dead if they were unable to

move after a gentle prodding with a fine brush.

For *Drosophila* larvae bioassay, we adopted a method described previously in our

lab with minor modifications (Huang et al., 2020). Briefly, twenty third instar larvae were

placed in fly vials containing fly food (based on corn powder, brown sugar, yeast and

agar) supplemented with EB of different concentrations. Four concentrations (LC_{10},

LC_{30} and LC_{50}) were tested together with a negative (no insecticide) control. For

Drosophila adult bioassays, we selected virgin females three days after eclosion.

Several concentrations were overlaid onto fly food in standard *Drosophila* vials and

allowed to dry overnight at room temperature. 15 adult flies (three days after eclosion)

were then added to each vial and mortality assessed after 2 d. Four replicates were

carried out for each concentration. Control mortality was assessed using vials

containing food with solvent minus insecticide.

Contact route: For topical bioassays working insecticide solutions were

prepared in acetone. 4^{th} instar nymphs or newly emerged males/females were

anesthetized with carbon dioxide for 5 s, and then 0.04 µl/insect test solution applied

topically to the dorsal plates with a capillary micro-applicator (Yao et al., 2017). Insects

were then placed in an artificial climate incubator with a temperature of 27±1°C, a

photoperiod of 16:8 h (L:D), and a humidity of 70%±10%. Mortality was determined 2

d after treatment. Data with over 20% mortality in the control treatment were

discarded, and the assay was repeated at least three times.

Fecundity assays

Fourth instar nymphs of BPH and 3^{rd} instar nymph of *L. striatellus* and *S. furcifera*

were treated with the LC_{15} and LC_{50} of EB, and then transferred to fresh rice seedlings.

After eclosion, the adults were used in the following experiment. Newly emerged

treated adults and untreated adults were paired to produce four groups: untreated

males and untreated females (♂ck×♀ck; ck indicates untreated); untreated males and

treated females (♂ck×♀t; t indicates insecticide treated); treated males and untreated

females (♂t×♀ck); treated males and treated females (♂t×♀t). Each group comprised

at least 10 mating pairs. All pairs were transferred to glass cylinders (diameter 5 cm

and height 20 cm) containing rice plants (25 days old almost 20 cm high) as a food

source for eleven days. The number of eggs and nymphs in plants were counted by a

microscope.

For *Drosophila* egg-laying assay, we adopted our previous method (<u>Wu et al.,</u>

<u>2019</u>). Briefly, insecticide-treated virgin females were paired with untreated males for

three days and then the mated females transferred into the *Drosophila* ovipositing

apparatus. The number of eggs were counted after 16 hours.

Fitness analysis

The fitness of EB-treated BPH were analyzed using methods reported previously

(<u>Zeng et al., 2023</u>). We selected two groups, (♂ck×♀ck) and (♂ck×♀t), to study the

effects of the LC_{50} concentration of EB on BPH fitness. In the case of systemic

exposure, 4^{th} instar nymphs of BPH were treated with the LC_{50} of EB for 4 days and

then transferred to tubes containing untreated rice plants for individual rearing. The

rice plants were replaced every three days with untreated plants. The emergence ratio,

female ratio, preoviposition period, female longevity, brachypterism female ratio and

female weight were calculated.

Examination of ovarian development

Adult females from ♂ck×♀ck control and ♂ck×♀t group on 1, 3, 5, 7 DAE were

dissected for observe the ovarian development. The mature eggs in ovary were

photographed and recorded. Each group has at least fifteen replicates.

To examination whether EB treated impairs egg maturation, we dissected

untreated or EB-treated ovaries and fixed them in 4% paraformaldehyde (PFA) in

phosphate buffered saline (PBS) for 30 min at room temperature. After four washes of

10 min (4 × 10 min) in PAT3 (PBS with 0.5% Triton X-100 and 0.5% bovine serum

albumin), the ovaries were then incubated with DAPI (4',6-diamidino-2-phenylindole,

100 nM) and Actin-stain 670 Fluorescent Phalloidin (200 nM). Imaging was performed

using Zeiss LSM980 confocal laser microscope.

Measurements of glycogen, triglyceride, total protein content, cholesterol and

four sugars

The content of glycogen, triglyceride, cholesterol and total protein was

determined by spectrophotometry at 620 nm, 510 nm, 510 nm and 562 nm

respectively using the glycogen assay kit (A043-1-1), triglyceride reagent kit

(A110-1-1), cholesterol assay kit (A111-1-1) and total protein assay kit (A045-2-2)

obtained from Nanjing Jiancheng Bioengineering Institute following to the

manufacturer's instructions. The determined results were normalized to the protein

content in the sample, which was determined using BCA Protein Assay Reagent Kit

(Thermo Scientific, Waltham, USA). Each sample contained tissue extracts from five

adult female BPH, with three biological replicates per sample.

To assess the content of four sugars (sucrose, glucose, fructose and trehalose) in

the extract of BPH tissue, the same extraction method was used as above. Sugar

content was quantified using the colorimetric method by scanning at 290 nm, 505 nm,

285 nm and 620 nm respectively using the sucrose reagent kit (A099-1-1), glucose

reagent kit (F006-1-1), fructose reagent kit (A085-1-1) and trehalose reagent kit

(A150-1-1) obtained from Nanjing Jiancheng Bioengineering Institute based on the

manufacturer's instructions. Each sample contained tissue extracts from five adult

female *N. lugens*, with three biological replicates per sample.

Determination of Juvenile hormone III and ecdysone titers of BPH by ELISA

The titer of Juvenile hormone III in BPH was measured using the Juvenile

hormone ELISA Kit (Lot Number: HLE92086, Shanghai HaLing Biotechnology Co.,

Ltd., Shanghai, China) which employs the dual-antibody sandwich ELISA method.

The titer of ecdysone in BPH were measured using the ecdysone ELISA Kit (Lot

Number: ZK14705, Shanghai ZhenKe Biotechnology Co., Ltd., Shanghai, China). At

least three biological replicates were employed for each treatment.

Determination of Juvenile hormone III titer in BPH using HPLC-MS/MS

The whole bodies of 5 individuals BPH were mixed with 1 ml of n-hexane,

followed by centrifugation at 10,000×g for 10 min, the upper hexane layer was then

dried with nitrogen, dissolved in methanol and sonicated for 10 min, after

centrifugation at 10000×g for 10 min, the supernatant was collected through the

organic filter membrane of 0.22 μm into 2 ml vials for JH Ⅲ determination. JH Ⅲ

standard sample (J912305-10 mg) purchased from (Shanghai McLean Biochemical

Technology Co. Ltd), dissolved in methyl alcohol as stock solution 10,000 mg/L was

diluted in a series of six concentration gradients to serve as a reference. Liquid

chromatography-tandem mass spectrometry (LC-MS/MS) was then carried out using

UPLC Xevo TQ-S Micro (Waters technology), quantitative method according to the

external standard, the chromatographic column was EC-C18 (4.6 mm×150 mm, 2.7

μm), column temperature was 30℃ , injection volume was 20 μl, elution flow rate was

0.3 ml/min, and the mobile phase was acetonitrile:formic acid water (90:10), detection

wavelength was 218 nm, the peak height was used for quantification.

Cloning, sequence and phylogenetic analysis

The NCBI database and BLAST program were used to carry out sequence

alignment and analysis. Open Reading Frames (ORFs) were predicted with EditSeq.

Primers were designed using the primer design tool in NCBI. Total RNA Extraction

was extracted from 30 adults BPH using TRIzol reagent (Invitrogen, Carlsbad, CA,

USA) according to the manufacturer's instructions. cDNA was synthesized using the

Biotech M-MLV reverse transcription kit. Full-length gene sequences were amplified

by PCR using cDNA as template and Rapid Taq Master Mix (Vazyme Biotech, Cat#

P222-02). The PCR product was purified on a 1% agarose gel, cloned into pClone007

Simple Vector Kit (Tsingke Biotech, Cat# TSV-007S), and then sequenced using the

3730 XL DNA analyzer (Applied Biosystems, Carlsbad, CA, USA). Table S2 contains

a list of the primers used in this study.

The exon and intron architectures of AT, AstA, AstB, AstCC and AstACCC were

predicted based on the alignments of putative ORFs against their corresponding

genomic sequences. Sequence similarity/annotations and orthologous gene searches

were performed using BLAST programs available in NCBI. Multiple alignments of the

complete amino acid sequences were performed with Clustal Omega

(http://www.ebi.ac.uk/Tools/msa/clustalo). Phylogeny was conducted using the

maximum likelihood technique to create phylogenetic trees and these were

bootstrapped with 1000 replications were used using the MEGA 6 software (Tamura et

al., 2013).

RNA interference

Double-stranded RNA (dsRNA) of *gfp* (green fluorescent protein), *JHAMT*

(juvenile hormone acid O-methyltransferase,), *Met* (methoprene-tolerant), *Kr-h1*

(krüppel homolog 1), *AstAR* (allatostatin-A receptor) and *GluCl* (glutamate-gated

chloride channel) was prepared using Ambion's MEGAscript T7 kit instructions

following the manufacturer's instructions. The primer sequences for double-stranded

RNA synthesis are listed in Table S2. Newly emerged females were injected with 40 nl

(5,000 ng/µl) of double-stranded RNA of *gfp* (*dsgfp*) or double-stranded RNA of the

target genes in the conjunctive part between prothorax and mesothorax of insects. In

the RNAi experiments, BPH were then treated with the LC_{50} of EB 24h after dsRNA

injection and the whole body sampled for qRT-PCR analysis.

Quantitative RT-PCR

Fourth instar nymphs of BPH were treated with EB after which total RNA was

extracted from 5^{th} instar nymphs and 1-7 day post-eclosion females of *N. lugens* using

the methods detailed above. The HiScript® II RT SuperMix for qPCR (+gDNA wiper)

kit from Vazyme, Nanjing, China, was used to create first-strand cDNA. Primer3

(http://bioinfo.ut.ee/primer3/) was used to design real-time quantitative PCR (qPCR)

primers listed in Table S2. mRNA levels of candidate genes were detected by qPCR

using the UltraSYBR Mixture (with ROX) Kit (CWBIO, Beijing, China). Each reaction

contained 2 µL of cDNA template (500 ng), 1 µL each forward and reverse qPCR

primer (10 µM), 10 µL of UltraSYBR mixture buffer, and 6 µL of RNase-free water.

Q-PCR was run on an ABI 7500 Real-Time PCR System (Applied Biosystems, Foster

City, CA) under the following conditions: 5 min at 95°C, followed by 40 cycles of 95°C

for 10 s and 60°C for 30 s. Three independent biological replicates and four technical

replicates were used in each qPCR experiment. The housekeeping genes 18S

ribosomal RNA of BPH were selected to normalize the expression of candidate genes.

The $2^{-\Delta\Delta Ct}$ method (Ct represents the cycle threshold) was used to measure relative

expression levels (Livak and Schmittgen, 2001). Three biological replicates were used for

statistical comparison between samples. Table S2 contains a list of the primers used

in this study.

Statistics

PoloPlus v2.0 (LeOra Sofware 2008) was used to calculate the lethal

concentration (LC_{50}) and 95% fiducial limits (95% F.L.). GraphPad Prism 8.0 software

(GraphPad Software Inc., San Diego, USA) was used to generate graphs and perform

statistical analysis of data. Data presented in this study were first verified for normal

distribution using the D'Agostino–Pearson normality test. One-way analysis of

variance (ANOVA) with Duncan's multiple range test was used to test differences

among multiple groups of normally distributed groups of data. Student's t test was

used to test the differences between two groups. If not normally distributed,

Mann–Whitney test was used for pairwise comparisons, and Kruskal–Wallis test was

used for comparisons among multiple groups, followed by Dunn's multiple

comparisons. All data are presented as mean ± s.e.m. The sample sizes and

statistical tests used for each experiment are stated in the figures or figure legends.

Table S2 Sequences of oligonucleotide primers used in this study.

Primers	Primer sequences (5' - 3')
For cDNA cloning	
Allatotropin :	
NlAT-F	TACGCGGCCAAACACACTTA
NlAT-R	AGGGAAAGAGGGCGAAATTCA
Allatostatins :	
NlAstA-F	TCGGCCGTCACAAGTCAAG
NlAstA-R	CCGAACCCGTACTTCATGCT
NlAstB-F	ACCGGGCTCACAGGAATTTT
NlAstB-R	TGTAGGCGCAGATCTTGAGG
NlAstCC-F	AACACAGCTCTACGAGGCAC
NlAstCC-R	CCAAGCAGGTGACTGCCATA
NlAstCCC-F	TTTGTGTGTGCTTGCAGGTG
NlAstCCC-R	GGATAGAAACGGTAGATTTGGTAGA
Allatostatins receptor :	
NlA16-F	CCTCATTGTGGAACCACCGA
NlA16-R	CGCAGCTGTAAGGTGGAAGA
NlA2-F	GAACGTAATGGGAGTCGGCA
NlA2-R	GTTTTTGTGAGCGCCGACTT
NlA10-F	ATGCAAAACACGGCCAGCCT
NlA10-R	TTAATCGTCTCTGCTCAACTCCAAAGGAAGGT
NlA1-F	CGACCAGACCACTCTACTGC
NlA1-R	ACGTGGACCTCACTATACCAAAAA
For Quantitative RT-PCR	
Q-Nl18S-F	CGCTACTACCGATTGAA
Q-Nl18S-R	GGAAACCTTGTTACGACTT
Q-vitellogenin-F	GTGGCTCGTTCAAGGTTATGG
Q-vitellogenin-R	GCAATCTCTGGGTGCTGTTG
Q-Vitellogenin receptor-F	AGGCAGCCACACAGATAACCGC
Q-Vitellogenin receptor-R	AGCCGCTCGCTCCAGAACATT
Q-JHE-F	GAGCCTCACATCCACAGC
Q-JHE-R	AATGGGAGCCCTACGC
Q-NlMet-F	GGTGGTAAACGGATTGGAAA
Q-NlMet-R	CATCGTCAGCCAACTCGATA
Q-JHAMT-F	GAACCTGCAGGCCAAACACA
Q-JHAMT-R	ACCACTCGGTTGGGCTGAAT

Q-NlKr-h1-F	TGATGAGGCACACGATGACT
Q-NlKr-h1-R	ATGGAAGGCCACATCAAGAG
Q-NlAT-F	CACGATACGTGGCTTCAAGA
Q-NlAT-R	ACGATCACTTTCGCCAATTC
Q-NlAstA-F	AGGACTTACTGGGCGAGGAT
Q-NlAstA-R	GGTGTCTCGTTTCCTGGTGT
Q-NlAstB-F	AGCGAGCTAGACGAGGACAA
Q-NlAstB-R	TCGTCTCTGCTCAACTCCAA
Q-NlAstCC-F	CTGCTCCCAGTGAAAAGGAG
Q-NlAstCC-R	GCTTCCAGTAACTGCGCTTC
Q-NlAstCCC-F	TGTGCTTGCAGGTGGTAGTC
Q-NlAstCCC-F	AGAAGCATGTGACTGCGTTG
Q-NlA2-F	TCCTGGTGCTGAAGAGTGTG
Q-NlA2-R	CTTTTCGGGCCCATTAATTT
Q-NLA1-F	ATATCGGCACCGAAGATGAG
Q-NLA1-R	GTCTGACCCGACAGGTTCTC
Q-NLA10-F	ACTGGGTGTCGACCAATCTC
Q-NLA10-R	TCGGTAGCGAGGAAGACAGT
Q-NLA16-F	TACCGTTCTGTGGGATGTCA
Q-NLA16-R	CCGGGATATCAAAGACGAGA
Q-NlGluCl-F	CACTGACTGAGGCCAACAGA
Q-NlGluCl-R	GCTGGCCATTCTTAGTGAGC

For double-stranded RNA synthesis

T7-EGFP-F	TAATACGACTCACTATAGGGCGTAAACGGCCACAAGTTCA
T7-EGFP-R	TAATACGACTCACTATAGGGGACTGGGTGCTCAGGTAGTG
T7-Kr-h1-F	TAATACGACTCACTATAGGGCGCCAGTGAAAGTGAGACCT
T7-Kr-h1-R	TAATACGACTCACTATAGGGGAGACCGCAAGTGGTTCTGA
T7-Met-F	TAATACGACTCACTATAGGGCCACCAACCAGCAGATGAACCTGA
T7-Met-R	TAATACGACTCACTATAGGGCCACGCAAAGCCTCGTACTCTTGG
T7-JHAMT-F	TAATACGACTCACTATAGGGCTCCAGGCCATTGTCCCTCA
T7-JHAMT-R	TAATACGACTCACTATAGGGTTGGCCTGCAGGTTCTTTGG
T7-AstA-R-F	TAATACGACTCACTATAGGGTACTGCCGTTCTGGCCTTTT
T7-AstA-R-R	TAATACGACTCACTATAGGGGTGATCTGGAAGAGCGGCTT
T7-Glucl-F	TAATACGACTCACTATAGGGACACATCACCTGCTCACCTG
T7-Glucl-R	TAATACGACTCACTATAGGGGTGTGTTTGCCTGCTGTCTG

Acknowledgments

This research was funded by the National Key R&D Program of China to SFW

(2022YFD1700200) (https://service.most.gov.cn/) and National Natural Science

Foundation of China to SFW (32022011) (https://www.nsfc.gov.cn/).

Funding: This work was supported by the National Natural Science Foundation of

China to SFW (No.32022011 & 31772205) (https://isisn.nsfc.gov.cn/

egrantindex/funcindex/prjsearch-list?locale=zh_CN).

The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

References

Ahmed, T.H., Saunders, T.R., Mullins, D., Rahman, M.Z., Zhu, J., 2020. Molecular action of pyriproxyfen: Role of the Methoprene-tolerant protein in the pyriproxyfen-induced sterilization of adult female mosquitoes. PLoS Negl Trop Dis. 14, e0008669.

Amezian, D., Fricaux, T., de Sousa, G., Maiwald, F., Huditz, H.-I., Nauen, R., Le Goff, G., 2023. Investigating the role of the ROS/CncC signaling pathway in the response to xenobiotics in Spodoptera frugiperda using Sf9 cells. Pesticide Biochemistry and Physiology 195, 105563.

Audsley, N., Vandersmissen, H.P., Weaver, R., Dani, P., Matthews, J., Down, R., Vuerinckx, K., Kim, Y.-J., Vanden Broeck, J., 2013. Characterisation and tissue distribution of the PISCF allatostatin receptor in the red flour beetle, *Tribolium castaneum*. Insect Biochem Mol Biol. 43, 65-74.

Bellés, X., Graham, L.A., Bendenab, W.G., Ding, Q., Edwards, J.P., Weaver, R.J., Tobe, S.S., 1999. The molecular evolution of the allatostatin precursor in cockroaches. Peptides 20, 11-22.

Bownes, M., Rembold, H., 1987. The titre of juvenile hormone during the pupal and adult stages of the life cycle of *Drosophila melanogaster*. Eur J Biochem. 164, 709-712.

Chen, B., Wen, L., Zhao, J., Liang, H., Jiao, X., 2017. Laboratory risk assessment of seven insecticides to the wolf spider *Pardosa pseudoannulata*. J. Plant Protect. 44, 1059-1060.

Cheng, J., Huang, L.-J., Zhu, Z.-F., Jiang, L.-B., Ge, L.-Q., Wu, J.-C., 2014. Heat-dependent fecundity enhancement observed in *Nilaparvata lugens* (Hemiptera: Delphacidae) after treatment with triazophos. Environ Entomol. 43, 474-481.

Chiang, A.-S., Lin, W.-Y., Liu, H.-P., Pszczolkowski, M.A., Fu, T.-F., Chiu, S.-L., Holbrook, G.L., 2002a. Insect NMDA receptors mediate juvenile hormone biosynthesis. Proceedings of the National Academy of Sciences 99, 37.

Chiang, A.-S., Pszczolkowski, M.A., Liu, H.-P., Lin, S.-C., 2002b. Ionotropic glutamate receptors mediate juvenile hormone synthesis in the cockroach, *Diploptera punctata*. Insect Biochemistry and Molecular Biology 32, 669-678.

Chintalapati, P., Katti, G., Puskur, R.R., Nagella Venkata, K., 2016. Neonicotinoid-induced resurgence of rice leaffolder, *Cnaphalocrocis medinalis* (Guénee). Pest Manag. Sci. 72, 155-161.

Du, B., Ding, D., Ma, C., Guo, W., Kang, L., 2022. Locust density shapes energy metabolism and oxidative stress resulting in divergence of flight traits. Proc Natl Acad Sci U S A. 119, e2115753118.

Fu, B., Liang, J., Hu, J., Du, T., Tan, Q., He, C., Wei, X., Gong, P., Yang, J., Liu, S., Huang, M., Gui, L., Liu, K., Zhou, X., Nauen, R., Bass, C., Yang, X., Zhang, Y., 2024. GPCR–MAPK signaling pathways underpin fitness trade-offs in whitefly. Proceedings of the National Academy of Sciences 121, e2404407121.

Gandara, L., Jacoby, R., Laurent, F., Spatuzzi, M., Vlachopoulos, N., Borst, N.O., Ekmen, G., Potel, C.M., Garrido-Rodriguez, M., Böhmert, A.L., Misunou, N., Bartmanski, B.J., Li, X.C., Kutra, D., Hériché, J.-K., Tischer, C., Zimmermann-Kogadeeva, M., Ingham, V.A., Savitski, M.M., Masson, J.-B., Zimmermann, M., Crocker, J., 2024. Pervasive sublethal effects of agrochemicals on insects at environmentally relevant concentrations. Science 386, 446-453.

Ge, L.-Q., Cheng, Y., Wu, J.-C., Jahn, G.C., 2011. Proteomic analysis of insecticide triazophos-induced mating-responsive proteins of *Nilaparvata lugens* Stål (Hemiptera: Delphacidae). J. Proteome Res. 10, 4597-4612.

Ge, L.-Q., Huang, L.-J., Yang, G.-Q., Song, Q.-S., Stanley, D., Gurr, G.M., Wu, J.-C., 2013. Molecular basis for insecticide-enhanced thermotolerance in the brown planthopper *Nilaparvata lugens* Stål (Hemiptera:Delphacidae). Mol. Ecol. 22, 5624-5634.

Ge, L.-Q., Wu, J.-C., Zhao, K.-F., Chen, Y., Yang, G.-Q., 2010. Induction of Nlvg and suppression of Nljhe gene expression in *Nilaparvata lugens* (Stål) (Hemiptera: Delphacidae) adult females and males exposed to two insecticides. Pestic. Biochem. Physiol. 98, 269-278.

Gospocic, J., Glastad, K.M., Sheng, L., Shields, E.J., Berger, S.L., Bonasio, R., 2021. Kr-h1 maintains distinct caste-specific neurotranscriptomes in response to socially regulated hormones. Cell 184, 5807-5823.e5814.

Guedes, R.N.C., Smagghe, G., Stark, J.D., Desneux, N., 2016. Pesticide-induced stress in arthropod pests for optimized integrated pest management programs. Annu Rev Entomol. 61, 43-62.

Guo, W., Song, J., Yang, P., Chen, X., Chen, D., Ren, D., Kang, L., Wang, X., 2020. Juvenile hormone suppresses aggregation behavior through influencing antennal gene expression in locusts. PLoS Genet. 16, e1008762.

Hardin, M.R., Benrey, B., Coll, M., Lamp, W.O., Roderick, G.K., Barbosa, P., 1995. Arthropod pest resurgence: an overview of potential
mechanisms. Crop Protection 14, 3-18.

Hasebe, M., Shiga, S., 2022. Clock gene-dependent glutamate dynamics in the bean bug brain regulate photoperiodic reproduction. PLOS
Biol. 20, e3001734.

Hentze, J.L., Carlsson, M.A., Kondo, S., Nässel, D.R., Rewitz, K.F., 2015. The neuropeptide allatostatin A regulates metabolism and feeding
decisions in *Drosophila*. Sci. Rep. 5, 11680.

Hu, B., Huang, H., Hu, S., Ren, M., Wei, Q., Tian, X., Esmail Abdalla Elzaki, M., Bass, C., Su, J., Reddy Palli, S., 2021. Changes in both trans- and
cis-regulatory elements mediate insecticide resistance in a lepidopteron pest, Spodoptera exigua. PLOS Genetics 17, e1009403.

Huang, J.M., Rao, C., Wang, S., He, L.F., Zhao, S.Q., Zhou, L.Q., Zhao, Y.X., Yang, F.X., Gao, C.F., Wu, S.F., 2020. Multiple target-site mutations
occurring in lepidopterans confer resistance to diamide insecticides. Insect Biochem. Mol. Biol. 121, 103367.

Hun, L.V., Okamoto, N., Imura, E., Maxson, R., Bittar, R., Yamanaka, N., 2022. Essential functions of mosquito ecdysone importers in
development and reproduction. Proc Natl Acad Sci U S A. 119, e2202932119.

Ishaaya, I., Kontsedalov, S., Horowitz, A.R., 2002. Emamectin, a novel insecticide for controlling field crop pests. Pest Manag. Sci. 58,
1091-1095.

Janssen, A., van Rijn, P.C.J., 2021. Pesticides do not significantly reduce arthropod pest densities in the presence of natural enemies. Ecol
Lett. 24, 2010-2024.

Jiang, H., Meng, X., Zhang, N., Ge, H., Wei, J., Qian, K., Zheng, Y., Park, Y., Reddy Palli, S., Wang, J., 2023. The pleiotropic AMPK–CncC signaling
pathway regulates the trade-off between detoxification and reproduction. Proceedings of the National Academy of Sciences 120,
e2214038120.

Jin, M.-n., Xue, J., Yao, Y., Lin, X.-d., 2014. Molecular characterization and functional analysis of krüppel-homolog 1 (Kr-h1) in the brown
planthopper, *Nilaparvata lugens* (Stål). J Integr Agric. 13, 1972-1981.

Jindra, M., Palli, S.R., Riddiford, L.M., 2013. The juvenile hormone signaling pathway in insect development. Annu Rev Entomol. 58, 181-204.

Jing, Y.-P., Wen, X., Li, L., Zhang, S., Zhang, C., Zhou, S., 2021. The vitellogenin receptor functionality of the migratory locust depends on its
phosphorylation by juvenile hormone. Proc Natl Acad Sci U S A. 118, e2106908118.

Kamita, S.G., Hinton, A.C., Wheelock, C.E., Wogulis, M.D., Wilson, D.K., Wolf, N.M., Stok, J.E., Hock, B., Hammock, B.D., 2003. Juvenile hormone
(JH) esterase: why are you so JH specific? Insect Biochem Mol Biol. 33, 1261-1273.

Kataoka, H., Toschi, A., Li, J.P., Carney, R.L., Schooley, D.A., Kramer, S.J., 1989. Identification of an allatotropin from adult *Manduca Sexta*.
Science 243, 1481-1483.

Ko, K.I., Root, C.M., Lindsay, S.A., Zaninovich, O.A., Shepherd, A.K., Wasserman, S.A., Kim, S.M., Wang, J.W., 2015. Starvation promotes
concerted modulation of appetitive olfactory behavior via parallel neuromodulatory circuits. eLife 4, e08298.

Kramer, S.J., Toschi, A., Miller, C.A., Kataoka, H., Quistad, G.B., Li, J.P., Carney, R.L., Schooley, D.A., 1991. Identification of an allatostatin from
the tobacco hornworm *Manduca sexta*. Proc Natl Acad Sci U S A. 88, 9458-9462.

Lin, X., Yao, Y., Wang, B., 2015. Methoprene-tolerant (Met) and Krüpple-homologue 1 (Kr-h1) are required for ovariole development and egg
maturation in the brown plant hopper. Sci. Rep. 5, 18064.

Ling, L., Raikhel, A.S., 2018. Serotonin signaling regulates insulin-like peptides for growth, reproduction, and metabolism in the disease
vector *Aedes aegypti*. Proc Natl Acad Sci U S A. 115, E9822.

Ling, L., Raikhel, A.S., 2021. Cross-talk of insulin-like peptides, juvenile hormone, and 20-hydroxyecdysone in regulation of metabolism in
the mosquito *Aedes aegypti*. Proc. Natl. Acad. Sci. U S A. 118, e2023470118.

Liu, H.-P., Lin, S.-C., Lin, C.-Y., Yeh, S.-R., Chiang, A.-S., 2005. Glutamate-gated chloride channels inhibit juvenile hormone biosynthesis in the
cockroach, *Diploptera punctata*. Insect Biochem Mol Biol. 35, 1260-1268.

Liu, S., Zhang, C., Yang, B., Gu, J., Liu, Z., 2010. Cloning and characterization of a putative farnesoic acid O -methyltransferase gene from the
brown planthopper, *Nilaparvata lugens*. J Insect Sci. 10.

Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. Methods
25, 402-408.

Lorenz, M.W., Kellner, R., Hoffmann, K.H., 1995. A family of neuropeptides that inhibit juvenile hormone biosynthesis in the cricket, *Gryllus*
bimaculatus. J. Biol. Chem. 270, 21103-21108.

Lu, K., Chen, X., Liu, W.-T., Zhou, Q., 2016. TOR pathway-mediated juvenile hormone synthesis regulates nutrient-dependent female
reproduction in *Nilaparvata lugens* (Stål). Int. J. Mol. Sci. 17, 438.

Luo, W., Liu, S., Zhang, W., Yang, L., Huang, J., Zhou, S., Feng, Q., Palli, S.R., Wang, J., Roth, S., Li, S., 2021. Juvenile hormone signaling promotes
ovulation and maintains egg shape by inducing expression of extracellular matrix genes. Proc. Natl. Acad. Sci. U S A. 118, e2104461118.

Meiselman, M.R., Alpert, M.H., Cui, X, Shea, J., Gregg, I., Gallio, M., Yapici, N., 2022. Recovery from cold-induced reproductive dormancy is
regulated by temperature-dependent AstC signaling. Current Biol. 32, 1362-1375.e1368.

Nässel, D.R., 2002. Neuropeptides in the nervous system of *Drosophila* and other insects: multiple roles as neuromodulators and
neurohormones. Prog Neurobiol. 68, 1-84.

Nouzova, M., Rivera-Pérez, C., Noriega, F.G., 2018. Omics approaches to study juvenile hormone synthesis. Curr Opin Insect Sci. 29, 49-55.

Peng, Y.C., Sheng, C.W., Casida, J.E., Zhao, C.Q., Han, Z.J., 2017. Ryanodine receptor genes of the rice stem borer, *Chilo suppressalis*: Molecular
cloning, alternative splicing and expression profiling. Pestic. Biochem. Physiol. 135, 69-77.

Riddiford, L.M., 2008. Juvenile hormone action: A 2007 perspective. J Insect Physiol. 54, 895-901.

Riddiford, L.M., 2020. Rhodnius, golden oil, and met: A history of juvenile hormone research. Front Cell Dev Biol. 8.

Root, Cory M., Ko, Kang I., Jafari, A., Wang, Jing W., 2011. Presynaptic facilitation by neuropeptide signaling mediates odor-driven food
search. Cell 145, 133-144.

Roy, S., Saha, T.T., Johnson, L., Zhao, B., Ha, J., White, K.P., Girke, T., Zou, Z., Raikhel, A.S., 2015. Regulation of gene expression patterns in
mosquito reproduction. PLoS Genet. 11, e1005450.

Santos, C.G., Humann, F.C., Hartfelder, K., 2019. Juvenile hormone signaling in insect oogenesis. Curr Opin Insect Sci. 31, 43-48.

Shah, S., Zhang, S.-S., Elgizawy, K.K., Yan, W.-H., Tang, N., Wu, G., Yang, F.-L., 2022. Diallyl trisulfide reduced the reproductive capacity of male
Sitotroga cerealella via the regulation of juvenile and ecdysone hormones. Ecotoxicol. Environ. Saf. 248, 114304.

Shen, Y., Chen, Y.-Z., Lou, Y.-H., Zhang, C.-X., 2019. Vitellogenin and vitellogenin-like genes in the brown planthopper. Front Physiol. 10.

Shinoda, T., Itoyama, K., 2003. Juvenile hormone acid methyltransferase: A key regulatory enzyme for insect metamorphosis. Proc Natl Acad
Sci U S A. 100, 11986.

Smykal, V., Raikhel, A.S., 2015. Nutritional control of insect reproduction. Curr Opin Insect Sci. 11, 31-38.

Sogawa, K., 2015. Planthopper outbreaks in different paddy ecosystems in Asia: Man-made hopper plagues that threatened the green
revolution in rice, in: Heong, K.L., Cheng, J., Escalada, M.M. (Eds.), Rice Planthoppers: Ecology, Management, Socio Economics and Policy.
Springer Netherlands, Dordrecht, pp. 33-63.

Sōgawa, K., 1982. The rice brown planthopper: Feeding physiology and host plant interactions. Annu Rev Entomol. 27, 49-73.

Sparks, T.C., Crossthwaite, A.J., Nauen, R., Banba, S., Cordova, D., Earley, F., Ebbinghaus-Kintscher, U., Fujioka, S., Hirao, A., Karmon, D.,
Kennedy, R., Nakao, T., Popham, H.J.R., Salgado, V., Watson, G.B., Wedel, B.J., Wessels, F.J., 2020. Insecticides, biologics and nematicides:
Updates to IRAC's mode of action classification - a tool for resistance management. Pestic. Biochem. Physiol. 167, 104587.

Sparks, T.C., Storer, N., Porter, A., Slater, R., Nauen, R., 2021. Insecticide resistance management and industry: the origins and evolution of
the Insecticide Resistance Action Committee (IRAC) and the mode of action classification scheme. Pest Manag. Sci. 77, 2609-2619.

Stay, B., Fairbairn, S., Yu, C.G., 1996. Role of allatostatins in the regulation of juvenile hormone synthesis. Arch Insect Biochem Physiol. 32,
287-297.

Stay, B., Tobe, S.S., 2007. The role of allatostatins in juvenile hormone synthesis in insects and crustaceans. Annu Rev Entomol. 52, 277-299.

Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S., 2013. MEGA6: Molecular evolutionary genetics analysis version 6.0. Mol Biol Evol.
30, 2725-2729.

Tanaka, Y., Suetsugu, Y., Yamamoto, K., Noda, H., Shinoda, T., 2014. Transcriptome analysis of neuropeptides and G-protein coupled receptors
(GPCRs) for neuropeptides in the brown planthopper *Nilaparvata lugens*. Peptides 53, 125-133.

Veenstra, J.A., 2009. Allatostatin C and its paralog allatostatin double C: The arthropod somatostatins. Insect Biochem Mol Biol. 39, 161-170.

Veenstra, J.A., 2016. Allatostatins C, double C and triple C, the result of a local gene triplication in an ancestral arthropod. Gen Comp

Endocrinol. 230-231, 153-157.

Verlinden, H., Gijbels, M., Lismont, E., Lenaerts, C., Vanden Broeck, J., Marchal, E., 2015. The pleiotropic allatoregulatory neuropeptides and

their receptors: A mini-review. J Insect Physiol. 80, 2-14.

Wan, D.-J., Chen, J., Jiang, L.-B., Ge, L.-Q., Wu, J.-C., 2013. Effects of the insecticide triazophos on the ultrastructure of the flight muscle of the

brown planthopper *Nilaparvata lugens* Stål (Hemiptera: Delphacidae). Crop Protection 43, 54-59.

Wang, C., Chin-Sang, I., Bendena, W.G., 2012. The FGLamide-Allatostatins influence foraging behavior in *Drosophila melanogaster*. PLOS ONE

7, e36059.

Wang, G., Vega-Rodríguez, J., Diabate, A., Liu, J., Cui, C., Nignan, C., Dong, L., Li, F., Ouedrago, C.O., Bandaogo, A.M., Sawadogo, P.S., Maiga, H.,

Alves e Silva, T.L., Pascini, T.V., Wang, S., Jacobs-Lorena, M., 2021. Clock genes and environmental cues coordinate *Anopheles* pheromone

synthesis, swarming, and mating. Science 371, 411.

Wang, L.-P., Shen, J., Ge, L.-Q., Wu, J.-C., Yang, G.-Q., Jahn, G.C., 2010. Insecticide-induced increase in the protein content of male accessory

glands and its effect on the fecundity of females in the brown planthopper *Nilaparvata lugens* Stål (Hemiptera: Delphacidae). Crop

Protection 29, 1280-1285.

Wegener, C., Chen, J., 2022. Allatostatin A signalling: Progress and new challenges from a paradigmatic pleiotropic invertebrate

neuropeptide family. Front Physiol. 13.

Woodhead, A.P., Stay, B., Seidel, S.L., Khan, M.A., Tobe, S.S., 1989. Primary structure of four allatostatins: neuropeptide inhibitors of juvenile

hormone synthesis. Proc Natl Acad Sci U S A. 86, 5997-6001.

Wu, J., Ge, L., Liu, F., Song, Q., Stanley, D., 2020. Pesticide-induced planthopper population resurgence in rice cropping systems. Annu Rev

Entomol. 65, 409-429.

Wu, S.F., Ja, Y.L., Zhang, Y.J., Yang, C.H., 2019. Sweet neurons inhibit texture discrimination by signaling TMC-expressing mechanosensitive

neurons in *Drosophila*. eLife 8, e46165.

Wu, S.F., Mu, X.C., Dong, Y.X., Wang, L.X., Wei, Q., Gao, C.F., 2017. Expression pattern and pharmacological characterisation of two novel

alternative splice variants of the glutamate-gated chloride channel in the small brown planthopper *Laodelphax striatellus*. Pest Manag.

Sci. 73, 590-597.

Wu, S.F., Zeng, B., Zheng, C., Mu, X.C., Zhang, Y., Hu, J., Zhang, S., Gao, C.F., Shen, J.L., 2018a. The evolution of insecticide resistance in the

brown planthopper (*Nilaparvata lugens* Stål) of China in the period 2012–2016. Sci. Rep. 8, 4586.

Wu, Y., Ding, J., Xu, B., You, L., Ge, L., Yang, G., Liu, F., Stanley, D., Song, Q., Wu, J., 2018b. Two fungicides alter reproduction of the small brown

planthopper *Laodelphax striatellus* by influencing gene and protein expression. J Proteome Res. 17, 978-986.

Xu, B., You, L.-L., Wu, Y., Ding, J., Ge, L.-Q., Wu, J.-C., 2016. Transmission electron microscopy (TEM) observations of female oocytes from

Nilaparvata lugens (Hemiptera: Delphacidae): Antibiotic jinggangmycin (JGM)-induced stimulation of reproduction and associated

changes in hormone levels. J Econ Entomol. 109, 1677-1682.

Xu, H.J., Xue, J., Lu, B., Zhang, X.C., Zhuo, J.C., He, S.F., Ma, X.F., Jiang, Y.Q., Fan, H.W., Xu, J.Y., Ye, Y.X., Pan, P.L., Li, Q., Bao, Y.Y., Nijhout, H.F., Zhang,

C.X., 2015. Two insulin receptors determine alternative wing morphs in planthoppers. Nature advance online publication.

Yagi, K.J., Kwok, R., Chan, K.K., Setter, R.R., Myles, T.G., Tobe, S.S., Stay, B., 2005. Phe-Gly-Leu-amide allatostatin in the termite *Reticulitermes*

flavipes: Content in brain and corpus allatum and effect on juvenile hormone synthesis. J Insect Physiol. 51, 357-365.

Yang, H., Zhou, C., Jin, D.-c., Gong, M.-f., Wang, Z., Long, G.-y., 2019. Sublethal effects of abamectin on the development, fecundity, and wing

morphs of the brown planthopper *Nilaparvata lugens*. J Asia Pac Entomol. 22, 1180-1186.

Yao, R., Zhao, D.D., Zhang, S., Zhou, L.Q., Wang, X., Gao, C.F., Wu, S.F., 2017. Monitoring and mechanisms of insecticide resistance in *Chilo*

suppressalis (Lepidoptera: Crambidae), with special reference to diamides. Pest Manag. Sci. 73, 1169-1178.

Yin, G.-L., Yang, J.-S., Cao, J.-X., Yang, W.-J., 2006. Molecular cloning and characterization of FGLamide allatostatin gene from the prawn,

Macrobrachium rosenbergii. Peptides 27, 1241-1250.

Yin, J.-L., Xu, H.-W., Wu, J.-C., Hu, J.-H., Yang, G.-Q., 2014. Cultivar and insecticide applications affect the physiological development of the

brown planthopper, *Nilaparvata lugens* (Stål) (Hemiptera: Delphacidae). Environ Entomol. 37, 206-212.

Zeng, B., Chen, F.-R., Liu, Y.-T., Di, G., Zhang, Y.-J., Feng, Z.-R., Wang, L.-X., Vontas, J., Wu, S.-F., Zhu, K.Y., Gao, C.-F., 2023. A chitin synthase

mutation confers widespread resistance to buprofezin, a chitin synthesis inhibitor, in the brown planthopper, *Nilaparvata lugens*. J.

Pest Sci. 96, 819–832.

Zhang, C., Daubnerova, I., Jang, Y.-H., Kondo, S., Žitňan, D., Kim, Y.-J., 2021. The neuropeptide allatostatin C from clock-associated DN1p

neurons generates the circadian rhythm for oogenesis. Proc Natl Acad Sci U S A. 118, e2016878118.

Zhang, C., Kim, A.J., Rivera-Perez, C., Noriega, F.G., Kim, Y.-J., 2022a. The insect somatostatin pathway gates vitellogenesis progression during

reproductive maturation and the post-mating response. Nat Commun. 13, 969.

Zhang, Y.-X., Zhu, Z.-F., Lu, X.-L., Li, X., Ge, L.-Q., Fang, J.-C., Wu, J.-C., 2014. Effects of two pesticides, TZP and JGM, on reproduction of three

planthopper species, *Nilaparvata lugens* Stål, *Sogatella furcifera* Horvath, and *Laodelphax striatella* Fallén. Pestic. Biochem. Physiol. 115,

53-57.

Zhang, Y., Ruan, Y., Gong, C., Zhang, S., Zhang, J., He, Y., Wang, Q., Liu, D., Pu, J., Liu, X., Jiang, C., Wang, X., 2022b. Reproductive outbreaks of

Sogatella furcifera mediated by overexpression of the nuclear receptor USP under pressure from triflumezopyrim. Int J Mol Sci. 23,

13769.

Zhang, Y., Wang, X., Yang, B., Hu, Y., Huang, L., Bass, C., Liu, Z., 2015. Reduction in mRNA and protein expression of a nicotinic acetylcholine

receptor $\alpha 8$ subunit is associated with resistance to imidacloprid in the brown planthopper, Nilaparvata lugens. Journal of

Neurochemistry 135, 686-694.

Zhao, K.-F., Shi, Z.-P., Wu, J.-C., 2011. Insecticide-induced enhancement of flight capacity of the brown planthopper *Nilaparvata lugens* Stål

(Hemiptera: Delphacidae). Crop Protection 30, 476-482.

Zheng, H., Wang, N., Yun, J., Xu, H., Yang, J., Zhou, S., 2022. Juvenile hormone promotes paracellular transport of yolk proteins via remodeling

zonula adherens at tricellular junctions in the follicular epithelium. PLoS Genet. 18, e1010292.

Zhu, G.-H., Jiao, Y., Chereddy, S.C.R.R., Noh, M.Y., Palli, S.R., 2019. Knockout of juvenile hormone receptor, Methoprene-tolerant, induces black

larval phenotype in the yellow fever mosquito, *Aedes aegypti*. Proc Natl Acad Sci U S A., 201905729.

1251

1252