Positive selection and relaxed purifying selection contribute to rapid evolution of

male-biased genes in a dioecious flowering plant

Lei Zhao[1#], Wei Zhou[1#], Jun He[1], De-Zhu Li[1, 2*], Hong-Tao Li[1, 2*]

[1]Germplasm Bank of Wild Species, Yunnan Key Laboratory of Crop Wild Relatives

Omics, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming,

Yunnan 650201, China

[2]Kunming College of Life Science, University of Chinese Academy of Sciences,

Kunming, Yunnan 650201, China

[#]These authors contributed equally to this article

Running title: Positive selection and relaxed selection driving rapid evolution of

male-biased genes

* Corresponding authors:

Hong-Tao Li lihongtao@mail.kib.ac.cn;

De-Zhu Li dzl@mail.kib.ac.cn

Abstract

Sex-biased genes offer insights into the evolution of sexual dimorphism. Sex-biased genes, especially those with male bias, show elevated evolutionary rates of protein sequences driving by positive selection and relaxed purifying selection in animals. Although rapid sequence evolution of sex-biased genes and evolutionary forces have been investigated in animals and brown algas, less is known about them in dioecious angiosperms. In this study, we separately compared the expression of sex-biased gene between female and male flower buds and between female and male mature flowers in dioecious *Trichosanthes pilosa* (Cucurbitaceae). In floral buds, sex-biased gene expression was pervasive, and had significantly different roles on sexual dimorphism such as physiology. We observed higher rates of sequence evolution for male-biased genes in floral buds compared to female-biased and unbiased genes. Male-biased genes under positive selection were mainly related to abiotic and biotic stress, suggesting that high evolutionary rates are driven by adaptive evolution. Additionally, relaxed purifying selection may contribute to accelerated evolution in male-biased genes generated by gene duplication. Our findings, for the first time in dioecious angiosperms, suggest evident rapid evolution of male-biased genes, advance our understanding of the patterns and forces driving the evolution of sexual dimorphism in dioecious plants.

Introduction

Sexual dimorphism presents different morphological, ecological and
physiological traits in gonochoristic animals and dioecious plants, despite male and
female individuals sharing the same genome except for sex chromosomes (or
sex-determining loci) (Mank, 2009; Barrett and Hough, 2013). Such sexual
dimorphisms usually arise from differential expression of genes in both sexes, i.e.,
sex-biased genes (including sex-specific genes exclusively in one sex) that are located
on autosomal chromosomes and sex chromosomes/or sex-determining regions
(Ellegren and Parsch, 2007; Parsch and Ellegren, 2013; Charlesworth, 2018). Recently,
some studies have begun to explore the strength and impact of evolutionary forces
that shape different sexually dimorphic traits through sex-biased gene expression
(Mank, 2009; Rowe et al., 2018). Previous studies revealed that sex-biased gene
expressions have facilitated the evolution of sexual dimorphisms in many animals.
However, the extent to which this occurs varies greatly among taxa, tissues, and
developmental stages (Mank, 2017; Khodursky et al., 2020; Hsu et al., 2020; Lichilin
et al., 2021; Toubiana et al., 2021; Djordjevic et al., 2022). Unlike most animals, the
vast majority (~90%) of angiosperms are hermaphroditic plants, while only a small
fraction (~5%) are dioecious plants in which individuals have exclusively male or
female reproductive organs (Renner, 2014). Most dioecious plants do not possess
heteromorphic sex chromosomes. Furthermore, sexual dimorphism in dioecious plants
is less common and less conspicuous than that in almost animals (Barrett and Hough,

2013). Hence, the study of sex-biased gene expression is of great interest to plant

evolutionary biologists, as it is essential to understand the evolution of sexual

dimorphism in dioecious plants.

A common pattern that has emerged from previous studies is that sex-biased

genes, particularly male-biased genes, tend to evolve rapidly in protein sequence (the

ratio of non-synonymous to synonymous substitutions, d_N/d_S) compared to unbiased

genes (Ellegren and Parsch, 2007; Grath and Parsch, 2016). The rapid evolution of

male-biased genes was first observed in *Drosophila melanogaster* (Zhang et al., 2004;

Zhang and Parsch, 2005) and has been supported by recent investigations in a wider

range of animals (Pröschel et al., 2006; Mank et al., 2007; Mank, 2017; Papa et al.,

2017; Catalan et al., 2018; Toubiana et al., 2021). In recent years, there have been

growing studies on the expression dynamics and molecular evolutionary rates of

sex-biased genes in flowering plants, including *Silene latifolia* (Zemp et al., 2016),

Salix viminalis (Darolti et al., 2018), *Mercurialis annua* (Cossard et al., 2019),

Populus balsamifera (Sanderson et al., 2019), and *Leucadendron* plants (Scharmann

et al., 2021). However, despite such advances, the molecular evolution pattern of

sex-biased genes in plants remains inconsistent among the studied plant species

(Muyle, 2019; Veltsos, 2019). In dioecious plants such as *Mercurialis annua* and

Leucadendron, Cossard et al., (2019) and Scharmann et al., (2021) found no

significant differences in evolutionary rates of proteins among female-biased,

male-biased and unbiased genes, although the expression of sex-biased genes was

highly different from unbiased genes in leaves. Similar patterns have also been

reported in dioecious *Populus balsamifera*, where evolutionary rates of male-biased,

female-biased and unbiased genes did not differ in reproductive tissues (Sanderson et

al., 2019). However, in the dioecious plant *Salix viminalis*, male-biased genes have

significantly lower evolutionary rates of proteins than female-biased and unbiased

genes in catkin tissues (Darolti et al., 2018). To our knowledge, only five studies have

investigated the expression differences and protein evolutionary rates of sex-biased

genes in dioecious angiosperms. Moreover, these studies only compared gene

expression in vegetative vs. vegetative tissues and vegetative vs. reproductive tissues.

Therefore, more studies and taxa are needed to explore the common patterns of

sequence evolution in sex-biased genes, with more focus on comparing sex-biased

gene expression in reproductive vs. reproductive tissues, e.g., different floral

development stages in dioecious angiosperms.

Evolutionary dynamic analyses indicate that different evolutionary forces impact

the rate of sequence evolution of sex-biased genes. These forces include positive

selection, which promotes the spread and adaptive fixation of beneficial alleles;

sexual selection, which results from male-male competition or female choice; and

relaxed purifying selection, which reduces the removal of deleterious mutations

(Grath and Parsch, 2016; Mank, 2017; Dapper and Wade, 2020). For example, in

animal systems, particularly in *Drosophia*, the elevated sequence divergence rates of

male-biased genes have often been interpreted as the signature of positive selection

due to adaptive evolution, suggesting that sexual selection is the primary evolutionary

forces (Pro□schel et al., 2006; Assis et al., 2012). However, studies in plants have

rarely reported elevated rates of sex-biased genes, except for brown alga, in which

female-biased and/or male-biased genes exhibiting higher evolutionary rates than

unbiased genes suggests that rapid evolution is partly driven by adaptive evolution or

sexual selection (Lipinska et al., 2015; Cossard et al., 2022; Hatchett et al., 2023).

114   An alternative explanation for the rapid evolution of sex-biased genes is a

relaxation of purifying selection due to reduced constraints (Lahti et al., 2009; Dapper

and Wade, 2020). In the model plant *Arabidopsis thaliana*, pollen genes were found to

be evolving faster than sporophyte-specific genes due to relaxed purifying selection

associated with the transition from outcrossing to selfing (Harrison et al., 2019).

These trends were recently confirmed in *Arabis alpina*, which exhibits mating system

variation across its distribution, suggesting that the efficacy of purifying selection on

male gametophyte-expressed genes was significantly weaker in inbred populations

(Gutierrez-Valencia et al., 2022). Together, these findings in plants reinforce the idea

that both adaptive (e.g., positive selection, sexual selection) and non-adaptive (e.g.,

relaxed selection) evolutionary processes differentially impact the sequence evolution

of sex-biased genes. Hence, investigating the potential contribution of selection forces

to the emergence of specific evolutionary patterns of sex-biased genes within a focal

species is of great interest.

128   In the family Cucurbitaceae, there are about 96 genera and 1,000 species, about

50% of species are dioecious, and 50% are monoecious (Schaefer and Renner, 2011).

Phylogenetic analyses of Cucurbitaceae suggest that dioecy is the ancestral state of

the family, but transitions frequently to monoecy (Zhang et al., 2006). *Trichosanthes*

pilosa (2n = 22, Cucurbitaceae) is mainly distributed from Southwest, Southeast

China to Japan, New Guinea and Western Australia. It was suggested to have

originated in the late Miocene (c.13 Ma) (de Boer et al., 2012; 2015). *Trichosanthes*

pilosa (synonym: *T. ovigera*) is a perennial dioecious vine that reproduces sexually

and possesses a pair of heteromorphic sex chromosomes XX/XY (Ming et al., 2011).

The species exhibits strong sexual dimorphism in the floral morphological structures

and the life-history traits, such as racemose male flower and solitary female flower

(Fig. 1), early-flowering, caducous male flower and late-flowering, long-lived female

flower (Wu et al., 2011).

To understand the evolution of sex-biased genes in dioecious *Trichosanthes*

pilosa, we collected floral buds and mature flowers tissues from male and female

individuals and characterized their expression profiles using Illumina RNA

sequencing. Our primary objectives are to 1) compare expression divergences

between males and females at two floral developmental stages; 2) explore whether

there are differences in the evolutionary rates of proteins among female-biased,

male-biased and unbiased genes; and if so, 3) determine the main selective forces that

contribute to the differentiation of sequence evolution rates among gene categories.

Results

Transcriptome sequencing, *de novo* assembly and annotation

Using whole transcriptome shotgun sequencing, we sequenced 12 floral buds and

mature flower tissues from females and males of dioecious *T. pilosa* and generated a

total of nearly 276 million clean reads (Table S1). We then performed *de novo*

assembly of transcripts from all the clean reads, followed by clustering and filtering

analysis, resulting in 59,051 unigenes (Fig. S1A). To evaluate the quality of the

assembled unigenes, we compared them against protein databases such as NR, KEGG,

Swissport, PFAM, and GO using BLASTP and nucleotide database NT using

BLASTN (Table S2). The e-value distribution of the best hits in NR database

suggested that 47,241 unigenes (80%) had strong homology, with an e-value smaller

than 1.0e-15 (Fig. S1B). The majority of unigenes were annotated by homologs in

other species of Cucurbitaceae, such as *Momordica charantia*, *Cucumis melo*,

Cucurbita pepo, *Cucurbita moschata* and *Cucurbita maxima* (Fig. S1C). Furthermore,

we used BUSCO assessments based on embryophyta_odb9 database, which showed

the completeness of the reference transcriptome at 78.8% (Table S2). Overall, our

analyses suggested that we have generated high-quality reference transcriptomes.

Expression characteristics of sex-biased genes

We mapped the RNA-Seq reads of floral buds and mature flowers onto the

reference transcriptome in dioecious *T. pilosa*, which resulted in approximately 75%

read mappings per sample (Table S3). In floral buds, we identified 5,096 (9.50%)

female-biased genes and 4,214 (7.86%) male-biased genes (Fig. 2A). In contrast, only

380 (0.70%) female-biased genes and 233 (0.43%) male-biased genes were detected

in mature flowers (Fig. 2B). Using hierarchical clustering analysis, we evaluated

different levels of gene expression across sexes and tissues, revealing strong sexual

dimorphism (Fig. 2C). Furthermore, we observed that the number of sex-biased genes

in floral buds was approximately 15 times higher than in mature flowers, indicating

that sex-biased genes associated with sex differentiation and sexually dimorphic traits

are predominantly expressed in floral buds. We also analyzed sex-specific genes that

were exclusively expressed in floral buds and mature flowers of one sex. In floral

buds, we found 253 out of 5,096 (4.96%) female-specific genes and 465 out of 4,214

(11.03%) male-specific genes. However, in mature flowers, we only identified 26 out

of 380 (6.84%) female-specific genes and 52 out 233 (22.32%) male-specific genes.

Tissue-biased gene expression

We compared the expression levels of transcripts in floral bud tissues and mature

flower tissues within each sex to identify genes with tissue-biased expression. In male

plant, the number of tissue-biased genes in mature flowers was 1,040 higher than that

in floral buds (Fig. 3A and 3B). However, in female plant, the number of tissue-biased

genes in mature flowers was only 536 more than that in floral buds (Fig. 3C and 3D).

Our results indicated that males had a higher tissue-bias relative to females. We also

identified sex-biased genes that were expressed in both types of tissues by comparing

tissue-biased genes with male-biased and female-biased genes, respectively. Few of

female-biased genes overlapped with tissue-biased genes, accounting for only 85 out

of 5,096 (1.67%) in floral buds (Fig. 3C) and 5 out of 380 (1.32%) in mature flowers

(Fig. 3D). However, a significant proportion of male-biased genes overlapped with

tissue-biased genes, with 1,010 out of 4,214 (23.97%) in floral buds (Fig. 3A) and 145

out of 233 (62.23%) in mature flowers (Fig. 3B).

Elevated protein evolutionary rates of male-biased genes in floral buds

We identified 1,145 female-biased, 343 male-biased, and 2,378 unbiased

one-to-one orthologous groups (OGs) from floral buds in three additional species of

Trichanthes, *T. anguina*, *T. pilosa*, *T. kirilowii*, together with *Luffa cylindrica*.

Additionally, we detected 45 female-biased, 13 male-biased, and 3,782 unbiased

one-to-one OGs from mature flowers in all four species. To quantify the rates of

protein sequences, we separately calculated ω values for each sex-biased and unbiased

orthologous gene using two branch models in floral buds and mature flowers (Figs. 4

and S2).

In the results of 'two-ratio' branch model, the median of ω values in

female-biased, male-biased and unbiased genes were 0.227, 0.257 and 0.230 in floral

buds, respectively (Fig. 4A and Table S4). We observed that male-biased genes had a

13.22% and 11.74% higher median than female-biased and unbiased genes in floral

buds, respectively. The difference in the distribution of ω values between

female-biased versus male-biased genes ($P = 0.0021$) and male-biased versus

unbiased genes ($P = 0.0051$) was statistically significant in Wilcoxon rank sum tests.

However, we did not find a significant difference in ω values between female-biased

and unbiased genes in floral buds (Wilcoxon rank sum test, $P = 0.4618$). In mature

flowers, the median of ω values for female-biased, male-biased, and unbiased genes

were 0.269, 0.177 and 0.231, respectively (Fig. 4B and Table S4). However, we found

that there was no statistically significant difference in the distribution of ω values

using Wilcoxon rank sum tests for female-biased versus male-biased genes ($P =$

0.0556), female-biased versus unbiased genes ($P = 0.0796$), and male-biased versus

unbiased genes ($P = 0.3296$).

The 'free-ratio' branch model yielded interesting results. In floral buds, the

median ω values for female-biased, male-biased, and unbiased genes were 0.222,

0.265 and 0.226, respectively (Fig. 4C and Table S5). Male-biased genes had a

significantly higher median relative to female-biased genes (19.37% higher, Wilcoxon

rank sum test, $P = 0.0009$) and unbiased genes (17.26% higher, Wilcoxon rank sum

test, $P = 0.0004$) in floral buds. However, there was no significant difference in ω

values between female-biased and unbiased genes (Wilcoxon rank sum test, $P =$

0.9862). In mature flowers, the median ω values for female-biased, male-biased, and

unbiased genes were 0.300, 0.148 and 0.227, respectively (Fig. 4D and Table S5).

Female-biased and unbiased genes had significantly higher ω values than male-biased

(Wilcoxon rank sum test, $P = 0.0101$, $P = 0.0146$, respectively). However, there was

no significant difference in ω values between female-biased and unbiased genes

(Wilcoxon rank sum test, $P = 0.2887$). Since the number and evolutionary rates of

male-biased genes in mature flowers are lower than those in flora buds, we decided to

focus on the latter in subsequent analyses.

Evidences and contributions of positive selection and relaxed selection for

male-biased genes in floral buds

After comparing the A model and null model, we discovered that 39 out of 343

OGs (11.34%) exhibited strong evidence of having certain sites that evolved under

positive selection based on foreground 2b ω value, likelihood ratio tests (LRTs, P <

0.05) and Bayes empirical Bayes (BEB) value (Fig. 5 and Table S6). As an additional

approach, we utilized the aBSREL and BUSTED methods that were implemented in

HyPhy v.2.5 software, and detected significant evidence of positive selection.

According to our findings, 84 out of 343 OGs (24.49%) were identified to be under

episodic positive selection in male-biased genes of floral buds with a site proportion

of 0.17%–26.44% based on aBSREL (Table S7). In addition, 69 out of 343 OGs

(20.01%) exhibited significant signs of positive selection with the site proportion of

0.28%–32.65% in male-biased genes of floral buds according to BUSTED (Table S8).

Among these, a total of 32 OGs (9.30%) were identified through our tests using

CodeML, aBSREL and BUSTED (Fig. 5).

Relaxed selection may occur when the efficiency of natural selection is

dramatically reduced or eliminated at both gene-wide and genome-wide levels. This

has been proposed as an explaination for the rapid evolution of sex-biased genes

(Lahti et al., 2009; Mank, 2017). Using the RELAX model, we conducted an analysis

and found that 18 out of 343 OGs (5.23%) showed significant evidence of relaxed

selection (K = 0.0184–0.6497) (Table S9). Interestingly, we also observed that 61 out

of 343 OGs (17.73%) exhibited significant evidence of intensified positive selection

(K = 2.3363–50, $\omega_2 \geq 1$) (Fig. 5 and Table S10), which is consistent with the results

obtained from CodeML, aBSREL and BUSTED.

According to previous studies (Ellegren and Parsch, 2007; Catalan et al., 2018),

genes that exhibit sex-biased expression with rapid evolutionary rates tend to display

a lower codon bias usage compared to unbiased genes. In our results, we found that

male-biased genes in floral buds had a significantly lower median effective number of

codons (ENCs) than both female-biased and unbiased genes (Wilcoxon rank sum test,

female-biased vs male-biased genes, $P = 0.0001$ and male-biased vs unbiased genes,

$P = 0.0123$). This suggested that male-biased genes in floral buds exhibit a stronger

codon bias usage than both female-biased and unbiased genes (Fig. S3). In summary,

our analyses indicated that rapid evolutionary rates of male-biased genes in floral

buds were not associated with a reduction in codon usage bias.

To confirm the contributions of positive selection and relaxed selection to rapid

rates of male-biased genes in floral buds, we generated three datasets of OGs by

excluding different sets of genes. Specifically, we excluded 18 relaxed selective

male-biased genes (5.23%), 98 positively selected male-biased genes (28.57%), and

112 male-biased genes (32.65%) under positive and relaxed selection from 343 OGs

(Fig. S4). We observed that after excluding relaxed selective male-biased genes, the

median (0.264) decreased by 0.34% compared to the median (0.265) of all OGs (Fig.

S4A-B). However, after excluding positively selected male-biased genes, the median

(0.236) reduced by 11% (Fig. S4A, C) in the results of 'free-ratio' branch model. This

pattern was consistene in the results of 'two-ratio' branch model as well (Fig. S4E-G).

Additionally, we analyzed female-biased and unbiased genes that underwent positive

and relaxed selection in floral buds (Tables S6-S10). We identified 216 (18.86%)

positively selected (Fig. S5A), and 69 (6.03%) relaxed selective female-biased genes

from 1,145 OGs (Fig. S5B), respectively. Similarly, we found 436 (18.33%)

positively selected (Fig. S6A), and 43 (1.81%) unbiased genes under relaxed selection

from 2,378 OGs (Fig. S6B), respectively. Notably, male-biased genes have a higher

proportion (10%) of positively selected genes compared to female-biased and

unbiased genes. However, relaxed selective male-biased genes have a higher

proportion (3.24%) than unbiased genes, but about 0.8% lower than that of

female-biased genes. In summary, our analyses suggested that positive selection and

relaxed selection likely drove the rapid evolutionary rates of male-biased genes

compared to female-biased and unbiased genes in floral buds, but positive selection

had an even greater impact.

Functional analysis of sex-biased genes in floral buds

We conducted KEGG pathway enrichment analysis on sex-biased genes in floral

buds. Our results showed that 699 genes were female-biased and 358 genes were

male-biased, with significant enrichment ($P < 0.05$) in 26 and 24 KEGG pathways,

respectively (Table S11). In the floral bud stage, we observed that female-biased

genes were mainly enriched in metabolic and signaling pathways, such as ribosome,

Fatty acid elongation, photosynthesis and plant hormone signal transduction (Fig. S7A

and Table S11). On the other hand, male-biased genes were significantly enriched in

metabolic and signaling pathways, including Inositol phosphate metabolism, starch

and sucrose metabolism, regulation of autophagy, plant hormone signal transduction,

and Toll-like receptor signaling pathway (Fig. S7B and Table S11).

We have also found that certain male-biased genes, which are evolving under

positive selection and relaxed selection (Table S12 and S13), were related to abiotic

and biotic stress. For instance, mitogen-activated protein kinase kinase kinase 18

(MAPKKK18) (Zhang and Zhang 2022,), zinc finger CCCH domain-containing

protein 20 (C3H20/TZF2) (Bogamuwa and Jang, 2014), and heat stress transcription

factor B-3 (HSFB3) (Scharf et al., 2012) have been linked to stress. Interestingly, ten

male-biased genes with rapid evolutionary rates were associated with anther and

pollen development. These genes include LRR receptor-like serine/threonine protein

kinase (LRR-RLK) (Cui et al., 2022), pollen receptor-like kinase 3 (*PRK3*)

(Muschietti and Wengier, 2018), autophagy-related protein 18f (*ATG18f*) (Zhou et al.,

2015; Li et al., 2020a), and plant homeodomain (PHD) finger protein 3 (*MALE*

STERILITY 3) (Hou et al., 2022) in floral buds of male plants.

Discussion

The Cucurbitaceae family is an excellent model for studying the evolution of

sexual systems of angiosperms, including sex-determination mechanism and sexual

dimorphism. In this study, we aimed to investigate whether sex-biased genes

exhibited the evidence of rapid evolutionary rates of protein sequences and identify

the evolutionary forces responsible for the observed patterns in the dioecious *T. pilosa*.

We compared the expression profiles of sex-biased gene between sexes and two tissue

types and examined the signatures of rapid sequence evolution for sex-biased genes,

as well as the contributions of potential evolutionary forces.

Sex-biased expression in floral buds

Numerous studies have shown that in dioecious plants, male-biased genes tend to

outnumber female-biased genes. For instance, insect-pollinated dioecious plants such

as *Asparagus officinalis* (Harkess et al., 2015) and *Silene latifolia* (Zemp et al., 2016),

exhibit a higher proportion of male-biased genes, whereas wind-pollinated dioecious

plants like *Populus balsamifera* (Sanderson et al., 2019) has twice as many

female-biased genes than male-biased genes. These differences in these studies could

be partly attributed to the impact of sexual selection on secondary sexual traits in

insect-pollinated dioecious plants, as opposed to wind-pollinated ones (Delph and

Herlihy, 2012; Muyle, 2019; Sanderson et al., 2019). In contrast to the above studies,

our findings revealed that the number of female-biased genes in floral buds of the

night-flowering, insect-pollinated dioecious plant *Trichosanthes pilosa* exceeded that

of male-biased genes by 882. This excess of female-biased expression could be due to

lower energy consumption needs and reduced chemical defense capability against

herbivores in short-lived male flowers (Sanderson et al., 2019). Indeed, functional

enrichment analysis in chemical defense pathways such as terpenoid backbone and

diterpenoid biosynthesis, indicated that female floral buds had more expressed genes

and were better equipped to defend against herbivores compared to male floral buds

(Fig. S7A and Table S11). Additionally, our enrichment analysis showed that the

photosynthesis, porphyrin and chlorophyll metabolism pathways were more active in

female floral buds compared to male floral buds (Fig. S7A and Table S11), enabling

them to acquire more resources such as carbon for fruit and seed production

(Delph,1999).

We identified functional enrichments in Toll-like receptor signaling, NF-kappa B

signaling, and Inositol phosphate metabolism pathways in male floral buds (Fig. S7B,

Tables S11 and S12). These findings suggest that male floral buds are better adapted

to biotic and abiotic environments. Moreover, the enrichment in regulation of

autophagy pathways could be associated with the senescence for male floral buds

(Table S14) (Liu and Bassham, 2012; Li et al., 2020). We also observed that

homologous genes of two male-biased genes (Table S14) that control the raceme

inflorescence development (Teo et al., 2014) were highly expressed in floral buds.

This suggests that sex-biased genes, rather than sex-specific genes, are responsible for

dimorphic traits of inflorescence. In short, these results indicate that changes in

sex-biased genes expression play different roles on sexual dimorphism in

physiological and morphological traits (Dawson and Geber, 1999).

Rapid Evolution of male-biased genes in floral buds

It has been observed that, in most animals, sex-biased genes, particularly those

biased towards males, often exhibit more rapid evolutionary rates than unbiased genes

(Parsch and Ellegren, 2013; Grath and Parsch, 2016; Mank, 2017; Toubiana et al.,

2021). However, in dioecious angiosperms, no evidence of rapid evolution in

sex-biased genes relative to unbiased genes has been found (Zemp et al., 2016; Darolti

et al., 2018; Cossard et al., 2019; Sanderson et al., 2019; Scharmann et al., 2021). In

contrast, our findings indicated that male-biased genes exhibited rapid evolutionary

rates than both female-biased and unbiased genes in floral buds of dioecious *T. pilosa.*

We proposed that positive selection and relaxed purifying selection may be

responsible for the rapid sequence evolution of male-biased genes.

After analyzing the data, we found that around 28.57% (98 genes) of male-biased

genes have undergone positive selection. Additionally, we observed that the

proportion of male-biased genes under positive selection was about 10% higher than

that of female-biased and unbiased genes. Furthermore, we discovered that some

male-biased genes under positive selection were linked to abiotic and biotic stress

responses (Table S12). Our findings are consistent with studies on *Drosophila* and

Ectocarpus (Zhang and Parsch, 2005; Lipinska et al., 2015), suggesting that adaptive

evolution is one of the primary evolutionary forces for rapid evolutionary rates.

Notably, we identified several male-biased genes under positive selection that are

functionally related to early flowering (*phyB*) (Stephenson and Bertin, 1983; Forrest,

2014; Hajdu et al., 2015) and pollen development (Skogsmyr and Lankinen, 2002;

Williams and Reese, 2019) (Tables S12- S14). These findings indicate that a small

fraction of male-biased genes may experience adaptive evolution due to sexual

selection, driven by male-male competition.

Alternatively, relaxed constraints could contribute to the rapid evolutionary rates

of sex-biased genes through three key characteristics (Dapper and Wade, 2020; Tosto

et al., 2023). First, sex-biased genes are often expressed solely in reproductive tissues

of one sex (e.g., sex-specific genes), particularly in the haploid phase (Immler, 2019;

Beaudry et al., 2020). Sex-specific selection (e.g., relaxed purifying selection) acting

on sex-specific genes could decrease the elimination of deleterious mutations (Mank,

2017), such as pollen-specific (Harrison et al., 2019) or testes-specific genes

(Gershoni and Pietrokovski, 2014). However, our results did not reveal male-specific

genes that underwent relaxed purifying selection. Second, sex-biased genes are often

expressed in few tissues (tissue-specific genes) (Meisel, 2011; Tosto et al., 2023).

These genes could evolve under positive selection or relaxed purifying selection due

to low pleiotropy (Congrains et al., 2018; Whittle et al., 2021; Tosto et al., 2023). In

our results, 343 male-biased genes with faster evolutionary rates relative to

female-biased and unbiased genes overlapped with tissue-biased genes in floral buds

(27 out of 343, 7.87%) (Fig. S8A). Furthermore, 27 out of 343 male-biased genes

(that is, tissue-biased genes) in floral buds overlapped with nine out of 98 (9.18%)

male-biased genes under positive selection (Fig. S8B), and one out of 18 (5.56%)

male-biased genes under relaxed purifying selection (Fig. S8C). Consistent with

male-biased genes in *Fucus* (Hatchett et al., 2023), our analyses suggested that

elevated evolutionary rates may partly be linked to low pleiotropy. Finally, gene

duplication has long been thought to promote functional divergences and phenotypic

novelties by relaxing the constraints of purifying selection on the duplicated gene

copy early in its history (Lynch and Conery, 2000; Lynch and Katju, 2004; Lahti et al.,

2009). For instance, the progesterone receptor gene family in the human lineage

(Marinić and Lynch, 2020) and the CYP98A9 clade in Brassicales (Liu et al., 2016)

have demonstrated rapid evolution and divergent function due to relaxed purifying

selection. In our results, we identified 18 out of 343 (5.25%) male-biased genes that

underwent relaxed purifying selection using RELAX model (Table S13). Interestingly,

the vast majority of genes under relaxed selection were members of different gene

families generated by gene duplication (including whole-genome duplication), such as

LOB domain-containing protein 18 (*LBD18*) (Zhang et al., 2020), WRKY

transcription factor 72 (*WRKY72*) (Chen et al., 2017), and pollen receptor-like kinase

3 (*PRK3*) (Muschietti and Wengier, 2018).

Additionally, although reducing codon usage bias could theoretically accelerate

evolutionary rates of sex-biased genes by decreasing synonymous substitution rates,

our results did not support this idea. In fact, we found that male-biased genes had

significantly higher median d_S values than female-biased and unbiased genes, both in

the 'free-ratio' analysis (Fig. S9A, female-biased vs male-biased genes, $P = 6.444e-12$

and male-biased vs unbiased genes, $P = 4.564e-13$) and 'two-ratio' branch model (Fig.

S9B, female-biased vs. male-biased genes, $P = 2.2e-16$ and male-biased vs. unbiased

genes, $P = 9.421e-08$, respectively), indicating higher synonymous substitution rates.

Several species have been observed to exhibit rapid evolutionary rates of

sequences on sex chromosome compared to autosomes, which has been related to the

evolutionary theories of fast-X or fast-Z (Meisel and Connallon, 2013; Wright et al.,

2015; Charlesworth et al., 2018). Furthermore, the quantification of gene expression

by bulk RNA-seq technology, relative to single-cell transcriptome analysis, has been

shown to potentially obfuscating true signals in the evolution of sex-biased gene

expression in complex aggregations of diverse cell types (Darolti and Mank, 2023;

Tosto et al., 2023). However, investigation of these interesting issues related to

sex-biased gene evolution in *T. pilosa* can only be conducted when whole genome

sequences become available in the near future.

Methods

Plant materials and RNA isolation

Floral buds (≤ 3 mm) and mature flowers were sampled from three female and

three male plants (Fig. 1) in the field of Qinglong Gorge of Anning, Yunnan Province.

Floral buds from female and male plants were named F1 and M1, respectively.

Similarly, mature flowers from female and male plants, there were named F2 and M2,

respectively (Table S1). To exclude possible bacterial contamination, all tissues were

sterilized with 75% alcohol and immediately rinsed with purified water. All samples

were then snap-frozen in liquid nitrogen, and stored at -80˚C. Total RNA was

extracted from each sample using TRIzol reagent (Life Technologies, CA, USA)

according to the manufacturer's instructions. The quantification and qualification of

RNA were assessed by the RNA Nano 6000 Assay Kit of the Bioanalyzer 2100

system (Agilent Technologies, CA, USA).

Illumina sequencing, *de novo* assembly and annotation

To construct the library, approximately 2 µg of total RNA was used with the

Illumina NEBNext UltraTM RNA Library Prep Kit. RNA sequencing was performed

on the Illumina NovaSeq 6000, generating 150 bp paired-end reads. The resulting

clean reads were obtained by removing adapters, reads containing N bases and

low-quality reads using Trimmomatic v.0.39 (Bolger et al., 2014). These reads were

deposited in the NCBI database (PRJNA899312).

De novo assembly for clean reads from all samples was performed using Trinity

v.2.10.0 (Haas et al., 2013) with min_kmer_cov: 3 and all other default parameters. To

eliminate contamination, all transcripts of *de novo* assembly were compared to

bacterial genomes downloaded from NCBI databases using BLASTN with e-value of

1.0e-05 in blast+ 2.12.0 software. We used Corset v.4.6 (Davidson and Oshlack, 2014)

to obtain high quality, non-redundant consensus transcripts (unigenes). TransDecoder

v.5.5.0 was run with -m 100 parameters, namely at least 100 amino acids, to predict

the coding DNA and protein sequences (Haas et al., 2013).

To evaluate the accuracy and completeness of reference transcriptomes, we

performed gene function annotations based on the following databases, using BLAST

with a cutoff e-value of 1.0e-05: NR, NT and Swissport (Shiryev et al., 2007). We

mapped the unigenes to Pfam database using InterProScan v.5.41 (Jones et al., 2014),

to the GO database using Blast2GO (Conesa et al., 2005), and to the KEGG database

using KEGG automatic annotation server (Moriya et al., 2007). Additionally, we

estimated the completeness of reference transcriptomes using BUSCO v.5.2.2 based

on embryophyta_odb9 database (Seppey et al., 2019).

Detection of sex-biased genes

Clean reads were mapped onto all unigenes using Bowtie2 (Langmead and

Salzberg, 2012). Read counts were normalized to FPKM (Fragments Per Kilobase

Million) value for each unigene using RSEM (Li and Dewey, 2011) in different male

and female samples. Genes with zero read counts (i.e., no expression) in both two

sexes and tissues were excluded. Differential expression analysis between sexes and

tissue types was performed using DESeq2 R package. Unigenes with an

FDR-adjusted $P < 0.05$ and an absolute value of $\log_2$ ratio ≥ 1 identified by DESeq2

were considered as sex-biased genes. To perform KEGG functional enrichment, we

use all KEGG annotation terms for all genes as the background and performed the

analyses using KOBAS v.2.0.12 (Mao et al., 2005).

Evolutionary rate analyses

To quantify the evolutionary rates of sex-biased genes, we download published

genome datasets for monoecious *Trichosanthes anguina* (Ma et al., 2020) and

monoecious *Luffa cylindrica* which has a closer phylogenetic relationship with

Trichosanthes (de Boer et al., 2012; Wu et al., 2020) from CuGenDB database (Zheng

et al., 2019). Additionally, we also download published RNA sequencing reads of

floral buds and flowers from CNCB (Accession CRA002313) and NCBI databases

(Accession SRR5259239) for dioecious plant *Trichosanthes kirilowii* (Hu et al., 2020),

and *de novo* assembled by previously described methods.

We identified one-to-one OGs using OrthoFinder v.2.3.3 with default parameters

from *T. anguina*, *T. pilosa*, *T. kirilowii*, and *Luffa cylindrica* (Emms and Kelly, 2019).

Then, we employed TranslatorX with -c 1 -p M -g -b5 n parameters (i.e., the multiple

alignment and the trimming using Muscle and GBlocks, respectively), translated

nucleotide sequences and back-translated amino acid alignments into nucleotide

alignments to ensure codon-to-codon alignment (Abascal et al., 2010). The remaining

gapless alignments ($\geq$100 bp in length) were retained.

To investigate the evolutionary rates of code sequences, we estimated

nonsynonymous substitution (d_N), synonymous substitution (d_S) rates, as well as

protein substitution rates (d_N/d_S, ω), using two branch models from CodeML package

in PAML v.4.9h with the F3x4 codon frequencies (CodonFreq = 2) (Yang, 2007).

According to the phylogenetic relationships of *Trichosanthes* (de Boer et al., 2012;

Guo et al., 2020), we set up tree structure ((*T. anguina*, *T. pilosa*), *T. kirilowii*, *L.*

cylindrica) in the control file of CodeML. First, we employed a 'two-ratio' branch

model (model = 2, Nssites = 0) that assumes the foreground (two dioecious species)

has a different ω value from the background (two monoecious species) to estimate and

compare the divergences of the foreground. Second, to reduce the potential bias of ω

value due to the conflation of two dioecious species, we also implemented a

'free-ratio' branch model (model = 1, Nssites = 0), which assumes an independent ω

ratio for each branch. Finally, we excluded all OGs with $\omega > 2$, plotted the distribution

of ω values and compared the median of ω values in female-biased, male-biased and

unbiased orthologous genes of floral buds and mature flowers. All comparisons

between sex-biased and unbiased genes were tested using Wilcoxon rank sum test in

R software.

Estimation of the strength of natural selection

The rapid evolutionary rates of sex-biased genes may be attributed by positive

selection, relaxed selection and lower codon usage bias (Catalan et al., 2018; Dapper

and Wade, 2020). Therefore, we conducted separate analyses using classical

branch-site models, the adaptive branch-site random effects likelihood (aBSREL)

model, the branch-site unrestricted statistical test for episodic diversification

(BUSTED) model, the RELAX model, and the effective number of codons (ENC) in

PAML v.4.9h, HyPhy v.2.5 and CodonW v.1.4.2 to distinguish which evolutionary

forces are driving the rapid evolutionary rates of sex-biased genes (Pond et al., 2020).

To determine if amino acid sites in the foreground, including the *T. pilosa* lineage

have undergone positive selection (foreground 2b ω >1) compared with the

background for each OGs, we used branch-site model A (model = 2, Nssite = 2,

fix_omega = 0, omega = 1.5) and branch-site model null (model = 2, Nssite = 2,

fix_omega = 1, omega = 1). We examined the significance of likelihood ratio tests

(LRTs, $P < 0.05$) to identify positively selected sites between model A and model null

by comparing LRTs to the Chi-square distribution with two degrees of freedom. We

adjusted the LRTs P value for multiple comparisons using Benjamini and Hochberg's

(FDR) algorithm. When the *P* value was significant, we used Bayes Empirical Bayes

(BEB) estimates to identify sites with a high posterior probability (pp $\geq$ 0.95) of being

under positive selection (Yang et al., 2005; Zhang et al., 2005).

To detect episodic positive selection at a proportion of sites on the foreground

branch, we employed the aBSREL method in the HyPhy v.2.5 packages to compare

the fully adaptive model ($\omega > 1$) to the null model that allows no positive selection

rate classes by LRTs, which is an improved algorithm of branch-site models in PAML

(Smith et al., 2015). For relatively small datasets, such as those with fewer than 10

taxa, the aBSREL method may not have enough power to detect positive selection.

Therefore, we also ran the BUSTED method to identify gene-wide evidence of

episodic positive selection at least one site on at least one branch (Murrell et al., 2015).

We set *T. pilosa* as the foreground and assessed the statistical significance ($P < 0.05$)

using LRTs with the Holm-Bonferroni correction.

To test the relaxation of selective strength, we utilized the RELAX model in the

HyPhy v.2.5 software (Wertheim et al., 2015; Schrader et al., 2021). The RELAX

model estimates three ω parameters ($\omega_0 \leq \omega_1 \leq 1 \leq \omega_2$), and determines the proportion

of sites in the test (foreground) and reference (background) branches using a

branch-site model. The first two ω classifications indicate that sites have undergone

purifying selection, and the third classification indicates that sites have been under

positive selection. Additionally, the model introduces a selection intensity parameter

(K value) to compare a null model (K = 1) with an alternative model, thereby

assessing the the strength of natural selection. When K > 1, it suggests intensified

natural selection, when K < 1, indicates relaxed natural selection in the test branch

relative to the reference branch. We quantified the statistical confidence of K value (P

< 0.05) using LRTs and the Holm-Bonferroni correction.

To investigate codon usage bias, which refers to the differences in the frequency

of occurrence of synonymous codons in coding DNA, we employed CodonW v.1.4.2.

This program considers the ENC values from 20 to 61 as a measure of the departure

of the genetic codes for a given gene (Wright, 1990), with lower ENC values

represent stronger codon usage bias (Hambuch and Parsch, 2005). We performed a

Wilcoxon rank sum test to determine if there were deviations in ENC values among

female-biased, male-biased, and unbiased genes in floral buds.

Acknowledgements

We are indebated to Ting Zhang, Zhi-Yun Yang, Jiang-Li Ma, Peng-Fei Ma,

Xu-Kun Wu, Shi-Yu Lv and other members of staff of Germplasm Bank of Wild

Species for sampling. We are very grateful to Spencer C. H. Barrett (University of

Toronto, Canada) for his critical reading and suggestions on the manuscript. We also

thank to the iFlora HPC Center (iFlora High Performance Computing Center) of

Germplasm Bank of Wild Species for computationoal support on data analysis. This

study was funded by the Strategic Priority Research Program of the Chinese Academy

of Sciences, China (XDB31000000), the Key R and D Program of Yunnan Province,

China (202103AC100003), the Key Basic Research Program of Yunnan Province,

China (202101BC070003), the Science and Technology Basic Resources

Investigation Program of China (2019FY100900), the National Natural Science

Foundation of China (31570333), and the open research project of

"Cross-Cooperative Team" of the CAS' Germplasm Bank of Wild Species, Kunming

Institute of Botany. The study was also supported by National Wild Plant Germplasm

Resource Center.

Author contributions

HTL, DZL and LZ conceived the project the study. HTL and LZ designed and

performed the experiment. LZ, JH and HTL collected the samples. LZ analyzed the

data and wrote the manuscript. HTL, DZL, LZ, and WZ revised the manuscript. All

authors approved the final manuscript.

Additional files

Supplementary files

Supplementary file 1. Supplementary Figures S1 to S9.

Supplementary Figure S1. The length distribution of unigenes and the results of

alignment by BLAST.

Supplementary Figure S2. Boxplot of d_N/d_S values of all female-biased,

male-biased and unbiased genes (including $\omega \geq 2$) in floral buds and mature

flowers of the dioecious *T. pilosa*. Significant differences are represented by * in

Wilcoxon rank sum tests (***$P < 0.0005$, **$P < 0.005$ and *$P < 0.05$).

Supplementary Figure S3. Violin plots of ENCs values of female-biased,

595 male-biased and unbiased genes in floral buds. Significant differences using

596 Wilcoxon rank sum tests are represented by * (***$P < 0.0005$ and *$P < 0.05$).

597 Supplementary Figure S4. Boxplot of d_N/d_S values of male-biased genes,

598 including all 343 OGs, excluding 18 OGs under relaxed selection, excluding 98

599 OGs under positive selection, and excluding 112 OGs under both positive and

600 relaxed selection, and boxplot of d_N/d_S values of all female-biased and unbiased

601 genes in floral buds.

602 Supplementary Figure S5. Venn diagrams of female-biased genes under positive

603 selection and relaxed selection in floral buds. Overlaps of female-biased genes

604 were detected to be under positive selection using aBSREL, BUSTED, CodeML

605 and RELAX, and to be under relaxed selection using RELAX.

606 Supplementary Figure S6. Venn diagrams of unbiased genes under positive

607 selection and relaxed selection in floral buds. Overlaps of unbiased genes were

608 identified to be under positive selection using aBSREL, BUSTED, CodeML and

609 RELAX, and to be under relaxed selection using RELAX.

610 Supplementary Figure S7. Scatterplots of KEGG pathway of sex-biased in floral

611 buds of the dioecious *T. pilosa*.

612 Supplementary Figure S8. The overlap between male-biased genes with faster

613 evolutionary rates and tissue-biased genes in floral buds.

614 Supplementary Figure S9. Boxplot of d_S values of female-biased, male-biased

615 and unbiased genes in floral buds of dioecious *T. pilosa*. Significant differences

616 are represented by * in Wilcoxon rank sum tests (***$P < 0.0005$).

Supplementary file 2. Supplementary Tables S1 to S14.

Supplementary Table S1. Overview of sequencing reads from 12 samples of male

and female plants in *T. pilosa*.

Supplementary Table S2. Numbers of unigenes annotated in public databases.

Supplementary Table S3. The mapping rate of reads for each sample in floral

buds and mature flowers of the dioecious *T. pilosa*.

Supplementary Table S4. d_N, d_S and ω values of each female-biased, male-biased,

unbiased orthologous genes of floral buds and mature flowers for each species

using 'two-ratio' branch model of CodeML in PAML.

Supplementary Table S5. d_N, d_S and ω values of each female-biased, male-biased,

unbiased orthologous genes of floral buds and mature flowers for each species

using 'free-ratio' branch model of CodeML in PAML.

Supplementary Table S6. Genes under positive selection identified by branch-site

model of CodeML in PAML and functions in NR, KEGG, Swissport and GO

databases for male-biased orthologous genes in floral buds.

Supplementary Table S7. Genes under episodic positive selection tested by

aBSREL model in HyPhy and functions in NR, KEGG, Swissport and GO

databases for male-biased orthologous genes in floral buds.

Supplementary Table S8. Genes under episodic positive selection found by

BUSTED model in HyPhy and functions in NR, KEGG, Swissport and GO

databases for male-biased orthologous genes in floral buds.

Supplementary Table S9. Genes under relaxed selection detected by RELAX

model in HyPhy and functions in NR, KEGG, Swissport and GO databases for

male-biased orthologous genes in floral buds.

Supplementary Table S10. Genes under intensified positive selection identified by

RELAX model in HyPhy and functions in NR, KEGG, Swissport and GO

databases for male-biased orthologous genes in floral buds.

Supplementary Table S11. KEGG pathway enrichment analysis of female-biased

and male-biased genes in floral buds of the dioecious *T. pilosa*.

Supplementary Table S12. Functions and references associated with biotic and

abiotic stress responses, organ developments of male-biased genes under

significant positive selection ($P < 0.05$) in floral buds.

Supplementary Table S13. Functions and references associated with biotic and

abiotic stress responses, organ developments of male-biased genes under

significant relaxed selection ($P < 0.05$) in floral buds.

Supplementary Table S14. The expressions and functions of some male-biased

genes associated with senescence, raceme inflorescence development and early

flowering in floral buds.

Data Availability

All RNA-Sequencing clean reads have been deposited in the databases of the

National Center for Biotechnology Information (NCBI) under BioProject ID

PRJNA899312.

References

Abascal F, Zardoya R, Telford MJ. 2010. TranslatorX: multiple alignment of

661   nucleotide sequences guided by amino acid translations. *Nucleic Acids*

662   *Research* **38**: W7-13.

663   **Arunkumar R**, Josephs EB, Williamson RJ, Wright SI. 2013. Pollen-specific, but not

664   sperm-specific, genes show stronger purifying selection and higher rates of

665   positive selection than sporophytic genes in *Capsella grandiflora*. *Molecular*

666   *Biology and Evolution* **30**: 2475-2486.

667   **Assis R**, Zhou Q, Bachtrog D. 2012. Sex-biased transcriptome evolution in

668   *Drosophila*. *Genome Biology and Evolution* **4**: 1189-1200.

669   **Barrett SCH**, Hough J. 2013. Sexual dimorphism in flowering plants. *Journal of*

670   *Experimental Botany* **64**: 67-82.

671   **Beaudry FEG**, Rifkin JL, Barrett SCH, Wright SI. 2020. Evolutionary genomics of

672   plant gametophytic selection. *Plant Communications* **1**: 100115.

673   **Bogamuwa SP**, Jang JC. 2014. Tandem CCCH zinc finger proteins in plant growth,

674   development and stress response. *Plant and Cell Physiology* **55**: 1367-1375.

675   **Bolger AM**, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

676   sequence data. *Bioinformatics* **30**: 2114-2120.

677   **Catalan A**, Macias-Munoz A, Briscoe AD. 2018. Evolution of sex-biased gene

678   expression and dosage compensation in the eye and brain of *Heliconius*

679   *Butterflies*. *Molecular Biology and Evolution* **35**: 2343-2343.

680   **Charlesworth B**, Campos JL, Jackson BC. 2018. Faster-X evolution: Theory and

681   evidence from *Drosophila*. *Molecular Ecology* **27**: 3753-3771.

682   **Charlesworth D**. 2018. Does sexual dimorphism in plants promote sex chromosome

683   evolution? *Environmental and Experimental Botany* **146**: 5-12.

Chen F, Hu Y, Vannozzi A, Wu K, Cai H, Qin Y, Mullis A, Lin Z, Zhang L. 2017. The

WRKY transcription factor family in model plants and crops. *Critical Reviews in*

Plant Sciences **36**: 311-335.

Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M. 2005. Blast2GO:

a universal tool for annotation, visualization and analysis in functional

genomics research. *Bioinformatics* **21**: 3674-3676.

Congrains C, Campanini EB, Torres FR, Rezende VB, Nakamura AM, de Oliveira JL,

Lima ALA, Chahad-Ehlers S, Sobrinho IS, de Brito RA. 2018. Evidence of

adaptive evolution and relaxed constraints in sex-biased genes of South

American and West Indies fruit flies (Diptera: Tephritidae). *Genome Biology*

and Evolution **10**: 380-395.

Cossard GG, Godfroy O, Nehr Z, Cruaud C, Cock JM, Lipinska AP, Coelho SM.

2022. Selection drives convergent gene expression changes during transitions

to co-sexuality in haploid sexual systems. *Nature Ecology and Evolution* **6**:

579-589.

Cossard GG, Toups MA, Pannell JR. 2019. Sexual dimorphism and rapid turnover in

gene expression in pre-reproductive seedlings of a dioecious herb. *Annals of*

Botany **123**: 1119-1131.

Cui YW, Lu XT, Gou XP. 2022. Receptor-like protein kinases in plant reproduction:

current understanding and future perspectives. *Plant Communications*

3 :100273.

Dapper AL, Wade MJ. 2020. Relaxed selection and the rapid evolution of

reproductive genes. *Trends in Genetics* **36**: 640-649.

Darolti I, Mank JE. 2023. Sex-biased gene expression at single-cell resolution: cause

and consequence of sexual dimorphism. *Evolution Letters*

https://doi.org/10.1093/evlett/qrad013

Darolti I, Wright AE, Pucholt P, Berlin S, Mank JE. 2018. Slow evolution of

sex-biased genes in the reproductive tissue of the dioecious plant *Salix*

viminalis. *Molecular ecology* **27**: 694-708.

Davidson NM, Oshlack A. 2014. Corset: enabling differential gene expression

analysis for *de novo* assembled transcriptomes. *Genome Biology* **15**: 410.

Dawson TE, Geber MA. 1999. Sexual dimorphism in physiology and morphology. In:

Geber MA, Dawson TE, Delph LF (Eds). *Gender and Sexual Dimorphism in*

Flowering Plants. Berlin, Heidelberg: Springer. p. 175-215.

de Boer HJ, Schaefer H, Thulin M, Renner SS. 2012. Evolution and loss of

long-fringed petals: a case study using a dated phylogeny of the snake gourds,

Trichosanthes (Cucurbitaceae). *BMC Ecology and Evolution* **12**: 108.

de Boer HJ, Steffen K, Cooper WE. 2015. Sunda to Sahul dispersals in *Trichosanthes*

(Cucurbitaceae): a dated phylogeny reveals five independent dispersal events

to Australasia. *Journal of Biogeography* **42**: 519-531.

Delph LF. 1999. Sexual dimorphism in life history. In: Geber MA, Dawson TE,

Delph LF (Eds). *Gender and Sexual Dimorphism in Flowering Plants*. Berlin,

Heidelberg: Springer. p. 149-173.

Delph LF, Herlihy CR. 2012. Sexual, fecundity, and viability selection on flower size

and number in a sexually dimorphic plant. *Evolution* **66**: 1154-1166.

Djordjevic J, Dumas Z, Robinson-Rechavi M, Schwander T, Parker DJ. 2022.

Dynamics of sex-biased gene expression during development in the stick

insect *Timema californicum*. *Heredity* **129**:113-122.

Ellegren H, Parsch J. 2007. The evolution of sex-biased genes and sex-biased gene

expression. *Nature Reviews Genetics* **8**: 689-698.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for

comparative genomics. *Genome Biology* **20**: 238.

Forrest JRK. 2014. Plant size, sexual selection, and the evolution of protandry in

dioecious plants. *American Naturalist* **184**: 338-351.

Gershoni M, Pietrokovski S. 2014. Reduced selection and accumulation of

deleterious mutations in genes exclusively expressed in men. *Nature*

Communications **5**: 4438.

Grath S, Parsch J. 2016. Sex-biased gene expression. *Annual Review of Genetics* **50**:

29-44.

Guo J, Xu W, Hu Y, Huang J, Zhao Y, Zhang L, Huang CH, Ma H. 2020.

Phylotranscriptomics in cucurbitaceae reveal multiple whole-genome

duplications and key morphological and molecular innovations. *Molecular*

Plant **13**: 1117-1133.

Gutierrez-Valencia J, Fracassetti M, Horvath R, Laenen B, Desamore A, Drouzas

AD, Friberg M, Kolar F, Slotte T. 2022. Genomic signatures of sexual

selection on pollen-expressed genes in *Arabis alpina. Molecular Biology and*

Evolution **39**: msab349.

Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger

MB, Eccles D, Li B, Lieber M, et al. 2013. *De novo* transcript sequence

reconstruction from RNA-seq using the Trinity platform for reference

generation and analysis. *Nature Protocols* **8**: 1494-1512.

Hajdu A, Adam E, Sheerin DJ, Dobos O, Bernula P, Hiltbrunner A, Kozma-Bognar L,

Nagy F. 2015. High-level expression and phosphorylation of phytochrome B

modulates flowering time in *Arabidopsis. Plant Journal* **83**: 794-805.

Hambuch TM, Parsch J. 2005. Patterns of synonymous codon usage in *Drosophila*

melanogaster genes with sex-biased expression. *Genetics* **170**: 1691-1700.

Harkess A, Mercati F, Shan HY, Sunseri F, Falavigna A, Leebens-Mack J. 2015.

Sex-biased gene expression in dioecious garden asparagus (*Asparagus*

officinalis). *New Phytologist* **207**: 883-892.

Harrison MC, Mallon EB, Twell D, Hammond RL. 2019. Deleterious mutation

accumulation in *Arabidopsis thaliana* pollen genes: a role for a recent

relaxation of selection. *Genome Biology and Evolution* **11**: 1939-1951.

Harrison PW, Wright AE, Zimmer F, Dean R, Montgomery SH, Pointer MA, Mank

JE. 2015. Sexual selection drives evolution and rapid turnover of male gene

expression. *Proceedings of the National Academy of Sciences of the United*

States of America **112**: 4393-4398.

Hatchett WJ, Jueterbock AO, Kopp M, Coyer JA, Coelho SM, Hoarau G, Lipinska

AP. 2023. Evolutionary dynamics of sex-biased gene expression in a young XY

system: Insights from the brown alga genus *Fucus*. *New Phytologist*

https://doi.org/10.1111/nph.18710.

Hou JJ, Fan WW, Ma RR, Li B, Yuan ZH, Huang WX, Wu YY, Hu Q, Lin CJ, Zhao

XQ, et al. 2022. *MALE STERILITY 3* encodes a plant homeodomain-finger

protein for male fertility in soybean. *Journal of Integrative Plant Biology* **64**:

1076-1086.

Hsu S-K, Jakšić AM, Nolte V, Lirakis M, Kofler R, Barghi N, Versace E, Schlötterer

C. 2020. Rapid sex-specific adaptation to high temperature in *Drosophila*.

Elife **9**: e53237.

Hu XQ, Liao ZY, Zhang B, Yue JJ, Wang Z, Jie X, Liu J. 2020. Transcriptome

sequencing and screening of genes related to sex determination of

Trichosanthes kirilowii Maxim. *PLoS One* **15**: e0239230.

Hudson M, Smith H. 1998. The phytochrome B encoded by the *HLG* locus of

Nicotiana plumbaginifolia is required for detection of photoperiod: *hlg*

mutants show altered regulation of flowering and circadian movement. *Plant*

Journal **15**: 281-287.

Immler S. 2019. Haploid selection in "diploid" organisms. *Annual Review of Ecology,*

Evolution, and Systematics **50**: 219-236.

Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J,

Mitchell A, Nuka G, et al. 2014. InterProScan 5: genome-scale protein

function classification. *Bioinformatics* **30**: 1236-1240.

Khodursky S, Svetec N, Durkin SM, Zhao L. 2020. The evolution of sex-biased gene

expression in the *Drosophila* brain. *Genome Research* **30**: 874-884.

Lahti DC, Johnson NA, Ajie BC, Otto SP, Hendry AP, Blumstein DT, Coss RG,

Donohue K, Foster SA. 2009. Relaxed selection in the wild. *Trends in Ecology*

and Evolution **24**: 487-496.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nature*

Methods **9**: 357-359.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data

with or without a reference genome. *BMC Bioinformatics* **12**: 323.

Li SH, Yan H, Mei WM, Tse YC, Wang H. 2020. Boosting autophagy in sexual

reproduction: a plant perspective. *New Phytologist* **226**: 679-689.

Lichilin N, El Taher A, Bohne A. 2021. Sex-biased gene expression and recent sex

chromosome turnover. *Philosophical Transactions of the Royal Society B:*

Biological Sciences **376**: 20200107.

Lipinska A, Cormier A, Luthringer R, Peters AF, Corre E, Gachon CM, Cock JM,

Coelho SM. 2015. Sexual dimorphism and the evolution of sex-biased gene

expression in the brown alga *ectocarpus*. *Molecular Biology and Evolution* **32**:

1581-1597.

Liu ZH, Tavares R, Forsythe ES, Andre F, Lugan R, Jonasson G, Boutet-Mercey S,

Tohge T, Beilstein MA, et al. 2016. Evolutionary interplay between sister

cytochrome P450 genes shapes plasticity in plant metabolism. *Nature*

Communications **7**: 13026.

Liu Y, Bassham DC. 2012. Autophagy: pathways for self-eating in plant cells. *Annual*

Review of Plant Biology **63**: 215-237.

Lynch M, Conery JS. 2000. The evolutionary fate and consequences of duplicate

genes. *Science* **290**: 1151-1155.

Lynch M, Katju V. 2004. The altered evolutionary trajectories of gene duplicates.

Trends in Genetics **20**: 544-549.

Ma LL, Wang Q, Mu JL, Fu AZ, Wen CL, Zhao XY, Gao LP, Li J, Shi K, Wang YX,

et al. 2020. The genome and transcriptome analysis of snake gourd provide

insights into its evolution and fruit development and ripening. *Horticulture*

Research **7**: 199.

Mank JE. 2009. Sex chromosomes and the evolution of sexual dimorphism: lessons

from the genome. *American Naturalist* **173**: 141-150.

Mank JE. 2017. The transcriptional architecture of phenotypic dimorphism. *Nature*

Ecology and Evolution **1**: 6.

Mank JE, Hultin-Rosenberg L, Axelsson E, Ellegren H. 2007. Rapid evolution of

female-biased, but not male-biased, genes expressed in the avian brain.

Molecular Biology and Evolution **24**: 2698-2706.

Mao XZ, Cai T, Olyarchuk JG, Wei LP. 2005. Automated genome annotation and

pathway identification using the KEGG Orthology (KO) as a controlled

vocabulary. *Bioinformatics* **21**: 3787-3793.

Marinić M, Lynch VJ. 2020. Relaxed constraint and functional divergence of the

progesterone receptor (*PGR*) in the human stem-lineage. *PLOS Genetics* **16**:

e1008666.

Meisel RP. 2011. Towards a more nuanced understanding of the relationship between

sex-biased gene expression and rates of protein-coding sequence evolution.

Molecular Biology and Evolution **28**: 1893-1900.

Meisel RP, Connallon T. 2013. The faster-X effect: integrating theory and data.

Trends in Genetics **29**: 537-544.

Ming R, Bendahmane A, Renner SS. 2011. Sex chromosomes in land plants. *Annual*

Review of Plant Biology **62**: 485-514.

Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M. 2007. KAAS: an automatic

846   genome annotation and pathway reconstruction server. *Nucleic Acids Research*

847   **35**: W182-185.

848   **Murrell B**, Weaver S, Smith MD, Wertheim JO, Murrell S, Aylward A, Eren K,

849   Pollner T, Martin DP, Smith DM, et al. 2015. Gene-wide identification of

850   episodic selection. *Molecular Biology and Evolution* **32**: 1365-1371.

851   **Muschietti JP**, Wengier DL. 2018. How many receptor-like kinases are required to

852   operate a pollen tube. *Current Opinion in Plant Biology* **41**: 73-82.

853   **Muyle A.** 2019. How different is the evolution of sex-biased gene expression between

854   plants and animals? A commentary on: 'Sexual dimorphism and rapid turnover

855   in gene expression in prereproductive seedlings of a dioecious herb'. *Annals of*

856   *Botany* **123**: iv-v.

857   **Papa F**, Windbichler N, Waterhouse RM, Cagnetti A, D'Amato R, Persampieri T,

858   Lawniczak MKN, Nolan T, Papathanos PA. 2017. Rapid evolution of

859   female-biased genes among four species of *Anopheles* malaria mosquitoes.

860   *Genome Research* **27**: 1536-1548.

861   **Parsch J**, Ellegren H. 2013. The evolutionary causes and consequences of sex-biased

862   gene expression. *Nature Reviews Genetics* **14**: 83-87.

863   **Pond SLK**, Poon AFY, Velazquez R, Weaver S, Hepler NL, Murrell B, Shank SD,

864   Magalis BR, Bouvier D, Nekrutenko A, et al. 2020. HyPhy 2.5–a customizable

865   platform for evolutionary hypothesis testing using phylogenies. *Molecular*

866   *Biology and Evolution* **37**: 295-299.

867   **Pro☐schel M**, Zhang Z, Parsch J. 2006. Widespread adaptive evolution of *Drosophila*

868   genes with sex-biased expression. *Genetics* **174**: 893-900.

869   **Renner SS**. 2014. The relative and absolute frequencies of angiosperm sexual

870   systems: dioecy, monoecy, gynodioecy, and an updated online database.

871   *American Journal of Botany* **101**: 1588-1596.

872   **Rowe L**, Chenoweth SF, Agrawal AF. 2018. The genomics of sexual conflict.

873   *American Naturalist* **192**: 274-286.

Sanderson BJ, Wang L, Tiffin P, Wu ZQ, Olson MS. 2019. Sex-biased gene

expression in flowers, but not leaves, reveals secondary sexual dimorphism in

Populus balsamifera. New Phytologist **221**: 527-539.

Schaefer H, Renner SS. 2011. Phylogenetic relationships in the order Cucurbitales

and a new classification of the gourd family (Cucurbitaceae). *Taxon* **60**:

122-138.

Scharf KD, Berberich T, Ebersberger I, Nover L. 2012. The plant heat stress

transcription factor (Hsf) family: structure, function and evolution. *Biochimica*

Et Biophysica Acta-Gene Regulatory Mechanisms **1819**: 104-119.

Scharmann M, Rebelo AG, Pannell JR. 2021. High rates of evolution preceded shifts

to sex-biased gene expression in *Leucadendron*, the most sexually dimorphic

angiosperms. *Elife* **10**: e67485.

Schrader L, Pan H, Bollazzi M, Schiott M, Larabee FJ, Bi X, Deng Y, Zhang G,

Boomsma JJ, Rabeling C. 2021. Relaxed selection underlies genome erosion

in socially parasitic ant species. *Nature Communications* **12**: 2918.

Seppey M, Manni M, Zdobnov EM. 2019. BUSCO: Assessing genome assembly and

annotation completeness. *Methods in Molecular Biology* **1962**: 227-245.

Shiryev SA, Papadopoulos JS, Schaffer AA, Agarwala R. 2007. Improved BLAST

searches using longer words for protein seeding. *Bioinformatics* **23**:

2949-2951.

Skogsmyr I, Lankinen A. 2002. Sexual selection: an evolutionary force in plants.

Biological Reviews **77**: 537-562.

Smith MD, Wertheim JO, Weaver S, Murrell B, Scheffler K, Pond SLK. 2015. Less is

more: an adaptive branch-site random effects model for efficient detection of

episodic diversifying selection. *Molecular Biology and Evolution* **32**:

1342-1353.

Stephenson AG, Bertin RI. 1983. Male competition, female choice, and sexual

selection in plants. In: Real L (Eds). *Pollination Biology*. New York, NY, USA:

Academic Press. p.109-151.

Teo ZWN, Song SY, Wang YQ, Liu J, Yu H. 2014. New insights into the regulation of

inflorescence architecture. *Trends in Plant Science* **19**: 158-165.

Tosto NM, Beasley ER, Wong BBM, Mank JE, Flanagan SP. 2023. The roles of

sexual selection and sexual conflict in shaping patterns of genome and

transcriptome variation. *Nature Ecology and Evolution*

https://doi.org/10.1038/s41559-023-02019-7

Toubiana W, Armisén D, Dechaud C, Arbore R, Khila A. 2021. Impact of male trait

exaggeration on sex-biased gene expression and genome architecture in a

water strider. *BMC Biology* **19**:89

Veltsos P. 2019. Not all sex-biased genes are the same. *New Phytologist* **221**: 10-11.

Wertheim JO, Murrell B, Smith MD, Kosakovsky Pond SL, Scheffler K. 2015.

RELAX: detecting relaxed selection in a phylogenetic framework. *Molecular*

Biology and Evolution **32**: 820-832.

Whittle CA, Kulkarni A, Extavour CG. 2021. Evolutionary dynamics of sex-biased

genes expressed in cricket brains and gonads. *Journal of Evolutionary Biology*

34: 1188-1211.

Williams JH, Reese JB. 2019. Evolution of development of pollen performance.

Current Topics in Developmental Biology **131**: 299-336.

Wright AE, Harrison PW, Zimmer F, Montgomery SH, Pointer MA, Mank JE. 2015.

Variation in promiscuity and sexual selection drives avian rate of Faster-Z

evolution. *Molecular Ecology* **24**: 1218-1235.

Wright F. 1990. The effective number of codons used in a gene. *Gene* **87**: 23-29.

Wu H, Zhao G, Gong H, Li J, Luo C, He X, Luo S, Zheng X, Liu X, Guo J, et al.

2020. A high-quality sponge gourd (*Luffa cylindrica*) genome. *Horticulture*

Research **7**: 128.

Wu Z, Raven P, Hong D. 2011. *Flora of China.* Beijing, China: Science Press, and St.
Louis, USA: Missouri Botanical Garden Press.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Molecular*
Biology and Evolution **24**: 1586-1591.

Yang ZH, Wong WSW, Nielsen R. 2005. Bayes empirical bayes inference of amino
acid sites under positive selection. *Molecular Biology and Evolution* **22**:
1107-1118.

Zemp N, Tavares R, Muyle A, Charlesworth D, Marais GA, Widmer A. 2016.
Evolution of sex-biased gene expression in a dioecious plant. *Nature Plants* **2**:
16168.

Zhang JZ, Nielsen R, Yang ZH. 2005. Evaluation of an improved branch-site
likelihood method for detecting positive selection at the molecular level.
Molecular Biology and Evolution **22**: 2472-2479.

Zhang MM, Zhang SQ. 2022. Mitogen-activated protein kinase cascades in plant
signaling. *Journal of Integrative Plant Biology* **64**: 301-341.

Zhang LB, Simmons MP, Kocyan A, Renner SS. 2006. Phylogeny of the
Cucurbitales based on DNA sequences of nine loci from three genomes:
implications for morphological and sexual system evolution. *Molecular*
Phylogenetics and Evolution **39**: 305-322.

Zheng Y, Wu S, Bai Y, Sun HH, Jiao C, Guo SG, Zhao K, Blanca J, Zhang ZH,
Huang SW, et al. 2019. Cucurbit Genomics Database (CuGenDB): a central
portal for comparative and functional genomics of cucurbit crops. *Nucleic*
Acids Research **47**: D1128-D1136.

Zhang YW, Li ZW, Ma B, Hou QC, Wan XY. 2020. Phylogeny and functions of LOB
domain proteins in plants. *International Journal of Molecular Sciences* **21**:
2278.

Zhang Z, Hambuch TM, Parsch J. 2004. Molecular evolution of sex-biased genes in
Drosophila. *Molecular Biology and Evolution* **21**: 2130-2139.

Zhang Z, Parsch J. 2005. Positive correlation between evolutionary rate and

recombination rate in *Drosophila* genes with male-biased expression.

Molecular Biology and Evolution **22**: 1945-1947.

Zhou XM, Zhao P, Wang W, Zou J, Cheng TH, Peng XB, Sun MX. 2015. A

comprehensive, genome-wide analysis of autophagy-related genes identified

in tobacco suggests a central role of autophagy in plant response to various

environmental cues. *DNA Research* **22**: 245-257.

Figure Legends

Fig. 1 Floral buds and mature flowers of females (A, B) and males (C) in dioecious

Trichosanthes pilosa.

Fig. 2 Sex-biased gene expression for floral buds and mature flowers in males and

females of *T. pilosa.* Volcano plots of average expression between female-biased,

male-biased and unbiased genes in floral buds (A) and mature flowers (B). Heatmap

of sex-biased gene expression (C) using hierarchical clustering analysis.

Fig. 3 The overlap between sex-biased and tissue-biased genes in two type sexes and

tissues. Male-biased genes in floral buds (M1BGs) (A) or mature flowers (M2BGs) (B)

overlapped with tissue-biased genes of floral buds (M1TGs) and mature flowers

(M2TGs). Female-biased genes in floral buds (F1BGs) (C) or mature flowers (F2BGs)

(D) overlapped with tissue-biased genes of floral buds (F1TGs) and mature flowers

(F2TGs).

Fig. 4 Violin plots of d_N/d_S values ($0 < \omega < 2$) of female-biased, male-biased and

unbiased genes in floral buds and mature flowers of the dioecious *T. pilosa*. White dot

indicates the median of d_N/d_S values for sex-biased and unbiased genes. Wilcoxon

rank sum tests are used to test for significant differences (***$P < 0.0005$, **$P < 0.005$

and *$P < 0.05$). The distributions of d_N/d_S values for female-biased, male-biased and

unbiased genes in floral buds (A) and mature flowers (B) using 'two-ratio' branch

model. The distributions of d_N/d_S values for female-biased, male-biased and unbiased

genes in floral buds (C) and mature flowers (D) using 'free-ratio' branch model.

Fig. 5 Venn diagrams of male-biased genes detected to be under positive selection

using aBSREL, BUSTED, CodeML and RELAX in floral buds.

A Two-Ratio Model: Floral buds

B Two-Ratio Model: Mature flowers

C Free-Ratio Model: Floral buds

D Free-Ratio Model: Mature flowers