1

Differentiation of gram-negative intermembrane phospholipid transporter function by intrinsic substrate preference

Ashutosh K. Rai[a], Katsuhiro Sawasato[b], Anastasiia Kozlova[b], Genevieve C. Sparagna[c], Mikhail Bogdanov[b], and Angela M. Mitchell[a,#]

[a] Department of Biology, Texas A&M University, College Station, Texas, USA

[b] Department of Biochemistry and Molecular Biology, McGovern Medical School at The University of Texas Health Science Center at Houston, Houston, Texas, USA

[c] Department of Medicine, Division of Cardiology, University of Colorado Anschutz Medical Campus, Aurora, Colorado, USA

Running title: Phospholipid transporter substrate preference

Address correspondence to Angela M. Mitchell, amitchell@bio.tamu.edu

Keywords: phospholipid trafficking, gram-negative envelope, outer membrane, lipid metabolism, cold sensitivity

Abstract: 250 words

Text: 5778 words

Abstract

The outer membrane of diderm Gram-negative bacteria acts as a barrier from chemical and physical stress. Anterograde phospholipid transport to the outer membrane has long been an area of intense investigation and, in *E. coli* K-12, it has recently been shown to be mediated by three related proteins: YhdP, TamB, and YdbH, which appear to provide hydrophobic channels for phospholipid diffusion, with YhdP and TamB playing the major roles. However, YhdP and TamB have different phenotypes suggesting distinct phospholipid transport functions. We investigated these functions using the synthetic cold sensitivity of a strain with Δ*yhdP* (but not Δ*tamB* or Δ*ydbH*) and Δ*fadR*, a transcriptional regulator allowing switching between fatty acid degradation and synthesis and regulating unsaturated fatty acid production. Deletion of *tamB*, forcing phospholipid transport to YdbH, suppresses the Δ*yhdP* Δ*fadR* cold sensitivity suggesting this phenotype is due to TamB dysfunction. Increased levels of cardiolipin and fatty acid saturation are necessary for cold sensitivity and lowering levels of either suppresses this sensitivity. Our data support a model where YhdP primarily transports more saturated phospholipids, TamB primarily transports phospholipids with more than one carbon unsaturation, and cardiolipin obstructs TamB by selectively clogging its channel. Thus, the multiple phospholipid transporters may allow the saturation state of the outer membrane to be regulated independently of the inner membrane by altering the YhdP-TamB activity ratio. Maintaining membrane physical integrity and function under changing conditions may require envelope remodeling including altered phospholipid composition and intermembrane trafficking. Our data provide a potential mechanism for this regulation.

Importance

Gram-negative bacteria possess an impermeable outer membrane protecting against environmental stress and antibiotics. Outer membrane phospholipid transport remained mysterious until recently when YhdP, TamB, and YdbH were implicated in phospholipid transport between the inner and outer membranes of *E. coli*. TamB also modulates phospholipid transport in a distantly related diderm Fermicute. Here, we investigate functional differentiation between YhdP and TamB. Our data strongly suggest YhdP's and TamB's functions are distinguished by the saturation state of phospholipids with YhdP preferentially transporting more saturated phospholipids and TamB transporting more unsaturated phospholipids. Cardiolipin headgroup specificity may also contribute to TamB inhibition, perhaps due to the bulky nature of cardiolipin inhibiting the passage of other phospholipids. Diversification of function between YhdP and TamB provides a mechanism for regulation of phospholipid composition, and possibly the mechanical strength and permeability of the outer membrane, and so the cell's intrinsic antibiotic resistance, in changing environmental conditions.

Introduction

The gram-negative bacterial cell envelope has an outer membrane (OM) that sits outside the aqueous periplasm and peptidoglycan cell wall. The OM provides a barrier against various environmental stresses including toxic molecules such as antibiotics and osmotic pressure (1-6). Unlike the inner membrane (IM), a phospholipid bilayer, the OM is largely composed of phospholipids (mainly phosphatidylethanolamine (PE) (7)) in its inner leaflet and LPS (lipopolysaccharide) in its outer leaflet (2, 8). However, both membranes are asymmetric as the IM has different phospholipid compositions in its inner versus outer leaflets (9). Outer membrane proteins (OMPs) form a network across the cell surface interspersed with phase separated LPS patches (10). Lipoproteins are generally anchored in the inner leaflet of the OM (11).

OM components are synthesized in the cytoplasm or IM and transported to the OM. OMP (12), lipoprotein (11), and LPS (13) transport pathways are well defined. However, until recently, intermembrane phospholipid transport (between the IM and OM), especially anterograde transport from the IM to the OM, has remained largely very poorly understood. Phospholipids are synthesized at the IM's inner leaflet (14) and rapid, bidirectional intermembrane phospholipid transport occurs (15), even without ATP (16). Phospholipid transport has a relaxed specificity, allowing transport of non-native lipids (17, 18); however, intermembrane phospholipid composition differences are maintained (7, 19-22). Establishment and maintenance of this defined lipid topography between the IM and the OM must involve the coordinated biosynthesis and balanced bidirectional trafficking of specific phospholipids across the IM and from the IM to the OM (23).

A recent study has demonstrated that anterograde phospholipid transport is mediated by a high-flux, diffusive system in *Escherichia coli* (24). The Mla system removes mislocalized phospholipids from the OM outer leaflet and returns them to the IM (25). A gain-of-function allele, *mlaA** (26, 27), opens a channel allowing phospholipids to mislocalize to the cell surface by flowing from the OM's inner to outer leaflet (26, 28), resulting in a PldA-mediated attempt to reestablish OM asymmetry by increasing LPS production, OM vesiculation, and increased flow of phospholipids to the OM (26, 27). When nutrients are depleted, inhibiting phospholipid production, cells lyse due to loss of IM integrity (16, 18). Single cell imaging of *mlaA** cells confirmed an aberrant lipid flow from the IM to the OM at a fast rate. Loss of YhdP, an inner membrane protein, could significantly slow phospholipid flow to the OM resulting in loss of OM integrity before IM rupture (24), suggesting YhdP might play a role in intermembrane phospholipid transport.

Very recently, YhdP and two homologs, TamB and YdbH, were demonstrated to be intermembrane phospholipid transporters (29, 30). Ruiz, *et al*. investigated AsmA family proteins and found Δ*yhdP* Δ*tamB* mutants showed synthetic OM permeability and stationary phase lysis (29). Moreover, when combined, Δ*yhdP*, Δ*tamB*, and Δ*ydbH* are synthetically lethal. Genetic interactions of these mutants with *mlaA* and *pldA* show their

involvement in OM phospholipid homeostasis. Predicted structures of YhdP, TamB, and
YdbH (31, 32), as well as a partial crystal structure of TamB (33), have a β-taco fold domain
with a hydrophobic pocket, resembling eukaryotic lipid transporters (34-37). These
structural findings are another clue that suggests YhdP, TamB, and YdbH are involved in
phospholipid transport. Douglass, *et al.* confirmed the synthetic lethality of Δ*yhdP*,
Δ*tamB*, and Δ*ydbH* and demonstrated a Δ*tamB* Δ*ydbH* mutant with reduced *yhdP*
expression had decreased amounts of OM phospholipids, directly demonstrating their
involvement in intermembrane phospholipid transport or its regulation (30), although *in*
vitro interaction of YhdP, TamB, and YdbH with phospholipids has yet to be investigated.
Nevertheless, the role of these proteins in phospholipid transport may be widely
conserved as recent evidence suggests TamB plays a role in anterograde phospholipid
transport in *Veillonella parvula*, a diderm Fermicute (38).

Nevertheless, why there are three separate intermembrane phospholipid transport
proteins and the functional interactions between these proteins remains unanswered,
although it is clear the proteins are not fully redundant or functionally equivalent. The
conditional expression of *yhdP* alone, without the presence of *ydbH* and *tamB*, fully
complements neither growth phenotypes nor a normal level of OM phospholipids,
suggesting transport function is still impaired in this strain (30). In addition, YdbH seems
to play a more minor role than YhdP and TamB (29, 30) and the screen identifying Δ*yhdP*
as slowing *mlaA**-dependent lysis did not identify Δ*tamB* or Δ*ydbH* (24). Moreover, we
previously identified a role for YhdP in stationary phase SDS (sodium dodecyl sulfate)
resistance and modulating cyclic enterobacterial common antigen activity (39, 40) not
shared by TamB and YdbH (**Fig. S1**) (29, 30, 39, 40).

Here, we investigated the differentiation of YhdP and TamB function and identified
synthetic cold sensitivity due to impaired TamB function in a strain with Δ*yhdP* and Δ*fadR*,
a transcriptional regulator of fatty acid biosynthesis and degradation, acting as a
transcriptional switch between these pathways in response to acyl-CoA (41). FadR is
necessary for normal levels of unsaturated fatty acids to be synthesized (42). We found
for the first time that the Δ*yhdP* Δ*fadR* mutant has increased levels of a specific
phospholipid, cardiolipin (CL), and decreasing CL suppresses cold sensitivity. Furthermore,
increasing unsaturated fatty acid levels suppresses cold sensitivity. Overall, our data lead
us to a model where TamB and YhdP functions are differentiated based on phospholipid
fatty acid saturation state, with TamB transporting more unsaturated phospholipids and
YhdP transporting more saturated phospholipids. Phospholipid species or head group can
also contribute to transport specificity, since higher levels of dianionic CL are necessary

for the cold sensitivity phenotype, potentially due to functional inhibition of TamB by
clogging of its hydrophobic groove.

Results

Loss of FadR and YhdP results in cold sensitivity due to impaired TamB function. To
investigate YhdP-TamB function differentiation, we created a ΔyhdP ΔfadR strain, as FadR
is a major regulator of lipid homeostasis (41). A deletion strain of *fadR* alone is expected
to have decreased fatty acid synthesis, increased fatty acid degradation, and increased
fatty acid saturation (41, 42). Surprisingly, ΔyhdP ΔfadR cultures grown at 30 °C lagged for
more than 8 hours, followed by highly variable growth, suggesting suppressor outgrowth
(**Fig.1A**). The growth defect was decreased at 37 °C and completely absent at 42 °C. To
confirm this cold sensitivity, we estimated efficiency of plating(EOP) and confirmed the
severe cold intolerance of the ΔyhdP ΔfadR strain (5 logs at 30 °C and 3 logs at 37 °C) (**Fig.
1B**). Although, cyclic enterobacterial common antigen is necessary for ΔyhdP's OM
permeability phenotypes (40), the ΔyhdP ΔfadR strain's cold sensitivity was unchanged
when enterobacterial common antigen was not present (ΔwecA) (43) (**Fig. 1C**).

Given their involvement in phospholipid transport, we expected ΔtamB and/or ΔydbH to
show similar synthetic phenotypes with ΔfadR; however, ΔfadR ΔtamB and ΔfadR ΔydbH
strains had no growth defects (**Fig 1D**). We, therefore, hypothesized the ΔyhdP ΔfadR cold
sensitivity was due to impairment of another phospholipid transporter(s). Although
ΔydbH in the ΔyhdP ΔfadR strain did not alter cold sensitivity (**Fig. 1E**), ΔtamB in the ΔyhdP
ΔfadR background completely suppressed the cold sensitivity (**Fig. 1F**). Consistent with
previous observations (29), the ΔyhdP ΔfadR ΔtamB strain was mucoid due to Rcs stress
response activation and colanic acid capsule production. Deleting *rcsB*, an Rcs response
regulator, prevented mucoidy but did not affect suppression (**Fig. 1F**). We observed only
minimal growth phenotypes in the ΔyhdP ΔfadR ΔtamB strain, similar to those of ΔtamB
and less severe than those of ΔfadR (**Fig. 1G**), eliminating slow growth as a suppression
mechanism. Together, these data demonstrate suppression from loss of TamB is due to
reprieve of impaired TamB function, shifting phospholipid transport to YdbH.

We next looked for envelope homeostasis alterations in the ΔyhdP ΔfadR strain with
TamB impairment. To identify cell lysis and/or severe OM permeability defects at 30 °C,
we assayed CPRG (chlorophenol red-β-D-galactopyranoside) processing. CPRG must
contact LacZ in the cytoplasm or supernatant to release chlorophenol red (44). Compared
to wild type or single mutants, the ΔyhdP ΔfadR strain had increased CPRG processing
(**Fig. S2A**), demonstrating lysis or increased envelope permeability. When we assayed
resistance to molecules excluded by the OM (1, 39, 40, 45, 46) at 37 °C, a semi-permissive
temperature, the ΔyhdP ΔfadR strain showed sensitivity to vancomycin, a large scaffold
antibiotic, and SDS EDTA compared to single mutants (**Fig. S2B**). However, with bacitracin
treatment or EDTA treatment alone, the ΔyhdP ΔfadR strain phenocopied the ΔfadR

single deletion (**Fig. S2CD**). LPS levels were not altered in the ΔyhdP, ΔfadR, or the ΔyhdP
ΔfadR strain (**Fig. S2E**). OM asymmetry mutations (i.e., ΔpldA, ΔmlaA) (25-27, 47) did not
have suppressive or synthetic effects on cold sensitivity in the ΔyhdP ΔfadR strain (**Fig.**
S3AB), suggesting against a causal loss of OM asymmetry. Overall, the changes in OM
permeability did not seem sufficient to cause lethality at lower temperature, suggesting
the phospholipid transport disruption affects both IM and OM integrity.

Increased cardiolipin is necessary for impaired TamB function. We investigated
phospholipid composition of the wild-type and ΔyhdP ΔfadR strains grown at the
permissive temperature (42 °C) and grown to early log phase at 42 °C, then downshifted
to 30 °C. *E. coli* phospholipid composition is generally 75% phosphatidylethanolamine
(PE), 20% phosphatidylglycerol (PG), and 5% cardiolipin (CL) with CL increasing in
stationary phase (14, 48). Of the three CL synthases, ClsA and ClsB synthesizes CL from
two PG molecules, while ClsC synthesizes CL from one PG and one PE molecule, so CL and
PG levels are generally reciprocally regulated(49-52). For both strains and conditions, the
levels of PE were similar (**Fig. 2A**). However, the ΔyhdP ΔfadR strain had increased CL and
concomitant decreased PG compared to the wild-type strain, especially at 30 °C. PgsA
synthesizes phosphatidylglycerol-phosphate and is the first enzyme differentiating PG
and PE biosynthesis. PgsA depletion would decrease both PG and CL levels, reversing
increased CL levels and exacerbating decreased PG levels in the ΔyhdP ΔfadR strain. We
tested the effect of PgsA depletion in a strain background (ΔrcsF Δlpp) where *pgsA* is non-
essential (53-56). When *pgsA* expression was induced (arabinose panel), the ΔyhdP ΔfadR
PgsA-depletion strain had equal cold sensitivity to the ΔyhdP ΔfadR strain (**Fig. 2B**). When
pgsA expression was repressed (glucose panel), the ΔyhdP ΔfadR cold sensitivity was
partial suppressed, demonstrating that decreased PG does not cause for cold sensitivity
and suggesting increased CL may be necessary for cold sensitivity. Indeed, the cold
sensitivity of the ΔyhdP ΔfadR strain was lost when CL was fully depleted due to deletion
of the genes for all cardiolipin syntheses (ΔclsABC) or of the gene for the main exponential
phase cardiolipin synthase (ΔclsA) (**Fig. 2ACD**). ΔclsB or ΔclsC caused a smaller decrease
in CL levels that was very similar between the two strains (11% to 7.4% vs. 7.6%,
respectively) (**Fig. 2A**). Surprisingly, ΔclsC completely suppressed the cold sensitivity,
while ΔclsB did not (**Fig. 2EF**). Thus, increased CL is necessary for impaired TamB function
leading to cold sensitivity but CL levels alone cannot explain the phenotype: another
factor, such as a specific molecular form (e.g., acyl chain length, saturation state,
symmetry of acyl chain arrangement) or localization (e.g., poles vs. midcell or inner vs.
outer leaflet)of CL, is required.

Increased *fabA* expression suppresses cold sensitivity. To identify other factors involved
in the cold sensitivity, we applied a forward genetic approach by isolating spontaneous
suppressor mutants capable of growing at 30 °C and identified a suppressor mutant
(Suppressor 1) that restored growth at 30 °C (**Fig. 3A**), decreased CPRG processing (**Fig.**
S2A), and had very similar CL to the ΔyhdP ΔfadR strain (**Fig. 3B**, **Fig. S4**). We identified a
point mutation in the *fabA* promoter region in Suppressor 1 (**Fig. 3C**) and confirmed this

mutation was sufficient for suppression (**Fig. 3D**). FabA is a dehydratase/isomerase that
introduces *cis* unsaturation into fatty acids (14, 57) and *fabA* is expressed from two
promoters controlled by FadR and FabR (58, 59). FadR activates *fabA* expression, while
FabR represses. Thus, *fadR* deletion would be expected to lower *fabA* expression. The
suppressor mutation was located in the *fabA* promoter FabR binding site (**Fig. 3C**) and we
hypothesized this mutation would increase *fabA* expression. *fabA* mRNA levels were more
than 4-fold lower in the Δ*yhdP* Δ*fadR* strain than the wild-type strain (**Fig. 3E**). While lower
than wild type, the suppressor mutant increased *fabA* mRNA 1.6-fold over the Δ*yhdP*
Δ*fadR* strain.

We constructed luciferase reporter plasmids with the wild-type *fabA* promoter region or
the *fabA* promoter region containing the suppressor mutation. The reporter with the
mutated promoter had higher activity in a wild-type, Δ*fadR,* and Δ*fabR* background (**Fig.**
3F). *fabA* transcription occurs from a promoter downstream of the FabR binding site (59-
61). However, when *fadR* is deleted, transcription shifts to the FabR-regulated promoter
within the FabR binding region (59-61). Our data indicate the Suppressor 1 mutation
regulates this promoter's constitutive activity rather than FabR binding. To confirm the
effect of changing the second promoter's activity, we tested the effect of Δ*fabR* on cold
sensitivity in the Δ*yhdP* Δ*fadR* strain and observed partial suppression (**Fig. S5**). The
smaller effect of Δ*fabR* compared to Suppressor 1 may result from other gene expression
changes or to the smaller relative effect of Δ*fabR* on *fabA* expression (**Fig. 3F**). Overall,
these data demonstrate increasing *fabA* expression, which is necessary for unsaturated
fatty acid biosynthesis, rescue TamB's functional deficiency in the Δ*yhdP* Δ*fadR* strain
without changing CL levels.

Increasing unsaturated fatty acids relieves impaired TamB function. Given the effect of
fabA expression on the Δ*yhdP* Δ*fadR* cold sensitivity, we used liquid chromatography-
electrospray ionization mass spectrometry (LC/MS) to characterize the strains'
phospholipid saturation state (**Dataset S1**, **Fig. 4A-C**, **Fig. S6**). We analyzed whole cell and
IM extracts (**Fig. S6A**) from cells grown to early log phase at 42 °C then downshifted to 30
°C. As expected, in whole cell extracts, Δ*fadR* caused increased fully saturated and
monounsaturated PE with a concomitant decrease of diunsaturated PE (**Fig. 4A**, **Fig. S6B**).
Levels of fully saturated PG were also increased with a decrease in mono- and
diunsaturated species (**Fig. 4B**, **Fig. S6C**). Similarly, the Δ*fadR* strain demonstrated
increased monounsaturated CL with a concomitant decrease in triunsaturated CL (**Fig. 4C**,
Fig. S6D). Compared to the Δ*fadR* strain, the Δ*yhdP* Δ*fadR* strain had slightly increased
saturation for all three phospholipids, with the largest increase in fully saturated PE.
Suppressor 1 demonstrated decreased saturation of all three phospholipids but especially
of PE compared to the Δ*yhdP* Δ*fadR* strain (**Fig. 4A-C**, **Fig. S6E-G**). The OM is enriched for
PE and its saturated species compared to the IM outer leaflet (7, 9, 20-22) making the
large PE changes intriguing. We wondered whether the suppression difference between
Δ*clsC* and Δ*clsB* resulted from the CL saturation state. However, the Δ*yhdP* Δ*fadR* Δ*clsB*
and Δ*yhdP* Δ*fadR* Δ*clsC* strains had very similar CL profiles (**Fig. 4C**, **Fig. S6G**), suggesting

another qualitative difference affecting suppression, perhaps CL lateral or inter-leaflet
distribution in the cell.

We hypothesized increasing unsaturated fatty acids suppresses Δ*yhdP* Δ*fadR* cold
sensitivity and performed EOPs for cold sensitivity on media containing an unsaturated
fatty acid (oleic acid, C18:1 *cis*-9). Exogenous phospholipids can be taken up by *E. coli*,
attached to acyl-CoA, and incorporated into phospholipids (62). Oleic acid addition
suppressed the Δ*yhdP* Δ*fadR* strain's cold sensitivity (**Fig. 4D**). Supplementing oleic acid
increases unsaturated phospholipids and provides an exogenous fatty acid source. To
differentiate between saturation state and fatty acid availability, we compared treatment
with oleic acid and saturated stearic acid (C18:0). Only oleic acid, and not stearic acid,
suppressed the Δ*yhdP* Δ*fadR* strain's cold sensitivity (**Fig. 4E**), confirming increasing
unsaturated fatty acids, not fatty acid availability suppresses.

These data suggested the Δ*yhdP* Δ*fadR* strain was impaired in intermembrane transport
of more saturated phospholipids. To investigate phospholipid transport defects, we
compared phospholipids extracted from whole cells and isolated pure IM (**Fig. S6A**). We
did not use OM extracts as they are generally contaminated with IM, complicating
interpretation. PG and CL showed small differences between whole cell and IM extracts
but were largely similar between strains (**Fig. 4BC**). However, PE showed larger strain
differences. In the wild-type strain, fully saturated PE was enriched in whole cells
compared to IM, demonstrating OM enrichment of saturated PE (**Fig. 4A**, **Fig. S6H**). In
the Δ*fadR* strain, whole cell and IM extracts had similar fully saturated PE levels, but IM
extracts had higher amounts of diunsaturated PE, still suggesting disproportional OM
enrichment with more saturated PE. However, in the Δ*yhdP* Δ*fadR* strain, while the IM
had higher levels of diunsaturated PE, the IM was underrepresented for fully saturated
PE (**Fig. 4A**, **Fig. S6H**). The Δ*yhdP* Δ*fadR* IM also had much lower levels of short chain PE
(30:0 and 30:1) than whole cells (**Fig. 4A**, **Fig. S6HI**). Short chain phospholipids decrease
membrane rigidity and remain fluid at lower temperatures (63). Thus, the short fatty acid
PE increase after short-term adaptation to 30 °C may be a cellular attempt to increase
fluidity and restore homeostatic balance between the membranes. In Suppressor 1, the
profile more closely resembled Δ*fadR* cells. Thus, the alteration of PE saturation and fatty
acid length in the Δ*yhdP* Δ*fadR* OM demonstrates a disportortial distribution of
saturated and unsaturated PE as a consequence of disruption in intermembrane
phospholipid trafficking due to impairment of TamB function.

Suppression causes fatty acid metabolism and stress response alterations. In wild-type
cells, *fabA* overexpression does not change phospholipid saturation levels, demonstrating
the FabB enzyme is rate limiting for unsaturated fatty acid biosynthesis (64). We were
intrigued that Suppressor 1 altered saturation without directly effecting *fabB* expression.
We also wondered whether CL levels in the Δ*yhdP* Δ*fadR* strain are transcriptionally
regulated. Thus, we investigated the transcriptional landscape of the wild type, the Δ*yhdP*
Δ*fadR*, and Suppressor 1 strains after a 30 minute downshift of growth to 30 °C to

determine: (i) whether the Δ*yhdP* Δ*fadR* strain had altered *cls* gene expression; (ii)
whether Suppressor 1 had other transcriptional changes; and (iii) why increased *fabA*
expression decreased saturation. All differentially expressed genes (>2-fold change,
$p < 0.05$) are listed in **Dataset S2**. Principle component analysis of these genes showed the
closest relation between Δ*yhdP* Δ*fadR* strain and Suppressor 1, while the wild-type strain
was more distantly related (**Fig. 5A**, **S7A**). Nevertheless, Suppressor 1 was more closely
related to wild type than was the Δ*yhdP* Δ*fadR* strain. Most differentially regulated genes
between Δ*yhdP* Δ*fadR* and wild type and Suppressor 1 and wild type were upregulated
(**Fig. 5BC**, **S7BC**). Many of the most highly regulated genes are FadR-regulon members.
The majority of differentially regulated genes in Suppressor 1 compared to Δ*yhdP* Δ*fadR*
are downregulated (**Fig. 5D, S7D**). These genes are involved in many cellular pathways
and likely reflect the decreased cellular stress (**Table S2**). Indeed, many external stress
response genes are more enriched in Δ*yhdP* Δ*fadR* than in Suppressor 1 (**Fig. S8A**).

No significant expression changes occurred in the CL biosynthesis pathway (**Fig. 5E**) or
other phospholipid biosynthesis genes (**Fig. S8B**). The increased cardiolipin in Δ*yhdP*
Δ*fadR* and Suppressor 1 may be post-transcriptional or due to a change in a regulatory
pathway not yet altered after 30 minutes at 30 °C (**Fig. S8C**). Levels of mitochondrial CL
can be altered due to differences in substrate binding affinity of the CL synthase based on
saturation, with increased saturation decreasing CL synthesis (65), and it may be that a
similar pathway operates here, albeit in a different direction. We examined the
expression of genes in fatty acid synthesis and degradation, many of which are members
of the FadR regulon. In Δ*yhdP* Δ*fadR* compared to wild type, many fatty acid biosynthesis
genes significantly decreased while many fatty acid degradation genes significantly
increased (**Fig. 5F**), consistent with Δ*fadR* effects. While *fabA* expression is decreased,
fabB expression is not changed, explaining why increased *fabA* expression causes
decreased saturation in Suppressor 1. Overall, our data demonstrate the Δ*yhdP* Δ*fadR*
strain's cold sensitivity is due to impaired TamB function, and increased phospholipid
saturation and cardiolipin are necessary for this functional impairment.

330
331

Discussion
The identification of YhdP, TamB, and YdbH as putative intermembrane phospholipid
transporters (29, 30) posed the tantalizing question: why is having three transporters
advantageous? Here, we demonstrate each protein plays a distinguishable role in
phospholipid transport that can be differentiated based on genetic interactions with lipid
metabolism. Disruption of FadR and YhdP function causes synthetic cold sensitivity not
shared by TamB or YdbH, due to impaired TamB function, which can be suppressed by
switching transport to YdbH. This phenotype requires both increased levels of CL and
saturated fatty acids, and decreasing amounts of either suppresses. In addition, our data
demonstrate CL synthesized by ClsB and ClsC is qualitatively different—resulting in
differential suppression phenotypes—likely due to differences in CL localization, CL-

mediated, leaflet-specific changes in lipid packing order (9, 23), localized membrane
fluidity (66), or TamB interaction.

In our model based on these data (**Fig. 6A-C**), the functions of YhdP and TamB are
diversified based on the acyl carbon saturation level preference of the phospholipids they
transport: YhdP transports mainly phospholipids with more saturated fatty acids, while
TamB transports phospholipids with more unsaturated fatty acids (**Fig. 6A**). Our data and
previous data (29, 30) agree YdbH plays a relatively minor role. In the Δ*yhdP* Δ*fadR* strain,
YhdP's absence forces more saturated phospholipids to be transported by TamB. At 30
°C, where membrane fluidity and lipid packing order are decreased and compensating
levels of CL molecules are high, TamB becomes "clogged" by phospholipids with more
saturated phospholipids, impeding transport (**Fig. 6B**). Bulky and intrinsically disordered
CL molecules could contribute to the clogging sterically by obstructing the lipid channel
and precluding it from translocating other lipid substrates (i.e., PE). . Impaired transport
explains the difference in OM PE saturation profiles between the Δ*yhdP* Δ*fadR* strain and
the other, non-cold-sensitive strains. When Δ*tamB* is deleted, phospholipid flow is shifted
to YdbH, bypassing clogging and relieving cold sensitivity (**Fig. 6C**). YdbH's ability to
successfully transport phospholipids in the Δ*yhdP* Δ*tamB* strain only when TamB is absent
may be because YhdP and TamB are primary transporters (e.g., preferred interaction
partners for other pathway members), due to direct toxicity caused by TamB clogging, or
because *ydbH* is upregulated.

Several pieces of evidence demonstrate dysregulation of phospholipid transport not just
changes to OM phospholipid composition, and so permeability or physical strength, cause
the cold sensitivity phenotype of the Δ*yhdP* Δ*fadR* strain. First, the cold sensitivity is
unaffected by removing enterobacterial common antigen, which suppresses the
membrane permeability of a Δ*yhdP* strain (**Fig. 1C**, (40)), and the OM permeability
phenotypes of the Δ*yhdP* Δ*fadR* are not severe (**Fig. S2**). Second, the cold sensitivity is
suppressed by deletion of Δ*tamB* (**Fig. 1FG**). Δ*yhdP* Δ*tamB* strains have much greater OM
permeability (29) than the Δ*yhdP* or Δ*yhdP* Δ*fadR* strains. Third, CL has been shown to
have a normalizing, bidirectional, and leaflet-specific effect on lipid packing order (67, 68)
and membrane fluidity (69-72) opposite of or similar to that of cholesterol, respectively,
and so is unlike to contribute to a phenotype dependent solely on OM membrane fluidity.
Thus, the cold sensitivity phenotype correlates with the impairment of phospholipid
trafficking and the likely imbalance of phospholipids between the IM and OM and not
with the permeability phenotypes of the OM.

Although both fatty acid saturation and high CL levels are required for the Δ*yhdP* Δ*fadR*
strain's cold sensitivity, two lines of evidence point to saturation rather than headgroup
as the main factor separating YhdP and TamB function. (i) YhdP loss slows phospholipid
transport from the IM to the OM in a *mlaA** mutant, which losses OM material due to
vesiculation and lysis when phospholipid biosynthesis is insufficient to replace lost
phospholipids (24, 26). CL only accounts for 5% of phospholipids in exponentially growing

cells making unlikely loss of CL transport could slow net phospholipid transport to the
extent required to slow lysis. (ii) Δ*tamB* has a small growth defect at 30 °C (**Fig. 1G**) not
observed at higher temperature (29, 30), perhaps due to increased unsaturated
phospholipids (73-75) resulting in decreased net OM transport by YhdP or to lower OM
fluidity. Together, these observations provide strong evidence YhdP has a substrate
preference for phospholipids with more saturated fatty acids, while TamB prefers
phospholipids with more unsaturated fatty acids.

Our model suggests the IM and OM saturation state could be independently regulated
based on YhdP and TamB activity or levels. IM saturation is regulated by altering the
composition of newly synthesized fatty acids, while OM saturation could be separately
regulated by allowing increased transport of more saturated or unsaturated
phospholipids as necessary. IM and OM acyl saturation has been studied in several strains
of *E. coli* and in *Yersinia pseudotuberculosis* (7, 19-21) and higher saturated fatty acid
levels were observed in the OM than in the IM, supporting discrimination of phospholipids
transport based on saturation. The IM needs to maintain an appropriate level of fluidity
to allow IM proteins to fold and diffuse, to allow diffusion of hydrophobic molecules, to
regulate cellular respiration, and to maintain membrane integrity (76, 77). However, the
OM is a permeability barrier, a load-bearing element that resists turgor pressure (1-6),
and has an outer leaflet consisting of LPS (2, 8). LPS has saturated acyl chains and is largely
immobile in *E. coli* (78) and a specific phospholipid saturation level may be necessary to
match or counterbalance LPS rigidity. OM phospholipids may also play a role in the
physical strength of the membrane. Interestingly, previous work has demonstrated that
the tensile strength of the OM is lessened in a *yhdP* mutant (24).

Possessing two different phospholipid transporters with different saturation state
preferences would allow the cell to adapt OM saturation to changing conditions including
temperature shifts, chemical insults, or changes synthesis of other OM components (e.g.,
LPS or OMPs). A recent study demonstrated in *Y. pseudotuberculosis* the difference in
saturation levels between the membranes is exaggerated at 37 °C compared to 8 °C (21).
Thus, conditions where IM and OM saturation are independently regulated exist and this
regulation is likely important for optimum fitness of the cells. Transcriptional regulation
of *yhdP* and *tamB* has not been thoroughly explored. It seems likely *yhdP* transcription is
quite complex with several promoters and transcriptional termination sites (79-81). *tamB*
is predicted to be in an operon with *tamA* (which codes for an OMP necessary for TamB
function) and *ytfP* (79, 82) and to have a σ^E-dependent promoter (83). The σ^E stress
response is activated by unfolded OMP accumulation, and downregulates the production
of OMPs and upregulates chaperones and folding machinery (68). It is intriguing to
speculate a change in OM phospholipid composition might aid in OMP folding due to
altered biophysical properties and/or lipoprotein diffusion.

The mechanism of phospholipid transfer between the IM and OM has been one of the
largest remaining questions in gram-negative envelope biogenesis and has remained an

area of intense investigation for decades. Furthermore, given the lack of knowledge of
this pathway, the role of phospholipid composition in the permeability barrier of the OM,
and so the intrinsic antibiotic resistance of gram-negative bacteria, remains unknown.
Recent work has begun to elucidate the intermembrane phospholipid transport pathway
by identifying YhdP, TamB, and YdbH as proteins capable of transporting phospholipids
between the membranes (24, 29, 30). Our work has now clarified that YhdP and TamB
may have separate roles in phospholipid transport that are differentiated by their
preference for phospholipid saturation states and possibly lipid headgroup. Homologs of
YhdP and TamB are found throughout *Enterobacterales* and more distantly related
members of the ASMA family are wide spread in gram-negative bacteria (84). In fact,
TamB has recently been shown to play a role in anterograde intermembrane phospholipid
transport in the diderm Firmicute *Veillonella parvula* (38). Structural predictions of YhdP
and TamB are strikingly similar between species (31, 32). Given this, it is quite likely similar
mechanisms of discrimination in intermembrane phospholipid transfer exist in other
species. The data we present here provide a framework for investigation of phospholipid
transport and mechanisms differentiating phospholipid transporter function in these
species.
440
441
Materials and Methods

Bacterial strains and Growth Conditions. Table S3 lists the strains used in this study. Most
knockout strains were constructed with Keio collection alleles using P1vir transduction
(85, 86). New alleles were constructed through λ-red recombineering (87), using the
primers indicated in **Table S4**. The FLP recombinase-FRT system was used to remove
antibiotic resistance cassettes as has been described (87). *E. coli* cultures were grown in
LB Lennox media at 42°C, unless otherwise mentioned. Where noted, LB was
supplemented with vancomycin (Gold Biotechnology), bacitracin (Gold Biotechnology),
SDS (Thermo Fisher Scientific), EDTA (Thermo Fisher Scientific), oleic acid (Sigma Aldrich),
stearic acid (Sigma Aldrich), arabinose (Gold Biotechnology), or glucose (Thermo
Scientific). For plasmid maintenance, media was supplemented with 25 µg/mL kanamycin
(Gold Biotechnology), 10 or 20 µg/mL chloramphenicol (Gold Biotechnology), or 10 µg/mL
tetracycline (Gold Biotechnology) as appropriate.

Plasmid construction. The pJW15-P$_{fabA}$, and pJW15-P$_{fabA}$(-19G>A) reporter plasmids were
constructed by amplifying the region upstream of *fabA* from -432 bp to +27 bp relative to
the annotated transcriptional start site from wild-type MG1655 or Suppressor 1,
respectively, using the Overlap-pJW15-*fabA* FP and RP primers (**Table S4**). pJW15 was
amplified using primers Overlap-pJW15-*fabA* FP1 and RP1. The pJW15 plasmid was a kind
of gift from Dr. Tracy Raivio (Univeristy of Alberta) (88, 89). The resulting fragments were
assembled using HiFi Assembly master mix (New England Biolabs) as per the
manufacturer's instructions.

Growth and Sensitivity Assays. For EOPs assays, cultures were grown overnight, serially diluted, and plated on the indicated media. For *pgsA* depletion, overnight cultures grown in arabinose were washed and diluted 1:100 into fresh LB with 0.2% glucose or arabinose as indicated and grown to OD_{600}=1 before diluting and plating on media with glucose or arabinose respective, as above. EOPs were incubated at 30 °C unless otherwise indicated. Growth curves were performed at the indicated temperature as has been described (40). For CPRG assays (44), strains were streaked on LB plates supplemented with CPRG (40µg/ml) and IPTG (100 µM) incubated overnight at 30°C.

Suppressor isolation and sequencing. Serially diluted cultures were spread on LB plates and incubated overnight at a 30°C. Colonies that grew at 30°C were isolated and rechecked for cold-resistance through EOP and growth curves. The suppressors were subjected to whole-genome sequencing to identify potential suppressor mutations. Genomic DNA was isolated from the parent and suppressors using a DNeasy Blood and Tissue Kit (Qiagen) as per the manufacturer's instructions. Library preparation and Illumina sequencing were performed by the Microbial Genome Sequencing Center (MiGS, Pittsburgh, PA, USA). Briefly, Illumina libraries were prepared and the libraries were sequenced with paired-end 151 bp reads on a NextSeq 2000 platform to a read depth of at least 200 Mbp. To identify variants from the reference genome (GenBank: U00096.3), reads were trimmed to remove adaptors and the *breseq* software (version 0.35.4) was used to call variants (90). Identified mutations were confirmed with Sanger sequencing. The consensus motif of the FabR binding site of *fabA* was determined using Multiple Em for Motif Elicitation (MEME) tool in the MEME Suite based on the sequence of putative FabR binding sites reported by Feng and Cronan (2011) (58, 91, 92). The ability of the identified P*fabA* mutation to suppress was confirmed by linking the mutant to *zcb*-3059::Tn*10* (93) and transducing to a clean background.

Lux reporter assays. Overnight cultures of strains harboring the pJW15-P*fabA* and pJW15-P*fabA*(-19G>A) plasmids were sub-cultured at (1:100) into 200 µl of fresh LB broth in a black 96-well plate. The plate was incubated in a BioTek Synergy H1 plate reader and the OD_{600} and luminescence intensity were recorded every 2.5 min for 6 hours as described in previously (89, 94). Each biological replicate was performed in technical triplicate.

RNA-seq. Overnight cultures grown at 42°C were subcultured 1:100 into fresh media and incubated at 42 °C until an OD_{600} of 0.4. Then, the cultures were shifted to a 30 °C water bath for 30 minutes. 500 µL of each culture was immediately fixed in the RNAprotect Bacteria Reagent (Qiagen) as per the manufacturer's instructions. RNA was purified using the RNeasy Kit (Qiagen) with on-column DNase digestion following the manufacturer's protocol for gram-negative bacteria. Library preparation, Illumina sequencing, and analysis of differentially expressed genes were performed by MiGS. Briefly, samples were treated with RNase free DNase (Invitrogen) and library preparation was performed with the Stranded Total RNA Prep Ligation with Ribo-Zero Plus Kit (Illumina). Libraries were sequenced with 2x50 bp reads on a NextSeq2000 (Illumina). Demultiplexing, quality

control, and adapter trimming carried out using bcl-convert (Illumina, v3.9.3) and bcl2fastq (Illumina). Read mapping and read quantification were performed with HISAT2 (95) and the featureCounts tool in Subreader (96), respectively. Descriptive statistics of the RNA-seq read data can be found in **Table S1**. The raw RNA-seq data is available in the Sequence Read Archive database (SRA) (https://www.ncbi.nlm.nih.gov/sra, BioProject ID PRJNA965821; Reviewer Access).

Raw read counts were normalized using the Trimmed Mean of M values algorithm in edgeR (97) and converted to counts per million. The Quasi-Linear F-Test function of edgeR was used for differential expression analysis. Normalized read values and for all genes as well as differential expression analysis statistics can be found in **Dataset S3**. KEGG pathway analysis was conducted using the *"kegga"* function of limma (98). Expression of genes differentially expressed (greater than 2-fold change and $p < 0.05$) was subjected to principle component analysis, and pathway and enrichment analysis using the EcoCyc Omics Dashboard (98).

Lipid extraction and TLC. For thin layer chromatography (TLC) experiments, cultures were grown in LB with 1 mCi/mL ^{32}P until OD_{600} of 0.6-0.8 at the indicated temperature. For temperature downshift experiments, cells were grown at 42 $^{\circ}$C until OD_{600} of 0.2 and then transferred to 30 $^{\circ}$C for 2 hours. For lipid extraction, cells were pelleted and resuspended in 0.2 mL of 0.5 M NaCl in 0.1 N HCl. Lipids were extracted by first adding 0.6 mL of chloroform:methanol (1:2) to create a single-phase solution. After vigorous vortexing for 30 minutes, 0.2 mL of 0.5 M NaCl in 0.1 N HCl was added to convert the single-phase solution to a two-phase solution. After centrifugation at 20 000 x g for 5 minutes at room temperature, the lower phase was recovered and used for TLC. Approximately 2,000 cpm of phospholipid extract was subjected to TLC analysis using a HPTLC 60 plate (EMD, Gibbstown, NJ)) developed with either solvent 1: chloroform/methanol/acetic acid [60/25/10 (vol/vol/vol) in **Fig. 2A** or by solvent 2: chloroform/methanol/ammonia/water [65/37.5/3/1] (vol/vol/vol/vol) in **Fig. S4**. Radiolabeled lipids were visualized and quantified using a Phosphoimager Typhoon FLA 9500 (GE). Images were processed and quantified using ImageQuant^{TM} software for scanning and analysis. The phospholipid content is expressed as molecular percentage of the total phospholipids (correcting for the two phosphates per molecule of CL).

Phospholipid composition analysis. Whole cell lipid extracts were isolated as for TLC temperature downshift experiments without radiolabelling. IMs were isolated and extracts were prepared as has been previously described (67). Briefly, cells from temperature downshifted cultures were lysed using a French press at 8,000 psi and whole cells were pelleted by centrifugation at 30 000 x g for 20 minutes. Then, the supernatant was centrifuged at 150 000 x g for 1 hour. The resulting pellet contained pure IM as demonstrated by immunoblot analysis (**Fig. S6A**). Relative amounts of CL, PG, and PE were determined in these lipid extracts using liquid chromatography coupled to electrospray ionization mass spectrometry (LC/MS) in an API 4000 mass spectrometer (Sciex,

543 Framingham, MA, USA) using normal phase solvents as previously published (99).
544 Individual molecular species of CL, PG, and PE were expressed as a percentage within their
545 individual phospholipid by summing all the species and dividing by that sum for each
546 phospholipid. The identities of CL species were confirmed using tandem mass
547 spectrometry.

548
549

Acknowledgments

We thank members of the Mitchell Lab for productive discussions. We would like to thank
Professor Tracy Raivio (University of Alberta) for the kind gift of pJW15 plasmid. This work
was funded by the National Institute of Allergy and Infectious Disease under Award
Number R01-AI155915 (to A.M.M.), the National Institute of General Medical Sciences
under Award Number R01-GM121493 (to M.B.), and Texas A&M University start-up funds
(to A.M.M.).

558
559

References

1. Nikaido H. 2003. Molecular basis of bacterial outer membrane permeability
revisited. Microbiology and Molecular Biology Reviews 67:593-656.
2. Silhavy TJ, Kahne D, Walker S. 2010. The bacterial cell envelope. Cold Spring
Harbor Perspectives in Biology 2.
3. Henderson JC, Zimmerman SM, Crofts AA, Boll JM, Kuhns LG, Herrera CM, Trent
MS. 2016. The power of asymmetry: Architecture and assembly of the Gram-
negative outer membrane lipid bilayer. Annual Review of Microbiology 70:255-
278.
4. Berry J, Rajaure M, Pang T, Young R. 2012. The spanin complex is essential for
lambda lysis. J Bacteriol 194:5667-74.
5. Rojas ER, Billings G, Odermatt PD, Auer GK, Zhu L, Miguel A, Chang F, Weibel DB,
Theriot JA, Huang KC. 2018. The outer membrane is an essential load-bearing
element in Gram-negative bacteria. Nature 559:617-621.
6. Navarro Llorens JM, Tormo A, Martinez-Garcia E. 2010. Stationary phase in gram-
negative bacteria. FEMS Microbiol Rev 34:476-95.
7. Lugtenberg EJJ, Peters R. 1976. Distribution of lipids in cytoplasmic and outer
membranes of *Escherichia coli* K12. Biochimica et Biophysica Acta (BBA) - Lipids
and Lipid Metabolism 441:38-47.
8. Kamio Y, Nikaido H. 1976. Outer membrane of *Salmonella typhimurium*:
accessibility of phospholipid head groups to phospholipase c and cyanogen
bromide activated dextran in the external medium. Biochemistry 15:2561-70.
9. Bogdanov M, Pyrshev K, Yesylevskyy S, Ryabichko S, Boiko V, Ivanchenko P,
Kiyamova R, Guan Z, Ramseyer C, Dowhan W. 2020. Phospholipid distribution in

the cytoplasmic membrane of Gram-negative bacteria is highly asymmetric,
dynamic, and cell shape-dependent. Science Advances 6:eaaz6333.

10. Benn G, Mikheyeva IV, Inns PG, Forster JC, Ojkic N, Bortolini C, Ryadnov MG,
Kleanthous C, Silhavy TJ, Hoogenboom BW. 2021. Phase separation in the outer
membrane of *Escherichia coli*. Proceedings of the National Academy of Sciences
118:e2112237118.

11. Konovalova A, Silhavy TJ. 2015. Outer membrane lipoprotein biogenesis: Lol is
not the end. Philosophical Transactions of the Royal Society B: Biological
Sciences 370:20150030.

12. Konovalova A, Kahne DE, Silhavy TJ. 2017. Outer Membrane Biogenesis. Annu
Rev Microbiol 71:539-556.

13. Lundstedt E, Kahne D, Ruiz N. 2021. Assembly and maintenance of lipids at the
bacterial outer membrane. Chemical Reviews 121:5098-5123.

14. John E. Cronan J, Rock CO. 2008. Biosynthesis of membrane lipids. EcoSal Plus 3.

15. Jones NC, Osborn MJ. 1977. Translocation of phospholipids between the outer
and inner membranes of *Salmonella typhimurium*. Journal of Biological
Chemistry 252:7405-7412.

16. Donohue-Rolfe AM, Schaechter M. 1980. Translocation of phospholipids from
the inner to the outer membrane of *Escherichia coli*. Proceedings of the National
Academy of Sciences 77:1867-1871.

17. Raetz CR, Kantor GD, Nishijima M, Newman KF. 1979. Cardiolipin accumulation in
the inner and outer membranes of *Escherichia coli* mutants defective in
phosphatidylserine synthetase. J Bacteriol 139:544-51.

18. Langley KE, Hawrot E, Kennedy EP. 1982. Membrane assembly: movement of
phosphatidylserine between the cytoplasmic and outer membranes of
Escherichia coli. J Bacteriol 152:1033-41.

19. White DA, Lennarz WJ, Schnaitman CA. 1972. Distribution of lipids in the wall
and cytoplasmic membrane subfractions of the cell envelope of *Escherichia coli*.
Journal of Bacteriology 109:686-690.

20. Ishinaga M, Kanamoto R, Kito M. 1979. Distribution of phospholipid molecular
species in outer and cytoplasmic membranes of *Escherichia coli*. The Journal of
Biochemistry 86:161-165.

21. Davydova L, Bakholdina S, Barkina M, Velansky P, Bogdanov M, Sanina N. 2016.
Effects of elevated growth temperature and heat shock on the lipid composition
of the inner and outer membranes of *Yersinia pseudotuberculosis*. Biochimie
123:103-109.

22. Osborn MJ, Gander JE, Parisi E, Carson J. 1972. Mechanism of Assembly of the
Outer Membrane of *Salmonella* typhimurium : Isolation and characterization of
cytoplasmic and outer membrane. Journal of Biological Chemistry 247:3962-
3972.

23. Bogdanov M. 2023. Renovating a double fence with or without notifying the next
door and across the street neighbors: why the biogenic cytoplasmic membrane

of Gram-negative bacteria display asymmetry? . Emerging Topics in Life Sciences
7:137-150.

24.   Grimm J, Shi H, Wang W, Mitchell AM, Wingreen NS, Huang KC, Silhavy TJ. 2020.
The inner membrane protein YhdP modulates the rate of anterograde
phospholipid flow in *Escherichia coli*. Proc Natl Acad Sci U S A 117:26907-26914.

25.   Malinverni JC, Silhavy TJ. 2009. An ABC transport system that maintains lipid
asymmetry in the Gram-negative outer membrane. Proceedings of the National
Academy of Sciences 106:8009-8014.

26.   Sutterlin HA, Shi H, May KL, Miguel A, Khare S, Huang KC, Silhavy TJ. 2016.
Disruption of lipid homeostasis in the Gram-negative cell envelope activates a
novel cell death pathway. Proceedings of the National Academy of Sciences
113:E1565-E1574.

27.   May KL, Silhavy TJ. 2018. The *Escherichia coli* phospholipase PldA regulates outer
membrane homeostasis via lipid signaling. mBio 9:e00379-18.

28.   Tang X, Chang S, Qiao W, Luo Q, Chen Y, Jia Z, Coleman J, Zhang K, Wang T,
Zhang Z, Zhang C, Zhu X, Wei X, Dong C, Zhang X, Dong H. 2021. Structural
insights into outer membrane asymmetry maintenance in Gram-negative
bacteria by MlaFEDB. Nat Struct Mol Biol 28:81-91.

29.   Ruiz N, Davis RM, Kumar S. 2021. YhdP, TamB, and YdbH are redundant but
essential for growth and lipid homeostasis of the gram-negative outer
membrane. mBio 12:e02714-21.

30.   Douglass MV, McLean AB, Trent MS. 2022. Absence of YhdP, TamB, and YdbH
leads to defects in glycerophospholipid transport and cell morphology in Gram-
negative bacteria. PLOS Genetics 18:e1010096.

31.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O,
Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl
SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T,
Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M,
Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K,
Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with
AlphaFold. Nature 596:583-589.

32.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D,
Stroe O, Wood G, Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S,
Jumper J, Clancy E, Green R, Vora A, Lutfi M, Figurnov M, Cowie A, Hobbs N,
Kohli P, Kleywegt G, Birney E, Hassabis D, Velankar S. 2021. AlphaFold Protein
Structure Database: massively expanding the structural coverage of protein-
sequence space with high-accuracy models. Nucleic Acids Research 50:D439-
D444.

33.   Josts I, Stubenrauch CJ, Vadlamani G, Mosbahi K, Walker D, Lithgow T, Grinter R.
2017. The structure of a conserved domain of TamB reveals a hydrophobic β taco
fold. Structure 25:1898-1906.e5.

34.   Kumar N, Leonzino M, Hancock-Cerutti W, Horenkamp FA, Li P, Lees JA, Wheeler
668   H, Reinisch KM, De Camilli P. 2018. VPS13A and VPS13C are lipid transport
669   proteins differentially localized at ER contact sites. Journal of Cell Biology
670   217:3625-3639.
671   35.   Valverde DP, Yu S, Boggavarapu V, Kumar N, Lees JA, Walz T, Reinisch KM, Melia
672   TJ. 2019. ATG2 transports lipids to promote autophagosome biogenesis. J Cell
673   Biol 218:1787-1798.
674   36.   Otomo T, Maeda S. 2019. ATG2A transfers lipids between membranes *in vitro*.
675   Autophagy 15:2031-2032.
676   37.   Osawa T, Kotani T, Kawaoka T, Hirata E, Suzuki K, Nakatogawa H, Ohsumi Y, Noda
677   NN. 2019. Atg2 mediates direct lipid transfer between membranes for
678   autophagosome formation. Nat Struct Mol Biol 26:281-288.
679   38.   Grasekamp KP, Beaud B, Taib N, Audrain B, Bardiaux B, Rossez Y, Izadi-Pruneyre
680   N, Lejeune M, Trivelli X, Chouit Z, Guerardel Y, Ghigo J-M, Gribaldo S, Beloin C.
681   2023. Bridges instead of boats? The Mla system of diderm Firmicute *Veillonella*
682   *parvula* reveals an ancestral transenvelope core of phospholipid trafficking.
683   bioRxiv doi:10.1101/2023.06.30.547184:2023.06.30.547184.
684   39.   Mitchell AM, Wang W, Silhavy TJ. 2017. Novel Rpos-dependent mechanisms
685   strengthen the envelope permeability barrier during stationary phase. J Bacteriol
686   199:e00708-16.
687   40.   Mitchell AM, Srikumar T, Silhavy TJ. 2018. Cyclic enterobacterial common
688   antigen maintains the outer membrane permeability barrier of *Escherichia coli* in
689   a manner controlled by Yhdp. mBio 9:e01321-18.
690   41.   Cronan JE. 2021. The *Escherichia coli* FadR transcription factor: Too much of a
691   good thing? Molecular Microbiology 115:1080-1085.
692   42.   Nunn WD, Giffin K, Clark D, Cronan JE. 1983. Role for *fadR* in unsaturated fatty
693   acid biosynthesis in *Escherichia coli*. Journal of Bacteriology 154:554-560.
694   43.   Rai AK, Mitchell AM. 2020. Enterobacterial common antigen: Synthesis and
695   function of an enigmatic molecule. mBio 11:e01914-20.
696   44.   Paradis-Bleau C, Kritikos G, Orlova K, Typas A, Bernhardt TG. 2014. A genome-
697   wide screen for bacterial envelope biogenesis mutants identifies a novel factor
698   involved in cell wall precursor metabolism. PLoS Genet 10:e1004056.
699   45.   Ruiz N, Falcone B, Kahne D, Silhavy TJ. 2005. Chemical conditionality: a genetic
700   strategy to probe organelle assembly. Cell 121:307-17.
701   46.   Nikaido H, Vaara M. 1985. Molecular basis of bacterial outer membrane
702   permeability. Microbiol Rev 49:1-32.
703   47.   Bishop RE. 2008. Structural biology of membrane-intrinsic β-barrel enzymes:
704   Sentinels of the bacterial outer membrane. Biochimica et Biophysica Acta (BBA) -
705   Biomembranes 1778:1881-1896.
706   48.   Rowlett VW, Mallampalli V, Karlstaedt A, Dowhan W, Taegtmeyer H, Margolin W,
707   Vitrac H. 2017. Impact of membrane phospholipid slterations in *Escherichia coli*
708   on cellular function and bacterial stress adaptation. J Bacteriol 199.

49. Tan BK, Bogdanov M, Zhao J, Dowhan W, Raetz CRH, Guan Z. 2012. Discovery of
a cardiolipin synthase utilizing phosphatidylethanolamine and
phosphatidylglycerol as substrates. Proceedings of the National Academy of
Sciences 109:16504-16509.
50. Pluschke G, Hirota Y, Overath P. 1978. Function of phospholipids in *Escherichia*
coli. Characterization of a mutant deficient in cardiolipin synthesis. Journal of
Biological Chemistry 253:5048-5055.
51. Nishijima S, Asami Y, Uetake N, Yamagoe S, Ohta A, Shibuya I. 1988. Disruption of
the *Escherichia coli cls* gene responsible for cardiolipin synthesis. Journal of
Bacteriology 170:775-780.
52. Guo D, Tropp BE. 2000. A second *Escherichia coli* protein with CL synthase
activity. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of
Lipids 1483:263-274.
53. Kikuchi S, Shibuya I, Matsumoto K. 2000. Viability of an *Escherichia coli pgsA* null
mutant lacking detectable phosphatidylglycerol and cardiolipin. Journal of
Bacteriology 182:371-376.
54. Suzuki M, Hara H, Matsumoto K. 2002. Envelope disorder of *Escherichia coli* cells
lacking phosphatidylglycerol. Journal of Bacteriology 184:5418-5425.
55. Shiba Y, Yokoyama Y, Aono Y, Kiuchi T, Kusaka J, Matsumoto K, Hara H. 2004.
Activation of the Rcs signal transduction system is responsible for the
thermosensitive growth defect of an *Escherichia coli* mutant lacking
phosphatidylglycerol and cardiolipin. J Bacteriol 186:6526-35.
56. Grabowicz M, Silhavy TJ. 2017. Redefining the essential trafficking pathway for
outer membrane lipoproteins. Proceedings of the National Academy of Sciences
114:4769-4774.
57. Cronan JE, Jr., Gelmann EP. 1973. An estimate of the minimum amount of
unsaturated fatty acid required for growth of *Escherichia coli*. J Biol Chem
248:1188-95.
58. Feng Y, Cronan JE. 2011. Complex binding of the FabR repressor of bacterial
unsaturated fatty acid biosynthesis to its cognate promoters. Molecular
Microbiology 80:195-218.
59. Feng Y, Cronan JE. 2009. *Escherichia coli* unsaturated fatty acid synthesis:
Complex transcription of the *fabA* gene and *in vivo* identification of the essential
reaction catalyzed by FabB. Journal of Biological Chemistry 284:29526-29535.
60. Henry MF, Cronan JE. 1991. *Escherichia coli* transcription factor that both
activates fatty acid synthesis and represses fatty acid degradation. Journal of
Molecular Biology 222:843-849.
61. Henry MF, Cronan JE. 1992. A new mechanism of transcriptional regulation:
Release of an activator triggered by small molecule binding. Cell 70:671-679.
62. Rock CO, Jackowski S. 1985. Pathways for the incorporation of exogenous fatty
acids into phosphatidylethanolamine in *Escherichia coli*. J Biol Chem 260:12720-
4.

63.   Français M, Bott R, Dargaignaratz C, Giniès C, Carlin F, Broussolle V, Nguyen-Thé
752   C. 2021. Short-Chain and Unsaturated Fatty Acids Increase Sequentially From the
753   Lag Phase During Cold Growth of *Bacillus cereus*. Frontiers in Microbiology 12.
754   64.   Heath RJ, Rock CO. 1996. Roles of the FabA and FabZ β-hydroxyacyl-acyl carrier
755   protein dehydratases in *Escherichia coli* fatty acid biosynthesis. Journal of
756   Biological Chemistry 271:27795-27801.
757   65.   Ostrander DB, Sparagna GC, Amoscato AA, McMillin JB, Dowhan W. 2001.
758   Decreased Cardiolipin Synthesis Corresponds with Cytochromec Release in
759   Palmitate-induced Cardiomyocyte Apoptosis. Journal of Biological Chemistry
760   276:38061-38067.
761   66.   Lewis RNAH, Zweytick D, Pabst G, Lohner K, McElhaney RN. 2007. Calorimetric,
762   X-Ray Diffraction, and Spectroscopic Studies of the Thermotropic Phase Behavior
763   and Organization of Tetramyristoyl Cardiolipin Membranes. Biophysical Journal
764   92:3166-3177.
765   67.   Bogdanov M, Sun J, Kaback HR, Dowhan W. 1996. A Phospholipid Acts as a
766   Chaperone in Assembly of a Membrane Transport Protein. Journal of Biological
767   Chemistry 271:11615-11618.
768   68.   Mitchell AM, Silhavy TJ. 2019. Envelope stress responses: balancing damage
769   repair and toxicity. Nature Reviews Microbiology 17:417-428.
770   69.   Shibata A, Ikawa K, Shimooka T, Terada H. 1994. Significant stabilization of the
771   phosphatidylcholine bilayer structure by incorporation of small amounts of
772   cardiolipin. Biochimica et Biophysica Acta (BBA) - Biomembranes 1192:71-78.
773   70.   Boscia AL, Treece BW, Mohammadyani D, Klein-Seetharaman J, Braun AR,
774   Wassenaar TA, Klösgen B, Tristram-Nagle S. 2014. X-ray structure,
775   thermodynamics, elastic properties and MD simulations of
776   cardiolipin/dimyristoylphosphatidylcholine mixed membranes. Chemistry and
777   Physics of Lipids 178:1-10.
778   71.   Pennington ER, Fix A, Sullivan EM, Brown DA, Kennedy A, Shaikh SR. 2017.
779   Distinct membrane properties are differentially influenced by cardiolipin content
780   and acyl chain composition in biomimetic membranes. Biochimica et Biophysica
781   Acta (BBA) - Biomembranes 1859:257-267.
782   72.   Lopes SC, Ivanova G, de Castro B, Gameiro P. 2018. Revealing cardiolipins
783   influence in the construction of a significant mitochondrial membrane model.
784   Biochimica et Biophysica Acta (BBA) - Biomembranes 1860:2465-2477.
785   73.   Marr AG, Ingraham JL. 1962. Effect of temperature on the composition of fatty
786   acids in *Escherichia coli*. Journal of Bacteriology 84:1260-1267.
787   74.   Garwin JL, Klages AL, Cronan JE. 1980. Beta-ketoacyl-acyl carrier protein
788   synthase II of *Escherichia coli*. Evidence for function in the thermal regulation of
789   fatty acid synthesis. Journal of Biological Chemistry 255:3263-3265.
790   75.   de Mendoza D, Klages Ulrich A, Cronan JE. 1983. Thermal regulation of
791   membrane fluidity in *Escherichia coli*. Effects of overproduction of beta-ketoacyl-
792   acyl carrier protein synthase I. Journal of Biological Chemistry 258:2098-2101.

793   76.   Knapp BD, Huang KC. 2022. The effects of temperature on cellular physiology.
794   Annual Review of Biophysics 51:499-526.
795   77.   Bogdanov M, Umeda M, Dowhan W. 1999. Phospholipid-assisted refolding of an
796   integral membrane protein: Minimum structural features for
797   phosphatidylethanolamine to act as a molecular chaperone. Journal of Biological
798   Chemistry 274:12339-12345.
799   78.   Raetz CR, Whitfield C. 2002. Lipopolysaccharide endotoxins. Annu Rev Biochem
800   71:635-700.
801   79.   Conway T, Creecy JP, Maddox SM, Grissom JE, Conkle TL, Shadid TM, Teramoto J,
802   Miguel PS, Shimada T, Ishihama A, Mori H, Wanner BL. 2014. Unprecedented
803   high-resolution view of bacterial operon architecture revealed by RNA
804   sequencing. mBio 5:e01442-14.
805   80.   Thomason MK, Bischler T, Eisenbart SK, Förstner KU, Zhang A, Herbig A, Nieselt
806   K, Sharma CM, Storz G. 2015. Global transcriptional start site mapping using
807   differential RNA sequencing reveals novel antisense rnas in *Escherichia coli*.
808   Journal of Bacteriology 197:18-28.
809   81.   Adams PP, Baniulyte G, Esnault C, Chegireddy K, Singh N, Monge M, Dale RK,
810   Storz G, Wade JT. 2021. Regulatory roles of *Escherichia coli* 5' UTR and ORF-
811   internal RNAs detected by 3' end mapping. eLife 10:e62438.
812   82.   Karp PD, Ong WK, Paley S, Billington R, Caspi R, Fulcher C, Kothari A,
813   Krummenacker M, Latendresse M, Midford PE, Subhraveti P, Gama-Castro S,
814   Muñiz-Rascado L, Bonavides-Martinez C, Santos-Zavaleta A, Mackie A, Collado-
815   Vides J, Keseler IM, Paulsen I. 2018. The EcoCyc database. EcoSal Plus
816   8:10.1128/ecosalplus.ESP-0006-2018.
817   83.   Huerta AM, Collado-Vides J. 2003. Sigma70 promoters in *Escherichia coli*: specific
818   transcription in dense regions of overlapping promoter-like signals. Journal of
819   Molecular Biology 333:261-278.
820   84.   Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, Doncheva NT,
821   Legeay M, Fang T, Bork P, Jensen LJ, von Mering C. 2021. The STRING database in
822   2021: customizable protein-protein networks, and functional characterization of
823   user-uploaded gene/measurement sets. Nucleic Acids Res 49:D605-d612.
824   85.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita
825   M, Wanner BL, Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame,
826   single-gene knockout mutants: the Keio collection. Mol Syst Biol 2:2006.0008-
827   2006.0008.
828   86.   Silhavy TJ, Berman ML, Enquist LW. 1984. Experiments with gene fusions. Cold
829   Spring Harbor Laboratory.
830   87.   Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in
831   *Escherichia coli* K-12 using PCR products. Proceedings of the National Academy
832   of Sciences 97:6640-6645.
833   88.   MacRitchie DM, Ward JD, Nevesinjac AZ, Raivio TL. 2008. Activation of the Cpx
834   envelope stress response down-regulates expression of several locus of

835   enterocyte effacement-encoded genes in enteropathogenic *Escherichia coli*.
836   Infection and Immunity 76:1465-1475.

837 89. Wong JL, Vogt SL, Raivio TL. 2013. Using reporter genes and the *Escherichia coli*
838   ASKA overexpression library in screens for regulators of the Gram negative
839   envelope stress response. Methods Mol Biol 966:337-57.

840 90. Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-
841   evolved microbes from next-generation sequencing data using breseq. Methods
842   Mol Biol 1151:165-88.

843 91. Bailey TL, Elkan C. 1994. Fitting a mixture model by expectation maximization to
844   discover motifs in biopolymers. Proc Int Conf Intell Syst Mol Biol 2:28-36.

845 92. Bailey TL, Johnson J, Grant CE, Noble WS. 2015. The MEME Suite. Nucleic Acids
846   Res 43:W39-49.

847 93. Singer M, Baker TA, Schnitzler G, Deischel SM, Goel M, Dove W, Jaacks KJ,
848   Grossman AD, Erickson JW, Gross CA. 1989. A collection of strains containing
849   genetically linked alternating antibiotic resistance elements for genetic mapping
850   of Escherichia coli. Microbiol Rev 53:1-24.

851 94. Rai AK, Carr JF, Bautista DE, Wang W, Mitchell AM. 2021. ElyC and cyclic
852   enterobacterial common antigen regulate synthesis of phosphoglyceride-linked
853   enterobacterial common antigen. mBio 12:e02846-21.

854 95. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome
855   alignment and genotyping with HISAT2 and HISAT-genotype. Nature
856   Biotechnology 37:907-915.

857 96. Liao Y, Smyth GK, Shi W. 2013. featureCounts: an efficient general purpose
858   program for assigning sequence reads to genomic features. Bioinformatics
859   30:923-930.

860 97. Robinson MD, McCarthy DJ, Smyth GK. 2009. edgeR: a Bioconductor package for
861   differential expression analysis of digital gene expression data. Bioinformatics
862   26:139-140.

863 98. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. limma
864   powers differential expression analyses for RNA-sequencing and microarray
865   studies. Nucleic Acids Research 43:e47-e47.

866 99. Sparagna GC, Johnson CA, McCune SA, Moore RL, Murphy RC. 2005.
867   Quantitation of cardiolipin molecular species in spontaneously hypertensive
868   heart failure rats using electrospray ionization mass spectrometry. J Lipid Res
869   46:1196-204.

870
871

   Figures and Tables

Figure 1. Deletion of *yhdP* and *fadR* causes synthetic cold sensitivity due to impairment
of TamB function. (A) The Δ*yhdP* Δ*fadR* strain had impaired growth at lower
temperatures. **(B)** EOPs were performed to confirm the growth defect of the Δ*yhdP* Δ*fadR*
strain. **(C-F)** EOPs were performed at 30 °C to assess cold sensitivity. **(C)** The growth of the
Δ*yhdP* Δ*fadR* strain was not affected by the loss of ECA (Δ*wecA*). **(D)** Neither a Δ*tamB*
Δ*fadR* strain nor a Δ*ydbH* Δ*fadR* strain exhibited cold sensitivity. **(E)** Deletion of *ydbH* did
not suppress cold sensitivity in the Δ*yhdP* Δ*fadR* strain. **(F)** Deletion of *tamB* completely
suppressed cold sensitivity in the Δ*yhdP* Δ*fadR* strain and the Rcs stress response was not
necessary for this suppression. **(G)** Growth curves were performed at 30 °C . The Δ*tamB*
and the Δ*yhdP* Δ*fadR* Δ*tamB* strain grew equally well, indicating suppression is unlikely to
result from slow growth. Quantitative data are shown as the mean of three biological
replicates ± the SEM. Images are representative of three independent experiments.

888
Figure 2. Increased cardiolipin levels are necessary for cold sensitivity of the Δ*yhdP*
Δ*fadR* strain. (A) Thin layer chromatography (TLC) of lipid extracts was performed for the
indicated cultures grown to OD_{600}=0.2 at 42 °C then transferred to 30 °C or 42 °C for 2
hours before analysis. Relative phospholipid levels are provided for 30 °C samples. The
Δ*yhdP* Δ*fadR* strain shows increased levels of CL and a concomitant decrease in PG levels.

(B-F) Growth effects of altered phospholipid levels were investigated by EOP. Images are
representative of three independent experiments. **(B)** PgsA depletion strains in a
background where *pgsA* is non-essential were used to assay the effect of lowering PG and
CL levels on cold sensitivity. Arabinose induces expression of *pgsA* while glucose
represses. Lowering PG and CL levels partially suppressed cold sensitivity. **(C)** Deletion of
the three CL synthases, *clsA*, *clsB*, and *clsC*, suppressed cold sensitivity of the Δ*yhdP* Δ*fadR*
strain. **(D)** Deletion of *clsA* also suppressed cold sensitivity. **(E)** Deletion of *clsB* does not
suppress cold sensitivity, while deletion of *clsC* (F) does.
902
903

Figure 3. Cold sensitivity can be suppressed by increasing *fabA* expression. (A) The cold
sensitivity of a spontaneous suppressor of Δ*yhdP* Δ*fadR* cold sensitivity was tested by EOP.
The suppressor returned growth to the level of the Δ*fadR* mutant. **(B)** TLC was performed
to assay phospholipid composition. Suppressor 1 showed very similar phospholipid
composition to that of its parent strain and did not decrease CL levels. **(C)** A diagram of
the promoter region of *fabA* is shown with its transcriptional start sites (blue arrows) and
regulator binding sites (boxes). FadR acts as an activator while FabR acts as a repressor.
The motif of the FabR binding site, generated using MEME, is shown with the sequences
of the parent strain and of Suppressor 1. **(D)** The P$_{fabA}$(-19G>A) mutation was linked to a
Tn*10* and transferred to the Δ*yhdP* Δ*fadR* strain by transduction. The Δ*yhdP* Δ*fadR* strain
with P$_{fabA}$(-19G>A) was not cold sensitivity, confirming suppression. **(E)** *fabA* RNA levels
are shown. Bars are the mean of three biological replicates ± the SEM. Individual data
points are shown as open circles. ** $p<0.005$, *** $p<0.0005$ by quasi-linear F-test. Data
are from the RNA-seq experiment in Figure 5 and *fabA* data are also shown in Figure 5F.
(F) Luciferase reporters of wild-type and mutant *fabA* promoter activity were constructed.
The mutation caused similar fold increases in P$_{fabA}$ activity in all strain backgrounds. Data
are luminescence relative to OD$_{600}$ and are shown as the average of two biological
replicates ± the SEM.

Figure 4. Decreasing fatty acid saturation suppresses Δ*yhdP* Δ*fadR* cold sensitivity. (A-
C) Lipid extracts were made from whole cells or from IM separated from OM by
differential centrifugation. Strains were grown to OD_{600} 0.2 at 42 °C then shifted to 30 °C
for 2 hours before harvest. Lipid extracts were quantified using LC/MS and the percentage
of each specific species of PE (A), PG (B), and CL (C) was calculated using the sum of all
molecular species detected. Suppressor 1 showed decreased saturation compared to its
parent strain (Δ*yhdP* Δ*fadR*). For PE, this difference was evident for whole cells but not
for IM. Data for phospholipids without unsaturations are shown in grey, with one
unsaturation in blue, with two unsaturations in green, with three unsaturations in orange,
and odd chain fatty acids (likely cyclopropyl-fatty acids) in yellow. Data for phospholipids
with longer acyl chains are shown in darker colors. **(D)** EOPS were performed on media
supplemented with oleic acid or a vehicle control. Oleic acid suppresses the cold
sensitivity of the Δ*yhdP* Δ*fadR* strain. **(E)** Growth curves of the indicated strains were
performed in the presence of oleic acid, stearic acid, or a vehicle control. Oleic acid
suppressed the cold sensitivity of the Δ*yhdP* Δ*fadR* strain while stearic acid did not. Data
are the average of three biological replicates ± the SEM.
941

Figure 5. Transcriptional landscape of the Δ*yhdP* Δ*fadR* and suppressed strain. Three biological replicates of the indicated strains were grown at 42 °C to mid-log then transferred to 30 °C for 30 minutes before harvesting RNA and performing RNA-seq. Differential expression was calculated based on a greater than 2-fold change in expression and a p value of less than 0.05 by quasi-linear F-test. **(A)** PCA was performed on the expression of all genes differentially expressed between any groups of samples. Suppressor 1 and the Δ*yhdP* Δ*fadR* strain grouped closer together than to the wild-type strain, although Suppressor 1 was closer to wild type than the Δ*yhdP* Δ*fadR* strain. **(B-D)** Volcano plots show the average fold change and significance of expression changes between the Δ*yhdP* Δ*fadR* strain and wild-type MG1655 (B), Suppressor 1 and wild-type MG1655 (C), and Suppressor 1 and the Δ*yhdP* Δ*fadR* strain (D). Genes with changes greater than 2-fold are shown in blue, while genes with changes less than 2-fold are shown in grey. Genes with a q-value (false discovery rate) of less than 0.05 are shown in yellow. Blue lines indicate a 2-fold change and dotted line indicates p>0.05. The number of up or down regulated genes is indicated in blue. The names of the 10 genes with the largest changes are called out. More genes were up-regulated then down regulated in the Δ*yhdP* Δ*fadR* strain and Suppressor 1 vs. wild type. More genes were down regulated in

Suppressor 1 compared to the Δ*yhdP* Δ*fadR* strain. **(E)** Relative expression of genes in the
CL synthesis pathway is shown as averages ± the SEM and individual data points. No
significant changes were evident. **(F)** Relative expression of genes in the fatty acid
synthesis and degradation pathways is shown as averages ± the SEM and individual data
points. A general normalizing of these pathways occurred in Suppressor 1 compared to
the Δ*yhdP* Δ*fadR* strain; however, the most significant change was in the expression of
fabA. * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$ by quasi-linear F-test.
967

Figure 6. Model for the role of YhdP and TamB in phospholipid transport. Our data can
be explained by the following model of phospholipid transport. **(A)** In normal conditions,
YhdP is mainly responsible for the transport of phospholipids with more saturated fatty
acid chains between the IM and OM. TamB shows a strong preference for transporting
phospholipids with more unsaturated fatty acids, with YdbH playing a minor role in
phospholipid transport. **(B)** In the Δ*yhdP* Δ*fadR* strain, YhdP is not available to transport
more saturated phospholipids and the proportion of both saturated fatty acids and
cardiolipin is increased, especially at low temperature. The fluidity of the membrane is
also lessened by the low temperature. In these conditions, transport of phospholipids
between the membranes is impeded, due to clogging of TamB by a combination of less
saturated phospholipids and/or bulky and highly disordered CL. This leads to a lethal lack
of phospholipids in the OM and overabundance of phospholipids in the IM. **(C)** When
tamB is deleted in the Δ*yhdP* Δ*fadR* strain, the clogging of TamB is relieved, and the cell
relies on YdbH for phospholipid transport overcoming the cold sensitivity. PL:
phospholipid