1

Functional diversification of gram-negative intermembrane phospholipid transporter function by intrinsic substrate preference

Ashutosh K. Rai[a], Katsuhiro Sawasato[b], Haley C. Bennett[a], Anastasiia Kozlova[b],
Genevieve C. Sparagna[c], Mikhail Bogdanov[b], and Angela M. Mitchell[a,#]

[a] Department of Biology, Texas A&M University, College Station, Texas, USA

[b] Department of Biochemistry and Molecular Biology, McGovern Medical School at The
University of Texas Health Science Center at Houston, Houston, Texas, USA

[c] Department of Medicine, Division of Cardiology, University of Colorado Anschutz
Medical Campus, Aurora, Colorado, USA

Running title: Phospholipid transporter substrate preference

Address correspondence to Angela M. Mitchell, amitchell@bio.tamu.edu

Keywords: phospholipid trafficking, gram-negative envelope, outer membrane, lipid
metabolism, cold sensitivity

Abstract

The outer membrane of Gram-negative bacteria is a barrier to chemical and physical stress. Phospholipid transport between the inner and outer membranes has been an area of intense investigation and, in *E. coli* K-12, it has recently been shown to be mediated by YhdP, TamB, and YdbH, which are suggested to provide hydrophobic channels for phospholipid diffusion, with YhdP and TamB playing the major roles. However, YhdP and TamB have different phenotypes suggesting distinct functions. We investigated these functions using synthetic cold sensitivity (at 30 °C) of a strain with deletion of *yhdP*, but not *tamB* or *ydbH*, and *fadR*, a transcriptional regulator controlling fatty acid degradation and unsaturated fatty acid production. Deletion of *tamB*, redirecting phospholipid transport to YdbH, suppresses the Δ*yhdP* Δ*fadR* cold sensitivity suggesting this phenotype is directly related to phospholipid transport. The Δ*yhdP* Δ*fadR* strain shows a greater increase in cardiolipin upon transfer to the non-permissive temperature and genetically lowering cardiolipin levels can suppress cold sensitivity. These data also reveal a qualitative difference between cardiolipin synthases in *E. coli*, as deletion of *clsA and clsC* suppresses cold sensitivity but deletion of *clsB* does not despite lower cardiolipin levels. In addition to increased cardiolipin, increased fatty acid saturation is necessary for cold sensitivity and lowering this level genetically or through supplementation of oleic acid suppresses the cold sensitivity of the Δ*yhdP* Δ*fadR* strain. A parsimonious explanation for our data is that YhdP and TamB have differential substrate transport preferences, most likely with YhdP preferentially transporting more saturated phospholipids and TamB preferentially transporting more unsaturated phospholipids. We envision cardiolipin contributing to this transport preference by sterically clogging TamB-mediated transport of saturated phospholipids. Thus, our data provide a potential mechanism for independent control of the phospholipid composition of the inner and outer membranes in response to changing conditions.

Author Summary

Gram-negative bacteria possess an impermeable outer membrane, which protects against environmental stress and antibiotics. Outer membrane phospholipid transport remained mysterious until recently when YhdP, TamB, and YdbH were implicated in phospholipid transport between the inner and outer membranes of *E. coli*. Similar roles for YhdP and/or TamB have been suggested in both closely and distantly related gram-negative bacteria. Here, given the transporters' apparent partial redundancy, we investigated functional differentiation between YhdP and TamB. Our data demonstrate that YhdP and TamB have differential involvement with fatty acid and phospholipid metabolism. In fact, transport of higher than normal levels of cardiolipin and saturated phospholipids by TamB at a non-permissive temperature is lethal. From these data, we suggest a model where the functions of YhdP and TamB are distinguished by their preference for phospholipid transport with YhdP preferentially transporting more saturated phospholipids and TamB transporting more unsaturated phospholipids. Cardiolipin headgroup specificity may contribute transport inhibition due to its bulky nature inhibiting the passage of other phospholipids. Diversification of function between YhdP and TamB provides a mechanism for regulation of phospholipid composition, and possibly the mechanical strength and permeability of the outer membrane, and so the cell's intrinsic antibiotic resistance, in changing environmental conditions.

Introduction

The gram-negative bacterial cell envelope has an outer membrane (OM) that sits outside the aqueous periplasm and peptidoglycan cell wall. The OM provides a barrier against various environmental stresses including toxic molecules such as antibiotics and osmotic pressure (1-6). Unlike the inner membrane (IM), a phospholipid bilayer, the OM is largely composed of phospholipids (mainly phosphatidylethanolamine (PE) (7)) in its inner leaflet and LPS (lipopolysaccharide) in its outer leaflet (2, 8). However, both membranes are asymmetric as the IM has different phospholipid compositions in its inner versus outer leaflets (9). Outer membrane proteins (OMPs) form a network across the cell surface interspersed with phase separated LPS patches (10). Lipoproteins are generally anchored in the inner (i.e., periplasmic) leaflet of the OM (11).

OM components are synthesized in the cytoplasm or IM and transported to the OM. OMP (12), lipoprotein (11), and LPS (13) transport pathways are well defined. However, until recently, intermembrane phospholipid transport (between the IM and OM), especially anterograde transport from the IM to the OM, has remained very poorly understood. Phospholipids are synthesized at the IM's inner leaflet (14) and rapid, bidirectional intermembrane phospholipid transport occurs (15), even without ATP (16). Phospholipid transport has a relaxed specificity, allowing transport of non-native lipids (17, 18); however, intermembrane phospholipid composition differences are maintained (7, 19-22). For instance, the OM is enriched for PE and its saturated species compared to the IM outer leaflet (7, 9, 20-22); however, PG and CL can be evenly distributed between the IM and OM when the cellular level of PE is greatly reduced (17), suggesting a maintenance of membrane charge balance. This headgroup redistribution was accompanied by a concomitant fatty acid redistribution evidenced by an accumulation of palmitic acid (C16:0) and cyclopropane derivatives of palmitoleic acid (C16:1) and *cis*-vaccenic acid (C18:1). In contrast, the 1,2-diglyceride that accumulates in mutants lacking DgkA (diglyceride kinase) predominantly associates with the IM (23), suggesting existence of a mechanism of discrimination of lipid transfer to the OM that does not recognize diglyceride molecules and contributes to the diversification of distribution of polar and acyl groups between the IM and OM. This establishment and maintenance of defined lipid topography between the IM and the OM must involve the coordinated biosynthesis and balanced bidirectional trafficking of specific phospholipids across the IM and from the IM to the OM (24).

It has been demonstrated that anterograde phospholipid transport is mediated by a high-flux, diffusive system in *Escherichia coli* (25). A retrograde trafficking pathway, the Mla system, removes mislocalized phospholipids from the OM outer leaflet and returns them to the IM (26). A gain-of-function allele, *mlaA** (27, 28), opens a channel allowing phospholipids to mislocalize to the cell surface by flowing from the OM's inner to outer leaflet (27, 29), resulting in a PldA (OM phospholipase A)-mediated attempt to reestablish OM asymmetry by increasing LPS production, OM vesiculation, and increased flow of

phospholipids to the OM (27, 28). When nutrients are depleted, inhibiting phospholipid
production, cells lyse due to loss of IM integrity (16, 18). Single cell imaging of *mlaA** cells
confirmed an aberrant lipid flow from the IM to the OM at a fast rate. Loss of YhdP, an
inner membrane protein, could significantly slow phospholipid flow to the OM resulting
in loss of OM integrity before IM rupture (25), suggesting YhdP might play a role in
intermembrane phospholipid transport.

Recently, YhdP and two homologs, TamB and YdbH, were demonstrated to be
intermembrane phospholipid transporters (30, 31). Ruiz, *et al*. investigated AsmA family
proteins and found Δ*yhdP* Δ*tamB* mutants showed synthetic OM permeability and
stationary phase lysis (30). Moreover, when combined, Δ*yhdP*, Δ*tamB*, and Δ*ydbH* are
synthetically lethal. Genetic interactions of these mutants with *mlaA* and *pldA* show their
involvement in OM phospholipid homeostasis. Predicted structures of YhdP, TamB, and
YdbH (32, 33), as well as a partial crystal structure of TamB (34), have a β-taco fold domain
with a hydrophobic pocket, resembling eukaryotic lipid transporters (35-38). These
structural findings are another clue that suggests YhdP, TamB, and YdbH are involved in
phospholipid transport. Douglass, *et al.* confirmed the synthetic lethality of Δ*yhdP*,
Δ*tamB*, and Δ*ydbH* and demonstrated a Δ*tamB* Δ*ydbH* mutant with reduced *yhdP*
expression had decreased amounts of OM phospholipids, directly demonstrating their
involvement in intermembrane phospholipid transport or its regulation (31). Structural
studies have shown that YhdP is long enough to span the periplasmic space and molecular
dynamics indicate that the C-terminus of YhdP can insert into the OM to allow
phospholipid transfer between the membranes (39). This study also directly
demonstrates that a phosphate containing substrate, putatively assigned to be
phospholipids, can be crosslinked to the hydrophobic groove of YhdP (39). The role of
YhdP, TamB, and YdbH in phospholipid transport may be widely conserved as recent
evidence implicates TamB in anterograde phospholipid transport in *Veillonella parvula*, a
diderm Fermicute (40) and YhdP, TamB, YdbH, and PA4735 in intermembrane
phospholipid transport in *Pseudomonas aeruginosa* (41).

Nevertheless, why there are three separate intermembrane phospholipid transport
proteins and the functional interactions between these proteins remains unanswered,
although it is clear the proteins are not fully redundant or functionally equivalent. The
conditional expression of *yhdP* alone, without the presence of *ydbH* and *tamB*, fully
complements neither growth phenotypes nor a normal level of OM phospholipids,
suggesting transport function is still impaired in this strain (31). In addition, YdbH seems
to play a more minor role than YhdP and TamB (30, 31) and the screen identifying Δ*yhdP*
as slowing *mlaA**-dependent lysis did not identify Δ*tamB* or Δ*ydbH* (25). Moreover, we
previously identified a role for YhdP in stationary phase SDS (sodium dodecyl sulfate)
resistance and modulating cyclic enterobacterial common antigen activity (42, 43) not
shared by TamB and YdbH (**Fig. S1**) (30, 31, 42, 43). Similarly, TamB, in conjunction with

TamA, has been suggested to play a role in OM insertion of some "complicated" β-barrel
OMPs such as autotransporters and usher proteins (34, 44-46).

Here, we investigated the differentiation of YhdP and TamB function and identified
synthetic cold sensitivity in a strain with Δ*yhdP* and Δ*fadR*, a transcriptional regulator of
fatty acid biosynthesis and degradation, acting as a transcriptional switch between these
pathways in response to acyl-CoA (47). FadR is necessary for normal levels of unsaturated
fatty acids to be synthesized (48). The cold sensitivity was unique to the Δ*yhdP* Δ*fadR*
strain and was suppressed by loss of TamB, demonstrating that the sensitivity is related
to phospholipid transport. We found that the Δ*yhdP* Δ*fadR* mutant had a greater increase
in levels of a specific phospholipid, cardiolipin (CL) during growth at the non-permissive
temperature, and decreasing CL suppressed cold sensitivity. Furthermore, increasing
unsaturated fatty acid levels suppressed cold sensitivity. Overall, our data are consistent
with a model where TamB and YhdP transport functions are differentiated based on
phospholipid fatty acid saturation state, with TamB transporting more unsaturated
phospholipids and YhdP transporting more saturated phospholipids. Phospholipid species
or head group can also contribute to transport specificity, since higher levels of dianionic
CL are necessary for the cold sensitivity phenotype, potentially due to functional
inhibition of TamB by clogging of its hydrophobic groove.

170
171

Results

Loss of FadR and YhdP results in cold sensitivity suppressed by loss of TamB. To
investigate YhdP-TamB function differentiation, we created a Δ*yhdP* Δ*fadR* strain, as FadR
is a major regulator of lipid homeostasis (47). A deletion strain of *fadR* alone is expected
to have decreased fatty acid synthesis, increased fatty acid degradation, and increased
fatty acid saturation (47, 48). Surprisingly, Δ*yhdP* Δ*fadR* cultures grown at 30 °C lagged for
more than 8 hours, followed by highly variable growth, suggesting suppressor outgrowth
(**Fig.1A**). The growth defect was decreased at 37 °C and completely absent at 42 °C. To
confirm this cold sensitivity, we estimated efficiency of plating (EOP) and confirmed the
severe cold intolerance of the Δ*yhdP* Δ*fadR* strain (5 logs at 30 °C and 3 logs at 37 °C) (**Fig.
1B**). Although, enterobacterial common antigen is necessary for Δ*yhdP*'s OM permeability
phenotypes (43), the Δ*yhdP* Δ*fadR* strain's cold sensitivity was unchanged when
enterobacterial common antigen was not present (Δ*wecA*) (49) (**Fig. 1C**).

We hypothesized the Δ*yhdP* Δ*fadR* strain's cold sensitivity was due to impairment of
phospholipid transport and expected Δ*tamB* and/or Δ*ydbH* to show similar synthetic
phenotypes with Δ*fadR*. However, Δ*fadR* Δ*tamB* and Δ*fadR* Δ*ydbH* strains had no growth
defects (**Fig 1D**). Although Δ*ydbH* in the Δ*yhdP* Δ*fadR* strain did not alter cold sensitivity
(**Fig. 1E**), Δ*tamB* in the Δ*yhdP* Δ*fadR* background completely suppressed the cold
sensitivity (**Fig. 1F**). Consistent with previous observations (30), the Δ*yhdP* Δ*fadR* Δ*tamB*
strain was mucoid due to Rcs (regulator of capsule synthesis) stress response activation

and colanic acid capsule production. Deleting *rcsB*, an Rcs response regulator, prevented
mucoidy but did not affect suppression (**Fig. 1F**). We observed only minimal growth
phenotypes in the Δ*yhdP* Δ*fadR* Δ*tamB* strain, similar to those of Δ*tamB* and less severe
than those of Δ*fadR* (**Fig. 1G**), making slow growth an unlikely suppression mechanism.
Together, these data demonstrate the Δ*yhdP* Δ*fadR* strain has impaired phospholipid
transport leading synthetic cold sensitivity, which can be relieved by deletion of Δ*tamB*,
shifting phospholipid transport to YdbH.

We next looked for envelope homeostasis alterations in the Δ*yhdP* Δ*fadR* strain. To
identify cell lysis and/or severe OM permeability defects at 30 °C, we assayed CPRG
(chlorophenol red-β-D-galactopyranoside) processing. CPRG must contact LacZ in the
cytoplasm or supernatant to release chlorophenol red (50). Compared to wild type or
single mutants, the Δ*yhdP* Δ*fadR* strain had increased CPRG processing (**Fig. S2A**),
demonstrating lysis or increased envelope permeability. When we assayed resistance to
molecules excluded by the OM (1, 42, 43, 51, 52) at 37 °C, a semi-permissive temperature,
the Δ*yhdP* Δ*fadR* strain showed sensitivity to vancomycin, a large scaffold antibiotic, and
SDS EDTA compared to single mutants (**Fig. S2B**). However, with bacitracin treatment or
EDTA treatment alone, the Δ*yhdP* Δ*fadR* strain phenocopied the Δ*fadR* single deletion
(**Fig. S2CD**). LPS levels were not altered in the Δ*yhdP*, Δ*fadR*, or the Δ*yhdP* Δ*fadR* strain
(**Fig. S2E**). OM asymmetry mutations (i.e., Δ*pldA*, Δ*mlaA*) (26-28, 53) did not have
suppressive or synthetic effects on cold sensitivity in the Δ*yhdP* Δ*fadR* strain (**Fig. S3AB**),
suggesting against a causal loss of OM asymmetry. Overall, the changes in OM
permeability did not seem sufficient to cause lethality at lower temperature, suggesting
the phospholipid transport disruption affects both IM and OM integrity.

Increased cardiolipin is necessary for cold sensitivity. We investigated phospholipid
composition of the wild-type, Δ*yhdP*, Δ*fadR*, and Δ*yhdP* Δ*fadR* strains grown at the
permissive temperature (42 °C) to log phase then downshifted to 30 °C. *E. coli*
phospholipid composition is generally 75% phosphatidylethanolamine (PE), 20%
phosphatidylglycerol (PG), and 5% cardiolipin (CL) with CL increasing in stationary phase
(14, 54). Of the three CL synthases, ClsA and ClsB synthesize CL from two PG molecules,
while ClsC synthesizes CL from one PG and one PE molecule, so CL and PG levels are
generally reciprocally regulated (55-58). For all strains, the levels of PE were similar (**Fig.
2A**, **Fig. S4A**). However, the Δ*yhdP* Δ*fadR* strain had increased CL and concomitant
decreased PG compared to the wild-type strain and compared to individual Δ*yhdP* and
Δ*fadR* mutants (**Fig. 2B**). PgsA synthesizes phosphatidylglycerol-phosphate and is the first
enzyme differentiating PG and PE biosynthesis. PgsA depletion would decrease both PG
and CL levels, reversing increased CL levels and exacerbating decreased PG levels in the
Δ*yhdP* Δ*fadR* strain. We tested the effect of PgsA depletion in a strain background (Δ*rcsF*
Δ*lpp*) where *pgsA* is non-essential (59-62). When *pgsA* expression was induced (arabinose
panel), the Δ*yhdP* Δ*fadR* PgsA-depletion strain had equal cold sensitivity to the Δ*yhdP*
Δ*fadR* strain (**Fig. 2C**). When *pgsA* expression was repressed (glucose panel), the Δ*yhdP*
Δ*fadR* cold sensitivity was partially suppressed, demonstrating that decreased PG does

not cause cold sensitivity and suggesting increased CL may be necessary for cold
sensitivity. Indeed, the cold sensitivity of the Δ*yhdP* Δ*fadR* strain was lost when CL was
fully depleted due to deletion of the genes for all cardiolipin syntheses (Δ*clsABC*) or of the
gene for the primary cardiolipin synthase (Δ*clsA*) responsible for CL synthesis in
exponential phase (**Fig. 2ADE**, **Fig. S4B**). Δ*clsB* caused a smaller decrease in CL levels, while
Δ*clsC* did not cause a decrease (**Fig. 2AB**, **Fig. 4B**). Surprisingly, however, Δ*clsC* completely
suppressed the cold sensitivity, while Δ*clsB* did not (**Fig. 2FG**). Thus, increased CL is
necessary for cold sensitivity but CL levels alone cannot explain the phenotype: another
factor, such as a specific molecular form (e.g., acyl chain length, saturation state,
symmetry of acyl chain arrangement) or localization (e.g., poles vs. midcell or inner vs.
outer leaflet) of CL, is required.

Increased *fabA* expression suppresses cold sensitivity. To identify other factors involved
in the cold sensitivity, we applied a forward genetic approach by isolating spontaneous
suppressor mutants capable of growing at 30 °C and identified a suppressor mutant
(Suppressor 1) that restored growth at 30 °C (**Fig. 3A**), decreased CPRG processing (**Fig.
S2A**), and had very similar CL to the Δ*yhdP* Δ*fadR* strain (**Fig. 3AB**, **Fig. S4A**). We identified
a point mutation in the *fabA* promoter region in Suppressor 1 (**Fig. 3B**) and confirmed this
mutation was sufficient for suppression (**Fig. 3C**). FabA is a dehydratase/isomerase that
introduces *cis* unsaturation into fatty acids (14, 63) and *fabA* is expressed from two
promoters controlled by FadR and FabR (64, 65). FadR activates *fabA* expression, while
FabR represses. Thus, *fadR* deletion would be expected to lower *fabA* expression. The
suppressor mutation was located in the *fabA* promoter FabR binding site (**Fig. 3B**) and we
hypothesized this mutation would increase *fabA* expression. *fabA* mRNA levels were more
than 4-fold lower in the Δ*yhdP* Δ*fadR* strain than the wild-type strain (**Fig. 3D**). While
lower than wild type, the suppressor mutant increased *fabA* mRNA 1.6-fold over the
Δ*yhdP* Δ*fadR* strain.

We constructed luciferase reporter plasmids with the wild-type *fabA* promoter region or
the *fabA* promoter region containing the suppressor mutation. The reporter with the
mutated promoter had higher activity in a wild-type, Δ*fadR,* and Δ*fabR* background (**Fig.
3E**). *fabA* transcription occurs from a promoter downstream of the FabR binding site (65-
67). However, when *fadR* is deleted, transcription shifts to the FabR-regulated promoter
within the FabR binding region (65-67). Our data indicate the Suppressor 1 mutation
regulates this promoter's constitutive activity rather than FabR binding. To confirm the
effect of changing the second promoter's activity, we tested the effect of Δ*fabR* on cold
sensitivity in the Δ*yhdP* Δ*fadR* strain and observed partial suppression (**Fig. S5**). The
smaller effect of Δ*fabR* compared to Suppressor 1 may result from other gene expression
changes or to the smaller relative effect of Δ*fabR* on *fabA* expression (**Fig. 3F**). Overall,
these data demonstrate increasing *fabA* expression, which is necessary for unsaturated

fatty acid biosynthesis, rescues cold sensitivity in the ΔyhdP ΔfadR strain without changing
CL levels.

Increasing unsaturated fatty acids relieves cold sensitivity. Given the effect of *fabA*
expression on the ΔyhdP ΔfadR cold sensitivity, we used liquid chromatography-
electrospray ionization mass spectrometry (LC/MS) to characterize the strains'
phospholipid saturation state (**Dataset S1**, **Fig. 4A-D**). Representitive spectra
demonstrating relative phospholipid composition are shown in **Fig. S6**. As expected,
Δ*fadR* caused increased fully saturated and monounsaturated PE and PG with a
concomitant decrease of diunsaturated PE (**Fig. 4AB**). Similarly, the Δ*fadR* strain
demonstrated increased monounsaturated CL with a concomitant decrease in
triunsaturated CL (**Fig. 4C**). Compared to the Δ*fadR* strain, the ΔyhdP Δ*fadR* strain had
slightly decreased saturation for all three phospholipids, while still displaying more
phospholipid saturation than wild type or ΔyhdP strains (**Fig. 4A-C**). Suppressor 1 trended
towards decreased saturation of all three phospholipids compared to the ΔyhdP Δ*fadR*
strain (**Fig. 4A-C**). We wondered whether the suppression difference between Δ*clsC* and
Δ*clsB* resulted from the CL saturation state. However, the ΔyhdP Δ*fadR* Δ*clsB* and ΔyhdP
Δ*fadR* Δ*clsC* strains had very similar CL profiles (**Fig. 4C**), suggesting another qualitative
difference affecting suppression, perhaps CL lateral or inter-leaflet distribution in the cell.
These strains also showed similar PE and PG saturation profiles to their ΔyhdP Δ*fadR*
parent (**Fig. 4AB**).

We hypothesized increasing unsaturated fatty acids suppresses ΔyhdP Δ*fadR* cold
sensitivity and performed EOPs for cold sensitivity on media containing an unsaturated
fatty acid (oleic acid, C18:1 *cis*-9). Exogenous phospholipids can be taken up by *E. coli*,
attached to acyl-CoA, and incorporated into phospholipids (68). Oleic acid addition
suppressed the ΔyhdP Δ*fadR* strain's cold sensitivity (**Fig. 4D**). Supplementing oleic acid
increases unsaturated phospholipids and provides an exogenous fatty acid source. To
differentiate between saturation state and fatty acid availability, we compared treatment
with oleic acid and saturated stearic acid (C18:0). Only oleic acid, and not stearic acid,
suppressed the ΔyhdP Δ*fadR* strain's cold sensitivity (**Fig. 4E**), confirming increasing
unsaturated fatty acids, not fatty acid availability suppresses.

Suppression causes fatty acid metabolism and stress response alterations. In wild-type
cells, *fabA* overexpression does not change phospholipid saturation levels, demonstrating
the FabB enzyme is rate limiting for unsaturated fatty acid biosynthesis (69). We were
intrigued that Suppressor 1 altered saturation without directly effecting *fabB* expression.
We also wondered whether CL levels in the ΔyhdP Δ*fadR* strain are transcriptionally
regulated. Thus, we investigated the transcriptional landscape of the wild type, the ΔyhdP
Δ*fadR*, and Suppressor 1 strains after a 30 minute downshift of growth to 30 °C to
determine: (i) whether the ΔyhdP Δ*fadR* strain had altered *cls* gene expression; (ii)
whether Suppressor 1 had other transcriptional changes; and (iii) why increased *fabA*
expression decreased saturation. All differentially expressed genes (>2-fold change,

p<0.05) are listed in **Dataset S2**. Principle component analysis of these genes showed the
closest relation between Δ*yhdP* Δ*fadR* strain and Suppressor 1, while the wild-type strain
was more distantly related (**Fig. 5A**, **S7A**). Nevertheless, Suppressor 1 was more closely
related to wild type than was the Δ*yhdP* Δ*fadR* strain. Most differentially regulated genes
between Δ*yhdP* Δ*fadR* and wild type and Suppressor 1 and wild type were upregulated
(**Fig. 5BC**, **S7BC**). Many of the most highly regulated genes are FadR-regulon members.
The majority of differentially regulated genes in Suppressor 1 compared to Δ*yhdP* Δ*fadR*
are downregulated (**Fig. 5D, S7D**). These genes are involved in many cellular pathways
and likely reflect the decreased cellular stress (**Table S2**). Indeed, many external stress
response genes are more enriched in Δ*yhdP* Δ*fadR* than in Suppressor 1 (**Fig. S8A**).

No significant expression changes occurred in the CL biosynthesis pathway (**Fig. 5E**) or
other phospholipid biosynthesis genes (**Fig. S8B**). The increased cardiolipin in Δ*yhdP*
Δ*fadR* and Suppressor 1 may be post-transcriptional or due to a change in a regulatory
pathway not yet altered after 30 minutes at 30 °C (**Fig. S8C**). Levels of mitochondrial CL
can be altered due to differences in substrate binding affinity of the CL synthase based on
saturation, with increased saturation decreasing CL synthesis (70), and it may be that a
similar pathway operates here, albeit in a different direction. We examined the
expression of genes in fatty acid synthesis and degradation, many of which are members
of the FadR regulon. In Δ*yhdP* Δ*fadR* compared to wild type, many fatty acid biosynthesis
genes significantly decreased while many fatty acid degradation genes significantly
increased (**Fig. 5F**), consistent with Δ*fadR* effects. While *fabA* expression is decreased,
fabB expression is not changed, explaining why increased *fabA* expression causes
decreased saturation in Suppressor 1. Overall, our data demonstrate the Δ*yhdP* Δ*fadR*
strain's cold sensitivity is due to impaired phospholipid trafficking, and increased
phospholipid saturation and cardiolipin are necessary for this functional impairment.

336
337

Discussion
The identification of YhdP, TamB, and YdbH as putative intermembrane phospholipid
transporters (30, 31) posed the tantalizing question: why is having three transporters
advantageous? Differential phenotypes between the transporters suggest that they
possess specialized functions (25, 30, 31, 34, 43-46). Here, we demonstrate each protein
plays a distinguishable role in phospholipid transport that can be differentiated based on
genetic interactions with lipid metabolism. Disruption of FadR and YhdP function causes
synthetic cold sensitivity not shared by TamB or YdbH, which can be suppressed by
removing TamB and switching transport to YdbH. This phenotype requires both increased
levels of CL and saturated fatty acids, and decreasing amounts of either suppresses. In
addition, our data demonstrate that CL synthesized by ClsB and ClsC is qualitatively
different—resulting in differential suppression phenotypes—likely due to differences in
CL localization, CL-mediated, leaflet-specific changes in lipid packing order (9, 24),
localized membrane fluidity (71), or TamB interaction.

Based on these data, we now suggest a model for the differentiation of function between
YhdP and TamB that leads to their independent phenotypes (**Fig. 6A-C**). In this model, the
functions of YhdP and TamB are diversified based on the acyl carbon saturation level
preference of the phospholipids they transport: YhdP transports mainly phospholipids
with more saturated fatty acids, while TamB transports phospholipids with more
unsaturated fatty acids (**Fig. 6A**). Our data and previous data (30, 31) agree YdbH plays a
relatively minor role. In the Δ*yhdP* Δ*fadR* strain, YhdP's absence forces more saturated
phospholipids to be transported by TamB. At 30 °C, TamB becomes "clogged" by
phospholipids having more saturated fatty acyl side chains, impeding transport (**Fig. 6B**).
This cold-specific effect may occur due to (i) temperature-dependent decreased
membrane fluidity and higher lipid packing order involving lower expression of FabF (**Fig.**
5F, **Dataset 3**), which is responsible for increased diunsaturated phospholipids at lower
temperature (14); (ii) higher relative levels of CL (72); (iii) higher amounts of per cell
phospholipids at lower temperatures (72) contributing to increased diffusive flow rate;
or, (iv) to thermodynamic properties altering TamB transport specificity and/or substrate
behavior contributing to efficiency of substrate loading or transport. Bulky and
intrinsically disordered CL molecules could contribute to the clogging sterically by
obstructing the lipid channel and precluding it from translocating other lipid substrates
(i.e., PE). YdbH's ability to successfully transport phospholipids in the Δ*yhdP* Δ*tamB* strain
only when TamB is absent may be because YhdP and TamB are primary transporters (e.g.,
preferred interaction partners for other pathway members), due to direct toxicity caused
by TamB clogging, or because *ydbH* is upregulated in the absence of the other
phospholipid transporters.

Several pieces of evidence demonstrate dysregulation of phospholipid transport not just
changes to OM phospholipid composition, and so permeability or physical strength, cause
the cold sensitivity phenotype of the Δ*yhdP* Δ*fadR* strain. First, the cold sensitivity is
unaffected by removing enterobacterial common antigen, which suppresses the
membrane permeability of a Δ*yhdP* strain (**Fig. 1C**, (43)), and the OM permeability
phenotypes of the Δ*yhdP* Δ*fadR* are not severe (**Fig. S2**). Second, the cold sensitivity is
suppressed by deletion of Δ*tamB* (**Fig. 1FG**). Δ*yhdP* Δ*tamB* strains have much greater OM
permeability (30) than the Δ*yhdP* or Δ*yhdP* Δ*fadR* strains. Third, CL has been shown to
have a normalizing, bidirectional, and leaflet-specific effect on lipid packing order (9, 73-
76) and membrane fluidity (77-80) opposite of or similar to that of cholesterol,
respectively, and so is unlikely to contribute to a phenotype dependent solely on OM
membrane fluidity. Thus, the cold sensitivity phenotype correlates with the impairment
of phospholipid trafficking and the likely imbalance of phospholipids between the IM and
OM and not with the permeability phenotypes of the OM.

Although both fatty acid saturation and high CL levels are required for the Δ*yhdP* Δ*fadR*
strain's cold sensitivity, two lines of evidence point to saturation rather than headgroup
as the main factor separating YhdP and TamB function. (i) YhdP loss slows phospholipid
transport from the IM to the OM in a *mlaA** mutant, which loses OM material due to

vesiculation and lysis when phospholipid biosynthesis is insufficient to replace lost
phospholipids (25, 27). CL only accounts for 5% of phospholipids in exponentially growing
cells making it unlikely that the loss of CL transport could slow net phospholipid transport
to the extent required to slow lysis. (ii) Δ*tamB* has a small growth defect at 30 °C (**Fig. 1G**)
not observed at higher temperature (30, 31), perhaps due to increased unsaturated
phospholipids (81-83) resulting in decreased net OM transport by YhdP or to lower OM
fluidity. Together, these observations provide strong evidence YhdP has a substrate
preference for phospholipids with more saturated fatty acids, while TamB prefers
phospholipids with more unsaturated fatty acids.

Our model suggests the IM and OM saturation state could be independently regulated
based on YhdP and TamB activity or levels. IM saturation is regulated by altering the
composition of newly synthesized fatty acids, while OM saturation could be separately
regulated by allowing increased transport of more saturated or unsaturated
phospholipids as necessary. IM and OM acyl saturation has been studied in several strains
of *E. coli* and in *Yersinia pseudotuberculosis* (7, 19-21) and higher saturated fatty acid
levels were observed in the OM than in the IM, supporting discrimination of phospholipid
transport based on saturation. The IM needs to maintain an appropriate level of fluidity
to allow IM proteins to fold and diffuse, to allow diffusion of hydrophobic molecules, to
regulate cellular respiration, and to maintain membrane integrity (84, 85). However, the
OM is a permeability barrier, a load-bearing element that resists turgor pressure (1-6),
and has an outer leaflet consisting of LPS (2, 8). LPS has saturated acyl chains and is largely
immobile in *E. coli* (86) and a specific phospholipid saturation level may be necessary to
match or counterbalance LPS rigidity. OM phospholipids may also play a role in the
physical strength of the membrane. Interestingly, previous work has demonstrated that
the tensile strength of the OM is lessened in a *yhdP* mutant (25).

Possessing two different phospholipid transporters with different saturation state
preferences would allow the cell to adapt OM saturation to changing conditions including
temperature shifts, chemical and physical insults, or changes in synthesis of other OM
components (e.g., LPS or OMPs). A recent study demonstrated in *Y. pseudotuberculosis*
the difference in saturation levels between the membranes is exaggerated at 37 °C
compared to 8 °C (21). Thus, conditions where IM and OM saturation are independently
regulated exist and this regulation is likely important for optimum fitness of the cells.
Transcriptional regulation of *yhdP* and *tamB* has not been thoroughly explored. It seems
likely *yhdP* transcription is quite complex with several promoters and transcriptional
termination sites (87-89). *tamB* is predicted to be in an operon with *tamA* (which codes
for an OMP necessary for TamB function) and *ytfP* (87, 90) and to have a σ^E-dependent
promoter (91). The σ^E stress response is activated by unfolded OMP accumulation, and
downregulates the production of OMPs and upregulates chaperones and folding
machinery (74). It is intriguing to speculate a change in OM phospholipid composition
might aid in OMP folding due to altered biophysical properties and/or lipoprotein
diffusion. In fact, this could explain the observations that some complicated OMPs fold

more slowly in the absence of TamB (34, 44-46). Proper phospholipid composition may
be necessary for efficient folding of these OMPs, as it is necessary for the correct folding
and topology of some IM proteins (73, 92-95). Recently, it has been demonstrated that
the *tamAB* operon is regulated by PhoPQ in *Salmonella enterica* Serovar Typhimurium
and that this regulation helps maintain OM integrity in the acidic phagosome
environment, perhaps due to indirect effects on OMP folding (96). Thus, changing
environments may necessitate changes in phospholipid composition that can be
accomplished through differential TamB and, perhaps YhdP, regulation.

The mechanism of phospholipid transfer between the IM and OM has been one of the
largest remaining questions in gram-negative envelope biogenesis and has remained an
area of intense investigation for decades. Furthermore, given the lack of knowledge of
this pathway, the role of phospholipid composition in the permeability barrier of the OM,
and so the intrinsic antibiotic resistance of gram-negative bacteria, remains unknown.
Recent work has begun to elucidate the intermembrane phospholipid transport pathway
by identifying YhdP, TamB, and YdbH as proteins capable of transporting phospholipids
between the membranes (25, 30, 31). Our work here suggests that YhdP and TamB may
have separate roles in phospholipid transport that are differentiated by their preference
for phospholipid saturation states and possibly lipid headgroup. Homologs of YhdP and
TamB are found throughout *Enterobacterales* and more distantly related members of the
AsmA family are widespread in gram-negative bacteria (97)*. In fact, TamB has recently
been shown to play a role in anterograde intermembrane phospholipid transport in the
diderm Firmicute *Veillonella parvula* (40, 98) and four AsmA members play a role in
phospholipid transport in *Pseudomonas aeruginosa* (41). Structural predictions of YhdP
and TamB are strikingly similar between species (32, 33). Given this, it is quite likely similar
mechanisms of discrimination in intermembrane phospholipid transfer exist in other
species. The data we present here provide a framework for investigation of phospholipid
transport and mechanisms differentiating phospholipid transporter function in these
species.

462
463
Materials and Methods

Bacterial strains and Growth Conditions. Table S3 lists the strains used in this study. Most
knockout strains were constructed with Keio collection alleles using P1vir transduction
(99, 100). New alleles were constructed through λ-red recombineering (101), using the
primers indicated in **Table S4**. The FLP recombinase-FRT system was used to remove
antibiotic resistance cassettes as has been described (101). *E. coli* cultures were grown in
LB Lennox media at 42°C, unless otherwise mentioned. Where noted, LB was
supplemented with vancomycin (Gold Biotechnology), bacitracin (Gold Biotechnology),
SDS (Thermo Fisher Scientific), EDTA (Thermo Fisher Scientific), oleic acid (Sigma Aldrich),
stearic acid (Sigma Aldrich), arabinose (Gold Biotechnology), or glucose (Thermo

Scientific). For plasmid maintenance, media was supplemented with 25 µg/mL kanamycin
(Gold Biotechnology), 10 or 20 µg/mL chloramphenicol (Gold Biotechnology), or 10 µg/mL
tetracycline (Gold Biotechnology) as appropriate.

Plasmid construction. The pJW15-P*fabA*, and pJW15-P*fabA*(-19G>A) reporter plasmids were
constructed by amplifying the region upstream of *fabA* from -432 bp to +27 bp relative to
the annotated transcriptional start site from wild-type MG1655 or Suppressor 1,
respectively, using the Overlap-pJW15-*fabA* FP and RP primers (**Table S4**). pJW15 was
amplified using primers Overlap-pJW15-*fabA* FP1 and RP1. The pJW15 plasmid was a kind
of gift from Dr. Tracy Raivio (Univeristy of Alberta) (102, 103). The resulting fragments
were assembled using HiFi Assembly master mix (New England Biolabs) as per the
manufacturer's instructions.

Growth and Sensitivity Assays. For EOPs assays, cultures were grown overnight, serially
diluted, and plated on the indicated media. For *pgsA* depletion, overnight cultures grown
in arabinose were washed and diluted 1:100 into fresh LB with 0.2% glucose or arabinose
as indicated and grown to OD_{600}=1 before diluting and plating on media with glucose or
arabinose respective, as above. EOPs were incubated at 30 °C unless otherwise indicated.
Growth curves were performed at the indicated temperature as has been described (43).
For CPRG assays (50), strains were streaked on LB plates supplemented with CPRG
(40µg/ml) and IPTG (100 µM) incubated overnight at 30°C.

Suppressor isolation and sequencing. Serially diluted cultures were spread on LB plates
and incubated overnight at a 30°C. Colonies that grew at 30°C were isolated and
rechecked for cold-resistance through EOP and growth curves. The suppressors were
subjected to whole-genome sequencing to identify potential suppressor mutations.
Genomic DNA was isolated from the parent and suppressors using a DNeasy Blood and
Tissue Kit (Qiagen) as per the manufacturer's instructions. Library preparation and
Illumina sequencing were performed by the Microbial Genome Sequencing Center (MiGS,
Pittsburgh, PA, USA). Briefly, Illumina libraries were prepared and the libraries were
sequenced with paired-end 151 bp reads on a NextSeq 2000 platform to a read depth of
at least 200 Mbp. To identify variants from the reference genome (GenBank: U00096.3),
reads were trimmed to remove adaptors and the *breseq* software (version 0.35.4) was
used to call variants (104). Identified mutations were confirmed with Sanger sequencing.
The consensus motif of the FabR binding site of *fabA* was determined using Multiple Em
for Motif Elicitation (MEME) tool in the MEME Suite based on the sequence of putative
FabR binding sites reported by Feng and Cronan (2011) (64, 105, 106). The ability of the
identified P*fabA* mutation to suppress was confirmed by linking the mutant to *zcb*-
3059::Tn*10* (107) and transducing to a clean background.

Lux reporter assays. Overnight cultures of strains harboring the pJW15-P*fabA* and pJW15-
P*fabA*(-19G>A) plasmids were sub-cultured at (1:100) into 200 µl of fresh LB broth in a black
96-well plate. The plate was incubated in a BioTek Synergy H1 plate reader and the OD_{600}

and luminescence intensity were recorded every 2.5 min for 6 hours as described in
previously (103, 108). Each biological replicate was performed in technical triplicate.

RNA-seq. Overnight cultures grown at 42°C were subcultured 1:100 into fresh media and
incubated at 42 °C until an OD_{600} of 0.4. Then, the cultures were shifted to a 30 °C water
bath for 30 minutes. 500 µL of each culture was immediately fixed in the RNAprotect
Bacteria Reagent (Qiagen) as per the manufacturer's instructions. RNA was purified using
the RNeasy Kit (Qiagen) with on-column DNase digestion following the manufacturer's
protocol for gram-negative bacteria. Library preparation, Illumina sequencing, and
analysis of differentially expressed genes were performed by MiGS. Briefly, samples were
treated with RNase free DNase (Invitrogen) and library preparation was performed with
the Stranded Total RNA Prep Ligation with Ribo-Zero Plus Kit (Illumina). Libraries were
sequenced with 2x50 bp reads on a NextSeq2000 (Illumina). Demultiplexing, quality
control, and adapter trimming carried out using bcl-convert (Illumina, v3.9.3) and
bcl2fastq (Illumina). Read mapping and read quantification were performed with HISAT2
(109) and the featureCounts tool in Subreader (110), respectively. Descriptive statistics of
the RNA-seq read data can be found in **Table S1**. The raw RNA-seq data is available in the
Sequence Read Archive database (SRA) (https://www.ncbi.nlm.nih.gov/sra, BioProject ID
PRJNA965821; Reviewer Access).

Raw read counts were normalized using the Trimmed Mean of M values algorithm in
edgeR (111) and converted to counts per million. The Quasi-Linear F-Test function of
edgeR was used for differential expression analysis. Normalized read values and for all
genes as well as differential expression analysis statistics can be found in **Dataset S3**.
KEGG pathway analysis was conducted using the *"kegga"* function of limma (112).
Expression of genes differentially expressed (greater than 2-fold change and p<0.05) was
subjected to principle component analysis, and pathway and enrichment analysis using
the EcoCyc Omics Dashboard (112).

Lipid extraction and TLC. For thin layer chromatography (TLC) experiments, cultures were
grown in LB with 1 mCi/mL ^{32}P until OD_{600} of 0.6-0.8 at the indicated temperature. For
temperature downshift experiments, cells were grown at 42 °C until OD_{600} of 0.2 and then
transferred to 30 °C for 2 hours. For lipid extraction, cells were pelleted and resuspended
in 0.2 mL of 0.5 M NaCl in 0.1 N HCl. Lipids were extracted by first adding 0.6 mL of
chloroform:methanol (1:2) to create a single-phase solution. After vigorous vortexing for
30 minutes, 0.2 mL of 0.5 M NaCl in 0.1 N HCl was added to convert the single-phase
solution to a two-phase solution. After centrifugation at 20 000 x g for 5 minutes at room
temperature, the lower phase was recovered and used for TLC. Approximately 2,000 cpm
of phospholipid extract was subjected to TLC analysis using a HPTLC 60 plate (EMD,
Gibbstown, NJ)) developed with either solvent 1: chloroform/methanol/acetic acid
[60/25/10 (vol/vol/vol) in **Fig. 2A** or by solvent 2: chloroform/methanol/ammonia/water
[65/37.5/3/1] (vol/vol/vol/vol) in **Fig. S4**. Radiolabeled lipids were visualized and
quantified using a Phosphoimager Typhoon FLA 9500 (GE). Images were processed and

quantified using ImageQuant^{TM} software for scanning and analysis. The phospholipid
content is expressed as molecular percentage of the total phospholipids (correcting for
the two phosphates per molecule of CL).

Phospholipid composition analysis. Whole cell lipid extracts were isolated as for TLC
temperature downshift experiments without radiolabelling. Absolute amounts of CL, PG,
and PE were determined in these lipid extracts using liquid chromatography coupled to
electrospray ionization mass spectrometry (LC/MS) in an API 4000 mass spectrometer
(Sciex, Framingham, MA, USA) using normal phase solvents as previously published (113).
The cardiolipin internal standard (14:0)$_4$CL (Avanti Polar Lipids, Alabaster, AL USA) was
added to all samples to quantify CL, PG, and PE with in a single run. Standard curves
containing reference standards of all three phospholipids, (18:1)$_4$CL (Avanti Polar Lipids),
16:0$_1$18:1$_1$PG and 16:0$_1$18:1$_1$PE (Cayman Chemicals, Ann Arbor, MI, USA) were used to
quantify the nmol/mg protein for each species (**Dataset 1**) as previously described (114).
Percentage values were calculated by dividing the absolute value of that species by the
sum of individual molecular species of CL, PG, or PE. The identities of CL species were
confirmed using tandem mass spectrometry.
569
570

571 **Acknowledgments**
572
We thank members of the Mitchell Lab for productive discussions. We would like to thank
Professor Tracy Raivio (University of Alberta) for the kind gift of pJW15 plasmid. This work
was funded by the National Institute of Allergy and Infectious Disease under Award
Number R01-AI155915 and R01-AI155915-S1 (to A.M.M.), the National Institute of
General Medical Sciences under Award Number R01-GM121493 (to M.B.), and Texas
A&M University start-up funds (to A.M.M.).

579
580

References

1.   Nikaido H. 2003. Molecular basis of bacterial outer membrane permeability
revisited. Microbiology and Molecular Biology Reviews 67:593-656.
2.   Silhavy TJ, Kahne D, Walker S. 2010. The bacterial cell envelope. Cold Spring
Harbor Perspectives in Biology 2.
3.   Henderson JC, Zimmerman SM, Crofts AA, Boll JM, Kuhns LG, Herrera CM, Trent
MS. 2016. The power of asymmetry: Architecture and assembly of the Gram-
negative outer membrane lipid bilayer. Annual Review of Microbiology 70:255-
278.
4.   Berry J, Rajaure M, Pang T, Young R. 2012. The spanin complex is essential for
lambda lysis. J Bacteriol 194:5667-74.

5.   Rojas ER, Billings G, Odermatt PD, Auer GK, Zhu L, Miguel A, Chang F, Weibel DB,
Theriot JA, Huang KC. 2018. The outer membrane is an essential load-bearing
element in Gram-negative bacteria. Nature 559:617-621.
6.   Navarro Llorens JM, Tormo A, Martinez-Garcia E. 2010. Stationary phase in gram-
negative bacteria. FEMS Microbiol Rev 34:476-95.
7.   Lugtenberg EJJ, Peters R. 1976. Distribution of lipids in cytoplasmic and outer
membranes of *Escherichia coli* K12. Biochimica et Biophysica Acta (BBA) - Lipids
and Lipid Metabolism 441:38-47.
8.   Kamio Y, Nikaido H. 1976. Outer membrane of *Salmonella typhimurium*:
accessibility of phospholipid head groups to phospholipase c and cyanogen
bromide activated dextran in the external medium. Biochemistry 15:2561-70.
9.   Bogdanov M, Pyrshev K, Yesylevskyy S, Ryabichko S, Boiko V, Ivanchenko P,
Kiyamova R, Guan Z, Ramseyer C, Dowhan W. 2020. Phospholipid distribution in
the cytoplasmic membrane of Gram-negative bacteria is highly asymmetric,
dynamic, and cell shape-dependent. Science Advances 6:eaaz6333.
10.   Benn G, Mikheyeva IV, Inns PG, Forster JC, Ojkic N, Bortolini C, Ryadnov MG,
Kleanthous C, Silhavy TJ, Hoogenboom BW. 2021. Phase separation in the outer
membrane of *Escherichia coli*. Proceedings of the National Academy of Sciences
118:e2112237118.
11.   Konovalova A, Silhavy TJ. 2015. Outer membrane lipoprotein biogenesis: Lol is
not the end. Philosophical Transactions of the Royal Society B: Biological
Sciences 370:20150030.
12.   Konovalova A, Kahne DE, Silhavy TJ. 2017. Outer Membrane Biogenesis. Annu
Rev Microbiol 71:539-556.
13.   Lundstedt E, Kahne D, Ruiz N. 2021. Assembly and maintenance of lipids at the
bacterial outer membrane. Chemical Reviews 121:5098-5123.
14.   John E. Cronan J, Rock CO. 2008. Biosynthesis of membrane lipids. EcoSal Plus 3.
15.   Jones NC, Osborn MJ. 1977. Translocation of phospholipids between the outer
and inner membranes of *Salmonella typhimurium*. Journal of Biological
Chemistry 252:7405-7412.
16.   Donohue-Rolfe AM, Schaechter M. 1980. Translocation of phospholipids from
the inner to the outer membrane of *Escherichia coli*. Proceedings of the National
Academy of Sciences 77:1867-1871.
17.   Raetz CR, Kantor GD, Nishijima M, Newman KF. 1979. Cardiolipin accumulation in
the inner and outer membranes of *Escherichia coli* mutants defective in
phosphatidylserine synthetase. J Bacteriol 139:544-51.
18.   Langley KE, Hawrot E, Kennedy EP. 1982. Membrane assembly: movement of
phosphatidylserine between the cytoplasmic and outer membranes of
Escherichia coli. J Bacteriol 152:1033-41.
19.   White DA, Lennarz WJ, Schnaitman CA. 1972. Distribution of lipids in the wall
and cytoplasmic membrane subfractions of the cell envelope of *Escherichia coli*.
Journal of Bacteriology 109:686-690.

20.   Ishinaga M, Kanamoto R, Kito M. 1979. Distribution of phospholipid molecular
635   species in outer and cytoplasmic membranes of *Escherichia coli*. The Journal of
636   Biochemistry 86:161-165.
637   21.   Davydova L, Bakholdina S, Barkina M, Velansky P, Bogdanov M, Sanina N. 2016.
638   Effects of elevated growth temperature and heat shock on the lipid composition
639   of the inner and outer membranes of *Yersinia pseudotuberculosis*. Biochimie
640   123:103-109.
641   22.   Osborn MJ, Gander JE, Parisi E, Carson J. 1972. Mechanism of Assembly of the
642   Outer Membrane of *Salmonella* typhimurium : Isolation and characterization of
643   cytoplasmic and outer membrane. Journal of Biological Chemistry 247:3962-
644   3972.
645   23.   Raetz CR, Newman KF. 1979. Diglyceride kinase mutants of *Escherichia coli*: inner
646   membrane association of 1,2-diglyceride and its relation to synthesis of
647   membrane-derived oligosaccharides. J Bacteriol 137:860-8.
648   24.   Bogdanov M. 2023. Renovating a double fence with or without notifying the next
649   door and across the street neighbors: why the biogenic cytoplasmic membrane
650   of Gram-negative bacteria display asymmetry? . Emerging Topics in Life Sciences
651   7:137-150.
652   25.   Grimm J, Shi H, Wang W, Mitchell AM, Wingreen NS, Huang KC, Silhavy TJ. 2020.
653   The inner membrane protein YhdP modulates the rate of anterograde
654   phospholipid flow in *Escherichia coli*. Proc Natl Acad Sci U S A 117:26907-26914.
655   26.   Malinverni JC, Silhavy TJ. 2009. An ABC transport system that maintains lipid
656   asymmetry in the Gram-negative outer membrane. Proceedings of the National
657   Academy of Sciences 106:8009-8014.
658   27.   Sutterlin HA, Shi H, May KL, Miguel A, Khare S, Huang KC, Silhavy TJ. 2016.
659   Disruption of lipid homeostasis in the Gram-negative cell envelope activates a
660   novel cell death pathway. Proceedings of the National Academy of Sciences
661   113:E1565-E1574.
662   28.   May KL, Silhavy TJ. 2018. The *Escherichia coli* phospholipase PldA regulates outer
663   membrane homeostasis via lipid signaling. mBio 9:e00379-18.
664   29.   Tang X, Chang S, Qiao W, Luo Q, Chen Y, Jia Z, Coleman J, Zhang K, Wang T,
665   Zhang Z, Zhang C, Zhu X, Wei X, Dong C, Zhang X, Dong H. 2021. Structural
666   insights into outer membrane asymmetry maintenance in Gram-negative
667   bacteria by MlaFEDB. Nat Struct Mol Biol 28:81-91.
668   30.   Ruiz N, Davis RM, Kumar S. 2021. YhdP, TamB, and YdbH are redundant but
669   essential for growth and lipid homeostasis of the gram-negative outer
670   membrane. mBio 12:e02714-21.
671   31.   Douglass MV, McLean AB, Trent MS. 2022. Absence of YhdP, TamB, and YdbH
672   leads to defects in glycerophospholipid transport and cell morphology in Gram-
673   negative bacteria. PLOS Genetics 18:e1010096.
674   32.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O,
675   Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl

SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T,
Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M,
Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K,
Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with
AlphaFold. Nature 596:583-589.

33.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D,
Stroe O, Wood G, Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S,
Jumper J, Clancy E, Green R, Vora A, Lutfi M, Figurnov M, Cowie A, Hobbs N,
Kohli P, Kleywegt G, Birney E, Hassabis D, Velankar S. 2021. AlphaFold Protein
Structure Database: massively expanding the structural coverage of protein-
sequence space with high-accuracy models. Nucleic Acids Research 50:D439-
D444.

34.   Josts I, Stubenrauch CJ, Vadlamani G, Mosbahi K, Walker D, Lithgow T, Grinter R.
2017. The structure of a conserved domain of TamB reveals a hydrophobic β taco
fold. Structure 25:1898-1906.e5.

35.   Kumar N, Leonzino M, Hancock-Cerutti W, Horenkamp FA, Li P, Lees JA, Wheeler
H, Reinisch KM, De Camilli P. 2018. VPS13A and VPS13C are lipid transport
proteins differentially localized at ER contact sites. Journal of Cell Biology
217:3625-3639.

36.   Valverde DP, Yu S, Boggavarapu V, Kumar N, Lees JA, Walz T, Reinisch KM, Melia
TJ. 2019. ATG2 transports lipids to promote autophagosome biogenesis. J Cell
Biol 218:1787-1798.

37.   Otomo T, Maeda S. 2019. ATG2A transfers lipids between membranes *in vitro*.
Autophagy 15:2031-2032.

38.   Osawa T, Kotani T, Kawaoka T, Hirata E, Suzuki K, Nakatogawa H, Ohsumi Y, Noda
NN. 2019. Atg2 mediates direct lipid transfer between membranes for
autophagosome formation. Nat Struct Mol Biol 26:281-288.

39.   Cooper BF, Clark R, Kudhail A, Bhabha G, Ekiert DC, Khalid S, Isom GL. 2023.
Phospholipid transport to the bacterial outer membrane through an envelope-
spanning bridge. bioRxiv doi:10.1101/2023.10.05.561070:2023.10.05.561070.

40.   Grasekamp KP, Beaud Benyahia B, Taib N, Audrain B, Bardiaux B, Rossez Y, Izadi-
Pruneyre N, Lejeune M, Trivelli X, Chouit Z, Guerardel Y, Ghigo J-M, Gribaldo S,
Beloin C. 2023. The Mla system of diderm Firmicute *Veillonella parvula* reveals
an ancestral transenvelope bridge for phospholipid trafficking. Nature
Communications 14:7642.

41.   Sposato D, Mercolino J, Torrini L, Sperandeo P, Lucidi M, Alegiani R, Varone I,
Molesini G, Leoni L, Rampioni G, Visca P, Imperi F. 2024. Redundant essentiality
of AsmA-like proteins in *Pseudomonas aeruginosa*. mSphere 9:e00677-23.

42.   Mitchell AM, Wang W, Silhavy TJ. 2017. Novel Rpos-dependent mechanisms
strengthen the envelope permeability barrier during stationary phase. J Bacteriol
199:e00708-16.

43.   Mitchell AM, Srikumar T, Silhavy TJ. 2018. Cyclic enterobacterial common
718   antigen maintains the outer membrane permeability barrier of *Escherichia coli* in
719   a manner controlled by Yhdp. mBio 9:e01321-18.
720   44.   Selkrig J, Mosbahi K, Webb CT, Belousoff MJ, Perry AJ, Wells TJ, Morris F, Leyton
721   DL, Totsika M, Phan M-D, Celik N, Kelly M, Oates C, Hartland EL, Robins-Browne
722   RM, Ramarathinam SH, Purcell AW, Schembri MA, Strugnell RA, Henderson IR,
723   Walker D, Lithgow T. 2012. Discovery of an archetypal protein transport system
724   in bacterial outer membranes. Nature Structural & Molecular Biology 19:506-
725   510.
726   45.   Stubenrauch CJ, Dougan G, Lithgow T, Heinz E. 2017. Constraints on lateral gene
727   transfer in promoting fimbrial usher protein diversity and function. Open Biology
728   7:170144.
729   46.   Stubenrauch C, Belousoff MJ, Hay ID, Shen H-H, Lillington J, Tuck KL, Peters KM,
730   Phan M-D, Lo AW, Schembri MA, Strugnell RA, Waksman G, Lithgow T. 2016.
731   Effective assembly of fimbriae in *Escherichia coli* depends on the translocation
732   assembly module nanomachine. Nature Microbiology 1:16064.
733   47.   Cronan JE. 2021. The *Escherichia coli* FadR transcription factor: Too much of a
734   good thing? Molecular Microbiology 115:1080-1085.
735   48.   Nunn WD, Giffin K, Clark D, Cronan JE. 1983. Role for *fadR* in unsaturated fatty
736   acid biosynthesis in *Escherichia coli*. Journal of Bacteriology 154:554-560.
737   49.   Rai AK, Mitchell AM. 2020. Enterobacterial common antigen: Synthesis and
738   function of an enigmatic molecule. mBio 11:e01914-20.
739   50.   Paradis-Bleau C, Kritikos G, Orlova K, Typas A, Bernhardt TG. 2014. A genome-
740   wide screen for bacterial envelope biogenesis mutants identifies a novel factor
741   involved in cell wall precursor metabolism. PLoS Genet 10:e1004056.
742   51.   Ruiz N, Falcone B, Kahne D, Silhavy TJ. 2005. Chemical conditionality: a genetic
743   strategy to probe organelle assembly. Cell 121:307-17.
744   52.   Nikaido H, Vaara M. 1985. Molecular basis of bacterial outer membrane
745   permeability. Microbiol Rev 49:1-32.
746   53.   Bishop RE. 2008. Structural biology of membrane-intrinsic β-barrel enzymes:
747   Sentinels of the bacterial outer membrane. Biochimica et Biophysica Acta (BBA) -
748   Biomembranes 1778:1881-1896.
749   54.   Rowlett VW, Mallampalli V, Karlstaedt A, Dowhan W, Taegtmeyer H, Margolin W,
750   Vitrac H. 2017. Impact of membrane phospholipid slterations in *Escherichia coli*
751   on cellular function and bacterial stress adaptation. J Bacteriol 199.
752   55.   Tan BK, Bogdanov M, Zhao J, Dowhan W, Raetz CRH, Guan Z. 2012. Discovery of
753   a cardiolipin synthase utilizing phosphatidylethanolamine and
754   phosphatidylglycerol as substrates. Proceedings of the National Academy of
755   Sciences 109:16504-16509.
756   56.   Pluschke G, Hirota Y, Overath P. 1978. Function of phospholipids in *Escherichia*
757   *coli*. Characterization of a mutant deficient in cardiolipin synthesis. Journal of
758   Biological Chemistry 253:5048-5055.

57.   Nishijima S, Asami Y, Uetake N, Yamagoe S, Ohta A, Shibuya I. 1988. Disruption of
the *Escherichia coli cls* gene responsible for cardiolipin synthesis. Journal of
Bacteriology 170:775-780.

58.   Guo D, Tropp BE. 2000. A second *Escherichia coli* protein with CL synthase
activity. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of
Lipids 1483:263-274.

59.   Kikuchi S, Shibuya I, Matsumoto K. 2000. Viability of an *Escherichia coli pgsA* null
mutant lacking detectable phosphatidylglycerol and cardiolipin. Journal of
Bacteriology 182:371-376.

60.   Suzuki M, Hara H, Matsumoto K. 2002. Envelope disorder of *Escherichia coli* cells
lacking phosphatidylglycerol. Journal of Bacteriology 184:5418-5425.

61.   Shiba Y, Yokoyama Y, Aono Y, Kiuchi T, Kusaka J, Matsumoto K, Hara H. 2004.
Activation of the Rcs signal transduction system is responsible for the
thermosensitive growth defect of an *Escherichia coli* mutant lacking
phosphatidylglycerol and cardiolipin. J Bacteriol 186:6526-35.

62.   Grabowicz M, Silhavy TJ. 2017. Redefining the essential trafficking pathway for
outer membrane lipoproteins. Proceedings of the National Academy of Sciences
114:4769-4774.

63.   Cronan JE, Jr., Gelmann EP. 1973. An estimate of the minimum amount of
unsaturated fatty acid required for growth of *Escherichia coli*. J Biol Chem
248:1188-95.

64.   Feng Y, Cronan JE. 2011. Complex binding of the FabR repressor of bacterial
unsaturated fatty acid biosynthesis to its cognate promoters. Molecular
Microbiology 80:195-218.

65.   Feng Y, Cronan JE. 2009. *Escherichia coli* unsaturated fatty acid synthesis:
Complex transcription of the *fabA* gene and *in vivo* identification of the essential
reaction catalyzed by FabB. Journal of Biological Chemistry 284:29526-29535.

66.   Henry MF, Cronan JE. 1991. *Escherichia coli* transcription factor that both
activates fatty acid synthesis and represses fatty acid degradation. Journal of
Molecular Biology 222:843-849.

67.   Henry MF, Cronan JE. 1992. A new mechanism of transcriptional regulation:
Release of an activator triggered by small molecule binding. Cell 70:671-679.

68.   Rock CO, Jackowski S. 1985. Pathways for the incorporation of exogenous fatty
acids into phosphatidylethanolamine in *Escherichia coli*. J Biol Chem 260:12720-
4.

69.   Heath RJ, Rock CO. 1996. Roles of the FabA and FabZ β-hydroxyacyl-acyl carrier
protein dehydratases in *Escherichia coli* fatty acid biosynthesis. Journal of
Biological Chemistry 271:27795-27801.

70.   Ostrander DB, Sparagna GC, Amoscato AA, McMillin JB, Dowhan W. 2001.
Decreased Cardiolipin Synthesis Corresponds with Cytochromec Release in
Palmitate-induced Cardiomyocyte Apoptosis. Journal of Biological Chemistry
276:38061-38067.

71. Lewis RNAH, Zweytick D, Pabst G, Lohner K, McElhaney RN. 2007. Calorimetric,
X-Ray Diffraction, and Spectroscopic Studies of the Thermotropic Phase Behavior
and Organization of Tetramyristoyl Cardiolipin Membranes. Biophysical Journal
92:3166-3177.
72. De Siervo AJ. 1969. Alterations in the phospholipid composition of *Escherichia*
coli B during growth at different temperatures. J Bacteriol 100:1342-9.
73. Bogdanov M, Sun J, Kaback HR, Dowhan W. 1996. A Phospholipid Acts as a
Chaperone in Assembly of a Membrane Transport Protein. Journal of Biological
Chemistry 271:11615-11618.
74. Mitchell AM, Silhavy TJ. 2019. Envelope stress responses: balancing damage
repair and toxicity. Nature Reviews Microbiology 17:417-428.
75. Luevano-Martinez LA, Kowaltowski AJ. 2015. Phosphatidylglycerol-derived
phospholipids have a universal, domain-crossing role in stress responses. Arch
Biochem Biophys 585:90-97.
76. Fernandez M, Paulucci NS, Peppino Margutti M, Biasutti AM, Racagni GE,
Villasuso AL, Agostini E, Gonzalez PS. 2019. Membrane Rigidity and Phosphatidic
Acid (PtdOH) Signal: Two Important Events in Acinetobacter guillouiae SFC 500-
1A Exposed to Chromium(VI) and Phenol. Lipids 54:557-570.
77. Shibata A, Ikawa K, Shimooka T, Terada H. 1994. Significant stabilization of the
phosphatidylcholine bilayer structure by incorporation of small amounts of
cardiolipin. Biochimica et Biophysica Acta (BBA) - Biomembranes 1192:71-78.
78. Boscia AL, Treece BW, Mohammadyani D, Klein-Seetharaman J, Braun AR,
Wassenaar TA, Klösgen B, Tristram-Nagle S. 2014. X-ray structure,
thermodynamics, elastic properties and MD simulations of
cardiolipin/dimyristoylphosphatidylcholine mixed membranes. Chemistry and
Physics of Lipids 178:1-10.
79. Pennington ER, Fix A, Sullivan EM, Brown DA, Kennedy A, Shaikh SR. 2017.
Distinct membrane properties are differentially influenced by cardiolipin content
and acyl chain composition in biomimetic membranes. Biochimica et Biophysica
Acta (BBA) - Biomembranes 1859:257-267.
80. Lopes SC, Ivanova G, de Castro B, Gameiro P. 2018. Revealing cardiolipins
influence in the construction of a significant mitochondrial membrane model.
Biochimica et Biophysica Acta (BBA) - Biomembranes 1860:2465-2477.
81. Marr AG, Ingraham JL. 1962. Effect of temperature on the composition of fatty
acids in *Escherichia coli*. Journal of Bacteriology 84:1260-1267.
82. Garwin JL, Klages AL, Cronan JE. 1980. Beta-ketoacyl-acyl carrier protein
synthase II of *Escherichia coli*. Evidence for function in the thermal regulation of
fatty acid synthesis. Journal of Biological Chemistry 255:3263-3265.
83. de Mendoza D, Klages Ulrich A, Cronan JE. 1983. Thermal regulation of
membrane fluidity in *Escherichia coli*. Effects of overproduction of beta-ketoacyl-
acyl carrier protein synthase I. Journal of Biological Chemistry 258:2098-2101.

84.　Knapp BD, Huang KC. 2022. The effects of temperature on cellular physiology.
Annual Review of Biophysics 51:499-526.
85.　Bogdanov M, Umeda M, Dowhan W. 1999. Phospholipid-assisted refolding of an
integral membrane protein: Minimum structural features for
phosphatidylethanolamine to act as a molecular chaperone. Journal of Biological
Chemistry 274:12339-12345.
86.　Raetz CR, Whitfield C. 2002. Lipopolysaccharide endotoxins. Annu Rev Biochem
71:635-700.
87.　Conway T, Creecy JP, Maddox SM, Grissom JE, Conkle TL, Shadid TM, Teramoto J,
Miguel PS, Shimada T, Ishihama A, Mori H, Wanner BL. 2014. Unprecedented
high-resolution view of bacterial operon architecture revealed by RNA
sequencing. mBio 5:e01442-14.
88.　Thomason MK, Bischler T, Eisenbart SK, Förstner KU, Zhang A, Herbig A, Nieselt
K, Sharma CM, Storz G. 2015. Global transcriptional start site mapping using
differential RNA sequencing reveals novel antisense rnas in *Escherichia coli*.
Journal of Bacteriology 197:18-28.
89.　Adams PP, Baniulyte G, Esnault C, Chegireddy K, Singh N, Monge M, Dale RK,
Storz G, Wade JT. 2021. Regulatory roles of *Escherichia coli* 5' UTR and ORF-
internal RNAs detected by 3' end mapping. eLife 10:e62438.
90.　Karp PD, Ong WK, Paley S, Billington R, Caspi R, Fulcher C, Kothari A,
Krummenacker M, Latendresse M, Midford PE, Subhraveti P, Gama-Castro S,
Muñiz-Rascado L, Bonavides-Martinez C, Santos-Zavaleta A, Mackie A, Collado-
Vides J, Keseler IM, Paulsen I. 2018. The EcoCyc database. EcoSal Plus
8:10.1128/ecosalplus.ESP-0006-2018.
91.　Huerta AM, Collado-Vides J. 2003. Sigma70 promoters in *Escherichia coli*: specific
transcription in dense regions of overlapping promoter-like signals. Journal of
Molecular Biology 333:261-278.
92.　Dowhan W, Bogdanov M. 2009. Lipid-dependent membrane protein
topogenesis. Annu Rev Biochem 78:515-40.
93.　Bogdanov M, Xie J, Heacock P, Dowhan W. 2008. To flip or not to flip: lipid-
protein charge interactions are a determinant of final membrane protein
topology. J Cell Biol 182:925-35.
94.　Bogdanov M, Dowhan W. 2012. Lipid-dependent generation of dual topology for
a membrane protein. J Biol Chem 287:37939-48.
95.　Bogdanov M, Dowhan W, Vitrac H. 2014. Lipids and topological rules governing
membrane protein assembly. Biochim Biophys Acta 1843:1475-88.
96.　Ramezanifard R, Golubeva YA, Palmer AD, Slauch JM. 2023. TamAB is regulated
by PhoPQ and functions in outer membrane homeostasis during *Salmonella*
pathogenesis. J Bacteriol 205:e0018323.
97.　Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, Doncheva NT,
Legeay M, Fang T, Bork P, Jensen LJ, von Mering C. 2021. The STRING database in

883   2021: customizable protein-protein networks, and functional characterization of
884   user-uploaded gene/measurement sets. Nucleic Acids Res 49:D605-d612.

98.   Grasekamp KP, Beaud B, Taib N, Audrain B, Bardiaux B, Rossez Y, Izadi-Pruneyre
N, Lejeune M, Trivelli X, Chouit Z, Guerardel Y, Ghigo J-M, Gribaldo S, Beloin C.
2023. Bridges instead of boats? The Mla system of diderm Firmicute *Veillonella*
parvula reveals an ancestral transenvelope core of phospholipid trafficking.
bioRxiv doi:10.1101/2023.06.30.547184:2023.06.30.547184.

99.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita
M, Wanner BL, Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame,
single-gene knockout mutants: the Keio collection. Mol Syst Biol 2:2006.0008-
2006.0008.

100.   Silhavy TJ, Berman ML, Enquist LW. 1984. Experiments with gene fusions. Cold
Spring Harbor Laboratory.

101.   Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. Proceedings of the National Academy
of Sciences 97:6640-6645.

102.   MacRitchie DM, Ward JD, Nevesinjac AZ, Raivio TL. 2008. Activation of the Cpx
envelope stress response down-regulates expression of several locus of
enterocyte effacement-encoded genes in enteropathogenic *Escherichia coli*.
Infection and Immunity 76:1465-1475.

103.   Wong JL, Vogt SL, Raivio TL. 2013. Using reporter genes and the *Escherichia coli*
ASKA overexpression library in screens for regulators of the Gram negative
envelope stress response. Methods Mol Biol 966:337-57.

104.   Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-
evolved microbes from next-generation sequencing data using breseq. Methods
Mol Biol 1151:165-88.

105.   Bailey TL, Elkan C. 1994. Fitting a mixture model by expectation maximization to
discover motifs in biopolymers. Proc Int Conf Intell Syst Mol Biol 2:28-36.

106.   Bailey TL, Johnson J, Grant CE, Noble WS. 2015. The MEME Suite. Nucleic Acids
Res 43:W39-49.

107.   Singer M, Baker TA, Schnitzler G, Deischel SM, Goel M, Dove W, Jaacks KJ,
Grossman AD, Erickson JW, Gross CA. 1989. A collection of strains containing
genetically linked alternating antibiotic resistance elements for genetic mapping
of Escherichia coli. Microbiol Rev 53:1-24.

108.   Rai AK, Carr JF, Bautista DE, Wang W, Mitchell AM. 2021. ElyC and cyclic
enterobacterial common antigen regulate synthesis of phosphoglyceride-linked
enterobacterial common antigen. mBio 12:e02846-21.

109.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. Nature
Biotechnology 37:907-915.

110. Liao Y, Smyth GK, Shi W. 2013. featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. Bioinformatics
30:923-930.
111. Robinson MD, McCarthy DJ, Smyth GK. 2009. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics
26:139-140.
112. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. limma
powers differential expression analyses for RNA-sequencing and microarray
studies. Nucleic Acids Research 43:e47-e47.
113. Sparagna GC, Johnson CA, McCune SA, Moore RL, Murphy RC. 2005.
Quantitation of cardiolipin molecular species in spontaneously hypertensive
heart failure rats using electrospray ionization mass spectrometry. J Lipid Res
46:1196-204.
114. Hryc CF, Mallampalli VKPS, Bovshik EI, Azinas S, Fan G, Serysheva II, Sparagna GC,
Baker ML, Mileykovskaya E, Dowhan W. 2023. Structural insights into cardiolipin
replacement by phosphatidylglycerol in a cardiolipin-lacking yeast respiratory
supercomplex. Nature Communications 14:2783.

940
941

Figures and Tables

Figure 1. Deletion of *yhdP* and *fadR* causes synthetic cold sensitivity due to impairment
of TamB function. (A) The Δ*yhdP* Δ*fadR* strain had impaired growth at lower
temperatures. **(B)** EOPs were performed to confirm the growth defect of the Δ*yhdP* Δ*fadR*
strain. **(C-F)** EOPs were performed at 30 °C to assess cold sensitivity. **(C)** The growth of the
Δ*yhdP* Δ*fadR* strain was not affected by the loss of ECA (Δ*wecA*). **(D)** Neither a Δ*tamB*
Δ*fadR* strain nor a Δ*ydbH* Δ*fadR* strain exhibited cold sensitivity. **(E)** Deletion of *ydbH* did
not suppress cold sensitivity in the Δ*yhdP* Δ*fadR* strain. **(F)** Deletion of *tamB* completely
suppressed cold sensitivity in the Δ*yhdP* Δ*fadR* strain and the Rcs stress response was not
necessary for this suppression. **(G)** Growth curves were performed at 30 °C . The Δ*tamB*
and the Δ*yhdP* Δ*fadR* Δ*tamB* strain grew equally well, indicating suppression is unlikely to
result from slow growth. Quantitative data are shown as the mean of three biological
replicates ± the SEM. Images are representative of three independent experiments.

Figure 2. Increased cardiolipin levels are necessary for cold sensitivity of the Δ*yhdP*
Δ*fadR* strain. (A-B) Thin layer chromatography (TLC) of lipid extracts was performed for
the indicated cultures grown to log phase at 42 °C then transferred to 30 °C for 2 hours
before analysis. **(A)** Relative phospholipid levels were calculated. The Δ*yhdP* Δ*fadR* strain
shows increased levels of CL and a concomitant decrease in PG levels, increasing the ratio

of CL to PG (B). Data are averages of two to three biological replicates ± the SEM. For
ratios individual data points are indicated with open circles. **(C-G)** Growth effects of
altered phospholipid levels were investigated by EOP. Images are representative of three
independent experiments. **(C)** PgsA depletion strains in a background where *pgsA* is non-
essential were used to assay the effect of lowering PG and CL levels on cold sensitivity.
Arabinose induces expression of *pgsA* while glucose represses. Lowering PG and CL levels
partially suppressed cold sensitivity. **(D)** Deletion of the three CL synthases, *clsA*, *clsB*, and
clsC, suppressed cold sensitivity of the Δ*yhdP* Δ*fadR* strain. **(E)** Deletion of *clsA* also
suppressed cold sensitivity. **(F)** Deletion of *clsB* does not suppress cold sensitivity, while
deletion of *clsC* (F) does.
974
975

976
Figure 3. Cold sensitivity can be suppressed by increasing *fabA* expression. (A) The cold
sensitivity of a spontaneous suppressor of Δ*yhdP* Δ*fadR* cold sensitivity was tested by EOP.
The suppressor returned growth to the level of the Δ*fadR* mutant. **(B)** A diagram of the
promoter region of *fabA* is shown with its transcriptional start sites (blue arrows) and
regulator binding sites (boxes). FadR acts as an activator while FabR acts as a repressor.
The motif of the FabR binding site, generated using MEME, is shown with the sequences
of the parent strain and of Suppressor 1. **(C)** The P$_{fabA}$(-19G>A) mutation was linked to a
Tn*10* and transferred to the Δ*yhdP* Δ*fadR* strain by transduction. The Δ*yhdP* Δ*fadR* strain
with P$_{fabA}$(-19G>A) was not cold sensitivity, confirming suppression. **(D)** *fabA* RNA levels
are shown. Bars are the mean of three biological replicates ± the SEM. Individual data
points are shown as open circles. ** $p<0.005$, *** $p<0.0005$ by quasi-linear F-test. Data
are from the RNA-seq experiment in Figure 5 and *fabA* data are also shown in Figure 5F.
(E) Luciferase reporters of wild-type and mutant *fabA* promoter activity were
constructed. The mutation caused similar fold increases in P$_{fabA}$ activity in all strain
backgrounds. Data are luminescence relative to OD$_{600}$ and are shown as the average of
two biological replicates ± the SEM.
993

Figure 4. Decreasing fatty acid saturation suppresses Δ*yhdP* Δ*fadR* cold sensitivity. (A-C) Lipid were extracted from cells grown to OD$_{600}$ 0.2 at 42 °C then shifted to 30 °C for 2 hours before harvest. LC/MS was performed on lipid extracts using absolute quantification and the percentage of each specific species of PE (A), PG (B), and CL (C) was calculated using the sum of all molecular species detected. CL is not shown for Δ*clsABC* and Δ*clsA* strains due to low CL levels. Suppressor 1 trends towards decreased saturation compared to its parent strain (Δ*yhdP* Δ*fadR*). Data for phospholipids without unsaturations are shown in grey, with one unsaturation in blue, with two unsaturations in green, with three unsaturations in orange. Data for phospholipids with longer acyl chains are shown in darker colors. Data are the average of three biological replicates ± the SEM. Phospholipids with odd chain lengths and unsaturations may have either odd chain or cyclopropyl fatty acids. **(D)** EOPS were performed on media supplemented with oleic acid or a vehicle control. Oleic acid suppresses the cold sensitivity of the Δ*yhdP* Δ*fadR* strain. **(E)** Growth curves of the indicated strains were performed in the presence of oleic

1010   acid, stearic acid, or a vehicle control. Oleic acid suppressed the cold sensitivity of the
1011   Δ*yhdP* Δ*fadR* strain while stearic acid did not. Data are the average of three biological
1012   replicates ± the SEM.
1013

Figure 5. Transcriptional landscape of the Δ*yhdP* Δ*fadR* and suppressed strain. Three
biological replicates of the indicated strains were grown at 42 °C to mid-log then
transferred to 30 °C for 30 minutes before harvesting RNA and performing RNA-seq.
Differential expression was calculated based on a greater than 2-fold change in expression
and a p value of less than 0.05 by quasi-linear F-test. **(A)** PCA was performed on the
expression of all genes differentially expressed between any groups of samples.
Suppressor 1 and the Δ*yhdP* Δ*fadR* strain grouped closer together than to the wild-type
strain, although Suppressor 1 was closer to wild type than the Δ*yhdP* Δ*fadR* strain. **(B-D)**
Volcano plots show the average fold change and significance of expression changes
between the Δ*yhdP* Δ*fadR* strain and wild-type MG1655 (B), Suppressor 1 and wild-type
MG1655 (C), and Suppressor 1 and the Δ*yhdP* Δ*fadR* strain (D). Genes with changes
greater than 2-fold are shown in blue, while genes with changes less than 2-fold are
shown in grey. Genes with a q-value (false discovery rate) of less than 0.05 are shown in
yellow. Blue lines indicate a 2-fold change and dotted line indicates p>0.05. The number
of up or down regulated genes is indicated in blue. The names of the 10 genes with the
largest changes are called out. More genes were up-regulated then down regulated in the
Δ*yhdP* Δ*fadR* strain and Suppressor 1 vs. wild type. More genes were down regulated in

Suppressor 1 compared to the Δ*yhdP* Δ*fadR* strain. **(E)** Relative expression of genes in the
CL synthesis pathway is shown as averages ± the SEM and individual data points. No
significant changes were evident. **(F)** Relative expression of genes in the fatty acid
synthesis and degradation pathways is shown as averages ± the SEM and individual data
points. A general normalizing of these pathways occurred in Suppressor 1 compared to
the Δ*yhdP* Δ*fadR* strain; however, the most significant change was in the expression of
fabA. * $p<0.05$, ** $p<0.005$, *** $p<0.0005$ by quasi-linear F-test.
1039

1040

Figure 6. Model for the role of YhdP and TamB in phospholipid transport. Our data can be explained by the following model of phospholipid transport. **(A)** In normal conditions, YhdP is mainly responsible for the transport of phospholipids with more saturated fatty acid chains between the IM and OM. TamB shows a strong preference for transporting phospholipids with more unsaturated fatty acids, with YdbH playing a minor role in phospholipid transport. **(B)** In the Δ*yhdP* Δ*fadR* strain, YhdP is not available to transport more saturated phospholipids and the proportion of both saturated fatty acids and cardiolipin is increased, especially at low temperature. The fluidity of the membrane is also lessened by the low temperature. In these conditions, transport of phospholipids between the membranes is impeded, due to clogging of TamB by a combination of less saturated phospholipids and/or bulky and highly disordered CL. This leads to a lethal lack of phospholipids in the OM and overabundance of phospholipids in the IM. **(C)** When *tamB* is deleted in the Δ*yhdP* Δ*fadR* strain, the clogging of TamB is relieved, and the cell relies on YdbH for phospholipid transport overcoming the cold sensitivity. PL: phospholipid