1

Genetic evidence for functional diversification of gram-negative intermembrane phospholipid transporters

Ashutosh K. Rai[a], Katsuhiro Sawasato[b], Haley C. Bennett[a], Anastasiia Kozlova[b], Genevieve C. Sparagna[c], Mikhail Bogdanov[b], and Angela M. Mitchell[a,#]

[a] Department of Biology, Texas A&M University, College Station, Texas, USA

[b] Department of Biochemistry and Molecular Biology, McGovern Medical School at The University of Texas Health Science Center at Houston, Houston, Texas, USA

[c] Department of Medicine, Division of Cardiology, University of Colorado Anschutz Medical Campus, Aurora, Colorado, USA

Running title: Phospholipid transporter functional differentiation

Address correspondence to Angela M. Mitchell, amitchell@bio.tamu.edu

Keywords: phospholipid trafficking, gram-negative envelope, outer membrane, lipid metabolism, cold sensitivity

Abstract

The outer membrane of Gram-negative bacteria is a barrier to chemical and physical stress. Phospholipid transport between the inner and outer membranes has been an area of intense investigation and, in *E. coli* K-12, it has recently been shown to be mediated by YhdP, TamB, and YdbH, which are suggested to provide hydrophobic channels for phospholipid diffusion, with YhdP and TamB playing the major roles. However, YhdP and TamB have different phenotypes suggesting distinct functions. We investigated these functions using synthetic cold sensitivity (at 30 °C) caused by deletion of *yhdP* and *fadR*, a transcriptional regulator controlling fatty acid degradation and unsaturated fatty acid production, but not by Δ*tamB* Δ*fadR* or Δ*ydbH* Δ*fadR*,. Deletion of *tamB* suppresses the Δ*yhdP* Δ*fadR* cold sensitivity suggesting this phenotype is related to phospholipid transport. The Δ*yhdP* Δ*fadR* strain shows a greater increase in cardiolipin upon transfer to the non-permissive temperature and genetically lowering cardiolipin levels can suppress cold sensitivity. These data also reveal a qualitative difference between cardiolipin synthases in *E. coli*, as deletion of *clsA and clsC* suppresses cold sensitivity but deletion of *clsB* does not despite lower cardiolipin levels. In addition to increased cardiolipin, increased fatty acid saturation is necessary for cold sensitivity and lowering this level genetically or through supplementation of oleic acid suppresses the cold sensitivity of the Δ*yhdP* Δ*fadR* strain. Although indirect effects are possible, we favor the parsimonious hypothesis that YhdP and TamB have differential substrate transport preferences, most likely with YhdP preferentially transporting more saturated phospholipids and TamB preferentially transporting more unsaturated phospholipids. We envision cardiolipin contributing to this transport preference by sterically clogging TamB-mediated transport of saturated phospholipids. Thus, our data provide a potential mechanism for independent control of the phospholipid composition of the inner and outer membranes in response to changing conditions.

Author Summary

Gram-negative bacteria possess a highly impermeable outer membrane, which protects against environmental stress and antibiotics. Outer membrane phospholipid transport remained mysterious until YhdP, TamB, and YdbH were recently implicated in phospholipid transport between the inner and outer membranes of *E. coli*. Similar roles for YhdP and/or TamB have been suggested in both closely and distantly related gram-negative bacteria. Here, given the transporters' apparent partial redundancy, we investigated functional differentiation between YhdP and TamB. Our data suggest YhdP and TamB have differential involvement with fatty acid and phospholipid metabolism. In fact, transport of higher than normal levels of cardiolipin and saturated phospholipids in the absence of YhdP and presence of TamB at a non-permissive temperature is lethal. We suggest a model where the functions of YhdP and TamB are distinguished by phospholipid transport preference with YhdP preferentially transporting more saturated phospholipids and TamB more unsaturated phospholipids. Cardiolipin headgroup specificity may contribute transport inhibition due to its bulky nature inhibiting the passage of other phospholipids. Diversification of function between YhdP and TamB provides a mechanism for regulation of phospholipid composition, and possibly the mechanical strength and permeability of the outer membrane, and so the cell's intrinsic antibiotic resistance, in changing environmental conditions.

Introduction

The gram-negative bacterial cell envelope has an outer membrane (OM) that sits outside the aqueous periplasm and peptidoglycan cell wall. The OM provides a barrier against various environmental stresses including toxic molecules such as antibiotics and osmotic pressure (1-6). Unlike the inner membrane (IM), a phospholipid bilayer, the OM is largely composed of phospholipids (mainly phosphatidylethanolamine (PE) (7)) in its inner leaflet and LPS (lipopolysaccharide) in its outer leaflet (2, 8). However, both membranes are asymmetric as the IM has different phospholipid compositions in its inner versus outer leaflets (9). Outer membrane proteins (OMPs) form a network across the cell surface interspersed with phase separated LPS patches (10). Lipoproteins are generally anchored in the inner (i.e., periplasmic) leaflet of the OM (11).

OM components are synthesized in the cytoplasm or IM and transported to the OM. OMP (12), lipoprotein (11), and LPS (13) transport pathways are well defined. However, until recently, intermembrane phospholipid transport (between the IM and OM), especially anterograde transport from the IM to the OM, has remained very poorly understood. Phospholipids are synthesized at the IM's inner leaflet (14) and rapid, bidirectional intermembrane phospholipid transport occurs (15), even without ATP (16). Phospholipid transport has a relaxed specificity, allowing transport of non-native lipids (17, 18); however, intermembrane phospholipid composition differences are maintained (7, 19-22). For instance, the OM is enriched for PE and its saturated species compared to the IM outer leaflet (7, 9, 20-22); however, phosphatidylglycerol (PG) and cardiolipin (CL) can be evenly distributed between the IM and OM when the cellular level of PE is greatly reduced (17), suggesting a maintenance of membrane charge balance. This headgroup redistribution is accompanied by a concomitant fatty acid redistribution evidenced by an accumulation of palmitic acid (C16:0) and cyclopropane derivatives of palmitoleic acid (C16:1) and *cis*-vaccenic acid (C18:1). In contrast, the 1,2-diglyceride that accumulates in mutants lacking DgkA (diglyceride kinase) predominantly associates with the IM (23), suggesting existence of a mechanism of discrimination of lipid transfer to the OM that does not recognize diglyceride molecules and contributes to the diversification of distribution of polar and acyl groups between the IM and OM. This establishment and maintenance of defined lipid topography between the IM and the OM must involve the coordinated biosynthesis and balanced bidirectional trafficking of specific phospholipids across the IM and from the IM to the OM (24).

It has been demonstrated that anterograde phospholipid transport is mediated by a high-flux, diffusive (i.e., concentration gradient, not energy dependent) system in *Escherichia coli* (25). A retrograde trafficking pathway, the Mla system, removes mislocalized phospholipids from the OM outer leaflet and returns them to the IM (26). A gain-of-function allele, *mlaA** (27, 28), opens a channel allowing phospholipids to mislocalize to the cell surface by flowing from the OM's inner to outer leaflet (27, 29), resulting in a PldA (OM phospholipase A)-mediated attempt to reestablish OM asymmetry by increasing LPS

production, OM vesiculation, and increased flow of phospholipids to the OM (27, 28).
When nutrients are depleted, inhibiting phospholipid production, cells lyse due to loss of
IM integrity (16, 18). Single cell imaging of *mlaA** cells confirmed an aberrant lipid flow
from the IM to the OM at a fast rate. Loss of YhdP, an inner membrane protein, could
significantly slow phospholipid flow to the OM resulting in loss of OM integrity before IM
rupture (25), suggesting YhdP might play a role in intermembrane phospholipid transport.

Recently, YhdP and two homologs, TamB and YdbH, were demonstrated to be
intermembrane phospholipid transporters (30, 31). Ruiz, *et al*. investigated AsmA family
proteins and found Δ*yhdP* Δ*tamB* mutants showed synthetic OM permeability and
stationary phase lysis (30). Moreover, when combined, Δ*yhdP*, Δ*tamB*, and Δ*ydbH* are
synthetically lethal. Genetic interactions of these mutants with *mlaA* and *pldA* show their
involvement in OM phospholipid homeostasis. Predicted structures of YhdP, TamB, and
YdbH (32, 33), as well as a partial crystal structure of TamB (34), have a β-taco fold domain
with a hydrophobic pocket, resembling eukaryotic lipid transporters (35-38). These
structural findings are another clue that suggests YhdP, TamB, and YdbH are involved in
phospholipid transport. Douglass, *et al.* confirmed the synthetic lethality of Δ*yhdP*,
Δ*tamB*, and Δ*ydbH* and demonstrated a Δ*tamB* Δ*ydbH* mutant with reduced *yhdP*
expression had decreased amounts of OM phospholipids, directly demonstrating their
involvement in intermembrane phospholipid transport or its regulation (31). Structural
studies have shown that YhdP is long enough to span the periplasmic space and molecular
dynamics indicate that the C-terminus of YhdP can insert into the OM to allow
phospholipid transfer between the membranes (39). This study also directly
demonstrates that a phosphate containing substrate, putatively assigned to be
phospholipids, can be crosslinked to the hydrophobic groove of YhdP (39). The role of
YhdP, TamB, and YdbH in phospholipid transport may be widely conserved as recent
evidence implicates TamB in anterograde phospholipid transport in *Veillonella parvula*, a
diderm Fermicute (40) and YhdP, TamB, YdbH, and PA4735 in intermembrane
phospholipid transport in *Pseudomonas aeruginosa* (41).

Nevertheless, why there are three separate intermembrane phospholipid transport
proteins and the functional interactions between these proteins remains unanswered,
although it is clear the proteins are not fully redundant or functionally equivalent. The
conditional expression of *yhdP* alone, without the presence of *ydbH* and *tamB*, fully
complements neither growth phenotypes nor a normal level of OM phospholipids,
suggesting transport function is still impaired in this strain (31). In addition, YdbH seems
to play a more minor role than YhdP and TamB (30, 31) and the screen identifying Δ*yhdP*
as slowing *mlaA**-dependent lysis did not identify Δ*tamB* or Δ*ydbH* (25). Moreover, we
previously identified a role for YhdP in stationary phase SDS (sodium dodecyl sulfate)
resistance and modulating cyclic enterobacterial common antigen activity (42, 43) not
shared by TamB and YdbH (**Fig. S1AB**) (30, 31, 42, 43). Similarly, TamB, in conjunction with

TamA, has been suggested to play a role in OM insertion of some "complicated" β-barrel
OMPs such as autotransporters and usher proteins (34, 44-46).

Here, we investigated the differentiation of YhdP and TamB function and identified
synthetic cold sensitivity in a strain with Δ*yhdP* and Δ*fadR*, a transcriptional regulator of
fatty acid biosynthesis and degradation, acting as a transcriptional switch between these
pathways in response to acyl-CoA (47). FadR is necessary for normal levels of unsaturated
fatty acids to be synthesized (48). The cold sensitivity was unique to the Δ*yhdP* Δ*fadR*
strain and was suppressed by loss of TamB, demonstrating that the sensitivity is related
to phospholipid transport. We found that the Δ*yhdP* Δ*fadR* mutant had a greater increase
in levels of a specific phospholipid, cardiolipin (CL) during growth at the non-permissive
temperature, and decreasing CL suppressed cold sensitivity. Furthermore, increasing
unsaturated fatty acid levels suppressed cold sensitivity. Overall, our data are consistent
with a model where TamB and YhdP transport functions are differentiated based on
phospholipid fatty acid saturation state, with TamB transporting more unsaturated
phospholipids and YhdP transporting more saturated phospholipids, although indirect
effects may contribute to the phenotype. Phospholipid species or head group can also
contribute to transport specificity, since higher levels of dianionic CL are necessary for the
cold sensitivity phenotype, potentially due to functional inhibition of TamB by clogging of
its hydrophobic groove.

170
171
Results

Loss of FadR and YhdP results in cold sensitivity suppressed by loss of TamB. To
investigate YhdP-TamB function differentiation, we created a Δ*yhdP* Δ*fadR* strain, as FadR
is a major regulator of lipid homeostasis (47). A deletion strain of *fadR* alone is expected
to have decreased fatty acid synthesis, increased fatty acid degradation, and increased
fatty acid saturation (47, 48). Surprisingly, Δ*yhdP* Δ*fadR* cultures grown at 30 °C lagged for
more than 8 hours, followed by highly variable growth, suggesting suppressor outgrowth
(**Fig.1A**). The growth defect was decreased at 37 °C and completely absent at 42 °C. To
confirm this cold sensitivity, we estimated efficiency of plating (EOP) and confirmed the
severe cold intolerance of the Δ*yhdP* Δ*fadR* strain (5 logs at 30 °C and 3 logs at 37 °C) (**Fig.**
1B). Although, enterobacterial common antigen is necessary for Δ*yhdP*'s OM permeability
phenotypes (43), the Δ*yhdP* Δ*fadR* strain's cold sensitivity was unchanged when
enterobacterial common antigen was not present (Δ*wecA*) (49) (**Fig. 1C**).

We hypothesized the Δ*yhdP* Δ*fadR* strain's cold sensitivity was due to impairment of
phospholipid transport and expected Δ*tamB* and/or Δ*ydbH* to show similar synthetic
phenotypes with Δ*fadR*. However, Δ*fadR* Δ*tamB* and Δ*fadR* Δ*ydbH* strains had no growth
defects (**Fig 1D**). Although Δ*ydbH* in the Δ*yhdP* Δ*fadR* strain did not alter cold sensitivity
(**Fig. 1E**), Δ*tamB* in the Δ*yhdP* Δ*fadR* background completely suppressed the cold
sensitivity (**Fig. 1F**). Consistent with previous observations for Δ*yhdP* Δ*tamB* (30), the

Δ*yhdP* Δ*fadR* Δ*tamB* strain was mucoid due to Rcs (regulator of capsule synthesis) stress
response activation and colanic acid capsule production. Deleting *rcsB*, an Rcs response
regulator, prevented mucoidy but did not affect suppression (**Fig. 1F**). We observed only
minimal growth phenotypes at 30 °C in the Δ*yhdP* Δ*fadR* Δ*tamB* strain (**Fig. 1G**), similar to
those of Δ*yhdP* Δ*tamB* (**Fig. S1C**). Thus, although Δ*fadR* does not suppress Δ*yhdP* Δ*tamB*
phenotypes, these data make slow growth an unlikely suppression mechanism. Together,
these data suggest the Δ*yhdP* Δ*fadR* strain has impaired phospholipid transport leading
synthetic cold sensitivity, which can be relieved by deletion of Δ*tamB*, shifting
phospholipid transport to YdbH or causing some regulatory change.

We next looked for envelope homeostasis alterations in the Δ*yhdP* Δ*fadR* strain. To
identify cell lysis and/or severe OM permeability defects at 30 °C, we assayed CPRG
(chlorophenol red-β-D-galactopyranoside) processing. CPRG must contact LacZ in the
cytoplasm or supernatant to release chlorophenol red (50). Compared to wild type or
single mutants, the Δ*yhdP* Δ*fadR* strain had increased CPRG processing (**Fig. S2A**),
demonstrating lysis or increased envelope permeability. When we assayed resistance to
molecules excluded by the OM (1, 42, 43, 51, 52) at 37 °C, a semi-permissive temperature,
the Δ*yhdP* Δ*fadR* strain showed sensitivity to vancomycin, a large scaffold antibiotic, and
SDS EDTA compared to single mutants (**Fig. S2B**). However, with bacitracin treatment or
EDTA treatment alone, the Δ*yhdP* Δ*fadR* strain phenocopied the Δ*fadR* single deletion
(**Fig. S2CD**). LPS levels were not altered in the Δ*yhdP*, Δ*fadR*, Δ*yhdP* Δ*fadR* or Δ*yhdP* Δ*fadR*
Δ*tamB* strains (**Fig. S2E**). OM asymmetry mutations (i.e., Δ*pldA*, Δ*mlaA*) (26-28, 53) did
not have suppressive or synthetic effects on cold sensitivity in the Δ*yhdP* Δ*fadR* strain (**Fig.**
S3AB), cold sensitivity is not due to OM asymmetry or the loss thereof. Overall, the
changes in OM permeability did not seem sufficient to cause lethality at lower
temperature, suggesting the phospholipid transport disruption affects both IM and OM
integrity.

Lowering cardiolipin suppresses cold sensitivity. We investigated phospholipid
composition of the wild-type, Δ*yhdP*, Δ*fadR*, and Δ*yhdP* Δ*fadR* strains grown at the
permissive temperature (42 °C) to log phase then downshifted to 30 °C. *E. coli*
phospholipid composition is generally 75% phosphatidylethanolamine (PE), 20%
phosphatidylglycerol (PG), and 5% cardiolipin (CL) with CL increasing in stationary phase
(14, 54). Of the three CL synthases, ClsA and ClsB synthesize CL from two PG molecules,
while ClsC synthesizes CL from one PG and one PE molecule, so CL and PG levels are
generally reciprocally regulated (55-58). For all strains, the levels of PE were similar (**Fig.**
2A, **Fig. S4A**). However, the Δ*yhdP* Δ*fadR* strain had increased CL and concomitant
decreased PG compared to the wild-type strain and compared to individual Δ*yhdP* and
Δ*fadR* mutants (**Fig. 2B**). PgsA synthesizes phosphatidylglycerol-phosphate and is the first
enzyme differentiating PG and PE biosynthesis. PgsA depletion would decrease both PG
and CL levels, reversing increased CL levels and exacerbating decreased PG levels in the
Δ*yhdP* Δ*fadR* strain. We tested the effect of PgsA depletion in a strain background (Δ*rcsF*
Δ*lpp*) where *pgsA* is non-essential (59-62). When *pgsA* expression was repressed (glucose

panel), the ΔyhdP ΔfadR cold sensitivity was partially suppressed (**Fig. 2C**), demonstrating
that decreased PG does not cause cold sensitivity and suggesting increased CL may be
necessary for cold sensitivity. The partial suppression may be due to incomplete depletion
of PG and CL or the adaptive response of a cell containing only PE in its membrane.
Induction of *pgsA* (arabinose panel) reversed the suppression. The cold sensitivity of the
ΔyhdP ΔfadR strain was lost when CL was fully depleted due to deletion of the genes for
all cardiolipin syntheses (ΔclsABC) or of the gene for the primary cardiolipin synthase
(ΔclsA) responsible for CL synthesis in exponential phase (**Fig. 2ADE**, **Fig. S4B**). A decrease
in LPS levels is not responsible for the suppression as LPS levels did not change in the
ΔyhdP ΔfadR ΔclsA strain (ΔyhdP ΔfadR ΔtamB). ΔclsB caused a smaller decrease in CL
levels, while ΔclsC did not cause a decrease (**Fig. 2AB**, **Fig. 4B**). Surprisingly, however,
ΔclsC completely suppressed the cold sensitivity, while ΔclsB did not (**Fig. 2FG**). Thus,
increased CL is necessary for cold sensitivity but CL levels alone cannot explain the
phenotype: another factor, such as a specific molecular form (e.g., acyl chain length,
saturation state, symmetry of acyl chain arrangement) or localization (e.g., poles vs.
midcell or inner vs. outer leaflet) of CL, is required.

Increased *fabA* expression suppresses cold sensitivity. To identify other factors involved
in the cold sensitivity, we applied a forward genetic approach by isolating spontaneous
suppressor mutants capable of growing at 30 °C and identified a suppressor mutant
(Suppressor 1) that restored growth at 30 °C (**Fig. 3A**), decreased CPRG processing (**Fig.**
S2A), and had very similar CL levels (**Fig. 2AB**, **Fig. S4A**) and LPS levels (**Fig. S2**) to the
ΔyhdP ΔfadR strain. We identified a point mutation in the *fabA* promoter region in
Suppressor 1 (**Fig. 3B**) and confirmed this mutation was sufficient for suppression (**Fig.**
3C). FabA is a dehydratase/isomerase that introduces *cis* unsaturation into fatty acids (14,
63) and *fabA* is expressed from two promoters controlled by FadR and FabR (64, 65). FadR
activates *fabA* expression, while FabR represses. Thus, *fadR* deletion would be expected
to lower *fabA* expression. The suppressor mutation was located in the *fabA* promoter
FabR binding site (**Fig. 3B**) and we hypothesized this mutation would increase *fabA*
expression. *fabA* mRNA levels were more than 4-fold lower in the ΔyhdP ΔfadR strain than
the wild-type strain (**Fig. 3D**). While lower than wild type, the suppressor mutant
increased *fabA* mRNA 1.6-fold over the ΔyhdP ΔfadR strain.

We constructed luciferase reporter plasmids with the wild-type *fabA* promoter region or
the *fabA* promoter region containing the suppressor mutation. The reporter with the
mutated promoter had higher activity in a wild-type, ΔfadR, and ΔfabR background (**Fig.**
3E). *fabA* transcription occurs from a promoter downstream of the FabR binding site (65-
67). However, when *fadR* is deleted, transcription shifts to the FabR-regulated promoter
within the FabR binding region (65-67). Thus, our data indicate the Suppressor 1 mutation
regulates the constitutive activity of the promoter located in the FabR binding site rather
than the affinity of FabR binding. To confirm the effect of changing the second promoter's
activity, we tested the effect of ΔfabR on cold sensitivity in the ΔyhdP ΔfadR strain and
observed partial suppression (**Fig. S5**). The smaller effect of ΔfabR compared to

Suppressor 1 may result from other gene expression changes or to the smaller relative
effect of Δ*fabR* on *fabA* expression (**Fig. 3E**). Overall, these data demonstrate increasing
fabA expression, which is necessary for unsaturated fatty acid biosynthesis, rescues cold
sensitivity in the Δ*yhdP* Δ*fadR* strain without changing CL levels.

Increasing unsaturated fatty acids relieves cold sensitivity. Given the effect of *fabA*
expression on the Δ*yhdP* Δ*fadR* cold sensitivity, we used liquid chromatography-
electrospray ionization mass spectrometry (LC/MS) to characterize the strains'
phospholipid saturation state (**Dataset S1**, **Fig. 4A-C, Fig. S6**). Representative spectra
demonstrating relative phospholipid composition are shown in **Fig. S7**. As expected,
Δ*fadR* caused increased fully saturated and monounsaturated PE and PG with a
concomitant decrease of diunsaturated PE (**Fig. 4AB, S6AB**). Similarly, the Δ*fadR* strain
demonstrated increased monounsaturated CL with a concomitant decrease in
triunsaturated CL (**Fig. 4C, S6C**). Compared to the Δ*fadR* strain, the Δ*yhdP* Δ*fadR* strain
had slightly decreased saturation for all three phospholipids, while still displaying more
phospholipid saturation than wild type or Δ*yhdP* strains (**Fig. 4A-C, S6C**). Suppressor 1
trended towards decreased saturation of all three phospholipids compared to the Δ*yhdP*
Δ*fadR* strain (**Fig. 4A-C, S6C**). We wondered whether the suppression difference between
Δ*clsC* and Δ*clsB* resulted from the CL saturation state. However, the Δ*yhdP* Δ*fadR* Δ*clsB*
and Δ*yhdP* Δ*fadR* Δ*clsC* strains had very similar CL profiles (**Fig. 4C, S6C**), suggesting
another qualitative difference affecting suppression, perhaps CL lateral or inter-leaflet
distribution in the cell. These strains also showed similar PE and PG saturation profiles to
their Δ*yhdP* Δ*fadR* parent (**Fig. 4AB, S6AB**).

We hypothesized increasing unsaturated fatty acids suppresses Δ*yhdP* Δ*fadR* cold
sensitivity and performed EOPs for cold sensitivity on media containing an unsaturated
fatty acid (oleic acid, C18:1 *cis*-9). Exogenous phospholipids can be taken up by *E. coli*,
attached to acyl-CoA, and incorporated into phospholipids (68). Oleic acid addition
suppressed the Δ*yhdP* Δ*fadR* strain's cold sensitivity (**Fig. 4D**). Supplementing oleic acid
increases unsaturated phospholipids and provides an exogenous fatty acid source. To
differentiate between saturation state and fatty acid availability, we compared treatment
with oleic acid and saturated stearic acid (C18:0). Only oleic acid, and not stearic acid,
suppressed the Δ*yhdP* Δ*fadR* strain's cold sensitivity (**Fig. 4E**), confirming increasing
unsaturated fatty acids, not fatty acid availability suppresses.

Suppression causes fatty acid metabolism and stress response alterations. In wild-type
cells, *fabA* overexpression does not change phospholipid saturation levels, demonstrating
the FabB enzyme is rate limiting for unsaturated fatty acid biosynthesis (69). We were
intrigued that Suppressor 1 altered saturation without directly effecting *fabB* expression.
We also wondered whether CL levels in the Δ*yhdP* Δ*fadR* strain are transcriptionally
regulated. Thus, we investigated the transcriptional landscape of the wild type, the Δ*yhdP*
Δ*fadR*, and Suppressor 1 strains after a 30-minute downshift of growth to 30 °C to
determine: (i) whether the Δ*yhdP* Δ*fadR* strain had altered *cls* gene expression; (ii)

whether Suppressor 1 had other transcriptional changes; and (iii) why increased *fabA*
expression decreased saturation. All differentially expressed genes (>2-fold change,
p<0.05) are listed in **Dataset S2**. Principle component analysis of these genes showed the
closest relation between Δ*yhdP* Δ*fadR* strain and Suppressor 1, while the wild-type strain
was more distantly related (**Fig. 5A**, **S8A**). Nevertheless, Suppressor 1 was more closely
related to wild type than was the Δ*yhdP* Δ*fadR* strain. Most differentially regulated genes
between Δ*yhdP* Δ*fadR* and wild type and Suppressor 1 and wild type were upregulated
(**Fig. 5BC**, **S8BC**). Many of the most highly regulated genes are FadR-regulon members.
The majority of differentially regulated genes in Suppressor 1 compared to Δ*yhdP* Δ*fadR*
are downregulated (**Fig. 5D, S8D**). These genes are involved in many cellular pathways
and likely reflect the decreased cellular stress (**Table S2**). Indeed, many external stress
response genes are more enriched in Δ*yhdP* Δ*fadR* than in Suppressor 1 (**Fig. S9A**).

No significant expression changes occurred in the CL biosynthesis pathway (**Fig. 5E**) or
other phospholipid biosynthesis genes (**Fig. S9B**). The increased cardiolipin in Δ*yhdP*
Δ*fadR* and Suppressor 1 may be post-transcriptional or due to a change in a regulatory
pathway not yet altered after 30 minutes at 30 °C (**Fig. S9C**). Levels of mitochondrial CL
can be altered due to differences in substrate binding affinity of the CL synthase based on
saturation, with increased saturation decreasing CL synthesis (70), and it may be that a
similar pathway operates here, albeit in a different direction. We examined the
expression of genes in fatty acid synthesis and degradation, many of which are members
of the FadR regulon. In Δ*yhdP* Δ*fadR* compared to wild type, many fatty acid biosynthesis
genes significantly decreased while many fatty acid degradation genes significantly
increased (**Fig. 5F**), consistent with Δ*fadR* effects. While *fabA* expression is decreased,
fabB expression is not changed, explaining why increased *fabA* expression causes
decreased saturation in Suppressor 1. Overall, our data demonstrate the Δ*yhdP* Δ*fadR*
strain's cold sensitivity is due to impaired phospholipid trafficking, and increased
phospholipid saturation and cardiolipin are necessary for this functional impairment.

340
341

Discussion
The identification of YhdP, TamB, and YdbH as putative intermembrane phospholipid
transporters (30, 31) posed the tantalizing question: why is having three transporters
advantageous? Differential phenotypes between the transporters suggest that they
possess specialized functions (25, 30, 31, 34, 43-46). Here, we demonstrate each protein
plays a distinguishable role in phospholipid transport that can be differentiated based on
genetic interactions with lipid metabolism. Disruption of *fadR* and *yhdP* function causes
synthetic cold sensitivity not shared by *tamB* or *ydbH*. This sensitivity can be suppressed
by removing TamB. This phenotype involves both increased levels of CL and saturated
fatty acids, and decreasing amounts of either suppresses. In addition, our data
demonstrate that CL synthesized by ClsB and ClsC is qualitatively different—resulting in
differential suppression phenotypes—likely due to differences in CL localization, CL-

mediated, leaflet-specific changes in lipid packing order (9, 24), localized membrane
fluidity (71), or phospholipid transporter interaction.

Based on the currently available data, we cannot rule out the possibility that the
suppressors we have identified for the Δ*yhdP* Δ*fadR* cold sensitivity act indirectly (e.g.,
through regulatory or adaptive changes). However, we now suggest a simple model for
the differentiation of function between YhdP and TamB that fully explains our data
without invoking additional, more complex interpretations (**Fig. 6A-C**). In this hypothesis,
the functions of YhdP and TamB are diversified based on the acyl carbon saturation level
preference of the phospholipids they transport: YhdP transports mainly phospholipids
with more saturated fatty acids, while TamB transports phospholipids with more
unsaturated fatty acids (**Fig. 6A**). Our data and previous data (30, 31) agree YdbH plays a
relatively minor role. In the Δ*yhdP* Δ*fadR* strain, YhdP's absence forces more saturated
phospholipids to be transported by TamB. At 30 °C, TamB becomes "clogged" by
phospholipids having more saturated fatty acyl side chains, impeding transport (**Fig. 6B**).
Bulky and intrinsically disordered CL molecules could contribute to the clogging sterically
by obstructing the lipid channel and precluding it from translocating other lipid substrates
(i.e., PE). We find it interesting that the phenotype of the Δ*yhdP* Δ*fadR* strain is most
severe at low temperature, leading to cold sensitivity. This cold-specific effect may occur
due to (i) temperature-dependent decreased membrane fluidity and higher lipid packing
order involving lower expression of FabF, which is responsible for increased diunsaturated
phospholipids at lower temperature (**Fig. 5F**, **Dataset 3**)(14); (ii) higher relative levels of
CL (72); (iii) higher amounts of per cell phospholipids at lower temperatures (72)
contributing to increased diffusive flow rate; or, (iv) to thermodynamic properties altering
TamB transport specificity and/or substrate behavior contributing to efficiency of
substrate loading or transport.

One interesting aspect of our data is that the presence of TamB, with perhaps inhibited
function, causes a worse phenotype than the absence of TamB. Furthermore, the
phenotype of a Δ*yhdP* Δ*fadR* Δ*ydbH* strain is not measurably worse than that of the Δ*yhdP*
Δ*fadR* strain. Together, these data suggest that YdbH is able to successfully transport
phospholipids in the Δ*yhdP* Δ*fadR* strain only when TamB is absent. There are several
possible mechanisms for this. The first is that YhdP and TamB are the primary transporters
and are the preferred interaction partners for other, as of yet unidentified interaction
partners (e.g., putative IM proteins facilitating phospholipid transfer to YhdP and TamB).
Another possibility is that there is direct toxicity caused by TamB clogging, perhaps
through leaking of micellar phospholipids into the periplasm or through stress response
overactivation. A third possibility is that YdbH levels are insufficient in the presence of
YhdP and TamB to facilitate phospholipid transport and *ydbH* is upregulated in the
absence of the other phospholipid transporters. This will be an interesting area for future
investigations.

Several pieces of evidence suggest dysregulation of phospholipid transport not just changes to OM phospholipid composition, and so permeability or physical strength, cause the cold sensitivity phenotype of the Δ*yhdP* Δ*fadR* strain. First, the cold sensitivity is unaffected by removing enterobacterial common antigen, which suppresses the membrane permeability of a Δ*yhdP* strain (**Fig. 1C**, (43)), and the OM permeability phenotypes of the Δ*yhdP* Δ*fadR* are not severe (**Fig. S2**). Second, the cold sensitivity is suppressed by deletion of Δ*tamB* (**Fig. 1FG**). Δ*yhdP* Δ*tamB* strains have much greater OM permeability (30) than the Δ*yhdP* or Δ*yhdP* Δ*fadR* strains. Third, CL has been shown to have a normalizing, bidirectional, and leaflet-specific effect on lipid packing order (9, 73-76) and membrane fluidity (77-80) opposite of or similar to that of cholesterol, respectively, and so is unlikely to contribute to a phenotype dependent solely on OM membrane fluidity. Thus, the cold sensitivity phenotype correlates with the impairment of phospholipid trafficking and the likely imbalance of phospholipids between the IM and OM and not with the permeability phenotypes of the OM. Nonetheless, it will be useful in the future to conduction comparative analysis of the phospholipid composition of pure IM and OM extracts (i.e., without cross contamination of the other membrane) at both permissive and non-permissive temperatures to directly link the cold sensitivity observed here with changes in phospholipid transport.

Although both fatty acid saturation and CL levels affect the Δ*yhdP* Δ*fadR* strain's cold sensitivity, evidence points to saturation rather than headgroup as the main factor separating YhdP and TamB function. Specifically, YhdP loss slows phospholipid transport from the IM to the OM in a *mlaA** mutant, which loses OM material due to vesiculation and lysis when phospholipid biosynthesis is insufficient to replace lost phospholipids (25, 27). CL only accounts for 5% of phospholipids in exponentially growing cells making it unlikely that the loss of CL transport could slow net phospholipid transport to the extent required to slow lysis.

Our model suggests the IM and OM saturation state could be independently regulated based on YhdP and TamB activity or levels. IM saturation is regulated by altering the composition of newly synthesized fatty acids, while OM saturation could be separately regulated by allowing increased transport of more saturated or unsaturated phospholipids as necessary. IM and OM acyl saturation has been studied in several strains of *E. coli* and in *Yersinia pseudotuberculosis* (7, 19-21) and higher saturated fatty acid levels were observed in the OM than in the IM, supporting discrimination of phospholipid transport based on saturation. The IM needs to maintain an appropriate level of fluidity to allow IM proteins to fold and diffuse, to allow diffusion of hydrophobic molecules, to regulate cellular respiration, and to maintain membrane integrity (84, 85). However, the OM is a permeability barrier, a load-bearing element that resists turgor pressure (1-6), and has an outer leaflet consisting of LPS (2, 8). LPS has saturated acyl chains and is largely immobile in *E. coli* (86) and a specific phospholipid saturation level may be necessary to match or counterbalance LPS rigidity. OM phospholipids may also play a role in the

physical strength of the membrane. Interestingly, previous work has demonstrated that
the tensile strength of the OM is lessened in a *yhdP* mutant (25).

Possessing two different phospholipid transporters with different saturation state
preferences would allow the cell to adapt OM saturation to changing conditions including
temperature shifts, chemical and physical insults, or changes in synthesis of other OM
components (e.g., LPS or OMPs). A recent study demonstrated in *Y. pseudotuberculosis*
the difference in saturation levels between the membranes is exaggerated at 37 °C
compared to 8 °C (21). Thus, conditions where IM and OM saturation are independently
regulated exist and this regulation is likely important for optimum fitness of the cells.
Transcriptional regulation of *yhdP* and *tamB* has not been thoroughly explored. It seems
likely *yhdP* transcription is quite complex with several promoters and transcriptional
termination sites (87-89). *tamB* is predicted to be in an operon with *tamA* (which codes
for an OMP necessary for TamB function) and *ytfP* (87, 90) and to have a σ^E-dependent
promoter (91). The σ^E stress response is activated by unfolded OMP accumulation, and
downregulates the production of OMPs and upregulates chaperones and folding
machinery (74). It is intriguing to speculate a change in OM phospholipid composition
might aid in OMP folding due to altered biophysical properties and/or lipoprotein
diffusion. In fact, this could explain the observations that some complicated OMPs fold
more slowly in the absence of TamB (34, 44-46). Proper phospholipid composition may
be necessary for efficient folding of these OMPs, as it is necessary for the correct folding
and topology of some IM proteins (73, 92-95). Recently, it has been demonstrated that
the *tamAB* operon is regulated by PhoPQ in *Salmonella enterica* Serovar Typhimurium
and that this regulation helps maintain OM integrity in the acidic phagosome
environment, perhaps due to indirect effects on OMP folding (96). Thus, changing
environments may necessitate changes in phospholipid composition that can be
accomplished through differential TamB and, perhaps YhdP, regulation.

The mechanism of phospholipid transfer between the IM and OM has been one of the
largest remaining questions in gram-negative envelope biogenesis and has remained an
area of intense investigation for decades. Furthermore, given the lack of knowledge of
this pathway, the role of phospholipid composition in the permeability barrier of the OM,
and so the intrinsic antibiotic resistance of gram-negative bacteria, remains unknown.
Recent work has begun to elucidate the intermembrane phospholipid transport pathway
by identifying YhdP, TamB, and YdbH as proteins capable of transporting phospholipids
between the membranes (25, 30, 31). Our work here suggests that YhdP and TamB may
have separate roles in phospholipid transport that are differentiated by their preference
for phospholipid saturation states and possibly lipid headgroup. Homologs of YhdP and
TamB are found throughout *Enterobacterales* and more distantly related members of the
AsmA family are widespread in gram-negative bacteria (97)*.* In fact, TamB has recently
been shown to play a role in anterograde intermembrane phospholipid transport in the
diderm Firmicute *Veillonella parvula* (40, 98) and four AsmA members play a role in
phospholipid transport in *Pseudomonas aeruginosa* (41). Structural predictions of YhdP

and TamB are strikingly similar between species (32, 33). Given this, it is quite likely similar
mechanisms of discrimination in intermembrane phospholipid transfer exist in other
species. The data we present here provide a framework for investigation of phospholipid
transport and mechanisms differentiating phospholipid transporter function in these
species.

479
480
Materials and Methods

Bacterial strains and Growth Conditions. Table S3 lists the strains used in this study. Most
knockout strains were constructed with Keio collection alleles using P1vir transduction
(99, 100). New alleles were constructed through λ-red recombineering (101), using the
primers indicated in **Table S4**. The FLP recombinase-FRT system was used to remove
antibiotic resistance cassettes as has been described (101). *E. coli* cultures were grown in
LB Lennox media at 42°C, unless otherwise mentioned. Where noted, LB was
supplemented with vancomycin (Gold Biotechnology), bacitracin (Gold Biotechnology),
SDS (Thermo Fisher Scientific), EDTA (Thermo Fisher Scientific), oleic acid (Sigma Aldrich),
stearic acid (Sigma Aldrich), arabinose (Gold Biotechnology), or glucose (Thermo
Scientific). For plasmid maintenance, media was supplemented with 25 µg/mL kanamycin
(Gold Biotechnology), 10 or 20 µg/mL chloramphenicol (Gold Biotechnology), or 10 µg/mL
tetracycline (Gold Biotechnology) as appropriate.

Plasmid construction. The pJW15-P$_{fabA}$, and pJW15-P$_{fabA}$(-19G>A) reporter plasmids were
constructed by amplifying the region upstream of *fabA* from -432 bp to +27 bp relative to
the annotated transcriptional start site from wild-type MG1655 or Suppressor 1,
respectively, using the Overlap-pJW15-*fabA* FP and RP primers (**Table S4**). pJW15 was
amplified using primers Overlap-pJW15-*fabA* FP1 and RP1. The pJW15 plasmid was a kind
of gift from Dr. Tracy Raivio (University of Alberta) (102, 103). The resulting fragments
were assembled using HiFi Assembly master mix (New England Biolabs) as per the
manufacturer's instructions.

Growth and Sensitivity Assays. For EOPs assays, cultures were grown overnight, serially
diluted, and plated on the indicated media. For *pgsA* depletion, overnight cultures grown
in arabinose were washed and diluted 1:100 into fresh LB with 0.2% glucose or arabinose
as indicated and grown to OD$_{600}$=1 before diluting and plating on media with glucose or
arabinose respective, as above. EOPs were incubated at 30 °C unless otherwise indicated.
Growth curves were performed at the indicated temperature as has been described (43),
except that starting cultures for growth curves were grown to early stationary phase
before inoculation of growth curves rather than overnight. For CPRG assays (50), strains
were streaked on LB plates supplemented with CPRG (40µg/ml) and IPTG (100 µM)
incubated overnight at 30°C.

Suppressor isolation and sequencing. Serially diluted cultures were spread on LB plates and incubated overnight at 30°C. Colonies that grew at 30°C were isolated and rechecked for cold resistance through EOP and growth curves. The suppressors were subjected to whole-genome sequencing to identify potential suppressor mutations. Genomic DNA was isolated from the parent and suppressors using a DNeasy Blood and Tissue Kit (Qiagen) as per the manufacturer's instructions. Library preparation and Illumina sequencing were performed by the Microbial Genome Sequencing Center (MiGS, Pittsburgh, PA, USA). Briefly, Illumina libraries were prepared and the libraries were sequenced with paired-end 151 bp reads on a NextSeq 2000 platform to a read depth of at least 200 Mbp. To identify variants from the reference genome (GenBank: U00096.3), reads were trimmed to remove adaptors and the *breseq* software (version 0.35.4) was used to call variants (104). Identified mutations were confirmed with Sanger sequencing. The consensus motif of the FabR binding site of *fabA* was determined using Multiple Em for Motif Elicitation (MEME) tool in the MEME Suite based on the sequence of putative FabR binding sites reported by Feng and Cronan (2011) (64, 105, 106). The ability of the identified P_{fabA} mutation to suppress was confirmed by linking the mutant to *zcb*-3059::Tn*10* (107) and transducing to a clean background.

Lux reporter assays. Overnight cultures of strains harboring the pJW15-P_{fabA} and pJW15-P_{fabA}(-19G>A) plasmids were sub-cultured at (1:100) into 200 µl of fresh LB broth in a black 96-well plate. The plate was incubated in a BioTek Synergy H1 plate reader and the OD_{600} and luminescence intensity were recorded every 2.5 min for 6 hours as described in previously (103, 108). Each biological replicate was performed in technical triplicate.

RNA-seq. Overnight cultures grown at 42°C were subcultured 1:100 into fresh media and incubated at 42 °C until an OD_{600} of 0.4. Then, the cultures were shifted to a 30 °C water bath for 30 minutes. 500 µL of each culture was immediately fixed in the RNAprotect Bacteria Reagent (Qiagen) as per the manufacturer's instructions. RNA was purified using the RNeasy Kit (Qiagen) with on-column DNase digestion following the manufacturer's protocol for gram-negative bacteria. Library preparation, Illumina sequencing, and analysis of differentially expressed genes were performed by MiGS. Briefly, samples were treated with RNase free DNase (Invitrogen) and library preparation was performed with the Stranded Total RNA Prep Ligation with Ribo-Zero Plus Kit (Illumina). Libraries were sequenced with 2x50 bp reads on a NextSeq2000 (Illumina). Demultiplexing, quality control, and adapter trimming carried out using bcl-convert (Illumina, v3.9.3) and bcl2fastq (Illumina). Read mapping and read quantification were performed with HISAT2 (109) and the featureCounts tool in Subreader (110), respectively. Descriptive statistics of the RNA-seq read data can be found in **Table S1**. The raw RNA-seq data is available in the Sequence Read Archive database (SRA) (https://www.ncbi.nlm.nih.gov/sra, BioProject ID PRJNA965821; Reviewer Access).

Raw read counts were normalized using the Trimmed Mean of M values algorithm in edgeR (111) and converted to counts per million. The Quasi-Linear F-Test function of

edgeR was used for differential expression analysis. Normalized read values and for all
genes as well as differential expression analysis statistics can be found in **Dataset S3**.
KEGG pathway analysis was conducted using the *"kegga"* function of limma (112).
Expression of genes differentially expressed (greater than 2-fold change and $p<0.05$) was
subjected to principle component analysis, and pathway and enrichment analysis using
the EcoCyc Omics Dashboard (112).

Lipid extraction and TLC. For thin layer chromatography (TLC) experiments, cultures were
grown in LB with 1 mCi/mL ^{32}P until OD_{600} of 0.6-0.8 at the indicated temperature. For
temperature downshift experiments, cells were grown at 42 °C until OD_{600} of 0.2 and then
transferred to 30 °C for 2 hours. For lipid extraction, cells were pelleted and resuspended
in 0.2 mL of 0.5 M NaCl in 0.1 N HCl. Lipids were extracted by first adding 0.6 mL of
chloroform:methanol (1:2) to create a single-phase solution. After vigorous vortexing for
30 minutes, 0.2 mL of 0.5 M NaCl in 0.1 N HCl was added to convert the single-phase
solution to a two-phase solution. After centrifugation at 20 000 x g for 5 minutes at room
temperature, the lower phase was recovered and used for TLC. Approximately 2,000 cpm
of phospholipid extract was subjected to TLC analysis using a HPTLC 60 plate (EMD,
Gibbstown, NJ)) developed with either solvent 1: chloroform/methanol/acetic acid
[60/25/10] (vol/vol/vol) in **Fig. S4A** or by solvent 2:
chloroform/methanol/ammonia/water [65/37.5/3/1] (vol/vol/vol/vol) in **Fig. S4B**.
Radiolabeled lipids were visualized and quantified using a Phosphoimager Typhoon FLA
9500 (GE). Images were processed and quantified using ImageQuant^{TM} software for
scanning and analysis. The phospholipid content is expressed as molecular percentage of
the total phospholipids (correcting for the two phosphates per molecule of CL).

Phospholipid composition analysis. Whole cell lipid extracts were isolated as for TLC
temperature downshift experiments without radiolabeling. Absolute amounts of CL, PG,
and PE were determined in these lipid extracts using liquid chromatography coupled to
electrospray ionization mass spectrometry (LC/MS) in an API 4000 mass spectrometer
(Sciex, Framingham, MA, USA) using normal phase solvents as previously published (113).
The cardiolipin internal standard $(14:0)_4$CL (Avanti Polar Lipids, Alabaster, AL USA) was
added to all samples to quantify CL, PG, and PE within a single run. Standard curves
containing reference standards of all three phospholipids, $(18:1)_4$CL (Avanti Polar Lipids),
$16:0_1 18:1_1$PG and $16:0_1 18:1_1$PE (Cayman Chemicals, Ann Arbor, MI, USA) were used to
quantify the nmol/mg protein for each species (**Dataset 1**) as previously described (114).
Percentage values were calculated by dividing the absolute value of that species by the
sum of individual molecular species of CL, PG, or PE. The identities of CL species were
confirmed using tandem mass spectrometry.
588
589
590   **Acknowledgments**
591

We thank members of the Mitchell Lab for productive discussions. We would like to thank
Professor Tracy Raivio (University of Alberta) for the kind gift of pJW15 plasmid. This work
was funded by the National Institute of Allergy and Infectious Disease under Award
Number R01-AI155915 and R01-AI155915-S1 (to A.M.M.), the National Institute of
General Medical Sciences under Award Number R01-GM121493 (to M.B.), and Texas
A&M University start-up funds (to A.M.M.).

598
599

References

1.   Nikaido H. Molecular basis of bacterial outer membrane permeability revisited.
Microbiology and Molecular Biology Reviews. 2003;67(4):593-656.
2.   Silhavy TJ, Kahne D, Walker S. The bacterial cell envelope. Cold Spring Harbor
Perspectives in Biology. 2010;2(5).
3.   Henderson JC, Zimmerman SM, Crofts AA, Boll JM, Kuhns LG, Herrera CM, Trent
MS. The power of asymmetry: Architecture and assembly of the Gram-negative outer
membrane lipid bilayer. Annual Review of Microbiology. 2016;70(1):255-78.
4.   Berry J, Rajaure M, Pang T, Young R. The spanin complex is essential for lambda
lysis. J Bacteriol. 2012;194(20):5667-74.
5.   Rojas ER, Billings G, Odermatt PD, Auer GK, Zhu L, Miguel A, et al. The outer
membrane is an essential load-bearing element in Gram-negative bacteria. Nature.
2018;559(7715):617-21.
6.   Navarro Llorens JM, Tormo A, Martinez-Garcia E. Stationary phase in gram-
negative bacteria. FEMS Microbiol Rev. 2010;34(4):476-95.
7.   Lugtenberg EJJ, Peters R. Distribution of lipids in cytoplasmic and outer
membranes of *Escherichia coli* K12. Biochimica et Biophysica Acta (BBA) - Lipids and
Lipid Metabolism. 1976;441(1):38-47.
8.   Kamio Y, Nikaido H. Outer membrane of *Salmonella typhimurium*: accessibility of
phospholipid head groups to phospholipase c and cyanogen bromide activated dextran
in the external medium. Biochemistry. 1976;15(12):2561-70.
9.   Bogdanov M, Pyrshev K, Yesylevskyy S, Ryabichko S, Boiko V, Ivanchenko P, et al.
Phospholipid distribution in the cytoplasmic membrane of Gram-negative bacteria is
highly asymmetric, dynamic, and cell shape-dependent. Science Advances.
2020;6(23):eaaz6333.
10.   Benn G, Mikheyeva IV, Inns PG, Forster JC, Ojkic N, Bortolini C, et al. Phase
separation in the outer membrane of *Escherichia coli*. Proceedings of the National
Academy of Sciences. 2021;118(44):e2112237118.
11.   Konovalova A, Silhavy TJ. Outer membrane lipoprotein biogenesis: Lol is not the
end. Philosophical Transactions of the Royal Society B: Biological Sciences.
2015;370(1679):20150030.
12.   Konovalova A, Kahne DE, Silhavy TJ. Outer Membrane Biogenesis. Annu Rev
Microbiol. 2017;71:539-56.

13.   Lundstedt E, Kahne D, Ruiz N. Assembly and maintenance of lipids at the
bacterial outer membrane. Chemical Reviews. 2021;121(9):5098-123.
14.   John E. Cronan J, Rock CO. Biosynthesis of membrane lipids. EcoSal Plus.
2008;3(1).
15.   Jones NC, Osborn MJ. Translocation of phospholipids between the outer and
inner membranes of *Salmonella typhimurium*. Journal of Biological Chemistry.
1977;252(20):7405-12.
16.   Donohue-Rolfe AM, Schaechter M. Translocation of phospholipids from the inner
to the outer membrane of *Escherichia coli*. Proceedings of the National Academy of
Sciences. 1980;77(4):1867-71.
17.   Raetz CR, Kantor GD, Nishijima M, Newman KF. Cardiolipin accumulation in the
inner and outer membranes of *Escherichia coli* mutants defective in phosphatidylserine
synthetase. J Bacteriol. 1979;139(2):544-51.
18.   Langley KE, Hawrot E, Kennedy EP. Membrane assembly: movement of
phosphatidylserine between the cytoplasmic and outer membranes of *Escherichia coli*. J
Bacteriol. 1982;152(3):1033-41.
19.   White DA, Lennarz WJ, Schnaitman CA. Distribution of lipids in the wall and
cytoplasmic membrane subfractions of the cell envelope of *Escherichia coli*. Journal of
Bacteriology. 1972;109(2):686-90.
20.   Ishinaga M, Kanamoto R, Kito M. Distribution of phospholipid molecular species
in outer and cytoplasmic membranes of *Escherichia coli*. The Journal of Biochemistry.
1979;86(1):161-5.
21.   Davydova L, Bakholdina S, Barkina M, Velansky P, Bogdanov M, Sanina N. Effects
of elevated growth temperature and heat shock on the lipid composition of the inner
and outer membranes of *Yersinia pseudotuberculosis*. Biochimie. 2016;123:103-9.
22.   Osborn MJ, Gander JE, Parisi E, Carson J. Mechanism of Assembly of the Outer
Membrane of *Salmonella* typhimurium : Isolation and characterization of cytoplasmic
and outer membrane. Journal of Biological Chemistry. 1972;247(12):3962-72.
23.   Raetz CR, Newman KF. Diglyceride kinase mutants of *Escherichia coli*: inner
membrane association of 1,2-diglyceride and its relation to synthesis of membrane-
derived oligosaccharides. J Bacteriol. 1979;137(2):860-8.
24.   Bogdanov M. Renovating a double fence with or without notifying the next door
and across the street neighbors: why the biogenic cytoplasmic membrane of Gram-
negative bacteria display asymmetry? . Emerging Topics in Life Sciences. 2023;7(1):137-
50.
25.   Grimm J, Shi H, Wang W, Mitchell AM, Wingreen NS, Huang KC, Silhavy TJ. The
inner membrane protein YhdP modulates the rate of anterograde phospholipid flow in
Escherichia coli. Proc Natl Acad Sci U S A 2020;117(43):26907-14.
26.   Malinverni JC, Silhavy TJ. An ABC transport system that maintains lipid
asymmetry in the Gram-negative outer membrane. Proceedings of the National
Academy of Sciences. 2009;106(19):8009-14.

27.   Sutterlin HA, Shi H, May KL, Miguel A, Khare S, Huang KC, Silhavy TJ. Disruption
of lipid homeostasis in the Gram-negative cell envelope activates a novel cell death
pathway. Proceedings of the National Academy of Sciences. 2016;113(11):E1565-E74.
28.   May KL, Silhavy TJ. The *Escherichia coli* phospholipase PldA regulates outer
membrane homeostasis via lipid signaling. mBio. 2018;9(2):e00379-18.
29.   Tang X, Chang S, Qiao W, Luo Q, Chen Y, Jia Z, et al. Structural insights into outer
membrane asymmetry maintenance in Gram-negative bacteria by MlaFEDB. Nat Struct
Mol Biol. 2021;28(1):81-91.
30.   Ruiz N, Davis RM, Kumar S. YhdP, TamB, and YdbH are redundant but essential
for growth and lipid homeostasis of the gram-negative outer membrane. mBio.
2021;12(6):e02714-21.
31.   Douglass MV, McLean AB, Trent MS. Absence of YhdP, TamB, and YdbH leads to
defects in glycerophospholipid transport and cell morphology in Gram-negative
bacteria. PLOS Genetics. 2022;18(2):e1010096.
32.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly
accurate protein structure prediction with AlphaFold. Nature. 2021;596(7873):583-9.
33.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al.
AlphaFold Protein Structure Database: massively expanding the structural coverage of
protein-sequence space with high-accuracy models. Nucleic Acids Research.
2021;50(D1):D439-D44.
34.   Josts I, Stubenrauch CJ, Vadlamani G, Mosbahi K, Walker D, Lithgow T, Grinter R.
The structure of a conserved domain of TamB reveals a hydrophobic β taco fold.
Structure. 2017;25(12):1898-906.e5.
35.   Kumar N, Leonzino M, Hancock-Cerutti W, Horenkamp FA, Li P, Lees JA, et al.
VPS13A and VPS13C are lipid transport proteins differentially localized at ER contact
sites. Journal of Cell Biology. 2018;217(10):3625-39.
36.   Valverde DP, Yu S, Boggavarapu V, Kumar N, Lees JA, Walz T, et al. ATG2
transports lipids to promote autophagosome biogenesis. J Cell Biol. 2019;218(6):1787-
98.
37.   Otomo T, Maeda S. ATG2A transfers lipids between membranes *in vitro*.
Autophagy. 2019;15(11):2031-2.
38.   Osawa T, Kotani T, Kawaoka T, Hirata E, Suzuki K, Nakatogawa H, et al. Atg2
mediates direct lipid transfer between membranes for autophagosome formation. Nat
Struct Mol Biol. 2019;26(4):281-8.
39.   Cooper BF, Clark R, Kudhail A, Bhabha G, Ekiert DC, Khalid S, Isom GL.
Phospholipid transport to the bacterial outer membrane through an envelope-spanning
bridge. bioRxiv. 2023:2023.10.05.561070.
40.   Grasekamp KP, Beaud Benyahia B, Taib N, Audrain B, Bardiaux B, Rossez Y, et al.
The Mla system of diderm Firmicute *Veillonella parvula* reveals an ancestral
transenvelope bridge for phospholipid trafficking. Nature Communications.
2023;14(1):7642.

41.   Sposato D, Mercolino J, Torrini L, Sperandeo P, Lucidi M, Alegiani R, et al.
Redundant essentiality of AsmA-like proteins in *Pseudomonas aeruginosa*. mSphere.
2024;9(2):e00677-23.
42.   Mitchell AM, Wang W, Silhavy TJ. Novel Rpos-dependent mechanisms
strengthen the envelope permeability barrier during stationary phase. J Bacteriol.
2017;199(2):e00708-16.
43.   Mitchell AM, Srikumar T, Silhavy TJ. Cyclic enterobacterial common antigen
maintains the outer membrane permeability barrier of *Escherichia coli* in a manner
controlled by Yhdp. mBio. 2018;9(4):e01321-18.
44.   Selkrig J, Mosbahi K, Webb CT, Belousoff MJ, Perry AJ, Wells TJ, et al. Discovery
of an archetypal protein transport system in bacterial outer membranes. Nature
Structural & Molecular Biology. 2012;19(5):506-10.
45.   Stubenrauch CJ, Dougan G, Lithgow T, Heinz E. Constraints on lateral gene
transfer in promoting fimbrial usher protein diversity and function. Open Biology.
2017;7(11):170144.
46.   Stubenrauch C, Belousoff MJ, Hay ID, Shen H-H, Lillington J, Tuck KL, et al.
Effective assembly of fimbriae in *Escherichia coli* depends on the translocation assembly
module nanomachine. Nature Microbiology. 2016;1(7):16064.
47.   Cronan JE. The *Escherichia coli* FadR transcription factor: Too much of a good
thing? Mol Microbiol. 2021;115(6):1080-5.
48.   Nunn WD, Giffin K, Clark D, Cronan JE. Role for *fadR* in unsaturated fatty acid
biosynthesis in *Escherichia coli*. Journal of Bacteriology. 1983;154(2):554-60.
49.   Rai AK, Mitchell AM. Enterobacterial common antigen: Synthesis and function of
an enigmatic molecule. mBio. 2020;11(4):e01914-20.
50.   Paradis-Bleau C, Kritikos G, Orlova K, Typas A, Bernhardt TG. A genome-wide
screen for bacterial envelope biogenesis mutants identifies a novel factor involved in
cell wall precursor metabolism. PLoS Genet. 2014;10(1):e1004056.
51.   Ruiz N, Falcone B, Kahne D, Silhavy TJ. Chemical conditionality: a genetic strategy
to probe organelle assembly. Cell. 2005;121(2):307-17.
52.   Nikaido H, Vaara M. Molecular basis of bacterial outer membrane permeability.
Microbiol Rev. 1985;49(1):1-32.
53.   Bishop RE. Structural biology of membrane-intrinsic β-barrel enzymes: Sentinels
of the bacterial outer membrane. Biochimica et Biophysica Acta (BBA) - Biomembranes.
2008;1778(9):1881-96.
54.   Rowlett VW, Mallampalli V, Karlstaedt A, Dowhan W, Taegtmeyer H, Margolin W,
Vitrac H. Impact of membrane phospholipid slterations in *Escherichia coli* on cellular
function and bacterial stress adaptation. J Bacteriol. 2017;199(13).
55.   Tan BK, Bogdanov M, Zhao J, Dowhan W, Raetz CRH, Guan Z. Discovery of a
cardiolipin synthase utilizing phosphatidylethanolamine and phosphatidylglycerol as
substrates. Proceedings of the National Academy of Sciences. 2012;109(41):16504-9.

56.   Pluschke G, Hirota Y, Overath P. Function of phospholipids in *Escherichia coli*.
Characterization of a mutant deficient in cardiolipin synthesis. Journal of Biological
Chemistry. 1978;253(14):5048-55.
57.   Nishijima S, Asami Y, Uetake N, Yamagoe S, Ohta A, Shibuya I. Disruption of the
Escherichia coli cls gene responsible for cardiolipin synthesis. Journal of Bacteriology.
1988;170(2):775-80.
58.   Guo D, Tropp BE. A second *Escherichia coli* protein with CL synthase activity.
Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids.
2000;1483(2):263-74.
59.   Kikuchi S, Shibuya I, Matsumoto K. Viability of an *Escherichia coli pgsA* null
mutant lacking detectable phosphatidylglycerol and cardiolipin. Journal of Bacteriology.
2000;182(2):371-6.
60.   Suzuki M, Hara H, Matsumoto K. Envelope disorder of *Escherichia coli* cells
lacking phosphatidylglycerol. Journal of Bacteriology. 2002;184(19):5418-25.
61.   Shiba Y, Yokoyama Y, Aono Y, Kiuchi T, Kusaka J, Matsumoto K, Hara H.
Activation of the Rcs signal transduction system is responsible for the thermosensitive
growth defect of an *Escherichia coli* mutant lacking phosphatidylglycerol and cardiolipin.
J Bacteriol. 2004;186(19):6526-35.
62.   Grabowicz M, Silhavy TJ. Redefining the essential trafficking pathway for outer
membrane lipoproteins. Proceedings of the National Academy of Sciences.
2017;114(18):4769-74.
63.   Cronan JE, Jr., Gelmann EP. An estimate of the minimum amount of unsaturated
fatty acid required for growth of *Escherichia coli*. J Biol Chem. 1973;248(4):1188-95.
64.   Feng Y, Cronan JE. Complex binding of the FabR repressor of bacterial
unsaturated fatty acid biosynthesis to its cognate promoters. Mol Microbiol.
2011;80(1):195-218.
65.   Feng Y, Cronan JE. *Escherichia coli* unsaturated fatty acid synthesis: Complex
transcription of the *fabA* gene and *in vivo* identification of the essential reaction
catalyzed by FabB. Journal of Biological Chemistry. 2009;284(43):29526-35.
66.   Henry MF, Cronan JE. *Escherichia coli* transcription factor that both activates
fatty acid synthesis and represses fatty acid degradation. Journal of Molecular Biology.
1991;222(4):843-9.
67.   Henry MF, Cronan JE. A new mechanism of transcriptional regulation: Release of
an activator triggered by small molecule binding. Cell. 1992;70(4):671-9.
68.   Rock CO, Jackowski S. Pathways for the incorporation of exogenous fatty acids
into phosphatidylethanolamine in *Escherichia coli*. J Biol Chem. 1985;260(23):12720-4.
69.   Heath RJ, Rock CO. Roles of the FabA and FabZ β-hydroxyacyl-acyl carrier protein
dehydratases in *Escherichia coli* fatty acid biosynthesis. Journal of Biological Chemistry.
1996;271(44):27795-801.
70.   Ostrander DB, Sparagna GC, Amoscato AA, McMillin JB, Dowhan W. Decreased
Cardiolipin Synthesis Corresponds with Cytochromec Release in Palmitate-induced
Cardiomyocyte Apoptosis. Journal of Biological Chemistry. 2001;276(41):38061-7.

71.   Lewis RNAH, Zweytick D, Pabst G, Lohner K, McElhaney RN. Calorimetric, X-Ray
Diffraction, and Spectroscopic Studies of the Thermotropic Phase Behavior and
Organization of Tetramyristoyl Cardiolipin Membranes. Biophysical Journal.
2007;92(9):3166-77.

72.   De Siervo AJ. Alterations in the phospholipid composition of *Escherichia coli* B
during growth at different temperatures. J Bacteriol. 1969;100(3):1342-9.

73.   Bogdanov M, Sun J, Kaback HR, Dowhan W. A Phospholipid Acts as a Chaperone
in Assembly of a Membrane Transport Protein. Journal of Biological Chemistry.
1996;271(20):11615-8.

74.   Mitchell AM, Silhavy TJ. Envelope stress responses: balancing damage repair and
toxicity. Nature Reviews Microbiology. 2019;17(7):417-28.

75.   Luevano-Martinez LA, Kowaltowski AJ. Phosphatidylglycerol-derived
phospholipids have a universal, domain-crossing role in stress responses. Arch Biochem
Biophys. 2015;585:90-7.

76.   Fernandez M, Paulucci NS, Peppino Margutti M, Biasutti AM, Racagni GE,
Villasuso AL, et al. Membrane Rigidity and Phosphatidic Acid (PtdOH) Signal: Two
Important Events in Acinetobacter guillouiae SFC 500-1A Exposed to Chromium(VI) and
Phenol. Lipids. 2019;54(9):557-70.

77.   Shibata A, Ikawa K, Shimooka T, Terada H. Significant stabilization of the
phosphatidylcholine bilayer structure by incorporation of small amounts of cardiolipin.
Biochimica et Biophysica Acta (BBA) - Biomembranes. 1994;1192(1):71-8.

78.   Boscia AL, Treece BW, Mohammadyani D, Klein-Seetharaman J, Braun AR,
Wassenaar TA, et al. X-ray structure, thermodynamics, elastic properties and MD
simulations of cardiolipin/dimyristoylphosphatidylcholine mixed membranes. Chemistry
and Physics of Lipids. 2014;178:1-10.

79.   Pennington ER, Fix A, Sullivan EM, Brown DA, Kennedy A, Shaikh SR. Distinct
membrane properties are differentially influenced by cardiolipin content and acyl chain
composition in biomimetic membranes. Biochimica et Biophysica Acta (BBA) -
Biomembranes. 2017;1859(2):257-67.

80.   Lopes SC, Ivanova G, de Castro B, Gameiro P. Revealing cardiolipins influence in
the construction of a significant mitochondrial membrane model. Biochimica et
Biophysica Acta (BBA) - Biomembranes. 2018;1860(11):2465-77.

81.   Marr AG, Ingraham JL. Effect of temperature on the composition of fatty acids in
Escherichia coli. Journal of Bacteriology. 1962;84(6):1260-7.

82.   Garwin JL, Klages AL, Cronan JE. Beta-ketoacyl-acyl carrier protein synthase II of
Escherichia coli. Evidence for function in the thermal regulation of fatty acid synthesis.
Journal of Biological Chemistry. 1980;255(8):3263-5.

83.   de Mendoza D, Klages Ulrich A, Cronan JE. Thermal regulation of membrane
fluidity in *Escherichia coli*. Effects of overproduction of beta-ketoacyl-acyl carrier protein
synthase I. Journal of Biological Chemistry. 1983;258(4):2098-101.

84.   Knapp BD, Huang KC. The effects of temperature on cellular physiology. Annual
Review of Biophysics. 2022;51(1):499-526.

85.   Bogdanov M, Umeda M, Dowhan W. Phospholipid-assisted refolding of an
integral membrane protein: Minimum structural features for phosphatidylethanolamine
to act as a molecular chaperone. Journal of Biological Chemistry. 1999;274(18):12339-
45.
86.   Raetz CR, Whitfield C. Lipopolysaccharide endotoxins. Annu Rev Biochem.
2002;71:635-700.
87.   Conway T, Creecy JP, Maddox SM, Grissom JE, Conkle TL, Shadid TM, et al.
Unprecedented high-resolution view of bacterial operon architecture revealed by RNA
sequencing. mBio. 2014;5(4):e01442-14.
88.   Thomason MK, Bischler T, Eisenbart SK, Förstner KU, Zhang A, Herbig A, et al.
Global transcriptional start site mapping using differential RNA sequencing reveals novel
antisense rnas in *Escherichia coli*. Journal of Bacteriology. 2015;197(1):18-28.
89.   Adams PP, Baniulyte G, Esnault C, Chegireddy K, Singh N, Monge M, et al.
Regulatory roles of *Escherichia coli* 5' UTR and ORF-internal RNAs detected by 3' end
mapping. eLife. 2021;10:e62438.
90.   Karp PD, Ong WK, Paley S, Billington R, Caspi R, Fulcher C, et al. The EcoCyc
database. EcoSal Plus. 2018;8(1):10.1128/ecosalplus.ESP-0006-2018.
91.   Huerta AM, Collado-Vides J. Sigma70 promoters in *Escherichia coli*: specific
transcription in dense regions of overlapping promoter-like signals. Journal of Molecular
Biology. 2003;333(2):261-78.
92.   Dowhan W, Bogdanov M. Lipid-dependent membrane protein topogenesis. Annu
Rev Biochem. 2009;78:515-40.
93.   Bogdanov M, Xie J, Heacock P, Dowhan W. To flip or not to flip: lipid-protein
charge interactions are a determinant of final membrane protein topology. J Cell Biol.
2008;182(5):925-35.
94.   Bogdanov M, Dowhan W. Lipid-dependent generation of dual topology for a
membrane protein. J Biol Chem. 2012;287(45):37939-48.
95.   Bogdanov M, Dowhan W, Vitrac H. Lipids and topological rules governing
membrane protein assembly. Biochim Biophys Acta. 2014;1843(8):1475-88.
96.   Ramezanifard R, Golubeva YA, Palmer AD, Slauch JM. TamAB is regulated by
PhoPQ and functions in outer membrane homeostasis during *Salmonella* pathogenesis. J
Bacteriol. 2023;205(10):e0018323.
97.   Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, et al. The STRING
database in 2021: customizable protein-protein networks, and functional
characterization of user-uploaded gene/measurement sets. Nucleic Acids Res.
2021;49(D1):D605-d12.
98.   Grasekamp KP, Beaud B, Taib N, Audrain B, Bardiaux B, Rossez Y, et al. Bridges
instead of boats? The Mla system of diderm Firmicute *Veillonella parvula* reveals an
ancestral transenvelope core of phospholipid trafficking. bioRxiv.
2023:2023.06.30.547184.

99.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of
Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol
Syst Biol. 2006;2:2006.0008-2006.0008.
100.   Silhavy TJ, Berman ML, Enquist LW. Experiments with gene fusions: Cold Spring
Harbor Laboratory; 1984.
101.   Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. Proceedings of the National Academy of
Sciences. 2000;97(12):6640-5.
102.   MacRitchie DM, Ward JD, Nevesinjac AZ, Raivio TL. Activation of the Cpx
envelope stress response down-regulates expression of several locus of enterocyte
effacement-encoded genes in enteropathogenic *Escherichia coli*. Infect Immun.
2008;76(4):1465-75.
103.   Wong JL, Vogt SL, Raivio TL. Using reporter genes and the *Escherichia coli* ASKA
overexpression library in screens for regulators of the Gram negative envelope stress
response. Methods Mol Biol. 2013;966:337-57.
104.   Deatherage DE, Barrick JE. Identification of mutations in laboratory-evolved
microbes from next-generation sequencing data using breseq. Methods Mol Biol.
2014;1151:165-88.
105.   Bailey TL, Elkan C. Fitting a mixture model by expectation maximization to
discover motifs in biopolymers. Proc Int Conf Intell Syst Mol Biol. 1994;2:28-36.
106.   Bailey TL, Johnson J, Grant CE, Noble WS. The MEME Suite. Nucleic Acids Res.
2015;43(W1):W39-49.
107.   Singer M, Baker TA, Schnitzler G, Deischel SM, Goel M, Dove W, et al. A
collection of strains containing genetically linked alternating antibiotic resistance
elements for genetic mapping of Escherichia coli. Microbiol Rev. 1989;53(1):1-24.
108.   Rai AK, Carr JF, Bautista DE, Wang W, Mitchell AM. ElyC and cyclic
enterobacterial common antigen regulate synthesis of phosphoglyceride-linked
enterobacterial common antigen. mBio. 2021;12(6):e02846-21.
109.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. Nature Biotechnology.
2019;37(8):907-15.
110.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2013;30(7):923-30.
111.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics.
2009;26(1):139-40.
112.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. limma powers
differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Research. 2015;43(7):e47-e.
113.   Sparagna GC, Johnson CA, McCune SA, Moore RL, Murphy RC. Quantitation of
cardiolipin molecular species in spontaneously hypertensive heart failure rats using
electrospray ionization mass spectrometry. J Lipid Res. 2005;46(6):1196-204.

114.   Hryc CF, Mallampalli VKPS, Bovshik EI, Azinas S, Fan G, Serysheva II, et al.
Structural insights into cardiolipin replacement by phosphatidylglycerol in a cardiolipin-
lacking yeast respiratory supercomplex. Nature Communications. 2023;14(1):2783.

924
925

Figures and Tables

Figure 1. Deletion of *yhdP* and *fadR* causes synthetic cold sensitivity for which TamB is necessary. **(A)** The Δ*yhdP* Δ*fadR* strain had impaired growth at lower temperatures. **(B)** EOPs were performed to confirm the growth defect of the Δ*yhdP* Δ*fadR* strain. **(C-F)** EOPs were performed at 30 °C to assess cold sensitivity. **(C)** The growth of the Δ*yhdP* Δ*fadR* strain was not affected by the loss of ECA (Δ*wecA*). **(D)** Neither a Δ*tamB* Δ*fadR* strain nor a Δ*ydbH* Δ*fadR* strain exhibited cold sensitivity. **(E)** Deletion of *ydbH* did not suppress cold sensitivity in the Δ*yhdP* Δ*fadR* strain. **(F)** Deletion of *tamB* completely suppressed cold sensitivity in the Δ*yhdP* Δ*fadR* strain and the Rcs stress response was not necessary for this suppression. **(G)** Growth curves were performed at 30 °C . The Δ*yhdP* Δ*fadR* Δ*tamB* strain grew well, indicating suppression was unlikely to result from slow growth.

940 Quantitative data are shown as the mean of three biological replicates ± the SEM. Images
941 are representative of three independent experiments.

Figure 2. Lowering cardiolipin levels can suppress cold sensitivity of the ΔyhdP ΔfadR
945
strain. (A-B) Thin layer chromatography (TLC) of lipid extracts was performed for the

indicated cultures grown to log phase at 42 °C then transferred to 30 °C for 2 hours before
analysis. **(A)** Relative phospholipid levels were calculated. The Δ*yhdP* Δ*fadR* strain shows
increased levels of CL and a concomitant decrease in PG levels, increasing the ratio of CL
to PG (B). Data are averages of two to three biological replicates ± the SEM. For ratios
individual data points are indicated with open circles. * p<0.05 vs. MG1655 by Student's
T-test; ‡ p<0.05 vs. the Δ*yhdP* Δ*fadR* strain by Student's T-test. **(C-G)** Growth effects of
altered phospholipid levels were investigated by EOP. Images are representative of three
independent experiments. **(C)** PgsA depletion strains in a background where *pgsA* is non-
essential were used to assay the effect of lowering PG and CL levels on cold sensitivity.
Arabinose induces expression of *pgsA* while glucose represses. Cultures were induced or
repressed for 4-5 generations before as well as during the EOP. Lowering PG and CL levels
partially suppressed cold sensitivity. **(D)** Deletion of the three CL synthases, *clsA*, *clsB*, and
clsC, suppressed cold sensitivity of the Δ*yhdP* Δ*fadR* strain. **(E)** Deletion of *clsA* also
suppressed cold sensitivity. **(F)** Deletion of *clsB* does not suppress cold sensitivity, while
deletion of *clsC* (F) does.
962
963

964
Figure 3. Cold sensitivity can be suppressed by increasing *fabA* expression. (A) The cold
sensitivity of a spontaneous suppressor of Δ*yhdP* Δ*fadR* cold sensitivity was tested by EOP.
The suppressor returned growth to the level of the Δ*fadR* mutant. **(B)** A diagram of the
promoter region of *fabA* is shown with its transcriptional start sites (blue arrows) and
regulator binding sites (boxes). FadR acts as an activator while FabR acts as a repressor.
The motif of the FabR binding site, generated using MEME, is shown with the sequences
of the parent strain and of Suppressor 1. **(C)** The P$_{fabA}$(-19G>A) mutation was linked to a
Tn*10* and transferred to the Δ*yhdP* Δ*fadR* strain by transduction. The Δ*yhdP* Δ*fadR* strain
with P$_{fabA}$(-19G>A) was not cold sensitivity, confirming suppression. **(D)** *fabA* RNA levels
are shown. Bars are the mean of three biological replicates ± the SEM. Individual data
points are shown as open circles. ** $p < 0.005$, *** $p < 0.0005$ by quasi-linear F-test. Data
are from the RNA-seq experiment in Figure 5 and *fabA* data are also shown in Figure 5F.
(E) Luciferase reporters of wild-type and mutant *fabA* promoter activity were
constructed. The mutation caused similar fold increases in P$_{fabA}$ activity in all strain
backgrounds, indicating the activity of the second *fabA* promoter and not FabR binding
were likely affected by the suppressor mutation. Data are luminescence relative to OD$_{600}$
and are shown as the average of two biological replicates ± the SEM.
982

Figure 4. Decreasing fatty acid saturation suppresses Δ*yhdP* Δ*fadR* cold sensitivity. (A-C) Lipid were extracted from cells grown to OD$_{600}$ 0.2 at 42 °C then shifted to 30 °C for 2 hours before harvest. LC/MS was performed on lipid extracts using absolute quantification and the percentage of each specific species of PE (A), PG (B), and CL (C) was calculated using the sum of all molecular species detected. Suppressor 1 trends towards decreased saturation compared to its parent strain (Δ*yhdP* Δ*fadR*). Data for phospholipids without unsaturations are shown in grey, with one unsaturation in blue, with two unsaturations in green, with three unsaturations in orange. Data are the average of three biological replicates ± the SEM. * $p < 0.05$ vs. MG1655 by the Mann-Whitney test; ‡ $p < 0.05$ vs. the Δ*yhdP* Δ*fadR* strain by the Mann-Whitney test. All molecular species are shown in Fig. S6. **(D)** EOPS were performed on media supplemented with oleic acid or a vehicle control. Oleic acid suppresses the cold sensitivity of the Δ*yhdP* Δ*fadR* strain. **(E)** Growth curves of the indicated strains were performed in the presence of oleic acid, stearic acid,

998   or a vehicle control. Oleic acid suppressed the cold sensitivity of the Δ*yhdP* Δ*fadR* strain

999   while stearic acid did not. Data are the average of three biological replicates ± the SEM.

1000

Figure 5. Transcriptional landscape of the Δ*yhdP* Δ*fadR* and suppressed strain. Three
biological replicates of the indicated strains were grown at 42 °C to mid-log then
transferred to 30 °C for 30 minutes before harvesting RNA and performing RNA-seq.
Differential expression was calculated based on a greater than 2-fold change in expression
and a p value of less than 0.05 by quasi-linear F-test. **(A)** PCA was performed on the
expression of all genes differentially expressed between any groups of samples.
Suppressor 1 and the Δ*yhdP* Δ*fadR* strain grouped closer together than to the wild-type
strain, although Suppressor 1 was closer to wild type than the Δ*yhdP* Δ*fadR* strain. **(B-D)**
Volcano plots show the average fold change and significance of expression changes
between the Δ*yhdP* Δ*fadR* strain and wild-type MG1655 (B), Suppressor 1 and wild-type
MG1655 (C), and Suppressor 1 and the Δ*yhdP* Δ*fadR* strain (D). Genes with changes
greater than 2-fold are shown in blue, while genes with changes less than 2-fold are
shown in grey. Genes with a q-value (false discovery rate) of less than 0.05 are shown in
yellow. Blue lines indicate a 2-fold change and dotted line indicates p>0.05. The number
of up or down regulated genes is indicated in blue. The names of the 10 genes with the
largest changes are called out. More genes were up-regulated then down regulated in the
Δ*yhdP* Δ*fadR* strain and Suppressor 1 vs. wild type. More genes were down regulated in

Suppressor 1 compared to the Δ*yhdP* Δ*fadR* strain. **(E)** Relative expression of genes in the
CL synthesis pathway is shown as averages ± the SEM and individual data points. No
significant changes were evident. **(F)** Relative expression of genes in the fatty acid
synthesis and degradation pathways is shown as averages ± the SEM and individual data
points. A general normalizing of these pathways occurred in Suppressor 1 compared to
the Δ*yhdP* Δ*fadR* strain; however, the most significant change was in the expression of
fabA. * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$ by quasi-linear F-test.
1026

1027

Figure 6. Hypothesized model for the role of YhdP and TamB in phospholipid transport.
Our data can be explained by the following model of phospholipid transport. **(A)** In normal
conditions, YhdP is mainly responsible for the transport of phospholipids with more
saturated fatty acid chains between the IM and OM. TamB shows a strong preference for
transporting phospholipids with more unsaturated fatty acids, with YdbH playing a minor
role in phospholipid transport. **(B)** In the Δ*yhdP* Δ*fadR* strain, YhdP is not available to
transport more saturated phospholipids and the proportion of both saturated fatty acids
and cardiolipin is increased, especially at low temperature. The fluidity of the membrane
is also lessened by the low temperature. In these conditions, transport of phospholipids
between the membranes is impeded, due to clogging of TamB by a combination of less
saturated phospholipids and/or bulky and highly disordered CL. This leads to a lethal lack
of phospholipids in the OM and overabundance of phospholipids in the IM. **(C)** When
tamB is deleted in the Δ*yhdP* Δ*fadR* strain, the clogging of TamB is relieved, and the cell
relies on YdbH for phospholipid transport overcoming the cold sensitivity. PL:
phospholipid