1 **Characterization of Tigurilysin, a Novel Human CD59-Specific Cholesterol-Dependent**
2 **Cytolysin, Reveals a Role for Host Specificity in Augmenting Toxin Activity**
3
4
5

6 Ifrah Shahi[a], Sophia A. Dongas[a], Juliana K. Ilmain[b], Victor J. Torres[b], Adam J. Ratner[a,b#]
7
8
9
10 [a]New York University Grossman School of Medicine, Department of Pediatrics, New York, NY
11 [b]New York University Grossman School of Medicine, Department of Microbiology, New York, NY
12
13
14 Running Title: Host Restriction and Toxin Activity
15
16
17 [#]Address correspondence to Adam J. Ratner, Adam.Ratner@nyulangone.org
18
19

ABSTRACT

Cholesterol dependent cytolysins (CDCs) are a large family of pore forming toxins, produced by

numerous gram-positive pathogens. CDCs depend on host membrane cholesterol for pore

formation; some CDCs also require surface associated human CD59 (hCD59) for binding,

conferring specificity for human cells. We purified a recombinant version of a putative CDC

encoded in the genome of *Streptococcus oralis subsp. tigurinus*, tigurilysin (TGY), and used

CRISPR/Cas9 to construct hCD59 knockout (KO) HeLa and JEG-3 cell lines. Cell viability assays

with TGY on WT and hCD59 KO cells showed that TGY is a hCD59-dependent CDC. Two variants

of TGY exist among *S. oralis subsp. tigurinus* genomes, only one of which is functional. We

discovered that a single amino acid change between these two TGY variants determines its

activity. Flow cytometry and oligomerization western blots revealed that the single amino acid

difference between the two TGY isoforms disrupts host cell binding and oligomerization.

Furthermore, experiments with hCD59 KO cells and cholesterol depleted cells demonstrated

that TGY is fully dependent on both hCD59 and cholesterol for activity, unlike other known

hCD59-dependent CDCs. Using full-length CDCs and toxin constructs differing only in the

binding domain, we determined that having hCD59-dependence leads to increased lysis

efficiency, conferring a potential advantage to organisms producing hCD59-dependent CDCs.

IMPORTANCE

Cholesterol dependent cytolysins (CDCs) are produced by a variety of disease-causing bacteria,

and may play a significant role in pathogenesis. Understanding CDC mechanisms of action

provides useful information for developing anti-virulence strategies against bacteria that utilize

CDCs and other pore-forming toxins in pathogenesis. This study describes for the first time a

novel human-specific CDC with an atypical pore forming mechanism compared to known CDCs.

In addition, this study demonstrates that human-specificity potentially confers increased lytic

efficiency to CDCs. These data provide a possible explanation for the selective advantage of

developing hCD59-dependency in CDCs and the consequent host restriction.

INTRODUCTION

Pore forming toxins (PFTs) are produced by many pathogenic bacteria and form a diverse class of bacterial protein toxins (1). PFTs likely contribute to bacterial pathogenesis in several ways, including enhancement of colonization (2-4), release of host cell nutrients (5), hijacking of host cell pathways (6, 7), and intracellular delivery of other virulence factors (8, 9). PFTs function by perforating cell surface membranes, often leading to osmotic lysis, or triggering other downstream effects, such as pro-inflammatory cell signaling cascades, membrane repair, or programmed cell death (10).

Cholesterol dependent cytolysins (CDCs) are a family of PFTs produced by many gram-positive bacteria (10, 11), as well as some gram-negative bacteria (12). CDCs are characterized by large pores and by a dependency on host membrane cholesterol (11, 13, 14). For most CDCs, the presence of cholesterol in eukaryotic cell plasma membranes is sufficient for targeting and lysing a cell (11). A subset of CDCs additionally requires human CD59 (hCD59), a cell surface molecule, for binding to host membranes.

CDC protein structures include four domains (15, 16). During pore formation, Domain 4 binds to the target cell membrane, anchoring the CDC to the cell surface. For most CDCs, this entails binding to host cell membrane cholesterol via a threonine-leucine pair conserved within the Cholesterol Recognition Motif (CRM), followed by insertion of several other structural loops including a conserved 11-amino acid sequence (undecapeptide [UDP]), providing anchorage and stability. For hCD59-dependent CDCs, the toxin first binds to hCD59 via a conserved sequence in Domain 4, before further being stabilized by cholesterol binding (16-18). Once a CDC is anchored to the host cell surface, >30 units oligomerize to form a beta-barrel pore (19-21). To date, four hCD59-dependent CDCs have been characterized: intermedilysin (ILY) produced by *Streptococcus intermedius*, lectinolysin (LLY) produced by *Streptococcus mitis*, vaginolysin (VLY) produced by *Gardnerella vaginalis*, and most recently, discoidinolysin (DLY) produced by *Streptococcus mitis* (17, 22-24). Since there are few established hCD59-dependent

CDCs, it is difficult to form a comprehensive overview of the evolution of hCD59-dependency

and how that contributes to virulence for the bacteria producing such CDCs.

Streptococcus oralis subsp. tigurinus is a commensal bacterium found in the human oral

microbiome that has been implicated in invasive infections such as infective endocarditis,

spondylodiscitis, and meningitis (25-28). It is an α-hemolytic, non-motile, and non-spore-

forming bacterium. The *Streptococcus oralis subsp. tigurinus* genome includes an open reading

frame with sequence similarity to other hCD59-dependent CDC genomes (17, 18). Using a

recombinant protein, we characterized this novel hCD59-dependent CDC, tigurilysin (TGY). Of

the two TGY alleles present in the NCBI database, one encodes a protein that is inactive at

physiological concentrations, and we identified a single amino acid change in Domain 4

responsible for its activity. Furthermore, we delved into the question of hCD59-dependency

and showed that while targeting cells with hCD59 restricts such CDCs to only one species, it also

increases their lytic efficiency on epithelial cells by several orders of magnitude.

RESULTS

There are two variants of the *Streptococcus oralis subsp. tigurinus* CDC gene

Whole genome sequencing of three *Streptococcus oralis subsp. tigurinus* strains has revealed

that some, but not all, strains have an open reading frame encoding a protein with similarity to

previously characterized CDCs (29). We compared the amino acid sequences of *Streptococcus*

oralis subsp. tigurinus CDC (tigurilysin [TGY]) from two strains: AZ_3a and AZ_14. Five amino

acid differences were found between the mature predicted amino acid sequences of the two

variants (**Fig 1A**). We also compared TGY with amino acid sequences from other CDCs. Previous

studies with ILY have revealed that the toxin binds to hCD59 via residues separate from those of

the cholesterol-recognizing motifs (17, 30). Comparison among CDC amino acid sequences has

shown hCD59-binding residues to be conserved among hCD59-binding CDCs, and they are also

found in the TGY amino acid sequence. Furthermore, the UDP of hCD59-binding CDCs harbors a

proline in the 9th position instead of a tryptophan (18); this proline is also seen in the TGY amino

acid sequence. Based on these observations, we predicted that TGY was likely a hCD59-

dependent toxin (**Fig 1B**).

Recombinant His-tagged versions of TGY based on the sequences from both strains were

created, expressed in *Escherichia coli*, and purified. In order to assess the role of hCD59 in host

cell susceptibility to TGY, we utilized HeLa cells, a human cell line that expresses hCD59 (**Fig S1**).

We used cells that were transduced with CRISPR-Cas9 single guide RNA targeting the hCD59

gene (HeLa hCD59 Knockout [KO]) and HeLa control cells that have been transduced with a non-

targeting sgRNA (HeLa NT). We have previously used these cell lines to successfully examine ILY

activity and hCD59-dependency (31). The two recombinant TGY variants were tested for lysis on

the HeLa NT cells and HeLa hCD59 KO cells, and IC_{50} values (i.e. concentration required for 50%

lysis of HeLa cells) were calculated. AZ_14 TGY (TGY[14]) lysed HeLa cells in a hCD59-dependent

manner (IC_{50}: 460 ng/ml), albeit at concentrations higher than those needed for lysis by ILY

(IC_{50}: 30 ng/ml). In contrast, the AZ_3a TGY (TGY[3a]) did not induce lysis at the concentrations

tested, appearing to be inactive (**Fig 1C and 1D**). When TGY[3a] was further tested on HeLa cells

at supraphysiological concentrations, it was found to cause lysis in a hCD59-dependent manner,

indicating that the toxin is functional but inefficient, requiring high dosages for functionality

(IC_{50}: 261 μg/ml) (**Fig 1E**). To confirm that this hCD59-dependent lysis is not specific to HeLa

cells only, both TGY variants were also tested for lysis on JEG-3 cells (a human choriocarcinoma

cell line) with and without hCD59 expression, with results similar to those on HeLa cells: TGY[14]

lysed the cells in a hCD59-dependent manner, while TGY[3a] remained inactive at the

concentrations tested (**Fig S2**).

TGY residue at position 454 is important for lytic efficiency

To understand the cause of TGY[3a] inefficiency, we turned our attention to the amino acid

sequence differences between the two TGY variants. Of the five single amino acid differences,

two are in Domain 4: residue 454 and residue 505 (**Fig 1A**). Residue 505 incorporates an

isoleucine in TGY[3a] and a leucine in TGY[14]. Based on amino acid sequences of other CDCs, this

residue position seems to be permissible for both amino acids – some CDCs have an isoleucine

while others have a leucine, without any correlation to the species specificity of the CDC.

Conversely, residue 454 is part of the short hydrophobic loop known as Loop 2 (L2), one of the

three conserved CDC loops that insert into the host membrane in a cholesterol-dependent

manner (32, 33). In most CDCs, the corresponding amino acid in L2 is an alanine; however, non-

functional TGY[3a] harbors a glutamic acid (E) at this position, while the functional TGY[14] has a

glycine (G). Additionally, both TGY variants have an isoleucine (I) instead of a leucine (L) at

position 544, which is part of the conserved CRM threonine-leucine pair.

Using site-directed mutagenesis, we made three amino acid substitutions in TGY[3a], resulting in

TGY[3a_E454G], TGY[3a_E454A] and TGY[3a_I544L]. These variants were tested on HeLa NT cells for

lysis. The results demonstrated that the single amino acid change at position 454 restored the

toxin's lytic activity (**Fig 2A**), while the amino acid change in the CRM had no effect.

TGY[3a_E454G] and TGY[3a_E454A] were further tested for lysis on HeLa hCD59 KO cells, confirming

their cytotoxicity to still be hCD59-dependent (**Fig 2B**).

TGY is fully dependent on both cholesterol and hCD59

Of the known hCD59-dependent CDCs, only ILY is completely non-functional in the absence of

hCD59 – it requires hCD59 to initiate binding to a host cell, though interaction with cholesterol

is still necessary for the subsequent steps of pore formation (34, 35). In contrast, VLY can bind

host cell membrane cholesterol in the absence of hCD59 (though less efficiently than binding

hCD59) (36), and LLY is similar (37). To assess whether TGY requires cholesterol in a manner

similar to ILY or VLY/LLY, HeLa cells were depleted of cholesterol using methyl-ß-cyclodextrin

(MßCD) and tested with TGY variants for lysis (**Fig 3A and 3B**). For comparison, similar assays

were conducted with ILY (fully hCD59-dependent), VLY (partially hCD59-dependent), and

pneumolysin (PLY; a non-specific CDC) (**Fig 3C, 3D, and 3E**).

Our results show that ILY does not lyse HeLa hCD59 KO cells but remains active on cholesterol

depleted HeLa cells at higher toxin concentrations. VLY, as a partially hCD59-dependent CDC,

lyses both HeLa hCD59 KO cells and cholesterol depleted HeLa cells at higher toxin

concentrations. PLY, as a non-specific CDC, lyses HeLa hCD59 KO cells at similar levels as HeLa

NT cells, but is inactive on cholesterol depleted HeLa cells. In contrast, TGY only lyses HeLa NT

cells and does not kill cells either lacking hCD59 or depleted of cholesterol. To ensure that

cholesterol depletion does not perturb hCD59 expression (and thereby indirectly affect lysis

results of hCD59-dependent toxins), hCD59 cell surface expression was analyzed using flow

cytometry (**Fig S1**). Results demonstrate that cholesterol depleted cells express similar levels of

hCD59 as HeLa NT cells. Taken together with the lysis data, our results suggest that TGY has a

distinct CDC lysis mechanism with an absolute requirement for both cholesterol and hCD59.

TGY binding to host cells is affected by both cholesterol and hCD59 presence

We set out to investigate sequential stages of CDC pore formation for TGY, starting with host

cell binding. HeLa cells were incubated with His-tagged TGY variants for flow cytometry analysis

(**Fig 4A**). The same experiment was also carried out with ILY, VLY and PLY (**Fig 4B**). The

percentage of PE-positive cells (indicating percentage of cells with detectable toxin bound to

their surface) and the mean fluorescence intensity (MFI – indicating the density of toxin bound)

for all conditions are listed in **Table 1**.

As expected, ILY bound well to cholesterol depleted cells, but not to hCD59 KO cells. VLY bound

cells in the absence of either cholesterol or hCD59, although more binding was detected on

cholesterol depleted cells than hCD59 KO cells. PLY bound well to hCD59 KO cells but showed

reduced binding to cholesterol depleted cells (**Fig 4B**). Since PLY is known to require cholesterol

for toxin function, the ~32% of PE-positive cholesterol depleted cells suggest that cholesterol

depletion is not completely effective; however, based on lysis data, the cholesterol depletion is

sufficient to block PLY-mediated lysis.

For the TGY variants (**Fig 4A**), the flow cytometry results suggest that TGY^{3a} is non-functional

due to a defect in host cell binding: only a small percentage of TGY^{3a}-exposed cells were PE-

positive. The results also indicate that the reduction in binding is due to the glutamic acid in L2

of TGY^{3a}, as there was a significant increase in PE fluorescence for TGY$^{3a\_E454G}$-incubated cells

compared to TGY^{3a}-incubated cells, both in the percentage of PE-positive cells and the MFI.

Additionally, L2 of TGY^{3a} seems to mediate interaction with the host cell in a cholesterol-

dependent manner, as there is minimal binding of TGY$^{3a\_E454G}$ to cholesterol depleted cells.

This is consistent with previously published findings, which show that L2 inserts into host cell

membranes in a cholesterol-dependent manner (32, 33, 35). In hCD59 KO cells, TGY$^{3a\_E454G}$

bound ~60% of the cells, although the significantly lower MFI (compared to TGY$^{3a\_E454G}$-

incubated control cells) suggests fewer toxin units per cell. This finding suggests that

TGY$^{3a\_E454G}$ may bind to cholesterol before interacting with hCD59, but that hCD59 is still

required for pore formation and lysis, possibly by playing a role in oligomerization – the lower

MFI could indicate a lack of concentration of toxin monomers on the surface that occurs during

oligomerization.

Of note, TGY^{14} and TGY$^{3a\_E454G}$ have similar lysis curves but differentially bind to cholesterol

depleted cells: TGY^{14} binds to a significantly higher percentage of cholesterol depleted cells

than TGY$^{3a\_E454G}$ (**Fig 4A**). This could imply a higher affinity for hCD59 or a higher rate of

oligomerization. Furthermore, even though TGY^{14} can bind to a large percentage of cholesterol

depleted cells and both functional TGY variants are able to bind to ~50-60% of hCD59 KO cells,

neither of those cell lines are lysed by these toxins (**Fig 3A and 3B**). Taken together, these data

suggest that differences in binding ability alone do not account fully for the lysis phenotypes of

TGY variants and that subsequent steps of pore formation also play key roles in lysis

differences.

Functional TGY requires both cholesterol and hCD59 for oligomerization

We next investigated the oligomerization of TGY on the host cell surface. HeLa cells were

incubated with toxins on ice, before the cells were lysed and prepared for SDS-AGE (20).

Glutaraldehyde was used as a cross-linker to preserve oligomer complexes and prevent them

from breaking down during sample preparation. For comparison, similar analyses were also

carried out with ILY, VLY and PLY.

The resulting western blots demonstrate oligomer "streaks" for ILY and VLY on HeLa NT and

HeLa hCD59 KO cells (**Fig S3A**). PLY oligomerization was only evident in both cell lines at higher

toxin concentrations (**Fig S3B**), which was unsurprising as the lower PLY concentration causes

only ~10-15% cell death in HeLa NT cells. Oligomerization was not detectable for ILY, VLY and

PLY on cholesterol depleted cells (**Fig S3C**).

SDS-AGE analysis with TGY showed visible oligomers for both functional TGY variants (TGY[14]

and TGY[3a_E454G]) on control cells but a lack of visible oligomers for TGY[3a]. Moreover, depleting

host cells of either cholesterol or hCD59 abrogated oligomerization for both TGY[14] and

TGY[3a_E454G] (**Fig 5**). Overall, our data indicate that TGY is dependent on both cholesterol and

hCD59 for binding as well as oligomerization.

hCD59-dependent CDCs show increased lysis activity, dependent on host cell hCD59

expression levels

We observed that the concentrations of hCD59-dependent CDCs ILY, VLY and TGY required for

lysis differ from those of the non-specific CDC PLY. To investigate how hCD59-dependency

affects the overall lytic ability of CDCs, we tested various CDCs for lysis on HeLa cells (**Fig 6A and**

6B). In the non-hCD59-dependent subset, we tested PLY and inerolysin (INY) (38). In the hCD59-

dependent subset, we tested ILY, VLY, TGY[3a], TGY[3a_E454G] and TGY[14]. To quantify lysis,

nonlinear regression analysis was performed on the data to obtain IC_{50} values for each toxin

(**Table 2**). All functional hCD59-dependent toxins had lower IC_{50} values than non-hCD59-

dependent toxins on HeLa NT control cells. VLY did not behave in a fully hCD59-dependent

manner, lysing cells in the absence of hCD59 expression but with lower efficiency (i.e. higher

IC_{50} value), as previously indicated in the literature (36). Quantitatively, VLY lyses HeLa control

cells at concentrations comparable to ILY (a hCD59-dependent CDC), and HeLa hCD59 KO cells

at concentrations similar to PLY and INY (non-hCD59-dependent CDCs) (**Table 2**). These data

indicate the possibility that VLY may form pores in a manner similar to either hCD59-dependent

CDCs or non-hCD59-dependent CDCs, depending on the characteristics of the target cell. In

addition, our data raise the possibility that hCD59-dependency, while restricting a CDC to a

single species, might increase lysis efficiency on some target cells.

To investigate if these lysis results are relevant to other human cell lines, we repeated the LDH

release assays with various CDCs on JEG-3 cells (**Fig 6C-6D**). Once again, hCD59-dependent

toxins (except for TGY variants) had lower IC_{50} values than non-hCD59-dependent toxins on

JEG-3 NT control cells. However, ILY consistently lysed JEG-3 NT cells at IC_{50} values ~2.5 times

higher than HeLa NT cells, and TGY variants lysed JEG-3 NT cells at ~14-16 times higher IC_{50}

values than HeLa NT cells (**Table 2**).

We hypothesized that the difference in lysis efficiency of hCD59-dependent CDCs on two

different human cell lines could be affected by the amount of hCD59 expressed on the surface

of these cells. To investigate this, we analyzed the cell surface expression of hCD59 on HeLa

cells and JEG-3 cells using flow cytometry (**Fig 6E**). The results show that HeLa cells express ~15

times more hCD59 than JEG-3 cells. These data suggest that changes in hCD59 expression levels

between different human cell lines can potentially affect the efficiency of hCD59-dependent

CDC pore formation and lysis.

hCD59-dependency results in increased efficiency of lysis by CDCs

Since Domain 4 is the key player that controls toxin specificity (16-18), we created hybrid CDCs

to isolate the role of hCD59-dependency in lysis efficiency. To do this, the Domain 4 of five

different CDCs was attached to Domains 1-3 of PLY, a non-hCD59 dependent CDC, resulting in

the following: PLY-INY, PLY-ILY, PLY-VLY, PLY-TGY[3a], PLY- TGY[3a_E454G] and PLY-TGY[14].

Lysis caused by each of the hybrid CDCs was measured in HeLa NT and HeLa hCD59 KO cells.

The presence of a functional hCD59-dependent Domain 4 increased lysis efficiency of PLY-ILY

and PLY-VLY when compared to full-length PLY, while the presence of a non-functional Domain

4 (PLY-TGY[3a]) rendered the hybrid inactive. PLY- TGY[3a_E454G] and PLY- TGY[14] also

demonstrated a decrease in IC_{50} compared to full-length PLY, albeit less than PLY-ILY and PLY-

VLY (**Fig 7A and Table 3**). All hybrid toxins exhibited hCD59-dependency as governed by each

Domain 4 (**Fig 7B**). PLY-INY (with a non-hCD59-dependent Domain 4) exhibited a slight

decrease in IC_{50} compared to full-length PLY, which could be expected considering full-length

INY has a lower IC_{50} than PLY (**Table 3**). PLY-VLY notably retained the 'intermediate' nature of its

Domain 4 parent: it lysed HeLa NT control cells at low concentrations, and HeLa hCD59 KO cells

at concentrations comparable to non-hCD59-dependent PLY, as evidenced by its IC_{50} values.

(**Fig 7A, 7B and Table 3**). Overall, our data suggest that in most cases, while hCD59-dependency

restricts CDC targets to a single species, it increases the efficiency of pore formation and lysis

for those toxins.

DISCUSSION

Cholesterol dependent cytolysins (CDCs) act as factors in pathogenesis for a wide range of

disease-causing bacteria. Understanding their mechanism of action is a key step towards

forming therapeutic strategies against such bacteria. Our study adds to the growing knowledge

regarding CDCs by characterizing a novel hCD59-dependent CDC (TGY) produced by

Streptococcus oralis subsp. tigurinus, a bacterial species implicated in various infections (25,

26). Our data demonstrate that of the two TGY variants, TGY[3a] is non-functional due to a

defective L2, which prevents it from binding to host cells and oligomerizing. Furthermore, TGY

is fully hCD59-dependent and fully cholesterol dependent, setting it apart from other known

CDCs which are generally grouped into categories based on hCD59-dependency (37, 39) but

none are known to be equally dependent on both cholesterol and hCD59 for lysis. Overall, TGY

seems to exhibit an atypical pore forming mechanism compared to CDCs such as PLY, VLY and

ILY.

Our flow cytometry data investigated host cell binding for various CDCs. Binding data with PLY

showed that cholesterol depletion for our cells was not complete, with ~30% cells still being

bound by PLY (a fully cholesterol dependent CDC). However, the large shift in PE fluorescence

for PLY on control cells versus cholesterol depleted cells, coupled with the PLY lysis data,

indicated the cholesterol depletion is sufficient to abolish lysis by nonspecific CDCs. For ILY and

VLY, ~100% of the cholesterol depleted cells are still bound by the toxins but there is a

reduction in MFI. This could be due to reduced oligomerization, as fewer oligomers would mean

less toxin detectable on the cell surfaces.

TGY[3a] appears to be non-functional largely due to a defect in binding to host cells. TGY[3a_E454G]

did not bind well to cholesterol depleted cells but could partially bind hCD59 KO cells. This is

unlike ILY, which is thought to bind hCD59 before cholesterol (18, 22) and, in our experiments,

bound well to cholesterol depleted cells but not to hCD59 KO cells. Notably, TGY[14] bound to

cholesterol depleted cells much better than TGY$^{3a\_E454G}$, though the reason for this aberration

is unclear.

328
We conducted SDS-AGE to visualize oligomerization of CDCs on host cell surfaces. Previously,

hCD59 dependent CDCs have been shown to require either cholesterol or hCD59 for

oligomerization, but not both factors together: ILY can oligomerize on cholesterol depleted cells

but requires hCD59 for oligomerization (22, 34, 40), while VLY has been shown to oligomerize

on liposomes lacking hCD59 (36). Our oligomerization assays demonstrated no visible oligomers

for ILY and VLY in the absence of either hCD59 or cholesterol. The oligomerization data for ILY

and VLY correspond to our binding data, which suggested loss of oligomers on cholesterol

depleted cells as discussed above. It is relevant to note, however, that all binding and

oligomerization data was based on a toxin concentration of 5 µg/ml, which was sufficient for

lysis of relevant cell lines when cells are incubated at 37°C for 1.5 hours, but might not be

enough for oligomerization when cells are incubated for a shorter time period on ice (as during

sample preparation for flow cytometry and SDS-AGE). Therefore, despite ILY and VLY lysing

cholesterol depleted cells, and VLY lysing hCD59 KO cells (**Fig 3 and 4**), we observe reduced

binding and no oligomerization.

343
344   SDS-AGE with TGY variants showed that both hCD59 and cholesterol are important for

345   oligomerization of functional TGY. Consequently, even though TGY$^{3a\_E454G}$ bound to about half

346   of hCD59 KO cells, and TGY^{14} bound partially to both cholesterol depleted cells and hCD59 KO

347   cells, no lysis was seen on these cell lines, likely due to a lack of sufficient oligomerization.

348
Overall, our data demonstrate that TGY residue 454 significantly affects the host cell binding,

modulating toxin activity. Furthermore, both cholesterol and hCD59 are required for

oligomerization. Our data also imply that the pore forming mechanism of TGY and its

interactions with hCD59 and cholesterol potentially vary from those of other hCD59-dependent

CDCs. It is prudent to note that the idea of an atypical CDC pore-formation mechanism is not

entirely unique. ILY and VLY (both hCD59-dependent CDCs) bind hCD59 differently: VLY has a

lower affinity for hCD59 than ILY, and its complex with hCD59 has a different conformation (18).

The UDP of VLY can also adopt two different conformations upon interacting with host cell

membrane, whereas ILY UDP consistently adopts one orientation (18). The CDC arcanolysin

(ALN) displays varying specificity for different mammalian cell lines in a non-hCD59-dependent

manner (41). There has also been recent evidence that CDCs might interact with glycans or

heparan sulfates on host cell surfaces (42-44). It is therefore possible that host cell factors other

than hCD59 and cholesterol, as well as conformational variations in the toxin structure, could

play roles in the TGY pore forming mechanism.

The second part of our study aimed to understand the selective pressure behind the evolution

of hCD59-dependency in some CDCs. Most CDCs are nonspecific, which means they can target

and lyse cells from different species. Conversely, hCD59-dependency restricts CDCs to targeting

only human cells. Furthermore, it has been speculated that as hCD59 is present on all human

cells, hCD59-dependency would not lead to greater specificity for a particular cell type (45). The

data presented in this study demonstrate that hCD59-dependent CDCs like ILY lyse HeLa cells

with higher efficiency than non-hCD59-dependent CDCs like PLY and INY. We also showed that

CDCs such as VLY behave like intermediate CDCs, lysing cells at low concentrations comparable

to ILY when hCD59 is accessible and at higher concentrations comparable to PLY/INY when

hCD59 is not available. Additionally, our data showed a difference in hCD59-dependent toxin

concentrations needed for lysis of HeLa cells than those needed for JEG-3 cells. This implied

that host factors other than the binary presence or absence of hCD59 affected lysis efficiency of

these toxins. Flow cytometry data for hCD59 expression on the surface of HeLa and JEG-3 cells

revealed a significantly lower amount of hCD59 expressed on JEG-3 cells. The fact that higher

concentrations of hCD59-dependent toxins are required to lyse JEG-3 cells leads to the theory

that while all human cells express hCD59, differing levels of hCD59 expression could allow

hCD59-dependent CDCs to preferentially target some cell types over others in the human body.

Understanding CDC pore-formation is an important step towards developing strategies that can

help mitigate PFT-driven bacterial virulence. To this end, our study adds to the growing

knowledge around human-specific CDCs by investigating the function of a novel hCD59-

dependent CDC. Furthermore, our comparative functional analysis on a range of CDCs

endeavors to help us understand the development of human-specificity within this broad family

of pore forming toxins.

MATERIALS AND METHODS

Cell Culture

HeLa cells (ATCC #CCL-2) and JEG-3 cells (ATCC #HTB-36) were cultivated in EMEM medium

supplemented with 10% heat-inactivated fetal bovine serum (FBS) and penicillin/streptomycin.

For cells transduced with lentivirus to create NT control cells and hCD59 KO cells, the media

was further supplemented with 0.9μg/ml puromycin. Cells were grown in humidified incubators

at 37°C and 5% CO_2.

Generation of Knockout Cell Lines

NT control and hCD59 KO cell lines were generated as previously described (31). Briefly, single

guide RNAs (sgRNAs) were ligated into an empty pLentiCRISPR plasmid backbone (gifted by Dr

Feng Zhang; *Addgene plasmid #52961* (46)) and packaged into lentivirus using XtremeGENE 9

DNA Transfection Reagent. Cells were transduced with the lentivirus using spinfection. To

generate single cell clones from polyclonal KO cell lines, limiting dilutions were performed.

Cell Viability Assays (LDH)

LDH Cytotoxicity assay (*Sigma-Aldrich 11644793001*) to measure lysis was conducted according

to the manufacturer's instructions. Briefly, 2.0 x 10^4 HeLa cells were seeded per well in a 96-

well plate overnight. For cholesterol depletion, cells were incubated with 4 mg/ml methyl-ß-

cyclodextrin (MßCD) for 1 hour and washed with PBS. Cells were incubated in appropriate

dilutions of toxin in fresh medium for 1.5 hours. 100 μl of the supernatant was transferred to a

new 96-well plate and 100 μl of Reaction mixture consisting of dye solution and catalyst

(prepared according to the manufacturer's protocol) added. Plates were incubated in the dark

for 30 mins. Absorbance of samples was measured at 492nm.

Recombinant Toxins and Hybrid Toxins

Hybrid toxins PLY- TGY^{3a} and PLY- TGY^{14} were created using Gibson assembly. Primers were

designed to isolate PLY Domains 1-3 and the Domain 4 of TGY variants using NEB Builder tool,

with overhangs to ensure the alignment of each Domain 4 with PLY Domain 3. The primers

were used to PCR amplify the relevant fragments, which were then gel-extracted. A pET28a

vector was double-digested using NdeI and XhoI, and gel-extracted. Gibson assembly was

carried out with the digested pET28a vector, PLY Domains 1-3, and TGY Domain 4. The reaction

product was transformed into NEB 5-alpha cells and minipreps prepared, before being

sequenced using T7 primers.

Recombinant PLY, INY, ILY, VLY, PLY-ILY, PLY-VLY, and PLY-INY were prepared as described (38).

Full-length and hybrid toxins (TGY^{3a}, TGY^{14}, TGY$^{3a\_E454G}$, TGY$^{3a\_E454A}$, TGY$^{3a\_I544L}$, PLY-

TGY^{3a}, PLY- TGY$^{3a\_E454G}$ and PLY- TGY^{14}) were purified as follows. Briefly, the gene encoding

each toxin was codon-optimized for *E. coli*, cloned into pET28a, and transformed into *E. coli*

T7Iq. Protein expression was induced by the addition of IPTG. Cells were lysed with a lysis

buffer (50 mM NaH$_2$PO$_4$, 300 mM NaCl, 10 mM imidazole). The toxin was purified from the

lysate using HisTrap columns with an FPLC unit into elution buffer (50mM NaH$_2$PO$_4$, 300mM

NaCl, 250mM imidazole), then buffer exchanged into PBS using Amicon Ultra-4 filters. The

concentration of each toxin was measured using a Bradford Assay.

Site-Directed Mutagenesis

For mutagenesis of TGY residue 454 and 544, the New England Biolabs Q5 Site Directed

Mutagenesis kit (*NEB E0552S*) was used according to the manufacturer's protocols. Briefly, the

NEBaseChanger tool was used to generate primers to make the relevant substitutions. These

primers were used to PCR amplify the mutated gene incorporated into a pET28a vector, and the

PCR product transformed into NEB 5-alpha cells for amplification. Minipreps were prepared and

sequenced to confirm the mutations, before transformation into T7Iq cells for protein

purification.

Flow cytometry

Flow cytometry with His-tagged toxins was carried out using direct staining. 1.0×10^6 cells were

washed and incubated with 5 μg/ml of each toxin for 10 minutes on ice. Cells were washed with

PBS three times and resuspended in 100 μl of ice-cold PBS + 10% FBS + 1% sodium azide.

Phycoerythrin (PE) anti-His tag antibody (*BioLegend 362603*) was added to the cell suspension

at a dilution of 1:20 and incubated for 30 minutes on ice in the dark. Cells were washed with

ice cold PBS, and resuspended in 300 μl of ice-cold PBS + 3% BSA + 1% sodium azide. Flow

cytometry to visualize hCD59 expression was carried out using indirect staining as described

previously (31). Stained cells were visualized on a CytoFlex Analyzer flow cytometer, and the

data analyzed using FlowJo software (version 10.8.1). Cells were gated as follows: live cells

using forward scatter area and side scatter area; single cells using forward scatter area and

forward scatter height; fluorescence-positive cells using PE-fluorescence intensity or AF647

fluorescence intensity, and forward scatter height. Samples contained at least 10,000 gated live

cells.

Oligomerization western blot

To visualize CDC oligomers, SDS-AGE analysis was carried out as described (20). Briefly, 1.0×10^6

cells were washed then incubated with 5 μg/ml of each toxin (or 30 μg/ml for PLY where

indicated) for 10 minutes on ice. Cells were washed twice with PBS and once with HEPES/NaCl

buffer (20 mM HEPES at pH 7.5, 150 mM NaCl), before being resuspended in 40 μl of 0.01%

glutaraldehyde in HEPES/NaCl as a cross-linker and incubated at 37°C for 15 minutes. 0.1M Tris

(pH 8.0) was added to stop the cross-linking reaction. Tris-Glycine SDS Sample buffer (*LC2676;*

ThermoFisher) was added to the cell solution at a 1:1 ratio and the mixture incubated at 95°C

for 20 minutes to lyse the cells.

The lysate was loaded onto a 1.5% agarose gel. Proteins from the gel were transferred to a

polyvinylidene difluoride (PVDF) membrane using semi-wet transfer, blocked in 5% nonfat dry

milk in Tris-buffered saline (TBS) plus 1% Tween-20, and incubated with 6x His-tag monoclonal

primary antibody (*MA1-135; ThermoFisher*) overnight at a dilution of 1:750. The membrane was

washed with TBS plus 1% Tween-20, and incubated with goat anti-mouse IgG secondary

antibody (*31430; ThermoFisher*) for 1 hour at a 1:10,000 dilution. The membrane was then

washed with several changes of TBS plus 1% Tween-20 for 3 hours before being imaged on

iBright CL1000 using Pierce ECL Western Blotting Substrate.

ACKNOWLEDGEMENTS

This work was supported by grant R01 AI155476 to AJR.

Figure 1: Two variants of novel CDC tigurilysin (TGY) exist, with differing lysis activity

(**A**) The predicted amino acid sequences of the two mature TGY variants. The signal sequence

comprising the first 44 amino acids, as predicted by SignalP 5.0 (SignalP 5.0 - DTU Health Tech -

Bioinformatic Services), was eliminated from the sequences shown here. There are 5 single

amino acid differences (shaded) between TGY^{3a} and TGY^{14}. The last two (amino acid 454 and

amino acid 544) are found in domain 4.

(**B**) Left panel: the conserved hCD59-binding sequence (Y-X-Y-X14-R-S-R; underlined) in Domain

4 of various hCD59-dependent CDCs, as compared to the same region of non-hCD59-dependent

CDCs. Right panel: The conserved 11-amino acid sequence (undecapeptide), necessary for CDC

interaction with cholesterol and subsequent pore formation, as found in various CDCs. The

proline at the 9th position is conserved among hCD59-dependent CDCs. The shaded regions in

both panels show the consensus sequence as determined by MacVector.

(**C-D**) Percentage cell death in HeLa NT control cells and HeLa hCD59 KO cells when exposed to

increasing concentrations of TGY^{3a} and TGY^{14}, with ILY for comparison as a known hCD59-

dependent CDC. Cells were incubated with toxins for 1.5 hours and cell death measured by an

LDH-release cytotoxicity assay. Each point is the mean of 3 replicates, and error bars represent

±SD.

(**E**) Percentage cell death in HeLa NT control cells and HeLa hCD59 KO cells when exposed to

supraphysical concentrations of TGY^{3a}. Cells were incubated with toxin for 1.5 hours and cell

death measured by an LDH-release cytotoxicity assay. Each point is the mean of 3 replicates,

and error bars represent ±SD.

506
507
Figure 2: A glutamic acid-to-glycine change at position 454 affects TGY activity
(**A**) Percentage cell death in HeLa NT control cells when exposed to increasing concentrations of
TGY^{3a} and its mutagenized counterparts. Only the amino acid change at position 454 (glutamic
acid to glycine or alanine) restored toxin functionality. Cells were incubated with toxins for 1.5
hours and cell death measured by an LDH-release cytotoxicity assay. Each point is the mean of 3
replicates, and error bars represent ±SD.
(**B**) Percentage cell death in HeLa hCD59 KO cells when exposed to increasing concentrations of
TGY^{3a} and TGY$^{3a\_E454G}$, to determine if the toxins are hCD59-dependent. Cells were incubated
with the toxins for 1.5 hours and cell death measured by an LDH-release cytotoxicity assay.
Each point is the mean of 3 replicates, and error bars represent ±SD.
518
519

520
521
522
523 **Figure 3: Functional TGY is fully cholesterol dependent.**
524 (**A-E**) Percentage cell death in HeLa NT control cells, HeLa hCD59 KO cells, HeLa cholesterol
525 depleted cells, and HeLa hCD59 KO + cholesterol depleted cells. Cells were exposed to a range
526 of concentrations of each toxin for 1.5 hours, and cell death measured by an LDH-release
527 cytotoxicity assay. Each point is the mean of 3 replicates, and error bars represent ±SD.
528

529
530
531
Figure 4: TGY3a is defective in host cell binding, while functional TGY is dependent on both
cholesterol and hCD59 for host cell binding.
(**A-B**) Flow cytometry analysis of His-tag expression on the cell surface after HeLa cells (NT
control, hCD59 KO, cholesterol depleted, and hCD59 KO + cholesterol depleted) were incubated
with His-tagged toxins, before addition of a PE anti-His-tag antibody. Histogram x-axis shows
fluorescence intensity of the PE-conjugated antibody for each cell population. Histogram y-axis
shows cell count normalized to mode. The dotted line demarcates the PE-negative cells and the
PE-positive cells, based on the fluorescence intensity of control cells not incubated with any
toxin.
541

543
544
545

Figure 5: Functional TGY is dependent on both cholesterol and hCD59 for oligomerization on host cell surface.

Oligomerization of TGY variants as assessed by SDS-AGE analysis. Cells were incubated with the indicated toxins on ice before being prepared for western blotting. 0.01% glutaraldehyde was used as a cross-linker to preserve oligomers during sample preparation. Western blot membrane was probed with an anti-His-tag HRP-conjugated antibody.

Figure 6: hCD59-dependency coupled with high hCD59 expression increases CDC ability to lyse cells at lower concentrations

(**A-B**) Percentage cell death in HeLa NT control cells and HeLa hCD59 KO cells when exposed to various full-length CDCs, both hCD59-dependent and non-hCD59-dependent. Cells were incubated with toxins for 1.5 hours and cell death measured by an LDH-release cytotoxicity assay. Each point is the mean of 3 replicates, and error bars represent ±SD.

(**C-D**) Percentage cell death in JEG-3 NT control cells and JEG-3 hCD59 KO cells when exposed to various full-length CDCs, both hCD59-dependent and non-hCD59-dependent. Cells were incubated with toxins for 1.5 hours and cell death measured by an LDH-release cytotoxicity assay. Each point is the mean of 3 replicates, and error bars represent ±SD.

(**E**) Flow cytometry analysis of hCD59 expression on the cell surface for HeLa cells and JEG-3 cells. Cells were incubated with a primary anti-hCD59 mouse IgG2a antibody and a secondary anti-mouse IgG2a AF647 antibody. Histogram x-axis shows fluorescence intensity of the APC-conjugated secondary antibody for each cell population. Histogram y-axis shows cell count normalized to mode.

Abbreviations: ILY, intermedilysin; VLY, vaginolysin; TGY, tigurilysin; INY, inerolysin; PLY, pneumolysin.

Figure 7: Incorporating a hCD59-dependent Domain 4 increases CDC ability to lyse cells at
lower concentrations
(**A-B**) Percentage cell death in HeLa NT control cells and HeLa hCD59 KO cells when exposed to
various PLY-hybrid CDCs, with both hCD59-dependent and non-hCD59-dependent D4s. Cells
were incubated with toxins for 1.5 hours and cell death measured by an LDH-release
cytotoxicity assay. Each point is the mean of 3 replicates, and error bars represent ±SD.
583

	% PE-positive cells	MFI values
HeLa NT cells only	1	112
HeLa hCD59 KO cells only	1	578
HeLa NT cholesterol depleted cells only	0	136
HeLa hCD59 KO cholesterol depleted cells only	1	138
HeLa NT + ILY	100	170246
HeLa hCD59 KO + ILY	11	6767
HeLa NT cholesterol depleted + ILY	99	9910
HeLa hCD59 KO cholesterol depleted + ILY	1	151
HeLa NT + VLY	100	150828
HeLa hCD59 KO + VLY	65	8167
HeLa NT cholesterol depleted + VLY	100	21252
HeLa hCD59 KO cholesterol depleted + VLY	2	181
HeLa NT + PLY	96	30816
HeLa hCD59 KO + PLY	95	23677
HeLa NT cholesterol depleted + PLY	32	697
HeLa hCD59 KO cholesterol depleted + PLY	24	625
HeLa NT + TGY[3a]	28	9735
HeLa hCD59 KO + TGY[3a]	6	335
HeLa NT cholesterol depleted + TGY[3a]	7	4261
HeLa hCD59 KO cholesterol depleted + TGY[3a]	1	598
HeLa NT + TGY[3a_E454G]	99	28460
HeLa hCD59 KO + TGY[3a_E454G]	60	3813
HeLa NT cholesterol depleted + TGY[3a_E454G]	34	3847
HeLa hCD59 KO cholesterol depleted + TGY[3a_E454G]	8	281
HeLa NT + TGY[14]	99	31109
HeLa hCD59 KO + TGY[14]	54	4126

HeLa NT cholesterol depleted + TGY[14]	79	2380
HeLa hCD59 KO cholesterol depleted + TGY[14]	2	232

584
585
586 **Table 1: Quantification of Flow Cytometry Data of CDC Binding on HeLa cells**
587 The percentage of PE-positive live cells were calculated by FlowJo based on gating of PE-
588 negative cells as determined by cells-only controls. The MFI for each sample was calculated by
589 FlowJo for cells gated as live cells → single cells.

	IC$_{50}$ on HeLa NT control cells (µg/ml)	IC$_{50}$ on HeLa hCD59 KO cells (µg/ml)	IC$_{50}$ ratio HeLa hCD59 KO : HeLa NT	IC$_{50}$ on JEG-3 NT control cells (µg/ml)	IC$_{50}$ on JEG-3 hCD59 KO cells (µg/ml)	IC$_{50}$ ratio JEG-3 hCD59 KO : JEG-3 NT
ILY	0.03 ± 0.01	*ND*	*NA*	0.08 ± 0.01	*ND*	*NA*
VLY	0.04 ± 0.01	8.92 ± 1.61	223.0	0.05 ± 0.03	10.73 ± 0.65	214.6
TGY[3a]	*ND*	*ND*	*NA*	*ND*	*ND*	*NA*
TGY[3a_E454G]	0.50 ± 0.18	*ND*	*NA*	7.04 ± 1.14	*ND*	*NA*
TGY[14]	0.46 ± 0.04	*ND*	*NA*	7.56 ± 4.41	*ND*	*NA*
INY	8.12 ± 3.74	8.26 ± 3.52	1.0	12.98 ± 1.95	9.99 ± 0.13	0.7
PLY	15.46 ± 4.41	17.26 ± 5.34	1.1	10.89 ± 0.95	10.48 ± 2.77	1.0

Table 2: IC$_{50}$ values of CDCs tested for lysis on HeLa and JEG-3 cells.

Full-length CDCs were tested for lysis on the indicated cell lines at a range of 0.08 µg/ml to 50

µg/ml, and IC$_{50}$ values indicate the concentrations required for 50% lysis on each cell line. Each

IC$_{50}$ value average with standard deviation is calculated from 3 experimental repeats. "ND"

indicates where the CDC did not cause any cell lysis; "NA" indicates where the IC$_{50}$ ratio could

not be calculated.

	IC$_{50}$ on HeLa NT control cells (µg/ml)	IC$_{50}$ on HeLa hCD59 KO cells (µg/ml)	IC$_{50}$ ratio HeLa hCD59 KO : HeLa NT
PLY	15.46 ± 4.41	17.26 ± 5.34	1.1
PLY-ILY	1.82 ± 0.69	*ND*	*NA*
PLY-VLY	0.67 ± 0.26	37.60 ± 16.80	56.1
PLY-TGY[3a]	*ND*	*ND*	*NA*
PLY-TGY[3a_E454G]	9.48 ± 0.79	*ND*	*NA*
PLY-TGY[14]	11.26 ± 3.73	*ND*	*NA*
PLY-INY	8.66 ± 0.53	9.69 ± 0.44	1.2

Table 3: IC$_{50}$ values of hybrid CDCs tested for lysis on HeLa cells.

Hybrid CDCs with PLY D1-3 were tested for lysis on the indicated cell lines at a range of 0.08

µg/ml to 50 µg/ml, and IC$_{50}$ values indicate the concentrations required for 50% lysis on each

cell line. Full-length PLY IC$_{50}$ values are included for comparison. Each IC$_{50}$ value average with

standard deviation is calculated from 3 experimental repeats. "ND" indicates where the CDC did

not cause any cell lysis; "NA" indicates where the IC$_{50}$ ratio could not be calculated.

REFERENCES

1. Gonzalez MR, Bischofberger M, Pernot L, van der Goot FG, Freche B. 2008. Bacterial pore-forming toxins: the (w)hole story? Cell Mol Life Sci 65:493-507.
2. Stanley SA, Raghavan S, Hwang WW, Cox JS. 2003. Acute infection and macrophage subversion by Mycobacterium tuberculosis require a specialized secretion system. Proc Natl Acad Sci U S A 100:13001-6.
3. Hensler ME, Liu GY, Sobczak S, Benirschke K, Nizet V, Heldt GP. 2005. Virulence role of group B Streptococcus beta-hemolysin/cytolysin in a neonatal rabbit model of early-onset pulmonary infection. J Infect Dis 191:1287-91.
4. Orihuela CJ, Gao G, Francis KP, Yu J, Tuomanen EI. 2004. Tissue-specific contributions of pneumococcal virulence factors to pathogenesis. J Infect Dis 190:1661-9.
5. Geny B, Popoff MR. 2006. Bacterial protein toxins and lipids: pore formation or toxin entry into cells. Biol Cell 98:667-78.
6. Lim JH, Stirling B, Derry J, Koga T, Jono H, Woo CH, Xu H, Bourne P, Ha UH, Ishinaga H, Xu H, Andalibi A, Feng XH, Zhu H, Huang Y, Zhang W, Weng X, Yan C, Yin Z, Briles DE, Davis RJ, Flavell RA, Li JD. 2007. Tumor suppressor CYLD regulates acute lung injury in lethal Streptococcus pneumoniae infections. Immunity 27:349-60.
7. Inoshima I, Inoshima N, Wilke GA, Powers ME, Frank KM, Wang Y, Bubeck Wardenburg J. 2011. A Staphylococcus aureus pore-forming toxin subverts the activity of ADAM10 to cause lethal infection in mice. Nat Med 17:1310-4.
8. Ratner AJ, Aguilar JL, Shchepetov M, Lysenko ES, Weiser JN. 2007. Nod1 mediates cytoplasmic sensing of combinations of extracellular bacteria. Cell Microbiol 9:1343-51.
9. Collier RJ. 2009. Membrane translocation by anthrax toxin. Mol Aspects Med 30:413-22.
10. Los FC, Randis TM, Aroian RV, Ratner AJ. 2013. Role of pore-forming toxins in bacterial infectious diseases. Microbiol Mol Biol Rev 77:173-207.
11. Tweten RK. 2005. Cholesterol-dependent cytolysins, a family of versatile pore-forming toxins. Infect Immun 73:6199-209.
12. Hotze EM, Le HM, Sieber JR, Bruxvoort C, McInerney MJ, Tweten RK. 2013. Identification and characterization of the first cholesterol-dependent cytolysins from Gram-negative bacteria. Infect Immun 81:216-25.
13. Olofsson A, Hebert H, Thelestam M. 1993. The projection structure of perfringolysin O (Clostridium perfringens theta-toxin). FEBS Lett 319:125-7.
14. Bhakdi S, Tranum-Jensen J, Sziegoleit A. 1985. Mechanism of membrane damage by streptolysin-O. Infect Immun 47:52-60.
15. Rossjohn J, Feil SC, McKinstry WJ, Tweten RK, Parker MW. 1997. Structure of a cholesterol-binding, thiol-activated cytolysin and a model of its membrane form. Cell 89:685-92.
16. Polekhina G, Giddings KS, Tweten RK, Parker MW. 2005. Insights into the action of the superfamily of cholesterol-dependent cytolysins from studies of intermedilysin. Proc Natl Acad Sci U S A 102:600-5.
17. Wickham SE, Hotze EM, Farrand AJ, Polekhina G, Nero TL, Tomlinson S, Parker MW, Tweten RK. 2011. Mapping the intermedilysin-human CD59 receptor interface reveals a

deep correspondence with the binding site on CD59 for complement binding proteins
C8alpha and C9. J Biol Chem 286:20952-62.

18. Lawrence SL, Gorman MA, Feil SC, Mulhern TD, Kuiper MJ, Ratner AJ, Tweten RK,
Morton CJ, Parker MW. 2016. Structural Basis for Receptor Recognition by the Human
CD59-Responsive Cholesterol-Dependent Cytolysins. Structure 24:1488-98.

19. Shatursky O, Heuck AP, Shepard LA, Rossjohn J, Parker MW, Johnson AE, Tweten RK.
1999. The mechanism of membrane insertion for a cholesterol-dependent cytolysin: a
novel paradigm for pore-forming toxins. Cell 99:293-9.

20. Shepard LA, Shatursky O, Johnson AE, Tweten RK. 2000. The mechanism of pore
assembly for a cholesterol-dependent cytolysin: formation of a large prepore complex
precedes the insertion of the transmembrane beta-hairpins. Biochemistry 39:10284-93.

21. Czajkowsky DM, Hotze EM, Shao Z, Tweten RK. 2004. Vertical collapse of a cytolysin
prepore moves its transmembrane beta-hairpins to the membrane. EMBO J 23:3206-15.

22. Giddings KS, Zhao J, Sims PJ, Tweten RK. 2004. Human CD59 is a receptor for the
cholesterol-dependent cytolysin intermedilysin. Nat Struct Mol Biol 11:1173-8.

23. Gelber SE, Aguilar JL, Lewis KL, Ratner AJ. 2008. Functional and phylogenetic
characterization of Vaginolysin, the human-specific cytolysin from Gardnerella vaginalis.
J Bacteriol 190:3896-903.

24. Tabata A, Matsumoto A, Fujimoto A, Ohkura K, Ikeda T, Oda H, Yokohata S, Kobayashi
M, Tomoyasu T, Takao A, Ohkuni H, Nagamune H. 2022. Dual functions of
discoidinolysin, a cholesterol-dependent cytolysin with N-terminal discoidin domain
produced from Streptococcus mitis strain Nm-76. J Oral Microbiol 14:2105013.

25. Zbinden A, Mueller NJ, Tarr PE, Eich G, Schulthess B, Bahlmann AS, Keller PM,
Bloemberg GV. 2012. Streptococcus tigurinus, a novel member of the Streptococcus
mitis group, causes invasive infections. J Clin Microbiol 50:2969-73.

26. Zbinden A, Mueller NJ, Tarr PE, Sproer C, Keller PM, Bloemberg GV. 2012. Streptococcus
tigurinus sp. nov., isolated from blood of patients with endocarditis, meningitis and
spondylodiscitis. Int J Syst Evol Microbiol 62:2941-2945.

27. Bourassa L, Clarridge JE, 3rd. 2015. Clinical Significance and Characterization of
Streptococcus tigurinus Isolates in an Adult Population. J Clin Microbiol 53:3574-9.

28. Kanamori H, Kakuta R, Yano H, Suzuki T, Gu Y, Oe C, Inomata S, Aoyagi T, Hatta M, Endo
S, Tokuda K, Weber DJ, Nakamura Y, Saiki Y, Kaku M. 2015. A case of culture-negative
endocarditis due to Streptococcus tigurinus. J Infect Chemother 21:138-40.

29. Diene SM, Francois P, Zbinden A, Entenza JM, Resch G. 2016. Comparative Genomics
Analysis of Streptococcus tigurinus Strains Identifies Genetic Elements Specifically and
Uniquely Present in Highly Virulent Strains. PLoS One 11:e0160554.

30. Johnson S, Brooks NJ, Smith RA, Lea SM, Bubeck D. 2013. Structural basis for recognition
of the pore-forming toxin intermedilysin by human complement receptor CD59. Cell Rep
3:1369-77.

31. Shahi I, Llaneras CN, Perelman SS, Torres VJ, Ratner AJ. 2022. Genome-Wide CRISPR-
Cas9 Screen Does Not Identify Host Factors Modulating Streptococcus agalactiae beta-
Hemolysin/Cytolysin-Induced Cell Death. Microbiol Spectr 10:e0218621.

32. Ramachandran R, Heuck AP, Tweten RK, Johnson AE. 2002. Structural insights into the
membrane-anchoring mechanism of a cholesterol-dependent cytolysin. Nat Struct Biol
9:823-7.
33. Soltani CE, Hotze EM, Johnson AE, Tweten RK. 2007. Structural elements of the
cholesterol-dependent cytolysins that are responsible for their cholesterol-sensitive
membrane interactions. Proc Natl Acad Sci U S A 104:20226-31.
34. Giddings KS, Johnson AE, Tweten RK. 2003. Redefining cholesterol's role in the
mechanism of the cholesterol-dependent cytolysins. Proc Natl Acad Sci U S A
100:11315-20.
35. Soltani CE, Hotze EM, Johnson AE, Tweten RK. 2007. Specific protein-membrane
contacts are required for prepore and pore assembly by a cholesterol-dependent
cytolysin. J Biol Chem 282:15709-16.
36. Zilnyte M, Venclovas C, Zvirbliene A, Pleckaityte M. 2015. The cytolytic activity of
vaginolysin strictly depends on cholesterol and is potentiated by human CD59. Toxins
(Basel) 7:110-28.
37. Tabata A, Ohkura K, Ohkubo Y, Tomoyasu T, Ohkuni H, Whiley RA, Nagamune H. 2014.
The diversity of receptor recognition in cholesterol-dependent cytolysins. Microbiol
Immunol 58:155-71.
38. Rampersaud R, Planet PJ, Randis TM, Kulkarni R, Aguilar JL, Lehrer RI, Ratner AJ. 2011.
Inerolysin, a cholesterol-dependent cytolysin produced by Lactobacillus iners. J Bacteriol
193:1034-41.
39. Tomoyasu T, Matsumoto A, Takao A, Tabata A, Nagamune H. 2023. A simple method to
differentiate three classes of cholesterol-dependent cytolysins. J Microbiol Methods
207:106696.
40. Boyd CM, Parsons ES, Smith RA, Seddon JM, Ces O, Bubeck D. 2016. Disentangling the
roles of cholesterol and CD59 in intermedilysin pore formation. Sci Rep 6:38446.
41. Jost BH, Lucas EA, Billington SJ, Ratner AJ, McGee DJ. 2011. Arcanolysin is a cholesterol-
dependent cytolysin of the human pathogen Arcanobacterium haemolyticum. BMC
Microbiol 11:239.
42. Shewell LK, Day CJ, Jen FE, Haselhorst T, Atack JM, Reijneveld JF, Everest-Dass A, James
DBA, Boguslawski KM, Brouwer S, Gillen CM, Luo Z, Kobe B, Nizet V, von Itzstein M,
Walker MJ, Paton AW, Paton JC, Torres VJ, Jennings MP. 2020. All major cholesterol-
dependent cytolysins use glycans as cellular receptors. Sci Adv 6:eaaz4926.
43. Shewell LK, Harvey RM, Higgins MA, Day CJ, Hartley-Tassell LE, Chen AY, Gillen CM,
James DB, Alonzo F, 3rd, Torres VJ, Walker MJ, Paton AW, Paton JC, Jennings MP. 2014.
The cholesterol-dependent cytolysins pneumolysin and streptolysin O require binding to
red blood cell glycans for hemolytic activity. Proc Natl Acad Sci U S A 111:E5312-20.
44. Drabavicius G, Daelemans D. 2021. Intermedilysin cytolytic activity depends on heparan
sulfates and membrane composition. PLoS Genet 17:e1009387.
45. Johnstone BA, Joseph R, Christie MP, Morton CJ, McGuiness C, Walsh JC, Bocking T,
Tweten RK, Parker MW. 2022. Cholesterol-dependent cytolysins: The outstanding
questions. IUBMB Life 74:1169-1179.
46. Sanjana NE, Shalem O, Zhang F. 2014. Improved vectors and genome-wide libraries for
CRISPR screening. Nat Methods 11:783-784.

768
769
770
<u>Figure S1: Cholesterol depletion does not affect hCD59 expression on HeLa cell surface</u>
Flow cytometry analysis of hCD59 expression on the cell surfaces of HeLa cells (NT control,
hCD59 KO, and cholesterol depleted) using an anti-hCD59 primary antibody and an Alexa-Fluor
647 (AF647) secondary antibody. Histogram x-axis shows APC fluorescence intensity of the
secondary antibody for each cell population. Histogram y-axis shows cell count normalized to
mode.
777

778
779
780
781 **<u>Figure S2: Functional TGY variants lyse JEG-3 cells in a hCD59-dependent manner</u>**
782 (**A-B**) Percentage cell death in JEG-3 NT control cells and JEG-3 hCD59 KO cells when exposed to
783 increasing concentrations of the three TGY variants. Cells were incubated with toxins for 1.5
784 hours and cell death measured by an LDH-release cytotoxicity assay. Each point is the mean of 3
785 replicates, and error bars represent ±SD.

786
787
Figure S3: Visualizing CDC oligomers for ILY, VLY and PLY on HeLa cells
(**A-C**) Oligomerization of CDCs as assessed by SDS-AGE analysis. All CDCs were used at a
concentration of 5 μg/ml, unless indicated otherwise. Cells were incubated with the indicated
toxins on ice for 10 minutes before being prepared for western blotting. 0.01% glutaraldehyde
was used as a cross-linker to preserve oligomers during sample preparation. Western blot
membrane was probed with an anti-His-tag HRP-conjugated antibody.
794

	Primer name	Sequence (5'-3')
1.	PLY_D1-3_F	CTGGTGCCGCGCGGCAGCCATATGGCAAATAAAGCAGTAAATG
2.	PLY_D1-3_R_tgy	CGTTTTTATAAGCTGTAACCTTAGTCTC
3.	TGY_D4_F_ply	GGTTACAGCTTATAAAAACGGCTACCTG
4.	TGY_D4_R	AGTGGTGGTGGTGGTGGTGCTTAGTTGTTTTCAATTTCTTCG
5.	TGY_E454G_F	CACAAGGGTGGTTATGTGGCGC
6.	TGY_E454G_R	ATGCAGATTCAGGTAGCC
7.	TGY_E454A_F	CACAAGGGTGCATATGTGGCG
8.	TGY_E454A_R	ATGCAGATTCAGGTAGCC
9.	TGY_I544L_F	CGGCACCACCCTGCGCCCGAAAT
10.	TGY_I544L_R	TAGTTGGTGATGGTGCGTTTTTG
11.	T7_F	TAATACGACTCACTATAGGG
12.	T7_Term	GCTAGTTATTGCTCAGCGG

Table S1: Primers and oligonucleotides

Primers and oligonucleotides used for construction of hybrids (primers 1-4), site-directed
mutagenesis (primers 5-10), and sequencing reactions (primers 11-12).