1

Spatial and temporal distribution of ribosomes in single cells reveals aging

differences between old and new daughters of *Escherichia coli*

Lin Chao[#*], Chun Kuen Chan[*], Chao Shi, and Camilla U. Rang

Department of Ecology, Behavior and Evolution, School of Biological Sciences,

University of California San Diego, La Jolla, CA, USA

*Co-first authors.

Corresponding author: lchao@ucsd.edu

Running title: Ribosome distribution in aging *E. coli* single cells.

ABSTRACT Lineages of rod-shaped bacteria such as *Escherichia coli* exhibit a

temporal decline in elongation rate in a manner comparable to cellular or biological

aging. The effect results from the production of asymmetrical daughters, one with a

lower elongation rate, by the division of a mother cell. The slower daughter compared

to the faster daughter, denoted respectively as the old and new daughters, has more

aggregates of damaged proteins and fewer expressed gene products. We have

examined further the degree of asymmetry by measuring the density of ribosomes

between old and new daughters and between their poles. We found that ribosomes

were denser in the new daughter and also in the new pole of the daughters. These

ribosome patterns match the ones we previously found for expressed gene products.

This outcome suggests that the asymmetry is not likely to result from properties unique

to the gene expressed in our previous study, but rather from a more fundamental

upstream process affecting distribution of ribosomal abundance. Because damage

aggregates and ribosomes are both more abundant at the poles of *E. coli* cells, we

suggest that competition for space between the two could explain the reduced

ribosomal density in old daughters. Using published values for aggregate sizes and the

relationship between ribosomal number and elongation rates, we show that the

aggregate volumes could in principle displace quantitatively the amount of ribosomes

needed to reduce the elongation rate of the old daughters.

IMPORTANCE Bacteria exhibit a growth decline in a manner comparable to cellular or

biological aging. When a mother bacterium reproduces by binary fission it allocates

more damage to one of the two daughters. The extra damage correlates with a slower

growth. Thus, a lineage of daughters successively acquiring more damage over

generations ages, sometimes even to death under stressful conditions. Aging lineages

also have lower levels of expressed gene products. Here we show that the aging

process also correlates with lower cellular levels of ribosomes. The identification of a

ribosomal effect shows that the aging process is acting at a much more fundamental

upstream level. While decreased gene products could have resulted from local

regulation of specific genes, a lower ribosomal density affects the entirety of cellular

metabolism. Understanding bacterial aging is important because biological aging may

have originated in single-celled organisms such as *E. coli.*

INTRODUCTION

A mother *Escherichia coli* cell divides asymmetrically into two daughter cells that have different elongation rates (1–6). The new daughter elongates faster while the old daughter is slower. The asymmetry and old-new designation of the daughters results from the fact that the rod-shaped mother divides by forming new poles at its midplane. As a result, all *E. coli* cells are polarized at birth with a new and an old pole (Fig. 1A). When a polarized cell in turn becomes a mother and divides, its daughters will also be polarized but they can also be identified as new and old daughters depending on whether they acquired the mother's new or old pole. Note that mothers can also be denoted as new or old, depending on whether they were new or old daughters at birth. The difference in elongation rate between old and new daughters was found by Lindner et al. (7) to correlate negatively with the amount of damaged proteins (inclusion bodies) in individual cells. They also found that protein aggregates were more often associated with the old pole of the mother, in which case damaged proteins are partitioned asymmetrically by the mother to its daughters. However, a study by Govers et al. (8) has reported a weaker and non-significant negative correlation. Because the two studies used different external stressors (to induce damage) and fluorescent reporters, they are not fully comparable and the relationship between elongation rate and the damaged proteins in *E. coli* remains unresolved (see Discussion for additional details). Despite the differences between Lindner et al. and Govers et al., protein aggregates often have a negative qualitative effect on cell health and viability (9–12), and more so in old daughters (13, 14).

The asymmetry of both the elongation rate difference between old and new

daughters and the damage partitioning by the mother in *E. coli* has evolutionary

consequences. The asymmetry creates fitness variance between daughters that

increases the effectiveness of natural selection (2, 15), and an asymmetrical lineage

has more progeny over time relative to a symmetrical one. An analogy comes from

economic models. Two asymmetrical $500 bank accounts at 10% and 6% yield more

money over time than two symmetrical $500 accounts at 8%. However, the asymmetry

comes with costs. While the higher elongation rate of the new daughter rises in the

lineage over generations, the rate of the old daughter declines (1, 2). Under more

benign culture conditions, the rise and the decline can achieve stable values and the

lineages are immortal (2, 4, 6, 16, 17). Under more stressful and damage inducing

conditions, the old daughter lineage can die because its elongation rate declines to zero

and the death probability is dosage dependent (4, 5). The bacterial population survives

despite the death of the old daughter lineage because the new daughter lineage is still

able to achieve a stable rate that is greater than zero (2, 4, 5). Because the elongation

rate decline and death of the old daughter lineage is a form of physiological aging,

bacteria can be used to investigate the evolutionary origins of biological aging. Despite

that a quantitative correlation between the amount of damaged proteins and elongation

rates remains unresolved (see above), the asymmetries here described for elongation

rates and death strongly suggest that damage stress in *E. coli* may play a key, if not

causal, role in cell aging. We note that the asymmetrical partitioning of damage is not

an alternative to protein repair, or vice-versa. Protein repair (including reprocessing) is

occurring (18, 19) and its effect is to lower the damage rate experienced by the cell.

However, if damage aggregates are present in cells, the damage rate must necessarily

exceed the repair rate. There is likely a cost to repair, in which case the repair rate may

become limited by diminishing returns. Thus, the asymmetry could be an evolutionary

adaptation for dealing with a low repair rate. If repair rates were sufficiently high to

negate damage, *E. coli* should not have evolved asymmetry.

To develop further our understanding bacterial aging, our previous study

examined the distribution of expressed native (not fused to another protein) GFP

molecules in *E. coli* (3). We found that the GFP density was higher in new poles than

in old ones and in cells with a faster elongation rate. To take the analysis one step

further, we present here results from our examination of the distribution of ribosomes in

the poles and daughters of single *E. coli* cells. We used the *yfp* (yellow fluorescent

protein) and S2 subunit (*rpsB* gene) fusion by Bakshi et al. (20) as a reporter for

ribosomal density. The S2-YFP construct is an effective fluorescent reporter. The S2

ribosomal subunit makes a desirable reporting partner because free subunits are

negligible and nearly all cellular S2-YFP is incorporated into ribosomes (20, 21). Almost

all of the detected fluorescent S2-YFP comes from copies are already incorporated in

ribosomes (20). Bakshi et al. (20) used S2-YFP to successfully make a more detailed

map of the nucleoid/ribosome segregation. Most importantly, because fluorescent

reporters can produce artifacts (8, 22, 23), YFP and S2-YFP were extensively tested to

validate that the reported fluorescence was not artifactual. These tests are detailed

next.

The tests validating YFP and S2-YFP followed the methods of Landgraf et al. (23)

by which validation requires that either fluorescent aggregation, clusters or foci are

reported when non-uniformity is expected by an assay or alternatively not reported

when uniformity is expected. An artifact occurs when uniformity is reported when non-

uniformity is expected (and vice versa). Because reporting artifacts depend on the

protein to which a fluorescent reporter is fused (23), a reporter that makes artifacts with

one protein does not necessarily make it with a second. Thus, many of the tests were

targeted specifically to the S2-YFP construct. (i) A first test focused on native YFP (not

fused to any protein) because an earlier study suggested that YFP produced artifactual

non-uniformity due to self-aggregation (8). This test was needed because the self-

aggregation results were based on a fusion construct of the chaperone IbpA and YFP.

However, was the self-aggregation a property of YFP or an interaction with IbpA?

Bakshi et al. (20) found that native YFP reported a uniform distribution, in which case

YFP is not inherently prone to self-aggregation artifacts (see also IbpA-YFP self-

aggregation in Discussion). (ii) If YFP is fused to OmpR, which is a regulator that binds

to the OmpC and OmpF loci, and the three loci (OmpR-YFP, OmpC and OmpF) are

expressed from the bacterial chromosome (hereafter strain A), the fluorescence

distribution is uniform (24). Because the chromosome has a low copy number, most

OmpR-YFP is distributed over the entire cell. Some OmpR-YFP is bound to OmpC and

OmpF but they are too few to produce a fluorescence focus. If OmpC and OmpF are

cloned into the multi-copy plasmid pBR322, which forms clusters in cells (25), and the

plasmid is inserted into strain A (making strain B), a fluorescence focus reporting the

plasmid cluster emerges (24). Additionally, if native OmpR (not fused to YFP or any

fluorescent reporter) is over-expressed from a second plasmid that is inserted into strain

B (making strain C), the fluorescence focus disappears because the more numerous

native OmpR outcompetes OmpR-YFP in binding to the pBR322/OmpC-OmpF cluster.

Thus, OmpR-YFP is responding correctly when either uniformity or non-uniformity is

expected. (iii) To confirm that the nucleoid/ribosome segregation they observed was

not due to YFP reporting artifacts, Bakshi et al. repeated their experiments by adding

rifampicin and chloramphenicol, two antibiotics that affect the distribution of the nucleoid

DNA (26). Whereas the nucleoid is expanded and rendered more uniform by rifampicin,

it is compacted and clumped by chloramphenicol. Bakshi et al. found that the

fluorescence reported by S2-YFP became more uniform with rifampicin but more

clumped and non-uniform with chloramphenicol (20). As the nucleoid expanded with

rifampicin, the ribosome distribution also expanded as it filled the spaces created by the

nucleoid expansion. On the other hand, as the nucleoid contracted with

chloramphenicol, the ribosomes were excluded from the space occupied by the more

clumped nucleoid and they in turn became less uniform. In summary, these YFP,

OmpR-, and S2- controls gives us confidence that S2-YFP is a reliable reporter for

ribosomes and that reported fluorescence foci are better explained by the non-uniform

properties of the cell rather than any inherent property or artifact of YFP. We find these

tests that functionally validate S2-YFP to be more convincing than examining S2 fused

with more fluorescent reporters. If two reporters disagree, which one is the artifact? If

they agree, are they both truthful or artifactual?

Our examination of the distribution of ribosomes with S2-YFP could be

particularly instructive. First, while Bakshi et al. have shown that ribosomes congregate

in the poles, it is not known if ribosomal densities differ between old and new daughters

or their poles. Second, our previous report of more expressed gene products in new

poles and daughters (3) could have resulted from regulatory factors acting on a specific

gene or at a higher upstream level such as ribosomal abundance. In the latter case, the

consequences would be much more fundamental as they could affect the entire

proteome. The fact that ribosomal density and elongation rates are highly correlated in

bacteria (27–31) additionally reinforces the importance of knowing the distribution of

ribosomes in the poles and daughters of *E. coli*. Lastly, given that ribosomes occupy

nearly 40% of the cytoplasmic volume of an *E. coli* cell (32) and that both ribosomes

and damage aggregates tend to reside in the poles, the possibility that the two could be

excluding each other at the poles becomes a probability. However, a space limiting

model as a hypothesis for explaining bacterial aging requires first that the distribution of

ribosomes in old and new poles or daughters be characterized. Here we present the

results of our study quantifying the temporal and spatial distribution of ribosomes in

single *E. coli* cells by time-lapse microscopy.

RESULTS

Following Materials and Methods, the distribution of ribosomes was measured in

the *E. coli* strain AFS55 with a fluorescent *yfp* reporter fused to the ribosomal subunit

S25 (20). Fluorescence levels were quantified from time-lapse images of single cells

cultured on agarose pads. Because the level recorded for a cell on an image is affected

by the light that the cell loses by diffraction and gains by diffraction from neighboring

cells, the diffractional scatter of all images was corrected by deconvolution. Red and

blue colors in figures represent old and new daughters or mothers, respectively. Ratio

data (Fig. 1C) were derived from estimates that were not normalized. All other data for

cell length, ribosomal density, and time were normalized when pooling and comparing

cells. All error bars are ± SEM. and statistical details are presented in figure legends.

We first examined the density of ribosomes in colonies starting from a single cell.

A quick visual inspection of the fluorescent images and their heat maps for a few

representative cells showed clearly that the ribosomal densities tended to be highest

near the poles (Fig. 1B). However, high densities also built up in the mid region of a

mother cell right before division. As the mid region leads to the formation of the new

poles of the daughters, the buildup explains how those poles acquired their higher

levels. These polar concentration of ribosomal densities matched the previous ones

reported by Bakshi et al (20). To obtain a more quantitative analysis of the changes in

ribosomal densities during cell growth, we collected a larger sample size and compared

measurements between old and new mothers, old and new daughters, between their

poles, along the length of their cells, and during their elongation from birth to division.

New daughters at birth from old mothers have more ribosomes. A first

comparison revealed that if a pair of old and new daughters descending from the same

mother were compared, ribosome density measured by the fluorescence was higher for

new daughters. The ratio of reported fluorescence between new/old daughters (the

daughter ratio) was 1.08±0.012, which was significantly greater than 1.00 (p = 4.38×10^{-}

^{6}). However, if the mothers were split into pools of old and new mothers (Fig. 2A), the

daughter ratio from old mothers was 1.11 ± 0.018 and significantly greater than 1.00 (p

= 6.89×10-6) and the ratio from new mothers was 1.04 ± 0.015 and barely significant (p

= 0.04). Thus, the higher ribosome density in new daughters resulted primarily from old

mothers. The difference between old and new mothers was further demonstrated by a

strong significant difference between their daughter ratios (1.11 vs.1.04; p=0.007).

Thus, new daughters have more ribosomes relative to old daughters produced by the

same mother, but asymmetry is higher in pairs from old mothers. This higher density of

ribosomes in new daughters correlates qualitatively with our previously reported higher

levels of expressed gene products and elongation rates in new daughters (2–5).

Ribosomal asymmetry between daughters are spatially in place in mothers

before division. To further explore our find of a ribosomal asymmetry between new

and old daughters, we examined the intracellular distribution of ribosomal fluorescence

in the new and old pole halves (see diagram in Fig. 1C) of old and new mothers right

before division. The two pole halves were transformed into a ratio (new/old) for analysis

(Fig. 2B). We found that pole ratio in old mothers was significantly greater than 1.00

$(1.10 \pm 0.015; p = 6.23 \times 10^{-5})$. The pole ratio in new mothers was much lower $(1.03 \pm$

$0.010)$ and not significantly greater than 1.00 (p = 0.08). The pole ratios between new

and old mothers were significantly different from each other $(p= 1.3 \times 10^{-4})$. Thus, the

density of ribosomes in the old and new poles of mothers matches well the density in

the old and new daughters of both old and new mothers (cf. Figs. 2A, 2B). This match

argues that the physical placement of ribosomes in the mother at the time of division

sets in place the ribosome distribution in the daughters. To test this argument, we

compared directly the ribosomal pole ratio of an individual mother against the ribosomal

daughter ratio of the same mother. No significant difference was found, in both old and

new mothers, when the ribosomal ratio of the poles and daughters was compared. With

new mothers, the pole and daughter ratios were respectively 1.03 versus 1.04 and not

different (cf. Figs 2A, 2B; p=0.27). With old mothers the respective ratios were 1.10

versus 1.1 and also not different (cf. Figs 2A, 2B; p=0.62). Thus, as the new and old

pole halves yield directly the new and old daughters upon the division of a mother, the

placement of ribosomes in the mother cell sets up the quantity, and hence asymmetry,

of ribosomes in the new and old daughters of a mother.

Stochastic and deterministic components of ribosomal variance in

daughter cells. Although new and old daughters can differ significantly in ribosome

density, there was also substantial variation within each daughter population. A plot of

the distributions shows clearly the within population variation, although the difference

between old and new daughters is also apparent (Figs. 3A, 3B). Old and new

daughters form their own distributions, but their mean values are displaced by a

difference D, which is larger when the daughters are derived from old mothers. The

variation within the old and new daughter distributions is presumably due to stochastic

noise, although the stochasticity could also result from an unknown deterministic source.

The latter is well illustrated by discovery that *E. coli* mothers produce asymmetrical

daughters. For example, without knowledge about the asymmetry, old and new

daughters would have been pooled into one population with a total variance V_T (Figs.

3A, 3B; dashed lines), which would have been interpreted to result from stochasticity.

In reality, V_T would actually have a deterministic component due to the difference

between old and new daughters. As we have shown for expressed gene products (3),

$V_T = (V_{Old} + V_{New})/2 + D^2/4$, when there is deterministic asymmetry (D>0) and V_{Old} and

V_{New} are the variances of the unpooled old and new daughter populations (Figs. 3A, 3B;

red and blue lines). If D=0, V_T is the average of V_{Old} and V_{New}. Thus, $D^2/4$ represents

the deterministic component of V_T, and the contribution of deterministic asymmetry

expressed as a percentage is $h^2 = (D^2/4) / V_T$. Our estimates of D showed that h^2 was

39.9% from old mothers and 2.4% from new mothers (Fig. 4A). The higher h^2 for

daughters from old mothers is consistent with our results in Figs. 3A and 3B that show

old mothers having more different daughters than new mothers. These results were

also consistent with our previous report that h^2 is higher for expressed gene products in

daughters of old than of new mothers (h^2 of 40.1 and 10.1%) (3). Thus, a claim of

stochasticity to explain the variance of ribosomal density of *E. coli* cells could have

introduced an error of nearly 40% without knowledge of the asymmetry between old and

new daughters. Without knowing more, we presume that the variances V_{Old} and V_{New}

result from stochasticity, but it may also be that a deterministic process still lies hidden.

268   **Asymmetry and variance components for elongation rates in daughter cells.**

To determine whether the observed ribosome patterns reported in Figs. 3A and 3B had

a downstream effect, we measured elongation rates of the same daughters and

mothers. When new and old daughters were pooled irrespective of whether they

originated from old or new mothers, new daughters were found to elongate faster than

old daughters (new/old ratio of 1.09 ± 0.011 ($p = 3.9 \times 10^{-16}$). When the old and new

daughters were separated as descending from old or new mothers, the ratio was

respectively 1.11 ± 0.017 ($p = 1.3 \times 10^{-10}$) and 1.07 ± 0.011 ($p = 2.8 \times 10^{-7}$) (Fig. 2C).

Although both ratios were significantly greater than 1.00, the asymmetry was greater

between daughters from old mothers and the two ratios were significantly different ($p =$

0.02). As we did for the ribosome data, the variance of elongation rate measurements

was also portioned into a deterministic and stochastic component to estimate the

percentage h^2 (Figs. 3C, 3D, 4B). The deterministic percentage of the variance was

found to be h^2 = 18.5% and 12.2% for daughters originating from old and new mothers,

respectively. As we observed for ribosome densities, the h^2 for old mothers was still

higher, but not to the same magnitude (see Discussion). The overall similarity between

the ribosome and elongation rate data reflects the fact that the two phenotypes are well

known to be positively correlated in *E. coli* (27–31). An examination of the relationship

between our measurements for ribosome and elongation yielded similar outcomes (Fig.

5). Daughters from both old and new mothers had ribosome densities that correlated

positively with elongation rates (r = 0.387, $p < 1 \times 10^{-5}$ and r = 0.233, $p = 4 \times 10^{-4}$,

respectively; see Fig. 5 for more details). The correlation from new mothers was

weaker because its deterministic component h^2 was smaller (Fig. 4B), in which case the

distributions of new and old daughters were less separated (cf. Figs. 5A and 5B).

Spatial distribution of ribosomes in mother cells from birth to division. Our

observation that the asymmetry of the pole halves of mothers determines the

distribution of ribosomes of daughters at birth (Figs. 2A, 2B) raises the question of how

the spatial distribution of ribosomes changes temporally in the mother cell, and whether

the changes provide insights or hypotheses for the origin of the asymmetry. Thus, we

increased the resolution of the process by sampling over a finer spatial (lengthwise) and

temporal (birth to division) grain of the ribosomal density in a mother cell. The length of

the mother cell was divided into four quartiles, which were termed NP (new pole), L2, L3,

and OP (old pole) (Fig. 1D). The time from birth to division of the mother cell was also

divided into four quartiles termed Birth (B), T2, T3, and Division (D). Ribosome density

and the quartiles of length and time were all normalized for pooling the cells. Note that

because ribosome densities are normalized, they indicate relative rather than absolute

amounts and the former can change even when the latter is constant.

We first examined whether the new length quartile results were consistent with

our previous ribosome pole ratios (Fig. 2B) for the new and old pole halves in the

mothers. These new ribosome pole ratio data were obtained by taking the ratio of the

length quartiles (NP+L2) / (L3+OP) and plotting the values over the time quartiles (Fig.

6). The ratios increased from birth to division in both old and new mothers, but increase

was significant only for the old mothers (p=0.026 vs. 0.59, respectively). The rapid rise

in the old mothers also began from a higher value at birth. More importantly, the final

ratio values at quartile D (1.11 and 1.02 for old and new mothers, respectively) were

comparable to Fig. 2C (1.11 and 1.04), which was also timed to division. Thus, old

mothers magnified the asymmetry of their ribosome distribution over time to achieve the

higher ratio that we used above to explain the larger asymmetry of their daughters. On

the other hand, the new mothers begun with a pole ratio close to 1.02 and did not

change over time, which explained the higher symmetry of their daughters.

By using the length into quartiles NP, L2, L3, and OP as individual

measurements, we were able to examine in more detail the trajectories for the changes

of ribosome densities of mothers from birth to division (Fig. 7A). Several key patterns

emerged. First, the trajectories decreased in NP and OP and increased in L2 and L3 in

both old and new mothers. Moreover, the density at the final time quartile D was L2>NP

and L3>OP. Because L2 and L3 are the new poles of the daughters after division (cf.

Fig. 1A and 7A), the daughters were already being provisioned by their mothers to be

biased to have more ribosomes at their future new poles at the expense of their future

old poles. Second, the trajectories of new mothers in NP-L2 were mirrored by L3-OP

(Fig. 7A; blue lines), i.e. NP was more similar to OP, and L2 more similar to L3. The

trajectory slopes of NP vs. OP (-0.00520 ± 0.00313 vs. -0.00809 ± 0.00340; p > .05)

and L2 vs. L3 (0.00675 ± 0.00200 vs. 0.00655 ± 0.00198; p > .05) were not significantly

different (Fig. 7B). Thus, ribosomes were being redistributed from NP and OP to L2 and

L3 in an equal manner and the lower asymmetry of new mothers was being maintained

(Fig. 2B). Third, the trajectories for the old mothers had a higher asymmetrical

signature (Fig. 7B; red lines). They resided above the new mother trajectories in NP

and declined until they were below in OP. Moreover, the trajectory slopes of NP vs. OP

(-0.00390 ± 0.00313 vs. -0.00803 ± 0.00347; p<.05) and L2 vs. L3 (0.00788 ± 0.00220

vs. 0.00405 ± 0.00217; p<.05) were now significantly different (Fig. 7B). Thus, relative

to L3 and OP, NP and L2 not only began at birth with more ribosomes, but also NP lost

fewer and L2 acquired more ribosomes over time. As a result, the old mothers became

even more asymmetric than new mothers at division.

DISCUSSION

The discovery that the division of mother *E. coli* cells produces two asymmetric

daughters that have different elongation rates was ground breaking (1). The old

daughter receiving the maternal old pole (Fig. 1A) elongates more slowly than the new

daughter receiving the maternal new pole. The difference is not due to genetic

mutations because the daughter with the slower rate can in turn produce a grand-

daughter with the elevated rate (1, 2, 4, 5, 16). These results have been replicated on

agar surfaces(1, 3, 12, 17) and in liquid media (4–6, 16), although the tracking of cells

by microscopy requires the cells to be a monolayer in both methods. Population growth

rates during microscopy are comparable to those in shaking liquid cultures (4–7, 33).

The causal trigger for the asymmetry between the daughters remains unknown,

although damaged proteins (7), expressed gene products (GFP), and now ribosomes

(Fig. 6; see below) have been shown to correlate with the differences in elongation rate.

Damage may have the strongest causal link because the addition of external damage

agents such as heat, phototoxicity, and antibiotics increases differentially the death

probability of old daughters (4, 5). Because new daughters are born with less damage,

they are not killed by the damage agents and allow for the survival of the population (2,

5). The elongation rate differences have an epigenetic effect because slow mothers

produce slower daughter pairs, although the new daughter is still quicker within the pair

(2, 4, 5, 16). If the causal factor were damaged proteins, the epigenetic interpretation is

that the slower mother has more damage than average, and therefore the old and new

daughter receive also more than their respective averages but the asymmetry is

preserved because the former still gets more than the latter.

Because old daughters are more vulnerable to cell death than new daughters

(see above), the elongation rate asymmetry can be considered a form biological aging

(1). Thus, *E. coli* is a model for the evolutionary origin of biological aging from single-

celled organisms. We would extend the argument to suggest that the old daughters are

more properly considered to be the continuation of mother, and the new daughters are

the true daughters. Given an epigenetic factor that slows the elongation rate of a cell,

the mother retains a larger pool during division and it ages. The daughter is allocated

less and it is rejuvenated. The larger pool that the mother does not transmit to the

372 daughter is the original soma. Aging is the price that a mother pays to produce healthy

373 daughters. Because the elongation rate asymmetry between mothers and daughters

374 maximizes the fitness of the lineage (see Introduction), aging is indirectly favored by

375 natural selection (2, 15). Because the origin of life required organization of genetic

376 material into protocells (34) in hazardous damage-inducing conditions (35, 36),

377 biological aging may have evolved in the first life forms.

378   The higher vulnerability of old daughters (or old mothers) in the presence of

379 added damage agents was quantified by estimating the mean a and the variance s^2 of

380 the regression (or mapping) coefficient of the doubling time (DT) of daughters onto

381 mothers (4). The vulnerability is high if $(a^2 + s^2) < 1$ because the daughter could then

382 acquire a very long DT and fail to divide. For example, if $a = 1.1$, a mother with DT = 40

383 min would have a daughter with a longer DT = 1.1*40 min, and the DT each subsequent

384 daughter in the lineage is even longer. If s^2 is large, then by chance the DT of a

385 daughter could also be vulnerably long. When damage agents were added, the old

386 daughters were the first ones to die because their values of a and s^2 increased above

387 the threshold (5). Thus, the asymmetry between old and new daughters is extended to

388 include the values of a and s^2.

389   A question that is often raised is why evolution by natural selection is not able to

390 minimize a and s^2 to diminish the window of vulnerability. An answer is that it does, but

391 the damage agents are pathological conditions beyond the physiological limits. We

392 suggest an additional answer for s^2 based on our variance measurements for ribosome

393 density and elongation rate (Fig. 4). From those measurements, we estimated the

394 stochastic and deterministic components of the variances. The deterministic

component, which resulted from the old and new daughter asymmetry, was $D^2/4$. The

stochastic component is therefore $V_E = V_T - D^2/4 = V_T (1 - h^2)$. V_E is comparable to s^2

because both describe phenotypic variation in a clonal population and are generated by

stochastic noise. From our measured values (Fig. 4C), V_E for ribosome density was

0.0038 and 0.0034 in old and new mothers and not significantly different (p=0.40). For

elongation rate, V_E was 0.074 and 0.051, significantly higher in old mothers (p=.007),

and comparable with the trend reported for s^2 (5). A cross comparison of V_E for

ribosome and elongation showed that the latter was significantly higher both in old

mothers (0.00381 vs. 0.00744; p=1x10^{-6}) and in new mothers (0.00338 vs. 0.00510;

p=0.0041). This inflation of the elongation V_E is consistent with the hypothesis that

downstream traits are inherently noisier because they are at the end of many

converging pathways or networks, each one with its own inherent noise (37, 38).

Because ribosomes are farther upstream, they have fewer inputs and a lower final V_E.

Thus, another answer is that although complex traits such as elongation are important

for a cell and under strong natural selection, they are most likely downstream and their

V_E are noisier and higher in value.

Previous studies have shown that old and new daughters in *E. coli* are

asymmetrical for elongation rates (1, 4–7, 15, 16), gene expression (3), and vulnerability

to dying in the presence of damage agents (5). We show here that the variances (V_E) of

ribosome density and elongation rates are also asymmetric. If asymmetry is a central

characteristic of bacterial aging, identifying its cause becomes an important goal. We

began our studies by examining asymmetry for elongation rates, and then to gene

products and now to ribosomes. Because ribosomes are positioned highly upstream in

pathway of these cells and their abundance correlates strongly with elongation rates in

E. coli (27–30), their physiological role is both fundamental and general. As we have

argued, our previous report on the asymmetry of gene products was instructive but left

uncertain whether the outcome was unique to the gene that was studied or a more

general phenomenon. Our present results on ribosome asymmetry moves the process

to a more fundamental level because ribosomes can affect more genes and more

downstream processes. More importantly, existing measurements of ribosome cytology

and cell growth in *E. coli*, in conjunction with our new results, allow us to quantitatively

evaluate, understandingly not to prove, our previous suggestion (3) that space

competition between damage aggregates and ribosomes could be responsible for the

asymmetry.

The nucleoid and ribosomes in a rod-shaped *E. coli* occupy non-overlapping

space along the length of the cell (20). At birth, the cells are smaller and the nucleiod

resides at the mid-region of the rod and the ribosomes are displaced to the poles. As

the cell elongates, the nucleoid splits to create the two nucleoids for the daughters. The

splitting opens space in mid-region and the void is then filled by the ribosomes. Cell

division ensues with a cleavage at the mid-region, and the ribosomes at the mid-region

are distributed to the new poles of the daughters. However, the data used to generate

this description did not distinguish between old and new poles or daughters. As a result,

the cells were oriented randomly, and the resulting profiles are generic averages and

necessarily symmetrical. When we distinguished between the old and new poles and

daughters, the ribosome distribution resembles the generic profile only in new mothers

and is asymmetrical in old mothers (Fig. 7). Because damage aggregates also tend to

reside at the poles (7), we need to evaluate whether the volume of added aggregates

displaced a volume of ribosomes that could quantitatively account for the elongation

rate difference between old and new daughters from old mothers (Fig. 2C). Such an

analysis is possible using published information.

Single cell measurements estimate elongation rates for *E. coli* to be 2.148 and

1.964 hr^{-1} for new and old daughters in rich media (4). The well-known linear

relationship between elongation rate and ribosomal content (27–29, 31) can be

quantified from the data (30) as $Y = 337.046 \cdot X + 87.167$, where Y is ribosomal content

(A.U.) and X is elongation rate (hr^{-1}). The relationship between ribosome density and

elongation rate in Fig. 5 could not be used because the values were normalized.

Letting $X_{New} = 2.148$ and $X_{Old} = 1.964$ hr^{-1} from above, $Y_{New} = 811.02$ and $Y_{Old} = 749.08$

A.U. The proportional reduction of ribosomes in the old daughter relative to the new

daughter is therefore $p = (811.02 - 749.08) / 811.02 = 0.0764$. The total volume

occupied by ribosomes in rich media is estimated to be $V_{ribo} = 0.79$ μm^3 (32). The

volume needed to reduce the elongate rate of the old daughter to 1.964 hr^{-1} becomes Δ

$= p \cdot V_{ribo} = 0.0764 \cdot 0.79 = 0.0603$ μm^3. Aggregate radii have been estimated from

diffusion constants for *E. coli* cells in rich media (39). By converting the radii to a

sphere as $(4\pi/3) \cdot radius^3$, the largest aggregates had mean values of 0.0852 μm^3,

which is well in the range of our estimate of $\Delta = 0.0603$ μm^3. Thus, space competition

between aggregates and ribosomes could quantitatively account for the lower

elongation rate of old daughters of old mothers.

Our above space competition model implies that elongation rates and the size of

damage aggregates should be negatively correlated. However, while the initial study by

Lindner et al. (7) found a strong and significant negative correlation (p<.0001), a later

study by Govers et al. (8) found one that was negative but non-significant (p=.08).

Lindner et al. and Govers et al., respectively, used different stressors to induce damage

(streptomycin and heat shock) and fluorescent reporters (YFP and msfGFP) to quantify

aggregate size. The YFP and msfGFP used in both studies was fused to the chaperone

IbpA, which associates to aggregates of damaged proteins. Because Govers et al.

believed that IbpA-YFP formed self-aggregates they suggested that the correlation of

Lindner et al. was an artifact of YFP self-aggregation. However, the rejection of Lindner

et al. may be premature. First, the suggestion of YFP self-aggregation by Govers et al.

was based on their demonstration that IbpA-YFP increased the expression of IbpA (at

another locus), which is induced by perturbations to protein homeostasis (8). The

expression was not increased in the IbpA-msfGFP control. Because the assay was

conducted in the absence of heat shock or another stressor, they surmised that it was

YFP self-aggregates that had perturbed the homeostasis. However, a recent report that

IbpA self-represses its own translation because over-expression is harmful in the

absence of stress (40) suggests an alternative explanation for why IbpA-YFP increased

expression. It could be that IbpA-YFP was unable to self-repress because of its larger

size. Second, msfGFP-YFP may have its own issues because super folding GFP's are

more prone to photobleaching (41, 42). Could photobleaching have weakened the

correlation of Govers et al. and rendered it non-significant? Third, most importantly, the

correlations of Lindner et al. and Govers et al. must be compared with caution. While

Lindner et al. induced cell damage with 10μg ml^{-1} of streptomycin, where the minimal

inhibitory concentration is 16-32 μg ml^{-1} (43, 44), Govers et al. used 47°C heat shock for

15 min, where 50°C at 15 min is lethal (8). Although both Lindner et al. and Govers et

al. grew their cells in LB broth, the elongation rate in the two studies, respectively,

centered about 0.35 hr^{-1} and 0.25 hr^{-1} (19 and 27 min doubling times). To have a

doubling time of 27 min in LB, the cells in Govers et al. must have experienced severe

stress and damage. In relative terms, the two treatments, respectively, were 10/16 to

10/32 = 63% to 31% vs. 47/50 = 94% of stopping growth. It is likely that at 94% lethality,

the stress level in Govers et al. is collapsing most cellular processes, including protein

aggregation. This possibility is supported by Govers et al.'s measurement of the mean

IbpA-msfGFP fluorescence per cell as a function of increasing heat shock temperature.

The mean response increased linearly from 42°C up to 48°C, and then from 48°C to

50°C it crashed abruptly back to the control level (shock-free) (8). If the accuracy of

laboratory incubators is ±1°C, Govers et al. were using IbpA-msfGFP dangerously near

the failure point of the reporter, which would could have underestimated (or estimated

with more error) the true amount of protein damage in the cells. If a 47°C heat shock

induced damage to other cellular components besides proteins (e.g. DNA), the IbpA-

msfGFP would not have reported them. High heat can damage DNA and elevate

mutation rates in *E. coli* (45, 46). The choice of 47°C was justified because by Govers

et al. wanted to test whether protein aggregates could protect cells from new damage.

However, if the underestimates or statistical analyses weakened by measurement

errors compromised the correlation of Govers et al., the two correlations are not

equivalent and cannot be used to reject each other.

Until new studies prove otherwise, we believe that the elongation rate and

aggregate correlations of Lindner et al. and Govers et al. should be viewed as

alternative models to be further tested. Although the correlations do not bear directly on

our ribosome results, they are relevant to our discussion of the relationship of

ribosomes, damage aggregates, and elongation rates. Our space competition model

(see above) was evaluated to test whether our data was (or was not) quantitatively

consistent with the correlation reported by Lindner et al. We look forward to other tests

or challenges as connecting the two correlations will be critical. The protection

uncovered by Govers et al. is an exciting new phenomenon. However, we are most

excited by the possibility that both correlations ar correct and protein aggregates

are harmful and protective under low and high stress, respectively. At low stress levels,

proteins could be the main target of damage and the amount of protein aggregates

correlates with elongation rate. The damaged proteins are aggregated and then

partitioned asymmetrically between the daughters. Protein repair is occurring (18, 19)

and its effect is to lower the damage rate experienced by the cell. If aggregates are

observed, damage rates must be exceeding repair rates. However, a full stress

response (e. g. SOS, efflux transporters, and RpoS) (47) is too costly and unnecessary.

At high stress levels and near lethality, proteins, DNA, and other cellular processes

(including protein aggregation and IbpA-msfGFP reporting) are all failing due to damage

and the correlation between elongation rate and protein aggregates is weakened. In

order to survive the cell needs a full stress response and aggregates become beneficial

because they are present to act as a trigger (or the "epigenetic memory" of Govers et

al.). In effect, protein aggregates at high levels are inducing persister states (48–51).

In summary, previous reports have documented in *E. coli* the asymmetrical

associations between bacterial aging, elongation rates, cell poles, damage aggregates,

and expressed gene products, and we have added one more phenotype, the density of

ribosomes. Because ribosomes have a fundamental role in determining cell function,

our results indicate that bacterial aging and the asymmetry are resulting from processes

that are central to cell function and not just a subset of genes or pathways. Moreover,

ribosomal function has been well characterized and the information allowed us to show

that space competition between ribosomes and damage aggregates, both of which tend

to reside in the poles, could quantitatively account for the elongation rate asymmetry

between old and new daughters. The latter calculation does not prove a causal role for

damage aggregates, but we hope it will motivate and guide new research to evaluate

hypotheses emanating from the work by us and others.

MATERIALS AND METHODS

Bacterial strain. *E. coli* AFS55 was used to quantify ribosome density. AFS55

has a translational fusion of *yfp* (yellow fluorescent protein) to the C-terminus of *rpsB*

(ribosomal subunit S2) that served as a reporter for ribosome density (20). AFS55 was

chosen for this study because it offers currently one of the best fluorescent reporters for

ribosomes (see Introduction for full details). For example, numerous controls have been

conducted to rule out non-uniform fluorescent distribution by YFP self-aggregation

artifacts (20, 24). Additionally, the S2 protein was picked from other possible ribosomal

proteins because it is known to have negligible number of free copies (not attached to

ribosomes) in the cell (20, 21) .

Growing cells and preparing microscopy slides. All cells were stored in 40%

glycerol at -80˚C and incubated at 37˚C. A glycerol stock was streaked onto LB agar

plates (52)to obtain single colonies. Liquid cultures were made by inoculating a single

colony into 10 ml M9 (52)media and incubating overnight. The overnight culture was

diluted 1:100 in M9 and grown for two more hours. One µl of two-hour culture was then

pipetted onto a 10 µl agarose pad with M9 and 15% agarose. The agarose pad was

flipped with the bacterial side down onto a 24 x 60 mm cover glass, which was then

placed over a 25 x 75 mm single depression slide. The contact between the cover

glass and slide was sealed with Vaseline to prevent desiccation. These methods were

modified from Rang et al. (16, 17) and Stewart et al. (1).

Microscopy and time-lapse movies. After cells were placed onto agar pads,

the microscopy slides imaged at 37˚C with an inverted microscope (Nikon Eclipse Ti-S),

equipped with Nikon NIS-Elements AR control software, 100X objective (CFI Plan APO

NA 1.4), external phase contrast rings for full intensity fluorescence imaging,

fluorescence light source (Prior Lumen 200) with motorized shutter (Lambda 10-B

Sutter SmartShutter), and camera (Retiga 2000R FAST 1394, mono, 12 bit). To be

able to quantify the fluorescence of growing single cells, movies were started and ended

following guidelines. To prevent crowing as cells grew into micro-colonies over time,

movies were started from microscope fields containing single cells with distant

neighbors. To avoid effects of crowding within the micro-colonies, movies were ended

when micro-colonies exceeded 128 cells. Phase contrast and fluorescence images

were recorded every 2 and 20 minutes, respectively. Thus, every tenth Phase image

has a matching fluorescence image taken at the same time and with the same X-Y

coordinates. Such fluorescence-phase pairs were used for our deconvolution protocols

(see next Section).

Cells to be measured were collected from colonies started by a single cell.

Measurements were made at generations 4, 5, 6, and 7, which correspond to colonies

with 16, 32, 64, and 128 cells. Smaller sizes (1, 2, 4, and 8 cells) were not measured

because a colony started from a single cell experiences a lag, despite being inoculated

with fresh exponentially cells. Colony sizes larger than 128 were also excluded

because cell growth slows down due to crowding within and between colonies. Thus, to

be safe and to ensure the most uniform physiological state for the cells that were

measured, we restricted our measurements to colonies with 16 to 128 cells.

Image processing, discounting light scatter from neighbors, outlining cells,

and deconvolution. Fluorescence measurements reporting ribosome density in single

cells were obtained from the fluorescence images. Because a cell in a micro-colony

receives scattered fluorescence light from neighboring cells, the value of a fluorescence

pixel inside a cell is the sum of the true emission of the pixel plus the scatter. Note that

background pixels, which are outside a cell and lacking a fluorescent reporter, should

have a true emission of zero (not be emitting) and only contain scatter from pixels in the

neighboring cells. The true emission can be estimated by dividing the image into all its

pixels, treating the pixels as light point sources, using a system of differential equations

to describe the scatter from every pixel to every other pixel, and solving the system.

Because every fluorescence image has a matching phase image (see above

fluorescence-phase pair), any single cell can be outlined in the Phase image. The

correct total amount of fluorescence in the cell is therefore the true emission of all pixels

contained by the outline. However, because in practice the value of a pixel is measured

with error (stochasticity), the system cannot be solved analytically and must be solved

numerically by the iterative process of deconvolution (53). Deconvolution introduces a

new problem because the number of iterations needs to be optimized. The true

emission is underestimated if too few iterations are conducted (54). Too many

iterations also underestimate, but by introducing ringing (55). Ringing results when a

background pixel is brighter than average and is treated as an emitting pixel by the

over-iteration. As a result, the over-iteration removes light from the true emission value

of the pixels inside cells and allocates them to such background pixels. By tracking and

plotting the sum of all pixels in a cell as a function of iterations, we have found that the

sum invariably produced a peak by increasing initially and then decreasing (3). The

number of iterations corresponding to the peak is chosen as the desired number that

balanced the opposing effects of under and over iterating. Because a detailed

description has been published in Shi et al. (3) we present next a more abbreviated

technical summary.

The background of the fluorescence-phase image pairs (see above) was first

subtracted using the "rolling ball" algorithm in ImageJ (NIH) with ball radius 20 pixels.

Noise created by heat overflow of single pixel was corrected by "remove outliers"

algorithm in ImageJ with threshold intensity difference 1000 and threshold radius 0.5

pixel. The resulting image was processed by iterative Lucy-Richardson deconvolution in

Matlab 2017b using an optimal number of iterations for each fluorescence-phase pair as

described above. The outline of a single cell was traced manually with ImageJ. Blind

replicates without any awareness of the history of the cells found no bias between

outlines measured for old and new mothers.

Statistical analyses, data measurements, and normalization. Standard t-test

and regression analyses were implemented with packaged R functions. All reported

error bars are standard error of the mean (SEM).

Following deconvolution, fluorescence images and the outlines of cells were

imported as data matrices into computational R-programs. For fluorescence

measurements reporting ribosomal density, the outlines were cut into the desired

number of partitions (halves or quartiles; see Fig. 1), the partitions were superimposed

onto the fluorescence images, and the fluorescence pixel values contained by each

partition were summed. The sum was then divided by the area of the partitions to

obtain the average pixel value or density. The elongation rate of a cell was estimated

as $r = \ln(L_T/L_0) / T$, where L_T and L_0 are the lengths of the cell when it was born and

divided, respectively, and T is the time to division (min).   Cell length measured

manually from the Phase images with ImageJ. A log transformation was used because

elongation rates are known to be exponential (1).

For Figs. 3A and 3B, ribosome densities (here denoted X) were normalized as (X

– u) / u, where $u = (X_{Old} + X_{New}) / 2$. For Figs. 3C and 3D, elongation rates were likewise

normalized. For Fig. 7, the cells were pooled by first dividing each cell into four length

quartiles (NP, L2, L3, and OP; see Fig. 1D). Each cell was then normalized by dividing

the ribosome density of each of the cell's quartiles by the sum of the ribosome density

of the cell's four quartiles. After normalization, the sum equals 1.0. For statistical

comparisons between two regression slopes (Fig. 7B), the SEM values and sample

sizes from the regression were used to estimate the resulting t-statistic and ensuing p-

value (56).

To pool cells for comparison over a time cycle from birth to division, time was

normalized into quartiles (B, T2, T3, and D; birth, quartile 2, quartile 3, and division).

For example, given a cell with a division time of 28 min and time-lapse images of the

cell from birth to division, the images from the window t= [0 to 7], [7 to 14], [14 to 21],

and [21 to 28] min. Because time quartiles are dimensionless, we provide as a

reference the mean division times that were observed. The means for old and new

daughters (depicted in Figs. 3C and 3D) were, respectively, 74.58 and 67.48 min (from

old mothers; n=216) and 69.76 and 65.71 min (from new mothers; n=198).

All statistical analyses were conducted with both length and ribosome densities

values normalized as halves or quartiles. We provide in Supplemental Materials (Fig.

S1) a plot of ribosomal density versus length as a continuous variable for visualization

and reference.

ACKNOWLEDGMENTS

We thank James Weisshaar for bacterial strain. This study was support by the NSF

grant (DEB 1354253) to L.C.

SUPPLEMENTAL MATERIALS

Figure S1

REFERENCES

1.   Stewart EJ, Madden R, Paul G, Taddei F, Burland V. 2005. Aging and death in an

organism that reproduces by morphologically symmetric division. PLoS Biol 3:e45.

2.   Chao L. 2010. A Model for Damage Load and Its Implications for the Evolution of

671   Bacterial Aging. PLoS Genet 6:1001076.

672   3.   Shi C, Chao L, Proenca AM, Qiu A, Chao J, Rang CU. 2020. Allocation of gene

673   products to daughter cells is determined by the age of the mother in single

674   Escherichia coli cells. Proceedings Royal Society B 287:20200569.

675   4.   Proenca AM, Rang CU, Buetz C, Shi C, Chao L. 2018. Age structure landscapes

676   emerge from the equilibrium between aging and rejuvenation in bacterial

677   populations. Nat Commun 9:1–11.

678   5.   Proenca Menegaz A, Rang CU, Qiu A, Shi C, Chao L. 2019. Cell aging preserves

679   cellular immortality in the presence of lethal levels of damage. PLoS Biol May 23.

680   6.   Urszula Ł, Glover G, Capilla-lasheras P, Young AJ, Pagliara S, Young AJ,

681   Pagliara S. 2019. Bacterial ageing in the absence of external stressors. Phil Trans

682   R Soc B 374:20180442.

683   7.   Lindner AB, Madden R, Demarez A, Stewart EJ, Taddei F. 2008. Asymmetric

684   segregation of protein aggregates is associated with cellular aging and

685   rejuvenation. Proceedings of the National Academy of Sciences 105:3076–3081.

686   8.   Govers SK, Mortier J, Adam A, Aertsen A. 2018. Protein aggregates encode

687   epigenetic memory of stressful encounters in individual Escherichia coli cells.

688   PLoS Biol 16:e2003853.

689   9.   Maisonneuve E, Ezraty B, Dukan S. 2008. Protein aggregates: An aging factor

690   involved in cell death. J Bacteriol 190:6070–6075.

10.   Gatti-Lafranconi P, Natalello A, Ami D, Doglia SM, Lotti M. 2011. Concepts and

tools to exploit the potential of bacterial inclusion bodies in protein science and

biotechnology. FEBS Journal 278:2408–2418.

11.   Schramm FD, Schroeder K, Jonas K. 2019. Protein aggregation in bacteria.

FEMS Microbiol Rev. Oxford University Press

https://doi.org/10.1093/femsre/fuz026.

12.   Rang CU, Proenca A, Buetz C, Shi C, Chao L. 2018. Minicells as a damage

disposal mechanism in Escherichia coli. mSphere 3:e00428-18.

13.   Vedel S, Nunns H, Košmrlj A, Semsey S, Trusina A. 2016. Asymmetric Damage

Segregation Constitutes an Emergent Population-Level Stress Response. Cell

Syst 3:187–198.

14.   Winkler J, Seybert A, König L, Pruggnaller S, Haselmann U, Sourjik V, Weiss M,

Frangakis AS, Mogk A, Bukau B. 2010. Quantitative and spatio-temporal features

of protein aggregation in Escherichia coli and consequences on protein quality

control and cellular ageing. EMBO J 29:910–923.

15.   Chao L, Rang CU, Proenca AM, Chao JU. 2016. Asymmetrical damage

partitioning in bacteria: a model for the evolution of stochasticity, determinism,

and genetic assimilation. PLoS Comput Biol 12:1–17.

16.   Rang CU, Peng AY, Chao L. 2011. Temporal dynamics of bacterial aging and

rejuvenation. Current Biology 21:1813–1816.

17.   Rang CU, Peng AY, Poon AF, Chao L. 2012. Ageing in Escherichia coli requires

damage by an extrinsic agent. Microbiology (United Kingdom) 158:1553–1559.

18.   Ezraty B, Gennaris A, Barras F, Collet JF. 2017. Oxidative stress, protein damage

and repair in bacteria. Nat Rev Microbiol https://doi.org/10.1038/nrmicro.2017.26.

19.   Merdanovic M, Clausen T, Kaiser M, Huber R, Ehrmann M. 2011. Protein quality

control in the bacterial periplasm. Annu Rev Microbiol

https://doi.org/10.1146/annurev-micro-090110-102925.

20.   Bakshi S, Siryaporn A, Goulian M, Weisshaar JC. 2012. Superresolution imaging

of ribosomes and RNA polymerase in live Escherichia coli cells. Mol Microbiol

85:21–38.

21.   Sanamrad A, Persson F, Lundius EG, Fange D, Gynnå AH, Elf J. 2014. Single-

particle tracking reveals that free ribosomal subunits are not excluded from the

Escherichia coli nucleoid. Proc Natl Acad Sci U S A 111:11413–11418.

22.   Swulius MT, Jensen GJ. 2012. The helical mreb cytoskeleton in Escherichia coli

MC1000/pLE7 is an artifact of the N-terminal yellow fluorescent protein tag. J

Bacteriol 194.

23.   Landgraf D, Okumus B, Chien P, Baker TA, Paulsson J. 2012. Segregation of

molecules at cell division reveals native protein localization. Nat Methods 9.

24.   Batchelor E, Goulian M. 2006. Imaging OmpR localization in Escherichia coli. Mol

Microbiol 59.

25.   Pogliano J, Ho TQ, Zhong Z, Helinski DR. 2001. Multicopy plasmids are clustered

and localized in Escherichia coli. Proc Natl Acad Sci U S A 98.

26.   Cabrera JE, Cagliero C, Quan S, Squires CL, Ding JJ. 2009. Active transcription

of rRNA operons condenses the nucleoid in Escherichia coli: Examining the effect

of transcription on nucleoid structure in the absence of transertion. J Bacteriol 191.

27.   Schaechter M, Maaløe O, Kjeldgaard NO. 1958. Dependency on medium and

temperature of cell size and chemical composition during balanced growth of

Salmonella typhimurium. J Gen Microbiol 19:592–606.

28.   Neidhardt FC, Magasanik B. 1960. Studies on the role of ribonucleic acid in the

growth of bacteria. Biochim Biophys Acta 42:99–116.

29.   Kjeldgaard NO, Kurland CG. 1963. The distribution of soluble and ribosomal RNA

as a function of growth rate. J Mol Biol 6:341–348.

30.   Poulsen LK, Licht TR, Rang C, Krogfelt K a, Molin S. 1995. Physiological state of

Escherichia Coli BJ4 growing in the large intestines of streptomycin-treated Mice.

J Bacteriol 177:5840–5845.

31.   Greulich P, Scott M, Evans MR, Allen RJ. 2015. Growth-dependent bacterial

susceptibility to ribosome-targeting antibiotics. Mol Syst Biol 11:796.

32.   Woldringh CL, Nanninga N. 1985. Structure of Nucleoid and Cytoplasm in the

Intact Cell., p. 161–197. *In* Nanninga, N (ed.), Molecular Cytology of Escherichia

coli. Academic Press New York and London, London, New York.

33.   Wang P, Robert L, Pelletier J, Dang WL, Taddei F, Wright A, Jun S. 2010. Robust
752   growth of escherichia coli. Current Biology 20:1099–1103.

753   34.   Chao L. 1991. Levels of selection, evolution of sex in RNA viruses, and the origin
754   of life. J Theor Biol 153:229–246.

755   35.   Miller SL, Urey HC. 1959. Organic compound synthesis on the primitive earth.
756   Science (1979) 130:245–251.

757   36.   Michaelian K. 2011. Thermodynamic dissipation theory for the origin of life. Earth
758   Syst Dynam 2:37–51.

759   37.   Price T, Schluter D. 1991. On the Low Heritability of Life-History Traits. Evolution
760   (N Y) 45:853–861.

761   38.   Houle D. 1992. Comparing Evolvability and Variability of Quantitative Traits.
762   Genetics 130:195–204.

763   39.   Coquel A-SS, Jacob J-PP, Primet M, Demarez A, Dimiccoli M, Julou T, Moisan L,
764   Lindner AB, Berry H. 2013. Localization of protein aggregation in Escherichia coli
765   is governed by diffusion and nucleoid macromolecular crowding effect. PLoS
766   Comput Biol 9:e1003038.

767   40.   Miwa T, Chadani Y, Taguchi H. 2021. Escherichia coli small heat shock protein
768   IbpA is an aggregation-sensor that self-regulates its own expression at
769   posttranscriptional levels. Mol Microbiol 115.

770   41.   Valbuena FM, Fitzgerald I, Strack RL, Andruska N, Smith L, Glick BS. 2020. A
771   photostable monomeric superfolder green fluorescent protein. Traffic 21.

772   42.   Pédelacq JD, Cabantous S, Tran T, Terwilliger TC, Waldo GS. 2006. Engineering

773   and characterization of a superfolder green fluorescent protein. Nat Biotechnol 24.

774   43.   Tashiro Y, Eida H, Ishii S, Futamata H, Okabe S. 2017. Generation of small

775   colony variants in biofilms by Escherichia coli harboring a conjugative F plasmid.

776   Microbes Environ 32.

777   44.   Regoes RR, Wiuff C, Zappala RM, Garner KN, Baquero F, Levin BR. 2004.

778   Pharmacodynamic functions: a multiparameter approach to the design of

779   antibiotic treatment regimens. Antimicrob Agents Chemother 48:3670–6.

780   45.   Zamenhof S, Greer S. 1958. Heat as an agent producing high frequency of

781   mutations and unstable genes in Escherichia coli. Nature 182.

782   46.   Chu XL, Zhang BW, Zhang QG, Zhu BR, Lin K, Zhang DY. 2018. Temperature

783   responses of mutation rate and mutational spectrum in an Escherichia coli strain

784   and the correlation with metabolic rate. BMC Evol Biol 18.

785   47.   Roemhild R, Bollenbach T, Andersson DI. 2022. The physiology and genetics of

786   bacterial responses to antibiotic combinations. Nat Rev Microbiol

787   https://doi.org/10.1038/s41579-022-00700-5.

788   48.   Leszczynska D, Matuszewska E, Kuczynska-Wisnik D, Furmanek-Blaszk B,

789   Laskowska E. 2013. The Formation of Persister Cells in Stationary-Phase

790   Cultures of Escherichia Coli Is Associated with the Aggregation of Endogenous

791   Proteins. PLoS One 8:e54737.

49.   Dewachter L, Bollen C, Wilmaerts D, Louwagie E, Herpels P, Matthay P,

Khodaparast L, Khodaparast L, Rousseau F, Schymkowitz J, Michiels J. 2021.

The dynamic transition of persistence toward the viable but nonculturable state

during stationary phase is driven by protein aggregation. mBio 12.

50.   Goode O, Smith A, Zarkan A, Cama J, Invergo BM, Belgami D, Caño-Muñiz S,

Metz J, O'neill P, Jeffries A, Norville IH, David J, Summers D, Pagliara S. 2021.

Persister escherichia coli cells have a lower intracellular ph than susceptible cells

but maintain their ph in response to antibiotic treatment. mBio 12.

51.   Balaban NQ, Helaine S, Lewis K, Ackermann M, Aldridge B, Andersson DI,

Brynildsen MP, Bumann D, Camilli A, Collins JJ, Dehio C, Fortune S, Ghigo JM,

Hardt WD, Harms A, Heinemann M, Hung DT, Jenal U, Levin BR, Michiels J,

Storz G, Tan MW, Tenson T, Van Melderen L, Zinkernagel A. 2019. Definitions

and guidelines for research on antibiotic persistence. Nature Reviews

Microbiology 2019 17:7 17:441–448.

52.   Sambrook J, Fritsch EF, Maniatis T. 1989. Molecular cloning: a laboratory manual.

Molecular cloning: a laboratory manual.

53.   Lucy LB. 1974. An iterative technique for the rectification of observed distributions.

Astron J 79:745–754.

54.   Biggs DSC, Andrews M. 1997. Acceleration of iterative image restoration

algorithms. Appl Opt 36.

55.   Mosleh A, Langlois JMP, Green P. 2014. Image deconvolution ringing artifact

detection and removal via PSF frequency analysisLecture Notes in Computer

Science (including subseries Lecture Notes in Artificial Intelligence and Lecture

Notes in Bioinformatics).

56. Sokal RR, Rohlf FJ. 1995. Biometry, 3rd ed. WH Freeman and Company, New

York.

Fig 1 Cell division and polarity in *E. coli.* (A) Assignment of old (red) and new (blue)

poles and daughters. The starting cell is white because it was randomly picked to start

the experiment and its polarity was unknown. Because the division plane (dashed line)

cuts the white cell at the midpoint of the long axis, the poles formed at the division point

are new and the distal poles are old. At the next division, the new daughter acquires the

new pole, while the old daughter receives the old pole. Note that two divisions must be

tracked to determine the old and new daughters from the starting white cell. The

outlines of the bottom four daughters in the figure are colored red and blue to identify

them as old and new daughters, while the intracellular red and blue colors identify the

old and new poles, also designated as O and N. Although old and new poles and

daughters are here tracked for only 2 generations, the notation and tracking methods

can be extended into generations 3, 4, 5, 6, 7, and further if needed. For example, if the

old and new poles of any of the four daughters after 2 divisions are known as in Fig. 1A,

and these daughters elongate, become a mother, and divide to produce two grand-

daugthers, the polarity of the grand-daughters can be determined by the same tracking

methods.(B) Time-lapse images of an *E. coli* bacterium dividing into two and four cells.

Top row: Phase contrast. Second row: Assignment of old (red) and new (blue) poles

from the top row cells. Third row: Fluorescence images matching phase images in time

and position. Bottom row: Heat-map of fluorescence images reporting ribosome density

in the top row cells, showing lesser intensity by the old poles (blue color spots) than the

new poles and inside the cells (purple color). Scale on the right shows intensity gradient.

(C) Schematic showing division of a cell into two halves containing either the new pole

or the old pole. The ratio of the fluorescence in the new half divided by that in the old

half (or pole ratio) was used to normalize and pool different old and new daughter pairs,

when each pair descended from a different mother. (D) Schematic showing division of

a cell into four length quartiles denoted NP, L2, L3 and OP in relation to the old and new

poles. These quartiles were used to quantify the distribution of fluorescence along the

length of a cell.

Fig 2 Ratios of ribosome density and elongation rates within and between cells.

Because ratios are a form of normalization, their values were derived from not

normalized data. Asterisks in figures indicate levels of significance from comparison

between ratios of new (blue bar) versus old (red bar) mothers (* ** ***, significance at p

< 0.05, 0.01, and 0.001). Error bars are SEM. Values of p in text below are for the

significance probability that a ratio is greater than 1.0, unless indicated otherwise.

Sample size for ratios is for pairs of data: n = 89 pairs correspond to 89 old/new

daughters of the same mother. Each pair is used to obtain one ratio to yield 89 ratios.

(A) Ribosome ratio of daughters (new/old) at birth from old (red) and new (blue) mother

cells. The ratio was 1.11 ± 0.018 from old mothers (p = 6.89×10^{-6}, one-tailed paired t-

test, n = 89 pairs) and 1.04 ± 0.015 from new mothers (p = 0.04, one-tailed paired t-test,

n = 91 pairs). The two ratios were also significantly different from each other (p= 0.007,

two-tailed non-paired t-test, ** in figure). (B) Ribosome ratio (new/old) of the two polar

halves (Fig. 1C) from old (red) and new (blue) mothers at division. The ratio in old

mothers was 1.10 ± 0.015 (p = 6.23×10^{-5}, one-tailed paired t-test, n = 89 pairs) and from

new mothers 1.03 ± 0.010 (p = 0.08, one-tailed paired t-test, n = 91 pairs). The

difference between the two ratios was significant (p= 1.3×10^{-4}, two-tailed non-paired t-

test, *** in figure). A comparison of the daughter and polar half ratios from old mothers

(Figs. 2A, 2B; red bars) found no significant difference (p = 0.62; two-tailed paired t-test).

A likewise comparison for new mothers (Figs. 2A, 2B; blue bars) also found no

significance (p = 0.27). (C) Elongation rate ratio of new over old daughters from new

(blue) and old (red) mothers. The ratio from old mothers was 1.11 ± 0 .017 (p = $1.3 \times 10^{-}$

^{10}, one-tailed paired t-test, n = 216 pairs) and from new mothers 1.07 ± 0.011 (p =

2.76×10^{-7}, one-tailed paired t-test, n = 198 pairs). The two ratios were significantly

different from each other (p= 0.02, two-tailed non-paired t-test; * in figure). Note that

these elongation rate ratios parallel the ribosomal pattern of a higher asymmetry in

daughters from old mothers (Fig. 2A).

Fig 3 Variation in ribosome density and elongation rate in old and new daughters from

old and new mothers. The measurements were made from the same cells in Fig. 2C.

All density and elongation values were normalized for this analysis. Means, variances,

and other parameters extracted from these data are presented in Fig. 4. (A) Top panel:

Normalized density distribution of ribosome density of old (red line) and new (blue line)

daughters from old mothers. Red and blue dots on the x-axis indicate the mean density

for each distribution. Dashed lines represent the density distribution of the old and new

daughters combined into one pooled total population. D (black arrow) shows the

distance between peaks of old and new daughter curves (*** $p=2.2 \times 10^{-16}$ for

significance of D>0; 1-tailed paired t-test; n=216 daughter pairs; see Fig. 4A). Bottom

panel: Normalized ribosome density of each old (red) and new (blue) daughter in pairs

from old mothers. The zero point is set as the average ribosome density for each pair.

As shown, the old daughter in each pair more often ends up on the minus side of the

pair's zero point, i.e. having lower ribosome density. (B) Same as (A), but from new

mothers (* p=0.014 for D>0; 1-tailed paired t-test; n=198 daughter pairs; see Fig. 4A).

(C) Normalized elongation rate distributions (*** $p=2.0 \times 10^{-11}$ for D>0; 1-tailed paired t-

test; n=216 daughter pairs; see Fig. 4B), but otherwise as (A). (D) Same as (C), but

from new mothers (*** $p=2.1 \times 10^{-7}$ for D>0; 1-tailed paired t-test; n=198 daughter pairs;

see Fig. 4B).

Fig 4 Variance components of ribosome density and elongation rates. Total (V_T),

stochastic (V_E) and deterministic ($D^2/4$) components of variances estimated from Fig. 3

(see Materials and Methods). The deterministic proportion of total variance V_T was $h^2 =$

$(D^2/4) / V_T$. The stochastic component of V_T is therefore $V_E = V_T (1 - h^2)$. Significance

testing for D>0 as described in Fig. 3 (*, **, and *** denote p<0.05, 0.01, and 0.001). (A)

Estimates for ribosomal density. (B) Estimates for elongation rates. (C) Comparison of

V_E estimates. Testing of significance difference between V_E was by a Bartlett's test of

homogeneity of variances. p-values[1] are for comparisons across rows (old versus new

mothers). p-values[2] are for significance of V_E down columns (elongation rate vs.

ribosome density).

Fig 5 Correlation between normalized elongation rate and ribosome density of old and

new daughters from old and new mothers. Because the old and new daughter values

were jointly normalized, they were not independent and their combination into a single

plot could not be assessed by statistical models assuming independence. Thus, all

analyses for the figure were conducted by randomizing the data and obtaining a null

distribution to estimate the significance values (see Materials and Methods). (A)

Ribosome density versus elongation rate between new and old daughters from old

mothers. Slope of linear regression = 0.498, correlation r = 0.387, $p < 1 \times 10^{-5}$, n = 216

daughter pairs. (B) Ribosome density versus elongation rate between new and old

daughters from new mothers. Slope of linear regression = 0.301, correlation r = 0.233, p

$= 4 \times 10^{-4}$, n = 198 daughter pairs.

Fig 6 Ratio of ribosome density in old and new mothers over quartile time from birth to

division. Ratios correspond to density in the new pole half divided by density in the old

pole half (Fig. 1C) of the mothers. Regression slopes were significantly greater than

zero for old (slope=0.0177, * p=0.026, df=278) but not for new mothers (slope=0.0039,

n.s. p=0.59, df=278). Regression was based on ratios that were binned by the four time

quartiles (Old mothers: n=71, 74, 46, 89; df=278. New mothers: n=68, 72, 49, 91;

df=278), although the plots only show the ribosome means of the four time bins for

illustration. Old mothers began with a higher asymmetry ratio and elevated to even

higher levels at division. On the other hand, new mothers had a lower and more

symmetrical ratio that was held more constant from birth to division.

Fig 7 Spatial and temporal distribution of ribosome density in old and new mothers. (A)

Density in length quartiles over time quartiles. Lines presented are from regressions of

ribosome densities for one length quartile that were binned by the four time quartiles.

All length quartiles for a mother type had time bins of the same sample size (old

mothers: n=71, 74, 46, 89; new mothers: n=68, 72, 49, 91). The single dots presented

are ribosome density means of the four time bins and are presented for illustration. (B)

Values of regression slopes of the 8 lines in Fig. 7A. All regression slopes were

significantly greater or less than, i.e. not equal to, zero (Old mothers: p=0.015, 1.46×10^{-11},

0.00029, 7.8×10^{-6}; df=278. New mothers: p=0.0012, 1.7×10^{-10}, 3.5×10^{-10}, 4.2×10^{-6};

df=278). To test whether old mothers were more asymmetrical than new ones, the

regression slopes of the pairs NP vs. OP and L2 vs. L3 for each mother type were

compared (see Materials and Methods for details). The comparisons test whether

length quartiles of the mothers on the left new pole side mirrors the right old pole side.

The horizontal brackets in the figure denote the comparisons and whether they were

significant. Old mothers were asymmetrical because their comparisons were all

significantly different (NP vs. OP and L2 vs. L3; p=0.04 and 0.008; df=158 and 112).

New mothers were symmetrical as the differences were not significant (NP vs. OP and

L2 vs. L3; p=0.11 and 0.44; df=156 and 115).

Fig S1 Fluorescence profile along new and old daughter pairs' cell lengths at birth. The fluorescence and length values were normalized (n = 89 pairs). Color designation is as in Fig. 1A for old (red) and new (blue) poles and daughters. Error bars show standard error of the mean (s.e.m.). These transects are shown only for visualization and were not used in the final analyses, which were based on halves and quartiles (Figs. 1C and 1D).

A

Cell of unknown polarity

Old pole O N New Pole   New pole N O Old Pole

O N   N O   O N   N O

Old daughter   New daughter   New daughter   Old daughter

B

1 μm

A.U.
5000

2500

C

| NP half | OP half |

New pole ⟵⟶ Old pole

D

| NP | L2 | L3 | OP |

New pole ⟵⟶ Old pole

Fig 1

Fig 2

Fig 3

A   Ribosome density

	V_T	D	$D^2/4$	h^2
Old Mothers	0.00634	0.101 ***	0.00253	0.399
New Mothers	0.00347	0.0184 *	0.00008	0.024

B   Elongation rate

	V_T	D	$D^2/4$	h^2
Old Mothers	0.00913	0.0821 ***	0.00168	0.185
New Mothers	0.00581	0.0533 ***	0.00071	0.122

C V_E estimates

	Old mothers	New mothers	p-value[1]
Elongation rate	0.00744	0.00510	0.0073 **
Ribosome density	0.00381	0.00338	0.40 n.s.
p-value[2]	0.000001 ***	0.0041 **	

Fig 4

Fig 5

Fig 6

Fig 7

Fig S1