Divergent Outcomes Of Myeloid Cell Infection By Gammaherpesvirus 68 Reveal Stepwise Regulation By Host And Viral Factors

Gabrielle Vragel[1], Kyra S. Noell[1,2], Ashley Tseng[1,2], Rachael E. Kostelecky[1], Elizabeth A. Alexander[1], Shirli Cohen[1], Catalina Galvan[1], Manaal Dalwadi[1], Brittany D. Gomez[1], Eva M. Medina[3], Darby G. Oldenburg[4], Andrew Goodspeed[5,6], Eric T. Clambey[2]*, Linda F. van Dyk[1]*#

Affiliations:

[1] Department of Immunology and Microbiology, [2] Department of Anesthesiology, [3] Department of Biochemistry and Molecular Genetics, University of Colorado Denver | Anschutz Medical Campus, School of Medicine, Aurora, CO, 80045, USA

[4] Gundersen Medical Foundation, Emplify Health System, La Crosse, Wisconsin, USA

[5] Department of Biomedical Informatics, [6] University of Colorado Cancer Center, University of Colorado Denver | Anschutz Medical Campus, School of Medicine, Aurora, CO, 80045, USA

* Co-Corresponding authors:

Eric T. Clambey, eric.clambey@cuanschutz.edu, 303-724-7783 (phone)

Linda F. van Dyk, linda.vandyk@cuanschutz.edu, 303-724-4207 (phone)

 # Lead author for MS correspondence

Linda F. van Dyk, linda.vandyk@cuanschutz.edu, 303-724-4207 (phone)

ORCIDs

https://orcid.org/0000-0001-9167-3013 (G.V.); 0000-0003-3623-8822 (K.S.N.); 0000-0002-3078-3718 (A.T.); 0000-0002-6529-3831 (R.E.K.); 0000-0002-4859-4122 (E.A.A.); 0000-0002-0253-786X (S.C.); 0009-0009-3439-7644 (C.G.); 0009-0005-3187-7803 (M.K.D); 0000-0002-3871-3773 (B.D.G.); 0000-0001-6899-0213 (E.M.M.); 0000-0003-2090-7489 (D.O.); 0000-0001-7055-7128 (A.G.); 0000-0002-7972-9544 (E.T.C.); 0000-0003-2662-5554 (L.F.v.D.)

Summary

The gammaherpesviruses establish a lifelong infection, with the cellular outcome of infection

intimately regulated by cell type. Though myeloid cells are an early infection target, infection

results in widely divergent outcomes without a clear explanation. Here we use murine

gammaherpesvirus 68 to demonstrate that macrophages are readily infectable, resulting in

three divergent infection outcomes dictated by viral and host factors. Infection in vivo and in a

cell culture model results in a high frequency of viable cells characterized by restricted viral

transcription and unexpected transcription of the ORF75 locus, with rare cells initiating but

failing to complete lytic replication. In contrast, infection with a high viral dose triggers abortive

infection and cell death. Restricted infection can be fully converted to productive lytic replication

by pre-treatment with the host cytokine IL-4. These studies demonstrate stepwise regulation of

virus infection and identify virus and host factors that dictate divergent outcomes in a single cell

type.

Introduction

The gammaherpesviruses (γHVs), including the human viruses Epstein-Barr Virus (EBV) and

Kaposi's sarcoma-associated herpesvirus (KSHV or HHV-8), are large double-stranded DNA

tumor viruses. The γHVs establish a lifelong infection and are associated with chronic

inflammation and cancer, particularly in immunosuppressed individuals [1-3]. Due to the strict

host specificity of the human γHVs, murine gammaherpesvirus 68 (MHV68, γHV68 or MuHV-4)

has been extensively used as a genetic and phenotypically relevant model to study γHV

infection and pathogenesis in vivo [4, 5]. The γHVs infect multiple cell types, resulting in a range

of outcomes including lytic infection characterized by active virus replication or quiescent latent

infection [1], dictated by target cell type-specific virus-host interactions. Infection outcomes at

the single-cell level can be heterogeneous, including abortive infection [6-10].

Myeloid cells, including macrophages, monocytes and dendritic cells are innate immune cells

that have critical functions during γHV infection: they promote the early antiviral response yet

are also targeted for infection and subject to viral manipulation. While there is clear evidence

that KSHV, EBV and MHV68 infect myeloid cells [11-23], the state of infection in these cells

remains ambiguous, with evidence for both lytic transcription [19, 24, 25] and latent infection

[15, 18]; infection also appears to be abortive or unstable over time [26]. In some cases, myeloid

cells are thought to facilitate transit to B cells where γHVs establish long-term latent infection

[14, 26, 27]. γHV infection has also been shown to alter myeloid cell function, including KSHV-

induced differentiation of dysfunctional macrophages [22], manipulation of innate immune

signaling [16, 28, 29], and impaired function [30].

The most detailed insights regarding macrophage infection in vivo have come from the MHV68

system. MHV68 lytically replicates in macrophages prior to B cell infection [14, 27, 31]; MHV68

also establishes long-term latent infection in macrophages [15, 32]. While MHV68 passage

through alveolar macrophages and marginal zone macrophages promotes establishment of B

cell latency [14, 33], subcapsular sinus macrophages limit [34]. Macrophage infection is further

regulated by viral (ORF4, ORF36 [24, 25]) and host factors (e.g. interleukin (IL)-4 and IL-13,

Liver X receptors, cholesterol synthesis, type I and II interferons [35-38]).

Here, we sought to define γHV macrophage infection outcomes with a new level resolution

using the MHV68 system. Early macrophage infection in vivo is dominated by cells with frequent

transcription of the ORF75 locus, a γHV-conserved gene expressed in Kaposi's sarcoma lesions

and in EBV-transformed latently-infected B cells [39-42]. Using a reductionist system, we find

that MHV68 robustly infects macrophages with three distinct outcomes regulated by virus and

host factors: i) a restricted form of infection with rare cells undergoing lytic replication, ii) a

transcriptionally distinct form of abortive replication accompanied by high rates of cell death, and

iii) fully productive lytic replication. These studies emphasize that macrophage infection is

subject to multiple levels of control, resulting in diverse infection outcomes in a single cell type.

RESULTS

Single cell RNA sequencing analysis of acute MHV68 macrophage infection reveals two

distinct transcriptional profiles.

MHV68 macrophage infection has been associated with both lytic replication and latent

infection [14, 15, 24, 32], potentially serving as an important intermediary prior to B-cell latency

[14]. Despite these insights, the exact nature of myeloid cell infection remains incompletely

characterized. To define the viral transcriptional signatures at the earliest stages of macrophage

infection in vivo, we sort-purified MHV68 infected peritoneal macrophages from

immunocompetent C57BL/6J mice at 16 hours post-infection, followed by single-cell RNA-

sequencing (scRNA-seq). MHV68 infected cells were purified based on expression of a viral

reporter gene encoded by the <u>l</u>atency-<u>a</u>ssociated <u>n</u>uclear <u>a</u>ntigen fused with the <u>beta</u>-<u>lac</u>tamase

enzyme (LANAβlac), a sensitive system that enables the direct identification and purification of

both lytic- and latently-infected cells [43, 44]. As expected, Seurat-based clustering and UMAP-

based visualization identified a dominant cluster of LANA+ peritoneal macrophages, with minor

cell subsets including: i) macrophages with a proliferative gene signature (i.e. "macrophage

proliferating"), ii) macrophages with an MHV68 lytic gene signature (i.e. "macrophage lytic"),

defined by expression of two canonical lytic genes ORF7 and ORF54, and infrequent iii) B cells,

iv) monocytes, v) dendritic cells and vi) T cells (Fig. 1A-C, Fig. S1A). Quality control metrics

were relatively comparable across all cell subsets, including the "macrophage lytic" subset (Fig.

S1B-E).

When we visualized and quantified the frequency of cells with detectable viral unique

molecular identifiers (UMIs), the majority (n=7,512; 82.54%) of LANA+ macrophages had

detectable viral reads accounting for <0.1% of total UMIs per cell (Fig. 1D-E). In contrast rare

macrophages with a lytic signature were characterized by a higher proportion (>0.1%) of viral

reads per cell (Fig. 1D-E). The presence of cells with no detectable viral UMIs could either be

due to the depth of sequencing or to the known sparseness of scRNA-seq data [45]. Increased

granularity of cell clustering identified 14 cell clusters (Fig. S2) but did not significantly change

defined cell populations nor the percentage of viral UMIs detected per cell subsets defined in

Figure 1.

Next, we quantified the frequency of macrophages that expressed each viral gene,

comparing the dominant macrophage population (n=9101 cells), with rare macrophages

characterized by a proliferative (n=48 cells) or lytic (n=25 cells) gene expression signature. We

detected ORF73-βlac mRNA, the transcript encoding the LANAβlac reporter by which cells were

purified, in 12.89% of macrophages, with equivalent or higher frequencies of detection for only 3

other viral genes: ORF75C (30.75%), ORF75B (27.83%) and ORF75A (12.54%) (Fig. 2A).

Additional transcripts were detected in a lower frequency of cells: ORF72 (5.3%), ORF58

(3.5%), with detectable RNA for M3, ORF7, ORF50, ORF57, ORF61, ORF74, M12, M13, or

M14 in 1-3% of cells (yellow circles, Fig. 2A). Remaining viral genes were only sporadically

detected (<1% of cells). A similar hierarchy, albeit reduced in magnitude, was observed in

proliferative macrophages (Fig. 2B). In contrast, rare lytic macrophages had a distinct viral gene

signature: multiple lytic cycle-associated genes (ORF7, ORF54, ORF57, ORF58, and ORF61)

were detected in 100% of cells (dark blue circles, Fig. 2C), with many more lytic cycle RNAs

(M3, ORF6, ORF21, ORF38, ORF50, ORF52, ORF59, ORF72 and ORF75C) detected in >75%

of cells (light blue circles, Fig. 2C). Detection of multiple, canonical lytic cycle RNAs in these

cells is consistent with these cells initiating lytic gene transcription in vivo.

The ORF75 locus is conserved among γHVs, with MHV68 containing triplicated ORF75

genes, ORF75A, ORF75B and ORF75C [4, 46]. Previous studies of this locus have revealed

multiple transcriptional units, including a polycistronic RNA that spans from the 5' end of

ORF75A through the 3' end of ORF75C [46]. Though these genes likely arose from gene

duplication, there is no significant similarity between the 3' ends of ORF75A and ORF75B or

ORF75A and ORF75C; only 22 of 26 nucleotides align between the final 150 nucleotides of

ORF75B and ORF75C (defined by blastn [47], data not shown). The nucleotide divergence

between ORF75 genes strongly suggests that UMIs mapping to each ORF75 gene were not

due to mapping errors. Notably, 50.89% of LANAβlac+ macrophages had detectable expression

of ORF75A, ORF75B and/or ORF75C (Fig. 2D). These studies identify two distinct

transcriptional signatures early after infection in vivo: 1) a high frequency of macrophages

expressing transcripts from the ORF75 locus, 2) with rare cells characterized by lytic

transcription.

MHV68 readily infects J774 macrophages with limited lytic transcription.

We next examined whether these viral outcomes could be observed in a reductionist cell

culture model, assessing MHV68 infection in the murine myeloid-like cell line, J774, compared

to 3T12 fibroblasts, a permissive murine fibroblast cell line [48]. As expected, MHV68 underwent

robust lytic replication in 3T12s with extensive cytopathic effect (CPE) by 72 hours post infection

(hpi). In contrast, MHV68-infected cultures of J774 cells showed minimal evidence of infection,

with no discernable CPE (Fig. 3A) and minimal viral genome replication or virus particle

production above input virus (Fig. 3B-C). Infection with WT MHV68.LANAβlac resulted in a

comparable frequency of LANAβlac+ cells in 3T12s fibroblasts and J774 cells (Fig. 3D-E, Fig.

S3A), demonstrating that J774 cells are fully susceptible to MHV68 infection, initiating

transcription and translation of ORF73 (an immediate early gene), with restricted infectious virus

particle production.

We next analyzed viral transcription in MHV68-infected J774 cells. ORF75A, ORF75B

and ORF75C were readily detected in MHV68-infected J774 cells, albeit at lower levels than

infected 3T12 cells (Fig. 3F, Fig. S3B). ORF75A expression was relatively comparable to

ORF73, with minimal change in expression between 24 and 48 hpi, suggesting constitutive

transcription of ORF73 and ORF75A (Fig. 3F). In contrast, ORF75B and ORF75C transcript

levels were moderately lower than ORF73 at 16 and 24 hpi, with ~4- and ~22 fold increase

between 24 and 48 hpi, respectively (right panel, Fig. 3F), potentially reflecting a subset of J774

cells initiating lytic infection. To assess lytic transcription, we quantified immediate early (IE)

(ORF73), early/late (ORF6), and late gene (M7) expression [49]. ORF73 (and Blac) expression

was reduced ~10 fold in J774 myeloid cells relative to 3T12 fibroblasts (Fig. 3G, Fig. S3C-D),

with ORF6 and M7 transcripts decreased >100 fold (Fig. 3G), consistent with greater deficits in

lytic gene expression at later stages of the lytic cycle in MHV68-infected J774 macrophages.

ORF6 and M7 expression increased ~10x between 24 hpi and 48 hpi (Fig. 3G, right panel),

consistent with some lytic transcription in MHV68-infected J774 cultures.

Only rare MHV68 infected macrophages display lytic gene expression.

We next defined the frequency of cells with lytic gene expression using a series of flow

cytometry-based assays. First, we infected 3T12 and J774 cells with a recombinant WT MHV68

virus, MHV68.HygroGFP (MHV68.H.GFP), that encodes a hygromycin resistance protein-GFP

fusion under the transcriptional control of the human cytomegalovirus IE promoter inserted

between ORF27 and 29b [50]. This virus robustly expresses GFP during active, lytic infection,

identifying cells that initiated lytic transcription and translation. 3T12 cells had a much higher

frequency of HygroGFP+ cells than J774 cells at 18 hpi (~8.2% vs. ~0.5% GFP+ respectively,

Fig. 4A-B, Fig. S3A); a similar low frequency of HygroGFP+ cells was observed in both J774

and RAW264.7 macrophage cell lines (Fig. S4A). Infection with two additional reporter viruses,

encoding either an ORF59GFP gene fusion [51] or a viral cyclin-beta lactamase gene fusion

(vCycβlac), further demonstrated that J774 cells had a greatly reduced frequency of reporter

gene positive events compared to 3T12 fibroblasts (Fig. 4C-D). The reduced frequency of lytic

gene expression in J774 cells was further observed when we analyzed two proteins associated

with lytic replication, the viral regulator of complement activation (vRCA), a protein encoded by

the ORF4 late gene, or phosphorylated histone H2AX (γH2AX), a host protein that undergoes

virus-induced phosphorylation [24]. Whereas ~45% of MHV68-infected 3T12 cells were

vRCA+, γH2AX+, or vRCA+γH2AX+ by 24 hpi, the frequency of MHV68-infected J774 cells that

expressed either of these proteins was significantly lower (~2% cells at 24 hpi, ~8% at 48 hpi)

(Fig. 4E-F, Fig. S4B). Given the low frequency of cells expressing lytic cycle genes, we tested

whether virally-infected J774 cells were capable of long-term growth and survival by purifying

LANAβlac+ J774 cells. Over 14 days, LANAβlac+ J774 cells showed a progressive growth

impairment relative to mock-infected cells, with limited evidence of viral replication (Fig. S4C-I).

These data indicate that MHV68-infected J774 cells rarely undergo lytic gene expression,

mirroring in vivo findings, and suggest the utility of this system to interrogate MHV68-myeloid

cell interactions.

High dose MHV68 infection results in abortive replication.

Our studies to this point focused on infection using an intermediate infectious dose, a

multiplicity of infection (MOI) of 1 plaque forming unit per cell. Since high MOI has been

implicated in overcoming intrinsic antiviral immunity [52, 53], we next compared infection

outcomes at an intermediate (MOI=1) or high inoculum (MOI=10). J774 cells infected with

intermediate MOI had a low frequency of HygroGFP+ cells (~4%); in contrast, infection with high

MOI was characterized by a dramatically increased frequency of HygroGFP+ cells (~92% cells)

(Fig. 5A-B). Similar increases in lytic gene expression were observed following high-dose

infection, quantifying: i) the frequency of ORF59GFP+ cells (Fig. 5C) and ii) the frequency of

cells characterized by actin RNA degradation (Fig. 5D), a hallmark of virus-induced host shutoff

that occurs during lytic replication [6, 54]. Despite these increased measures of lytic infection,

high-dose cultures were characterized by only modest increases in viral RNAs using bulk

cultures (Fig. 5E), with high rates of cell death (Fig. 5F), and minimal evidence of viral DNA

replication or infectious virus production over input (Fig. 5G-H). These data suggest that high-

dose infection of J774 cells results in an abortive lytic infection, characterized by increased early

signs of lytic gene expression, with a failure to complete viral DNA replication or infectious virus

production, potentially as a result of high rates of cell death.

MHV68 infection of macrophages is modulated by viral and host factors.

214   Given that interleukin (IL)-4 induces viral replication and reactivation from latency in

MHV68-infected primary macrophages [37, 55], we tested whether IL-4 pre-treatment was

sufficient to overcome the blocks on lytic replication following infection with an intermediate

dose. J774 cells pre-treated with IL-4 had a greatly enhanced frequency of HygroGFP+ cells

with ~60% GFP+ cells by 48 hpi (Fig. 6A-B), resulting in a time- and concentration-dependent

increase in viral DNA replication relative to untreated cells (Fig. 6C), culminating in increased

infectious virus production (Fig. 6D). These data demonstrate that IL-4 pre-treatment is

sufficient to overcome restricted MHV68 replication in macrophages and demonstrate that J774

cells are fully competent to support MHV68 lytic replication.

223   To begin to define how viral factors regulate macrophage infection, we examined two

viral genes: the viral cyclin (vCyc), dispensable for lytic replication in fibroblasts but critical for

reactivation from latency [56], and the immediate early transactivator Rta, which is essential for

lytic replication and capable of initiating reactivation from latency [57-59]. vCyc-deficient MHV68

infection was associated with a greatly reduced frequency of cells that expressed either vRCA

and/or γH2AX relative to WT MHV68-infected cells (Fig. 6E). Conversely, infection with a virus

that overexpresses Rta (CA-RTA MHV68) was associated with a greatly increased frequency of

cells expressing these lytic cycle indicators (Fig. 6E). These data demonstrate that the vCyc is

necessary to promote lytic gene expression in macrophages and that Rta is sufficient to drive

lytic gene expression in macrophages, illustrating how host and viral factors shape the outcome

of macrophage infection.

DISCUSSION

Myeloid cells, including macrophages, monocytes and dendritic cells, are well-documented

targets of γHV infection [11-27], though there remains little consensus on how infection is

regulated in these cells or why infection results in widely divergent outcomes. Here, we present

an in-depth analysis of MHV68 infection of macrophages, documenting multi-layered regulation

of outcomes. First, macrophages harvested early after MHV68 infection in vivo are

characterized by frequent transcription of the ORF75 locus in the absence of other detectable

viral RNAs. Second, MHV68 induces three divergent outcomes in a reductionist system of

macrophage infection: i) a restricted state of infection, reminiscent of latency, with infrequent

cells initiating lytic infection, ii) an abortive form of lytic replication or iii) full lytic replication.

Third, divergent infection outcomes are readily observed by manipulating virus (multiplicity of

infection, viral mutations) and host factors (cytokine exposure), illustrating how divergent

infection outcomes can occur in a single cell type. These data suggest that MHV68 replication in

macrophages is blocked at two stages, first a block in early transcription, overcome by high MOI

infection, and second, a block at virus DNA replication, overcome by IL-4 pre-treatment (Fig 4).

These studies further suggest a previously unappreciated role for the ORF75 locus as a

potentially important regulator of macrophage infection, and suggest re-evaluation of previous

assumptions (e.g. that macrophages poorly support lytic replication).

By examining the heterogeneity of macrophage infection at single cell resolution, our studies

provide a refined perspective of MHV68 infection in macrophages. These studies unequivocally

identify that MHV68 robustly infects macrophages and are capable of initiating transcription and

translation of at least the ORF73 gene, with only a minor fraction of these cells showing

hallmarks of lytic replication. Whether the majority of LANA+ cells reflect a bona fide latent

population or a previously undescribed, restricted form of infection remains an open question.

Although selective ORF75 transcription has not been reported in the context of MHV68

macrophage infection, transcripts mapping to the ORF75 locus have been observed in some in

vitro latency models, a finding previously attributed to potential detection of unspliced ORF73

transcripts [60-62]. Here, we detected individual ORF75 gene products (i.e. ORF75A, ORF75B,

and ORF75C) by polyA-based scRNA-seq, strongly suggesting that these RNAs represent

ORF75-specific transcripts, not unspliced ORF73 RNA. These findings further suggest that

ORF75 transcription is regulated in additional ways beyond those observed for lytic replication

[46]. Notably, ORF75 transcription has been reported in contexts beyond lytic replication for

EBV BNRF1 (the ORF75 ortholog), where it is expressed in lymphoblastoid cell lines [41], and

for KSHV ORF75, where it is detected in Kaposi's sarcoma lesions [63, 64]. Recent studies

have shown that the KSHV ORF75 promoter is regulated in a cell type-specific manner, active

in endothelial cells but repressed in B cells [40]. Collectively, these findings suggest that the

ORF75 locus may be subject to macrophage-specific regulation, that ORF75 gene products

may have unique functions in macrophages, and that ORF75-targeted interventions (e.g.

vaccination against ORF75-derived T cell epitopes [65-67]) may selectively disrupt macrophage

infection.

Through use of a reductionist system of macrophage infection, we find that MHV68 infection

can result in three divergent outcomes. Using infection with an intermediate infectious dose

(MOI=1), MHV68 readily infects the majority of cells, defined by LANAβlac expression, with only

a minor fraction of cells characterized by lytic protein expression and no measurable viral DNA

replication or infectious virus production; this finding parallels our in vivo analysis. While we can

detect LANAβlac+ J774 cells for up to 14 dpi, whether this in vitro model represents bona fide

latency remains unknown at this time. In striking contrast to intermediate dose infection,

infection with a high virus dose (MOI=10) greatly increases the frequency of cells initiating lytic

cycle transcription, with minimal viral DNA replication infectious virus production and a high rate

of cell death, findings strongly suggestive of abortive replication. The significant induction of

early transcription that occurs with a high MOI suggests that macrophage infection is subject to

at least two sequential levels of regulation: i) a block that limits entry into the lytic cycle

(observed with an intermediate infection dose) that can be overcome by a high MOI, and ii) an

additional block that prevents viral DNA replication and lytic replication following infection with a

high MOI, potentially due to activation of cell death pathways. Despite these barriers to lytic

replication, it is notable that pre-treatment with the cytokine IL-4 induces a high frequency of

lytic transcription, viral DNA replication and infectious virus production, consistent with previous

reports that IL-4 can promote virus replication in macrophages [55].

By investigating the interplay between viral and host factors in macrophage infection, our

studies demonstrate significant heterogeneity in infection outcome in macrophages. These

findings provide an important new perspective on previously published findings and help to

resolve previously ambiguities in the literature. First, the magnitude of MHV68 lytic replication

within bone marrow macrophages or myeloid cells is relatively modest compared to fibroblasts

[25, 68], with lytic antigen detected in a subset of bone marrow-derived macrophages and

RAW264.7 macrophages [30, 37, 55]. Our data are consistent with this and emphasize that

cells initiating lytic transcription represents only a fraction of infected cells. Second, there are

numerous studies that have identified positive and negative regulators of macrophage infection.

MHV68 lytic replication in macrophages is supported by multiple viral proteins (viral protein

kinase [24], vRCA [25], vCyclin, Rta) and host factors (H2AX [38], ATM [69] and apolipoprotein

E [70]). Parasitic infection can further promote macrophage infection through two distinct

mechanisms, either by eliciting IL-4 or IL-13, STAT6-inducing cytokines that directly

transactivate the MHV68 gene 50 promoter [37], or by expanding the number of large peritoneal

macrophages allowing for an increased frequency of infection targets in vivo [71, 72].

Conversely, MHV68 replication in macrophages is limited by type I and type II interferons [73],

IRF1 [74], HDAC1 and 2 [75], liver X receptors [35], and Low-Density Lipoprotein Receptor [36].

In vivo, IFNγ constrains MHV68 reactivation from latency within peritoneal macrophages [76]

and CD4 T cells constrain chronic macrophage infection in the lung [68]. Based on our current

findings, we postulate that these diverse regulatory mechanisms may affect the relative

proportion of cells initiating lytic replication with MHV68, or may fundamentally alter the

susceptibility of infection across all cells. Defining how these regulatory mechanisms influence

the frequency and outcome of macrophage infection at the single-cell level remains an

important future goal.

This study has important limitations. First, our studies of primary in vivo infection focus on a

single, early timepoint. Conservatively, we can conclude that the high frequency of LANA+

macrophages with detectable ORF75 transcripts is not explained by the current paradigm of

lytic or latent transcription (e.g. exemplified by [49]) and suggests additional mechanisms of

transcriptional regulation within the ORF75 locus beyond that in fibroblasts [46]. Second, our in

vitro studies use the J774 mouse myeloid cell line, a reductionist in vitro system to study

MHV68-macrophage interactions. Though these cells demonstrate parallels with primary

macrophages (e.g. infrequent lytic replication at baseline that can be potently enhanced by IL-4

pretreatment), whether the precise molecular mechanisms engaged in these cells reflect one or

more primary macrophage subsets remains to be determined. This is an important, but

challenging question, given that macrophage subsets in vivo are known to respond differently to

infection (e.g. large versus small peritoneal macrophages differ in their infection frequency with

MHV68 early after infection [71, 72]; infection of marginal zone macrophages is thought to

precede B cell latency [14, 33], whereas subcapsular sinus macrophages limit infection [34]).

Despite these limitations, our results are consistent with previous studies demonstrating lytic

protein expression in a subset of bone marrow-derived macrophages and RAW264.7

macrophages [30, 37, 55], and revealing that high MOI can overcome genetic requirements for

lytic gene expression at a lower MOI [38]. Finally, it is important to recognize that the outcome

of γHV-myeloid cell infection may be affected by the mode of virus entry. While our experiments

test macrophage infection using cell-free virus, additional modes of infection have been reported

(i.e. virus internalization via epithelial cell presentation [31], or infection through an Fc receptor-

dependent mechanism [30]). These factors would likely be influenced by the route and stage of

infection and host immune status.

In total, our studies provide critical new insights into how γHV infection of myeloid cells is

regulated, demonstrating unanticipated transcription during primary infection in vivo and three

divergent infection outcomes which can be readily manipulated by virus and host factors. Our

studies highlight how bulk analyses and subsequent conclusions drawn from these analyses are

insufficient to describe the heterogeneity of MHV68 infection in myeloid cells – a conclusion that

could apply to γHV interactions beyond the myeloid compartment. These data redefine a

conceptual framework to study myeloid cells as a γHV target and emphasize the knowable

complexity of γHV-myeloid cell interactions.

METHODS

Viruses and Tissue Culture

Mouse 3T12 fibroblasts (ATCC CCL-164), mouse J774A.1 macrophages ("J774"; ATCC TIB-67), and mouse RAW264.7 macrophages (ATCC, TIB-71) were cultured in complete DMEM (cDMEM): Dulbecco's modified Eagle medium (DMEM; Life Technologies) supplemented with 5% and 10% fetal bovine serum (FBS; Atlanta Biologicals) respectively, 2 mM L-glutamine, 10 U/mL penicillin, and 10 µg/mL streptomycin sulfate. Cells were cultured at 37°C with 5% CO_2. Wild-type (WT) MHV68, MHV68.LANA::βlac, MHV68.H.GFP [50], MHV68.ORF59.GFP [51], MHV68.vCycKO [56], MHV68.CA-RTA [77] and MHV68.vCyc::blac viruses were grown and prepared as described previously [43, 44, 50], with virus titers determined from at least three independent plaque assays. MHV68.vCyc::blac was created through BAC recombineering, introducing a C-terminal fusion with the blac gene downstream of the endogenous vCyc (ORF72) gene (manuscript in submission, Noell et al).

Infection

All in vitro infections were based on live cell counts with the TC20 Automated Cell Counter (Bio-Rad) with Trypan Blue dye (Bio-Rad, Cat. No. 145-0021). Viral stock was diluted with cDMEM for viral inoculum with a multiplicity of infection (MOI) of 1 or 10 PFU/cell. Cells were incubated at 37°C with 5% CO_2 for 1 hour, with rocking every 15 min, before viral inoculum was removed and replaced with 1mL of cDMEM. For qPCR analysis of viral DNA replication and for beta-lactamase staining and flow cytometric analysis of infected cells, viral inoculum was left on cells after 1 hr infection time. For IL-4 treatments, media containing recombinant mouse IL-4 (PeproTech, Cat. No. 214-14) was added to cells 16 hours before infection. Following infection, media containing 0, 10 or 100ng/mL of fresh IL-4 containing media were added respectively. Samples were harvested at the indicated hours post-infection (hpi) through either cell scraping (for DNA isolation) or through incubation in TRIzol (for RNA isolation).

Plaque Assay

Plaque assay quantification of viral titer was performed using 3T12 cells. Cells were

plated in 12-well plates at 8.5×10^4 cells per well one day prior to infection. Viral samples were

diluted 10-fold in 5% cDMEM. An internal standard was included for each infection to ensure

reproducible sensitivity for each plaque assay. Cells were incubated with virus for 1 hour at

37°C with 5% CO_2. Plates were rocked every 15 min. Cells were then covered with an overlay

composed of a 1:1 mix of 10% cDMEM and carboxymethyl cellulose (CMC; Sigma, Cat. No. C-

4888) supplemented with Gibco™ Amphotericin B (Thermo Fisher Scientific, Cat. No.

15290018). Cells were incubated for 8 days before staining with 0.5% methylene blue and

plaques were counted.

RT-qPCR

RNA was isolated from infected cells harvested at indicated times by 10-minute

incubation in TRIzol® reagent (Thermo Fisher Scientific, Cat. No. 15596026), followed by

TURBO™ DNase (Invitrogen, Cat. No. AM2238) treatment according to manufacturer's

protocols. RNA amplification and removal of DNA was confirmed by PCR amplification of the

control host gene, 18S, in the presence or absence of reverse transcriptase. RNA presence and

absence of DNA was confirmed with RT-PCR and PCR respectively. RT-PCR was performed

using the OneStep RT-PCR kit (Qiagen, Cat. No. 210212) with the following conditions: (i) 50°C

for 30 min, (ii) 95°C for 15 min, (iii) 40 cycles of 94°C for 30 sec, 52°C for 30 sec, and 72°C for

30 sec, (iv) 72°C for 10 min, and (v) hold at 4°C. PCR was performed using *Taq* DNA

polymerase (Qiagen, Cat No. 201205) with the following conditions: (i) 95°C for 5 min, (ii) 40

cycles of 94°C for 30 sec, 52°C for 30 sec, and 72°C for 30 sec, (iii) 72°C for 10 min, and (iv)

hold at 4°C. RNA samples that showed no product following PCR amplification were deemed

DNA-free, and then converted to cDNA using random primers (250ng/uL) (Invitrogen, Cat. No.

48190011) and SuperScript II reverse transcriptase (Invitrogen, Cat. No. 18064014) following

the manufacturer's protocol. 100 nanograms of cDNA was used for qPCR analysis of the

identified genes (Quantitect Primer Assay, Qiagen) using the iQ SYBR green supermix (BioRad

Cat. No.1708880) with the following conditions: (i) 95°C for 3 min, (ii) 40 cycles of 95°C for 15

sec, 60°C for 1 min, and (iii) 95°C for 15 sec, 60°C for 1 min, and 95°C for 15 sec. Amplification

of viral genes was normalized to murine 18S expression to calculate the relative difference of

target gene expression using the Pfaffl method, as previously described [6, 78] : Target primer

efficiency^TargetΔCt/18S primer efficiency^$18S\Delta Ct$. A single product for each target was

confirmed by melt curve analysis. PCR primers are listed in **Table 1.**

Quantitative PCR for quantification of viral DNA replication

Infected cells were harvested by cell scraping at time (hpi) indicated. Harvested cells

underwent three freeze/thaw cycles and DNA was isolated using the DNeasy Blood and Tissue

Kit (Qiagen, Cat. No. 69504), with an overnight proteinase K incubation and heat inactivation at

56°C. DNA was normalized to a concentration of 20 ng/uL in molecular grade water. qPCR

analysis was done on 100ng of DNA using a LightCycler 480 Probe Master-Mix kit (Roche, Cat.

No. 04707494001) and a primer and probe set specific to MHV68 glycoprotein B (gB) to

quantify the number of viral genome copies. Viral genome copies/cell concentration was

determined by dividing gB quantity over NFAT5 quantity. Primers and probes listed in Table 1. A

gB or nuclear factor of activated T cells 5 (NFAT5) [79, 80] standard curve was generated using

a gB or NFAT5 plasmid dilution series ranging from 10^2 to 10^{10} copies diluted in background

DNA, with a limit of detection (LOD) of 100 copies.

Single-cell RNA Sequencing

To characterize MHV68 transcription during primary, acute peritoneal macrophage

infection *in vivo*, female C57BL/6 mice were infected with 1x10^6 PFU by intraperitoneal injection.

Peritoneal cells were harvested 16 hours post-infection, stained with the live/dead discrimination

dye Zombie Near-IR (Zombie NIR™, BioLegend, Cat. No. 423105), F4/80-AlexaFluor 674

(clone BM8, 1:200 dilution) in the presence of Fc receptor blockade (clone 2.4G2), and CCF2-

AM (Thermo Fisher, Cat. No. K1023). Cells were sorted on a MoFlo Astrios, purifying live,

single, F4/80+ cells that either had CCF2 cleavage (indicating virus infection, defined by

expression of the LANA::blac fusion protein) or lacking CCF2 cleavage (indicating cells that

were not infected or failed to express the LANA::blac fusion protein). Single cell RNA-seq was

prepared using the Chromium 3' V3 library kit (10X Genomics). Cell processing and sequencing

was performed by the University of Colorado Cancer Center Genomics Shared Resource

(RRID: SCR_021984). Resulting data were processed as follows: Cell Ranger (v6.0.1) [81] was

used to process the fastq files to cell and gene count tables using unique molecule identifiers

(UMIs) with the include-introns parameter. Because of the difficulties counting viral reads, this

was performed in a two-pass manner. In the first pass, reads were aligned to a chimeric

genome of mouse mm10 (GENCODE M23 gene annotations) and MH636806.1 (no gene

annotations, only positive and negative strand alignment). The viral counts were stored in

metadata and removed from the counts matrix. Reads mapping to MH636806.1 in step 1 were

aligned to the same chimeric genome but in this case, the viral transcriptome as well as specific

intergenic regions were annotated. The intergenic regions were determined as the strand-

specific sequences between genes with 1 base pair of padding on both ends. To minimize reads

overlapping with multiple viral genes, only the first 70bp of each read were aligned and counted.

The resulting counts matrix of viral alignment was appended to the host gene counts from step

1.

The Seurat (v4.0.4) [82] pipeline was used for downstream quality control and analysis.

Cellranger-filtered data was read into Seurat. Host genes were removed if identified in fewer

than 10 cells while viral gene and intergenic regions were removed if found in no cells. Cells

were removed if they expressed < 50 genes or viral regions, < 5000 UMIs, > 5 % total UMIs

from mitochondria, or > 70000 UMIs. The filtered data was normalized by dividing gene counts

by total counts per cell and multiplied by 10,000 followed by natural-log transformation. The top

2,000 most variable genes were scaled with total UMI and percentage mitochondria regressed

out. These were used to calculate principal components (PCs) with the top 20 used to perform

Uniform Manifold Approximation and Projection (UMAP) and determining the k-nearest

neighbors and clustering.

Clusters were identified by over-representation analysis and examining the top enriched

markers. Canonical cell type markers were plotting in UMAP space and using dot plots. Dot

plots, bar plots, scatter plots, and violin plots were generated using Seurat and ggplot2 R

packages (v3.4.0) [83].

Flow Cytometry

3T12 and J774 cells were infected with MHV68 LANAβlac or MHV68.H.GFP at an MOI

of 1 PFU/cell and harvested at the indicated time points followed by staining for flow cytometric

analysis. Samples treated with the recombinant murine IL-4 were given 10% cDMEM with

10ng/mL of IL-4 eighteen hours prior to infection and replaced with fresh media containing IL-4

immediately following infection. Cell suspensions were stained with LIVE/DEAD fixable Near-IR

Dead Cell stain kit (Invitrogen, Cat. No. L10119) at a 1:1000 dilution in BSS wash (111mM

Dextrose, 2mM KH_2PO_4, 10mM Na_2HPO_4, 25.8 $CaCl_2 \cdot 2H_2O$, 2.7mM KCl, 137mM NaCl, 19.7mM

$MgCl \cdot 6H_2O$, 16.6mM $MgSO_4$). Cells were then stained for CCF2 or protein detection. Cells for

CCF2 detection were stained with CCF2-AM (Thermo Fisher, Cat. No. K1023) at a 1:10 dilution

following the manufacturer's protocol. To detect proteins, cells were stained with a Fc receptor

block (Fc Shield, Tonbo Biosciences, Cat. No. 70-0161), a rabbit antibody against the MHV68

ORF4 protein (vRCA, 1:400 dilution) (gift of the Virgin Lab, Washington University St. Louis [6,

84]) labeled with a Zenon R-phycoerythrin anti-rabbit IgG reagent (Invitrogen, Cat. No. Z25307),

per manufacturer's recommendations, and a AF647-conjugated anti-mouse phospho-histone

H2AX antibody (1:800 dilution) (clone JBW30). Samples were fixed in 1% paraformaldehyde

prior to analysis using the Bio-Rad ZE5/YETI Cell Analyzer or an Agilent Novocyte Penteon flow

cytometer. To detect actin mRNA, cells were stained with buffers and probes using the

PrimeFlow RNA assay kit (ThermoFisher), as previously published [6]. All flow cytometry

experiments included unstained, single-stain and full minus one controls to define background

fluorescence, fluorescent signal spread and compensation.

Light microscopy

At the indicated time points, 6-well plates of 3T12 or J774 cells were imaged on a Nikon

Eclipse Ti2 inverted light microscope equipped with Iris 15 camera (Photometris) and use of NIS

elements software. Plates were mounted on a stage heated to 37°C. A 10x phase objective was

used to capture brightfield images. Immediately following image acquisition, samples were

harvested and processed for flow cytometric analysis.

Statistical Analysis and Software

Data analysis, graphing and statistical analyses were performed using GraphPad Prism

(version 9.4.0; GraphPad Software, San Diego, California USA, www.graphpad.com). Flow

cytometry data were analyzed using FlowJo (version 10.8.1. Ashland, OR: Becton, Dickinson

and Company; 2022). scRNA-seq data processing and analysis were done as described above.

Statistical significance was tested by unpaired, nonparametric, Mann-Whitney t test or 1-way

ANOVA for comparing replicate values from experiments. Raw and processed single cell RNA-

Seq data is available through NCBI GEO (GSE233202).

Cell Sorting

J774 fibroblasts were infected at an MOI=1 with for one hour, after which inoculum was

removed. Cells were harvested at 48 hours post-infection (hpi) and stained with β-lactamase

substrate as previously reported [44] using the CCF2-AM substrate (Invitrogen, Life

Technologies). Cells were subjected to fluorescence activated cell sorting (FACS), with cells

gated on singlet cells that were positive for the cleaved CCF2-AM substrate (subsequently

referred to LANAβlac+ sorted cells). Cleavage of CCF2-AM was defined as positive

fluorescence in the 405-448/59nm emission channel obtained in WT MHV68.LANAβlac,

compared to parental WT MHV68 infected cultures with no beta-lactamase fusion gene, used to

define background fluorescence. β-lactamase positive cells were sorted using a Astrios EQ

(Beckham Coulter). Cells were either plated onto 6-well plates at 5×10^5 cells/mL, collected for

quantitative PCR or plaque assay measurement. The plated CCF2+ J774 cells were cultured at

37°C with 5% CO_2, grown in complete DMEM (cDMEM): Dulbecco's modified Eagle medium

(DMEM; Life Technologies) supplemented with 10% fetal bovine serum and passaged upon

confluency (typically every 3-4 days). Upon passaging, cells and supernatant were collected

followed by resuspension in 1mL of 10% cDMEM. Cell counts were measured by BIORAD

TC20 automated cell counter. At 7 days post sort (dps), one 6-well of cells was collected and

resuspended in 1mL of 10% cDMEM and flow cytometry for CCF2 detection was performed with

700uL of the resuspended cells. At 14 dps, one 6-well of cells was collected and resuspended in

1mL of 10% cDMEM and flow cytometry for CCF2 detection and viral protein vRCA and H2AX

was performed with 700uL of the resuspended cells (See Flow Cytometry Methods Section). At

both 7 and 14 dps 300uL of the 1mL collected cells were used for qPCR and plaque assay

analysis.

Ethics Statement

All animal studies were performed in accordance with the recommendations in the Guide

for the Care and Use of Laboratory Animals of the National Institutes of Health. Studies were

conducted in accordance with the University of Colorado Denver/Anschutz Institutional Animal

Use and Care Committee under the Animal Welfare Assurance of Compliance policy

(assurance no. D16-00171). All procedures were performed under isoflurane anesthesia, and all

efforts were made to minimize suffering.

Figure Legends

Figure 1. Single cell RNA sequencing analysis of acute MHV68 macrophage infection.

C57BL/6J mice were infected with 1x10^6 PFU of WT MHV68.LANAblac virus. MHV68-infected,

LANAblac+ peritoneal cells were sort-purified at 16 hpi and subjected to scRNA-seq analysis.

(A-B) Visualization of cell types based on lineage marker genes depicted in panel B. **(C)**

Quantification of cell subsets defined by scRNA-seq. **(D)** UMAP visualization of cells with at

least 1 viral UMI detected per cell identified by purple. **(E)** Frequency of cells stratified by the

fraction of viral UMIs out of total UMIs detected per cell. 56 cells had >0.1% of total UMIs

contributed by viral UMIs, including 23 of 9101 macrophages (0.25%), 25 of 25 macrophages

with lytic gene expression (100%) and 8 of 70 dendritic cells (11.43%). Data are from 9,537

cells. Supporting data in **Fig S1-S2**.

Figure 2. Single cell RNA sequencing analysis of acute MHV68 macrophage infection

reveals two distinct transcriptional profiles. Analysis of viral gene expression defined by

scRNA-seq in macrophage subsets from **Figure 1. (A-C)** Frequency of cells with detectable

reads mapping to MHV68 genes, comparing (A) macrophages (82.79% viral UMI+), (B)

macrophages with proliferative gene signature (52.08% viral UMI+), or (C) macrophages with

lytic gene expression (100% viral UMI+). Viral genes are indicated on the x axis, arranged from

left to right based on the published MHV68 genome, with the frequency of cells positive for each

viral gene depicted on the y axis. To facilitate comparison of gene expression across the viral

genome and cell populations, circles are color-coded to identify values that are in the indicated

ranges (defined in key, panel A). Dotted lines indicate different thresholds of viral gene

positivity, empirically defined. For panel C, only genes with >75% positivity were color-coded,

given the limited number of cells and high viral positivity. **(D)** Frequency among macrophages

defined by detection of ORF73, ORF75A, ORF75B, and ORF75C RNAs, enumerating all

potential gene combinations for macrophages from panels A-C.

Figure 3. MHV68 readily infects J774 macrophages with limited lytic transcription.

Analysis of MHV68 infection (WT MHV68.LANAblac, MOI=1 PFU/cell) comparing 3T12

fibroblasts and J774 macrophages. **(A)** Representative brightfield images of infected 3T12 (top)

or J774 cells (bottom) early (24 hpi, left) or late (72 hpi, right) post-infection. 3T12 cells showed

cytopathic effect (CPE), including cell rounding (24 hpi) and monolayer destruction (72 hpi), in

contrast to J774 cells. **(B)** Viral DNA quantitation defined by qPCR. **(C)** Virus replication defined

by plaque assay. **(D-E)** Frequency of MHV68-infected, LANAβlac+ cells defined by flow

cytometry, detected by CCF2 substrate cleavage, analyzing single, viable cells. Samples

infected with WT MHV68 (left, panel E) define background signal and do not encode LANAβlac.

(F-G) Quantitation of viral gene expression defined by qRT-PCR, comparing 3T12 and J774

cells at 24 hpi (left) or J774 cells at 24 and 48 hpi (right). Flow cytometric analysis focused on

single, viable cells. Data depict mean ± SEM, from 3 independent experiments with 3 biological

replicates per experiment (B, D, F-G) or 2 (t=0) or 6 (t=24, 48 hpi) independent experiments (C),

with individual symbols representing independent biological samples. Statistical analysis was

done using unpaired t test, with statistically significant differences indicated, *$p<0.05$, **$p<0.01$,

***$p<0.001$, ****$p<0.0001$.

Figure 4. Only rare MHV68 infected macrophages display lytic gene expression. Analysis

of MHV68 infection and lytic cycle profiles with either MHV68.H.GFP or MHV68.LANAblac

(MOI=1 PFU/cell) comparing 3T12 fibroblasts and J774 macrophages. **(A)** Flow cytometric

analysis of virus-expressed GFP in 3T12 or J774 cells 18 hpi. Data depict the frequency of cells

with GFP expression, an indicator of MHV68.H.GFP infection. Background fluorescence defined

by WT MHV68.LANAblac-infected cells (left column), without the addition of the CCF2

fluorescent substrate. **(B)** Frequency of HygroGFP+ cells defined by flow cytometry, as in panel

A. **(C)** Frequency of vCycblac+ cells defined by flow cytometry. **(D)** Flow cytometric analysis of

lytic infection-associated proteins, vRCA and gH2AX in 3T12 (top) or J774 (bottom) cells,

comparing mock (left) or WT MHV68.LANAblac-infected cells. Data for 3T12 cells at 48 hpi was

not assessed due to extensive cell destruction. **(E)** Quantification of the frequency of infected

cells with protein expression associated with lytic infection defined by flow cytometry, as in

panel D. Flow cytometry data focus on single, viable cells defined using gating strategies in **Fig.**

S3A, S4B. Data show mean ± SEM from 3 experiments, with 2 (B,E) or 3 samples (C) per

experiment. Statistical analysis done using Mann-Whitney test, with statistically significant

differences as indicated, **p<0.01, ****p<0.0001.

Figure 5. High dose MHV68 infection results in abortive replication. MHV68 outcomes in

J774 macrophages comparing intermediate (MOI=1 PFU/cell) or high dose (MOI=10) infection.

(A) Flow cytometric analysis of GFP in MHV68.LANAblac- (left; negative control with no CCF2

substrate) or MHV68.H.GFP-infected cells (right) at MOI=1 (top) or MOI=10 (bottom). **(B)**

Frequency of GFP+ cells quantified by flow cytometry, as in A. **(C)** Frequency of ORF59GFP+

J774 cells quantified by flow cytometry at 24 or 72 hpi. **(D)** Flow cytometric analysis of actin

mRNA levels 24 hpi in mock or WT MHV68-infected cells with individual rows showing

independent biological samples. **(E)** Quantitation of viral RNA expression by qRT-PCR

standardized to 18S RNA, comparing intermediate or high MOI (MHV68.LANAblac) at the

indicated timepoints. **(F)** Frequency of MHV68.HygroGFP-infected single cells that are viable,

quantified by flow cytometry. **(G)** Quantitation of viral DNA by qPCR in J774 (left) or 3T12 cells

(right) as a function of MOI (MHV68.LANAblac). **(H)** Infectious virus production quantified by

plaque assay, comparing 3T12 cells (MOI=1) with J774 cells (MOI=1 or 10). Flow cytometry

data analyzed single, viable cells (A-D) or single cells (F), with gating strategy in **Fig. S3**. Data

show mean ± SEM, with individual symbols showing independent biological samples from 2-3

independent experiments with 2-3 biological replicates per experiment. Statistical analysis was

done using Mann-Whitney test with statistically significant differences indicated, **p<0.01,

***p<0.001.

Figure 6. MHV68 infection of macrophages is modulated by host and viral factors. All

infections done with MOI=1. **(A-D)** IL-4 effect on MHV68 infection and genome replication in

J774 macrophages. IL-4 cultures were pretreated with IL-4 for 16-18 hours at the indicated

concentration, with fresh IL-4 replenished after infection. **(A)** Flow cytometric analysis of

MHV68.H.GFP infection in 3T12 (top), untreated J774 (middle) or IL4-treated J774 cells at 24

(left) or 48 hpi (right), analyzing single, viable cells. **(B)** Frequency of GFP+ cells quantified by

flow cytometry, as in A. **(C-D)** IL-4 effect on viral DNA replication **(C)** and infectious virus

production **(D)** in WT MHV68-infected J774 cells. **(E)** Frequency of J774 cells expressing

markers of lytic infection (gH2AX, vRCA) quantified by flow cytometry, comparing WT, vCycKO

or CA-RTA MHV68 at 24 or 48 hpi. Flow cytometric analysis focused on single, viable cells.

Data depict mean ± SEM, with individual symbols representing independent biological samples

from 2-3 independent experiments with 2-3 biological replicates per experiment (B,C,E) or from

1-2 independent experiments with 3 biological replicates measured (D). Statistical analysis used

(B) unpaired t test, (C) one-way ANOVA comparing samples collected at the same timepoint

untreated versus cytokine treated, or (E) one-way ANOVA comparing samples collected at the

same timepoint with WT infection (nonparametric Kruskal-Willis test subjected to Dunn's

multiple comparison test), with statistically significant differences indicated, *p<0.05, **p<0.01,

****p<0.0001.

Table 1: *Oligonucleotides for RT-PCR, PCR, and qPCR analysis*

Name	Accession number and Genome Position *	Sequence	Purpose
gB Forward Primer	MHV68: 17873…17892	5'—GGC CCA AAT TCA ATT TGC CT—3'	TaqMan qPCR for MHV68 genome
gB Reverse Primer	MHV68: 17924…17943	5'—CCC TGG ACA ACT CCT CAA GC—3'	TaqMan qPCR for MHV68genome
gB Probe	MHV68: 17896…17921	5'—ACA AGC TGA CCA CCA GCG TCA ACA AC—3'	TaqMan qPCR for MHV68genome
NFAT5 Probe	NC_000074.7 (108020072..108106149)	5'—ACG AGG TAC CTC AGT GTT—3'	TaqMan qPCR for murine genome
NFAT5 Forward	NC_000074.7 (108020072..108106149)	5'—CAT GAG CAC CAG TTC CTA CAA TGA T—3'	TaqMan qPCR for murine genome
NFAT5 Reverse	NC_000074.7 (108020072..108106149)	5'—TGC TTT GGA TTT CGT TTT CGT GAT T—3'	TaqMan qPCR for murine genome
18S Forward Primer	Accession Number: NR_003278	5'— AGA TCA AAA CCA ACC CGG TGA—3'	RT-PCR and PCR for murine 18S gene
18S Reverse Primer	Accession Number: NR_003278	5'—GGT AAG AGC ATC GAG GGG GC—3'	RT-PCR and PCR for murine 18S gene
ORF73 SYBR G – Forward Primer	MHV68: 104092…104073	5'— GAG CCC CCT ACA GAG CCC CC —3'	SYBR Green qPCR for MHV68ORF73 gene
ORF73 SYBR G- Reverse Primer	MHV68: 103981…104000	5'— CAC CTT GCT CTG CAC CGG CA —3'	SYBR Green qPCR for MHV68ORF73 gene
ORF6 SYBR G – Forward Primer	MHV68: 11699…11783	5'— ATG TTG TCA GCA CCC ATG AA—3'	SYBR Green qPCR for MHV68ORF6 gene
ORF6 SYBR G – Reverse Primer	MHV68: 11802…11783	5'— AAG GGC AGT AGC TGG TCA GA—3'	SYBR Green qPCR for MHV68ORF6 gene
M7 SYBR G – Forward Primer	MHV68: 70109…70128	5'—GCT CCT GCT GAC ACA TCA GA—3'	SYBR Green qPCR for MHV68M7 gene
M7 SYBR G – Reverse Primer	MHV68: 70185…70166	5'—TGG GGT TTG GAC TCT GTA GG—3'	SYBR Green qPCR for MHV68M7 gene
MHV68 ORF75A iQ SYBR - Forward Primer	MHV68: 114964…114945	5'—ACC TGA CAC CCC AAG AAC AG—3'	SYBR Green qPCR for MHV68ORF75A gene
MHV68 ORF75A iQ SYBR - Reverse Primer	MHV68: 114834…114853	5'—GAG CAC TTT TGG TGG AGA GC—3'	SYBR Green qPCR for MHV68ORF75A gene
MHV68 ORF75B iQ SYBR - Forward Primer	MHV68: 113267…113248	5'—CAG CCT CTC AAC CTT TCC AG—3'	SYBR Green qPCR for MHV68ORF75B gene

MHV68 ORF75B iQ SYBR - Reverse Primer	MHV68: 113093…113112	5'—ATA GGA GCC ACC GTT GAT TG—3'	SYBR Green qPCR for MHV68.ORF73βla ORF75B
MHV68 ORF 75C iQ SYBR - Forward Primer	MHV68: 109930…109911	5'—AAG ACA CAG AGG CTG GGA GA—3'	SYBR Green qPCR for MHV68ORF75C gene
MHV68 ORF75C iQ SYBR - Reverse Primer	MHV68: 109758…109777	5'—GAC GGG TAG ACG TGG TGA CT—3'	SYBR Green qPCR for MHV68ORF75C gene
BLAC SYBR G – Forward Primer	MHV68.ORF73βla: 104867…104886	5'—GCT ATG TGG CGC GGT ATT AT—3'	SYBR Green qPCR for MHV68.ORF73βla Blac gene
BLAC SYBR G – Reverse Primer	MHV68.ORF73βla: 104701…104720	5'—AAG TTG GCC GCA GTG TTA TC—3'	SYBR Green qPCR for MHV68.ORF73βla Blac gene

632   MHV68 genome coordinates refer to NC_001826. In the case of MHV68.ORF73βla Bla gene,

633   sequences correspond to NCBI GenBank: MH636806.1, the complete genome sequence of

634   MHV68 containing the ORF73bla gene fusion.

Supplemental Figure Legends

Figure S1. Single cell RNA sequencing analysis of acute MHV68 macrophage infection.
Data correspond to **Figure 1**. (A) Visualization of gene expression based on UMAP topography in Fig. 1A. Maximal expression is designated by purple, with absence of detection designated by gray. Each panel depicts a distinct lineage-defining gene used to identify cell types in Fig. 1, with signature genes for each cell type as follows. Macrophage: Adgre1 (encodes the F4/80 protein), Itgam (encodes the CD11b protein). Macrophage proliferative: Top2a, Mki67. Macrophage lytic: ORF54, ORF7. B cell: Cd19, CD79a. Monocyte: Ly6c2, Ccr2. Dendritic cell: Itgax, H2-Ab1. T cell: Cd3d, Thy1. (B-E) scRNA-seq quality control metrics across cell subsets, including (B) UMI Count, (C) Features, (D) % mitochondrial reads, and (E) S score, a proliferation-associated gene signature.

Figure S2. Single cell RNA sequencing analysis of acute MHV68 macrophage infection analyzed using a higher number of cell clusters. Data correspond to **Figure 1**, with scRNA-seq analyzed as 14 clusters. **(A)** UMAP visualization of clusters. **(B)** Visualization of the frequency and magnitude of host gene expression across clusters, with cell type designation indicated on right. **(C-F)** Quality control metrics across the 14 clusters, including (C) UMI Count, (D) Features, (E) % mitochondrial reads, and (F) S score, a proliferation-associated gene signature. **(G)** UMAP visualization of cells with at least 1 detectable viral UMI per cell identified in red. **(H)** Frequency of cell subsets stratified by the frequency of viral UMIs of total UMIs detected per cell. **(I-J)** Percentage of viral UMIs out of total UMIs across clusters (panel I, y axis range from 0-50%), or (J) which is the same as S2I but excluding 56 cells with viral UMIs > 0.1%. Data are from 9,537 cells, with clusters containing between 25-2,532 cells. Panel G is included to provide a frame of reference and depicts the same data as in Figure 1D.

Figure S3. Flow cytometry gating strategy and viral gene expression during MHV68

infection. (A) Flow cytometry gating strategy to analyze MHV68-infected cells, in support of

Figure 3E. 3T12 cells infected with MHV68 LANAblac or WT.MHV68.H.GFP were analyzed by

flow cytometry, using sequential focused analysis (i.e. "gating") to consider cells that are single

cells (defined sequentially as "Singlets by SSC" and "Singlets by FSC") that are viable ("Live

cells") defined by exclusion of a dye that identifies dead cells. Live cells were then analyzed for

GFP fluorescence as measured by the GFP detection channel. Cells infected with

MHV68.LANAblac, without additional of the CCF2 fluorescent substrate, identified background

fluorescence, in contrast to samples infected with MHV68.H.GFP which encoded GFP. Controls

included single-stained samples with only one stain to define the bounds of a gate for the

fluorophore of interest. **(B-D)** Viral gene expression defined by qRT-PCR in 3T12 or J774 cells

infected with WT MHV68.LANAblac (MOI=1) at 16 hpi (B,D) or 24 hpi (C) in support of **Figure**

3F-G. Data show mean ± SEM from 3 independent experiments with 3 biological replicates

measured in technical triplicates per experiment (depicted by individual symbols) (B,D) or from 1

experiment with 3 biological replicates measured in technical triplicates, with ORF73 data from

Fig. 2B (C). Statistical analysis was done using unpaired t test with statistically significant

differences as indicated. ** p<0.01, ***p<0.001, ****p<0.0001. ns, no statistical significance

detected.

Figure S4. Gating of flow cytometry analysis of MHV68 LANAβlac infected J774 cells for

lytic protein markers. (A) Analysis of GFP expression in 3T12 fibroblasts, J774 and

RAW264.7 macrophage cell lines at 24 or 48 hpi, following infection with MHV68.H.GFP

(MOI=1). **(B)** Gating strategy to analyze vRCA and gH2AX expression in MHV68-infected J774

cells, in support of **Figure 4C**. Cells were analyzed by flow cytometry, using sequential focused

analysis (i.e. "gating") to consider cells that are single cells (defined sequentially as "Singlets by

SSC" and "Singlets by FSC") that are viable ("Live cells") defined by exclusion of a dye that

identifies dead cells. Live cells were then analyzed for antibody binding for lytic protein markers.

Gate boundaries were defined using a combination of "single stain" samples, containing only the

fluorophore of interest, and "full minus one" controls, lacking the antibody in the fluorescent

channel of the excluded fluorophore (not shown). **(C-I)** Analysis of MHV68-infected, FACS-

purified J774 cells over time. **(C)** Experimental overview with J774 macrophages infected with

MHV68.LANAblac (MOI=1), followed by sorting for LANAblac+ cells defined based on CCF2

substrate cleavage. Mock and LANAblac+ sorted cells were cultured and analyzed over 14

days. **(D)** Total cell counts over the course of the study, with each vertical line indicate a

subsequent cell passage. **(E)** Flow cytometric analysis of LANAβlac expression defined by

cleavage of the CCF2 fluorescent beta-lactamase substrate, analyzing single, viable cells, with

values in the right gate defining the frequency of cells with CCF2 substrate cleavage. **(F)**

Quantification of LANAβlac+ cell frequency defined by flow cytometry, as in panel E. **(G)** Flow

cytometric analysis of MHV68 lytic cycle-associated proteins (vRCA, gH2AX) at 14 dps,

comparing mock (left) or CCF2-sorted, LANAblac+ cells (right). Data depict events that were

single, viable cells, with values defining the frequency of cells with vRCA and/or phosphorylated

host γH2AX, indicative of lytic MHV68 infection, as in panel B. **(H)** Frequency of vRCA+

gH2AX+ cells defined by flow cytometry, as in panel G. **(I)** Quantitation of viral DNA by qPCR.

Dotted line indicates mean 3T12 viral genome copies/cell at 72 hpi. Data show mean ± SEM

from 3 independent experiments with 3 biological replicates per experiment (depicted by

individual symbols) (A) or from 2 independent experiments.

Acknowledgments:

We thank Dr. Brian Russo's lab, especially Jenna Vickery for the use and training of their Nikon

Eclipse Ti2 inverted light microscope, and Christine Childs, Kristina Terrell and Dmitry Baturin of

the University of Colorado Cancer Center Flow Cytometry Shared Resource for training and

assistance with flow cytometry. We thank Dr. Laurie Krug and Dr. Craig Forrest for generously

sharing the ORF59.GFP recombinant virus. R.E.K. was supported by the Molecular Biology T32

pre-doctoral training grant (NIH 5T32GM136444). C.G. was supported by the University of

Colorado School of Medicine RNA Bioscience Initiative Summer Internship Program. This study

was supported by National Institutes of Health grant R01 AI157201 awarded to LvD and ETC,

with additional support by the National Institutes of Health Cancer Center Support Grant

(P30CA046934), including assistance from the Bioinformatics and Biostatistics Shared

Resource (RRID: SCR_021983), Flow Cytometry Shared Resource (RRID: SCR_022035), and

Genomics Shared Resource (RRID: SCR_021984). Generative AI was not used in preparation

of this manuscript.

Contributor Roles

Conceptualization: G.V., E.T.C., L.F.v.D.

Methodology: D.G.O.

Validation: G.V., K.S.N., A.T., R.E.K., E.A.A., S.C., C.G., M.D., B.D.G.

Formal analysis: A.G., E.T.C.

Investigation: G.V., K.S.N., A.T., R.E.K., E.A.A., S.C., C.G., M.D., B.D.G., E.M.M.

Data curation: G.V., A.G., E.T.C.

Writing – original draft: G.V., E.T.C., L.F.v.D.

Writing – review & editing: G.V., E.T.C., L.F.v.D.

Visualization: G.V., A.G., E.T.C., L.F.v.D.

Supervision: E.T.C., L.F.v.D.

Project administration: G.V., E.T.C., L.F.v.D.

Funding acquisition: E.T.C., L.F.v.D.

References:

1. Longnecker, R. and F. Neipel, *Introduction to the human gamma-herpesviruses*, in *Human Herpesviruses: Biology, Therapy, and Immunoprophylaxis*, A. Arvin, et al., Editors. 2007: Cambridge.

2. Mariggio, G., S. Koch, and T.F. Schulz, *Kaposi sarcoma herpesvirus pathogenesis.* Philos Trans R Soc Lond B Biol Sci, 2017. **372**(1732).

3. Young, L.S., L.F. Yap, and P.G. Murray, *Epstein-Barr virus: more than 50 years old and still providing surprises.* Nat Rev Cancer, 2016. **16**(12): p. 789-802.

4. Virgin, H.W.t., et al., *Complete sequence and genomic analysis of murine gammaherpesvirus 68.* J Virol, 1997. **71**(8): p. 5894-904.

5. Wang, Y., S.A. Tibbetts, and L.T. Krug, *Conquering the Host: Determinants of Pathogenesis Learned from Murine Gammaherpesvirus 68.* Annu Rev Virol, 2021. **8**(1): p. 349-371.

6. Oko, L.M., et al., *Multidimensional analysis of Gammaherpesvirus RNA expression reveals unexpected heterogeneity of gene expression.* PLoS Pathog, 2019. **15**(6): p. e1007849.

7. Landis, J.T., et al., *Evidence for Multiple Subpopulations of Herpesvirus-Latently Infected Cells.* mBio, 2022: p. e0347321.

8. Jung, K.L., et al., *Single-cell analysis of Kaposi's sarcoma-associated herpesvirus infection in three-dimensional air-liquid interface culture model.* PLoS Pathog, 2022. **18**(8): p. e1010775.

9. Berger, J.N., et al., *Redefining De Novo Gammaherpesvirus Infection Through High-Dimensional, Single-Cell Analysis of Virus and Host.* bioRxiv, 2020.

10. SoRelle, E.D., et al., *Epstein-Barr virus reactivation induces divergent abortive, reprogrammed, and host shutoff states by lytic progression.* PLoS Pathog, 2024. **20**(10): p. e1012341.

11. Salek-Ardakani, S., S.A. Lyons, and J.R. Arrand, *Epstein-Barr virus promotes human monocyte survival and maturation through a paracrine induction of IFN-alpha.* J Immunol, 2004. **173**(1): p. 321-31.

12. Tugizov, S., et al., *Epstein-Barr virus (EBV)-infected monocytes facilitate dissemination of EBV within the oral mucosal epithelium.* J Virol, 2007. **81**(11): p. 5484-96.

13. Torii, Y., et al., *Epstein-Barr virus infection-induced inflammasome activation in human monocytes.* PLoS One, 2017. **12**(4): p. e0175053.

14. Frederico, B., et al., *Myeloid infection links epithelial and B cell tropisms of Murid Herpesvirus-4.* PLoS Pathog, 2012. **8**(9): p. e1002935.

15. Weck, K.E., et al., *Macrophages are the major reservoir of latent murine gammaherpesvirus 68 in peritoneal cells.* J Virol, 1999. **73**(4): p. 3273-83.

16. West, J. and B. Damania, *Upregulation of the TLR3 pathway by Kaposi's sarcoma-associated herpesvirus during primary infection.* J Virol, 2008. **82**(11): p. 5440-9.

17. Gregory, S.M., et al., *Discovery of a viral NLR homolog that inhibits the inflammasome.* Science, 2011. **331**(6015): p. 330-4.

18. Gregory, S.M., et al., *Latent Kaposi's sarcoma-associated herpesvirus infection of monocytes downregulates expression of adaptive immune response costimulatory receptors and proinflammatory cytokines.* J Virol, 2012. **86**(7): p. 3916-23.

19. Host, K.M., et al., *Kaposi's Sarcoma-Associated Herpesvirus Increases PD-L1 and Proinflammatory Cytokine Expression in Human Monocytes.* mBio, 2017. **8**(5).

20. Kerur, N., et al., *Characterization of entry and infection of monocytic THP-1 cells by Kaposi's sarcoma associated herpesvirus (KSHV): role of heparan sulfate, DC-SIGN, integrins and signaling.* Virology, 2010. **406**(1): p. 103-16.

21. Savard, M., et al., *Infection of primary human monocytes by Epstein-Barr virus.* J Virol, 2000. **74**(6): p. 2612-9.

22. Shimoda, M., et al., *Virally encoded interleukin-6 facilitates KSHV replication in*
monocytes and induction of dysfunctional macrophages. PLoS Pathog, 2023. **19**(10): p.
e1011703.
23. Li, H., et al., *A replication-defective gammaherpesvirus efficiently establishes long-term*
latency in macrophages but not in B cells in vivo. J Virol, 2008. **82**(17): p. 8500-8.
24. Tarakanova, V.L., et al., *Gamma-herpesvirus kinase actively initiates a DNA damage*
response by inducing phosphorylation of H2AX to foster viral replication. Cell Host
Microbe, 2007. **1**(4): p. 275-86.
25. Tarakanova, V.L., et al., *MHV68 complement regulatory protein facilitates MHV68*
replication in primary macrophages in a complement independent manner. Virology,
2010. **396**(2): p. 323-8.
26. Szymula, A., et al., *Macrophages drive KSHV B cell latency.* Cell Rep, 2023. **42**(7): p.
112767.
27. Gaspar, M., et al., *Murid herpesvirus-4 exploits dendritic cells to infect B cells.* PLoS
Pathog, 2011. **7**(11): p. e1002346.
28. Lange, P.T., M.C. White, and B. Damania, *Activation and Evasion of Innate Immunity by*
Gammaherpesviruses. J Mol Biol, 2022. **434**(6): p. 167214.
29. Bussey, K.A., et al., *The gammaherpesviruses Kaposi's sarcoma-associated herpesvirus*
and murine gammaherpesvirus 68 modulate the Toll-like receptor-induced
proinflammatory cytokine response. J Virol, 2014. **88**(16): p. 9245-59.
30. Rosa, G.T., et al., *IgG fc receptors provide an alternative infection route for murine*
gamma-herpesvirus-68. PLoS One, 2007. **2**(6): p. e560.
31. Lawler, C., et al., *Rhadinovirus host entry by co-operative infection.* PLoS Pathog, 2015.
11(3): p. e1004761.
32. Flano, E., et al., *Latent murine gamma-herpesvirus infection is established in activated B*
cells, dendritic cells, and macrophages. J Immunol, 2000. **165**(2): p. 1074-81.
33. Frederico, B., et al., *A murid gamma-herpesviruses exploits normal splenic immune*
communication routes for systemic spread. Cell Host Microbe, 2014. **15**(4): p. 457-70.
34. Frederico, B., et al., *Subcapsular sinus macrophages limit acute gammaherpesvirus*
dissemination. J Gen Virol, 2015. **96**(8): p. 2314-2327.
35. Lange, P.T., et al., *Liver X Receptors Suppress Activity of Cholesterol and Fatty Acid*
Synthesis Pathways To Oppose Gammaherpesvirus Replication. mBio, 2018. **9**(4).
36. Aurubin, C.A., et al., *Low-Density Lipoprotein Receptor Suppresses the Endogenous*
Cholesterol Synthesis Pathway To Oppose Gammaherpesvirus Replication in Primary
Macrophages. J Virol, 2021. **95**(17): p. e0064921.
37. Reese, T.A., et al., *Helminth infection reactivates latent gamma-herpesvirus via cytokine*
competition at a viral promoter. Science, 2014. **345**(6196): p. 573-7.
38. Mounce, B.C., et al., *Gammaherpesvirus gene expression and DNA synthesis are*
facilitated by viral protein kinase and histone variant H2AX. Virology, 2011. **420**(2): p.
73-81.
39. Ramaswami, R., et al., *Transcriptional landscape of Kaposi sarcoma tumors identifies*
unique immunologic signatures and key determinants of angiogenesis. J Transl Med,
2023. **21**(1): p. 653.
40. Nair, A., et al., *The elevated expression of ORF75, a KSHV lytic gene, in Kaposi*
sarcoma lesions is driven by a GC-rich DNA cis element in its promoter region. PLoS
Pathog, 2025. **21**(3): p. e1012984.
41. Abbott, R.J., et al., *CD8+ T cell responses to lytic EBV infection: late antigen specificities*
as subdominant components of the total response. J Immunol, 2013. **191**(11): p. 5398-
409.
42. Sagou, K., et al., *Epstein-Barr virus lytic gene BNRF1 promotes B-cell lymphomagenesis*
via IFI27 upregulation. PLoS Pathog, 2024. **20**(2): p. e1011954.

43. Nealy, M.S., et al., *Use of a virus-encoded enzymatic marker reveals that a stable
fraction of memory B cells expresses latency-associated nuclear antigen throughout
chronic gammaherpesvirus infection.* J Virol, 2010. **84**(15): p. 7523-34.
44. Diebel, K.W., et al., *Gammaherpesvirus small noncoding RNAs are bifunctional
elements that regulate infection and contribute to virulence in vivo.* mBio, 2015. **6**(1): p.
e01670-14.
45. Kulkarni, A., et al., *Beyond bulk: a review of single cell transcriptomics methodologies
and applications.* Curr Opin Biotechnol, 2019. **58**: p. 129-136.
46. Van Skike, N.D., et al., *Viral FGARAT ORF75A promotes early events in lytic infection
and gammaherpesvirus pathogenesis in mice.* PLoS Pathog, 2018. **14**(2): p. e1006843.
47. Zhang, Z., et al., *A greedy algorithm for aligning DNA sequences.* J Comput Biol, 2000.
7(1-2): p. 203-14.
48. Weck, K.E., et al., *Mature B cells are required for acute splenic infection, but not for
establishment of latency, by murine gammaherpesvirus 68.* J Virol, 1996. **70**(10): p.
6775-80.
49. Cheng, B.Y., et al., *Tiled microarray identification of novel viral transcript structures and
distinct transcriptional profiles during two modes of productive murine
gammaherpesvirus 68 infection.* J Virol, 2012. **86**(8): p. 4340-57.
50. Forrest, J.C. and S.H. Speck, *Establishment of B-cell lines latently infected with
reactivation-competent murine gammaherpesvirus 68 provides evidence for viral
alteration of a DNA damage-signaling cascade.* J Virol, 2008. **82**(15): p. 7688-99.
51. Smith, K.R., et al., *Uracil-DNA glycosylase of murine gammaherpesvirus 68 binds
cognate viral replication factors independently of its catalytic residues.* mSphere, 2023.
8(5): p. e0027823.
52. Bieniasz, P.D., *Intrinsic immunity: a front-line defense against viral attack.* Nat Immunol,
2004. **5**(11): p. 1109-15.
53. Yan, N. and Z.J. Chen, *Intrinsic antiviral immunity.* Nat Immunol, 2012. **13**(3): p. 214-22.
54. Covarrubias, S., et al., *Host shutoff is a conserved phenotype of gammaherpesvirus
infection and is orchestrated exclusively from the cytoplasm.* J Virol, 2009. **83**(18): p.
9554-66.
55. Wang, G., et al., *Th2 Cytokine Modulates Herpesvirus Reactivation in a Cell Type
Specific Manner.* J Virol, 2021.
56. van Dyk, L.F., H.W.t. Virgin, and S.H. Speck, *The murine gammaherpesvirus 68 v-cyclin
is a critical regulator of reactivation from latency.* J Virol, 2000. **74**(16): p. 7451-61.
57. Wu, T.T., et al., *Rta of murine gammaherpesvirus 68 reactivates the complete lytic cycle
from latency.* J Virol, 2000. **74**(8): p. 3659-67.
58. Pavlova, I.V., H.W.t. Virgin, and S.H. Speck, *Disruption of gammaherpesvirus 68 gene
50 demonstrates that Rta is essential for virus replication.* J Virol, 2003. **77**(10): p. 5731-
9.
59. Wu, T.T., et al., *Function of Rta is essential for lytic replication of murine
gammaherpesvirus 68.* J Virol, 2001. **75**(19): p. 9262-73.
60. Martinez-Guzman, D., et al., *Transcription program of murine gammaherpesvirus 68.* J
Virol, 2003. **77**(19): p. 10488-503.
61. Gunther, T., et al., *A comparative epigenome analysis of gammaherpesviruses suggests
cis-acting sequence features as critical mediators of rapid polycomb recruitment.* PLoS
Pathog, 2019. **15**(10): p. e1007838.
62. Coleman, H.M., S. Efstathiou, and P.G. Stevenson, *Transcription of the murine
gammaherpesvirus 68 ORF73 from promoters in the viral terminal repeats.* J Gen Virol,
2005. **86**(Pt 3): p. 561-574.

63. Rose, T.M., et al., *Quantitative RNAseq analysis of Ugandan KS tumors reveals KSHV*
gene expression dominated by transcription from the LTd downstream latency promoter.
PLoS Pathog, 2018. **14**(12): p. e1007441.
64. Ramaswami, R., et al., *Transcriptional analysis identifies overlapping and tissue-distinct*
profiles between Kaposi sarcoma tumors of the skin and gastrointestinal tract. bioRxiv,
2022: p. 2022.03.18.484923.
65. Freeman, M.L., et al., *CD4 T cells specific for a latency-associated gamma-herpesvirus*
epitope are polyfunctional and cytotoxic. J Immunol, 2014. **193**(12): p. 5827-34.
66. Gredmark-Russ, S., et al., *The CD8 T-cell response against murine gammaherpesvirus*
68 is directed toward a broad repertoire of epitopes from both early and late antigens. J
Virol, 2008. **82**(24): p. 12205-12.
67. Freeman, M.L., et al., *Two kinetic patterns of epitope-specific CD8 T-cell responses*
following murine gammaherpesvirus 68 infection. J Virol, 2010. **84**(6): p. 2881-92.
68. Tan, C.S.E., C. Lawler, and P.G. Stevenson, *CD8+ T cell evasion mandates CD4+ T cell*
control of chronic gamma-herpesvirus infection. PLoS Pathog, 2017. **13**(4): p. e1006311.
69. Darrah, E.J., et al., *ATM supports gammaherpesvirus replication by attenuating type I*
interferon pathway. Virology, 2017. **510**: p. 137-146.
70. Schmalzriedt, D.L., et al., *Apolipoprotein E selectively supports gammaherpesvirus*
replication in macrophages. J Virol, 2025. **99**(6): p. e0048025.
71. Riggs, J.B., et al., *Optimized Detection of Acute MHV68 Infection With a Reporter*
System Identifies Large Peritoneal Macrophages as a Dominant Target of Primary
Infection. Front Microbiol, 2021. **12**: p. 656979.
72. Zarek, C.M., et al., *Preexisting helminth challenge exacerbates infection and reactivation*
of gammaherpesvirus in tissue resident macrophages. PLoS Pathog, 2023. **19**(10): p.
e1011691.
73. Goodwin, M.M., et al., *Murine gammaherpesvirus 68 has evolved gamma interferon and*
stat1-repressible promoters for the lytic switch gene 50. J Virol, 2010. **84**(7): p. 3711-7.
74. Mboko, W.P., et al., *Interferon Regulatory Factor 1 and Type I Interferon Cooperate To*
Control Acute Gammaherpesvirus Infection. J Virol, 2017. **91**(1).
75. Mounce, B.C., et al., *A conserved gammaherpesvirus protein kinase targets histone*
deacetylases 1 and 2 to facilitate viral replication in primary macrophages. J Virol, 2013.
87(13): p. 7314-25.
76. Steed, A., et al., *Gamma interferon blocks gammaherpesvirus reactivation from latency*
in a cell type-specific manner. J Virol, 2007. **81**(11): p. 6134-40.
77. Rickabaugh, T.M., et al., *Generation of a latency-deficient gammaherpesvirus that is*
protective against secondary infection. J Virol, 2004. **78**(17): p. 9215-23.
78. Pfaffl, M.W., *A new mathematical model for relative quantification in real-time RT-PCR.*
Nucleic Acids Res, 2001. **29**(9): p. e45.
79. Coomes, S.M., et al., *Severe gammaherpesvirus-induced pneumonitis and fibrosis in*
syngeneic bone marrow transplant mice is related to effects of transforming growth
factor-beta. Am J Pathol, 2011. **179**(5): p. 2382-96.
80. Knox, A.N., et al., *Lytic Infection with Murine Gammaherpesvirus 68 Activates Host and*
Viral RNA Polymerase III Promoters and Enhances Noncoding RNA Expression. J Virol,
2021. **95**(14): p. e0007921.
81. Zheng, G.X., et al., *Massively parallel digital transcriptional profiling of single cells.* Nat
Commun, 2017. **8**: p. 14049.
82. Hao, Y., et al., *Integrated analysis of multimodal single-cell data.* Cell, 2021. **184**(13): p.
3573-3587 e29.
83. Wickham, H., *ggplot2: Elegant Graphics for Data Analysis*. 2016: Springer-Verlag New
York.

84.   Kapadia, S.B., et al., *Murine gammaherpesvirus 68 encodes a functional regulator of*
complement activation. J Virol, 1999. **73**(9): p. 7658-70.
937

Highlights

• MHV68 efficiently infects macrophages

• Infection results in three divergent outcomes regulated by discrete inputs

• Macrophages limit lytic replication at two distinct stages

• Macrophages are capable of lytic replication, promoted by the host cytokine IL-4 and viral genes.

Vragel et al
Fig. 1

Figure 1. Single cell RNA sequencing analysis of acute MHV68 macrophage infection.
C57BL/6J mice were infected with 1×10^6 PFU of WT MHV68.LANAβlac virus, with MHV68-infected, LANAβlac+ peritoneal cells sort-purified at 16 hpi and subjected to scRNA-seq analysis. **(A-B)** Visualization of cell types based on lineage marker genes depicted in panel B. **(C)** Quantification of cell subsets defined by scRNA-seq. **(D)** UMAP visualization of cells with at least 1 viral UMI detected per cell identified by purple. **(E)** Frequency of cells stratified by the fraction of viral UMIs out of total UMIs detected per cell. 56 cells had >0.1% of total UMIs contributed by viral UMIs, including 23 of 9101 macrophages (0.25%), 25 of 25 macrophages with lytic gene expression (100%) and 8 of 70 dendritic cells (11.43%). Data are from 9,537 cells, with cell subsets enumerated in panel C. Supporting data in **Fig S1-S2**.

Vragel et al
Fig. 2

Figure 2. Single cell RNA sequencing analysis of acute MHV68 macrophage infection reveals two distinct transcriptional profiles. Analysis of viral gene expression defined by scRNA-seq in macrophage subsets from **Figure 1. (A-C)** Frequency of cells with detectable reads mapping to MHV68 genes, comparing (A) macrophages (82.79% viral UMI+), (B) macrophages with proliferative gene signature (52.08% viral UMI+), or (C) macrophages with lytic gene expression (100% viral UMI+). Viral genes are indicated on the x axis, arranged from left to right based on the published MHV68 genome, with the frequency of cells positive for each viral gene depicted on the y axis. To facilitate comparison of gene expression across the viral genome and cell populations, circles are color-coded to identify values that are in the indicated ranges (defined in key, panel A). Dotted lines indicate different thresholds for viral gene positivity, empirically defined. For panel C, only genes with >75% positivity were color-coded, given the limited number of cells. **(D)** Frequency among macrophages defined by detection of ORF73, ORF75A, ORF75B, and ORF75C RNAs, enumerating all potential gene combinations macrophages from panels A-C.

Figure 3. MHV68 readily infects J774 macrophages with limited lytic transcription. Analysis of MHV68 infection (WT MHV68.LANAβlac, MOI=1 PFU/cell) comparing 3T12 fibroblasts and J774 macrophages. **(A)** Representative brightfield images of infected 3T12 (top) or J774 cells (bottom) early (24 hpi, left) or late (72 hpi, right) post-infection. 3T12 cells showed cytopathic effect (CPE), including cell rounding (24 hpi) and monolayer destruction (72 hpi), in contrast to J774 cells. **(B)** Viral DNA quantitation defined by qPCR. **(C)** Virus replication defined by plaque assay. **(D-E)** Frequency of MHV68-infected, LANAβlac+ cells defined by flow cytometry, detected by CCF2 substrate cleavage, analyzing single, viable cells. Samples infected with WT MHV68 (left, panel E) define background signal and do not encode LANAβlac. **(F-G)** Quantitation of viral gene expression defined by qRT-PCR, comparing 3T12 and J774 cells at 24 hpi (left) or J774 cells at 24 and 48 hpi (right). Flow cytometric analysis focused on single, viable cells. Data depict mean ± SEM, from 3 independent experiments with 3 biological replicates per experiment (B, D, F-G) or 2 (t=0) or 6 (t=24, 48 hpi) independent experiments (C), with individual symbols representing independent biological samples. Statistical analysis was done using unpaired t test, with statistically significant differences indicated, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

A.

WT MHV68.LANAβlac   WT MHV68.H.GFP

3T12

GFP Neg. 99.9%   GFP Neg. 83.9%
GFP Pos. 0.059%   GFP Pos. 16.0%

J774

GFP Neg. 99.8%   GFP Neg. 99.3%
GFP Pos. 0.20%   GFP Pos. 0.56%

FSC → GFP

B.

GFP+ Frequency

**

WT   WT.H.GFP   WT   WT.H.GFP
3T12   J774

C.

% vCycβlac+ cells

****   **

Time (hpi): 24 48   24 48
Cells: 3T12   J774

D.

Mock   WT MHV68.LANAβlac
24hpi   24hpi   48hpi

3T12

0.21 0.76   2.70 13.5
87.9 11.1   55.0 28.8

J774

0.012 0   0.041 0.041   0.22 0.38
99.9 0.048   99.7 0.19   98.7 0.73

vRCA (PE) →   γH2AX (AF647) →

E.

Protein+ Frequency

****   ns

3T12   J774   J774
24 hpi   48 hpi

3T12:
☐ pH2AX+, vRCA-
▨ pH2AX-, vRCA+
■ pH2AX+, vRCA+

J774:
☐ pH2AX+, vRCA-
▨ pH2AX-, vRCA+
■ pH2AX+, vRCA+

Vragel et al
Fig. 4

Figure 4. Only rare MHV68 infected macrophages display lytic gene expression. Analysis of MHV68 infection and lytic cycle profiles with either MHV68.H.GFP or MHV68.LANAβlac (MOI=1 PFU/cell) comparing 3T12 fibroblasts and J774 macrophages. **(A)** Flow cytometric analysis of virus-expressed GFP in 3T12 or J774 cells 18 hpi. Data depict the frequency of cells with GFP expression, an indicator of MHV68.H.GFP infection. Background fluorescence defined by WT MHV68.LANAβlac-infected cells (left column), without the addition of the CCF2 fluorescent substrate. **(B)** Frequency of HygroGFP+ cells defined by flow cytometry, as in panel A. **(C)** Frequency of vCycβlac+ cells defined by flow cytometry. **(D)** Flow cytometric analysis of lytic infection-associated proteins, vRCA and γH2AX in 3T12 (top) or J774 (bottom) cells, comparing mock (left) or WT MHV68.LANAβlac-infected cells. Data for 3T12 cells at 48 hpi was not assessed due to extensive cell destruction. **(E)** Quantification of the frequency of infected cells with protein expression associated with lytic infection defined by flow cytometry, as in panel D. Flow cytometry data focus on single, viable cells defined using gating strategies in **Supp. Fig. S3A** and **S4B**. Data show mean ± SEM from 3 experiments, with 2 (B,E) or 3 samples (C) per experiment. Statistical analysis done using Mann-Whitney test, with statistically significant differences as indicated, **$p<0.01$, ****$p<0.0001$.

Vragel et al
Fig. 4

Figure 5. High dose MHV68 infection results in abortive replication. MHV68 outcomes in J774 macrophages comparing intermediate (MOI=1 PFU/cell) or high dose (MOI=10) infection. **(A)** Flow cytometric analysis of GFP in MHV68.LANAβlac- (left; negative control with no CCF2 substrate) or MHV68.H.GFP-infected cells (right) at MOI=1 (top) or MOI=10 (bottom). **(B)** Frequency of GFP+ cells quantified by flow cytometry, as in A. **(C)** Frequency of ORF59GFP+ J774 cells quantified by flow cytometry at 24 or 72 hpi. **(D)** Flow cytometric analysis of actin mRNA levels 24 hpi in mock or WT MHV68-infected cells with individual rows showing independent biological samples. **(E)** Quantitation of viral RNA expression by qRT-PCR standardized to 18S RNA, comparing intermediate or high MOI (MHV68.LANAβlac) at the indicated timepoints. **(F)** Frequency of MHV68.HygroGFP-infected single cells that are viable, quantified by flow cytometry. **(G)** Quantitation of viral DNA by qPCR in J774 (left) or 3T12 cells (right) as a function of MOI (MHV68.LANAβlac). **(H)** Infectious virus production quantified by plaque assay, comparing 3T12 cells (MOI=1) with J774 cells (MOI=1 or 10). Flow cytometry data analyzed single, viable cells (A-D) or single cells (F), with gating strategy in **Supp. Fig. 3**. Data show mean ± SEM, with individual symbols showing independent biological samples from 2-3 independent experiments with 2-3 biological replicates per experiment. Statistical analysis was done using Mann-Whitney test with statistically significant differences indicated, **p<0.01, ***p<0.001.

Vragel et al
Fig. 6

Figure 6. MHV68 infection of macrophages is modulated by host and viral factors. All infections done with MOI=1. **(A-D)** IL-4 effect on MHV68 infection and genome replication in J774 macrophages. IL-4 cultures were pretreated with IL-4 for 16-18 hours at the indicated concentration, with fresh IL-4 replenished after infection. **(A)** Flow cytometric analysis of MHV68.H.GFP infection in 3T12 (top), untreated J774 (middle) or IL4-treated J774 cells at 24 (left) or 48 hpi (right), analyzing single, viable cells. **(B)** Frequency of GFP+ cells quantified by flow cytometry, as in A. **(C-D)** IL-4 effect on viral DNA replication **(C)** and infectious virus production **(D)** in WT MHV68-infected J774 cells. **(E)** Frequency of J774 cells expressing markers of lytic infection (γH2AX, vRCA) quantified by flow cytometry, comparing WT, vCycKO or CA-RTA MHV68 at 24 or 48 hpi. Flow cytometric analysis focused on single, viable cells. Data depict mean $\pm$ SEM, with individual symbols representing independent biological samples from 2-3 independent experiments with 2-3 biological replicates per experiment (B,C,E) or from 1-2 independent experiments with 3 biological replicates measured (D). Statistical analysis used (B) unpaired t test, (C) one-way ANOVA comparing samples collected at the same timepoint untreated versus cytokine treated, or (E) one-way ANOVA comparing samples collected at the same timepoint with WT infection (nonparametric Kruskal-Willis test subjected to Dunn's multiple comparison test), with statistically significant differences indicated, *p<0.05, **p<0.01, ****p<0.0001.

Vragel et al
Fig. S1

Supplemental Figure 1 (Figure S1). Single cell RNA sequencing analysis of acute MHV68 macrophage infection. Data correspond to **Figure 1**. (A) Visualization of gene expression based on UMAP topography in Fig. 1A. Maximal expression is designated by purple, with absence of detection designated by gray. Each panel depicts a distinct lineage-defining gene used to identify cell types in Fig. 1, with signature genes for each cell type as follows. Macrophage: Adgre1 (encodes the F4/80 protein), Itgam (encodes the CD11b protein). Macrophage proliferative: Top2a, Mki67. Macrophage lytic: ORF54, ORF7. B cell: Cd19, CD79a. Monocyte: Ly6c2, Ccr2. Dendritic cell: Itgax, H2-Ab1. T cell: Cd3d, Thy1. (B-E) scRNA-seq quality control metrics across cell subsets, including (B) UMI Count, (C) Features, (D) % mitochondrial reads, and (E) S score, a proliferation-associated gene signature.

Vragel et al
Fig. S2

Figure S2. Single cell RNA sequencing analysis of acute MHV68 macrophage infection analyzed using a higher number of cell clusters. Data correspond to **Figure 1**, with scRNA-seq subjected to Seurat-based designation of 14 clusters. **(A)** UMAP visualization of Seurat-defined clusters. **(B)** Visualization of the frequency and magnitude of host gene expression across clusters, with cell type designation indicated on right. **(C-F)** Quality control metrics across the 14 clusters, including (C) UMI Count, (D) Features, (E) % mitochondrial reads, and (F) S score, a proliferation-associated gene signature. **(G)** UMAP visualization of cells with at least 1 detectable viral UMI per cell identified in red. **(H)** Frequency of cell subsets stratified by the frequency of viral UMIs of total UMIs detected per cell. **(I-J)** Percentage of viral UMIs out of total UMIs across clusters (panel I, y axis range from 0-50%), or (J) focused only on the majority of virus positive cells (y axis range, 0-0.1%; excluding 56 cells with % viral UMI > 0.1). Data are from 9,537 cells, with clusters containing between 25-2,532 cells. Panel G is included to provide a frame of reference and depicts the same data as in Figure 1D.

Vragel et al
Fig. S3

Figure S3. Flow cytometry gating strategy and viral gene expression during MHV68 infection. (A) Flow cytometry gating strategy to analyze MHV68-infected cells, in support of **Figure 3E**. 3T12 cells infected with MHV68 LANAblac or WT.MHV68.H.GFP were analyzed by flow cytometry, using sequential focused analysis (i.e. "gating") to consider cells that are single cells (defined sequentially as "Singlets by SSC" and "Singlets by FSC") that are viable ("Live cells") defined by exclusion of a dye that identifies dead cells. Live cells were then analyzed for GFP fluorescence as measured by the GFP detection channel. Cells infected with MHV68.LANAβlac, without additional of the CCF2 fluorescent substrate, identified background fluorescence, in contrast to samples infected with MHV68.H.GFP which encoded GFP. Controls included single-stained samples with only one stain to define the bounds of a gate for the fluorophore of interest. **(B-D)** Viral gene expression defined by qRT-PCR in 3T12 or J774 cells infected with WT MHV68.LANAblac (MOI=1) at 16 hpi (B,D) or 24 hpi (C) in support of **Figure 3F-G**. Data show mean ± SEM from 3 independent experiments with 3 biological replicates measured in technical triplicates per experiment (depicted by individual symbols) (B,D) or from 1 experiment with 3 biological replicates measured in technical triplicates, with ORF73 data from Fig. 2B (C). Statistical analysis was done using unpaired t test with statistically significant differences as indicated. ** $p<0.01$, ***$p<0.001$, ****$p<0.0001$. ns, no statistical significance detected.

Figure S4. Gating of flow cytometry analysis of MHV68 LANAβlac infected J774 cells for lytic protein markers. (A) Analysis of GFP expression in 3T12 fibroblasts, J774 and RAW264.7 macrophage cell lines at 24 or 48 hpi, following infection with MHV68.H.GFP (MOI=1). **(B)** Gating strategy to analyze vRCA and γH2AX expression in MHV68-infected J774 cells, in support of **Figure 4C**. Cells were analyzed by flow cytometry, using sequential focused analysis (i.e. "gating") to consider cells that are single cells (defined sequentially as "Singlets by SSC" and "Singlets by FSC") that are viable ("Live cells") defined by exclusion of a dye that identifies dead cells. Live cells were then analyzed for antibody binding for lytic protein markers. Gate boundaries were defined using a combination of "single stain" samples, containing only the fluorophore of interest, and "full minus one" controls, lacking the antibody in the fluorescent channel of the excluded fluorophore (not shown). **(C-I)** Analysis of MHV68-infected, FACS-purified J774 cells over time. **(C)** Experimental overview with J774 macrophages infected with MHV68.LANAβlac (MOI=1), followed by sorting for LANAblac+ cells defined based on CCF2 substrate cleavage. Mock and LANAblac+ sorted cells were cultured and analyzed over 14 days. **(D)** Total cell counts over the course of the study, with each vertical line indicate a subsequent cell passage. **(E)** Flow cytometric analysis of LANAβlac expression defined by cleavage of the CCF2 fluorescent beta-lactamase substrate, analyzing single, viable cells, with values in the right gate defining the frequency of cells with CCF2 substrate cleavage. **(F)** Quantification of LANAβlac+ cell frequency defined by flow cytometry, as in panel E. **(G)** Flow cytometric analysis of MHV68 lytic cycle-associated proteins (vRCA, γH2AX) at 14 dps, comparing mock (left) or CCF2-sorted, LANAblac+ cells (right). Data depict events that were single, viable cells, with values defining the frequency of cells with vRCA and/or phosphorylated host γH2AX, indicative of lytic MHV68 infection, as in panel B. **(H)** Frequency of vRCA+ γH2AX+ cells defined by flow cytometry, as in panel G. **(I)** Quantitation of viral DNA by qPCR. Dotted line indicates mean 3T12 viral genome copies/cell at 72 hpi. Data show mean ± SEM from 3 independent experiments with 3 biological replicates per experiment (depicted by individual symbols) (A) or from 2 independent experiments.