Murine Gammaherpesvirus 68 Efficiently Infects Myeloid Cells Resulting In An

Atypical, Restricted Form Of Infection

Gabrielle Vragel[1], Brittany D. Gomez[1], Rachael E. Kostelecky[1], Kyra S. Noell[1,2], Ashley

Tseng[1,2], Shirli Cohen[1], Manaal Dawaldi[1], Eva M. Medina[3], Elizabeth A. Nail[1], Andrew

Goodspeed[4,5], Eric T. Clambey[2]*, Linda F. van Dyk[1]*#

Affiliations:

[1] Department of Immunology and Microbiology, [2] Department of Anesthesiology, [3]

Department of Neurology, [4] Department of Pharmacology, [5] University of Colorado

Cancer Center, University of Colorado Denver | Anschutz Medical Campus, School of

Medicine, Aurora, CO, 80045, USA

* Co-Corresponding authors:

Eric T. Clambey, eric.clambey@cuanschutz.edu, 303-724-7783 (phone)

Linda F. van Dyk, linda.vandyk@cuanschutz.edu, 303-724-4207 (phone)

 # Lead author for MS correspondence

Linda F. van Dyk, linda.vandyk@cuanschutz.edu, 303-724-4207 (phone)

ORCIDs

https://orcid.org/0000-0001-9167-3013 (G.V.); 0000-0002-3871-3773 (B.D.G.); 0000-
0002-6529-3831 (R.E.K.); 0000-0003-3623-8822 (K.S.N); 0000-0002-3078-3718 (A.T.);
0000-0002-0253-786X (S.C.); 0009-0005-3187-7803 (M.K.D); 0000-0001-6899-0213
(E.M.M.);0000-0002-4859-4122 (E.A.N); 0000-0001-7055-7128 (A.G.); 0000-0002-
7972-9544 (E.T.C.); 0000-0003-2662-5554 (L.F.v.D.)

Abstract

The gammaherpesviruses (γHVs) establish a lifelong infection in their hosts, with the

cellular outcome of infection intimately regulated by target cell type. Murine

gammaherpesvirus 68 (MHV68), a small animal model of γHV infection, infects

macrophages in vivo, resulting in a range of outcomes, from lytic replication to latent

infection. Here, we have further investigated the nature of MHV68 macrophage infection

using reductionist and primary in vivo infection studies. While MHV68 readily infected the

J774 macrophage cell line, viral gene expression and replication were significantly

impaired relative to a fully permissive fibroblast cell line. Lytic replication only occurred in

a small subset of MHV68-infected J774 cells, despite the fact that these cells were fully

competent to support lytic replication following pre-treatment with interleukin-4, a known

potentiator of replication in macrophages. In parallel, we harvested virally-infected

macrophages at 16 hours after MHV68 infection *in vivo* and analyzed gene expression

by single cell RNA-sequencing. Among virally infected macrophages, only rare (0.25%)

cells had lytic cycle gene expression, characterized by detection of multiple lytic cycle

RNAs. In contrast, ~50% of virally-infected macrophages were characterized by

expression of ORF75A, ORF75B and/or ORF75C, in the absence of other detectable viral

RNAs. Selective transcription of the ORF75 locus also occurred in MHV68-infected J774

cells. In total, these studies indicate that MHV68 efficiently infects macrophages, with the

majority of cells characterized by an atypical state of restricted viral transcription, and only

rare cells undergoing lytic replication.

Importance:

The human gammaherpesviruses Epstein-Barr virus and Kaposi's sarcoma associated

herpesvirus are DNA viruses that cause lifelong infection and are associated with

multiple diseases, especially in immunocompromised individuals. Murine

gammaherpesvirus 68 (MHV68) is a powerful mouse model that permits close

examination of these viruses. Previous studies of MHV68 identified that macrophages

are an important in vivo target of infection; how infection within these cells is regulated

remains incompletely understood. Here, we demonstrate that MHV68 infection of

macrophages is characterized by two divergent outcomes across a population of

infected cells: while a small subset of cells undergo lytic replication, to make new virus

progeny, the majority of cells are characterized by an atypical, restricted form of

infection characterized by a distinct viral gene transcription program not previously

reported. These studies highlight important cell-type specific outcomes of

gammaherpesvirus infection and identify a potential alternate program by which these

viruses usurp macrophages.

Introduction

The gammaherpesviruses (γHVs) are large double-stranded DNA tumor viruses that establish a lifelong infection in their host and are associated with a variety of diseases, particularly in immunosuppressed individuals. These viruses include the human γHVs Epstein-Barr Virus (EBV) and Kaposi's sarcoma-associated herpesvirus (KSHV or HHV-8), and infect multiple cell types, typically resulting in either lytic infection or latent infection on a cellular level (1). The outcome of infection at this level is influenced by multiple factors, including target cell type and cell-type specific virus-host interactions. The γHVs are associated with the development of several malignancies especially in immunocompromised individuals. (2, 3). Due to the strict host specificity of the human γHVs, we and others have used the mouse γHV, murine gammaherpesvirus 68 (MHV68, γHV68 or MuHV-4), as a genetic and phenotypically relevant model to study γHV infection and pathogenesis (4, 5).

MHV68 infection of mice is a multi-step process, from lytic infection at mucosal surfaces in epithelial cells, to dissemination to the B-cell compartment where it can establish lifelong, systemic latency (6). While B cells are a major latent reservoir for MHV68 (7-9), EBV (10) and potentially KSHV (11), MHV68 has also been shown to establish latency in lung, spleen and peritoneal macrophages (7, 12, 13). Macrophage infection may occur by multiple mechanisms, from direct virus binding and entry, to internalization of the virus via epithelial cell presentation of MHV68 (14), to potential infection through an Fc receptor-dependent mechanism (15). MHV68 has also been reported to undergo lytic replication in macrophages in vitro and in vivo, albeit at lower levels than replication

within fibroblasts (16, 17). While MHV68 passage through myeloid cells has been

reported to be important for the establishment of B cell latency, a process potentially

involving both alveolar macrophages and marginal zone macrophages (12, 18), not all

macrophage subsets are created equal: subcapsular sinus macrophages have been

reported to be poorly permissive for MHV68 infection, serving an antiviral role (19).

These data emphasize the complexity of MHV68-macrophage interactions and the need

for further analysis of these interactions.

Here, we sought to better understand the regulation of MHV68 infection in macrophages

in cell culture and primary infection compared with fibroblasts, a known target of lytic

replication. We found efficient infection of both cell types, but significantly reduced DNA

synthesis and gene expression in macrophages relative to fibroblasts. Lytic protein

induction only occurred in rare macrophages. In our *in vivo* analysis, we found a rare

population of macrophages undergoing lytic replication, with the major population of

infected peritoneal macrophages demonstrating a unique transcriptional profile

characterized by expression of ORF75A, ORF75B, and/or ORF75C, in the absence of

other lytic genes. These studies suggest that macrophages are efficiently infected but

characterized by atypical, restricted viral transcription that is not representative of lytic

nor latent gene expression patterns described to date.

RESULTS

An in vitro model of MHV68-macrophage infection shows limited lytic replication despite robust viral entry.

The outcome of MHV68 on a cell level varies depending on host cell type and immune status (5, 20, 21). MHV68 infection of macrophages has been associated with both lytic replication and latent infection (7, 12, 16) with data suggesting that myeloid cells may be an important intermediary prior to B-cell latency (12). Despite these insights, the exact nature of myeloid cell infection remains incompletely characterized. We characterized MHV68 infection in the mouse myeloid-like cell line, J774, compared with infection of the 3T12 mouse fibroblast cell line, which fully supports lytic virus replication (22).

First, we compared the ability of MHV68 to replicate in both cell lines. As expected, MHV68 underwent robust lytic replication in 3T12s, characterized by rapid virus production with extensive cytopathic effect (CPE) by 72 hours post infection (hpi). In contrast, MHV68-infected cultures of J774 cells showed minimal evidence of infection, with no discernable CPE (Fig. 1A) and minimal viral replication (Fig. 1B).

The limited replication of MHV68 in J774 cells could potentially be due to inefficient infection. To define infection frequency, we infected cells with WT MHV68 LANAβlac, a virus that contains a gene fusion between the ORF73 gene, an immediate early gene encoding the latency-associated nuclear antigen (LANA) and the beta-lactamase (βlac) gene (23, 24). As ORF73, and LANA, are expressed during latent and lytic infection (23), βlac activity serves as an effective indicator of multiple states of infection by flow cytometry, defined by cleavage of a fluorescent substrate. As

predicted, infection of cells with WT MHV68 lacking the beta-lactamase reporter had no

detectable βlac+ cells (left panel, Fig. 1C). In contrast, infection with WT MHV68

LANAβlac resulted in a high frequency of LANAβlac+ cells in both 3T12s fibroblasts

(~77%) and J774 cells (~45%) at 24 hpi, that increased to ~81% LANAβlac+ cells by 72

hpi (Fig. 1C, D). These data demonstrate that J774 cells are fully susceptible to MHV68

infection and are capable of initiating transcription and translation of ORF73 (an

immediate early gene), despite restricted production of infectious virus.

J774 macrophages are deficient in viral DNA replication and lytic viral gene

expression.

We next quantified viral DNA replication between the two cell lines. MHV68

underwent large (>2.5 $\log_{10}$) increases in viral DNA replication in 3T12 fibroblasts; in

contrast, MHV68-infected J774 myeloid cells had minimal changes in viral DNA

replication over input virus (Fig. 2A).

We next analyzed MHV68 transcription in J774 cells, quantifying immediate early

(ORF73), early, early/late (ORF6), and late gene (M7) expression (25). Viral gene

expression was decreased in J774 myeloid cells relative to 3T12 fibroblasts at both 16

and 24 hpi (Fig. 2B). Whereas ORF73 (and Blac) expression was reduced ~10 fold,

both ORF6 and M7 transcripts were decreased >100 fold (Fig. 2B, Supp. Fig 1),

consistent with greater deficits in lytic gene expression for early and late genes.

When we quantified viral gene expression within J774 cells over time, we found

that ORF73 was relatively comparable between 24-48 hpi (bottom panel, Fig. 2B). In

contrast, ORF6 and M7 expression levels increased ~10x between 24 hpi and 48 hpi

(Fig. 2B). As ORF6 and M7 are lytic cycle transcripts, these changes are consistent with

some degree of lytic replication in MHV68-infected J774 cultures.

MHV68-infected J774 macrophages contain a small subset of cells that

demonstrate a lytic infection profile.

To assess lytic replication within MHV68-infected J774 cells, we used two

approaches that afforded single-cell resolution of infection. First, we infected 3T12 and

J774 cells with a recombinant WT MHV68 virus, MHV68.H.GFP, that encodes GFP

fused to a hygromycin resistance protein, under the transcriptional control of the human

cytomegalovirus IE promoter inserted between ORF27 and 29b (26). This virus robustly

expresses GFP during active, lytic infection, allowing identification of cells that have

initiated transcription via flow cytometry. When we infected 3T12 and J774 cells with

MHV68.H.GFP virus and assessed GFP expression at 18 hpi, 3T12 cells had a much

higher frequency of GFP+ cells than J774 cells (~8.2% vs. ~0.5% GFP+ respectively,

Fig. 3A-B, Supp. Fig. 2). For these studies, background fluorescence was defined by

infection with WT MHV68.LANAβlac, without addition of a fluorescent substrate. These

data indicate that early in infection, a rare population of J774 cells initiate robust

transcriptional induction from the viral genome.

We next quantified expression of proteins expressed or modified during lytic

MHV68 replication via flow cytometry (as in (27)). To do this, we measured the

frequency of cells that express the viral regulator of complement activation (vRCA), a

protein encoded by the ORF4 late gene, or demonstrated phosphorylated Histone

H2AX (γH2AX), a host protein that undergoes virus-induced phosphorylation (16). Mock

infected cells had minimal expression of either of these proteins ("Mock", Fig. 3C),

whereas ~45% of MHV68-infected 3T12 cells were either vRCA+, γH2AX+, or

vRCA+γH2AX+ by 24hpi (Fig. 3C-D, Supp. Fig. 3). In contrast, MHV68-infected J774

cells only contained ~2% cells at 24hpi and ~8% at 48hpi that expressed either

vRCA+, γH2AX+, or vRCA+γH2AX+ (Fig. 3C-D). These data indicate that only a minor

fraction of MHV68-infected J774 cells undergo lytic replication at these timepoints.

IL-4 pretreatment of J774 macrophages reveals a robust capacity of these cells to

support lytic replication.

The minor frequency of MHV68 infected J774 cells undergoing lytic replication

raised the possibility that J774 cells may poorly support lytic replication. To test whether

J774 cells had the capacity to support lytic replication, we pretreated cells with IL-4, a

known inducer of viral replication and reactivation from latency in primary macrophages

(28, 29). First, we quantified the frequency of GFP+ cells following MHV68.H.GFP

infection comparing 3T12s with untreated or IL-4 treated J774 cells, where cells were

both pre-treated with IL-4 and supplied with fresh IL-4 after infection. As expected, 3T12

cells had a significant fraction of GFP+ cells with near uniform expression by 48 hpi

(Fig. 4A). While untreated J774 cells had a minor frequency of GFP+ cells (~1% at 24

hpi, ~9% at 48 hpi), IL-4 treated J774 cells had a greatly enhanced frequency of GFP+

cells both early and late, with ~45% GFP+ cells by 48 hpi (Fig. 4A-B). We next

quantified viral DNA replication under similar conditions, treating J774 cells with either

10 or 100 ng/mL of IL-4 for 16 hours prior to infection with fresh IL-4 provided after

infection. IL-4 treatment resulted in a concentration-dependent increase in viral DNA

replication by 48 hpi after treatement with either 10 or 100 ng/mL IL-4, compared to the

minimal viral DNA replication observed in untreated cells, consistent with our earlier

observations (Fig. 4C). These data demonstrate that J774 cells have the capacity to

support lytic replication following treatment with IL-4, a known inducer of virus

replication in macrophages.

In vivo analysis of primary MHV68 macrophage infection by single-cell RNA-seq

reveals unanticipated viral gene expression.

To examine how our *in vitro* results reflect *in vivo* biology, we infected

immunocompetent, C57BL/6J mice by intraperitoneal injection with WT

MHV68.LANAβlac and sort purified peritoneal macrophages early after infection (16

hpi). Virally-infected peritoneal macrophages were purified based on cells that were

viable, expressed the F4/80+ macrophage-specific cell surface protein and

demonstrated LANAβlac+ expression by CCF-2 cleavage (as in Fig. 1). Cells were then

subjected to 3' based single-cell RNA-seq (scRNA-seq), analyzing host and viral gene

expression to identify infected cell subsets and viral gene expression respectively. We

identified 14 clusters of cell subsets, dominantly comprised of macrophages (Fig. 5A-C).

This analysis further identified minor frequencies of: i) macrophages with a proliferative

gene signature (i.e. "macrophage proliferating"), ii) macrophages with an MHV68 lytic

gene signature (i.e. "macrophage lytic"), defined by expression of two canonical lytic

genes ORF7 and ORF54, and infrequent iii) B cells, iv) T cells, v) dendritic cells and vi)

monocytes (Fig. 5A-C, Supplemental Fig. 4A-B, Supplemental Fig. 5A-B). Data were

characterized by high quality control metrics including high reads (unique molecular

identifiers, UMIs) and genes per cell, with low mitochondrial reads and only

"macrophage proliferating" cells characterized by a proliferative "S score" (Fig. 5D-G,

Supplemental Fig. 4C-F). When we visualized (Fig. 5H) and quantified the frequency of

cells with detectable viral UMIs (Fig. 5I), we found that the majority (n=7,512; 82.54%)

of macrophages had detectable viral reads accounting for <0.1% of total UMIs per cell,

with few (n=23; 0.25%) macrophages containing a higher proportion (>0.1%) of viral

reads (Fig. 5I). Similar results were found at the cluster level, with cluster 13, identified

as macrophages with lytic gene expression, characterized both by 100% viral UMI+

(including ORF7 and ORF54) and a higher proportion of viral UMIs per cell

(Supplemental Fig. 4G-J). The presence of cells with no detectable viral UMIs could

either be due to the depth of sequencing or to the known sparseness of scRNA-seq

data (30). Because these limitations of scRNA-seq could further extend to the lack of

detection of viral genes, below, subsequent discussion focuses on those genes that we

can definitively detect and avoids conclusion about lack of expression.

Next, we analyzed viral gene expression within macrophages, quantifying the

frequency of cells that expressed each viral gene, comparing the dominant macrophage

population (n=9101 cells), with minor macrophage populations with either a proliferative

gene signature (n=48 cells) or lytic gene expression (n=25 cells). While we detected

ORF73-βlac mRNA, the transcript encoding the LANAβlac reporter by which cells were

purified, in 12.89% of macrophages, there were equivalent or higher frequencies of cells

that expressed 3 other viral genes: ORF75C (30.75%), ORF75B (27.83%) and ORF75A

(12.54%) (Fig. 6A). Additional transcripts were detected in a lower frequency of cells:

ORF72 (5.3%), ORF58 (3.5%), with 1-3% cells containing detectable RNA for M3,

ORF7, ORF50, ORF57, ORF61, ORF74, M12, M13, or M14 (Fig. 6A). In contrast, the

remaining viral genes were only sporadically detected in less than 1% of cells. A similar

hierarchy, albeit reduced in magnitude, was also observed in the proliferating

macrophage cell subset (Fig. 6B).

In contrast to the dominant macrophage subset, rare lytic macrophages had a

distinct viral gene signature. Here, we detected multiple lytic cycle-associated genes

including ORF7, ORF54, ORF57, ORF58, and ORF61 in 100% of these rare cells (dark

blue circles, Fig. 6C). Further, >75% of cells contained detectable RNAs for M3, ORF6,

ORF21, ORF38, ORF50, ORF52, ORF59, ORF72 and ORF75C (light blue circles, Fig.

6C). The high frequency of cells that expressed multiple, canonical lytic cycle transcripts

is consistent these cells initiating lytic gene transcription in vivo.

We next sought to understand the high frequency detection of ORF75 RNAs

within the dominant macrophage population (Fig. 6A). Previous studies of the ORF75

locus have revealed multiple transcriptional units, including a polycistronic RNA that

spans from the 5' end of ORF75A through the 3' end of ORF75C (31). Notably, this

polycistronic RNA would map to ORF75C based on polyA-based, oligo-dT primed

scRNA-seq, and would not generate sequencing reads that would map to ORF75B or

ORF75A. We also considered whether a high-degree of homology at the 3' ends of the

ORF75 genes might complicate mapping of sequencing reads to each individual

ORF75. Though these genes likely arose from gene duplication, these genes have

limited nucleotide identity within the final 150 nucleotides of each gene: using blastn

(32), there is no significant similarity between the 3' ends of either ORF75A and

ORF75B or ORF75A and ORF75C. The closest homology across the final nucleotides

of ORF75B and ORF75C remained extremely low, with only a stretch of 22 of 26

nucleotides aligning across the final 150 nucleotides of ORF75B and ORF75C (data not

shown). This nucleotide divergence at the 3' end of these genes strongly suggests that

UMIs mapping to each gene were not due to errors in mapping. When we examined the

intersection of ORF73, ORF75A, ORF75B, and ORF75C detection within single cells,

considering all combinations of expression for these 4 viral genes, we found that

50.89% of macrophages had detectable expression of ORF75A, ORF75B and/or

ORF75C; in many of these cells, we did not detect ORF73 (Fig. 6D). In total, these

studies identify a high frequency of macrophages expressing transcripts from the

ORF75 locus, and further corroborate that a low frequency of primary infected

peritoneal macrophages are characterized by lytic transcription early after primary

infection.

MHV68-infected J774 macrophages show expression of the ORF75 locus in the

absence of full lytic gene transcription.

We next tested analyzed the expression of the ORF75 genes in MHV68-infected

J774 cells. ORF75A, ORF75B and ORF75C were readily detected in MHV68-infected

J774 cells from 16-48 hpi, albeit lower than in MHV68-infected 3T12 cells (Fig. 7A).

Among these genes, we observed two different patterns. ORF75A expression was

relatively comparable to ORF73 at all timepoints measured, with minimal change in

expression between 24 and 48 hpi, suggesting constitutive transcription for both ORF73

and ORF75A (Fig. 7A). In contrast, ORF75B and ORF75C transcript levels were

moderately lower than ORF73 at 16 and 24 hpi and showed a ~4 and ~22 fold increase

between 24 and 48 hpi, respectively (right panel, Fig. 7A). The induction of these

transcripts suggested that these transcripts may be at least, in part, induced during lytic

replication in J774 cells (akin to the increased signal observed for ORF6, M7, Fig. 2B).

To test the impact of viral DNA replication, a key process required for late gene

transcription, we treated MHV68-infected J774 or 3T12 cells with cidofovir (CDV), an

inhibitor of the viral DNA polymerase (33, 34). In parallel, we treated cells with the viral

DNA synthesis inhibitor phosphonoacetic acid (PAA) at 200 μg/mL (35); this treatment

was toxic to J774 cells (data not shown). CDV and PAA would inhibit lytic cycle

transcription of both early/late and late genes with minimal impact on gene expression

during latent (or restricted) infection. As expected, infected 3T12 cells had higher

baseline expression for ORF73, ORF75A, ORF75B, ORF75C than infected J774 cells

(Fig. 7B). When 3T12 cells were treated with CDV or PAA, there was a pronounced

reduction across viral genes compared to untreated samples, ranging from ~10-fold

reduction in CDV- or PAA-treated cultures for ORF73 and ORF75A, to ~1000-fold

reduction for ORF75C and M7 (Fig. 7B). In contrast, the impact of CDV on gene

expression in J774 cells was much more modest, with ~2-fold reduction in RNA

detected for ORF73 and ORF75A in CDV treated cultures relative to untreated cultures

(Fig. 7B), compared with ~5-10 fold reduction in ORF75B, ORF75C and M7 RNAs,

consistent with these transcripts being induced, in part, in a lytically-replicating subset of

cells. These data are consistent with expression of the ORF75 locus in MHV68-infected

J774 cells and suggest that at least the ORF75A gene shows comparable regulation to

that observed for ORF73.

DISCUSSION

DISCUSSION

The cross-talk between MHV68 and macrophages has been a significant area of

ongoing investigation, with early studies identifying macrophages as a significant latent

reservoir beyond B cells (7, 13). Since then, macrophages have been identified as an

early target of MHV68 infection after both intraperitoneal (36) and intranasal infection

(12, 19), where a significant fraction of virus passes through macrophages (as defined

by cre-dependent lineage tracing studies) (12). Whether MHV68 directly infects

macrophages, or macrophage infection is expedited by intercellular handoff from

infected epithelial cells (14) or by antibody-dependent entry through Fc receptors (15)

would likely be influenced by the route of infection, stage of infection and host immune

status. Regardless, there is now clear evidence that MHV68 can undergo replication

within a subset of macrophages, a process revealed by lineage-tracing studies (12),

lytic antigen expression within these cells (e.g. (16, 17)), and the studies presented

here. MHV68 replication in macrophages can further be enhanced in the context of

certain parasitic infections that both antagonize IFNγ-dependent suppressive

mechanisms and induce expression of the Th2 cytokines IL-4 or IL-13, STAT6-inducing

cytokines that can directly transactivate the gene 50 promoter, to promote lytic

replication (29).

By examining the heterogeneity of infection at single cell resolution, through both

flow cytometric analysis for markers of entry and LANA gene expression, lytic

replication and single-cell RNA sequencing, our studies provide a refined perspective of

MHV68 infection in macrophages. These studies unequivocally identify that MHV68

robustly infects macrophages and is capable of initiating transcription and translation of

at least the ORF73 gene, with only a minor fraction of these cells showing hallmarks of

lytic replication. Whether the majority of LANA+ cells reflect a bona fide latent

population or reflect a previously undescribed, restricted form of infection remains an

open question. Although our identification of ORF75 expression has not been previously

observed in the context of MHV68 macrophage infection, recent studies of KSHV gene

expression in Kaposi's sarcoma lesions identified ORF75 transcription in the absence of

many lytic genes, expression that also did not clearly correspond with expression from

the ORF73 locus (37), suggesting undiscovered modes of transcriptional regulation of

the ORF75 genes that may span across at least MHV68 and KSHV.

Broadly our findings are consistent with various aspects of published literature. On

the one hand, the magnitude of MHV68 lytic replication within bone marrow

macrophages or myeloid cells is relatively modest compared to fibroblasts (17, 38), and

lytic antigen expression is reported to occur within a subset of bone marrow-derived

macrophages and RAW264.7 macrophages (15, 28, 29). Despite the relatively modest

lytic replication of MHV68 in macrophages, MHV68 encodes multiple mechanisms that

facilitate this process, including ORF4 which encodes the vRCA and promotes

replication in a complement-independent manner (38) and ORF36 which encodes a

viral protein kinase that phosphorylates H2AX and antagonizes HDAC1 and HDAC2

(16, 39). Conversely, multiple inhibitory mechanisms have been reported to limit lytic

replication in macrophages, including IFNγ-dependent inhibition of the gene 50 promoter

(40), and cell-intrinsic mechanisms mediated in part by HDACs and NCoR (39, 41)

MHV68 lytic replication in macrophages is further regulated by cellular modulators of

metabolism, constrained both by Liver X receptor- and Low-Density Lipoprotein

Receptor-dependent mechanisms (42, 43). We postulate that these inhibitory

mechanisms may affect the relative proportion of cells initiating lytic replication with

MHV68, may constrain the magnitude of virus produced by lytic cells or may

fundamentally alter the susceptibility of infection across all cells. Future studies will

require single cell-based investigation of MHV68 to differentiate between these two

potentially distinct modes of regulation.

Despite major inroads from the above studies, in vitro models of macrophage

infection to date have fallen short of defining the dominant outcome of infection

described here as being truly latent, an important future question that will only be

resolved by focused analysis on cells lacking lytic protein expression. We anticipate that

this could be done by purifying infected macrophages that lack lytic gene expression,

followed by characterization of the state of the viral genome (i.e. episomal or linear) and

the ability of these cells to enter productive virus replication (i.e. reactivation from

latency). Alternatively, it is possible that this analysis may identify alternate regulation of

the viral genome, or a failure to produce virus particles, which would suggest a

restricted form of infection. Ultimately, these studies should seek to afford single-cell

resolution, potentially combined with temporal fate-mapping strategies, to better

differentiate between heterogeneous states of infection that may exist across a

population of macrophages, and how the nature of infection may evolve over time. This

is particularly relevant in the context of in vivo infection, where IFNγ constrains MHV68

reactivation from latency within peritoneal macrophages (44) and CD4 T cells constrain

chronic macrophage infection in the lung (17). Recent studies of parasitic infection have

further revealed that parasite-dependent expansion of the large peritoneal macrophage

compartment, a primary target of MHV68 infection (36), can dramatically enhance the

pool of MHV68-infected macrophages (45) indicating that MHV68 infection of

macrophages is subject to multiple positive and negative regulatory pathways.

This study has important limitations that constrain our interpretations. First, many of

the studies focus on infection of the J774 mouse myeloid cell line, a system we have

leveraged as a reductionist in vitro system to study MHV68-macrophage interactions.

Second, our studies of primary in vivo infection focus on a single, early timepoint of

macrophage infection. Thus, it remains possible that a time course of in vivo

macrophage infection may reveal a delayed form of lytic cycle transcription.

Conservatively, we can conclude at least that the high frequency of ORF75 genes

(ORF75A, B, and/or C) within primary macrophages is not explained by either the

current paradigm of lytic or latent transcription (e.g. exemplified by (25)). The

expression of individual ORF75 gene products, as defined by polyA-based scRNA-seq

suggests that there are additional mechanisms of transcriptional regulation within the

ORF75 locus than previously identified in lytically infected fibroblasts (31). These

studies may further identify macrophage-specific regulation of the ORF75 locus and/or

unique roles for the ORF75 gene products within macrophages. The ORF75 genes are

known to encode proteins with a number of functions that could disproportionally impact

macrophage infection.

The frequent expression of ORF75 genes within macrophages suggests that

ORF75-targeted interventions (e.g. vaccination against ORF75 gene products, which

contain known T cell epitopes (46-48) may selectively disrupt macrophage infection

over infection in other cell types. Additionally, the distinct gene expression patterns of

infected macrophages indicates variable susceptibility to treatment with current

antivirals.

In total, our studies demonstrate both heterogeneity of MHV68 infection

outcomes within macrophages and reveal an alternate transcriptional state at least early

after primary macrophage infection in vivo. These studies further suggest a previously

unappreciated role for the ORF75 locus as a potentially important regulator of

macrophage infection, or a marker of an alternate outcome of infection that is neither

lytic nor latent.

METHODS

Viruses and Tissue Culture

Mouse 3T12 fibroblasts (ATCC CCL-164) and mouse J774A.1 macrophages (subsequently referred to as "J774"; ATCC TIB-67) were cultured in complete DMEM (cDMEM): Dulbecco's modified Eagle medium (DMEM; Life Technologies) supplemented with 5% and 10% fetal bovine serum (FBS; Atlanta Biologicals) respectively, 2 mM L-glutamine, 10 U/mL penicillin, and 10 µg/mL streptomycin sulfate. Cells were cultured at 37°C with 5% CO_2. Wild-type (WT) MHV68, MHV68.LANA::βlac, and MHV68.H.GFP viruses were grown and prepared as described previously (23, 24, 26), with all virus titers determined from at least three independent plaque assays.

Infection

All in vitro infections were based on live cell counts with the TC20 Automated Cell Counter (Bio-Rad) with Trypan Blue dye (Bio-Rad, Cat. No. 145-0021). Viral stock was diluted with cDMEM for viral inoculum with a multiplicity of infection (MOI) of 1 PFU/cell. Cells were incubated at 37°C with 5% CO_2 for 1 hour, with rocking every 15 min, before viral inoculum was removed and replaced with 1mL of cDMEM. For qPCR analysis of viral DNA replication and for beta-lactamase staining and flow cytometric analysis of infected cells, viral inoculum was left on cells after 1 hr infection time. For IL-4 treatments, media containing recombinant mouse IL-4 (PeproTech, Cat. No. 214-14) was added to cells 16 hours before infection. Following infection, media containing 0, 10 or 100ng/mL of fresh IL-4 containing media were added respectively. Samples were harvested at the indicated hours post-infection (hpi) through either cell scraping (for DNA isolation) or through incubation in TRIzol (for RNA isolation).

Plaque Assay

Plaque assay quantification of viral titer was performed using 3T12 cells. Cells were plated in 12-well plates at 8.5×10^4 cells per well one day prior to infection. Viral samples were diluted 10-fold in 5% cDMEM. An internal standard was included for each infection to ensure reproducible sensitivity for each plaque assay. Cells were incubated with virus for 1 h at 37°C at 5% CO_2. Plates were rocked every 15 min. Cells were then covered with an overlay composed of a 1:1 mix of 10% cDMEM and carboxymethyl cellulose (CMC; Sigma, Cat. No. C- 4888) supplemented with Gibco™ Amphotericin B (Thermo Fisher Scientific, Cat. No. 15290018). Cells were incubated for 8 days before staining with 0.5% methylene blue and plaques were counted.

RT-qPCR

RNA was isolated from infected cells harvested at indicated times by 10-minute incubation in TRIzol® reagent (Thermo Fisher Scientific, Cat. No. 15596026), followed by TURBO™ DNase (Invitrogen, Cat. No. AM2238) treatment according to manufacturer's protocols. RNA amplification and removal of DNA was confirmed by PCR amplification of the control host gene, 18S, in the presence or absence of reverse transcriptase. RNA presence and absence of DNA was confirmed with RT-PCR and PCR respectively. RT-PCR was performed using the OneStep RT-PCR kit (Qiagen, Cat. No. 210212) with the following conditions: (i) 50°C for 30 min, (ii) 95°C for 15 min, (iii) 40 cycles of 94°C for 30 sec, 52°C for 30 sec, and 72°C for 30 sec, (iv) 72°C for 10 min, and (v) hold at 4°C. PCR was performed using *Taq* DNA polymerase (Qiagen, Cat No. 201205) with the following conditions: (i) 95°C for 5 min, (ii) 40 cycles of 94°C for 30 sec, 52°C for 30 sec, and 72°C for 30 sec, (iii) 72°C for 10 min, and (iv) hold at 4°C.

RNA samples that showed no product following PCR amplification were deemed DNA-

free, and then converted to cDNA using random primers (250ng/uL) (Invitrogen, Cat.

No. 48190011) and SuperScript II reverse transcriptase (Invitrogen, Cat. No. 18064014)

following the manufacturer's protocol. 100 nanograms of cDNA was used for qPCR

analysis of the identified genes (Quantitect Primer Assay, Qiagen) using the iQ SYBR

green supermix (BioRad Cat. No.1708880) with the following conditions: (i) 95°C for 3

min, (ii) 40 cycles of 95°C for 15 sec, 60°C for 1 min, and (iii) 95°C for 15 sec, 60°C for

1 min, and 95°C for 15 sec. Amplification of viral genes were normalized to murine 18S

expression to calculate the relative difference of target gene expression using the Pfaffl

method, as previously described (49, 50) : Target primer efficiency^TargetΔCt/18S

primer efficiency^18SΔCt. A single product for each target was confirmed by melt curve

analysis. PCR primers are listed in **Table 1**.

Quantitative PCR for quantification of viral DNA replication

Infected cells were harvested by cell scraping at time (hpi) indicated. Harvested

cells underwent three freeze/thaw cycles and DNA was isolated using the DNeasy

Blood and Tissue Kit (Qiagen, Cat. No. 69504), with an overnight proteinase K

incubation and heat inactivation at 56°C. DNA was normalized to a concentration of 20

ng/uL in molecular grade water. qPCR analysis was done on 100ng of DNA using a

LightCycler 480 Probe Master-Mix kit (Roche, Cat. No. 04707494001) and a primer and

probe set specific to MHV68 gB to quantify the number of viral genome copies. Primers

and probes listed in Table 1. A glycoprotein B (gB) standard curve (51) was generated

using a gB plasmid dilution series ranging from 10^2 to 10^{10} copies diluted in background

DNA, with a limit of detection (LOD) of 100 copies.

Ethics Statement

All animal studies were performed in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. Studies were conducted in accordance with the University of Colorado Denver/Anschutz Institutional Animal Use and Care Committee under the Animal Welfare Assurance of Compliance policy (assurance no. D16-00171). All procedures were performed under isoflurane anesthesia, and all efforts were made to minimize suffering.

Single-cell RNA Sequencing

To characterize MHV68 transcription during primary, acute peritoneal macrophage infection *in vivo*, C57BL/6 mice were infected with $1x10^6$ PFU by intraperitoneal injection. Peritoneal cells were harvested 16 hours post-infection, stained with the live/dead discrimination dye Zombie Near-IR (Zombie NIR™, BioLegend, Cat. No. 423105), F4/80-AlexaFluor 674 (clone BM8, 1:200 dilution) in the presence of Fc receptor blockade (clone 2.4G2), and CCF2-AM (Thermo Fisher, Cat. No. K1023). Cells were sorted on a MoFlo Astrios, purifying live, single, F4/80+ cells that either had CCF-2 cleavage (indicating virus infection, defined by expression of the LANA::blac fusion protein) or lacking CCF-2 cleavage (indicating cells that were not infected or failed to express the LANA::blac fusion protein). Cells were subjected to standard 3'-based single cell RNA-seq analysis (10x Genomics), with cell processing and sequencing done in the University of Colorado Cancer Center Genomics Shared Resource (RRID: SCR_021984). Resulting data were processed as follows: Cell Ranger (v6.0.1) (52) was used to process the fastq files to cell and gene count tables

using unique molecule identifiers (UMIs) with the include-introns parameter. Because of

the difficulties counting viral reads, this was performed in a two-pass manner. In the first

pass, reads were aligned to a chimeric genome of mouse mm10 (GENCODE M23 gene

annotations) and MH636806.1 (no gene annotations, only positive and negative strand

alignment). The viral counts were stored in metadata and removed from the counts

matrix. Reads mapping to MH636806.1 in step 1 were aligned to the same chimeric

genome but in this case, the viral transcriptome as well as specific intergenic regions

were annotated (data not shown). The intergenic regions were determined as the

strand-specific sequences between genes with 1 base pair of padding on both ends. To

minimize reads overlapping with multiple viral genes, only the first 70bp of each read

were aligned and counted. The resulting counts matrix of viral alignment was appended

to the host gene counts from step 1.

The Seurat (v4.0.4) (53) pipeline was used for downstream quality control and

analysis. Cellranger-filtered data was read into Seurat. Host genes were removed if

identified in fewer than 10 cells while viral gene intergenic regions were removed if

found in no cells. Cells were removed if they expressed < 50 genes or viral regions, <

5000 UMIs, > 5 % total UMIs from mitochondria, or > 70000 UMIs. The filtered data was

normalized by dividing gene counts by total counts per cell and multiplied by 10,000

followed by natural-log transformation. The top 2,000 most variable genes were scaled

with total UMI and percentage mitochondria regressed out. These were used to

calculate principal components (PCs) with the top 20 used to perform Uniform Manifold

Approximation and Projection (UMAP) and determining the k-nearest neighbors and

clustering.

Clusters were identified by over-representation analysis and examining the top

enriched markers. Canonical cell type markers were plotting in UMAP space and using

dot plots. Heatmaps, dot plots, bar plots, scatter plots, and violin plots were generated

using Seurat and ggplot2 R packages (v3.4.0) (54). Groups of cells in heatmaps were

randomly subset to the size of the smallest group.

Flow Cytometry

3T12 and J774 cells were infected with MHV68 LANAβlac or MHV68.H.GFP at

an MOI of 1 PFU/cell and harvested at the indicated time points followed by staining for

flow cytometric analysis. Samples treated with the recombinant murine IL-4 were given

10% cDMEM with 10ng/mL of IL-4 eighteen hours prior to infection and replaced with

fresh media containing IL-4 immediately following infection. Cell suspensions were

stained with LIVE/DEAD fixable Near-IR Dead Cell stain kit (Invitrogen, Cat. No.

L10119) at a 1:1000 dilution in BSS wash (111mM Dextrose, 2mM KH_2PO_4, 10mM

Na_2HPO_4, 25.8 $CaCl_2 \cdot 2H_2O$, 2.7mM KCl, 137mM NaCl, 19.7mM $MgCl \cdot 6H_2O$, 16.6mM

$MgSO_4$). Cells were then stained for CCF2 or protein detection. Cells for CCF2

detection were stained with CCF2-AM (Thermo Fisher, Cat. No. K1023) at a 1:10

dilution following the manufacturer's protocol. To detect proteins, cells were stained with

a Fc receptor block (Fc Shield, Tonbo Biosciences, Cat. No. 70-0161), a rabbit antibody

against the MHV68 ORF4 protein (vRCA, 1:400 dilution) (gift of the Virgin Lab,

Washington University St. Louis (35, 49)) labeled with a Zenon R-phycoerythrin anti-

rabbit IgG reagent (Invitrogen, Cat. No. Z25307), per manufacturer's recommendations,

and a AF647-conjugated anti-mouse phospho-histone H2AX antibody (1:800 dilution)

(clone JBW30). Samples were fixed in 1% paraformaldehyde prior to analysis using the

Bio-Rad ZE5/YETI Cell Analyzer or an Agilent Novocyte Penteon flow cytometer. All

flow cytometry experiments included unstained, single-stain and full minus one controls

to define background fluorescence, fluorescent signal spread and compensation.

Light microscopy

At the indicated time points, 6-well plates of 3T12 or J774 cells were imaged on a

Nikon Eclipse Ti2 inverted light microscope equipped with Iris 15 camera (Photometris)

and use of NIS elements software. Plates were mounted on a stage heated to 37°C. A

10x phase objective was used to capture brightfield images. Immediately following

image acquisition, samples were harvested and processed for flow cytometric analysis.

Statistical Analysis and Software

Data analysis, graphing and statistical analyses were performed using GraphPad

Prism (version 9.4.0; GraphPad Software, San Diego, California USA,

www.graphpad.com). Flow cytometry data were analyzed using FlowJo (version 10.8.1.

Ashland, OR: Becton, Dickinson and Company; 2022). scRNA-seq data processing and

analysis were done as described above. Statistical significance was tested by unpaired,

nonparametric, Mann-Whitney t test or 1-way ANOVA for comparing replicate values

from experiments. RNA-Seq data have been deposited to NCBI GEO and will be made

publicly available upon manuscript acceptance.

Table 1: *Oligonucleotides for RT-PCR, PCR, and qPCR analysis*

Name	Accession number and Genome Position *	Sequence	Purpose
gB Forward Primer	MHV68: 17873…17892	5'—GGC CCA AAT TCA ATT TGC CT—3'	TaqMan qPCR for MHV68 genome
gB Reverse Primer	MHV68: 17924…17943	5'—CCC TGG ACA ACT CCT CAA GC—3'	TaqMan qPCR for MHV68genome

gB Probe	MHV68: 17896…17921	5'—ACA AGC TGA CCA CCA GCG TCA ACA AC—3'	TaqMan qPCR for MHV68genome
18S Forward Primer	Accession Number: NR_003278	5'— AGA TCA AAA CCA ACC CGG TGA—3'	RT-PCR and PCR for murine 18S gene
18S Reverse Primer	Accession Number: NR_003278	5'—GGT AAG AGC ATC GAG GGG GC—3'	RT-PCR and PCR for murine 18S gene
ORF73 SYBR G – Forward Primer	MHV68: 104092…104073	5'— GAG CCC CCT ACA GAG CCC CC—3'	SYBR Green qPCR for MHV68ORF73 gene
ORF73 SYBR G- Reverse Primer	MHV68: 103981…104000	5'— CAC CTT GCT CTG CAC CGG CA—3'	SYBR Green qPCR for MHV68ORF73 gene
ORF6 SYBR G – Forward Primer	MHV68: 11699…11783	5'— ATG TTG TCA GCA CCC ATG AA—3'	SYBR Green qPCR for MHV68ORF6 gene
ORF6 SYBR G – Reverse Primer	MHV68: 11802…11783	5'— AAG GGC AGT AGC TGG TCA GA—3'	SYBR Green qPCR for MHV68ORF6 gene
M7 SYBR G – Forward Primer	MHV68: 70109…70128	5'—GCT CCT GCT GAC ACA TCA GA—3'	SYBR Green qPCR for MHV68M7 gene
M7 SYBR G – Reverse Primer	MHV68: 70185…70166	5'—TGG GGT TTG GAC TCT GTA GG—3'	SYBR Green qPCR for MHV68M7 gene
MHV68 ORF75A iQ SYBR - Forward Primer	MHV68: 114964…114945	5'—ACC TGA CAC CCC AAG AAC AG—3'	SYBR Green qPCR for MHV68ORF75A gene
MHV68 ORF75A iQ SYBR - Reverse Primer	MHV68: 114834…114853	5'—GAG CAC TTT TGG TGG AGA GC—3'	SYBR Green qPCR for MHV68ORF75A gene
MHV68 ORF75B iQ SYBR - Forward Primer	MHV68: 113267…113248	5'—CAG CCT CTC AAC CTT TCC AG—3'	SYBR Green qPCR for MHV68ORF75B gene
MHV68 ORF75B iQ SYBR - Reverse Primer	MHV68: 113093…113112	5'—ATA GGA GCC ACC GTT GAT TG—3'	SYBR Green qPCR for MHV68.ORF73βla ORF75B
MHV68 ORF 75C iQ SYBR - Forward Primer	MHV68: 109930…109911	5'—AAG ACA CAG AGG CTG GGA GA—3'	SYBR Green qPCR for MHV68ORF75C gene
MHV68 ORF75C iQ SYBR - Reverse Primer	MHV68: 109758…109777	5'—GAC GGG TAG ACG TGG TGA CT—3'	SYBR Green qPCR for MHV68ORF75C gene

BLAC SYBR G – Forward Primer	MHV68.ORF73βla: 104867…104886	5'—GCT ATG TGG CGC GGT ATT AT—3'	SYBR Green qPCR for MHV68.ORF73βla Blac gene
BLAC SYBR G – Reverse Primer	MHV68.ORF73βla: 104701…104720	5'—AAG TTG GCC GCA GTG TTA TC—3'	SYBR Green qPCR for MHV68.ORF73βla Blac gene

570   MHV68 genome coordinates refer to NC_001826. In the case of MHV68.ORF73βla Bla

571   gene, sequences correspond to NCBI GenBank: MH636806.1, the complete genome

572   sequence of MHV68 containing the ORF73bla gene fusion.

Acknowledgments:

We thank Dr. Brian Russo's lab, especially Jenna Vickery for the use and training of

their Nikon Eclipse Ti2 inverted light microscope, and Christine Childs, Kristina Terrell

and Dmitry Baturin of the University of Colorado Cancer Center Flow Cytometry Shared

Resource for training and assistance with flow cytometry. This study was supported by

National Institutes of Health grant R01 AI157201 awarded to LvD and ETC, with

additional support by the National Institutes of Health Cancer Center Support Grant

(P30CA046934), including assistance from the Bioinformatics and Biostatistics Shared

Resource (RRID: SCR_021983), Flow Cytometry Shared Resource (RRID:

SCR_022035), and Genomics Shared Resource (RRID: SCR_021984).

References:

1. Longnecker R, Neipel F. 2007. Introduction to the human gamma-herpesviruses. *In* Arvin A, Campadelli-Fiume G, Mocarski E, Moore PS, Roizman B, Whitley R, Yamanishi K (ed), Human Herpesviruses: Biology, Therapy, and Immunoprophylaxis, Cambridge.
2. Mariggio G, Koch S, Schulz TF. 2017. Kaposi sarcoma herpesvirus pathogenesis. Philos Trans R Soc Lond B Biol Sci 372.
3. Young LS, Yap LF, Murray PG. 2016. Epstein-Barr virus: more than 50 years old and still providing surprises. Nat Rev Cancer 16:789-802.
4. Virgin HWt, Latreille P, Wamsley P, Hallsworth K, Weck KE, Dal Canto AJ, Speck SH. 1997. Complete sequence and genomic analysis of murine gammaherpesvirus 68. J Virol 71:5894-904.
5. Wang Y, Tibbetts SA, Krug LT. 2021. Conquering the Host: Determinants of Pathogenesis Learned from Murine Gammaherpesvirus 68. Annu Rev Virol 8:349-371.
6. Cieniewicz B, Santana AL, Minkah N, Krug LT. 2016. Interplay of Murine Gammaherpesvirus 68 with NF-kappaB Signaling of the Host. Front Microbiol 7:1202.
7. Flano E, Husain SM, Sample JT, Woodland DL, Blackman MA. 2000. Latent murine gamma-herpesvirus infection is established in activated B cells, dendritic cells, and macrophages. J Immunol 165:1074-81.
8. Flano E, Kim IJ, Moore J, Woodland DL, Blackman MA. 2003. Differential gamma-herpesvirus distribution in distinct anatomical locations and cell subsets during persistent infection in mice. J Immunol 170:3828-34.
9. Sunil-Chandra NP, Efstathiou S, Nash AA. 1992. Murine gammaherpesvirus 68 establishes a latent infection in mouse B lymphocytes in vivo. J Gen Virol 73 (Pt 12):3275-9.
10. Babcock GJ, Decker LL, Volk M, Thorley-Lawson DA. 1998. EBV persistence in memory B cells in vivo. Immunity 9:395-404.
11. Aalam F, Nabiee R, Castano JR, Totonchy J. 2020. Analysis of KSHV B lymphocyte lineage tropism in human tonsil reveals efficient infection of CD138+ plasma cells. PLoS Pathog 16:e1008968.
12. Frederico B, Milho R, May JS, Gillet L, Stevenson PG. 2012. Myeloid infection links epithelial and B cell tropisms of Murid Herpesvirus-4. PLoS Pathog 8:e1002935.
13. Weck KE, Kim SS, Virgin HI, Speck SH. 1999. Macrophages are the major reservoir of latent murine gammaherpesvirus 68 in peritoneal cells. J Virol 73:3273-83.
14. Lawler C, Milho R, May JS, Stevenson PG. 2015. Rhadinovirus host entry by co-operative infection. PLoS Pathog 11:e1004761.
15. Rosa GT, Gillet L, Smith CM, de Lima BD, Stevenson PG. 2007. IgG fc receptors provide an alternative infection route for murine gamma-herpesvirus-68. PLoS One 2:e560.
16. Tarakanova VL, Leung-Pineda V, Hwang S, Yang CW, Matatall K, Basson M, Sun R, Piwnica-Worms H, Sleckman BP, Virgin HWt. 2007. Gamma-herpesvirus

kinase actively initiates a DNA damage response by inducing phosphorylation of
H2AX to foster viral replication. Cell Host Microbe 1:275-86.
17.   Tan CSE, Lawler C, Stevenson PG. 2017. CD8+ T cell evasion mandates CD4+
T cell control of chronic gamma-herpesvirus infection. PLoS Pathog
13:e1006311.
18.   Frederico B, Chao B, May JS, Belz GT, Stevenson PG. 2014. A murid gamma-
herpesviruses exploits normal splenic immune communication routes for
systemic spread. Cell Host Microbe 15:457-70.
19.   Frederico B, Chao B, Lawler C, May JS, Stevenson PG. 2015. Subcapsular sinus
macrophages limit acute gammaherpesvirus dissemination. J Gen Virol 96:2314-
2327.
20.   Suarez AL, van Dyk LF. 2008. Endothelial cells support persistent
gammaherpesvirus 68 infection. PLoS Pathog 4:e1000152.
21.   Sunil-Chandra NP, Efstathiou S, Nash AA. 1993. Interactions of murine
gammaherpesvirus 68 with B and T cell lines. Virology 193:825-33.
22.   Weck KE, Barkon ML, Yoo LI, Speck SH, Virgin HI. 1996. Mature B cells are
required for acute splenic infection, but not for establishment of latency, by
murine gammaherpesvirus 68. J Virol 70:6775-80.
23.   Nealy MS, Coleman CB, Li H, Tibbetts SA. 2010. Use of a virus-encoded
enzymatic marker reveals that a stable fraction of memory B cells expresses
latency-associated nuclear antigen throughout chronic gammaherpesvirus
infection. J Virol 84:7523-34.
24.   Diebel KW, Oko LM, Medina EM, Niemeyer BF, Warren CJ, Claypool DJ,
Tibbetts SA, Cool CD, Clambey ET, van Dyk LF. 2015. Gammaherpesvirus small
noncoding RNAs are bifunctional elements that regulate infection and contribute
to virulence in vivo. mBio 6:e01670-14.
25.   Cheng BY, Zhi J, Santana A, Khan S, Salinas E, Forrest JC, Zheng Y, Jaggi S,
Leatherwood J, Krug LT. 2012. Tiled microarray identification of novel viral
transcript structures and distinct transcriptional profiles during two modes of
productive murine gammaherpesvirus 68 infection. J Virol 86:4340-57.
26.   Forrest JC, Speck SH. 2008. Establishment of B-cell lines latently infected with
reactivation-competent murine gammaherpesvirus 68 provides evidence for viral
alteration of a DNA damage-signaling cascade. J Virol 82:7688-99.
27.   Berger JN, Sanford B, Kimball AK, Oko LM, Kaspar RE, Niemeyer BF, Jones KL,
Clambey ET, van Dyk LF. 2020. Redefining De Novo Gammaherpesvirus
Infection Through High-Dimensional, Single-Cell Analysis of Virus and Host.
bioRxiv doi:10.1101/2020.08.11.203117:2020.08.11.203117.
28.   Wang G, Zarek C, Chang T, Tao L, Lowe A, Reese TA. 2021. Th2 Cytokine
Modulates Herpesvirus Reactivation in a Cell Type Specific Manner. J Virol
doi:10.1128/JVI.01946-20.
29.   Reese TA, Wakeman BS, Choi HS, Hufford MM, Huang SC, Zhang X, Buck MD,
Jezewski A, Kambal A, Liu CY, Goel G, Murray PJ, Xavier RJ, Kaplan MH,
Renne R, Speck SH, Artyomov MN, Pearce EJ, Virgin HW. 2014. Helminth
infection reactivates latent gamma-herpesvirus via cytokine competition at a viral
promoter. Science 345:573-7.

30.   Kulkarni A, Anderson AG, Merullo DP, Konopka G. 2019. Beyond bulk: a review
674   of single cell transcriptomics methodologies and applications. Curr Opin
675   Biotechnol 58:129-136.
676   31.   Van Skike ND, Minkah NK, Hogan CH, Wu G, Benziger PT, Oldenburg DG, Kara
677   M, Kim-Holzapfel DM, White DW, Tibbetts SA, French JB, Krug LT. 2018. Viral
678   FGARAT ORF75A promotes early events in lytic infection and
679   gammaherpesvirus pathogenesis in mice. PLoS Pathog 14:e1006843.
680   32.   Zhang Z, Schwartz S, Wagner L, Miller W. 2000. A greedy algorithm for aligning
681   DNA sequences. J Comput Biol 7:203-14.
682   33.   Lalezari JP, Stagg RJ, Jaffe HS, Hitchcock MJ, Drew WL. 1996. A preclinical and
683   clinical overview of the nucleotide-based antiviral agent cidofovir (HPMPC). Adv
684   Exp Med Biol 394:105-15.
685   34.   Andrei G, Topalis D, De Schutter T, Snoeck R. 2015. Insights into the
686   mechanism of action of cidofovir and other acyclic nucleoside phosphonates
687   against polyoma- and papillomaviruses and non-viral induced neoplasia. Antiviral
688   Res 114:21-46.
689   35.   Kapadia SB, Molina H, van Berkel V, Speck SH, Virgin HWt. 1999. Murine
690   gammaherpesvirus 68 encodes a functional regulator of complement activation. J
691   Virol 73:7658-70.
692   36.   Riggs JB, Medina EM, Perrenoud LJ, Bonilla DL, Clambey ET, van Dyk LF, Berg
693   LJ. 2021. Optimized Detection of Acute MHV68 Infection With a Reporter System
694   Identifies Large Peritoneal Macrophages as a Dominant Target of Primary
695   Infection. Front Microbiol 12:656979.
696   37.   Ramaswami R, Tagawa T, Mahesh G, Serquina A, Koparde V, Lurain K, Dremel
697   S, Li X, Mungale A, Beran A, Ohler ZW, Bassel L, Warner A, Mangusan R, Widell
698   A, Ekwede I, Krug LT, Uldrick TS, Yarchoan R, Ziegelbauer JM. 2022.
699   Transcriptional analysis identifies overlapping and tissue-distinct profiles between
700   Kaposi sarcoma tumors of the skin and gastrointestinal tract. bioRxiv
701   doi:10.1101/2022.03.18.484923:2022.03.18.484923.
702   38.   Tarakanova VL, Molleston JM, Goodwin M, Virgin HWt. 2010. MHV68
703   complement regulatory protein facilitates MHV68 replication in primary
704   macrophages in a complement independent manner. Virology 396:323-8.
705   39.   Mounce BC, Tsan FC, Droit L, Kohler S, Reitsma JM, Cirillo LA, Tarakanova VL.
706   2011. Gammaherpesvirus gene expression and DNA synthesis are facilitated by
707   viral protein kinase and histone variant H2AX. Virology 420:73-81.
708   40.   Goodwin MM, Canny S, Steed A, Virgin HW. 2010. Murine gammaherpesvirus
709   68 has evolved gamma interferon and stat1-repressible promoters for the lytic
710   switch gene 50. J Virol 84:3711-7.
711   41.   Goodwin MM, Molleston JM, Canny S, Abou El Hassan M, Willert EK, Bremner
712   R, Virgin HW. 2010. Histone deacetylases and the nuclear receptor corepressor
713   regulate lytic-latent switch gene 50 in murine gammaherpesvirus 68-infected
714   macrophages. J Virol 84:12039-47.
715   42.   Lange PT, Schorl C, Sahoo D, Tarakanova VL. 2018. Liver X Receptors
716   Suppress Activity of Cholesterol and Fatty Acid Synthesis Pathways To Oppose
717   Gammaherpesvirus Replication. mBio 9.

718   43.   Aurubin CA, Knaack DA, Sahoo D, Tarakanova VL. 2021. Low-Density
719   Lipoprotein Receptor Suppresses the Endogenous Cholesterol Synthesis
720   Pathway To Oppose Gammaherpesvirus Replication in Primary Macrophages. J
721   Virol 95:e0064921.
722   44.   Steed A, Buch T, Waisman A, Virgin HWt. 2007. Gamma interferon blocks
723   gammaherpesvirus reactivation from latency in a cell type-specific manner. J
724   Virol 81:6134-40.
725   45.   Zarek CM, Dende C, Coronado J, Pendse M, Dryden P, Hooper LV, Reese TA.
726   2022. Coinfection with Intestinal Parasite Expands Resident Macrophages and
727   Impairs Control of Chronic Herpesvirus Infection. bioRxiv
728   doi:10.1101/2022.10.05.510926:2022.10.05.510926.
729   46.   Freeman ML, Lanzer KG, Cookenham T, Peters B, Sidney J, Wu TT, Sun R,
730   Woodland DL, Sette A, Blackman MA. 2010. Two kinetic patterns of epitope-
731   specific CD8 T-cell responses following murine gammaherpesvirus 68 infection. J
732   Virol 84:2881-92.
733   47.   Gredmark-Russ S, Cheung EJ, Isaacson MK, Ploegh HL, Grotenbreg GM. 2008.
734   The CD8 T-cell response against murine gammaherpesvirus 68 is directed
735   toward a broad repertoire of epitopes from both early and late antigens. J Virol
736   82:12205-12.
737   48.   Freeman ML, Burkum CE, Cookenham T, Roberts AD, Lanzer KG, Huston GE,
738   Jensen MK, Sidney J, Peters B, Kohlmeier JE, Woodland DL, van Dyk LF, Sette
739   A, Blackman MA. 2014. CD4 T cells specific for a latency-associated gamma-
740   herpesvirus epitope are polyfunctional and cytotoxic. J Immunol 193:5827-34.
741   49.   Oko LM, Kimball AK, Kaspar RE, Knox AN, Coleman CB, Rochford R, Chang T,
742   Alderete B, van Dyk LF, Clambey ET. 2019. Multidimensional analysis of
743   Gammaherpesvirus RNA expression reveals unexpected heterogeneity of gene
744   expression. PLoS Pathog 15:e1007849.
745   50.   Pfaffl MW. 2001. A new mathematical model for relative quantification in real-
746   time RT-PCR. Nucleic Acids Res 29:e45.
747   51.   Coomes SM, Farmen S, Wilke CA, Laouar Y, Moore BB. 2011. Severe
748   gammaherpesvirus-induced pneumonitis and fibrosis in syngeneic bone marrow
749   transplant mice is related to effects of transforming growth factor-beta. Am J
750   Pathol 179:2382-96.
751   52.   Zheng GX, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, Ziraldo SB,
752   Wheeler TD, McDermott GP, Zhu J, Gregory MT, Shuga J, Montesclaros L,
753   Underwood JG, Masquelier DA, Nishimura SY, Schnall-Levin M, Wyatt PW,
754   Hindson CM, Bharadwaj R, Wong A, Ness KD, Beppu LW, Deeg HJ, McFarland
755   C, Loeb KR, Valente WJ, Ericson NG, Stevens EA, Radich JP, Mikkelsen TS,
756   Hindson BJ, Bielas JH. 2017. Massively parallel digital transcriptional profiling of
757   single cells. Nat Commun 8:14049.
758   53.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, Lee MJ,
759   Wilk AJ, Darby C, Zager M, Hoffman P, Stoeckius M, Papalexi E, Mimitou EP,
760   Jain J, Srivastava A, Stuart T, Fleming LM, Yeung B, Rogers AJ, McElrath JM,
761   Blish CA, Gottardo R, Smibert P, Satija R. 2021. Integrated analysis of
762   multimodal single-cell data. Cell 184:3573-3587 e29.

763   54.   Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag
764   New York.
765

Figure 1. MHV68-infected J774 macrophages are deficient in virus replication relative to permissive 3T12 fibroblasts, despite evidence of equivalent virus entry and LANA gene expression. Analysis of MHV68 infection (MOI=1 PFU/cell) comparing 3T12 fibroblasts and J774 macrophages. (A) Representative brightfield images of infected 3T12 or J774 cells. 3T12 cells show early signs of CPE, including cell rounding (24 hpi) with extensive monolayer destruction (72 hpi), whereas J774 cells show minimal CPE. (B) Virus replication as defined by plaque assay. Data depict mean +/- SEM, with limit of detection at 10^2 PFU/mL (t=0, 24, and 48 are from 2, 6, and 6 independent experiments with one biological sample per experiment). (C) Flow cytometric analysis of LANAβlac expression defined by cleavage of the CCF2 fluorescent beta-lactamase substrate. Data depict events that were single, viable cells, with values in the right gate defining the frequency of cells with CCF2 substrate cleavage, an indicator of LANAβlac expression. Samples infected with WT MHV68 (left panel) do not encode LANAβlac and define background cleavage. Representative flow cytometry plots were defined as samples that were closest to the mean frequency in panel D. (D) Quantification of LANAβlac + cell frequency defined by flow cytometry, showing mean +/- SEM, as in panel C. Data are from 3 independent experiments with 3 biological replicates per experiment. Data for 3T12 cells at 72 hpi was not assessed due to extensive cell destruction. Statistical analysis was performed using unpaired t test with statistically significant differences as indicated, * $p<0.05$, ** $p<0.01$, **** $p<0.0001$.

Vragel et al
Fig. 1

Figure 2. MHV68-infected J774 macrophages are deficient in viral DNA replication and viral gene expression relative to infected fibroblasts. Analysis of MHV68 replication and gene expression (MOI=1 PFU/cell) comparing 3T12 fibroblasts and J774 macrophages at the indicated times. (A) Viral infection quantified by viral genome copies via q-PCR. (B-C) Quantitative analysis of viral RNA expression of ORF73 (immediate early gene), ORF6 (an early/late gene), and M7 (late gene) by qRT-PCR, comparing (top row) 3T12 and J774 cells or (bottom row) time-dependent changes within J774s. "Relative Difference" defined as quantification of target gene transcript in comparison to reference gene transcript (host 18S) as described in Methods. Data are from 3 independent experiments with 3 biological replicates per experiment, measured in technical triplicates. The value shown is the mean +/- SEM of each experiment. Statistical analysis used unpaired t test with statistically significant differences as indicated. ** p<0.01, ****p<0.0001. ns, not statistically significant.

Figure 3. MHV68-infected J774 macrophages contain a small subset of cells that demonstrate lytic infection profiles. Analysis of MHV68 infection and lytic cycle profiles with either MHV68.H.GFP or MHV68.LANAβlac (MOI=1 PFU/cell) comparing 3T12 fibroblasts and J774 macrophages. (A) Flow cytometric analysis of virus-expressed GFP in 3T12 or J774 cells. Data depict events that were single, viable cells, with values defining the frequency of cells with GFP expression, an indicator of MHV68.H.GFP infection, with gating strategy in Supp. Fig. 2. Samples infected with WT MHV68.LANAβlac (left column), without the addition of the CCF2 fluorescent substrate, defined background fluorescence. (B) Quantification of GFP + cell frequency defined by flow cytometry. Symbols depict individual experimental values, with data showing mean +/- SEM. (C) Flow cytometric analysis of MHV68 lytic viral protein vRCA and phosphorylated host γH2AX in either 3T12 (top) or J774 (bottom) cells at the indicated timepoints, comparing mock (left) or WT MHV68.LANAβlac infected cells. Data depict events that were single, viable cells, with values defining the frequency of cells with vRCA and/or phosphorylated host γH2AX, indicative of lytic MHV68 infection, with gating strategy presented in Supp. Fig. 3. Data for 3T12 cells at 48 hpi was not assessed due to extensive cell destruction. (D) Quantification of positive protein expression defined by flow cytometry, showing mean +/- SEM, as in panel C. Representative images were defined as samples that were closest to the mean frequency. Data are from 3 independent experiments with 2 biological experiments per experiment. Statistical analysis used unpaired t test with statistically significant differences as indicated. ****p<0.0001, **p<0.01.

Vragel et al
Fig. 3

Figure 4. J774 macrophages are capable of supporting lytic MHV68 replication following pre-treatment with IL4, a known potentiator of virus replication in macrophages. Analysis of the effects of cytokine IL-4 on MHV68 infection and genome replication (MOI=1 PFU/cell) in 3T12 fibroblasts and J774 macrophages. (A) Flow cytometric analysis of MHV68.H.GFP infection in 3T12 (top row), untreated J774 (middle) or IL4-treated J774 cells (10 ng/mL, 18 hour pre-treatment with fresh IL4 added post-infection; bottom row). Data depict events that were single, viable cells, with values defining the frequency of cells with GFP expression, an indicator of viral infection, transcription and translation. Representative images were defined as samples that were closest to the mean frequency. (B) Quantification of GFP+ cell frequency defined by flow cytometry, as in panel A, with individual symbols representing independent biological samples and data showing mean +/- SEM. Data are from 3 independent experiments with 2 biological replicates per experiment, measured in technical duplicates. (C) Quantitation of the impact of IL4 treatment on MHV68.LANAβlac viral DNA replication in MHV68-infected J774 cells quantified by qPCR. Cells were pre-treated with 0, 10 or 100 ng/mL murine IL-4 for 16 hours, with fresh IL4 added post-infection, and cells harvested for DNA at 0, 24, or 48hpi. Data depict mean +/- SEM, with individual symbols representing independent biological samples. Data are from 3 independent experiments with 3 biological replicates per experiment measured in triplicate. Statistical analysis used (B) unpaired t test or (C) 1-way ANOVA comparing samples collected at the same timepoint untreated versus cytokine treated, with statistically significant differences as indicated, **p<0.01, ****p<0.0001.

Vragel et al
Fig. 4

Vragel et al
Fig. 5

Figure 5. Single cell RNA-seq analysis of primary MHV68-infected LANAβlac+ peritoneal cells harvested 16 hours post infection in vivo. C57BL/6J mice were infected with 1x10^6 PFU of WT MHV68.LANAblac virus, with MHV68-infected, LANAblac+ peritoneal macrophages harvested and sort-purified at 16 hpi and subjected to scRNA-seq analysis. Data focus on cell types identified based on Seurat-designated clusters as outlined in Supplemental Figures 4-5. (A-B) Visualization of cell types present within scRNA-seq dataset, with cell identities based on lineage marker genes depicted in panel B (C) Quantification of cell subsets defined by scRNA-seq identifies macrophages as the dominant population. (D-F) Quality control metrics across cell subsets, including UMI Count (D), Features (E), % mitochondrial reads (F), and S score, a proliferation-associated gene signature (G). (H) UMAP visualization of cells that contain at least 1 viral UMI per cell identified by red. (I) % of cell subsets stratified by extent of viral UMIs detected per cell, including cells with >0.1% total reads that are viral UMIs (bright red), cells with detectable viral UMIs accounting for <0.1% of total reads (pale red), or no viral UMIs detected (gray). In total, 56 cells had >0.1% of total UMIs contributed by viral UMIs, including 23 of 9101 macrophages (0.25%), 25 of 25 macrophages with lytic gene expression (100%) and 8 of 70 dendritic cells (11.43%). Data are from 9,537 cells, with cell subsets containing between 36-9,101 cells (as in panel I).

Vragel et al
Fig. 6

Figure 6. scRNA-seq analysis of MHV68 transcription within LANAβlac+ peritoneal macrophages harvested ex vivo 16 hours post-infection. C57BL/6J mice were infected with 1×10^6 PFU of WT MHV68.LANAβlac virus, with MHV68-infected, LANAβlac+ peritoneal macrophages harvested and sort-purified at 16 hpi and subjected to scRNA-seq analysis. Analysis focuses on macrophage populations identified in Figure 5. (A-C) Frequency of cells with detectable reads mapping to MHV68 genes, comparing (A) macrophages, (B) macrophages with proliferative gene signature, or (C) macrophages with lytic gene expression, with number of cells for each population indicated. Viral genes are indicated on the x axis, arranged from left to right based on the gene arrangement for the published MHV68 genome, with the frequency of cells positive for each viral gene depicted on the y axis. To facilitate comparison of gene expression across the viral genome and cell populations, circles are color-coded to identify values that are in the indicated ranges (as defined in the key, panel A). Dotted lines indicate different threshholds for viral gene positivity, with these threshold values arbitrarily based on data. For panel C, only genes with >75% positivity were color-coded, given the limited number of events in this cell subset. (D) Frequency of macrophages based on viral gene expression, with a focus on the inter-relationship between ORF73, ORF75A, ORF75B, and ORF75C expression within single cells. Data in panel D shows data from 9,174 macrophages including both proliferating and lytic subsets. The frequency of viral UMI+ events differed between cell subsets in panels A-C (macrophages, 82.79% viral UMI+; macrophage proliferating (52.08% viral UMI+; macrophage lytic, 100% viral UMI+).

Figure 7. MHV68-infected J774 macrophages express multiple ORF75 genes with ORF73 and ORF75A characterized by similar expression characteristics. Analysis of MHV68.LANAβlac gene expression (MOI=1 PFU/cell) comparing 3T12 fibroblasts and J774 macrophages. (A) Quantitation of viral RNAs for ORF73 (immediate early), ORF75A (immediate early/early-late), ORF75B (early) and ORF75C (late) genes during MHV68 infection by qRT-PCR, comparing expression in 3T12 and J774 cells at the indicated timepoints. Data are from 3 independent experiments with 3 biological replicates per experiment (depicted by individual symbols), measured in technical triplicates, with data showing mean +/- SEM for each timepoint and sample. Statistical analysis was done using unpaired t test with statistically significant differences as indicated. ** p<0.01,***p<0.001 ****p<0.0001. (B) Quantitative RNA analysis of indicated genes in 3T12 and J774 cells, with or without treatment using a viral inhibitor (CDV or PAA). For panels A-B, "Relative Difference" defined as quantification of target gene transcript in comparison to reference gene transcript (host 18S) as described in Methods. Data for 3T12 cell line are from 1 independent experiment, J774 cell line data are from 2 independent experiments, with three biological replicates per experiment, measured in technical triplicates.

A.

Supplementary Figure 1. Blac expression is equivalent to ORF73 expression during infection with WT MHV68 LANAβlac. Analysis of MHV68 gene expression after WT MHV68.LANAβlac infection (MOI=1 PFU/cell). Samples are from experiments detailed in Fig. 2B at 24hpi, with ORF73 data presenting one replicate contained in Fig. 2B. Viral RNA analysis of ORF73 (immediate early gene) and beta-lactamase (blac) genes during MHV68 infection in J774 and 3T12 cell lines by qRT-PCR. "Relative Difference" defined as quantification of target gene transcript in comparison to reference gene transcript (host 18S) as described in Methods. Data are from 1 independent experiments with 3 biological replicates per experiment, measured in technical triplicates. Symbols depict values from independent biological samples, with mean +/- SEM for each gene and condition depicted. Statistical analysis was done using unpaired t test with statistically significant differences as indicated. ns, no statistical significance detected.

A.

Supplementary Figure 2. Gating of flow cytometry analysis of MHV68 LANAβlac and WT.MHV68.H.GFP. 3T12 cells infected with MHV68 LANAblac or WT.MHV68.H.GFP were analyzed by flow cytometry, as in Figure 3A. Samples were first analyzed by looking at the area of the forward scatter vs. the area of the side scatter for all cells. A gate was drawn around a majority of cells to exclude debris and named "generous cells." Doublet discrimination was then done by analyzing the cells from the "generous cells" gate by height vs. area for both forward and side scatter. Gates along the diagonal define single cells since single cells express a 1:1 height-to-area ratio. These singlets were then analyzed via a Live/Dead Fixable stain. A gate was drawn around live cells which uptake less of the fixable stain compared to dead cells. Live cells were then analyzed for GFP fluorescence as measured by the GFP detection channel. Cells infected with MHV68.LANAβlac, without additional of the CCF2 fluorescent substrate, identified background fluorescence, in contrast to samples infected with WT.MHV68.HG which encoded GFP. Controls included single-stained samples with only one stain to define the bounds of a gate for the fluorophore of interest.

A.

pH2AX single stain   vRCA single stain   pH2AX/vRCA

Supplementary Figure 3. Gating of flow cytometry analysis of MHV68 LANAβlac infected J774 cells for lytic protein markers. J774 cells were infected as detailed in Figure 3C. Samples were first analyzed by looking at the area of the forward scatter vs. the area of the side scatter for all cells. A gate was drawn around a majority of cells to exclude debris and named "generous cells." Doublet discrimination was then done by analyzing the cells in the "generous cells" by height vs. area for both forward and side scatter. Gates along the diagonal indicated single cells since single cells express a 1:1 height-to-area ratio. These singlets were then analyzed via a Live/Dead Fixable stain. A gate was drawn around live cells which uptake less of the fixable stain compared to dead cells. Live cells were then analyzed for antibody binding for lytic protein markers. Controls included single-stained samples with only one stain to define the bounds of a gate for the fluorophore of interest (i.e. "single stain"). Further verification for fluorophores vRCA:PE and pH2AX:AF647 were done with full minus one (FMO) controls. FMO control samples included the whole panel of fluorophores except for one to define the background fluorescence in the fluorescent channel of the excluded fluorophore.

Vragel et al
Supp. Fig. 4

Supplemental Figure 4. Seurat-defined clustering of primary LANAβlac + peritoneal cells harvested 16 hours post-infection and subjected to scRNA-seq. Data focus on Seurat-based designation of 14 clusters of these data, corresponding to data shown in Figure 5. (A) UMAP visualization of Seurat-defined clusters. (B) Visualization of the frequency and magnitude of host gene expression across clusters, with cell type designation indicated on right side. (C-F) Quality control metrics across the 14 clusters, including (C) UMI Count, (D) Features, (E) % mitochondrial reads, and (F) S score, a proliferation-associated gene signature. (G) UMAP visualization of cells that contain at least 1 viral UMI per cell identified by red. (H) % of cells per cluster that contain at least 1 viral UMI. (I-J) % viral UMIs of total UMIs across clusters (panel I, y axis range from 0-50%), or (J) focused only on the majority of virus positive cells which have <0.1% viral UMIs per cell (excluding 56 cells with viral UMI > 0.1). Data are from 9,537 cells, with clusters containing between 25-2,532 cells. Panel G is included to provide a frame of reference and depicts the same data as in Figure 5, panel H.

Supplemental Figure 5. Visualization of genes associated with distinct cell type designations designations in single-cell RNAseq study. Data correspond to that shown in Figure 4-5, Supplemental Figure 4. (A) Seurat-defined clustering of primary peritoneal LANAβlac+ cells harvested 16 hours post-infection and subjected to scRNA-seq, identifying 14 clusters, visualized by UMAP dimensionality reduction. (B) Gene expression visualization on UMAP dimensionality reduction. Genes depicted are lineage defining genes used to identify cell types in Figure 4. Maximal expression is designated by purple, with absence of expression designated by gray. (A) is provided as a reference and is also shown as Supplemental Figure 4A. Adgre1 encodes the F4/80 protein; Itgam encodes the CD11b protein. Both of these proteins are markers of peritoneal macrophages. Proliferative gene signature includes expression of Top2a, Mki67. Lytic gene expression characterized by ORF54, ORF7. B cell signature defined by Cd19, CD79a. Monocyte cell signature defined by Ly6c2, Ccr2. Dendritic cell signature defined by Itgax, H2-Ab1. T cell signature defined by Cd3d, Thy1.