Hip thrust and back squat training elicit similar gluteus muscle hypertrophy and transfer similarly to the deadlift

Daniel L. Plotkin[1,*], Merlina A. Rodas[2], Andrew D. Vigotsky[3,4], Mason C. McIntosh[1], Emma Breeze[1], Rachel Ubrik[1], Cole Robitzsch[1], Anthony Agyin-Birikorang[1], Madison L. Mattingly[1], J. Max Michel[1], Nicholas J. Kontos[1], Andrew D. Frugé[5], Christopher M. Wilburn[1], Wendi H. Weimar[1], Adil Bashir[6], Ronald J. Beyers[6], Menno Henselmans[7], Bret M. Contreras[8], Michael D. Roberts[1,*]

Affiliations: [1]School of Kinesiology, Auburn University, Auburn, AL, USA; [2]Department of Psychological Sciences, Auburn, AL, USA; [3]Departments of Biomedical Engineering and Statistics, Evanston, IL, USA; [4]Department of Neuroscience, Northwestern University, Chicago, IL, USA; [5]College of Nursing, Auburn University, Auburn, AL, USA; [6]MRI Research Center, Auburn University, Auburn AL, USA; [7]International Scientific Research Foundation for Fitness and Nutrition, Amsterdam, Netherlands; [8]BC Strength, San Diego, CA, USA

[*]Co-correspondence to:

Daniel L. Plotkin, MS
PhD student, School of Kinesiology
Auburn University
E-mail: dzp0092@auburn.edu

Michael D. Roberts, PhD
Professor, School of Kinesiology
Auburn University
E-mail: mdr0024@auburn.edu

ABSTRACT

Purpose: We examined how set-volume equated resistance training using either the back squat (SQ) or hip thrust (HT) affected hypertrophy and various strength outcomes.

Methods: Untrained college-aged participants were randomized into HT or SQ groups. Surface electromyograms (sEMG) from the right gluteus maximus and medius muscles were obtained during the first training session. Participants completed nine weeks of supervised training (15–17 sessions), before and after which we assessed muscle cross-sectional area (mCSA) via magnetic resonance imaging and strength via three-repetition maximum (3RM) testing and an isometric wall push test.

Results: Glutei mCSA growth was similar across both groups. Estimates [(−) favors HT; (+) favors SQ] modestly favored the HT compared to SQ for lower [effect ± SE, -1.6 ± 2.1 cm^2], mid [-0.5 ± 1.7 cm^2], and upper [-0.5 ± 2.6 cm^2], but with appreciable variance. Gluteus medius+minimus [-1.8 ± 1.5 cm^2] and hamstrings [0.1 ± 0.6 cm^2] mCSA demonstrated little to no growth with small differences between groups. Thigh mCSA changes were greater in SQ for the quadriceps [3.6 ± 1.5 cm^2] and adductors [2.5 ± 0.7 cm^2]. Squat 3RM increases favored SQ [14 ± 2.5 kg] and hip thrust 3RM favored HT [-26 ± 5 kg]. 3RM deadlift [0 ± 2 kg] and wall push strength [-7 ± 13 N] similarly improved. All measured gluteal sites showed greater mean sEMG amplitudes during the first bout hip thrust versus squat set, but this did not consistently predict gluteal hypertrophy outcomes.

Conclusion: Nine weeks of squat versus hip thrust training elicited similar gluteal hypertrophy, greater thigh hypertrophy in SQ, strength increases that favored exercise allocation, and similar strength transfers to the deadlift and wall push.

Keywords: Hip thrust, back squat, gluteus maximus, strength

INTRODUCTION

Resistance training (RT) presents potent mechanical stimuli that produce robust biological responses (1). However, RT responses vary considerably depending on several training variables. One such variable is exercise selection—different exercises have varying mechanical demands that can lead to differences in muscle growth, strength, and other related outcomes (2-5). Practitioners and researchers often rely on functional anatomy, basic biomechanics, and acute physiological measurements to surmise what adaptations different exercises may elicit. The degree to which such surmises can meaningfully predict outcomes remains an open question, and recent work casts some doubt on their fidelity.

The reliance on theory and acute measures to guide exercise selection is especially evident in the hip extension exercise literature, an area of particular interest with applications in rehabilitation, performance, injury prevention, and bodybuilding. The roles of various hip extensor muscles during different hip extension tasks have been studied in several ways, including surface electromyography (sEMG), nerve blocks, and musculoskeletal modeling (6-8). Based on these acute measures, investigators infer stimulus potency or exercise superiority. For instance, previous work investigated sEMG amplitudes during two common and contentiously contrasted hip extension exercises—the hip thrust and squat—to compare muscle function, implying that this relates to subsequent adaptations (9-11). Although mean and peak sEMG amplitudes favored hip thrusts, sEMG's ability to predict longitudinal strength and hypertrophy outcomes from resistance training interventions was recently challenged (12). To help overcome some of sEMG's limitations, more sophisticated investigations integrate excitation into musculoskeletal models(8). Yet, more comprehensive analyses of muscle contributions are still limited by their underlying assumptions (13), and even perfect modeling of muscle contributions presumes a one-to-one relationship between tension and adaptations.

Muscle tension is the primary driver of muscle hypertrophy but is unlikely to be its sole determinant. Recent evidence demonstrates that RT at long muscle lengths and long-duration static stretching can augment hypertrophic outcomes (14, 15), suggesting other factors may modulate anabolic signaling. It is unknown to what extent muscle tension may interact with position-specific anabolic signaling and other variables to contribute to the anabolic response and how this interaction may change under different conditions. Regarding the squat and hip thrust, the former has a steeper hip extension resistance curve with a relatively greater emphasis in hip flexion(7, 16), which may confer a more potent gluteal training stimulus. However, this notion assumes proportional force sharing among the hip extensors, but contributions shift throughout the range of motion, clouding inferences (17). This highlights that longitudinal predictions necessitate assumptions about how motor systems satisfy the mechanical demands imposed by each exercise and subsequent biological responses, making it difficult to infer the potency of the hypertrophic stimulus using indirect measures. We ultimately need longitudinal data to understand and accurately forecast longitudinal outcomes from individual movements.[1]

Direct evidence is presently needed to compare the outcomes of various exercises. Therefore, the purpose of this study was to examine how RT using either the barbell squat or barbell hip thrust on a set-volume equated basis affected gluteus maximus, medius, and minimus muscle hypertrophy (determined by MRI) and various strength outcomes including the back squat, hip thrust, deadlift, and isometric wall push. As a secondary outcome, we sought to determine how

---

[1] We acknowledge but will not further discuss the squat versus hip thrust paper by Barbalho et al. (18). These and other data from this laboratory were scrutinized for being improbable, resulting in several retractions (19).

102 these exercises affected gluteus maximus/medius muscle excitation patterns using sEMG and if
103 sEMG amplitudes forecasted hypertrophy.
104
METHODS
Ethical considerations and participant recruitment
Before commencing study procedures with human participants, this study was approved by the
Auburn University Institutional Review Board (protocol #: 22-588). All approved study
procedures followed the latest revisions to the Declaration of Helsinki (2013) except for being pre-
registered as a clinical trial on an online repository. Inclusion criteria were as follows: (a) between
the ages of 18-30 years old with a body mass index (body mass/height^2) of less than 30 kg/m^2; (b)
have minimal experience with resistance training, averaging less than or equivalent to one day per
week for the last five years; (c) have not been actively participating in any structured endurance
training program (e.g. running or cycling) for more than two days per week over the past six
months; (d) free of any known overt cardiovascular or metabolic disease; (e) have not consumed
supplemental creatine, and/or agents that affect hormones (testosterone boosters, growth hormone
boosters, etc.) within the past two months, (f) free of any medical condition that would
contraindicate participation in an exercise program, (g) do not have conditions which preclude
performing an MRI scan (e.g., medically-implanted devices), (h) and free of allergies to lactose or
intolerances to milk derived products that would contraindicate ingestion of whey protein. Eligible
participants who provided verbal and written consent partook in the testing and training procedures
outlined in the following paragraphs.
123
Study design overview
An overview of the study design can be found in Figure 1. Participants performed two pre-
intervention testing visits, one in a fasted state for body composition and MRI assessments and the
other in a non-fasted state for strength assessments. These visits occurred in this sequence ~48
hours apart; after the pre-intervention strength visit, participants were randomly assigned to one of
two experimental groups, including the barbell back squat (SQ) or barbell hip thrust (HT) groups.
Two days following the pre-intervention strength testing, all participants partook in their first
workout, which served to record right gluteal muscle excitation via sEMG during one set of 10
repetitions for both the SQ and HT exercises. Thereafter, participants engaged in 9 weeks of
resistance training (two days per week). Seventy-two hours following the last training bout,
participants performed two post-intervention testing visits with identical timing and protocols as
pre-testing.
136

Figure 1. Study design overview
Legend: Figure depicts study design overview described in-text. Abbreviations: PRE, pre-intervention testing visit; POST, post-intervention testing visit; HT, barbell hip thrust; SQ, barbell squat; body comp., body composition testing using bioelectrical impedance spectroscopy; MRI, magnetic resonance imaging; sEMG, electromyography.

Body composition and MRI assessments

Body composition. Participants were told to refrain from eating for 8 h prior to testing, eliminate alcohol consumption for 24 h, abstain from strenuous exercise for 24 h, and to be well hydrated for testing. Upon arrival participants submitted a urine sample (~50 mL) for urine specific gravity assessment (USG). Measurements were performed using a handheld refractometer (ATAGO; Bellevue, WA, USA), and USG levels in all participants were ≤ 1.020, indicating sufficient hydration. Participants' heights were measured using a stadiometer and body mass was assessed using a calibrated scale (Seca 769; Hanover, MD, USA) with body mass being collected to the nearest 0.1 kg and height to the nearest 0.5 cm. Body composition was then measured by bioelectrical impedance spectroscopy (BIS) using a 4-lead (two hands, two feet) SOZO device (ImpediMed Limited, Queensland, Australia) according to the methods described by Moon et al. (20). Our laboratory has previously shown these methods to produce test-retest intraclass correlation coefficients ($ICC_{3,1}$) >0.990 for whole body intracellular and extracellular water metrics on 24 participants (21), and this device provided estimates of fat free mass, skeletal muscle mass, and fat mass.

MRI Measurements. MRI testing assessed the muscle cross-sectional area (mCSA) of both glutei maximi. Upon arriving to the Auburn University MRI Research Center, participants were placed onto the patient table of the MRI scanner (3T SkyraFit system; Siemens, Erlangen, Germany) in a prone position with a ~5-minute latency period before scanning was implemented. A T1-weighted turbo spin echo pulse sequence (1400 ms repetition time, 23 ms echo time, in-plane resolution of 0.9×0.9 mm^2) was used to obtain transverse image sets. 71 slices were obtained with a slice thickness of 4 mm with no gap between slices. Measurements were taken by the same investigator (R.J.B.) for all scans who did not possess knowledge of the training conditions for each participant.

Following the conclusion of the study, MRI DICOM files were preprocessed using Osirix MD software (Pixmeo, Geneva, Switzerland), and these images were imported into ImageJ (National Institutes of Health; Bethesda, MD, USA) whereby the polygon function was used to manually trace the borders of muscles of interest to obtain mCSA. For all participants, image standardization was as follows: (a) the middle of the gluteus maximus was standardized at the image revealing the top of the femur, (b) the image that was 10 slices upward from this mark was considered to be the upper gluteus maximus, (c) the image that was 18 slices downward from the top of the femur was considered lower gluteus maximus, (d) gluteus medius and minimus mCSAs were ascertained at the upper gluteus maximus image, and (e) combined quadriceps (vastii and rectus femoris), adductors (brevis, longus, and magnus), and combined hamstrings (biceps femoris, semitendinosus, semimembranosus) mCSAs were ascertained at the first transverse slice distal to the last portion of the lower gluteus maximus. When drawing borders to quantify muscles of interest, care was taken to avoid fat and connective tissue. Certain muscles were grouped (i.e., gluteus medius + minimus, combined quadriceps muscles, combined adductor muscles, combined hamstrings muscles) due to inconsistent and poorly delineated muscle borders within participants. All left- and right-side gluteus muscles were summed to provide bilateral mCSA values at each site. Alternatively, thigh musculature mCSA values were yielded from the averages of the left and right legs. This method was performed on the thigh because ~10% of participants yielded either

185   left or right thigh images that presented visual artifacts from the edge of the MRI receiving coil.
186   In these situations, thigh musculature from only one of the two legs was quantified.
187
Strength assessments
Isometric muscle strength (wall push). Participants reported to the laboratory (non-fasted)
having refrained from any exercise other than activities of daily living for at least 48 h before
baseline testing. A tri-axial force plate (Bertec FP4060-10-2000; Columbus, OH, USA) with an
accompanying amplifier (Bertec model # AM6800) sampling at 1000 Hz was used to measure
horizontal force production in newtons (N) during a wall push test. The distance from the force
plate to the wall was positioned such that when the subjects' forearms parallel with the ground,
the torso was at a ~45º angle with the ground, and one rear foot was in contact with the force plate.
Hand placement was standardized by distance from the ground and foot placement was
standardized by distance from the wall. The subject was instructed to push, using the dominant
leg, as hard as possible into the wall while keeping the torso at 45º (Figure 2). Two wall pushes
were performed for three seconds each, with each repetition being separated by two minutes of
rest. The highest peak horizontal force from these two tests was used for analysis.
201

202
203   **Figure 2. Wall push demonstration**
204   Legend: Figure depicts the wall push test with one of the co-authors (M.D.R.) and shows force tracing.

Dynamic muscle strength. Following wall push testing, dynamic lower body strength was assessed by three-repetition maximum (3RM) testing for the barbell back squat, barbell hip thrust, and barbell deadlift exercises. Notably, our laboratory has extensively performed 3RM dynamic strength testing on numerous occasions in untrained and trained participants (22-25). Briefly, specific warm-up sets for each exercise consisted of coaching participants through the movement patterns and gauging comfort and movement proficiency. Subsequent warm-ups for each exercise were chosen with an attempt at approximating 5 repetitions at ~50% 1RM for one set and 2–3 repetitions at ~60–80% 1RM for two additional sets. Participants then performed sets of three repetitions with incremental increases in load for 3RM determinations for each exercise and three minutes of rest was given between each successive attempt. For all exercises, participants were instructed to perform repetitions in a controlled fashion, with a concentric action of approximately 1 s and an eccentric action of approximately 2 s. All three exercises were performed with feet spaced 1–1.5-times shoulder width apart. For the barbell squat, depth was set to when the femur was parallel to the floor, with all but one participant achieving a depth near, at, or below this point. Silhouettes of individual squat form are provided in supplementary material for ultimate transparency. For the barbell hip thrust, the hip thrust apparatus (Thruster 3.0, BC Strength; San Diego, CA, USA) was set to a height at which participants could make brief contact with the ground with the weight plate (21") and hips at the bottom of each repetition. Repetitions were considered properly executed when the participant's tibia was perpendicular to the floor and the femur was parallel to the floor. Torso position was sufficiently maintained to avoid excessive motion through the pelvis. For the barbell deadlift, participants began repetitions from the floor and were prompted to maintain the torso position throughout the execution of the lift. A lift was deemed successful once participants stood upright with full knee and hip extensions.

sEMG measurements during the first training bout

Subjects were asked to wear loose athletic attire to access the EMG electrode placement sites. Before placing the electrodes on the skin, if necessary, excess hair was removed with a razor, and the skin was cleaned and abraded using an alcohol swab. After preparation, double-sided adhesives were attached to wireless sEMG electrodes (Trigno system; Delsys, Natick, MA, USA), where were placed in parallel to the fibers of the right upper gluteus maximus, mid gluteus maximus, lower gluteus maximus, and gluteus medius (see Fig. 4a in *Results*). Upper and middle gluteus maximus electrodes were placed based on the recommendations of Fujisawa and colleagues (26), albeit we considered the lower gluteus maximus as middle. The upper gluteus maximus electrodes were placed superior and lateral to the shortest distance between the posterior superior iliac spine (PSIS) and the posterior greater trochanter, and the middle gluteus maximus electrodes were placed inferior and medial to the shortest distance between the PSIS and the posterior greater trochanter. Lower gluteus maximus electrodes were placed one inch (2.54 cm) above the most medial presentation of the gluteal fold. If it was ambiguous as to whether an appreciable amount of muscle tissue existed in this lower region, the participant was asked to contract the area and palpation was used to confirm proper placement. Gluteus medius electrodes were placed over the proximal third of the distance between the iliac crest and the greater trochanter. After the electrodes were secured, a quality check was performed to ensure sEMG signal validity. Following electrode placement, maximum voluntary isometric contraction (MVIC) testing was performed immediately prior to 10RM testing. For the gluteus maximus, the MVIC reference was a prone bent-leg hip extension against manual resistance applied to the distal thigh, as used by Boren and colleagues

(6). For the gluteus medius MVIC, participants laid on their side with a straight leg and abducted against manual resistance. Care was taken not to depress the joint of interest during manual testing. In all MVIC positions, participants were instructed to contract the tested muscle as hard as possible. After five minutes of rest following MVIC testing, all participants performed one set of ten repetitions utilizing estimated 10RM loads for both the barbell back squat and the barbell hip thrust exercises. The exercise form and tempo used were the same as described in the strength testing section above. During both sets, muscle excitation of the upper/middle/lower gluteus maximus and gluteus medius were recorded with the wireless sEMG system whereby electrodes were sampled at 1000 Hz. Participants allocated to HT training performed the squat set first followed by the hip thrust set. Participants allocated to SQ training performed the hip thrust set first followed by the back squat set. Following these two sEMG sets, the wireless sEMG electrodes were removed. Participants finished the session with two more sets of 8–12 repetitions using the calculated 10RM load for the exercise allocated to them for the intervention.

Signal processing was performed using software associated with the sEMG system (Delsys EMGworks Analysis v4.7.3.0; Delsys). sEMG signals from the MVICs and 10RM sets of back squat and hip thrust were first rectified. Signals were then processed with a second-order digital low-pass Butterworth filter, with a cutoff frequency of 10 Hz, and further smoothed using a root mean square moving window of 250 ms. The average of the middle 3 seconds of the filtered MVIC time series was then used to normalize the squat and hip thrust data for each site. Data were then visually inspected for fidelity before calculating the mean and peak sEMG values. Partial sequences of sEMG data were removed in the rare event that tempo was irregular or not maintained, or if a brief artifact was introduced. Final EMG data are presented as mean and peak sEMG amplitudes during the hip thrust and back squat 10RM sets. sEMG issues were only evident for a small portion (see *Results*) of the 34 participants who finished the intervention. Data were dropped from analyses due to artifacts produced through either electrode slippage or sEMG electrode jarring during the 10RM sets, leading to persistent clipping. In this regard, sample sizes for each muscle site are presented in the results section.

Resistance training procedures

The RT protocol consisted of 3–6 sets per session of barbell hip thrusts for HT participants or barbell back squats for SQ participants. Excluding the first week, which consisted of one session, all remaining weeks consisted of two sessions per week on non-consecutive days for 9 weeks. Week-to-week set schemes per session were as follows: week 1, 3 sets; week 2, 4 sets; weeks 3–6, 5 sets; weeks 6–9, 6 sets. The repetition range was set to 8–12 repetitions; if a participant performed less than 8 repetitions or more than 12 repetitions, the load was adjusted accordingly. D.L.P. and 1–2 other co-authors supervised all sessions, during which participants were verbally encouraged to perform all sets to the point of volitional muscular failure, herein defined as the participants being unable to volitionally perform another concentric repetition while maintaining proper form. Again, the exercise form and tempo used were the same as described in the strength testing section above; however, squat repetitions were not limited to a depth corresponding to the femur parallel to the floor but rather the lowest depth achievable. Outside of these supervised training sessions, participants were instructed to refrain from performing any other lower-body RT for the duration of the study. Participants could miss a maximum of 2 sessions and still be included in the analysis.

Dietary instructions during the study

297 Participants were given containers of a whey protein supplement (Built with Science; Richmond,
298 BC, Canada) and were instructed to consume one serving per day (per serving: 29 g protein, 1 g
299 carbohydrate, 0.5 g fat, 130 kcal). This was done in the hope of diminishing inadequate protein
300 intake as a confounding variable. Other than this guidance, participants were advised to maintain
301 their customary nutritional regimens to avoid other potential dietary confounders.
302
Notes on randomization and blinding
Investigators were blinded to group allocation during the MRI scan and its analysis. Participants
were not blinded to group allocation as exercise comparisons were not amenable to blinding. Due
to logistical constraints investigators were not blinded to group allocation during strength testing
and, thus, bias cannot be completely ruled out in this context. Randomization into SQ and HT
groups was performed via a random number generator in blocks of 2 or 4 as participants consented.
309
Statistics and figure construction
Data were analyzed in Jamovi v2.3 (https://www.jamovi.org) and R (version 4.3.0). We performed
three different sets of analyses. First, we compared mean and peak HT and SQ sEMG amplitudes
from the first training session, for which we performed paired *t*-tests.
Second, we compared the longitudinal effects of HT and SQ training on mCSA and
strength. Notably, baseline and within-group inferential statistics were not calculated, as baseline
significance testing is inconsequential (27) and within-group outcomes are not the subject of our
research question (28). However, we descriptively present within-group changes to help
contextualize our findings. The effect of group (SQ versus HT) on each outcome variable was
estimated using linear regression, in which post-intervention scores were the response variable,
group was dummy-coded 0 for SQ and 1 for HT, and the pre-intervention score was included as a
covariate of no interest (29). The model output can thus be interpreted as the expected difference
in post-intervention (or mathematically equivalently, change) scores between the SQ and HT
groups for a given pre-intervention score. We used the bias-corrected and accelerated stratified
bootstrap with 10,000 replicates to calculate 95% compatibility intervals (CIs).
Third, we investigated the extent to which sEMG amplitudes from the first session
forecasted growth. There are multiple ways this question could be posed, and since claims
surrounding sEMG amplitude's predictive power are ambiguous, we addressed each of the
following questions: i) Do individuals with greater sEMG amplitudes grow more than individuals
with lower sEMG amplitudes? For this, we calculated a Pearson correlation for each muscle using
changes in mCSA and the sEMG amplitudes. ii) Do regions or muscles with greater sEMG
amplitudes grow more than regions or muscles with lower sEMG amplitudes? For this, we used a
linear mixed-effects model in which $\ln(mCSA_{post}/mCSA_{pre})$ was the response variable; sEMG
amplitude, group, and their interaction were fixed effects; and we permitted intercepts and slopes
for sEMG amplitude to vary across subjects. Since we are interested in generalizable predictions,
we calculated prediction intervals for the slopes by calculating a Wald interval using the sum of
the parameter variance and random effects variance. iii) Can the differences in growth elicited
from different exercises be accounted for by sEMG amplitude? For this, we calculated the so-
called "indirect effect" of sEMG amplitude, which represents the extent to which the group effect
on hypertrophy can be explained by sEMG amplitudes. This was done the same way a typical
"mediation analysis" is done (although, this should not be viewed as causal here)—we
bootstrapped the difference between the group effect (SQ vs. HT) when sEMG was not in the
model and when sEMG was added to the model. If group-based sEMG differences accounted for

group-based hypertrophy differences, then the effect of group on growth would shrink towards 0
and sEMG would absorb the variance in growth.
Figures were constructed using Microsoft PowerPoint and through paid site licenses for
BioRender (https://www.biorender.com), GraphPad Prism v9.2.0 (San Diego, CA, USA), and
ggplot2.
348
RESULTS
CONSORT and general baseline participant characteristics
The CONSORT diagram is presented in Figure 3. In total, 18 HT and 16 SQ participants completed
the study and were included in data analyses unless there were technical issues precluding the
inclusion of data (e.g., sEMG clipping).
General baseline characteristics of the 18 HT participants who finished the intervention
were as follows: age: 22 ± 3 years old, 24 ± 3 kg/m^2, 5 M and 13 F. Baseline characteristics of the
16 SQ participants who finished the intervention were as follows: age: 24 ± 4 years old, 23 ± 3
kg/m^2, 6 M and 10 F. Also notable, the HT participants missed an average of 0.8 ± 0.4 workouts
during the study, and the SQ participants missed 0.8 ± 0.5.
359

360
361 **Figure 3. CONSORT diagram**
362 Figure depicts participant numbers through various stages of the intervention. All participants were included in data
363 analysis unless there were technical issues precluding the inclusion of data (e.g., EMG clipping).
364
365

First bout sEMG results

sEMG data obtained from the right gluteus muscles during the first workout bout, based on one set of 10RM hip thrust and one set of 10RM sqaut, are presented in Figure 4. All sites showed greater mean sEMG values during the hip thrust versus squat set ($p < 0.01$ for all; Fig. 4b). Peak sEMG values were greater for the upper and middle gluteus maximus ($p < 0.001$ and $p = 0.015$, respectively), whereas small differences existed for the lower gluteus maximus or gluteus medius sites (Fig. 4b). The number of repetitions completed during the 10RM sets used for sEMG recordings were not different between exercises (back squat: 9±1 repetitions, hip thrust: 9±2 repetitions).

Figure 4. Surface electromyogram (sEMG) amplitudes during the back squat and barbell hip thrust
Legend: During the first session, all participants performed both back squats and barbell hip thrusts while we recorded sEMG amplitudes. (**a**) Representative sEMG electrode placement is depicted on a co-author in panel. (**b**) Data depict mean (left) and peak (right) sEMG amplitudes during one 10RM set of hip thrusts and one 10RM set of back squats. As 34 participants partook in this test, sample sizes vary due to incomplete data from electrode slippage or clipping. Bars are mean ± SD, and individual participant values are depicted as dots. (**c**) Representative data from one participant.

Gluteus musculature mCSAs according to MRI

The effect of SQ relative to HT for left+right mCSA was negligible across gluteal muscles (Figure 6). Point estimates modestly favored HT for lower [effect ± SE, -1.6 ± 2.1 cm^2; CI$_{95\%}$ (-6.1, 2.0)], mid [-0.5 ± 1.7 cm^2; CI$_{95\%}$ (-4.0, 2.6)], and upper [-0.5 ± 2.6 cm^2; CI$_{95\%}$ (-5.8, 4.1)] gluteal mCSAs; these point estimates were dwarfed by the variance. Left+right mCSA values for the gluteus medius + minimus demonstrated a lesser magnitude of growth (see *Table 1*), with a point estimate that also modestly favored HT albeit with appreciable variance [-1.8 ± 1.5 cm^2; CI$_{95\%}$ (-4.6, 1.4)].

Figure 6. Gluteus musculature mCSA changes following back squat and barbell hip thrust training, assessed using MRI

Legend: Figure depicts pre-to-post intervention MRI-derived muscle cross-sectional area (mCSA) summed values for (**a**) left + right (L+R) upper gluteus maximus, (**b**) L+R middle gluteus maximus, (**c**) L+R lower gluteus maximus, (**d**) L+R gluteus medius+minimus. Data include 18 participants in the hip thrust group and 16 participants in the back squat group. Graphs contain change scores with individual participant values depicted as dots. (**e**) Three pre and post representative MRI images are presented from the same participant with white polygon tracings of the L+R upper gluteus maximus and gluteus medius+minimus (top), L+R middle gluteus maximus (middle), and L+R lower gluteus maximus (bottom).

Thigh musculature mCSAs according to MRI

Compared to HT, SQ produced greater mCSA growth for quadriceps [3.6 ± 1.5 cm^2; CI$_{95\%}$ (0.7, 6.4)] and adductors [2.5 ± 0.7 cm^2; CI$_{95\%}$ (1.2, 3.9)] (Figure 7). However, hamstrings growth was fairly equivocal across both conditions, yielding negligible between-group effects [0.1 ± 0.6 cm^2; CI$_{95\%}$ (−0.9, 1.4)] (Figure 7).

Figure 7. Thigh musculature mCSA changes following back squat and barbell hip thrust training, assessed using MRI

Legend: Figure depicts MRI-derived muscle cross-sectional area (mCSA) average change scores for left and/or right (**a**) quadriceps, (**b**) adductors, and (**c**) hamstrings. Data include 18 participants in the hip thrust group and 16 participants in the back squat group. Bar graphs contain change scores with individual participant values depicted as dots. (**d**) A representative pre- and post-intervention MRI image is presented with white polygon tracings of the quadriceps (denoted as Q), adductors (denoted as ADD), and hamstrings (denoted as H).

Strength outcomes

Strength outcomes of SQ relative to HT favored respective group allocation for specific lift 3RM values. Specifically, Squat 3RM favored SQ [14 ± 2 kg; $CI_{95\%}$ (9, 18)], and hip thrust 3RM favored HT [-26 ± 5 kg; $CI_{95\%}$ (-34, -16)] (Figure 8). Results were more equivocal for the deadlift 3RM [0 ± 2 kg; $CI_{95\%}$ (-4, 3)] and wall push [-7 ± 12 N; $CI_{95\%}$ (-32, 17)] (Figure 8).

Figure 8. Strength outcomes following back squat and barbell hip thrust training
Legend: Figure depicts change scores for (**a**) 3RM barbell back squat values, (**b**) 3RM barbell hip thrust values, (**c**) 3RM barbell deadlift values, and (**d**) horizontal ground reactive forces (GRF) during the wall push as demonstrated in Figure 2. Data include 18 participants in the hip thrust group and 16 participants in the back squat group.

Forecasting training-induced gluteus muscle mCSA changes with sEMG amplitudes

across-subject correlations. sEMG amplitude's ability to forecast muscle growth across-subjects was generally poor and variable. Mean sEMG amplitudes produce negligible to moderate correlations for lower [$r = 0.18$ (-0.30, 0.57)], middle, [$r = -0.03$ (-0.32, 0.25)], upper [$r = 0.50$ (0.03, 0.81)], and medius+minimus [$r = 0.28$ (0, 0.53)]. We observed similar results for peak sEMG amplitudes from the lower [$r = 0.13$ (-0.16, 0.46)], middle [$r = -0.03$ (-0.33, 0.21)], upper [$r = 0.32$ (-0.05, 0.62)], and medius+minimus [$r = 0.24$ (-0.02, 0.48)].

Across-region correlations. We fit two linear mixed-effects models to assess how differences in sEMG amplitudes across muscles can account for regional growth. Since the response variable was relative muscle size on the log scale, the exponentiated coefficients can be interpreted as the increase in muscle relative to baseline for each additional %MVIC; notably, this effect is multiplicative rather than additive. The first model, which used mean sEMG amplitudes, produced small and variable estimates for both SQ [1.003, $PI_{95\%}$ (0.998, 1.008)] and HT [1.002, $PI_{95\%}$ (0.997, 1.006)] groups. The second model, which used peak sEMG amplitudes, produced even more

451 modest results for both the SQ [1.0003, PI$_{95\%}$ (0.9997, 1.0009)] and HT [1.0002, PI$_{95\%}$ (0.9996,
452 1.0007)] groups.
453
Across-exercise variance. Mean sEMG amplitude's ability to capture the group effects was
inconsistent for lower [indirect effect = −0.55, CI$_{95\%}$ (−3.87, 0.58)], middle [0.06, CI$_{95\%}$ (−0.82,
1.56)], upper [−2.98, CI$_{95\%}$ (−8.73, −0.38)], and medius+minimus [−0.73, CI$_{95\%}$ (−2.70, 0.14)].
We observed similar results for peak sEMG amplitudes for lower [−0.08, CI$_{95\%}$ (−2.27, 0.59)],
middle [0.22, CI$_{95\%}$ (−1.63, 1.89)], upper [−3.04, CI$_{95\%}$ (−8.32, 0.15)], and medius+minimus
[−0.86, CI$_{95\%}$ (−2.47, 0)]. These estimates can be compared to the group effects ("total effects")
earlier in the Results.
461
DISCUSSION
To further our understanding of hip extensor exercises and the validity of relying on theory and
acute physiological measures for exercise selection, here we acutely (sEMG) and longitudinally
(hypertrophy, strength) compared two common hip extension exercises: the back squat and barbell
hip thrust. Acutely, HT sEMG amplitudes were generally greater for the HT. However, this did
not appear to translate and accurately capture longitudinal adaptations. Across all gluteus muscle
hypertrophy outcomes, SQ and HT training yielded modest differences but meaningful growth
occurring, except in the gluteus medius and minimus. Thigh hypertrophy outcomes favored SQ in
the adductors and quadriceps, with no meaningful growth in either group in the hamstrings.
Strength outcomes indicated that hip thrust 3RM changes favored HT, back squat 3RM changes
favored SQ, and other strength measures similarly increased in both groups. sEMG amplitudes
could not reliably predict hypertrophic outcomes across several analytical approaches. In the
following paragraphs, we discuss these results in the context of available evidence and speculate
on their potential implications for exercise prescription. Moreover, a summary of findings is
provided here in tabular form for convenience to the reviewer (Table 1).
477
478 **Table 1. Descriptive scores for each training variable**

Variable	SQ PRE	SQ POST	SQ Δ	HT PRE	HT POST	HT Δ
SMM (kg)	21.6 (5.0)	22.2 (5.3)	0.7 (0.8)	21.9 (4.8)	22.4 (5.0)	0.5 (0.9)
FM (kg)	20.3 (5.0)	19.5 (4.2)	-0.7 (1.7)	19.7 (6.2)	19.4 (6.0)	-0.4 (1.5)
Squat 3RM (kg)	49.8 (17.6)	71.9 (22.2)	22.1 (8.4)	53.2 (15.7)	61.9 (15.4)	8.68 (5.2)
Hip Thrust 3RM (kg)	79.8 (24.0)	106.7 (31.9)	26.9 (11.7)	81.8 (25.3)	134.4 (27.7)	52.7 (15.4)
Deadlift 3RM (kg)	61.5 (17.5)	70.7 (21.1)	9.2 (5.7)	59.0 (17.0)	68.2 (15.6)	9.2 (5.5)
Wall push (N)	299.3 (97.2)	322.1 (101.1)	22.8 (39.1)	298.1 (80.9)	327.9 (84.3)	29.8 (36.7)
Gmax Upper CSA (cm²)	52.0 (17.9)	58.5 (16.7)	6.5 (4.9)	50.9 (13.9)	58.0 (15.7)	7.1 (9.8)
Gmax Middle CSA (cm²)	92.2 (22.9)	101.3 (23.1)	9.16 (4.4)	88.71 (16.6)	98.31 (19.2)	9.6 (5.7)
Gmax Lower CSA (cm²)	72.4 (21.0)	86.2 (23.9)	13.8 (4.8)	71.0 (17.2)	86.3 (18.3)	15.3 (7.6)
MED+MIN CSA (cm²)	79.1 (16.4)	79.6 (14.9)	0.5 (4.6)	76.4 (14.1)	79.0 (14.1)	2.6 (4.8)
QUAD CSA (cm²)	61.8 (16.4)	69.8 (17.7)	7.9 (4.8)	63.8 (12.5)	68.1 (12.8)	4.3 (3.4)
ADD CSA (cm²)	41.4 (9.4)	45.6 (9.5)	4.2 (1.7)	40.6 (8.9)	42.2 (9.5)	1.7 (2.3)

479 Abbreviations: SMM, skeletal muscle mass; RM, repetition maximum; GRF, ground reaction force; mCSA, muscle
480 cross-sectional area; Gmax, Gluteus Maximus; MED+MIN, Gluteus medius and minimus; QUAD, quadriceps; ADD,
481 adductors; HAM, hamstring. Symbol: Δ, pre-to-post intervention change score. Note: all data are presented as mean
482 (standard deviation).
483
Hypertrophy Outcomes
The primary finding of interest was that upper, middle, and lower gluteus maximus muscle
hypertrophy was similar after nine weeks of training with either the squat or hip thrust. This may
seem to run counter to recent evidence suggesting muscle tension in lengthened positions augments

growth (14) since the sticking region for the squat occurs in greater hip flexion as compared to the
hip thrust. Importantly, much of the previous work on this topic is in muscles being worked in a
more isolated fashion (2, 4, 30). Thus, the equivocal findings may suggest that the context in which
the muscle is experiencing lengthened loading critically determines subsequent adaptations.
Muscle contributions, and not just positions, may need to be jointly considered in determining
whether superior hypertrophy outcomes would be achieved. This idea is loosely supported by
sEMG and musculoskeletal modeling research, suggesting the gluteus maximus may not be
strongly recruited toward the bottom of the squat (9, 17). This notion would suggest the nervous
system does not strongly recruit the gluteus maximus while at its longest length in the squat,
precluding one from maximizing the benefits of stretch-augmented hypertrophy.
In addition to motor control governing how the gluteus maximus contributes to and adapts
from the squat, there are study-specific considerations. Both exercises may stimulate similar
muscle hypertrophy in untrained populations given that RT in general elicits rapid growth early
on, creating a ceiling effect on growth rate and thus observed growth. Alternatively stated, skeletal
muscle hypertrophy in novice trainees may be less influenced by nuances in exercise selection.
Notwithstanding, our results suggest that a nine-week set-equated training program with either the
hip thrust, or squat elicits similar gluteal muscle hypertrophy in novice trainees.
Finally, our data show that thigh hypertrophy favored the squat, whereas thigh hypertrophy
was minimal in the hip thrust. This is perhaps unsurprising and is consistent with previous
literature. The adductors, particularly the adductor magnus, have the largest extension moment
contribution at the bottom of a squat (17). Thus, the nervous system may favor its recruitment for
this purpose. In line with this finding, adductor magnus mCSA changes favor a greater squat depth
(31). Hamstring mCSA changes did not occur in either group, in accordance with previous work
(31). Critically, these data imply that the hip thrust exercise primarily targets gluteus muscle
hypertrophy while limiting non-gluteal thigh muscle hypertrophy; in other words, the hip thrust
appears to be more gluteus maximus-specific.
514
Strength Outcomes
Both groups effectively increased strength outcomes for all exercises tested. However, HT RT
better increased hip thrust strength and SQ RT better increased back squat strength, which is to be
expected due to training specificity (32). Baseline adjusted increases in back squat 3RM increased
by 17% in the HT group and 43% in the SQ group, while hip thrust strength increased by 65% in
HT group and 33% in SQ group. In contrast, deadlift and wall push outcomes increased similarly
in both groups. Deadlift increased by 15% in both SQ and HT, and wall push increased by 8% in
SQ and 10% in HT.
523
Using acute first bout sEMG to Predict Hypertrophy
Our secondary aim was to evaluate the ability of sEMG to forecast longitudinal adaptations. In
agreement with previous work (9), gluteus muscle sEMG amplitudes during the hip thrust exercise
were greater across all measured gluteal sites. However, these sEMG amplitude differences did
not reliably translate to greater hypertrophy, no matter what analytical approach we took.
Specifically, i) individuals with greater sEMG amplitudes did not consistently experience greater
growth; ii) regions with greater sEMG amplitudes did not consistently experience greater growth;
iii) differences in sEMG amplitudes between exercises could not consistently explain differences
in growth, in large part since the hypertrophy results were equivocal. This finding implies that
acute sEMG readings during a workout bout are not predictive of hypertrophic outcomes, and this

viewpoint is supported by a recent review by Vigotsky et al. (12). As indicated by the authors,
inconsistent relationships between EMG amplitudes and muscle growth have been previously
reported, which may be due to one or several reasons, ranging from biases in the sEMG recordings
to assumptions about how adaptations occur (12). Evidently, the reliance on acute sEMG
measurements may in fact be an over-reliance, but more work is needed in this realm.
Finally, we also verbally asked participants which exercise they "felt more" in the gluteal
muscles after testing both exercises. All participants indicated they felt the hip thrust more in the
gluteal region. However, these data were not quantified and, despite these anecdotal sensations
and sEMG differences indicating more gluteus muscle excitation during HT, hip thrust RT and
squat RT elicited similar applied outcomes. These findings highlight the importance of
longitudinal investigations.
545

Limitations
Our study has a few limitations to consider. First, our participants were young untrained men and
women; thus, results cannot necessarily be generalized to other populations including adolescents,
older individuals, or trained populations. Additionally, like most training studies, this study was
limited in duration. It should also be noted that gluteal hypertrophy was the main outcome, and the
MRI coil was placed over this region as subjects were lying prone. Thus, compression may have
affected the thigh musculature, and distal measures were not obtained for the thigh. Finally,
training volume was equated, and frequency was set at two training days per week. Therefore,
results can only be generalized to this protocol.
Although we did not consider female participants' menstrual cycle phase or contraceptive
usage, we do not view this as a limitation. In this regard, several reports indicate that contraceptives
have no meaningful impact on muscle hypertrophy in younger female participants during periods
of resistance training (33-38). Likewise, well-controlled trials indicate that the menstrual cycle
phase does not affect strength characteristics (39), and that variations in female hormones during
different phases do not affect muscle hypertrophy and strength gains during 12 weeks of resistance
training (40).
562

Future Directions
Future research should aim to examine a group that performs both exercises on a volume-equated
basis to determine if there are synergetic effects. Comparing these exercises with different
volumes/frequencies is also warranted as exercises may have differing volume tolerances. From a
mechanistic standpoint, future studies should characterize anabolic signaling between different
points on the length-tension curve as well as ascertain where a muscle exists on this curve with
more clarity.
570

Conclusions
Squat and hip thrust RT elicited similar gluteal hypertrophy, whereas quadriceps and adductors
hypertrophy was superior with squat training. Further, although strength increases were specific
to exercise allocation, both forms of RT elicited similar strength transfer to the deadlift and wall
push. Importantly, these results could not be reliably predicted from acute data (sEMG). These
current data provide trainees with valuable insight concerning two widely popular hip-specific
exercise modalities, and this information can be leveraged for exercise selection based on specific
structural or functional goals.

ACKNOWLEDGEMENTS
We thank the participants who volunteered and participated in the study. We also thank Bradley
Ruple, Josh Godwin, and C. Brooks Mobley for their assistance and insight throughout the project.
We also thank Jeremy Ethier for donating whey protein to the study. B.M.C. and M.H. disclose
that they sell exercise related products and services. However, neither was involved in any aspect
of the study beyond assisting with the study design and providing funds to partially cover
participant and MRI costs through a gift to the laboratory of M.D.R. All other co-authors have no
apparent conflicts of interest in relation to these data.
587
588   DATA AVAILABILITY
589   Raw data related to the current study outcomes are provided in the supplementary materials.
590
591
FUNDING INFORMATION
Funding for this study was made possible through gift funds (some of which were donated by the
International Scientific Research Foundation for Fitness and Nutrition and B.M.C.) to the M.D.R.
laboratory. Other financial sources included indirect cost sharing (generated from various
unrelated contracts) from the School of Kinesiology, M.C.M. being fully supported through a T32
NIH grant (T32GM141739), and D.L.P. being fully supported by a Presidential Graduate Research
Fellowship (fund cost-sharing from Auburn University's President's office, the College of
Education, and the School of Kinesiology).

REFERENCES

1. Egan B, Sharples AP. Molecular responses to acute exercise and their relevance for adaptations in skeletal muscle to exercise training. *Physiological Reviews*. 2023;103(3):2057-170.

2. Maeo S, Huang M, Wu Y et al. Greater Hamstrings Muscle Hypertrophy but Similar Damage Protection after Training at Long versus Short Muscle Lengths. *Medicine and Science in Sports and Exercise*. 2021;53(4):825-37.

3. Zabaleta-Korta A, Fernández-Peña E, Torres-Unda J, Garbisu-Hualde A, Santos-Concejero J. The role of exercise selection in regional Muscle Hypertrophy: A randomized controlled trial. *J Sports Sci*. 2021;39(20):2298-304.

4. Maeo S, Wu Y, Huang M et al. Triceps brachii hypertrophy is substantially greater after elbow extension training performed in the overhead versus neutral arm position. *European Journal of Sport Science*. 2022:1-11.

5. Waters R, Perry J, McDaniels J, House K. The Relative Strength of the Hamstrings during Hip Extension : JBJS. *The Journal of Bone & Joint Surgery*. 1974.

6. Boren K, Conrey C, Le Coguic J, Paprocki L, Voight M, Robinson TK. ELECTROMYOGRAPHIC ANALYSIS OF GLUTEUS MEDIUS AND GLUTEUS MAXIMUS DURING REHABILITATION EXERCISES. *Int J Sports Phys Ther*. 2011;6(3):206-23.

7. Brazil A, Needham L, Palmer JL, Bezodis IN. A comprehensive biomechanical analysis of the barbell hip thrust. *PLoS ONE*. 2021;16(3):e0249307.

8. Collings TJ, Bourne MN, Barrett RS et al. Gluteal Muscle Forces during Hip-Focused Injury Prevention and Rehabilitation Exercises. *Medicine and Science in Sports and Exercise*. 2023;55(4):650-60.

9. Contreras B, Vigotsky AD, Schoenfeld BJ, Beardsley C, Cronin J. A Comparison of Gluteus Maximus, Biceps Femoris, and Vastus Lateralis Electromyographic Activity in the Back Squat and Barbell Hip Thrust Exercises. *J Appl Biomech*. 2015;31(6):452-8.

10. Delgado J, Drinkwater EJ, Banyard HG, Haff GG, Nosaka K. Comparison Between Back Squat, Romanian Deadlift, and Barbell Hip Thrust for Leg and Hip Muscle Activities During Hip Extension. *J Strength Cond Res*. 2019;33(10):2595-601.

11. Williams MJ, Gibson NV, Sorbie GG, Ugbolue UC, Brouner J, Easton C. Activation of the Gluteus Maximus During Performance of the Back Squat, Split Squat, and Barbell Hip Thrust and the Relationship With Maximal Sprinting. *J Strength Cond Res*. 2021;35(1):16-24.

12. Vigotsky AD, Halperin I, Trajano GS, Vieira TM. Longing for a Longitudinal Proxy: Acutely Measured Surface EMG Amplitude is not a Validated Predictor of Muscle Hypertrophy. *Sports medicine (Auckland, N.Z.)*. 2022;52(2):193-9.

13. Herzog W, Leonard TR. Validation of optimization models that estimate the forces exerted by synergistic muscles. *Journal of Biomechanics*. 1991;24:31-9.

14. Wolf M, Androulakis-Korakakis P, Fisher J, Schoenfeld B, Steele J. Partial Vs Full Range of Motion Resistance Training: A Systematic Review and Meta-Analysis. *International Journal of Strength and Conditioning*. 2023;3(1).

15. Warneke K, Wirth K, Keiner M et al. Comparison of the effects of long-lasting static stretching and hypertrophy training on maximal strength, muscle thickness and flexibility in the plantar flexors. *Eur J Appl Physiol*. 2023.

645   16.   Lahti J, Hegyi A, Vigotsky A, Ahtiainen J. Effects of barbell back squat stance width on
646   sagittal and frontal hip and knee kinetics. *Scandinavian journal of medicine & science in
647   sports*. 2019;29(1).
648   17.   Vigotsky ABM. Relative Muscle Contributions to Net Joint Moments in the Barbell Back
649   Squat. In: 2016.
650   18.   Barbalho M, Coswig V, Souza D, Serrão JC, Hebling Campos M, Gentil P. Back Squat
651   vs. Hip Thrust Resistance-training Programs in Well-trained Women. *International
652   journal of sports medicine*. 2020;41(5):306-10.
653   19.   Vigotsky A, Nuckols GL, Fisher J et al. Improbable data patterns in the work of Barbalho
654   et al. In: SportRxiv; 2020.
655   20.   Moon JR, Tobkin SE, Roberts MD et al. Total body water estimations in healthy men and
656   women using bioimpedance spectroscopy: a deuterium oxide comparison. *Nutr Metab
657   (Lond)*. 2008;5:7.
658   21.   Haun CT, Vann CG, Mobley CB et al. Effects of Graded Whey Supplementation During
659   Extreme-Volume Resistance Training. *Front Nutr*. 2018;5:84.
660   22.   Smith MA, Sexton CL, Smith KA et al. Molecular predictors of resistance training
661   outcomes in young untrained female adults. *Journal of applied physiology (Bethesda,
662   Md. : 1985)*. 2023;134(3).
663   23.   Godwin JS, Sexton CL, Kontos NJ et al. Extracellular matrix content and remodeling
664   markers do not differ in college-aged men classified as higher and lower responders to
665   resistance training. *Journal of applied physiology (Bethesda, Md. : 1985)*. 2023;134(3).
666   24.   Mobley CB, Haun CT, Roberson PA et al. Biomarkers associated with low, moderate,
667   and high vastus lateralis muscle hypertrophy following 12 weeks of resistance training.
668   *PLoS ONE*. 2018;13(4).
669   25.   Vann CG, Osburn SC, Mumford PW et al. Skeletal Muscle Protein Composition
670   Adaptations to 10 Weeks of High-Load Resistance Training in Previously-Trained Males.
671   *Frontiers in Physiology*. 2020;11.
672   26.   Fujisawa H, Suzuki H, Yamaguchi E, Yoshiki H, Wada Y, Watanabe A. Hip Muscle
673   Activity during Isometric Contraction of Hip Abduction. *J Phys Ther Sci*.
674   2014;26(2):187-90.
675   27.   Senn S. Testing for baseline balance in clinical trials. *Statistics in Medicine*.
676   1994;13(17):1715-26.
677   28.   Bland JM, Altman DG. Comparisons against baseline within randomised groups are often
678   used and can be highly misleading. *Trials*. 2011;12(1):264.
679   29.   Vickers AJ, Altman DG. Statistics notes: Analysing controlled trials with baseline and
680   follow up measurements. *BMJ*. 2001;323(7321):1123-4.
681   30.   Kassiano W, Costa B, Kunevaliki G et al. Greater Gastrocnemius Muscle Hypertrophy
682   After Partial Range of Motion Training Performed at Long Muscle Lengths. *Journal of
683   strength and conditioning research*. 2023.
684   31.   Kubo K, Ikebukuro T, Yata H. Effects of squat training with different depths on lower
685   limb muscle volumes. *European journal of applied physiology*. 2019;119(9):1933-42.
686   32.   Morrissey MC, Harman EA, Johnson MJ. Resistance training modes: specificity and
687   effectiveness. *Medicine and science in sports and exercise*. 1995;27(5).
688   33.   Dalgaard LB, Jorgensen EB, Oxfeldt M et al. Influence of Second Generation Oral
689   Contraceptive Use on Adaptations to Resistance Training in Young Untrained Women. *J
690   Strength Cond Res*. 2020.

34.   Romance R, Vargas S, Espinar S et al. Oral Contraceptive Use does not Negatively
Affect Body Composition and Strength Adaptations in Trained Women. *Int J Sports Med*.
2019;40(13):842-9.
35.   Dalgaard LB, Dalgas U, Andersen JL et al. Influence of Oral Contraceptive Use on
Adaptations to Resistance Training. *Front Physiol*. 2019;10:824.
36.   Ruzic L, Matkovic BR, Leko G. Antiandrogens in hormonal contraception limit muscle
strength gain in strength training: comparison study. *Croat Med J*. 2003;44(1):65-8.
37.   Riechman SE, Lee CW. Oral Contraceptive Use Impairs Muscle Gains in Young
Women. *J Strength Cond Res*. 2021.
38.   Oxfeldt M, Dalgaard LB, Jorgensen EB et al. Molecular markers of skeletal muscle
hypertrophy following 10 wk of resistance training in oral contraceptive users and
nonusers. *J Appl Physiol (1985)*. 2020;129(6):1355-64.
39.   Romero-Moraleda B, Coso JD, Gutiérrez-Hellín J, Ruiz-Moreno C, Grgic J, Lara B. The
Influence of the Menstrual Cycle on Muscle Strength and Power Performance. *Journal of
human kinetics*. 2019;68.
40.   Sakamaki-Sunaga M, Min S, Kamemoto K, Okamoto T. Effects of Menstrual Phase-
Dependent Resistance Training Frequency on Muscular Hypertrophy and Strength.
Journal of strength and conditioning research. 2016;30(6).
709