Past family selection in *Populus nigra* leads to an amplification of rust damages when neighbors are related

Authors: Mathieu Tiret[1,3*], Leopoldo Sánchez[2], Martin Lascoux[1].

Affiliations:

- 1: Program in Plant Ecology and Evolution, Department of Ecology and Genetics, EBC and SciLife Lab, Uppsala University, Uppsala, Sweden
- 2: BioForA, INRAE, ONF, Orléans, France.
- 3: IGEPP, INRAE, Institut Agro, Université de Rennes, Domaine de la Motte, Le Rheu, France.

* corresponding author: mathieu.tiret@inrae.fr

Abstract: (191 words)

Main text: 4,936 words.

keywords: group selection, genomic selection, Populus nigra, quantitative genetics.

Abstract:

Selfish individuals might prevail in the wild, but they often cause waste of nutritional

resources and decrease the population yield in breeding schemes. Re-orienting past

evolution towards less competitive paths (e.g., group selection) can limit negative

interactions between elite material and leads to a more efficient transformation of resources

into economically interesting traits. In particular, there are reasons to believe that, in the

presence of kin, individuals either will cooperate (kin selection) or over-compete (niche

partitioning). In either case, local relatedness might reveal interaction that can enhance the

predictive abilities of genomic models when accounted for. However, recurrent family

selection during domestication can have blurred the effect of relatedness on neighbors'

phenotypes. To overcome this difficulty, we analyzed data from the French breeding

program of *Populus nigra* L., where 1,452 genotypes were replicated six to eight times, each

time encountering a different neighborhood. We assessed local relatedness and investigated

genomic estimated breeding values on tree height and rust vulnerability with a single-step

GBLUP incorporating local relatedness as a covariate. We found that, when surrounded by

related individuals, rust damage is significantly more important, whereas no interaction was

visible on tree height.

Introduction

Breeding is redirecting evolutionary trade-offs towards new equilibria that often decrease fitness in favor of other economically interesting traits, or towards paths that were insufficiently explored in the wild, such as group selection (Weiner, 2019). When implemented artificially, group selection would favor candidates with a high performance in a group (e.g., robust productivity, tolerance to biotic or abiotic stress, or long-term response to selection). It is then not surprising that overall performance can be higher when performing group selection than when selecting candidates on their own individual performances (Wade et al, 2010). This is especially true when intraspecific competition is strong among candidates (Weiner et al., 2017). Indeed, selfish strategies, although increasing individual fitness, are often costly for the population - and breeding circumventing such strategies ended up with very successful results (e.g., Donald, 1981; Donald and Hamblin, 1983; Weiner et al., 2010). Despite the early success of some ideotypes optimizing group performance, current breeding programs do not necessarily focus on integrating this dimension (e.g., Denison et al., 2003; Murphy et al., 2017; Montazeaud et al., 2020). Consequently, the pool of elite genotypes used nowadays might be mostly composed of selfish genotypes, since early mass selection, bypassing group dynamic, favored them for their vigorous phenotypes (Murphy et al., 2017). It is therefore increasingly important for a sustainable breeding program to counteract the effect of natural selection, and, perhaps, a bit of past artificial selection.

In the specific case of plant breeding, resource competition between candidates occurs with the nearest plants (Casper et al., 1997; Milbau et al., 2007; File et al., 2011). The intensity of competition among neighbors may depend on the relatedness among them, or local relatedness, either because of kin selection, which predicts that relatives may cooperate

(Hamilton, 1964), or niche partitioning, which predicts, on the contrary, that similar

individuals may compete for similar resources (Silvertown, 2004). There are therefore

reasons to believe that local relatedness impacts phenotypes in one way or the other (Cahill

et al., 2011), so that understanding plasticity to the presence of related neighbors might

provide valuable information for the implementation of group selection in plant breeding.

Although kin recognition was reported to be not particularly relevant in crop species because

of their already high relatedness (e.g., Murphy et al., 2017), we might be able to detect it

nonetheless in breeding schemes that have been less intensive, such as forest tree

breeding. Group selection is in fact expected to give better genetic progress for perennial

species, as (i) inbreeding coefficient is much lower than in crop species, and (ii) positive or

negative interactions between neighbors lasts a lifetime.

Local relatedness creates micro-environment to which genotypes can respond to, so that,

even within a plot, strong genotype x environment interaction (GxE) can be expected.

Breeding on GxE is, however, known to be challenging (Diouf et al., 2020), especially for

non-model species where small SNP chips can lead to ascertainment bias or low heritability.

To meet this challenge, we chose to conduct our analyses on *Populus nigra* L.. Black poplar

is a Eurasian riparian forest tree that contributes, as a parent along with *Populus deltoides*

Bartr. ex Marsh., to one of the most widely used hybrid tree in forest breeding (*Populus x*

canadensis). The dataset of the French breeding program with 1,452 genotypes replicated

six to eight times created different neighborhoods for each replicate of a genotype (Pégard

et al., 2020). To detect any signal of group selection, we focused on two quantitative traits

relevant for breeding, tree height and rust vulnerability. Growth of Black poplar is particularly

susceptible to foliar rust, caused by a fungus (*Melampsora larici-populina* Kleb.) that leads to

a reduction in photosynthesis efficiency, and might lead to the death of susceptible clones

(Legionnet et al., 1999). Rust development is strongly influenced by vulnerability among

neighbors, whereas tree height is a poor proxy of competition among neighbors and a good

one of the fertility of the trial. In order to assess the contribution of local relatedness in

genomic selection (Meuwissen et al., 2001), we performed genomic evaluations with a

multitrait single-step Genomic Best Linear Unbiased Predictor (ssGBLUP; Legarra et al.,

2009; Christensen et al., 2010). Estimating breeding values with ssGBLUP is a solid

baseline in forest trees as recently shown (e.g., Cappa et al., 2019; Ratcliffe et al., 2017),

especially its multitrait version that is known to give higher accuracy (Calus et al., 2011). We

evaluated the ability of ssGBLUP models to explain phenotypic variance when including

local relatedness as a way of measuring the extent to which the phenotype of an individual

was impacted by the relatedness with its neighbors. We assessed local relatedness with

different subsets of SNPs, each maximizing or minimizing one of the following features:

Minor Allele Frequency, Ancestry Informativeness Coefficient, significance in a Genome

Wide Association Study, or heritability. We show that local relatedness significantly impacts

the phenotype of an individual and, when surrounded by related individuals, rust damage is

significantly more important, whereas no interaction was visible on tree height.

Material and Methods

Plant material. The study was based on previously published data (Pégard et al., 2019, 2020). Seventeen male and 17 female *Populus nigra* were sampled in France, from 21 natural populations and 13 from breeding programs. These individuals were used as parents in a factorial crossing plan. Clones of both parents and progeny, and a hundred additional individuals from breeding programs, were grown in four experimental trials at the same location (Guéméné-Penfao, France, 47°37'59"N, 1°49'59"W), each with a random block design. In total, the experiment comprised 10,301 trees (including 7,169 trees with genotype information), 42 full-sib families, 5 half-sib families, 1,452 genotypes (each replicated on average 7.09 times), and an Unknown Parent Group (UPG) of 105 genotypes (Table S1). Family size ranged from 1 to 119, with an average of 30.2 genotypes per family. Twenty genotypes and 17 families were shared across the four trials.

Phenotype measurements. We focused on two phenotypes of one-year-old trees: height and rust vulnerability (both measured on field). Height was assessed with a graduation rod (in cm), and rust vulnerability was assessed on a 1 to 9 scale (Legionnet et al., 1999). Across the trials, at least 95.29%, 96.05%, and 95.13% of the trees were phenotyped for height, rust vulnerability, or both traits, respectively. Given the low missing rate, we assumed that missing values only marginally unbalanced the block design, and so we used default parameters of the software packages for handling missing values when analyzing the data.

In each trial, every phenotype was first corrected for micro-environmental heterogeneity with a multitrait mixed model, including as explanatory variables a random block effect and a B-spline spatial autocorrelation correction. We used the function *remlf90* (R package *breedR*; Munoz and Sanchez, 2014) with the parameters 'model' set to 'splines' and 'method' set to

'em'. All further analyses were performed on the residuals of this model, i.e., on spatially

adjusted phenotypes.

Genomic data. Out of the 1,452 genotypes, 1,034 were genotyped (25 parents and 1,009

offspring), using a *Populus nigra* 12K custom Infinium Bead-Chip (Illumina, San Diego, CA)

(Faivre-Rampant et al., 2016). Details of the DNA extraction protocol, bioinformatic pipelines,

and SNP mapping on the 19 chromosomes are given in Faivre-Rampant et al. (2016) and

Pégard et al. (2019, 2020). It is worth noting that the SNP chip array included SNP

associated with, among other traits, rust resistance, but not with vertical growth. We further

filtered for a Minor Allele Frequency (MAF) of 5% − corresponding to a threshold of 50

individuals for a genotyping error; in total, 7,129 SNPs were retained for further analyses

(out of 7,513 initial SNPs).

Genomic relationship matrix. In order to build the relationship matrix necessary for

inferring breeding values, we combined pedigree and genomic information (Legarra et al.,

2009; Christensen et al., 2010). The pedigree relationship matrix A^Γ was built using the

tabular rule method (Emik and Terrill, 1949), modified to account for a single metafounder

(assuming a single base population for the Unkown Parent Group or UPG; Legarra et al.,

2015), with a self-relationship of $\gamma = 8\sigma_p^2$, with σ_p^2 the variance of allele frequencies across

markers (Garcia-Baccino et al., 2017). The combined relationship matrix H was summarized

in Aguilar et al. (2019), and is:

$H^{-1} = A^{\Gamma-1} + (0,0;0; G^{-1} - A_{22}^{\Gamma-1})$ (1),

where A_{22}^Γ was the pedigree relationship matrix for genotyped individuals, $G = (1-\alpha) (\omega_a +$

$\omega_b \times G_V) + \alpha A_{22}^\Gamma$, with G_V the genomic relationship matrix estimated from the SNPs and

scaled following the first scaling method of Van Raden (2008), α a scaling parameter (here

equal to 0.05), and ω_a and ω_b chosen to equate the average inbreeding and the average

relationships in G_v and A_{22}^Γ as in Christensen et al. (2012). As the sample size was relatively

small, the matrix H^{-1} was obtained by simply inverting H with the function *solve* (R package

base).

Population structure. In order to control for population structure and avoid spurious

associations, we computed a distance matrix $1 - H*$, where H* is the correlation matrix

obtained from H using the function *cov2cor* (R package *stats*), and the minus sign is to

convert similarities to distances. We then used Multidimensional Scaling (MDS) on the

distance matrix with the function *cmdscale_lanczos* (R package *refund*; Goldsmith et al.,

2022; Miller, 2022) with the parameter 'k' set to 5, and 'eig' set to 'TRUE'. Negative

eigenvalues were set to null, and the variance explained by an axis was computed as the

ratio between its squared eigenvalue and the sum of the squared eigenvalues. Additionally,

to assess how well the family structure was captured, we measured the variable V_F defined

as the average within family variance on the first five axes (the lower the V_F, the better the

family structure is captured). To make MDSs comparable, we normalized V_F by the sum of

squares of the first five eigenvalues.

Breeding value inferences. The spatially adjusted phenotypes were then used to infer the

breeding values with a multitrait single-step Genomic Best Linear Unbiased Predictor

(ssGBLUP), as follows:

$Y_{ijk} = \mu + mds1_j + mds2_j + TRIAL_i + ANIMAL_j + e_{ijk}$,   (2)

where 'Y_{ijk}' is the phenotype of the k-th replicate of the j-th genotype in the i-th trial, μ is the

grand mean, '$mds1_j$' and '$mds2_j$' the fixed effects on the j-th genotype of the first and second

axes of the MDS respectively, '$TRIAL_i$' the random effect of the i-th trial, '$ANIMAL_j$' the

polygenic effect of the j-th genotype, and 'e_{ijk}' the residuals. Both the trial effect and the

residuals follow a Gaussian distribution with diagonal covariance matrices, and the polygenic

effect follows a Gaussian distribution with a covariance matrix equal to k.H x Σ, where k is a

scaling parameter equal to 1 - γ/2 (Legarra et al., 2015), H the combined relationship matrix,

x the Kronecker product, and Σ the 2x2 covariance matrix between height and rust

vulnerability. Estimates of coefficients and variance components of model (2) were obtained

with the function *remlf90* (R package *breedR*) through one run with the 'method' parameter

set to 'em' until convergence, then another run with the parameter 'method' set to 'ai' and the

parameter 'progsf90.option' set to 'maxrounds 1'. The output estimate of 'ANIMAL$_j$' is the

Estimated Breeding Value (EBV) of the j-th genotype. The heritability of the traits was

estimated as the diagonal of Σ divided by the sum of the diagonals of Σ and the estimated

variance of the residuals. The predictive ability (PA) of the model was defined as Pearson's

correlation coefficient between the breeding values re-estimated from the effect sizes (see

below) and the adjusted phenotypes corrected by the trial effect (i.e., Y_{ijk} − TRIAL$_i$).

Genome-Wide Association Study (GWAS). Following the equations of Aguilar et al.

(2019), the effect size of the SNPs was backsolved from the EBVs, along with their standard

error and their p-values. This process was performed trait by trait (i.e., without accounting for

the multitrait dimension) without loss of information, as the Pearson's correlation between

the ssGBLUP's EBV and the breeding values re-estimated by the effect sizes was greater

than 0.999.

With the estimated effect sizes and the standard errors, we also computed an alternative to

the p-value accounting for multiple testing: the local false sign rate (lfsr) estimated with an

empirical Bayes approach for adaptive shrinkage using the function *ash* (R package *ashr*;

Stephens, 2017). In order to detect GWAS peaks (regions in the chromosomes with a high

density of significant SNPs), we used a custom peak detection script on the profile of lfsr

(Tiret & Milesi, 2021) with default parameters.

Neighborhood. In order to account for the effect of the neighborhood, we estimated the breeding values with an additional fixed effect, the local relatedness, defined as the average relationship (from the matrix H) between a focal individual and its neighbors (eight or less if on boundaries; as the king's moves on chess). Local relatedness was orthogonal to any micro-environmental effect, because of the random block design. The re-estimated breeding values were denoted nEBV.

SNP subsets. We investigated the effect of sampling SNPs on the ssGBLUP (i.e., weighted ssGBLUP). We selected subsets based on population and quantitative genetics features computed after a first run of ssGBLUP: MAF, Ancestry Informativeness Coefficient (AIM; Rozenberg et al., 2003), and for each trait, significance in the GWAS, and heritability. For each feature, we kept one-tenth of the whole set (713 SNPs, a sample size close to previous studies on weighted GBLUP, e.g., Li et al., 2018) with either the smallest or the largest values, ending up with 12 subsets. AIM was estimated per family with the R script provided by Cappa et al. (2022), and SNP heritability was estimated as $2p(1-p)a^2$, where p is the MAF, and a the effect size (assuming no dominance nor interaction deviation of the genetic variance; p.129, Falconer, 1989). We re-estimated the combined relationship matrix for each SNP subset by re-estimating in the equation (1) the genomic relationship matrix G_V with the SNP subset. With the re-estimated matrices H, we re-performed the MDSs, and re-estimated the PAs of the EBVs and nEBVs (where local relatedness were measured with the re-estimated H).

Statistical tests. For each SNP subset, we computed the confidence interval of the Predictive Ability (PA) with a 1000 iteration bootstrap (re-sampling individuals), without re-estimating the covariance matrix Σ. Here, the bootstrap was only used to assess the variance, so that we centered the bootstrapped PA around the true PA. To assess the effect of sampling, these subsets were compared to 1000 random samples of 713 SNPs (the size of the SNP subsets). Comparisons were performed with a Student's one-sample *t*-test

(denoted t_1 with a degree of freedom or df of 999 corresponding to the number of bootstrap

iterations), or a Welch two-sample t-test (denoted t_2 with a varying df), both using the

function *t.test* (R package *stats*). The sign of the t statistic is arbitrarily reported as positive

when greater than a baseline (e.g., the true PA), and negative when not. Models were

compared with a likelihood ratio test with a df of 1 using the function *lrtest* (R package *lmtest*;

Zeileis & Hothorn, 2002). All scripts were written in R v4.1.3 (R Core Team, 2022).

Results

Once relatedness among neighbors is actually shown to be correlated with focal

phenotypes, the difficulty in telling apart kin selection from niche competition lies in finding

outwhether relatedness is the cause, as the expression of kin selection or niche competition,

or a consequence, as the expression of a confusion between relatedness and phenotypic

interaction. In the latter case, the focal phenotype would be modified by that of its neighbors

through phenotypic interaction, and if phenotypic interaction increases when phenotypes are

similar (i.e., directional selection, e.g., competition for resource uptake), it can be confused

with the effect of relatedness, as genetically related individuals tend to have similar

phenotypes. Tree height and rust vulnerability have been the choice of our focus; both

exhibit high heritabilities (0.469 and 0.666 respectively), high predictive abilities (0.677 and

0.755 respectively), and a strong independence, with an estimated genetic correlation of

0.022. The low correlation is consistent with some previous reports (e.g., Beaulieu et al.,

2020; Liu et al., 2022), though not all (e.g., Lenz et al., 2021; Cappa et al., 2022); we will

assume in our case that orthogonality is strong enough to show that the traits are non-

redundant. Whether investigating highly heritable and orthogonal traits can help detecting kin

selection is yet unknown, and will be developed further on, but these preliminary results

make a solid case for tree height and rust vulnerability, and the choice of the SNP chip array.

Kin selection is a local relationship that can be hard to evaluate accurately (e.g., blurred by

micro-environmental effects); we posit that this problem can be overcome by investigating

the extent to which SNPs structuring families (i.e., kins) control phenotypes of the focal

individual, compared to other subsets of SNPs. Genetic variance explained by the markers

was low for both tree height (0.051) and rust vulnerability (0.058), displaying an expected

missing heritability, because of the size of the SNP chip (Yang et al., 2010); association

studies on our dataset was then expected at best to underestimate SNP contributions. On

the one hand, the GWAS was not solely artefactual, as it did not overfit phenotypic variation:

swapping phenotypes among family members of the same block (i.e., only randomly

mismatching Mendelian sampling) significantly reduced the average significance of the

SNPs ($t_1 < -2.59 \times 10^2$, $p < 0.001$). On the other hand, only one genetic variant was

significantly associated with rust vulnerability, located on the chromosome 1, and none for

tree height (Fig. 1A, 1B). The Q-Q plot on rust vulnerability revealed that some SNPs were

significantly associated with a level of significance well above the 95% confidence interval

(Fig. 1D); most of them being in the identified peak. None of the SNPs were significantly

associated with tree height after Bayesian shrinkage, and Q-Q plot revealed a classic

population structure overcorrection (Fig. 1C). We ascribe the difference in significance

between tree height and rust vulnerability to the SNP chip that was designed over QTLs and

expressional candidate genes related to rust resistance among other traits, but not tree

height (Faivre-Rampant et al., 2016). Altogether, although the genomic evaluation (i.e., when

using all SNPs) seems robust and consistent, SNP-wise approaches such as GWAS were

noticeably not robust to the SNP chip design. From here onwards, to account for this design

bias, we will only work with the subsets of SNPs described above, not single SNPs.

293 Common and most heritable alleles structured families

The population under study is structured in a hierarchical way: first into population structure,

then in family structure. Population structure, captured by the first eigenvalues (Patterson et

al., 2006), substantially shaped genetic diversity, as the first two eigenvalues alone

explained more than 50% of the variance (40.0% and 24.5% respectively). Because the

population underwent family selection during breeding in the past, family structure is strong

and can even be apprehended visually (Fig. 2). Several statistics significantly supports the

strong family structure: (i) randomly shuffling the family ID increased by 9.96 times the within

"family" variance ($t_1 = 3.70 \times 10^3$, $p < 0.001$); (ii) when restricting to common alleles, the first

two eigenvalues of the PCA still explained a large part of the variance (56.2%), showing that

family structure is bound to selection, as expected under family selection.

Population structure is generally assumed to be reflected by polymorphism atneutral genes

scattered across the genome, to the point where sufficiently large samples of SNPs can

reveal population structure quite accurately. That is not the case for family structure in our

dataset, as random samples of SNPs performed quite poorly compared to the whole set of

SNPs ($t_1 = 2.12 \times 10^2$, $p < 0.001$), suggesting a non-random distribution of the family

structure information across the genome. Actually, the "most" neutral and the "most"

selected SNPs had the strongest structuring power on the family level (Fig. 3): SNPs with

the largest AIM, performing better than random samples ($t_1 = 3.86 \times 10^2$, $p < 0.001$), and

even than the whole set as can be expected under neutrality; and SNPs putatively under

selection, i.e., with the largest MAF and the largest heritability, performing better than

random samples ($t_1 > 1.09 \times 10^2$, $p < 0.001$). Additionally, and supporting the aforementioned

bias of SNP-wise approaches, both significant and non-significant SNPs (in the GWAS) were

only slightly different from random samples in regards to family structure though significant (|

$t_1| > 5.52$, $p < 0.001$).

Local relatedness caused an amplification of rust damage

Although the growth of focal individuals was not affected by their neighbors, damage due to

rust was more important the more related the focal individuals were with their neighbors.

Indeed, the genomic models incorporating local relatedness as a covariate had significantly

higher likelihood than without for rust vulnerability ($\chi^2 = 9.83$, $p = 0.002$), but not for tree

height (χ^2 = 0.09, p = 0.236). As for family structure which is driven by some SNPs, different

categories of SNPs had a different impact on focal phenotypes. When accounting for

relatedness via the different subsets of SNPs, the Akaike Information Criterion (AIC)

generally decreased (Fig. S2). However, for rust vulnerability, only the subset of SNPs with

the largest MAF significantly increased the likelihood of the model (χ^2 = 4.02, p = 0.04), that

is the one that best captured the family structure. Likelihood was not significantly improved

by SNPs with the largest AIM (χ^2 = 0.15, p = 0.703) or the largest heritability (χ^2 = 2.65, p =

0.104, respectively), and was significantly worse with other subsets (χ^2 > 6.66, p < 0.010).

For tree height, no subset of SNPs improved the likelihood (χ^2 < 3.48, p > 0.062). In other

words, when the probability of focal individuals and neighbors to be in the same family was

high, focal phenotypes were more affected by their neighbors; in this case, family

neighboring caused significantly more damage by rust.

One of the differences between tree height and rust vulnerability is the genetic architecture:

the GWAS shows a monogenic architecture of rust vulnerability as opposed to a putative

polygenic architecture of tree height, hence a "simpler" architecture for rust vulnerability.

Even if it should be taken cautiously, it is not uncommon to encounter a simpler genetic

architecture for response to biotic constraints compared to complex quantitative traits (e.g.,

gene-for-gene resistance; Flor, 1971). Consequently, many haplotype combinations can lead

to tall individuals, but few to rust resistance, because a polygenic architecture is redundant;

in other words, when individuals are related, their probability of both being resistant is higher

than that of both being tall. Local relatedness thus reveals a shared vulnerability within the

neighborhood. Results suggest that shared vulnerability is the cause of rust damage

amplification in related neighborhood. Supporting this claim, the covariance between random

pairs of neighboring individuals' rust vulnerability was enriched by local relatedness (from

0.28 for the whole set to 0.43 for the highest quartile), suggesting again that local

relatedness was a proxy of phenotypes being correlated due to a correlation of genotypes.

In spite of genomic models being sensitive to local relatedness, improvement of predictive

abilities (PA) was limited - probably due to a lack of power. Indeed, incorporating local

relatedness did not significantly change the PA neither with the whole set of SNPs ($|t_2|$ <

0.594, p > 0.553), nor with any of the subset ($|t_2|$ < 1.03, p > 0.301), with a slightly better PA

when incorporating local relatedness for SNPs with the smallest MAF on rust vulnerability (t_2

= 1.73, df = 2.00 x 10^3, p = 0.0841), and SNPs with the smallest tree height significance on

rust vulnerability (t_2 = 1.97, df = 2.00 x 10^3, p = 0.0491). See the Supplementary materials for

further details.

Discussion

Genomic selection in forest tree breeding has successfully predicted growth and wood quality for many species (Lenz et al., 2020), such as Eucalyptus (e.g. Resende et al., 2012), pines (e.g., Isik et al., 2016), or spruces (e.g., Beaulieu et al., 2014). The predictive ability (PA) of the underlying genomic models relies on the ability of genotypic data to capture the covariance between candidate loci, thereby reflecting how much genetic similarity explains phenotypic similarities (Powell et al., 2010). An additional source of phenotypic covariances is micro-environmental effects, such as local relatedness, and results clearly demonstrates that in some cases, interaction between neighbors cause an individual to grow differently. In the case of the Black poplars under study, for rust vulnerability the interaction was more likely due to phenotypic interaction rather than kin selection/niche competition.

SNP chip designs strongly impact the analysis

In order to understand what drove the phenotypic covariance between individuals, we subset 10% of the SNPs according to their MAF, AIM, significance in a GWAS, or heritability, thus redefining local relatedness with a different set of linkage disequilibrium. Subsetting SNPs is commonplace in genomic selection, mainly to reduce costs when, for instance, commodity value per individual is low (e.g., Luo et al., 2021), or when imputation is powerful enough to make genomic predictions possible with low-density markers (e.g., Kijas et al., 2017; Sanchez et al., 2017; Tsai et al., 2017). So far, different subsetting techniques have been proposed: AM-BLUP whereby selecting relevant regions of chromosomes (Astle & Balding, 2009; but see critiques from Gualdrón Duarte et al., 2020), TA-BLUP equivalent to our set of heritable SNPs (Zhang et al., 2010), from GWAS (Luo et al., 2021), or from machine learning methods (Li et al., 2018). In the present paper, we tried to integrate this large spectrum of methods in a common framework to assess their relative efficiency.

The use of subsets of SNPs in genomic selection is an ongoing debate, as doing so ended

in some studies with better predictive abilities (e.g., Dong et al., 2016; Luo et al., 2020;

Cappa et al., 2022), whereas, as our results showed, some studies showed otherwise (e.g.,

Lu et al., 2020; Thistlethwaite et al., 2020). The second option can be understood when

following the infinitesimal model (Fisher, 1918), or its recent development, the omnigenic

model (e.g., Boyle et al., 2017). In particular, subsetting reduces the scope of linkage

disequilibrium (Thistlethwaite et al., 2020), and deleting common SNPs ends up with a

considerable loss of heritability (Yang et al., 2010; Biddanda et al., 2020). Our results

suggest that subsetting can be successful, as it was here for rust vulnerability, only because

the SNP chip design matched the trait or past selection. Genomic selection should therefore

be carefully designed in accordance with the forecasted usage of a low-density marker chips

if any, and acknowledging that extending its usage can be fraught with difficulties.

409 Selection of relatedness or co-selection of relatives?

Interactions between neighbors may depend on local relatedness, and whether this

interaction is positive or negative depends on the relative strength of kin selection (Hamilton,

1964) and niche partitioning (Silvertown, 2004). Although the importance of group selection

in nature is controversial (Maynard Smith, 1964; Lewontin, 1970; Wilson & Wilson, 2007;

Okasha, 2020), local relatedness has been shown to be perceived through kin recognition in

plant species (e.g., Dudley et al., 2007; Murphy et al., 2009). Competition can be accounted

for in a quantitative genetic model by assigning to an individual one direct genetic effect

(DGE) as the effect of its genes on its own phenotype, and one indirect genetic effect (IGE)

as the effect of its genes on phenotypes of surrounding individuals (Griffing, 1967; potential

applications in Muir, 2005). Although adding complexity to the model, including competition

proved to reduce bias in covariance estimation (Costa e Silva et al., 2013). In our case, on

top of all our attempts to incorporate interactions, IGE resulted in a lower predictive ability

than with the ssGBLUP baseline (results not shown).

The simpler genetic architecture of rust vulnerability compared to that of tree height might

play a role in shaping the phenotypes. Family structure was driving similarity of rust

vulnerability, and combination of similar vulnerability resulted in increased rust damage. The

amplifying effect is even stronger in perennials as shared vulnerability can be perceived by

successive generations of pathogens. In agreement with the higher resilience in regards to

diseases confered by varietal association in a field (e.g., Smithson and Lenne, 1996; Burdon

and Thrall, 2009), locally pyramiding vulnerability amplified rust damage. Resistance

similarity and relatedness were strongly connected, but it is likely that the more complex the

genetic architecture, the easier it is to uncorrelate phenotypic similarity and relatedness, and

so profitable for breeding schemes when trying to limit local pyramiding while using family

selection.

Whether there was kin selection/competition on top of phenotypic interaction or we simply

lacked statistical power is hard to decide. Selection on local relatedness due to kin selection

or niche partitioning - namely, selection on relatedness - might be confused with the fact that

selection leads to phenotypic convergence, that in turn leads to a genotypic convergence -

namely, a co-selection of related. The random block design and repetition can partly limit this

confusion, but it is evident that interpreting the data is fraught with difficulties, as the tallest

individuals showed also a higher relatedness with the rest of the population. The fact that

family structure was mainly carried by SNPs likely associated with selection (following the

hypothesis of Yang et al., 2010 and Biddanda et al., 2020) suggest that relatives were co-

selected, as is often the case in breeding programs implementing family selection. An

inevitable consequence is that, as seen in crop selection, interesting genotypes for kin

selection/cooperation were lost during domestication (Fréville et al., 2022).

18   18

Acknowledgements

The authors acknowledge the Uppsala University and the European Union's Horizon 2020 B4EST for basic functioning and postdoctoral grant for MT. The computations and data handling were enabled by resources provided by the Swedish National Infrastructure for Computing (SNIC 2017-7-296) at UppMax partially funded by the Swedish Research Council through grant agreement no. 2018-05973.

Author Contributions

MT was responsible for writing the report, conducting the search, extracting and analysing data, interpreting results, updating reference lists. LS was responsible for designing the experiments and screening potentially eligible studies. ML and LS contributed to interpreting results and provided feedback on the report.

Competing Interests

The authors declare that there is no conflict of interest.

Data Archiving

The data that support the findings of this study are openly available on DATA INRAE at https://data.inrae.fr/privateurl.xhtml?token=b79ab1ca-ebb9-47c6-9272-1568c0c33d70.

Rerefences

Aguilar, I., Legarra, A., Cardoso, F., Masuda, Y., Lourenco, D., & Misztal, I. (2019). Frequentist p-values for large-scale-single step genome-wide association, with an application to birth weight in American Angus cattle. *Genetics Selection Evolution*, *51*(1), 1-8.

Astle, W., & Balding, D. J. (2009). Population structure and cryptic relatedness in genetic association studies. *Statistical Science*, *24*(4), 451-471.

Beaulieu, J., Doerksen, T., Clément, S., MacKay, J., & Bousquet, J. (2014). Accuracy of genomic selection models in a large population of open-pollinated families in white spruce. *Heredity*, *113*(4), 343-352.

Beaulieu, J., Nadeau, S., Ding, C., Celedon, J. M., Azaiez, A., Ritland, C., et al. (2020). Genomic selection for resistance to spruce budworm in white spruce and relationships with growth and wood quality traits. *Evolutionary applications*, *13*(10), 2704-2722.

Biddanda, A., Rice, D. P., & Novembre, J. (2020). A variant-centric perspective on geographic patterns of human allele frequency variation. *Elife*, *9*, e60107.

Burdon, J. J., & Thrall, P. H. (2009). Coevolution of plants and their pathogens in natural habitats. *Science*, *324*(5928), 755-756.

Boyle, E. A., Li, Y. I., & Pritchard, J. K. (2017). An expanded view of complex traits: from polygenic to omnigenic. *Cell*, *169*(7), 1177-1186.

Cahill Jr, J. F., & McNickle, G. G. (2011). The behavioral ecology of nutrient foraging by

plants. *Annual Review of ecology, evolution, and systematics*, *42*, 289-311.

Calus, M. P., & Veerkamp, R. F. (2011). Accuracy of multi-trait genomic selection using

different methods. *Genetics Selection Evolution*, *43*(1), 1-14.

Cappa, E. P., de Lima, B. M., da Silva-Junior, O. B., Garcia, C. C., Mansfield, S. D., &

Grattapaglia, D. (2019). Improving genomic prediction of growth and wood traits in

Eucalyptus using phenotypes from non-genotyped trees by single-step GBLUP. *Plant*

Science, *284*, 9-15.

Cappa, E. P., Ratcliffe, B., Chen, C., Thomas, B. R., Liu, Y., Klutsch, J., et al. (2022).

Improving lodgepole pine genomic evaluation using spatial correlation structure and SNP

selection with single-step GBLUP. *Heredity*, *128*(4), 209-224.

Casper, B. B., & Jackson, R. B. (1997). Plant competition underground. *Annual review of*

ecology and systematics, 545-570.

Christensen, O. F., & Lund, M. S. 2010. Genomic prediction when some animals are

not genotyped. Genetics Selection Evolution 42, 2.

Christensen, O. F., Madsen, P., Nielsen, B., Ostersen, T., & Su, G. (2012). Single-step

methods for genomic evaluation in pigs. *animal*, *6*(10), 1565-1571.

Costa e Silva, J., Kerr, R.J. (2013) Accounting for competition in genetic analysis, with

particular emphasis on forest genetic trials. *Tree Genetics & Genomes, 9,* 1-17.

Denison, R. F., Kiers, E. T., & West, S. A. (2003). Darwinian agriculture: when can humans

find solutions beyond the reach of natural selection? *The quarterly review of biology*, *78*(2),

145-168.

Diouf, I., Derivot, L., Koussevitzky, S., Carretero, Y., Bitton, F., Moreau, L., & Causse, M.

(2020). Genetic basis of phenotypic plasticity and genotype× environment interactions in a

multi-parental tomato population. Journal of Experimental Botany, 71(18), 5365-5376

Donald, C.M. (1981). Competitive plants, communal plants, and yields in wheat crops.

L. Evans, W.J. Peacock (Eds.), Wheat Science – Today and Tomorrow, Cambridge

University Press.

Donald, C.M., Hamblin, J. (1983). The convergent evolution of annual seed crops in

agriculture. In Advances in agronomy (ed. NC Brady), pp. 97–143. New York, NY:

Academic Press.

Dong, L., Xiao, S., Chen, J., Wan, L., & Wang, Z. (2016). Genomic selection using extreme

phenotypes and pre-selection of SNPs in large yellow croaker (Larimichthys crocea). *Marine*

Biotechnology, *18*(5), 575-583.

Fisher, R.A. (1918). The correlation between relatives under the supposition of Mendelian

inheritance. *Trans. Roy. Soc. Edinburgh 52*, 399-433.

Gualdrón Duarte, J. L., Gori, A. S., Hubin, X., Lourenco, D., Charlier, C., Misztal, I., & Druet,

T. (2020). Performances of Adaptive MultiBLUP, Bayesian regressions, and weighted-

GBLUP approaches for genomic predictions in Belgian Blue beef cattle. *BMC genomics*,

21(1), 1-18.

Dudley, S. A., & File, A. L. (2007). Kin recognition in an annual plant. *Biology Letters*, *3*(4),

435-438.

Emik, L. O., & Terrill, C. E. (1949). Systematic procedures for calculating inbreeding

coefficients. *Journal of Heredity*, *40*(2), 51-55.

Faivre-Rampant, P., Zaina, G., Jorge, V., Giacomello, S., Segura, V., Scalabrin, S., et al.

(2016). New resources for genetic studies in Populus nigra: Genome-wide SNP discovery

and development of a 12k Infinium array. *Molecular ecology resources*, *16*(4), 1023-1036.

Falconer, D.S. (1989). Introduction to quantitative genetics (3rd edition, p. 129). Pearson

Education India.

File, A. L., Murphy, G. P., & Dudley, S. A. (2012). Fitness consequences of plants growing

with siblings: reconciling kin selection, niche partitioning and competitive ability. *Proceedings*

of the Royal Society B: Biological Sciences, *279*(1727), 209-218.

Flor, H. H. (1971). Current status of the gene-for-gene concept. *Annual review of*

phytopathology, *9*(1), 275-296.

Fréville, H., Montazeaud, G., Forst, E., David, J., Papa, R., & Tenaillon, M. I. (2022). Shift in

beneficial interactions during crop evolution. *Evolutionary Applications*, *15*(6), 905-918.

Garcia-Baccino, C. A., Legarra, A., Christensen, O. F., Misztal, I., Pocrnic, I., Vitezica, Z. G.,

& Cantet, R. J. (2017). Metafounders are related to F st fixation indices and reduce bias in

single-step genomic evaluations. *Genetics Selection Evolution*, *49*(1), 1-14.

Goldsmith, J., Scheipl, F., Huang, L., Wrobel, J., Di, C., Gellar, J., et al. (2022). refund:

Regression with Functional Data. R package version 0.1-26.

https://CRAN.R-project.org/package=refund

Griffing, B. (1967). Selection in reference to biological groups I. Individual and group

selection applied to populations of unordered groups. *Australian Journal of Biological*

Sciences, *20*(1), 127-140.

Hamilton, W. D. (1964). The genetical evolution of social behaviour. II. *Journal of theoretical*

biology, *7*(1), 17-52.

Isik, F., Bartholomé, J., Farjat, A., Chancerel, E., Raffin, A., Sanchez, L., et al. (2016).

Genomic selection in maritime pine. *Plant Science*, *242*, 108-119.

Kijas, J., Elliot, N., Kube, P., Evans, B., Botwright, N., King, H., et al. (2017). Diversity and

linkage disequilibrium in farmed Tasmanian Atlantic salmon. *Animal genetics*, *48*(2), 237-

241.

Legarra, A., Aguilar, I., & Misztal, I. (2009). A relationship matrix including full pedigree and

genomic information. *Journal of dairy science*, *92*(9), 4656-4663.

Legarra, A., Christensen, O. F., Vitezica, Z. G., Aguilar, I., & Misztal, I. (2015). Ancestral

relationships using metafounders: finite ancestral populations and across population

relationships. *Genetics*, *200*(2), 455-468.

Legionnet, A., Muranty, H., & Lefèvre, F. (1999). Genetic variation of the riparian pioneer

tree species Populus nigra. II. Variation in susceptibility to the foliar rust Melampsora larici-

populina. *Heredity*, *82*(3), 318-327.

Lenz, P. R., Nadeau, S., Mottet, M. J., Perron, M., Isabel, N., Beaulieu, J., & Bousquet, J.

(2020). Multi-trait genomic selection for weevil resistance, growth, and wood quality in

Norway spruce. *Evolutionary applications*, *13*(1), 76-94.

Lewontin, R. C. (1970). The units of selection. *Annual review of ecology and systematics*, 1-

18.

Li, B., Zhang, N., Wang, Y. G., George, A. W., Reverter, A., & Li, Y. (2018). Genomic

prediction of breeding values using a subset of SNPs identified by three machine learning

methods. *Frontiers in genetics*, *9*, 237.

Liu, Y., Erbilgin, N., Ratcliffe, B., Klutsch, J. G., Wei, X., Ullah, A., et al. (2022). Pest

defences under weak selection exert a limited influence on the evolution of height growth

and drought avoidance in marginal pine populations. *Proceedings of the Royal Society B*,

289(1982), 20221034.

Lu, S., Liu, Y., Yu, X., Li, Y., Yang, Y., Wei, M., et al. (2020). Prediction of genomic breeding

values based on pre-selected SNPs using ssGBLUP, WssGBLUP and BayesB for

Edwardsiellosis resistance in Japanese flounder. *Genetics Selection Evolution*, *52*(1), 1-10.

Luo, Z., Yu, Y., Xiang, J., & Li, F. (2021). Genomic selection using a subset of SNPs

identified by genome-wide association analysis for disease resistance traits in aquaculture

species. *Aquaculture*, *539*, 736620.

Maynard Smith, J. (1964). Group selection and kin selection. *Nature*, *201*(4924), 1145-1147.

Meuwissen, T. H. E. (1997). Maximizing the response of selection with a predefined rate of

inbreeding. *Journal of animal science*, *75*(4), 934-940.

Meuwissen, T. H., Hayes, B. J., & Goddard, M. (2001). Prediction of total genetic value using

genome-wide dense marker maps. *genetics*, *157*(4), 1819-1829.

Milbau, A., Reheul, D., De Cauwer, B., & Nijs, I. (2007). Factors determining plant–

neighbour interactions on different spatial scales in young species-rich grassland

communities. *Ecological research*, *22*(2), 242-247.

Miller, D. L. (2022). porridge: Principal Co-Ordinate Ridge Regression. R package version

1.2.

Montazeaud, G., Rousset, F., Fort, F., Violle, C., Fréville, H., & Gandon, S. (2020). Farming

plant cooperation in crops. *Proceedings of the Royal Society B*, *287*(1919), 20191290.

Muir, W. M. (2005). Incorporation of competitive effects in forest tree or animal breeding

programs. *Genetics*, *170*(3), 1247-1259.

Muñoz, F., and Sanchez, L. (2020). breedR: Statistical Methods for Forest Genetic

Resources Analysts. R package version 0.12-5. https://github.com/famuvie/breedR

Murphy, G. P., & Dudley, S. A. (2009). Kin recognition: competition and cooperation in

Impatiens (Balsaminaceae). *American journal of botany*, *96*(11), 1990-1996.

Murphy, G. P., Swanton, C. J., Van Acker, R. C., & Dudley, S. A. (2017). Kin recognition,

multilevel selection and altruism in crop sustainability. *Journal of Ecology (Oxford)*, *105*(4),

930-934.

Okasha, S. (2020). Altruism, group selection and correlated interaction. *The British journal*

for the philosophy of science.

Patterson, N., Price, A. L., & Reich, D. (2006). Population structure and eigenanalysis. *PLoS*

genetics, *2*(12), e190.

Pégard, M., Rogier, O., Bérard, A., Faivre-Rampant, P., Paslier, M. C. L., Bastien, C., et al.

(2019). Sequence imputation from low density single nucleotide polymorphism panel in a

black poplar breeding population. *BMC genomics*, *20*(1), 1-16.

Pégard, M., Segura, V., Muñoz, F., Bastien, C., Jorge, V., & Sanchez, L. (2020). Favorable

conditions for genomic evaluation to outperform classical pedigree evaluation highlighted by

a proof-of-concept study in poplar. *Frontiers in plant science*, *11*, 581954.

Powell, J. E., Visscher, P. M., & Goddard, M. E. (2010). Reconciling the analysis of IBD and

IBS in complex trait studies. *Nature Reviews Genetics*, *11*(11), 800-805.

R Core Team (2022). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL. https://www.R-project.org/

Ratcliffe, B., El-Dien, O. G., Cappa, E. P., Porth, I., Klápště, J., Chen, C., & El-Kassaby, Y.

A. (2017). Single-step BLUP with varying genotyping effort in open-pollinated Picea glauca.

G3: Genes, Genomes, Genetics, *7*(3), 935-942.

Resende, M. D., Resende Jr, M. F., Sansaloni, C. P., Petroli, C. D., Missiaggia, A. A.,

Aguiar, A. M., et al. (2012). Genomic selection for growth and wood quality in Eucalyptus:

capturing the missing heritability and accelerating breeding for complex traits in forest trees.

New Phytologist, *194*(1), 116-128.

Rosenberg, N. A., Li, L. M., Ward, R., & Pritchard, J. K. (2003). Informativeness of genetic

markers for inference of ancestry. *The American Journal of Human Genetics*, *73*(6), 1402-

1422.

Sanchez, M. P., Govignon-Gion, A., Croiseau, P., Fritz, S., Hozé, C., Miranda, G., et al.

(2017). Within-breed and multi-breed GWAS on imputed whole-genome sequence variants

reveal candidate mutations affecting milk protein composition in dairy cattle. *Genetics*

Selection Evolution, *49*(1), 1-16.

Silvertown, J. (2004). Plant coexistence and the niche. *Trends in Ecology & evolution*,

19(11), 605-611.

Smithson, J. B., & Lenne, J. M. (1996). Varietal mixtures: a viable strategy for sustainable

productivity in subsistence agriculture. *Annals of applied biology*, *128*(1), 127-158.

Stephens, M. (2017). False discovery rates: a new deal. *Biostatistics*, *18*(2), 275-294.

Thistlethwaite, F. R., Gamal El-Dien, O., Ratcliffe, B., Klápště, J., Porth, I., Chen, C., et al.

(2020). Linkage disequilibrium vs. pedigree: Genomic selection prediction accuracy in

conifer species. *PLoS One*, *15*(6), e0232201.

Tiret, M., & Milesi, P. Statistical Peak Detection for GWAS. Zenodo.

https://doi.org/10.5281/zenodo.5079556. Deposited 7 July 2021.

Tsai, H. Y., Matika, O., Edwards, S. M., Antolín–Sánchez, R., Hamilton, A., Guy, D. R., et al.

(2017). Genotype imputation to improve the cost-efficiency of genomic selection in farmed

Atlantic salmon. *G3: Genes, genomes, genetics*, *7*(4), 1377-1383.

VanRaden, P. M. (2008). Efficient methods to compute genomic predictions. *Journal of dairy*

science, *91*(11), 4414-4423.

Wade, M. J., Bijma, P., Ellen, E. D., & Muir, W. (2010). Group selection and social evolution

in domesticated animals. *Evolutionary applications*, *3*(5-6), 453-465.

Weiner, J., Andersen, S. B., Wille, W. K. M., Griepentrog, H. W., & Olsen, J. M. (2010).

Evolutionary Agroecology: the potential for cooperative, high density, weed-suppressing

cereals. *Evolutionary Applications*, *3*(5-6), 473-479.

Weiner, J., Du, Y. L., Zhang, C., Qin, X. L., & Li, F. M. (2017). Evolutionary agroecology:

individual fitness and population yield in wheat (Triticum aestivum).

Weiner, J. (2019). Looking in the wrong direction for higher-yielding crop genotypes. *Trends*

in Plant Science, *24*(10), 927-933.

Wilson, D. S., & Wilson, E. O. (2007). Rethinking the theoretical foundation of sociobiology.

The Quarterly review of biology, *82*(4), 327-348.

Yang, J., Benyamin, B., McEvoy, B. P., Gordon, S., Henders, A. K., Nyholt, D. R., et al.

(2010). Common SNPs explain a large proportion of the heritability for human height. *Nature*

genetics, *42*(7), 565-569.

Zeileis, A., & Hothorn, T. (2002). Diagnostic Checking in Regression Relationships. R News

2(3), 7-10.

Zhang, Z., Liu, J., Ding, X., Bijma, P., de Koning, D. J., & Zhang, Q. (2010). Best linear

unbiased prediction of genomic breeding values using a trait-specific marker-derived

relationship matrix. *PloS one*, *5*(9), e12648.

Figures

Figure 1. GWAS on tree height and rust vulnerability in Populus nigra. **(A)**. Manhattan plot of

the GWAS on tree height. The dots are the log significance of the markers. The dashed

horizontal line stands for the 5% significance level, and the solid horizontal line stands for

the 5% FDR level. **(B)**. Idem as panel A, for rust vulnerability. **(C)**. Q-Q plot of the GWAS on

tree height. The dots were the quantile of the markers; the solid line is the 1-slope segment;

the gray cross is the position of the median; and the gray area is the 95% confidence

interval. **(D)**. Idem as panel C, for rust vulnerability.

Figure 2. The first two axes of the Multidimensional Scaling (MDS) of Populus nigra. **(A)**.

MDS with labels of the fathers (with black large shapes). Colored and smaller shapes are the

offspring, the shape corresponding to its father. **(B)**. The same MDS as in panel A., but

labeled for the mothers.

Figure 3. Within family variance (V_F) for each subset of SNPs. The black dashed horizontal

line is the average V_F of randomly sampled SNPs, the gray dashed horizontal lines delimit

the 95% confidence interval, and the black horizontal dotted line is V_F of the entire set of

SNPs. Red dots are when selected for tree height, and blue for rust vulnerability. SNP

subsets: largest or largest MAF (+ or - MAF), largest or smallest AIM (+ or - AIM), most or

least significant (+ or - $\log_{10}(p)$), and most or least heritable for tree height (+ or - h^2).