The role of relatedness within a neighborhood in plant-to-plant interaction: effect on height growth and rust damage in *Populus nigra*

Authors: Mathieu Tiret[1,3*], Leopoldo Sánchez[2], Martin Lascoux[1].

Affiliations:

- 1: Program in Plant Ecology and Evolution, Department of Ecology and Genetics, EBC and SciLife Lab, Uppsala University, Uppsala, Sweden
- 2: BioForA, INRAE, ONF, Orléans, France.
- 3: IGEPP, INRAE, Institut Agro, Université de Rennes, Domaine de la Motte, Le Rheu, France.

* corresponding author: mathieu.tiret@inrae.fr

Abstract: (208 words)

Main text: 5,471 words.

keywords: group selection, local relatedness, competition, genomic selection, Populus nigra.

Abstract:

The competitive ability of domesticated plants, which may have conferred a fitness advantage in the wild, may result in a reduction of yield in agricultural and forestry contexts, as what matters is the group rather than the individual performance. Traits related to competitive ability can be affected by the presence or absence of related individuals in their neighborhood. Consequently, local relatedness might reveal plant-to-plant interaction that can enhance the predictive abilities of genomic models when accounted for, though it remains difficult to measure. To overcome this difficulty, we analyzed data from the French breeding program of *Populus nigra* L., where 1,452 genotypes were replicated six to eight times, each time encountering a different neighborhood. We assessed local relatedness and investigated genomic estimated breeding values on tree height and vulnerability to rust with a single-step GBLUP incorporating local relatedness as a covariate. The results indicate that incorporating local relatedness as an additional factor in GBLUP models has a significantly greater influence on resistance to rust than on tree height, though its overall effect on genomic predictions themselves was limited. The influence of local relatedness is small but likely trait-specific, and the genetic architecture of the trait under selection could attenuate or improve the efficacy of breeding for group performance.

Introduction

Breeding generally consists in redirecting evolutionary trade-offs towards new equilibria that often decrease fitness in favor of other economically interesting traits, or towards paths that were insufficiently explored by evolution in natural populations, such as group selection (Weiner, 2019). The importance of group selection, defined as the cooperation within groups or populations that would confer a selective advantage (Leigh, 2010), is controversial in nature (Maynard Smith, 1964; Lewontin, 1970; Wilson & Wilson, 2007; Okasha, 2020). However, in a breeding program, selecting for individuals that improve the overall performance (e.g., robust productivity, tolerance to biotic or abiotic stress, or average yield), by limiting unfavourable plant-to-plant interactions can be interesting (also known as 'communal ideotype'; Biernaskie, 2022).

Separating an individual's performance into what is due to its own genetics and to its neighbors' has a long history of theoretical development, especially in quantitative genetics (Griffing, 1967; Bijma & Wade, 2008; Ellen et al., 2016; Bailey et al., 2018). The performance of an individual is decomposed into direct and indirect breeding values (also called direct and indirect genetic effects or DGE/IGE), where the former is the intrinsic value of an individual without any competitors, and the latter the ability of the individual to increase or decrease performance due to competition. The estimation of the direct and indirect values is based on the variation of the neighborhood of a focal individual, at the cost of a quadratic increase in data requirements. Recent theoretical developments showed that accounting for IGE can limit the detrimental competitiveness of individuals (Montazeaud et al., 2020; Bourke et al., 2021; Lemoine et al., 2023).

Breeding can especially benefit from the IGE framework when intraspecific competition is strong among elite candidates (Weiner et al., 2017). Indeed, selfish strategies in the sense of selfish genes, promoting individual fitness at the expense of the population's collective interest (Dawkins, 1976), are often costly for the populations in breeding schemes - and breeding circumventing such strategies ended up with very successful results (e.g., Donald, 1981; Donald and Hamblin, 1983; Weiner et al., 2010). Competitive behavior can, therefore, be a burden that breeders might want to select against. Nonetheless, despite the early success of some ideotypes optimizing group performance, breeding programs do not necessarily focus on integrating this dimension (e.g., Denison et al., 2003; Murphy et al., 2017; Montazeaud et al., 2020). Consequently, the pool of elite genotypes used nowadays might be mostly composed of selfish genotypes, since early mass selection, bypassing group dynamics, favored selfish genotypes for their vigorous phenotypes (Murphy et al., 2017). It is therefore increasingly important in order to reach sustained genetic gain in breeding programs to counteract the effect of natural selection acting at the deployment phase, and to correct past artificial selection trajectories focussed solely on competitive (i.e., selfish) individuals.

In the specific case of plant breeding, resource competition between candidates occurs with the nearest plants (Casper et al., 1997; Milbau et al., 2007; File et al., 2012). The intensity of competition among neighbors may depend on their relatedness, or local relatedness (average relatedness in the vicinity of a given individual), either because of kin selection (explaining cooperation within a group because of relatedness; Hamilton, 1964; File et al., 2012; Ehlers & Bilde, 2019), or niche partitioning (explaining, on the contrary, that similar individuals may compete for similar resources; Silvertown, 2004; Guisan et al., 2005; Bengtsson et al., 2019). Local relatedness, which usually varies across trials due to varying neighborhood, can create spatial heterogeneity that, when not accounted for, biases genetic estimates. Therefore, plant-to-plant interactions can partly be due to local relatedness in one way or the other (Cahill et al., 2011), so that BLUP and derived models can be improved by

incorporating the information of neighboring genotypes. Although kin recognition was

reported to be not particularly relevant in crop species because of their already high

relatedness (e.g., Murphy et al., 2017), we might be able to detect it in breeding schemes

that have been less intensive, such as forest tree breeding with shorter linkage

disequilibrium and starting with a much wider genetic base (Grattapaglia & Resende, 2011).

Group selection is in fact expected to give better genetic progress for perennial species,

since (i) inbreeding coefficient is much lower than in crop species (even though inbreeding

depression can be higher; Lesaffre & Billiard, 2021), and (ii) positive or negative interactions

between neighbors last a lifetime.

Black poplar (*Populus nigra* L.) is a Eurasian riparian forest tree that contributes as a parent,

along with *Populus deltoides* Bartr. ex Marsh., to one of the most widely used hybrid tree in

forest breeding (*Populus x canadensis*), and is widely deployed as clones, constituting a

powerful case study of kin recognition. The dataset, stemming from the French breeding

program with 1,452 genotypes replicated six to eight times, created different neighborhoods

for each replicated genotype (Pégard et al., 2020), all resulting from controlled crosses of 34

parents from natural populations, some of which were already used in the breeding

programme for their performance. We focused on two quantitative traits relevant for

breeding, tree height and vulnerability to rust. Growth of Black poplar is particularly

susceptible to foliar rust, caused by the fungus *Melampsora larici-populina* Kleb.. In order to

assess the contribution of local relatedness in the context of genomic selection (Meuwissen

et al., 2001), we performed genomic evaluations with a multitrait single-step Genomic Best

Linear Unbiased Predictor (ssGBLUP; Legarra et al., 2009; Christensen et al., 2010).

Estimating breeding values with ssGBLUP is a solid baseline in forest trees as recently

shown (e.g., Cappa et al., 2019; Ratcliffe et al., 2017), and has successfully predicted

growth and wood quality for many species (Lenz et al., 2020), such as Eucalyptus (e.g.

Resende et al., 2012), pines (e.g., Isik et al., 2016), or spruces (e.g., Beaulieu et al., 2014).

We propose to assess the indirect genetic effect with local relatedness. More specifically, the

objective of the study was to measure the extent to which local relatedness impacts the focal

individuals' phenotypes, and to quantify the benefit of including local relatedness in genomic

prediction models. We also investigated the level of association between local relatedness

and the traits: rust development is influenced by susceptibility of nearby trees (Pinon & Frey,

1997; Burdon & Thrall, 1999), hence is expected to be associated to local relatedness,

whereas one-year-old tree height is expected to be a poor indicator of competition (as

seedlings are too small to overshadow their neighbors), and is most likely related to the

environmental quality of the trial. We show that (i) as expected, incorporating local

relatedness as an additional factor in GBLUP models has a significantly greater influence on

estimates of resistance to rust than on tree height, but (ii) its overall effect on genomic

predictions is limited. Differences in traits genetic architecture and its role in plant-to-plant

interactions is discussed.

139 Material and Methods

Plant material. The study is based on previously published data (Pégard et al., 2019, 2020).

Seventeen male and 17 female *Populus nigra* were sampled in France, from 21 natural

populations with no a priori selection and 13 from the French poplar breeding program for

their performance on a range of traits, including growth and resistance to rust. These

individuals were used as parents in a factorial crossing plan and double-pair mating scheme.

Clones of both parents and progeny, and a hundred additional individuals from breeding

programs, were grown in four sequential experimental trials (parallel strips of two lines of

trees, with an intra-spacing of 1m and an inter-spacing of 2 metres) at the same location

(Guéméné-Penfao, France, 47°37'59"N, 1°49'59"W), each with a randomized incomplete

block design of single tree-plots (1.0 m x 2.0 m) with six blocks each, except for one trial with

eight blocks. In total, the experiment comprised 10,301 trees (including 7,169 trees with

genotype information), 42 full-sib families, 5 half-sib families, 1,452 genotypes (each

replicated on average 7.09 times), and an Unknown Parent Group (UPG) of 105 genotypes

(Table S1). Family size ranged from 1 to 119, with an average of 30.2 genotypes per family.

Twenty genotypes and 17 families were shared across the four trials.

Phenotype measurements. We focused on two phenotypic traits on one-year-old trees:

height and vulnerability to rust (both measured on field). Height was assessed with a

graduation rod (in cm), and vulnerability to rust (natural infection) was assessed on a scale

of 1 to 9 : 1 when no rust was observed on the tree and 9 when more than 75% of lamina

was covered by rust on more than 25% of the leaves (Legionnet et al., 1999). The

homogeneity of rust pressure across the trial was assured by the level of infection measured

on control individuals, being equally infected across blocks (the control cv. BDG that is

infected with an average score of 4.37 ± 0.65 across trials, with scores ranging from 3 to 5).

Some missing values were present across the trials (dead trees or unexploitable data), but

at least 95.29%, 96.05%, and 95.13% of the trees were phenotyped for height, vulnerability

to rust, or both traits, respectively. Given the low missing rate, we assumed that missing

values only marginally unbalanced the block design, and so we used default parameters of

the software packages for handling missing values when analyzing the data.

In each trial, every phenotype was first corrected for micro-environmental heterogeneity with

a multitrait mixed model as follows:

$Y_{ijkl} = \mu + BLOCK_i + SPLINE + e_{ijkl}$

where 'Y_{ijkl}' is the l-th phenotype of the k-th replicate of the j-th genotype in the i-th trial, μ is

the grand mean, '$BLOCK_i$' the random effect of the i-th block, 'SPLINE' B-spline spatial

autocorrelation correction over the surface of rows and columns (Cappa & Cantet, 2007),

and 'e_{ijkl}' the residuals. Both the block effect and the residuals follow a Gaussian distribution

with diagonal covariance matrices, and the model was fitted separately by trial. We used the

function *remlf90* (R package *breedR* v0.12; Munoz and Sanchez, 2014) with the parameters

'model' set to 'splines' and 'method' set to 'em'. All further analyses were performed on the

residuals of this model, i.e., on spatially adjusted phenotypes.

Genomic data. Out of the 1,452 genotypes, 1,034 were genotyped (25 parents and 1,009

offspring), using a *Populus nigra* 12K custom Infinium Bead-Chip (Illumina, San Diego, CA)

(Faivre-Rampant et al., 2016). Details of the DNA extraction protocol, bioinformatic pipelines,

and SNP mapping on the 19 chromosomes are given in Faivre-Rampant et al. (2016) and

Pégard et al. (2019, 2020). It is worth noting that the SNP chip array included SNP in QTLs

and expression candidate genes associated with, among other traits, resistance to rust, but

not with vertical growth. We retained SNPs with a Minor Allele Frequency (MAF) > 5%; in

total, 7,129 SNPs were retained for further analyses (out of 7,513 initial SNPs).

Genomic relationship matrix. In order to build the relationship matrix necessary for inferring breeding values, pedigree and genomic information were used as specified in the framework of the single steps estimation method (Legarra et al., 2009; Christensen et al., 2010). The pedigree relationship matrix A^Γ (following the notation of Legarra et al., 2009) was built using the tabular rule method (Emik and Terrill, 1949), modified to account for a single metafounder (assuming a single base population for the Unkown Parent Group or UPG; Legarra et al., 2015), with a self-relationship of $\gamma = 8\sigma_p^2$, with σ_p^2 the variance of allele frequencies across markers (Garcia-Baccino et al., 2017). The combined relationship matrix H was summarized in Aguilar et al. (2019), and is:

$$H^{-1} = (A^\Gamma)^{-1} + (0,0;0; G^{-1} - (A_{22}^\Gamma)^{-1}) \qquad (1),$$

where A_{22}^Γ is the submatrix of A^Γ corresponding to the genotyped individuals, $G = (1-\alpha)(\omega_a + \omega_b \times G_V) + \alpha A_{22}^\Gamma$, with G_V the genomic relationship matrix estimated from the SNPs and scaled following the first scaling method of Van Raden (2008), α a scaling parameter (here equal to 0.05), and ω_a and ω_b chosen to equate the average inbreeding and the average relationships in G_V and A_{22}^Γ as in Christensen et al. (2012). As the sample size was relatively small, the matrix H^{-1} was obtained by simply inverting H with the function *solve* (R package *base*).

Population structure. In order to control for population structure and avoid spurious associations, we computed a distance matrix $1 - H^*$, where H^* is the correlation matrix obtained from H using the function *cov2cor* (R package *stats*), and the minus sign is to convert similarities to distances. We then used Multidimensional Scaling (MDS) on the distance matrix with the function *cmdscale_lanczos* (R package *refund* v0.1; Goldsmith et al., 2022; Miller, 2022) with the parameter 'k' set to 5 (hence explaining > 90% of the

variance), and 'eig' set to 'TRUE'. Negative eigenvalues were set to null, and the variance

explained by an axis was computed as the ratio between its squared eigenvalue and the

sum of the squared eigenvalues. Additionally, to assess how well the family structure was

captured, we measured the variable V_F defined as the average within family variance on the

first five axes (the lower the V_F, the better the family structure is captured). To make each

MDS derived from the different subsets comparable with each other, we normalized V_F by

the sum of squares of the first five eigenvalues.

Breeding value inferences. The spatially adjusted phenotypes were then used to infer the

breeding values with a multitrait single-step Genomic Best Linear Unbiased Predictor

(ssGBLUP), as follows:

$Y_{ijkl} = \mu + mds1_j + mds2_j + TRIAL_i + ANIMAL_j + e_{ijkl}$, (2)

where 'Y_{ijkl}' is the l-th phenotype of the k-th replicate of the j-th genotype in the i-th trial, μ is

the grand mean, '$mds1_j$' and '$mds2_j$' the fixed effects on the j-th genotype of the first and

second axes of the MDS, respectively, '$TRIAL_i$' the random effect of the i-th trial, '$ANIMAL_j$'

the polygenic effect of the j-th genotype, and 'e_{ijkl}' the residuals. Both the trial effect and the

residuals follow a Gaussian distribution with diagonal covariance matrices, and the polygenic

effect follows a Gaussian distribution with a covariance matrix equal to $k.H \times \Sigma$, where k is a

scaling parameter equal to 1 - γ/2 (Legarra et al., 2015), H the combined relationship matrix,

x the Kronecker product, and Σ the 2x2 covariance matrix between height and vulnerability

to rust. Estimates of coefficients and variance components of model (2) were obtained with

the function *remlf90* (R package *breedR* v0.12) through one run with the 'method' parameter

set to 'em' until convergence, then another run with the parameter 'method' set to 'ai' and the

parameter 'progsf90.option' set to 'maxrounds 1'. The output estimate of '$ANIMAL_j$' is the

Estimated Breeding Value (EBV) of the j-th genotype. The additive variance was obtained

from the diagonal elements of Σ, and used to estimate heritability (the denominator of which

also comprises the residual variance). The predictive ability (PA) of the model was defined

as Pearson's correlation coefficient between the breeding values re-estimated from the

effect sizes (see below) and the adjusted phenotypes corrected by the trial effect (i.e., $Y_{ijkl} -$

$TRIAL_i$).

Genome-Wide Association Study (GWAS). Following the equations of Aguilar et al.

(2019), the effect size of the SNPs was back-solved from the EBVs, along with their

standard error and their p-values. This process was performed trait by trait (i.e., without

accounting for the multitrait dimension) without loss of information, and taking all effects as

fixed, as the Pearson's correlation between the ssGBLUP's EBV and the breeding values re-

estimated by the effect sizes was greater than 0.999.

With the estimated effect sizes and the standard errors, we also computed an alternative to

the p-value accounting for multiple testing: the local false sign rate (lfsr) estimated with an

empirical Bayes approach for adaptive shrinkage using the function *ash* (R package *ashr*

v2.2; Stephens, 2017). In order to detect GWAS peaks (regions in the chromosomes with a

high density of significant SNPs), we used a custom peak detection script on the profile of

lfsr (Tiret & Milesi, 2021) with default parameters. The script filtered out significant hits that

are not statistically detected as a peak which is defined here as a statistically improbable

concentration of significant SNPs in a small chromosomic area.

Neighborhood. In order to account for the effect of the neighborhood, we estimated the

breeding values with an additional fixed effect, the local relatedness, defined as the average

relationship (from the matrix H) between a focal individual and its neighbors (eight or less if

on boundaries; as the king's moves on a chessboard). Local relatedness was orthogonal to

any micro-environmental effect, because of the random block design. The re-estimated

breeding values were denoted nEBV. To assess the goodness of fit of the different models,

we assessed the Akaike Information Criterion (AIC) with the function *AIC* of the R package

stats. To assess the extent to which local relatedness enriched phenotypic covariance, we

compared the phenotypic covariance between random pairs of neighboring individuals with

random pairs from the most related neighbors (highest quartile of local relatedness),

sampling for both calculation 10,000 pairs.

SNP subsets. We investigated different subsets of SNPs, each reflecting different aspects

of past selection or demography. We then computed (genomic) local relatedness with the

different subsets, so to assess which aspect of the past history is responsible for the

association between local relatedness and focal individuals' phenotype. We selected SNP

subsets based on Minor Allele Frequency (MAF), Ancestry Informativeness Coefficient (AIM;

Rosenberg et al., 2003), and for each trait, significance in the GWAS, and additive variance.

For each feature, we kept one-tenth of the whole set (713 SNPs, a sample size close to

previous studies on weighted GBLUP, e.g., Li et al., 2018) with either the smallest or the

largest values, ending up with 12 subsets. High MAF, low AIM, high significance in the

GWASs, and high additive variance were interpreted as reflecting past selection, and the

opposite as reflecting drift and neutral demographic dynamics. AIM was estimated per family

with the R script provided by Cappa et al. (2022). Additive variance explained by a SNP was

estimated as $2p(1-p)a^2$, where p is the MAF, and a the effect size (assuming no dominance

nor interaction deviation of the genetic variance; p.129, Falconer, 1989). We re-estimated

the combined relationship matrix for each SNP subset by re-estimating in the equation (1)

the genomic relationship matrix G_V with the SNP subset. With the re-estimated matrices H,

we re-performed the MDSs, and re-estimated the PAs of the EBVs and nEBVs (where local

relatedness were measured with the re-estimated H).

Statistical tests. For each SNP subset, we computed the confidence interval of the

Predictive Ability (PA) with a 1000 iteration bootstrap (re-sampling individuals), without re-

estimating the covariance matrix Σ. Here, the bootstrap was only used to assess the

variance, so that we centered the bootstrapped PA around the true PA. To assess the effect

of sampling, these subsets were compared to 1000 random samples of 713 SNPs (the size

of the SNP subsets). Comparisons were performed with a Student's one-sample t-test

(denoted t_1 with a degree of freedom or df of 999 corresponding to the number of bootstrap

iterations), or a Welch two-sample t-test (denoted t_2 with a varying df), both using the

function $t.test$ (R package $stats$). The sign of the t statistic is arbitrarily reported as positive

when greater than a baseline (e.g., the true PA), and negative when not. Models were

compared with a likelihood ratio test with a df of 1 using the function $lrtest$ (R package $lmtest$

v0.9; Zeileis & Hothorn, 2002). All scripts were written in R v4.1.3 (R Core Team, 2022).

Results

Tree height and vulnerability to rust both exhibited high heritabilities (0.469 and 0.666, respectively), high predictive abilities (0.677 and 0.755 respectively), and a strong independence, with an estimated genetic correlation of 0.022. We validated that GWAS peaks were likely not artefactual by swapping phenotypes (permutation test) among family members of the same block (i.e., only randomly mismatching Mendelian sampling): the permutation significantly reduced the average significance of the SNPs ($t_1 < -2.59 \times 10^2$, $p < 0.001$). Only one genetic variant was significantly associated with vulnerability to rust, located on chromosome 1, and none for tree height (Fig. 1A, 1B). The Q-Q plot on vulnerability to rust revealed that some SNPs were significantly associated with a level of significance well above the 95% confidence interval (Fig. 1D); most of them being in the identified peak. None of the SNPs were significantly associated with tree height after Bayesian shrinkage, and Q-Q plot revealed a classic population structure overcorrection (Fig. 1C).

Local relatedness improves ssGBLUP fit for vulnerability to rust

Although the growth of focal individuals was not affected by their neighbors, damage due to rust was more severe the more related the focal individuals were with their neighbors. Indeed, the genomic models incorporating local relatedness as a covariate had significantly higher likelihood than without for vulnerability to rust ($\chi^2 = 9.83$, $p = 0.002$), but not for tree height ($\chi^2 = 0.09$, $p = 0.236$). When computing relatedness with the different subsets of SNPs, the Akaike Information Criterion (AIC) decreased by 0.12 for height (when excluding the smallest MAF) and by 7.95 for vulnerability to rust (Fig. S2), suggesting a better fit when including local relatedness in the model for vulnerability to rust. However, for vulnerability to rust, only the subset of SNPs with the largest MAF significantly increased the likelihood of the model ($\chi^2 = 4.02$, $p = 0.04$). Likelihood was not significantly improved by SNPs with the largest AIM ($\chi^2 = 0.15$, $p = 0.703$) or the largest additive variance ($\chi^2 = 2.65$, $p = 0.104$,

respectively), and was significantly worse with other subsets ($\chi^2 > 6.66$, $p < 0.010$). For tree

height, no subset of SNPs improved the likelihood ($\chi^2 < 3.48$, $p > 0.062$). In other words,

focal individuals were more affected by rust when they had related individuals in their

vicinity. The covariance between random pairs of neighboring individuals' vulnerability to rust

compared to the most related neighbors' pairs increased from an average covariance of 0.28

to 0.43, showcasing a systematic association despite the randomized block design.

In spite of genomic models being sensitive to local relatedness, improvement of predictive

abilities (PA) was limited (Fig. S3; see Supplementary materials for further details) - probably

due to a lack of power. Indeed, incorporating local relatedness did not significantly change

the PA neither with the whole set of SNPs ($|t_2| < 0.594$, $p > 0.553$), nor with any of the

alternative subsets ($|t_2| < 1.03$, $p > 0.301$), with a slightly better, but non-significant PA when

incorporating local relatedness for SNPs with the smallest MAF on vulnerability to rust ($t_2 =$

1.73, df = 2.00×10^3, $p = 0.0841$), and SNPs with the smallest tree height significance on

vulnerability to rust ($t_2 = 1.97$, df = 2.00×10^3, $p = 0.0491$).

The impact of local relatedness on vulnerability to rust is <u>likely</u> due to

past family selection

The objective here is to assess which SNP subset reveals which aspect of the past history

(selection or drift), knowing that local relatedness is mainly associated to vulnerability to rust

through SNPs with large MAF. The population under study is structured in a hierarchical

way: first into populations, then into families. Population structure, captured by the first

eigenvalues (Patterson et al., 2006), substantially shaped genetic diversity, as the first two

eigenvalues alone explained more than 50% of the variance (40.0% and 24.5% respectively;

Fig. S1). Because the population underwent family selection during breeding in the past,

family structure is strong and can even be apprehended visually (Fig. 2). Several statistics

significantly supports the strong family structure: (i) randomly shuffling the family ID

increased by 9.96 times the within "family" variance ($t_1 = 3.70 \times 10^3$, p < 0.001); (ii) when

restricting to common alleles, the first two eigenvalues of the PCA still explained a large part

of the variance (56.2%), showing that family structure is bound to selection, as expected

under family selection.

In the case of the investigated populations, family structure can reveal neutral past

dynamics, or past family selection. In order to understand what drove the family structure,

similarly to using annotation to focus on some SNPs, we subset 10% of the SNPs first

randomly and then according to their MAF, AIM, significance in a GWAS, or additive

variance, thus redefining relatedness with a different set of linkage disequilibrium. In our

dataset, random samples of SNPs, performed quite poorly compared to the whole set of

SNPs ($t_1 = 2.12 \times 10^2$, p < 0.001). However, SNPs with the largest AIM, performing better

than random samples ($t_1 = 3.86 \times 10^2$, p < 0.001; Fig. 3), and even than the whole set; and

SNPs with the largest MAF and the largest additive variance, performing better than random

samples ($t_1 > 1.09 \times 10^2$, p < 0.001). The improvement in goodness of fit due to incorporating

local relatedness in the genomic models is thus due to any forces structuring the family,

likely past family selection since it increases the MAF.

Discussion

The success of prediction based on genomic models relies on the ability of genotypic data to
capture the covariance between candidate loci, thereby reflecting how much genetic
similarity explains phenotypic similarities (Powell et al., 2010). An additional source of
phenotypic covariances are micro-environmental effects, and accounting for spatial
autocorrelation can enhance genomic predictions (Cappa et al, 2022). In our case, on top of
spatial autocorrelation, we accounted for local relatedness, another source of heterogeneity.
The significant increase of the likelihood when adding local relatedness as a covariate
shows that part of the plant-to-plant interaction can be ascribed to the relatedness with
neighbors, at least for vulnerability to rust.

Different genetic architectures and local relatedness

Height is known to have a highly polygenic genetic architecture (Du et al., 2016), and
resistance to rust to have a mono- or oligogenic architecture (e.g., Jorge et al., 2005),
implying that relatedness – as the amount of shared alleles – may have different
consequences on those two traits: genetic redundancy (Nowak et al., 1997; Láruson et al.,
2020) of height probably making it less sensitive to local relatedness than resistance to rust.
The low genetic correlation between both traits is consistent with some previous reports
(e.g., Beaulieu et al., 2020; Liu et al., 2022), though not all (e.g., Lenz et al., 2021; Cappa et
al., 2022); we will assume in our case that orthogonality is strong enough to show that the
traits are non-redundant, with independent architectures.

The differences in genetic architecture between the two traits studied, as well as the design
of the SNP array, may have determined the results in terms of association, with only one
major gene detected (as a peak) for rust, in contrast to the absence of associations for the
more polygenic trait. Indeed, a major caveat of our analysis is that the SNP chip array we
used is enriched for QTLs and candidate expression genes related to resistance to rust

during its development (Faivre-Rampant et al., 2016), but not for height. While there may be

a risk of ascertainment bias due to the frequency criteria used and the design of the array,

this array was constructed using a very large diversity panel that is representative of the

western distribution of the species. Several studies have compared different technologies

(e.g., SNP arrays, genotyping-by-sequencing, etc.) and found that ascertainment bias was

negligible for monitoring diversity trends and estimating relatedness (Torkamaneh et al.,

2015; Negro et al., 2019). Although biases may exist, the effects are unlikely to be large

enough to invalidate the results. However, further investigation with alternative sequencing

technologies would be needed to validate this conclusion. Another element that may have

influenced the association for rust is its phenotypic scoring, which is based on a qualitative

scale of nine grades, which might artificially grade an inoculum pressure linearly. Overall,

even if the results should be taken cautiously, it is not uncommon to encounter a simpler

genetic architecture for response to biotic constraints compared to complex quantitative

traits (e.g., gene-for-gene resistance; Flor, 1971; Jorge et al., 2005). Consequently, many

haplotype combinations can lead to tall individuals, but few to resistance to rust, because a

polygenic architecture is redundant; in other words, when individuals are related, their

probability of both being resistant is higher than that of both being tall (individuals sharing

rare alleles are closer than when sharing common alleles; Van Raden, 2008). This reasoning

is specifically adapted in the case of factorial crossing where all possible crosses have been

performed numerous times. Therefore, local relatedness would increase with a vulnerability

to rust that is shared within the neighborhood, even if it is not inducing vulnerability to rust

per se.

Is local relatedness the cause of local phenotypic similarities?

Interactions between neighbors may depend on local relatedness, and whether this

interaction is positive or negative depends on the relative strength of kin selection (Hamilton,

1964) and niche partitioning (Silvertown, 2004). As mentioned above, the importance of

group selection in nature is still debated, but local relatedness has been shown to be

perceived through kin recognition in plant species (e.g., Dudley et al., 2007; Murphy et al.,

2009). Such recognition could then lead to favorable (kin selection) or unfavorable (niche

partitioning) interactions. One such detrimental interaction is competition, which can be

accounted for in a quantitative genetic model in the DGE/IGE framework (Griffing, 1967;

potential applications in Muir, 2005). Competition occurs when DGE and IGE are negatively

correlated. A positive correlation from such a model would suggest a cooperative synergistic

interaction. Although adding complexity to the model, including competition has been shown

to reduce bias in covariance estimation (Costa e Silva et al., 2013). In our case, in addition

to attempts to include interactions, IGE resulted in a lower predictive ability than with the

ssGBLUP baseline (results not shown).

Once it has been shown that neighbourhood relatedness is indeed correlated with similarity

at focal phenotypes, the difficulty in disentangling kin selection from niche partitioning is to

determine whether the relatedness is a cause, as an expression of kin selection or niche

competition, or a co-product, as an expression of a confounding between relatedness and

interaction through phenotypes, where the focal phenotype would be modified by that of its

neighbors through plant-to-plant interactions. If phenotypic interaction increases when

phenotypes are similar (e.g., competition for resource uptake), it can be confused with the

effect of relatedness, as genetically related individuals tend to have similar phenotypes.

However, if local relatedness was only a proxy for highly related resistant to rust

neighborhoods, average vulnerability to rust would increase with local relatedness but not

covariance of vulnerability to rust (i.e., a systematic and coherent increase in vulnerability to

rust), because of the randomized block design.

The results hence suggest that shared vulnerability is likely the cause of rust damage

amplification in the related neighborhood. The amplifying effect might be even stronger in

perennials as shared vulnerability can be perceived by successive generations of

pathogens. This phenomenon is consistent with the increased disease resilience conferred

by varietal association in a field (e.g., Smithson and Lenne, 1996; Burdon and Thrall, 2009),

or the detrimental effect of lack of diversity in several plant species (Zhu et al., 2000; Liu et

al., 2020), but would need further investigations. On the other hand, the fact that family

structure was mainly resulting from allele identity sharing at SNPs likely associated with

selection (following the hypothesis of Yang et al., 2010 and Biddanda et al., 2020) suggest

that relatives were co-selected (relatedness as a confounding effect), as is often the case in

breeding programs implementing family selection. An inevitable consequence is that, as

seen in crop selection, interesting genotypes for kin selection/cooperation have been lost

during domestication (Fréville et al., 2022), making it difficult for genomic selection to

account for local relatedness (as predictive abilities show).

Conclusion

The effect of the relatedness of a focal individual with its neighbors was investigated on tree

height and vulnerability to rust, with 10,301 trees of 1,452 genotypes across four trials. The

results show that incorporating local relatedness as an additional factor in GBLUP models

has a significantly greater influence on estimation of resistance to rust than on tree height,

though its overall effect on genomic predictions was limited. Whether there was kin

selection/competition in addition to the phenotypic interaction, or whether we simply lacked

statistical power, is difficult to assess. Selection on local relatedness due to kin selection or

niche partitioning - selection on relatedness (favoring specific plant-to-plant interaction due

to relatedness) - might be confused with the fact that selection leads to phenotypic

convergence, that in turn leads to a genotypic convergence - a co-selection of related

(selecting for phenotypes that accidentally but inevitably selected for related individuals).

Further investigation is required to elucidate the relationship between genetic architecture

and the impact of relatedness on phenotypes.

Acknowledgements

The authors acknowledge the Uppsala University and the European Union's Horizon 2020 B4EST for basic functioning and postdoctoral grant for MT. The authors would like to thank Véronique Jorge and Remy Gobin of the BioForA unit (INRAE, ONF, Orléans, France) for their help in compiling the data. The computations and data handling were enabled by resources provided by the Swedish National Infrastructure for Computing (SNIC 2017-7-296) at UppMax partially funded by the Swedish Research Council through grant agreement no. 2018-05973.

Author Contributions

MT was responsible for writing the report, conducting the search, extracting and analysing data, interpreting results, updating reference lists. LS was responsible for designing the experiments and screening potentially eligible studies. ML and LS contributed to interpreting results and provided feedback on the report.

Competing Interests

The authors declare that there is no conflict of interest.

Data Archiving

The data that support the findings of this study are openly available on DATA INRAE at https://data.inrae.fr/privateurl.xhtml?token=b79ab1ca-ebb9-47c6-9272-1568c0c33d70.

Rerefences

Aguilar, I., Legarra, A., Cardoso, F., Masuda, Y., Lourenco, D., & Misztal, I. (2019). Frequentist p-values for large-scale-single step genome-wide association, with an application to birth weight in American Angus cattle. *Genetics Selection Evolution*, *51*(1), 1-8.

Astle, W., & Balding, D. J. (2009). Population structure and cryptic relatedness in genetic association studies. *Statistical Science*, *24*(4), 451-471.

Bailey, N. W., Marie-Orleach, L., & Moore, A. J. (2018). Indirect genetic effects in behavioral ecology: does behavior play a special role in evolution?. *Behavioral Ecology*, *29*(1), 1-11.

Beaulieu, J., Doerksen, T., Clément, S., MacKay, J., & Bousquet, J. (2014). Accuracy of genomic selection models in a large population of open-pollinated families in white spruce. *Heredity*, *113*(4), 343-352.

Beaulieu, J., Nadeau, S., Ding, C., Celedon, J. M., Azaiez, A., Ritland, C., et al. (2020). Genomic selection for resistance to spruce budworm in white spruce and relationships with growth and wood quality traits. *Evolutionary applications*, *13*(10), 2704-2722.

Bengtsson, J., Bullock, J. M., Egoh, B., Everson, C., Everson, T., O'connor, T., ... & Lindborg, R. (2019). Grasslands—more important for ecosystem services than you might think. *Ecosphere*, *10*(2), e02582.

Biddanda, A., Rice, D. P., & Novembre, J. (2020). A variant-centric perspective on geographic patterns of human allele frequency variation. *Elife*, *9*, e60107.

Biernaskie, J. M. (2022). Kin selection theory and the design of cooperative crops.

Evolutionary Applications, *15*(10), 1555-1564.

Bijma, P., & Wade, M. J. (2008). The joint effects of kin, multilevel selection and indirect

genetic effects on response to genetic selection. *Journal of evolutionary biology*, *21*(5),

1175-1188.

Bourke, P. M., Evers, J. B., Bijma, P., van Apeldoorn, D. F., Smulders, M. J., Kuyper, T.

W., ... & Bonnema, G. (2021). Breeding beyond monoculture: Putting the "intercrop" into

crops. *Frontiers in Plant Science*, *12*, 734167.

Burdon, J. J., & Thrall, P. H. (1999). Spatial and temporal patterns in coevolving plant and

pathogen associations. *The American Naturalist*, *153*(S5), S15-S33.

Burdon, J. J., & Thrall, P. H. (2009). Coevolution of plants and their pathogens in natural

habitats. *Science*, *324*(5928), 755-756.

Boyle, E. A., Li, Y. I., & Pritchard, J. K. (2017). An expanded view of complex traits: from

polygenic to omnigenic. *Cell*, *169*(7), 1177-1186.

Cahill Jr, J. F., & McNickle, G. G. (2011). The behavioral ecology of nutrient foraging by

plants. *Annual Review of ecology, evolution, and systematics*, *42*, 289-311.

Cappa, E. P., & Cantet, R. J. (2007). Bayesian estimation of a surface to account for a

spatial trend using penalized splines in an individual-tree mixed model. *Canadian Journal of*

Forest Research, *37*(12), 2677-2688.

Cappa, E. P., de Lima, B. M., da Silva-Junior, O. B., Garcia, C. C., Mansfield, S. D., &

Grattapaglia, D. (2019). Improving genomic prediction of growth and wood traits in

Eucalyptus using phenotypes from non-genotyped trees by single-step GBLUP. *Plant*

Science, *284*, 9-15.

Cappa, E. P., Ratcliffe, B., Chen, C., Thomas, B. R., Liu, Y., Klutsch, J., et al. (2022).

Improving lodgepole pine genomic evaluation using spatial correlation structure and SNP

selection with single-step GBLUP. *Heredity*, *128*(4), 209-224.

Casper, B. B., & Jackson, R. B. (1997). Plant competition underground. *Annual review of*

ecology and systematics, 545-570.

Christensen, O. F., & Lund, M. S. 2010. Genomic prediction when some animals are

not genotyped. Genetics Selection Evolution 42, 2.

Christensen, O. F., Madsen, P., Nielsen, B., Ostersen, T., & Su, G. (2012). Single-step

methods for genomic evaluation in pigs. *animal*, *6*(10), 1565-1571.

Costa e Silva, J., Kerr, R.J. (2013) Accounting for competition in genetic analysis, with

particular emphasis on forest genetic trials. *Tree Genetics & Genomes, 9,* 1-17.

Dawkins, R. (1976). *The selfish gene*. Oxford: Oxford University Press.

Denison, R. F., Kiers, E. T., & West, S. A. (2003). Darwinian agriculture: when can humans

find solutions beyond the reach of natural selection? *The quarterly review of biology*, *78*(2),

145-168.

Diouf, I., Derivot, L., Koussevitzky, S., Carretero, Y., Bitton, F., Moreau, L., & Causse, M.

(2020). Genetic basis of phenotypic plasticity and genotype× environment interactions in a

multi-parental tomato population. Journal of Experimental Botany, 71(18), 5365-5376

Donald, C.M. (1981). Competitive plants, communal plants, and yields in wheat crops.

L. Evans, W.J. Peacock (Eds.), Wheat Science – Today and Tomorrow, Cambridge

University Press.

Donald, C.M., Hamblin, J. (1983). The convergent evolution of annual seed crops in

agriculture. In Advances in agronomy (ed. NC Brady), pp. 97–143. New York, NY:

Academic Press.

Dong, L., Xiao, S., Chen, J., Wan, L., & Wang, Z. (2016). Genomic selection using extreme

phenotypes and pre-selection of SNPs in large yellow croaker (Larimichthys crocea). *Marine*

Biotechnology, *18*(5), 575-583.

Du, Q., Gong, C., Wang, Q., Zhou, D., Yang, H., Pan, W., et al. (2016). Genetic architecture

of growth traits in Populus revealed by integrated quantitative trait locus (QTL) analysis and

association studies. *New Phytologist*, *209*(3), 1067-1082.

Ehlers, B. K., & Bilde, T. (2019). Inclusive fitness, asymmetric competition and kin selection

in plants. *Oikos*, *128*(6), 765-774.

Ellen, E. D., Peeters, K., Verhoeven, M., Gols, R., Harvey, J. A., Wade, M. J., ... & Bijma, P.

(2016). Direct and indirect genetic effects in life-history traits of flour beetles (Tribolium

castaneum). *Evolution*, *70*(1), 207-217.

Fisher, R.A. (1918). The correlation between relatives under the supposition of Mendelian

inheritance. *Trans. Roy. Soc. Edinburgh 52*, 399-433.

Gualdrón Duarte, J. L., Gori, A. S., Hubin, X., Lourenco, D., Charlier, C., Misztal, I., & Druet,

T. (2020). Performances of Adaptive MultiBLUP, Bayesian regressions, and weighted-

GBLUP approaches for genomic predictions in Belgian Blue beef cattle. *BMC genomics*,

21(1), 1-18.

Dudley, S. A., & File, A. L. (2007). Kin recognition in an annual plant. *Biology Letters*, *3*(4),

435-438.

Emik, L. O., & Terrill, C. E. (1949). Systematic procedures for calculating inbreeding

coefficients. *Journal of Heredity*, *40*(2), 51-55.

Faivre-Rampant, P., Zaina, G., Jorge, V., Giacomello, S., Segura, V., Scalabrin, S., et al.

(2016). New resources for genetic studies in Populus nigra: Genome-wide SNP discovery

and development of a 12k Infinium array. *Molecular ecology resources*, *16*(4), 1023-1036.

Falconer, D.S. (1989). Introduction to quantitative genetics (3rd edition, p. 129). Pearson

Education India.

Feau, N., Joly, D. L., & Hamelin, R. C. (2007). Poplar leaf rusts: model pathogens for a

model tree. *Botany*, *85*(12), 1127-1135.

File, A. L., Murphy, G. P., & Dudley, S. A. (2012). Fitness consequences of plants growing

with siblings: reconciling kin selection, niche partitioning and competitive ability. *Proceedings*

of the Royal Society B: Biological Sciences, *279*(1727), 209-218.

Flor, H. H. (1971). Current status of the gene-for-gene concept. *Annual review of*

phytopathology, *9*(1), 275-296.

Fréville, H., Montazeaud, G., Forst, E., David, J., Papa, R., & Tenaillon, M. I. (2022). Shift in

beneficial interactions during crop evolution. *Evolutionary Applications*, *15*(6), 905-918.

Garcia-Baccino, C. A., Legarra, A., Christensen, O. F., Misztal, I., Pocrnic, I., Vitezica, Z. G.,

& Cantet, R. J. (2017). Metafounders are related to F st fixation indices and reduce bias in

single-step genomic evaluations. *Genetics Selection Evolution*, *49*(1), 1-14.

Goldsmith, J., Scheipl, F., Huang, L., Wrobel, J., Di, C., Gellar, J., et al. (2022). refund:

Regression with Functional Data. R package version 0.1-26.

https://CRAN.R-project.org/package=refund

Grattapaglia, D., & Resende, M. D. (2011). Genomic selection in forest tree breeding. *Tree*

Genetics & Genomes, *7*, 241-255.

Griffing, B. (1967). Selection in reference to biological groups I. Individual and group

selection applied to populations of unordered groups. *Australian Journal of Biological*

Sciences, *20*(1), 127-140.

Guisan, A., & Thuiller, W. (2005). Predicting species distribution: offering more than simple

habitat models. *Ecology letters*, *8*(9), 993-1009.

Hamilton, W. D. (1964). The genetical evolution of social behaviour. II. *Journal of theoretical*

biology, *7*(1), 17-52.

Isik, F., Bartholomé, J., Farjat, A., Chancerel, E., Raffin, A., Sanchez, L., et al. (2016).

Genomic selection in maritime pine. *Plant Science*, *242*, 108-119.

Jorge, V., Dowkiw, A., Faivre-Rampant, P., & Bastien, C. (2005). Genetic architecture of

qualitative and quantitative *Melampsora larici-populina* leaf rust resistance in hybrid poplar:

genetic mapping and QTL detection. *New Phytologist*, *167*(1), 113-127.

Kijas, J., Elliot, N., Kube, P., Evans, B., Botwright, N., King, H., et al. (2017). Diversity and

linkage disequilibrium in farmed Tasmanian Atlantic salmon. *Animal genetics*, *48*(2), 237-

241.

Láruson, Á. J., Yeaman, S., & Lotterhos, K. E. (2020). The importance of genetic

redundancy in evolution. *Trends in ecology & evolution*, *35*(9), 809-822.

Legarra, A., Aguilar, I., & Misztal, I. (2009). A relationship matrix including full pedigree and

genomic information. *Journal of dairy science*, *92*(9), 4656-4663.

Legarra, A., Christensen, O. F., Vitezica, Z. G., Aguilar, I., & Misztal, I. (2015). Ancestral

relationships using metafounders: finite ancestral populations and across population

relationships. *Genetics*, *200*(2), 455-468.

Legionnet, A., Muranty, H., & Lefèvre, F. (1999). Genetic variation of the riparian pioneer

tree species Populus nigra. II. Variation in susceptibility to the foliar rust Melampsora larici-

populina. *Heredity*, *82*(3), 318-327.

Leigh Jr, E. G. (2010). The group selection controversy. *Journal of evolutionary biology*,

23(1), 6-19.

Lemoine, T., Violle, C., Montazeaud, G., Isaac, M. E., Rocher, A., Fréville, H., & Fort, F.

(2023). Plant trait relationships are maintained within a major crop species: lack of artificial

selection signal and potential for improved agronomic performance. *New Phytologist*, *240*(6),

2227-2238.

Lenz, P. R., Nadeau, S., Mottet, M. J., Perron, M., Isabel, N., Beaulieu, J., & Bousquet, J.

(2020). Multi-trait genomic selection for weevil resistance, growth, and wood quality in

Norway spruce. *Evolutionary applications*, *13*(1), 76-94.

Lesaffre, T., & Billiard, S. (2021). On Deleterious Mutations in Perennials: Inbreeding

Depression, Mutation Load, and Life-History Evolution. *The American Naturalist*, *197*(5),

E143-E155.

Lewontin, R. C. (1970). The units of selection. *Annual review of ecology and systematics*, 1-

18.

Li, B., Zhang, N., Wang, Y. G., George, A. W., Reverter, A., & Li, Y. (2018). Genomic

prediction of breeding values using a subset of SNPs identified by three machine learning

methods. *Frontiers in genetics*, *9*, 237.

Liu, X., Chen, L., Liu, M., García-Guzmán, G., Gilbert, G. S., & Zhou, S. (2020). Dilution

effect of plant diversity on infectious diseases: latitudinal trend and biological context

dependence. Oikos, 129(4), 457-465.

Liu, Y., Erbilgin, N., Ratcliffe, B., Klutsch, J. G., Wei, X., Ullah, A., et al. (2022). Pest

defences under weak selection exert a limited influence on the evolution of height growth

and drought avoidance in marginal pine populations. *Proceedings of the Royal Society B*,

289(1982), 20221034.

Lu, S., Liu, Y., Yu, X., Li, Y., Yang, Y., Wei, M., et al. (2020). Prediction of genomic breeding

values based on pre-selected SNPs using ssGBLUP, WssGBLUP and BayesB for

Edwardsiellosis resistance in Japanese flounder. *Genetics Selection Evolution*, *52*(1), 1-10.

Luo, Z., Yu, Y., Xiang, J., & Li, F. (2021). Genomic selection using a subset of SNPs

identified by genome-wide association analysis for disease resistance traits in aquaculture

species. *Aquaculture*, *539*, 736620.

Maynard Smith, J. (1964). Group selection and kin selection. *Nature*, *201*(4924), 1145-1147.

Meuwissen, T. H. E. (1997). Maximizing the response of selection with a predefined rate of

inbreeding. *Journal of animal science*, *75*(4), 934-940.

Meuwissen, T. H., Hayes, B. J., & Goddard, M. (2001). Prediction of total genetic value using

genome-wide dense marker maps. *genetics*, *157*(4), 1819-1829.

Milbau, A., Reheul, D., De Cauwer, B., & Nijs, I. (2007). Factors determining plant–

neighbour interactions on different spatial scales in young species-rich grassland

communities. *Ecological research*, *22*(2), 242-247.

Miller, D. L. (2022). porridge: Principal Co-Ordinate Ridge Regression. R package version

1.2.

Montazeaud, G., Rousset, F., Fort, F., Violle, C., Fréville, H., & Gandon, S. (2020). Farming

plant cooperation in crops. *Proceedings of the Royal Society B*, *287*(1919), 20191290.

Muir, W. M. (2005). Incorporation of competitive effects in forest tree or animal breeding

programs. *Genetics*, *170*(3), 1247-1259.

Muñoz, F., and Sanchez, L. (2020). breedR: Statistical Methods for Forest Genetic

Resources Analysts. R package version 0.12-5. https://github.com/famuvie/breedR

Murphy, G. P., & Dudley, S. A. (2009). Kin recognition: competition and cooperation in

Impatiens (Balsaminaceae). *American journal of botany*, *96*(11), 1990-1996.

Murphy, G. P., Swanton, C. J., Van Acker, R. C., & Dudley, S. A. (2017). Kin recognition,

multilevel selection and altruism in crop sustainability. *Journal of Ecology (Oxford)*, *105*(4),

930-934.

Negro, S. S., Millet, E. J., Madur, D., Bauland, C., Combes, V., Welcker, C., ... & Nicolas, S.

D. (2019). Genotyping-by-sequencing and SNP-arrays are complementary for detecting

quantitative trait loci by tagging different haplotypes in association studies. BMC plant

biology, 19, 1-22.

Nowak, M. A., Boerlijst, M. C., Cooke, J., & Smith, J. M. (1997). Evolution of genetic

redundancy. *Nature*, *388*(6638), 167-171.

Okasha, S. (2020). Altruism, group selection and correlated interaction. *The British journal*

for the philosophy of science.

Patterson, N., Price, A. L., & Reich, D. (2006). Population structure and eigenanalysis. *PLoS*

genetics, *2*(12), e190.

Pégard, M., Rogier, O., Bérard, A., Faivre-Rampant, P., Paslier, M. C. L., Bastien, C., et al.

(2019). Sequence imputation from low density single nucleotide polymorphism panel in a

black poplar breeding population. *BMC genomics*, *20*(1), 1-16.

Pégard, M., Segura, V., Muñoz, F., Bastien, C., Jorge, V., & Sanchez, L. (2020). Favorable

conditions for genomic evaluation to outperform classical pedigree evaluation highlighted by

a proof-of-concept study in poplar. *Frontiers in plant science*, *11*, 581954.

Pinon, J., & Frey, P. (1997). Structure of Melampsora larici-populina populations on wild and

cultivated poplar. *European Journal of Plant Pathology*, *103*, 159-173.

Powell, J. E., Visscher, P. M., & Goddard, M. E. (2010). Reconciling the analysis of IBD and

IBS in complex trait studies. *Nature Reviews Genetics*, *11*(11), 800-805.

R Core Team (2022). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL. https://www.R-project.org/

Ratcliffe, B., El-Dien, O. G., Cappa, E. P., Porth, I., Klápště, J., Chen, C., & El-Kassaby, Y.

A. (2017). Single-step BLUP with varying genotyping effort in open-pollinated Picea glauca.

G3: Genes, Genomes, Genetics, *7*(3), 935-942.

Resende, M. D., Resende Jr, M. F., Sansaloni, C. P., Petroli, C. D., Missiaggia, A. A.,

Aguiar, A. M., et al. (2012). Genomic selection for growth and wood quality in Eucalyptus:

capturing the missing heritability and accelerating breeding for complex traits in forest trees.

New Phytologist, *194*(1), 116-128.

Rosenberg, N. A., Li, L. M., Ward, R., & Pritchard, J. K. (2003). Informativeness of genetic

markers for inference of ancestry. *The American Journal of Human Genetics*, *73*(6), 1402-

1422.

Sanchez, M. P., Govignon-Gion, A., Croiseau, P., Fritz, S., Hozé, C., Miranda, G., et al.

(2017). Within-breed and multi-breed GWAS on imputed whole-genome sequence variants

reveal candidate mutations affecting milk protein composition in dairy cattle. *Genetics*

Selection Evolution, *49*(1), 1-16.

Silvertown, J. (2004). Plant coexistence and the niche. *Trends in Ecology & evolution*,

19(11), 605-611.

Smithson, J. B., & Lenne, J. M. (1996). Varietal mixtures: a viable strategy for sustainable

productivity in subsistence agriculture. *Annals of applied biology*, *128*(1), 127-158.

Stephens, M. (2017). False discovery rates: a new deal. *Biostatistics*, *18*(2), 275-294.

Thistlethwaite, F. R., Gamal El-Dien, O., Ratcliffe, B., Klápště, J., Porth, I., Chen, C., et al.

(2020). Linkage disequilibrium vs. pedigree: Genomic selection prediction accuracy in

conifer species. *PLoS One*, *15*(6), e0232201.

Tiret, M., & Milesi, P. Statistical Peak Detection for GWAS. Zenodo.

https://doi.org/10.5281/zenodo.5079556. Deposited 7 July 2021.

Torkamaneh, D., & Belzile, F. (2015). Scanning and filling: ultra-dense SNP genotyping

combining genotyping-by-sequencing, SNP array and whole-genome resequencing data.

PloS one, 10(7), e0131533.

Tsai, H. Y., Matika, O., Edwards, S. M., Antolín–Sánchez, R., Hamilton, A., Guy, D. R., et al.

(2017). Genotype imputation to improve the cost-efficiency of genomic selection in farmed

Atlantic salmon. *G3: Genes, genomes, genetics*, *7*(4), 1377-1383.

Van Raden, P. M. (2008). Efficient methods to compute genomic predictions. *Journal of*

dairy science, *91*(11), 4414-4423.

Weiner, J., Andersen, S. B., Wille, W. K. M., Griepentrog, H. W., & Olsen, J. M. (2010).

Evolutionary Agroecology: the potential for cooperative, high density, weed-suppressing

cereals. *Evolutionary Applications*, *3*(5-6), 473-479.

Weiner, J., Du, Y. L., Zhang, C., Qin, X. L., & Li, F. M. (2017). Evolutionary agroecology:

individual fitness and population yield in wheat (Triticum aestivum).

Weiner, J. (2019). Looking in the wrong direction for higher-yielding crop genotypes. *Trends*

in Plant Science, *24*(10), 927-933.

Wilson, D. S., & Wilson, E. O. (2007). Rethinking the theoretical foundation of sociobiology.

The Quarterly review of biology, *82*(4), 327-348.

Yang, J., Benyamin, B., McEvoy, B. P., Gordon, S., Henders, A. K., Nyholt, D. R., et al.

(2010). Common SNPs explain a large proportion of the heritability for human height. *Nature*

genetics, *42*(7), 565-569.

Zeileis, A., & Hothorn, T. (2002). Diagnostic Checking in Regression Relationships. R News

2(3), 7-10.

Zhang, Z., Liu, J., Ding, X., Bijma, P., de Koning, D. J., & Zhang, Q. (2010). Best linear

unbiased prediction of genomic breeding values using a trait-specific marker-derived

relationship matrix. *PloS one*, *5*(9), e12648.

Zhu, Y., Chen, H., Fan, J., Wang, Y., Li, Y., Chen, J., ... & Mundt, C. C. (2000). Genetic

diversity and disease control in rice. Nature, 406(6797), 718-722.

Figures

Figure 1. GWAS on tree height and vulnerability to rust in Populus nigra. **(A)**. Manhattan plot of the GWAS on tree height. The dots are the log significance of the markers. The solid horizontal line stands for the 5% significance level, and the dashed horizontal line stands for the 5% FDR level. The vertical dashed lines are position where significant hits were detected as being part of a peak. **(B)**. Idem as panel A, for vulnerability to rust. **(C)**. Q-Q plot of the GWAS on tree height. The dots were the quantile of the markers; the solid line is the 1-slope segment; the gray cross is the position of the median; and the gray area is the 95% confidence interval. **(D)**. Idem as panel C, for vulnerability to rust.

Figure 2. The first two axes of the Multidimensional Scaling (MDS) of Populus nigra's genotyping data. **(A)**. MDS with labels of the fathers (with black large shapes). Colored and smaller shapes are the offspring, the shape corresponding to its father. **(B)**. The same MDS as in panel A., but labelled for the mothers.

Figure 3. Within family variance (V_F) for each subset of SNPs. The black dashed horizontal line is the average V_F of randomly sampled SNPs, the gray dashed horizontal lines delimit the 95% confidence interval, and the black horizontal dotted line is V_F of the entire set of SNPs. Red dots are when selected for tree height, and blue for vulnerability to rust (green dots are when this distinction is not relevant). SNP subsets: largest or smallest MAF (+ or - MAF), largest or smallest AIM (+ or - AIM), most or least significant (+ or - $\log_{10}(p)$), and most or least heritable for tree height (+ or - h^2).