A Sex specific homologue of snake Waprin is essential for Embryonic Development in
the Red Flour Beetle, *Tribolium castaneum*

Running Title: Egg development in *Tribolium* requires Waprin

Chhavi Choudhary, Divyanshu Kishore, Keshav Kumar Meghwanshi, Vivek Verma[#], Jayendra Nath
Shukla*

Department of Biotechnology, School of Life Sciences, Central University of Rajasthan, Ajmer, 305817
Rajasthan, India.

#Present Address: Gujarat Biotechnology University, Gandhinagar, Gujarat, India.

*Corresponding author, email ID: jayendrashukla@curaj.ac.in

Keywords: Waprin, *Tribolium castaneum*, Sex-specific, Parental-RNAi, Embryonic
development.

Summary statement:

Tribolium castaneum possesses a female-specific homologue of snake Waprin (Tc_Wap^F)
which is essential for its embryonic development. Expression of *Tc_Wap^F* is independent of
Tribolium sex-determination cascade.

1. Abstract:

Waprin, an extracellular secretory protein, is widely known for its antibacterial activity. Through the analysis of sex-specific transcriptomes of adult *Tribolium castaneum*, we identified a homologue of snake *waprin* (Tc_Wap^F) with significantly higher level of its transcripts in females. Developmental- and tissue-specific profiling revealed the widespread expression of Tc_Wap^F in adult female tissues. Interestingly, Tc_Wap^F expression is not regulated by the classical sex-determination cascade of *T. castaneum*, as we fail to get any attenuation in Tc_Wap^F transcript levels in *Tcdsx* and *Tctra* (key players of sex determination cascade of *T. castaneum*) knockdown females. The eggs laid by Tc_Wap^F knockdown females never hatched, indicating the involvement of Tc_Wap^F in the embryonic development in *T. castaneum*. This is the first report of identification a sex-specific *waprin* homologue in an insect and its involvement as a novel candidate in the embryonic development. Future investigations on the functional conservation of insect *waprins* and their mechanistic role in embryonic development can be exploited for improving pest management strategies.

2. Introduction:

Waprin, a WAP (Whey Acidic Protein)-domain containing secretory protein was first identified in the venom of Black-necked spitting cobra, *Naja nigricollis*, and was named Nawaprin (Torres et al., 2003). Subsequent studies demonstrated the ubiquitous nature of Waprin proteins across genera (Alam et al., 2019). These proteins are mostly found in the extracellular secretions of mammals such as whey protein in milk (Piletz et al., 1981), as well as in snakes (St. Pierre et al., 2008; Reeks et al., 2015), insects (Bouzid et al., 2014), scorpions (Romero-Gutierrez et al., 2017; Romero-Gutiérrez et al., 2018), cnidaria (Zhang et al., 2007) and frogs (Ali et al., 2002; Liu et al., 2013) among others. Additionally, *waprin*-like genes have also been reported in the transcriptomes of several other organisms, like cnidarians (Klompen et al., 2020) and ants (Bouzid et al., 2014). A thorough structural and functional characterization of the Waprin proteins discovered in the venoms of *N. nigricollis* (Torres et al., 2003), Inland taipan snake (*Oxyuranus microlepidotus*) (Nair et al., 2007) and Venezuelan horned frog (*Ceratophrys calcarata*) (Liu et al., 2013) has been done. This led to the identification of eight cysteine residues at the core of Waprin protein, which facilitate the formation of four disulphide bonds (Suntravat et al., 2016; Alam et al., 2019).

The WAP-domain containing proteins are involved in multifarious activities including antiviral
activity (Chen et al., 2008), antimicrobial property (Ali et al., 2002; Hagiwara et al., 2003;
Hauton et al., 2006; Smith et al., 2008; Imjongjirak et al., 2009; Li et al., 2013), Na-K ATPase
inhibitory activity, protein and proteinase inhibitory activities (Ali et al., 2002;Li et al.,
2013),and immunomodulatory functions (Gao et al., 2021). The most studied function of
Waprin proteins include its antimicrobial activity against various gram-positive and gram-
negative bacteria and its antifungal activity against certain fungi (Liu et al., 2013). Ceratoxin
isolated from the skin secretions of *C. calcarata*, has been shown to inhibit *Staphylococcus*
aureus, a gram-positive bacterium (Liu et al., 2013), whereas Waprin protein (Omwaprin) from
O. microlepidotus specifically targets gram-negative bacterial strains like *Staphylococcus*
warneri and *Bacillus megaterium* (Nair et al., 2007). The antimicrobial activity of Omwaprin
remains stable under high salt concentration (up to 350mM NaCl) and is dependent upon its
eight cysteine residues or its six N-terminal residues, four of which are positively charged. This
suggested the critical role of the four disulphide bonds within the core of Waprin structure for
its antibacterial activity (Nair et al., 2007). In a recent study, Thankappan *et al*. (2019) used
Omwaprin to derive two cationic antimicrobial peptides (Omw1 and Omw2) with enhanced
antimicrobial, antifungal, and antibiofilm activities (Thankappan and Angayarkanni, 2019).
Identification of transcripts encoding Waprin-like proteins is, so far, limited only to a few insect
species, such as, guinea ant, *Tetramorium bicarinatum* (Bouzid et al., 2014), wood-boring
woodwasp, *Sirex nitobei* (Gao et al., 2021), and honeybee, *Apis mellifera* (Lee et al., 2023).
While the function of *waprin* in *T. bicarinatum* remains elusive, *waprin* of *S. nitobei* is
hypothesized to promote the growth of a symbiotic fungus, *Amylosterium areolatum*, by
exercising antibacterial activities. During oviposition, female *S. nitobei* injects its venom along
with arthrospores of *A. areolatum* into the tree trunk. *A. areolatum* helps in the growth of *S.*
nitobei eggs and larvae by providing nutrition while *waprin* in the venom inhibits the bacterial
growth, facilitating the growth of *A. areolatum* (Coutts, 1969a, 1969b; Gao et al., 2021).
However, the recent report of Waprin (Amwaprin) in *A. mellifera* is the first direct evidence
about the antibacterial potential of an insect Waprin. Recombinant Amwaprin produced in
baculovirus-infected Sf9 cells exhibited microbicidal and anti-elastolytic activities (Lee et al.,
2023).
The red flour beetle, *Tribolium castaneum*, is a pest of stored food grains. It's wide
geographical range and infestation potential in a variety of stored grains makes it an
economically important pest. Additionally, the extent of RNAi efficiency in *T. castaneum*
makes it an excellent model for reverse genetics studies (Campbell et al., 2021). Recently,

while analysing sex-specific transcriptomes (unpublished data) of six-day old adult *T. castaneum*, our group identified a female-biased (FPKM value-179.20 and 0.35 for females and males, respectively) transcript of *waprin* (*Tc_Wap^F*) gene. *Tc_Wap^F* attracted our attention as no sex-specific function of *waprin* has been reported in any organism till date. Therefore, in this study, we characterized *Tc_Wap^F* to unravel the significance of its female-exclusive expression. Developmental- and tissue-specific profiling of *Tc_Wap^F* confirmed its exclusive expression in adult female tissues. Furthermore, our study revealed that the expression of *Tc_Wap^F* is independent of the classical sex determination pathway of *T. castaneum*. Although, we fail to observe any phenotypic alteration in *Tc_Wap^F* knockdown females, eggs laid by these females failed to hatch, indicating a direct involvement of *Tc_Wap^F* in embryonic development. This pioneering study, reveals a novel aspect of sex-specifically regulated Waprin protein to contribute towards the embryonic development. Further insights into the mechanistic roles of Waprin proteins in the insect kingdom holds promise for their utilization in pest management strategies.

3. Materials and methods

3.1 Rearing of *Tribolium castaneum*

Wild-type strain of *T. castaneum* was reared on whole-wheat flour supplemented with 5% yeast at 30°C temperature and 60-70% relative humidity. Sexing of pupae and adults were done based on the presence or absence of sexually dimorphic characters (Shukla and Palli, 2012a) with the help of Leica s9i stereomicroscope (Leica microsystems, Germany).

3.2 RNA Isolation and RT-PCR

Total RNA was isolated using Trizol reagent (Invitrogen) from different developmental stages of *Tribolium castaneum* including eggs, 6th instar larvae, Q stage, sex-specific pupae and sex-specific adults. Additionally, total RNA was also isolated from gut, ovary, head and fatbody tissues of female adults. Isolated RNA samples were DNAse (Promega,) treated, and the concentrations of DNA-free RNA samples were determined using NanoDrop (Jenway Genova Nano, USA). Subsequently the RNA samples were stored at -20°C until further use. For cDNA synthesis, 3μg of RNA per sample was denatured at 70°C for 5 min and then immediately chilled on ice. First strand cDNA was synthesized using GoScript Reverse Transcriptase (Promega) with 18-mer oligo(dT) primer, according to the manufacturer's instructions. The prepared cDNA samples were diluted ten folds with nuclease-free water and used for subsequent PCR studies. GoTaq polymerase (Promega) was used for all the RT-PCR reactions

using gene-specific primers (Table S1) designed using Primer-3 software (Primer3 Input, version 0.4.0). PCR conditions comprised initial denaturation at 94°C for 2 min followed by 32 cycles of denaturation at 94°C for 30 s, annealing (temperatures in Table S1) for 30 s, extension at 72°C for 2 min and a final extension at 72°C for 10 min. The amplified products were electrophoresed on 1.5% agarose gels and visualised using a BioRad gel documentation system.

3.3 Sequence analysis

Following the identification of Tc_Wap^F gene from our transcriptome data (unpublished), its sequence was validated through BLAST (BLASTn) searches in the EST database of *T. castaneum*. Genomic DNA corresponding to *waprin* transcript was identified by performing BLAST (BLASTn) searches in the genomic database of *T. castaneum*. Subsequently, the exons and introns of Tc_Wap^F were identified by aligning the gene-specific EST sequences with their corresponding genomic DNA sequence using Splign software (https://www.ncbi.nlm.nih.gov/sutils/splign/splign.cgi).

3.4 Double stranded RNA (dsRNA) synthesis and injections

PCR was performed using cDNA from adult females as template and gene-specific primers (Table S1) containing the T7 promoter sequences at their 5' ends to amplify the specific regions of Tc_Wap^F (Fig. S1), *Tcdsx* and *Tctra* genes (Shukla and Palli, 2012a, 2012b). Purified amplicons were used as template in in-vitro transcription reaction using T7 RiboMAX™ Express RNAi System (Promega) according to the manufacturer's instructions. The control dsRNA was synthesized using an amplified fragment of the *GFP* gene sequence.

Newly emerged (0 day) female adults (5 h post adult emergence) were immobilized by keeping them at 4°C for 8–10 min prior to injections. Subsequently, anesthetized insects were then injected either with ds*GFP* (control) or with dsRNA targeting specific genes (Tc_Wap^F, *Tcdsx* or *Tctra*) using an aspirator tube assembly (Sigma) attached to a 3.50 mm glass capillary tube (Drummond Scientific) drawn by a micropipette puller (Narishige, PC-100). Approximately, 500–600 ng dsRNA was injected per insect. Injected insects were starved and allowed to recover at room temperature for around 5 to 6 hours before transferring them to the incubator under standard rearing conditions. Knockdown efficacy of Tc_Wap^F, *Tcdsx* and *Tctra* were analyzed after 5 days of injections. The ratio of gene expression between beetles injected with

168   target dsRNA and those injected with *GFP* dsRNA was used to calculate the efficacy of gene

169   knockdown in the RNAi insects.

3.5 Parental RNAi of *Tc_Wap^F*

To investigate the impact of *Tc_Wap^F* knockdown in the subsequent generation, target gene

specific dsRNA or control dsRNAs were injected into virgin adult females (0 day) following

the protocol described in the previous section (3.4). Subsequently, on 5th day, injected females

were allowed to mate with their counterparts (uninjected males of the same age) maintaining

one mating pair per cup. After one day of mating, each mating pair was transferred to fresh cup

containing fine flour and eggs laid were harvested after every 2 hours to obtain embryos of

different developmental stages. The developmental progression of staged eggs was tracked by

staining and visualising the dechorionated eggs and, the females (injected with either

ds*Tc_Wap^F* or ds*GFP*) were dissected to compare their tissue morphology.

3.6 Embryo collection, egg staining and imaging

A series of treatments were given to the staged eggs for embryo staining: initially staged eggs

were cleaned by rinsing them in water. Cleaned eggs were treated with 25% bleach solution for

1.5 minutes with continuous shaking for their dichlorination. Subsequently, eggs were washed

with double distilled water and then fixed in 4% paraformaldehyde solution for 15-20 minutes

followed by the removal of aqueous phase. Further, the eggs were vigorously shaken in

methanol to remove the vitelline membrane. The devitellinized eggs were then washed with

PBST buffer (PBS buffer containing 0.1% Triton X-100). At last, the eggs were stained with

Hoechst 33258 (Thermo Fisher) dye and imaged using Leica DMI6000 fluorescence

microscope.

3.7 qRT-PCR Analysis

Total RNA isolation and first strand cDNA synthesis from control and knockdown females was

done as described previously. qRT-PCR reactions were performed using synthesized cDNAs

as template and gene-specific primers (Table S1) to quantify the relative mRNA levels of

Tc_Wap^F with GoTaq® qPCR Master Mix (Promega) according to the manufacturer's

instructions.Three independent biological replicates were used for each quantitation. *T.*

castaneum ribosomal protein (*RPS-18*) gene was used as an endogenous control to normalize

the expression data (Toutges et al., 2010), and $2^{-\Delta\Delta CT}$ method was employed to analyze the gene

expression levels (Livak and Schmittgen, 2001). The qPCR reactions were carried out in a

CFX96 touch real-time PCR detection system (Bio-Rad) with following cycling parameters:
initial denaturation at 95 °C for 30 s, followed by 40 cycles of 95 °C for 15 s, 60 °C for 45 s
and 72 °C for 5s. Melt curve analysis was carried out in the temperature range of 60–95°C, in
steps of 0.5 °C increments per 5s.

3.8 Phylogenetic Analysis and Domain Architecture

Protein sequence of *T. castaneum* Waprin (XP_008198656.1) was used to perform BLAST
searches (BLAST-p) in the non-redundant (nr) protein database of NCBI. The protein
sequences of different insects and few other organisms (including bacteria, crustaceans,
scorpions, reptiles, amphibians and mammals) were aligned by MUSCLE and the aligned
sequences were used to construct the phylogenetic tree using the Maximum likelihood method
in MEGA 11.0. Phylogenetic analysis was performed using the iTOL tool which recognizes the
Newick file. Domain searches and motif scan were conducted using online softwares,
Conserved domain database (https://www.ncbi.nlm.nih.gov/cdd/) and motif scan
(https://myhits.sib.swiss/cgi-bin/motif_scan), respectively. Eight conserved cysteine residues
were used as a signature to identify the conserved domains and establish the relationship
between their associations and divergence.

3.9 Statistical Analysis

The statistical analysis was performed using GraphPad Prism (9.5.1). *t-test* was performed to
analyse the *Tc_Wap^F* gene expression and its knockdown efficiency while one way ANOVA
were used to compare the *TcTra* and *Tcdsx* gene expression between the control and knockdown
samples. 3-4 biological replicates per group were used in study and all experiments were
performed thrice.

4. Results

4.1. *Tc_Wap^F* is exclusively expressed in adult female tissues

Tc_Wap^F (XM_008200434.2) was identified in the sex-specific transcriptome of six-day old
adult *T. castaneum*. The *Tc_Wap^F* gene harbours 2 exons and 1 intron with an ORF of 240 bases
encoding 80 amino acids (aa) (Fig. S1A).

Tc_Wap^F was found to express only in female adults in our stage wise developmental profiling
(Fig. 1A). No expression of *Tc_Wap^F* was detected during larval or pupal stages. Further it was
found to express in the ovary, gut, and fatbody tissues of female adults (Fig. 1B). Females were
found to possesses 700-fold higher expression of *Tc_Wap^F* as compared to males in our qRT-

PCR analysis (Fig. 1C). Increasing expression pattern of Tc_Wap^F was observed in the
developing embryos (Fig. S3).

4.2. Expression of Tc_Wap^F is not regulated by sex-determination cascade of *T. castaneum*

To analyze whether the sex-specific expression of Tc_Wap^F is regulated by the sex-
determination cascade of *T. castaneum*, we scanned the promoter sequence of Tc_Wap^F for the
presence of Doublesex (Dsx) binding site. The *dsx* (a transcription factor) is the most
downstream gene of insect sex determination cascade and is known to regulate the transcription
of sex-specifically expressed genes (Shukla and Palli, 2012b; Clough et al., 2014). A conserved
Dsx-binding motif (ACAATGT) was found about 4.5 Kb upstream to the Tc_Wap^F transcription
start site.

dsRNA (ds*Tcdsx*) injections (in male and female adults of *T. castaneum*) targeting common
region of *Tcdsx* led to a significant reduction in the gene expression in both male and female
(Fig. 2A). However, no change in the expression of Tc_Wap^F was observed in *Tcdsx* knockdown
adults (females or males) as compared to control adults (females or males) clearly indicating
that sex-specific expression of Tc_Wap^F is not regulated by *Tcdsx* (Fig. 2B).

We further examined if female-specific expression of Tc_Wap^F is regulated by some unknown
transcription factor (other than TcDsx) controlled by *TcTra*. TcTra is a splicing factor
responsible for female specific splicing of *Tcdsx* (Shukla and Palli, 2012b). Similar to the
strategy employed earlier dsRNA mediated knockdown of *Tctra* was done in females. *dsTctra*
injections led to an efficient knockdown of the target gene as evident by the detection of both,
female and male *Tctra* isoforms in *dsTctra* injected females (Shukla and Palli, 2014) in our RT-
PCR results (Fig. S4). Surprisingly, no change in the expression levels of Tc_Wap^F was
observed in *Tctra* knockdown females (Fig. 2C). Our results suggest that female-specific
expression of Tc_Wap^F is independent of the classical sex determination cascade of *T.*
castaneum.

4.3. Development of eggs laid by Tc_Wap^F knockdown females is arrested

dsRNA injections targeting Tc_Wap^F in females lead to a ~12-fold reduction in the expression
of Tc_Wap^F (Fig. 3A). Eggs laid by Tc_Wap^F knockdown females developed normally until 45
hours from their laying. These eggs showed underdeveloped germband, shrivelled and hollow
appearance of embryos in their later stages of development (Fig.3B). Further, none of the eggs
laid by Tc_Wap^F knockdown females hatch even after 96 hours whereas the eggs laid by

females injected with control dsRNA developed normally and hatched into larvae within 65-
72 hours of egg laying.

4.4 Waprin is conserved across insecta
A total of 65 Waprin and Waprin-related protein sequences from seven different orders of class
insecta and six outgroups were used for phylogenetic analysis. This list also included two
additional *waprin* genes (*Tc_Wap²* and *Tc_Wap³*) identified from the database of *T. castaneum*.
The tree highlighted that *Tc_Wap^F* and its homologs *Tc_Wap²* and *Tc_Wap³* started from the
same parent branch but diverged into different sister branches (Fig. 4A). Tc_Wap^F was found
to be most closely related to Waprin of *T. madens*, while Tc_Wap² protein sequence was found
to be more similar to *Sandaracinus amylolyticus*. On the other hand, Tc_Wap³ protein sequence
showed similarity to the Waprin of *Hermetia illucens*.
Domain architecture analysis using TB tools predicted the presence of WAP domains in all the
protein sequences analysed (Fig. 4B). The WAP domain was found to be conserved in the
Waprin proteins of all the species included in our analysis. The number of these WAP domains
varied (ranging from one to four) in different Waprin proteins. Tc_Wap^F possesses a single
WAP domain whereas Tc_Wap³ has two WAP domains. Besides, some of the Waprin proteins
were found to possess additional domains also, predominantly PLAC and Kunitz domain.

5. Discussion
Waprin (WAP domain containing) proteins are found in diverse groups of organisms (Alam et
al., 2019) and are known to possess antiviral (Chen et al., 2008), antimicrobial (Ali et al., 2002;
Hagiwara et al., 2003; Hauton et al., 2006; Smith et al., 2008; Imjongjirak et al., 2009; Li et
al., 2013), Na-K ATPase inhibitory, protein and proteinase inhibitory (Ali et al., 2002; Li et al.,
2013) as well as immunomodulatory functions (Gao et al., 2021). In addition to their presence
in snake venom, Waprin proteins have also been reported in a few insect species, such as the
wood-boring woodwasp, *S. nitobei* (Gao et al., 2021), the guinea ant, *T. bicarinatum* (Bouzid
et al., 2014) and honeybees (*A. mellifera*) (Lee et al., 2023). Ironically, except for the
microbicidal and anti-elastolytic activities of Amwaprin, a recently discovered Waprin protein
from *A. mellifera*, the biological significance and evolutionary history of Waprin proteins in
the insect kingdom remain largely unknown (Lee et al., 2023).

Our study presents a detail analysis of *Tc_Wap^F*, a female-specific homologue of snake Waprin
in *Tribolium castaneum* including its unique expression patterns and crucial role in embryonic

development. Tc_Wap^F, which possesses eight conserved cysteine residues (Fig. S1B), a typical feature of Waprin proteins is exclusively expressed in adult female tissues, including the ovary, gut, and fatbody (Fig. 1B). Besides, we also identified two additional *waprin* genes (Tc_Wap^2 and Tc_Wap^3) in *Tribolium* (Fig. S2); however, these genes were found to express in a non-sex specific manner (Fig. S2). Sexually dimorphic expression of Tc_Wap^F indicated its function in female specific physiology or reproduction. To deduce the biological relevance of sex-specific expression of Tc_Wap^F, dsRNA mediated knockdown experiments were performed in adult females. Despite the significant knockdown in the expression of Tc_Wap^F (Fig. 3A), no observable changes were noticed in the morphology or anatomy of the knockdown individuals (data not shown). Furthermore, the fecundity of Tc_Wap^F knockdown females were unaffected (data not shown). Interestingly, none of the eggs laid by Tc_Wap^F knockdown females hatch. We hypothesized the failure of fertilisation or the growth arrest of fertilised embryos as the reason for the same. To test this hypothesis, time course tracking of staged eggs either laid by Tc_Wap^F knockdown females (knockdown eggs/embryos) or by control females (control eggs/embryos) was done. Head and leg appendices became prominent in the control embryos within 72 hours of their development whereas only a few larval structures were noticed in the knockdown embryos (Fig. 3B). Growth of knockdown embryos appeared arrested at the germband retraction stage and ultimately these eggs failed to hatch and started desiccating after 60-65 hours of egg laying. The expression of Tc_Wap^F in female adults (Fig. 1A), its gradual increase in developing eggs (Fig. S3) and the failure of hatching of eggs laid by Tc_Wap^F knockdown females points towards the crucial role of Tc_Wap^F in embryonic development in *T. castaneum*.

Since the effect of Tc_Wap^F injections were pronounced in the embryos laid by injected females, there could be two possibilities; 1) Tc_Wap^F transcript (or protein) may be maternally supplied to the embryos for their proper development. Injections of dsTc_Wap^F in the mother lead to a reduced transfer of Tc_Wap^F transcript (or protein) in the embryos resulting in the adverse effect on their development. This hypothesis is supported by the detection of Tc_Wap^F transcripts in abundance during 0-2 hours of egg stage (unfertilised) and its subsequent absence in the eggs of 2-4 hours development. Tc_Wap^F transcripts were again detected in 6-8 hours old eggs which was found to be gradually increased in later stages (Fig. S3). The expression profile of Tc_Wap^F during the embryonic development of *T. castaneum* suggests the switching of Tc_Wap^F transcription from maternal to zygotic. 2) Alternatively, the dsRNA injected into female adults might have been passed to the embryos, leading to a reduction of Tc_Wap^F

proteins in embryos and their growth arrest. In either of the cases, Tc_Wap^F seems to be
essentially required for the development of *Tribolium* embryos.

Our hypothesis aligns with several previous studies demonstrating strategies adopted by
mother to invest parental resources for the enhanced survival of their offspring. For example,
in *Blatella germanica* the maternal supply of hydrocarbons is required to prevent the water loss
from eggs (Fan et al., 2008). Deas *et al*. (2012), reported the plasticity in the egg laying
behaviour of *Mimosestes amicus* (a seed beetle) female to protect its eggs from parasitic wasps
(Deas and Hunter, 2012). Another evidence of maternal strategies for the enhanced survival of
offsprings comes from mice, where high expression levels of *Wap* gene in the mammary glands
of lactating mice play an essential role in nourishing their offspring (Triplett et al., 2005;
Oftedal, 2012). Interestingly, the authors noticed a reduced embryonic development but failed
to observe any phenotypic abnormalities in the mammary glands of *Wap*-knockout females
which has striking resemblance to the results obtained from our knockdown studies.

Furthermore, the expression of *Tc_Wap^F* in the ovary and fatbody has a relevance as the ovary
is the site of egg development, and the fatbody is the major site of vitellogenin synthesis, which
facilitates yolk deposition in eggs (Fig. 1B). At present, we are unable to justify the reason for
high expression of *Tc_Wap^F* in the *Tribolium* gut but a possible antimicrobial role of Tc_Wap^F
similar to the case of other Waprin proteins, cannot be denied.

The central axis (*tra-dsx*) of insect sex determination cascade is also conserved in *T. castaneum*
(Shukla and Palli, 2012a). Sex-specifically spliced *Tcdsx* produces sex-specific proteins,
TcDsx^M in males and TcDsx^F in females, which are sex-specific transcription factors regulating
all the genes expressed in a sex-specific manner (Shukla and Palli, 2012b). The expression
levels of sex specific Dsx targets have been shown to increase in one sex while decrease in
another sex upon Dsx knockdown in the two sexes (Verhulst and van de Zande, 2015). To
explore whether the sex-specific expression of *Tc_Wap^F* gene is regulated by TcDsx^F, we
examined the presence of TcDsx binding sites in the *Tc_Wap^F* gene. Although a conserved Dsx-
binding motif (ACAATGT) was identified 4.5 Kbp upstream to the *Tc_Wap^F* transcription start
site, we did not observe a decrease in the expression of *Tc_Wap^F* in *Tcdsx* knockdown females
(Fig. 2B). Additionally, no upregulation in the expression of *Tc_Wap^F* was seen in *Tcdsx*
knockdown males. We further hypothesized that *Tc_Wap^F* transcription might be regulated by
an undiscovered transcription factor under the control of TcTra, a splicing regulator of *Tcdsx*.
To test this hypothesis, we performed the knockdown of *Tctra* in female adults, but no change

in the expression Tc_Wap^F was observed in *Tctra* knockdown females (Fig. 2C). These results

suggest that the female-specific expression of Tc_Wap^F is not governed by the established sex

determination cascade of *T. castaneum*. It is possible that the sex-specific expression of

Tc_Wap^F is regulated by a pathway diverging upstream of *Tctra* (Shukla and Palli, 2012a;

Netschitailo et al., 2023). Alternatively, the sex-specific expression of Tc_Wap^F gene could

result from differences in chromatin confirmation between the sexes. This is supported by a

recent study highlighting the differential chromatin confirmation as the regulator of sex-biased

gene expression in two *Drosophila* species, *D. melanogaster* and *D. simulans* (Nanni et al.,

2023).

In conclusion, we identified a sex-specific homologue of snake *Waprin* (Tc_Wap^F) from *T.*

castaneum and analysed its function. Our study suggested the crucial role of Tc_Wap^F in the

embryogenesis of *T. castaneum* which provides the foundation for further investigations into

the molecular functions of *waprin* genes in other insects. The sex-specific expression of

Tc_Wap^F is not governed by the classical sex-determination cascade of *T. castaneum*. *waprin*

genes were found to be conserved across insect species which indicates its functional integrity

throughout evolution. Future research focusing sex-specific *waprin* genes and its regulation in

insects could provide valuable insights into their potential applications in enhancing the insect

control strategies utilising sterile insect technology.

6. Author's Contribution:

388 CC and JNS conceived the idea of manuscript. CC performed knockdown, PCR and qRT-PCR

389 experiments. DK did the phylogenetic analysis. KKM maintained the insect cultures and helped

390 CC in preparation of the manuscript. CC, VV and JNS wrote the manuscript and all others

391 agreed to it. All authors contributed to the article and approved the submitted version.

7. Acknowledgements:

394 JNS and VV acknowledge their Ramalingaswami fellowships (BT/RLF/Re-entry/10/2015 and

395 BT/RLF/Re-entry/22/2017, respectively) from the Department of Biotechnology, Ministry of

396 Science and Technology, India. We are thankful for the qRT-PCR facility of the DBT Builder

397 project [BT/INF/22/SP44383/2021]. CC and KKM are the Ph.D. students and supported by

fellowship program of CURAJ. JNS also acknowledges Dr. Satnam Singh (PAU, Ludhiana) and Dr. Deepa Agashe (NCBS, Bangalore) for providing the *Tribolium castaneum* cultures.

8.Competing interests: The authors have 'No competing interests' to declare.

9.Funding: This work was supported by the grant (EMR/2017/001378/AS) from the Department of Science and Technology, Ministry of Science and Technology, India**.**

8. References:

Alam, Md.I., R, O., MA, A., H, Q., O, A., 2019. Therapeutic potential of snake venoms as antimicrobial agents. Front. drug chem. clin. 2. https://doi.org/10.15761/fdccr.1000136

Ali, M.F., Lips, K.R., Knoop, F.C., Fritzsch, B., Miller, C., Conlon, J.M., 2002. Antimicrobial peptides and protease inhibitors in the skin secretions of the crawfish frog, Rana areolata. Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics 1601, 55–63. https://doi.org/10.1016/S1570-9639(02)00432-6

Bouzid, W., Verdenaud, M., Klopp, C., Ducancel, F., Noirot, C., Vétillard, A., 2014. De Novo sequencing and transcriptome analysis for tetramorium bicarinatum: A comprehensive venom gland transcriptome analysis from an ant species. BMC Genom 15. https://doi.org/10.1186/1471-2164-15-987

Campbell, J.F., Athanassiou, C.G., Hagstrum, D.W., Zhu, K.Y., 2021. Annual Review of Entomology Tribolium castaneum: A Model Insect for Fundamental and Applied Research. Annu. Rev. Entomol. 67, 347–365. https://doi.org/https://doi.org/10.1146/annurev-ento-080921-075157

Chen, D., He, N., Xu, X., 2008. Mj-DWD, a double WAP domain-containing protein with antiviral relevance in Marsupenaeus japonicus. Fish Shellfish Immunol. 25, 775–781. https://doi.org/10.1016/j.fsi.2008.02.017

Clough, E., Jimenez, E., Kim, Y.A., Whitworth, C., Neville, M.C., Hempel, L.U., Pavlou, H.J., Chen, Z.X., Sturgill, D., Dale, R.K., Smith, H.E., Przytycka, T.M., Goodwin, S.F., VanDoren, M., Oliver, B., 2014. Sex- and tissue-specific functions of drosophila doublesex transcription factor target genes. Dev. Cell. 31, 761–773. https://doi.org/10.1016/j.devcel.2014.11.021

Coutts, M.P., 1969a. The Mechanism of Pathogenicity of Sirex Noctilio on Pinus Radiata II. Effects of S. Noctilio Mucus. Aust. J. Biol. Sci. 22, 1153–1162. https://doi.org/https://doi.org/10.1071/BI9691153

Coutts, M.P., 1969b. The Mechanism of Pathogenicity of Sirex Noctilio on Pinus Radiata I. Effects of the Symbiotic Fungus Amylostereum Sp. (Thelophoraceae). Aust. J. Biol. Sci. 22, 915–924. https://doi.org/https://doi.org/10.1071/BI9690915

Deas, J.B., Hunter, M.S., 2012. Mothers modify eggs into shields to protect offspring from parasitism. Proc. R. Soc. B: Biol. Sci. 279, 847–853. https://doi.org/10.1098/rspb.2011.1585

Fan, Y., Eliyahu, D., Schal, C., 2008. Cuticular hydrocarbons as maternal provisions in embryos and nymphs of the cockroach Blattella germanica. J. Exp. Biol. 211, 548–554. https://doi.org/10.1242/jeb.009233

Gao, C., Ren, L., Wang, M., Wang, Z., Fu, N., Wang, H., Wang, X., Ao, T., Du, W., Zheng, Z., Li, H.,
Shi, J., 2021. Proteo-transcriptomic characterization of sirex nitobei (Hymenoptera: Siricidae)
venom. Toxins (Basel) 13. https://doi.org/10.3390/toxins13080562

Hagiwara, K., Kikuchi, T., Endo, Y., Huqun, Usui, K., Takahashi, M., Shibata, N., Kusakabe, T., Xin,
H., Hoshi, S., Miki, M., Inooka, N., Tokue, Y., Nukiwa, T., 2003. Mouse SWAM1 and SWAM2
Are Antibacterial Proteins Composed of a Single Whey Acidic Protein Motif. J. Immunol. 170,
1973–1979. https://doi.org/10.4049/jimmunol.170.4.1973

Hauton, C., Brockton, V., Smith, V.J., 2006. Cloning of a crustin-like, single whey-acidic-domain,
antibacterial peptide from the haemocytes of the European lobster, Homarus gammarus, and its
response to infection with bacteria. Mol. Immunol. 43, 1490–1496.
https://doi.org/10.1016/j.molimm.2005.07.029

Imjongjirak, C., Amparyup, P., Tassanakajon, A., Sittipraneed, S., 2009. Molecular cloning and
characterization of crustin from mud crab Scylla paramamosain. Mol. Biol. Rep. 36, 841–850.
https://doi.org/10.1007/s11033-008-9253-0

Klompen, A.M.L., Macrander, J., Reitzel, A.M., Stampar, S.N., 2020. Transcriptomic analysis of four
cerianthid (Cnidaria, Ceriantharia) venoms. Mar. Drugs 18.
https://doi.org/10.3390/MD18080413

Lee, K.S., Kim, B.Y., Kim, Y.H., Choi, Y.S., Jin, B.R., 2023. Identification of waprin and its
microbicidal activity: A novel protein component of honeybee (Apis mellifera) venom. Comp.
Biochem. Physiol. Part - C: Toxicol. Pharmacol. 266.
https://doi.org/10.1016/j.cbpc.2023.109561

Li, S., Jin, X.K., Guo, X.N., Yu, A.Q., Wu, M.H., Tan, S.J., Zhu, Y.T., Li, W.W., Wang, Q., 2013. A
Double WAP Domain-Containing Protein Es-DWD1 from Eriocheir sinensis Exhibits
Antimicrobial and Proteinase Inhibitory Activities. PLoS One 8.
https://doi.org/10.1371/journal.pone.0073563

Liu, D., Wang, Y., Wei, L., Ye, H., Liu, H., Wang, L., Liu, R., Li, D., Lai, R., 2013. Snake venom-like
waprin from the frog of Ceratophrys calcarata contains antimicrobial function. Gene 514, 99–
104. https://doi.org/10.1016/j.gene.2012.11.007

Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time
quantitative PCR and the 2-$\Delta\Delta$CT method. Methods 25, 402–408.
https://doi.org/10.1006/meth.2001.1262

Nair, D.G., Fry, B.G., Alewood, P., Kumar, P.P., Kini, R.M., 2007. Antimicrobial activity of
omwaprin, a new member of the waprin family of snake venom proteins. Biochem. J. 402, 93–
104. https://doi.org/10.1042/BJ20060318

Nanni, A. V, Martinez, N., Graze, R., Morse, A., Newman, J.R.B., Jain, V., Vlaho, S., Signor, S.,
Nuzhdin, S. V, Renne, R., McIntyre, L.M., 2023. Sex-Biased Expression Is Associated With
Chromatin State in Drosophila melanogaster and Drosophila simulans. Mol. Biol. Evol. 40,
msad078. https://doi.org/10.1093/molbev/msad078

Netschitailo, O., Wang, Y., Wagner, A., Sommer, V., Verhulst, E.C., Beye, M., 2023. The function and
evolution of a genetic switch controlling sexually dimorphic eye differentiation in honeybees.
Nat. Commun. 14, 463. https://doi.org/10.1038/s41467-023-36153-4

Oftedal, O.T., 2012. The evolution of milk secretion and its ancient origins. Animal 6, 355–368.
https://doi.org/10.1017/S1751731111001935

Piletz, J.E., Heinlen, M., Ganschow, R.E., 1981. Biochemical characterization of a novel whey protein from murine milk. J. Biol. Chem. 256, 11509–11516. https://doi.org/10.1016/S0021-9258(19)68430-9

Reeks, T.A., Fry, B.G., Alewood, P.F., 2015. Privileged frameworks from snake venom. Cell. Mol. Life Sci. 72, 1939–1958. https://doi.org/10.1007/s00018-015-1844-z

Romero-Gutiérrez, M.T., Santibáñez-López, C.E., Jiménez-Vargas, J.M., Batista, C.V.F., Ortiz, E., Possani, L.D., 2018. Transcriptomic and proteomic analyses reveal the diversity of venom components from the vaejovid scorpion Serradigitus gertschi. Toxins (Basel) 10. https://doi.org/10.3390/toxins10090359

Romero-Gutierrez, T., Peguero-Sanchez, E., Cevallos, M.A., Batista, C.V.F., Ortiz, E., Possani, L.D., 2017. A deeper examination of thorellius atrox scorpion venom components with omic techonologies. Toxins (Basel) 9. https://doi.org/10.3390/toxins9120399

Shukla, J.N., Palli, S.R., 2014. Production of all female progeny: Evidence for the presence of the male sex determination factor on the y chromosome. J. Exp. Biol. 217, 1653–1655. https://doi.org/10.1242/jeb.100438

Shukla, J.N., Palli, S.R., 2012a. Sex determination in beetles: Production of all male progeny by Parental RNAi knockdown of transformer. Sci. Rep. 2. https://doi.org/10.1038/srep00602

Shukla, J.N., Palli, S.R., 2012b. Doublesex target genes in the red flour beetle, Tribolium castaneum. Sci. Rep. 2. https://doi.org/10.1038/srep00948

Smith, V.J., Fernandes, J.M.O., Kemp, G.D., Hauton, C., 2008. Crustins: Enigmatic WAP domain-containing antibacterial proteins from crustaceans. Dev. Comp. Immunol. https://doi.org/10.1016/j.dci.2007.12.002

St. Pierre, L., Earl, S.T., Filippovich, I., Sorokina, N., Masci, P.P., De Jersey, J., Lavin, M.F., 2008. Common evolution of waprin and kunitz-like toxin families in Australian venomous snakes. Cell. Mol. Life Sci. 65, 4039–4054. https://doi.org/10.1007/s00018-008-8573-5

Suntravat, M., Uzcategui, N.L., Atphaisit, C., Helmke, T.J., Lucena, S.E., Sánchez, E.E., Acosta, A.R., 2016. Gene expression profiling of the venomgland from the Venezuelan mapanare (Bothrops colombiensis) using expressed sequence tags (ESTs). BMC Mol. Biol. 17. https://doi.org/10.1186/s12867-016-0059-7

Thankappan, B., Angayarkanni, J., 2019. Biological characterization of omw1 and omw2: antimicrobial peptides derived from omwaprin. 3 Biotech 9. https://doi.org/10.1007/s13205-019-1801-x

Torres, A.M., Wong, H.Y., Desai, M., Moochhala, S., Kuchel, P.W., Kini, R.M., 2003. Identification of a novel family of proteins in snake venoms. Purification and structural characterization of nawaprin from Naja nigricollis snake venom. J. Biol. Chem. 278, 40097–40104. https://doi.org/10.1074/jbc.M305322200

Toutges, M.J., Hartzer, K., Lord, J., Oppert, B., 2010. Evaluation of reference genes for quantitative polymerase chain reaction across life cycle stages and tissue types of Tribolium castaneum. J. Agric. Food Chem. 58, 8948–8951. https://doi.org/10.1021/jf101603j

Triplett, A.A., Sakamoto, K., Matulka, L.A., Shen, L., Smith, G.H., Wagner, K.U., 2005. Expression of the whey acidic protein (Wap) is necessary for adequate nourishment of the offspring but not functional differentiation of mammary epithelial cells. Genesis (United States) 43, 1–11. https://doi.org/10.1002/gene.20149

Verhulst, E.C., van de Zande, L., 2015. Double nexus--Doublesex is the connecting element in sex
determination. Brief Funct Genomics 14, 396–406. https://doi.org/10.1093/bfgp/elv005

Zhang, X., Boot-Handford, R.P., Huxley-Jones, J., Forse, L.N., Mould, A.P., Robertson, D.L., Li, L.,
Athiyal, M., Sarras, M.P., 2007. The collagens of hydra provide insight into the evolution of
metazoan extracellular matrices. J. Biol. Chem. 282, 6792–6802.
https://doi.org/10.1074/jbc.M607528200

Figure Legends:

Fig.1A: Developmental expression profiling of Tc_Wap^F of *T. castaneum*. RT-PCR analysis
was performed using stage specific cDNAs (6th Instar Larvae, Q Stage, Larvae, female pupae,
male pupae, female adults and male adults) as templates. **B)** Tissue (ovary, gut, head and
fatbody of female adults) specific expression profile of Tc_Wap^F. **C)** Relative expression of
Tc_Wap^F in 6⁻day old adult male and female. The error bars represent the standard deviation
(**p value=0.0035).

Fig.2A: Graphical representation of the knockdown efficiency of *Tcdsx*. ~ 6-fold and 9-fold
reduction was observed in *Tcdsx* expression in female and males respectively. Fold change (*p
value≤ 0.0311; **p value ≤ 0.0003). **B)** The relative expression of Tc_Wap^F was quantified in
response to *Tcdsx* knockdown in female and male individuals. Error bars represent standard
deviation. **C)** Graphical representation of Tc_Wap^F expression upon *TcTra* knockdown. The
knockdown efficiency was analysed on 5th day after knockdown. Five biological replicates
were used in experiment.

Fig.3A: Graphical representation of the relative expression of Tc_Wap^F in the knockdown and
control females. Three biological replicates of both control and knockdown individuals were
taken for analysis. Error bars represent standard deviation (**p value≤ 0.0020). **B)** Microscopic
images of different stages of *Tribolium* embryos laid by the ds*GFP* and dsTc_Wap^F injected
females. Eggs laid by dsTc_Wap^F injected females developed normally until 45 hours, but
uneven germband was formed subsequently. White arrow represents the region of abnormality.
Embryos were stained by Hoechst dye and imaged using fluorescence microscope (Scale bar
100µm, magnification 10X).

Fig.4A: Circular phylogenetic tree of WAP proteins from different species. The protein
sequences were aligned using MEGA11 and the tree was constructed using the neighbour-

joining method. **B)** Domain architecture of various WAP domain containing proteins across

different species. Different domains are colour coded.

Fig.S1A: Schematics representing the structures of different *waprin* genes (*Tc_Wap^F^, Tc_Wap²*

and Tc_Wap³) of *T.castaneum. Tc_Wap^F^* consists of two exons and one intron, *Tc_Wap²* contains

three exons and two introns whereas *Tc_Wap³* possess two exons and one intron. Boxes

represent exons and lines represent the introns. Numbers represent the length of respective exon

or intron. Start and stop codons are represented by green and red bars (vertical). Horizontal

arrows represent the primer positions; orange- dsRNA primers, black- semi quantitative PCR;

pink- qRT PCR. **B)** Sequence alignment of *Tribolium* Waprin proteins. Conserved cysteine

residues are shown in yellow boxes.

Fig. S2: Expression profiling of *Tc_Wap²* and *Tc_Wap³. Tc_RPS-18* was used as an endogenous

control.

Fig.S3: Expression profile of *Tc_Wap^F^* in staged embryos. The time mentioned in the figure

represents the stage of eggs used for RNA isolation. *Tc_RPS-18* gene was used as an

endogenous control.

Fig.S4: Gel image showing the splicing status of *Tctra* in control (*dsGFP* injected) and

knockdown (ds*Tctra* injected) female adults. RT-PCR was performed using cDNA synthesised

from RNA isolated from control females (C1-C4) and knockdown females (T1-T4).

Table S1: List of primers (and their details) used in the study.

Table S2: *waprin* genes of *T.castaneum* and their respective accession ID's.

A)

B)

C)

Expression of *Tc_Wap^F*

A) **Knockdown efficacy of *Tcdsx***

B) ***Tc_Wap^F* expression upon *Tcdsx* knockdown**

C) ***Tc_Wap^F* expression upon *TcTra* knockdown**

A) **Knockdown efficiency of *Tc_Wap^F***

	0-2 Hrs.	5 Hrs.	15 Hrs.
CONTROL			
KNOCKDOWN			

	20 Hrs.	25 Hrs.	45 Hrs.
CONTROL			
KNOCKDOWN			

100 µm

50 Hrs.   **65 Hrs.**   **72 Hrs.**

CONTROL

KNOCKDOWN

96 Hrs.

KNOCKDOWN

100 µm

		WAP superfamily
		Abhydrolase superfamily
		LamG superfamily
		EGF_CA superfamily
		FN3 superfamily
		P12 superfamily
		Ig superfamily
		Kunitz-type superfamily
		NTR_like superfamily
		KAZAL_FS superfamily
		ADAM_spacer1 superfamily
		TSP1_ADAMTS superfamily
		ADAM_CR_3 superfamily
		PLAC superfamily
		TSP_1 superfamily
		PHA02826 superfamily
		Smc superfamily

Legionella jordanis — WP_238583468.1
Centruroides sculpatus — XP_023240390.1
Tribolium castaneum (Tc_Wap^F) — XP_008198656.1
Apis florea — XP_012345872.1
Gryllus bimaculatus — GLG98105.1
Tribolium castaneum (Tc_Wap^2) — XP_969756.1
Schistocerca serialis cubense — XP_049962616.1
Anopheles stephensi — XP_035900555.1
Xenopus laevis — XP_018114486.2
Zootermopsis nevadensis — KDR08241.1
Artibeus jamaicensis — XP_037020609.1
Apis mellifera — XP_006570013.1
Hermetia illucens — XP_037920036.1
Tribolium madens — XP_044262500.1
Coptotermes formosanus — GFG40022.1
Apis mellifera caucasica — KAG6800588.1
Penaeus monodon — XP_037777091.1
Apis laboriosa — XP_043792881.1
Rhabdophis tigrinus — sp|A7X4J4.1|
Schistocerca gregaria — XP_049828788.1
Firmicutes bacterium — CDC43220.1
Homalodisca vitripennis — XP_046675194.1
Ceratophrys calcarata — AFX83953.1
Limnonectes fujianensis — AUD53200.1
Xenopus tropicalis — XP_031750632.1
Procambarus clarkii — QER91040.1
Pollicipes pollicipes — XP_037089176.1
Daktulosphaira vitifoliae — XP_050528637.1
Diprion similis — XP_046745300.1
Bombus affinis — XP_050588546.1
Nilaparvata lugens — XP_039282956.1
Oxyuranus microlepidotus sp|B5G6G7.1|(Omwaprin)
Rana temporaria — XP_040203993.1
Sturnira hondurensis — XP_036900052.1
Tenebrio molitor — KAH0813828.1
Legionella longbeachae — EEZ96907.1
Odontomachus brunneus — XP_032664332.1
Holotrichia oblita — KAI4460211.1
Ooceraea biroi — XP_011335694.1
Onthophagus taurus — XP_022911597.1
Schistocerca cancellata — XP_049768279.1
Onychomys torridus — XP_036051172.1
Schistocerca americana — XP_046988404.1
Formica exsecta — XP_029661547.1
Rousettus aegyptiacus — XP_036085246.1
Mus musculus — DAA00309.1
Pseudomyrmex gracilis — XP_020281693.1
Holocephalus stephensii sp|B5G6H3.1|(Stewaprin)
Cryptotermes secundus — XP_023726130.1
Drosophila melanogaster — ADX64765.1
Molossus molossus — XP_036101544.1
Phyllostomus discolor — XP_035865778.1
Homalodisca vitripennis — XP_046675007.1
Teleopsis dalmanni — XP_037945583.1
Amphibalanus amphitrite — KAF0306937.1
Sandaracinus amylolyticus — WP_236607153.1
Polypterus senegalus — KAG2461723.1
Rattus norvegicus — NP_598221.2
Tribolium castaneum (Tc_Wap^3) — XP_969824.1
Philodryas olfersii — sp|A7X4K7.1|
Naja nigricollis — pdb|1UDK|
Phodopus roborovskii — CAH6997122.1
Drosophila subobscura — XP_034662907.1
Harpegnathos saltator — EFN85384.1
Bemisia tabaci — XP_018917702.1

5'   3'

0   500   1000   1500   2000   2500   3000