A sex-specific homologue of snake *waprin* is required for embryonic development in the
Red Flour Beetle, *Tribolium castaneum*.

Chhavi Choudhary, Saloni Sharma, Keshav Kumar Meghwanshi, Divyanshu Kishore, Vivek Verma,
Jayendra Nath Shukla*

#Department of Biotechnology, School of Life Sciences, Central University of Rajasthan, Ajmer, 305817
Rajasthan, India.

***Corresponding author, email ID: jayendrashukla@curaj.ac.in**

12 1. Abstract:

Waprin, first discovered in snake venom, is an extracellular secretory protein known for its
antibacterial activity. Analysis of sex-specific transcriptomes of adult *Tribolium castaneum*
led to the identification of a female-specific Waprin protein gene (Tc_Wap^F) which expresses
only in female adults. Knockdown of Tc_Wap^F didn't show any effect in the same generation,
but the eggs laid by Tc_Wap^F knockdown females failed to develop/hatch. No expression of
Tc_Wap^F was detected in *Tribolium* embryos, hence considering the antibacterial properties of
Waprin proteins and the failure of hatching of eggs laid by Tc_Wap^F knockdown females, we
assume the involvement of Tc_Wap^F in protecting the embryos from the environmental
pathogens. Tc_Wap^F is not regulated by sex determination pathway of *T. castaneum* as the
expression of Tc_Wap^F was not changed upon the knockdown of sex determination pathway
genes of *T. castaneum*. Further, we also identified two more *waprin* genes of *T. castaneum*
expressing in a non-sex specific manner at all the developmental stages. To the best of our
knowledge this study is the first attempt to characterize the developmental role of *waprin*
genes in any insect. The study presented here provides an opportunity to probe the potential
of *Tribolium* Waprin proteins as non-conventional antimicrobials.

29 2. Introduction:

Waprin, a WAP (Whey Acidic Protein) domain containing secretory protein, was first
identified in the venom of Black-necked spitting cobra, *Naja nigricollis*, and was named
Nawaprin (Torres et al. 2003). Later studies demonstrated the ubiquitous nature of Waprin
proteins across genera (Alam et al. 2019). Waprin proteins were primarily identified in the

extracellular secretions from mammals (whey protein in milk), snakes (St. Pierre et al. 2008;

Reeks, Fry, and Alewood 2015), insects (Bouzid et al. 2014), scorpions (Romero-Gutierrez et

al. 2017; Romero-Gutiérrez et al. 2018), cnidaria (Zhang et al. 2007) and frogs (Ali et al.

2002; Liu et al. 2013) etc. Waprin proteins discovered in the venoms of *N. nigricollis* (Torres

et al. 2003), Inland taipan snake (*Oxyuranus microlepidotus*) (Nair et al. 2007) and

Venezuelan horned frog (*Ceratophrys calcarata*) (Liu et al. 2013) has been structurally and

functionally characterized. Besides, the presence of *waprin* like gene has been reported in the

transcriptome of several other organisms like scorpions (Romero-Gutierrez et al. 2017;

Romero-Gutiérrez et al. 2018), cnidarians (Klompen et al. 2020) and ants (Bouzid et al.

2014). The peculiar feature of Waprin protein is the presence of eight cysteine residues at the

core of the protein, which are involved in the formation of four disulphide bonds (Suntravat

et al. 2016; Alam et al. 2019).

Interestingly, the WAP domain proteins have been reported to have multifarious activities

including antiviral activity (Chen, He, and Xu 2008), antimicrobial activity (Ali et al. 2002;

Hagiwara et al. 2003; Hauton, Brockton, and Smith 2006; Smith et al. 2008; Imjongjirak et

al. 2009; Li et al. 2013), Na-K ATPase inhibitory activity, protein and proteinase inhibitory

activities (Ali et al. 2002; Li et al. 2013), as well as immunomodulatory functions (Gao et al.

2021). Antimicrobial activity against various gram-positive and gram-negative bacteria and

antifungal activity against certain fungi are the most widely studied functions of Waprin

proteins (Liu et al. 2013). For instance, ceratoxin from the skin secretions of *C. calcarata*, has

been shown to inhibit *Staphylococcus aureus*, a gram-positive bacterium (Liu et al. 2013)

whereas Waprin protein (Omwaprin) of *O. microlepidotus* works selectively against the

strains of *Staphylococcus warneri* and *Bacillus megaterium*, gram negative bacteria (Nair et

al. 2007). The antimicrobial activity of Omwaprin is salt tolerant up to 350mM NaCl and this

activity is lost upon reduction and alkylation of its eight cysteine residues or upon disruption

of its six N-terminal residues, four of which are positively charged. This suggested that the

four disulphide bonds are crucial for the antibacterial activity of Waprin proteins (Nair et al.

2007). In a recent study, Thankappan et al (2019) used Omwaprin to derive two cationic

antimicrobial peptides (Omw1 and Omw2) with enhanced antimicrobial, antifungal, and

antibiofilm activities (Thankappan and Angayarkanni 2019).

Identification of transcripts related to *waprin* is limited only to a few insect species i.e., in the

wood-boring woodwasp, *Sirex nitobei* (Gao et al. 2021) and in guinea ant, *Tetramorium*

bicarinatum (Bouzid et al. 2014), but the first ever study to show the antibacterial potential of

any insect Waprin has recently been done for the Waprin (Amwaprin) of honeybees (*Apis*

millefera). Recombinant Amwaprin produced in baculovirus-infected Sf9 cells was shown to

exhibit microbicidal and anti-elastolytic activities (Lee et al. 2023). The function of *waprin* in

T. bicarinatum is not known but the *waprin* of *S. nitobei* has been hypothesized to promote

the growth of a symbiotic fungus, *Amylosterium areolatum*. During oviposition, *S. nitobei*

female injects its venom along with arthrospores of *A. areolatum* into the tree trunk. *A.*

areolatum helps in the growth of *S. nitobei* eggs and larvae by providing nutrition to them

and in return, Waprin present in the venom of *S. nitobei* inhibits the bacterial growth, inturn

facilitating the growth of *A. areolatum* (Coutts 1969b; 1969a; Gao et al. 2021).

The red flour beetle, *Tribolium castaneum*, is a pest of stored food grains. Its wide

geographical range and infestation potential in a variety of food products makes it an

economically important pest. Besides, the extent of RNAi efficiency in *T. castaneum* makes it

an excellent model for reverse genetics studies (Campbell et al. 2021). Screening of sex-

specific transcriptomes (unpublished) of six-day old adult *T. castaneum* led to the

identification of a female-biased (FPKM value-179.20 and 0.35 for females and males

respectively) transcript of *waprin* (Tc_Wap^F) gene. qRT-PCR using sex-specific (female and

male adults) *Tribolium* cDNAs and Tc_Wap^F specific-primers confirmed the female-exclusive

expression of Tc_Wap^F. Developmental and tissue-specific profiling suggests the expression

of Tc_Wap^F exclusively in all the tissues of female adults. RNAi mediated knockdown in the

expression of Tc_Wap^F in adult females did not produce any phenotypic alteration in the

injected individuals whereas its parental RNAi resulted in the failure of hatching of eggs laid

by Tc_Wap^F knockdown females. Since, Tc_Wap^F is not expressed in embryos and parental

RNAi of Tc_Wap^F leads to the failure in the hatching of eggs, we hypothesize that Tc_Wap^F is

maternally supplied for the development of embryos. Considering the antimicrobial potential

of Waprin proteins (Amwaprin in particular) (Lee et al. 2023), we further hypothesize that the

maternal transfer of Tc_Wap^F has evolved to provide antimicrobial properties to the embryos.

Sex-specific expression and identification of a consensus Doublesex (Dsx), the most

downstream gene of insect sex-determination cascade binding site in the upstream region of

Tc_Wap^F prompted us to explore whether the expression of Tc_Wap^F is controlled by the sex

determination pathway genes of *T. castaneum*. For this we performed the RNAi mediated

knockdown of *T. castaneum* transformer (*Tctra*) and *dsx* (*Tcdsx*) genes and compared the

expression of Tc_Wap^F between control (ds_GFP injected) and knockdown (*dsTctra* or

dsTcdsx injected) individuals (Shukla and Palli 2012a; 2012b; 2013; 2014). Surprisingly, the

expression of Tc_Wap^F was unaltered upon the knockdown of *Tctra* or *Tcdsx* suggesting *Tcdsx*

and *Tctra* doesn't have any role in regulating the female-specific expression of Tc_Wap^F. We

identified two more *waprin* genes (Tc_Wap^2 and Tc_Wap^3) of *T. castaneum*; however, they

were found to express constitutively at all the developmental stages and in both the sexes. To

the best of our knowledge this study is the first to identify and characterize the biological

significance of a sex-specific insect *waprin* gene. Besides this is the first report of presence of

multiple *waprin* genes in any insect.

3. Materials and methods:

3.1 *Tribolium castaneum* strain and rearing:

111   Wild type strain of *T. castaneum* was reared on whole-wheat flour supplemented with 5%

112   yeast in an incubator set at 30^oC temperature and 60-70% relative humidity. Sexing of pupae

113   and adults were done by observing the presence or absence of sexually dimorphic characters

114   (Shukla and Palli 2012b) with the help of Leica s9i stereomicroscope (Leica microsystems,

115   Germany).

3.2 RNA Isolation and RT-PCR:

Total RNA from different developmental stages of *Tribolium* eggs, 6th instar larvae, Q stage,

sex-specific pupae, sex-specific adults and dissected tissues (gut, ovary, head and fatbody)

from female adults were isolated using Trizol reagent (Invitrogen, USA) and were DNAse

(Promega, USA) treated. The concentrations of DNA free RNA samples were taken using

NanoDrop and the RNA samples were stored at -20°C, until its further use. RNA samples

(3µg) were incubated at 70°C for 5 min and were immediately chilled on ice. First strand

cDNA was synthesized using GoScript Reverse Transcriptase (Promega, USA) using 18-mers

polyT primer, according to the manufacturer's instructions. The prepared cDNA was diluted

10 folds with nuclease free water and used for subsequent PCR studies. GoTaq polymerase

(Promega, USA) was used for all the RT-PCR reactions using gene-specific primers

(Supplementary Table-1) which were designed using Primer-3 software (Primer3 Input

(version 0.4.0). PCR conditions were as follows- Initial denaturation at 94°C for 2 min, 32

cycles of 94°C for 30 s, annealing temperature (supplementary table 1) 30 s, 72°C for 2 min

and the final extension at 72°C for 10 min. The amplified products were electrophoresed on

1.5% agarose gels and the gels were scanned in a gel documentation system (BioRad, USA).

3.3 Sequence analysis:

After identification of Tc_Wap^F gene from our transcriptome data (unpublished), its sequence was confirmed using BLAST (BLASTn) searches in the EST database (https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE_TYPE=BlastSearch&PROG_DEF=blastn&BLAST_SPEC=GDH_GCF_000002335.3) of *T. castaneum*. Sequences of other *waprin* genes (Tc_Wap^2 and Tc_Wap^3) of *T. castaneum* were obtained by tBLASTn searches in the EST database of *T. castaneum* using Tc_Wap^F protein as a query sequence. Exons and introns of all the *Tc_waprin* genes (Tc_Wap^F, Tc_Wap^2 and Tc_Wap^3) were identified by aligning the gene-specific EST sequences with their corresponding genomic DNA sequences using Splign software (https://www.ncbi.nlm.nih.gov/sutils/splign/splign.cgi).

3.4 Double stranded RNA (dsRNA) synthesis and injections:

cDNA from adult females and gene-specific primers (supplementary Table-1) containing the T7 promoter sequences at their 5' ends were used to PCR amplify the specific regions of Tc_Wap^F, *Tcdsx* and *Tctra* genes. Purified amplicons were used as template for in-vitro transcription reaction using T7 RiboMAX™ Express RNAi System (Promega, USA) according to the manufacturer's instructions. The control dsRNA was made using an amplified fragment of the GFP gene sequence.

Newly emerged (0 day) female adults (5 h PAE) were immobilized by keeping them at 4°C for 8–10 minutes prior to injections. Unconscious adults were injected with dsGFP (control) or with dsRNA specific to one of the genes (Tc_Wap^F, *Tcdsx* or *Tctra*) using an aspirator tube assembly (Sigma) attached to a 3.50 glass capillary tube (Drummond Scientific) drawn by a micropipette puller (Narishige, PC-100). Approximately, 500–600 ng dsRNA was injected per insect. Injected insects were starved and given time to recover at room temperature for around 5 to 6 hours before transferring them to the incubator under normal rearing conditions. Knockdown efficiency of Tc_Wap^F and *Tcdsx* were analyzed after 5 days of injections whereas knockdown efficiency of *Tctra* and its effect on Tc_Wap^F was analyzed on 3 days and 5 days after injections. The ratio of gene expression between beetles injected with target dsRNA and those injected with GFP dsRNA was used to calculate the efficacy of gene knockdown in the RNAi insects.

3.5 Parental RNAi:

To see the effect of Tc_Wap^F knockdown in the next generation, control or gene specific dsRNA was injected to virgin adult females (0 day) as described in the previous section (3.4). Injected females were allowed to mate with their counterparts (uninjected males of the same age) for 6 days; one mating pair per cup was maintained. All the adults were removed from the cups and females (Tc_Wap^F or dsGFP injected) were dissected to compare their tissue morphology. Isolated tissues were examined under microscope and their images were captured. The number of eggs in each cup was counted, the development of eggs was monitored and eggs were photographed periodically.

3.6 qRT-PCR Analysis:

Total RNA isolation from control and knockdown females, and first strand cDNA synthesis was carried out as described in section 3.2 Synthesized cDNAs and gene-specific primers (Supplementary Table 1) were used in qRT-PCR reactions to determine the relative mRNA levels of Tc_wap genes using GoTaq® qPCR Master Mix (Promega) according to the manufacturer's instructions. Three independent biological replicates were used for each treatment. *T. castaneum* ribosomal protein (RPS-18) gene was used as an endogenous control to normalize the expression data (Toutges et al. 2010), and 2-ΔΔCT method was employed to assess the gene expression levels (Livak and Schmittgen 2001). The qPCR reactions were carried out in a CFX96 touch real-time PCR detection system (Bio-Rad, USA) with following cycling parameters: initial denaturation at 95 °C for 30 s, followed by 40 cycles of 95 °C for 15 s, 60 °C for 45 s and 72 °C for 5s. Melt curve analysis was carried out at 60–95°C, in steps of 0.5 °C increments per 5s. The software GraphPad Prism (9.5.1) was used to analyze statistical data. Two tailed t-test and one way ANOVA were used to compare the gene expression between different samples.

3.7 Phylogenetic Analysis and Domain Architecture:

Phylogenetic analysis of Waprin proteins of different organisms was performed using the iTOL tool which recognizes the Newick file. Predicted protein sequence of *T. castaneum*

Waprin (XP_008198656.1) was retrieved from NCBI and BLAST (blast-p) analysis was

performed in the non-redundant (nr) protein database. The obtained protein sequences

(FASTA) were aligned using MUSCLE and were further used to construct the phylogenetic

tree using the Maximum likelihood method. The FASTA sequences of Tc_Wap^2, Tc_Wap^3,

Nawaprin and Omwaprin (1,11) were also included in the analysis. Domain searches in the

Waprin proteins of different organism were conducted using Conserved domain database

(https://www.ncbi.nlm.nih.gov/cdd/) and motif scan (https://myhits.sib.swiss/cgi-

bin/motif_scan), an online software. Eight conserved cysteine residues were used as a

signature to identify the conserved domains and establish the relationship between their

associations and divergence.

4. Results:

4.1. RNA Isolation, cDNA synthesis and RT-PCR:

Transcripts of *waprin* were analyzed by semiquantitative RT-PCR at all developmental stages

(eggs, 6th instar larva, Q stage larva, female pupa, male pupa, female pupa, female adult, and

male adult) of *Tribolium* using gene-specific primers. *Tc_Wap^F* was found to express

exclusively in adult females whereas *Tc_Wap^2* and *Tc_Wap^3* express at all the developmental

stages of *Tribolium* (Figure 1A). In addition, *Tc_Wap^F* transcripts were detected in all tissues

of females, including the ovary, gut, head, and fatbody (Figure 1B). Interestingly, expression

of *Tc_Wap^F* was not observed (a faint band at 40hrs egg stage was not reproduced) at any of

the egg stages whereas the expression of other two *waprin* genes (*Tc_Wap^2* and *Tc_Wap^3*)

gradually increased with the embryonic development (Figure 1C). Consistent with the result

of semi-quantitative PCR, we observed approximately 700-fold high expression of *Tc_Wap^F*

in females compared to males in qRT-PCR analysis (Figure 1D).

4.2 Gene structure and sequence analysis:

The mRNA of *Tc_Wap^F* (XM_008200434.2) is 573 nucleotides long and harbors 2 exons and

1 intron. The ORF of *Tc_Wap^F* is 240 bases encoding 80 amino acids (aa) polypeptide. The

mRNAs of *Tc_Wap^2* (XM_964663.4) and *Tc_Wap^3* (XM_964731.3) are of 578 and 512 bases

respectively. *Tc_Wap^2* contains 3 exons and 2 introns while *Tc_Wap^3* contains 2 exons and 1

intron and their ORFs encode for 116 aa and 122 aa polypeptides respectively

(Supplementary figure 1). Eight conserved cysteine residues, a distinctive feature of the WAP
domain, were identified in all three Waprin proteins of *T. castaneum*.

4.3. Knockdown of *Tc_Wap^F* and its Phenotype:

Knockdown of *Tc_Wap^F* was confirmed through semi-quantitative RT-PCR using cDNA of
control (ds*GFP* injected) and knockdown females (ds *Tc_Wap^F* injected) (Supplementary
figure 2). Further ~12-fold reduction in the expression of *Tc_Wap^F* in knockdown females as
compared to control females was observed in qRT-PCR analysis (Figure 2A). We observed a
slightly higher number of egg laying by *Tc_Wap^F* knockdown females as compared to that of
control females (Figure 2B). However, the eggs from knockdown females failed to
develop/hatch, whereas eggs from control females developed normally and eggs were
hatched (within 65hrs) into larvae (Figure 2C and Supplementary figure 3).

4.4 Effect of knockdown of *Tcdsx* and *Tctra* genes on the expression of *Tc_Wap^F*:

A conserved *dsx*-binding motif (ACAATGT) was found in the upstream region (4.5 Kbp
upstream of the *Tc_Wap^F* transcription start site) of *Tc_Wap^F*. This prompted us to investigate
whether the sex-specific expression of *Tc_Wap^F* is regulated by the TcDsx transcription factor
of *T. castaneum*. Injection of *Tcdsx* dsRNA (ds*Tcdsx*) lead to the significant knockdown in the
expression of *Tcdsx* in both male and female adults (Figure 3A) but no change in the
expression of *Tc_Wap^F* was observed in *Tcdsx* knockdown adults (females or males) as
compared to control adults (females or males) (Figure 3B).

We further hypothesized that female-specific expression of *Tc_Wap^F* could be regulated by
some other sex-specific transcription factor (other than TcDsx) controlled by *TcTra*, a splicing
factor responsible for female specific splicing of *Tcdsx* pre-mRNA (Shukla and Palli 2012b).
Hence, knockdown of *Tctra* was done by injecting the dsRNA of *Tctra* in female *Tribolium*
adults and the expression of *Tc_Wap^F* was compared between control and *Tctra* knockdown
females. Both female and male isoforms of *Tctra* were obtained in RT-PCR of *Tctra*
Knockdown females (Supplementary figure 4), suggesting a successful knockdown of *Tctra*
in the ds*Tctra* injected females (Shukla and Palli 2014). Surprisingly, there was no effect in
the expression of *Tc_Wap^F* in *Tctra* knockdown females; neither at 3 nor at 5 days after
ds*Tctra* injections (Figure 3C).

4.5 Phylogenetic Analysis and Domain Architecture:

A total of 65 Wap and Wap related protein sequences from seven different orders of class
insecta and six outgroups (including bacteria, crustaceans, scorpions, reptiles, amphibians and
mammals) were retrieved using the BLASTp tool. The phylogenetic tree generated using the
Maximum likelihood method grouped the proteins into three clusters. The relationships
between all three Waprin of *Tribolium* (Tc_Wap[F], Tc_Wap[2], Tc_Wap[3]) was established; the
three Wap diverged into different sister branches despite belonging to the same parent branch.
Tc_Wap[F] was closely related to waprin of *Tribolium madens* while *Sandaracinus*
amylolyticus and Tc_Wap[2] have similar protein sequences and were closely related. Similarly,
Tc_Wap[3] has a protein sequence similar to that of *Hermetia illucens*. Figure 4 depicts that
Tc_Wap[F] and Tc_Wap[3] are distantly related to each other and Tc_Wap[2] lies intermediate to
both, Tc_Wap[F] and Tc_Wap[3] (Figure 4).

The hallmark of Waprin and Waprin-related proteins throughout the animal kingdom is the
presence of eight conserved cysteine residues that produce four disulfide linkages along with
the WAP domain (Alam et al. 2019; Suntravat et al. 2016). Domain architecture analysis
using TB tools predicted the presence of WAP domains across protein sequences collected
from seven different orders of insects. Analysis of the domain organization of all the
predicted WAP proteins across major orders and representative species showed that the WAP
domain is conserved across all members. WAP protein of few species were found to possess
multiple WAP domains as well as additional domains, other than the WAP domain. Tc_Wap[F]
showed the presence of single WAP domain while Tc_Wap[2] consists of two WAP domains.
Although Tc_Wap[3] contains eight conserved cysteine residues, this tool indicated the absence
of the WAP domain in Tc_Wap[3]. Kunitz domain, a serine protease inhibitor found in the
venom of several animals including snakes, lizards, spiders, ticks, and sea anemones is the
second most conserved domain across 65 species (Figure 5).

5. Discussion:

Waprin (WAP domain containing proteins) are found in diverse groups of organisms (Alam et
al. 2019) and are known to possess antiviral (Chen, He, and Xu 2008) , antimicrobial(Ali et
al. 2002; Hagiwara et al. 2003; Hauton, Brockton, and Smith 2006; Smith et al. 2008;
Imjongjirak et al. 2009; Li et al. 2013), Na-K ATPase inhibitory, protein and proteinase
inhibitory(Ali et al. 2002; Li et al. 2013), as well as immunomodulatory functions (Gao et al.
2021). Waprin protein has been reported from a few insect species also for e.g., wood-boring
woodwasp, *S. nitobei* (Gao et al. 2021), guinea ant, *T. bicarinatum*(Bouzid et al. 2014) and

honeybees (*A. millefera*). Besides the microbicidal and anti-elastolytic activities of Amwaprin (Lee et al. 2023), a recently discovered Waprin protein from *A. melifera* not much is known about the biological significance and evolution of Waprin proteins in insect kingdom.

We identified three Waprin proteins from the red flour beetle, *T. castaneum*. Most importantly, one of the *Tribolium* Waprin proteins (Tc_Wap^F) was found to express only in adult females. Since Tc_Wap^F has the typical feature of Waprin protein i.e., the presence of eight conserved cysteine residues that produce four disulfide linkages along with the WAP domain (Alam et al. 2019; Suntravat et al. 2016) we assumed its functional conservation. To deduce the biological relevance of *Tc_Wap^F*, RNAi mediated knockdown of *Tc_Wap^F* was done in adult females. Despite the significant knockdown in the expression of *Tc_Wap^F* no observable change was noticed in the morphology or anatomy of the knockdown individuals. Considering the antiviral and antimicrobial activities of Waprin protein one can expect enhanced microbial infections in the *Tc_Wap^F* knockdown individuals but since our *Tribolium* cultures were maintained in a clean and optimal environmental condition we failed to notice any such change in the *Tc_Wap^F* knockdown females. Eggs are known to possess their own defence systems against microbial infection (Gantois et al. 2009), we further questioned if *Tc_Wap^F* acts as a line of defense for the eggs laid by *Tribolium* female adults. For this the development of eggs laid by *Tc_Wap^F* knockdown and control females was tracked and compared. Slightly higher number of eggs were laid by *Tc_Wap^F* knockdown females as compared to that of control females. At this moment we don't know the reason for the same, but it is possible that there is a tradeoff between female fecundity and immunity provided by Tc_Wap^F protein. Interestingly, the eggs laid by *Tc_Wap^F* knockdown females failed to hatch and started desiccating after 60-65hrs (normal hatching time of *Tribolium* eggs). Together, the female specific expression of *Tc_Wap^F* (in adults), no expression of *Tc_Wap^F* in *Tribolium* eggs and failure of hatching of eggs laid by knockdown females suggests the maternal transfer of Tc_Wap^F proteins to shield the embryos from environmental pathogens and is crucial for embryonic development. The schematics shown in figure 6 depicts the mode of action of *Tc_Wap^F* in *T. castaneum*. Constitutive expression of other two *waprin* genes (*Tc_Wap^2* and *Tc_Wap^3*) might be involved in protecting all the *Tribolium* stages from the environmental pathogens.

The central axis (*tra-dsx*) of insect sex determination cascade is conserved in *T. castaneum* also (Shukla and Palli 2012b). Sex-specifically spliced *Tcdsx* produces sex-specific proteins (sex-specific transcription factors- TcDsx^m in males and TcDsx^f in females) in the two sexes

which is known to regulate all the genes expressed in a sex-specific manner (Shukla and Palli

2012a). To explore whether the sex-specific expression of Tc_Wap^F gene is regulated by

female-specific TcDsx, we examined the presence of TcDsx binding sites in upstream region

of Tc_Wap^F gene. Even though the upstream region (4.5 Kbp upstream of the Tc_Wap^F

transcription start site) of Tc_Wap^F harbors a conserved Dsx-binding motif (ACAATGT),

neither the decreased expression of Tc_Wap^F was found in $Tcdsx$ knockdown females nor

there was increased expression of Tc_Wap^F in $Tcdsx$ knockdown males suggesting that TcDsx

is not involved in regulating the female specific expression of Tc_Wap^F.

We further hypothesized that the transcription of Tc_Wap^F might be regulated by another (yet

to discover) transcription factor regulated by TcTra, a splicing regulator of $Tcdsx$. To rule out

this possibility, we performed the knockdown of $Tctra$ in female adults but still no change in

the expression of Tc_Wap^F was noticed in $Tctra$ knockdown females. Hence, we conclude that

the sex-specific expression of Tc_Wap^F is not under the control of the established sex

determination pathway of $T.\ castaneum$. It is possible that the sex-specific expression of

Tc_Wap^F is regulated by a pathway bifurcated at the most upstream gene (yet to be

discovered) of $Tribolium$ sex determination cascade. Alternatively, we also presume the

involvement of chromatin remodelling genes in the female-specific expression of Tc_Wap^F.

Our hypothesis is in line with a recent report by Nanni et al., 2023 which suggests the

association of sex-biased expression of genes with differential state (open or closed state) of

chromatin in the two sexes of $Drosophila\ melanogaster$ and $D.\ simulans$ (Nanni et al. 2023).

Therefore, the sex-specific expression of Tc_Wap^F gene could be because of open chromatin

in female and closed chromatin in males, in $T.\ castaneum$.

In conclusion, we identified Tc_Wap^F, a distinct member of WAP family. To our knowledge,

this study represents the first report of identification of sex-specific $waprin$ from any insect

species besides deducing its biological function in the development of $Tribolium$.

Interestingly, Tc_Wap^F genes has the potential to be used in dsRNA mediated pesticide. It will

be interesting to study the comparative microbicidal properties of Tc_Wap genes in future.

The properties like anti-microbial can be exploited further to develop new strategies to

counteract the antimicrobial resistance which will add value to the pharmacological

applications.

6. Author's Contribution:

CC, VV and JNS conceived the idea of manuscript. CC performed experiments of knockdown, PCR and qRT-PCR. SS assisted CC in experiments and manuscript writing. DK did the phylogenetic analysis and domain search. KKM helped with all the figures and tables in the manuscript besides maintaining the insect cultures. CC and JNS wrote the manuscript and all others agreed to it. All authors contributed to the article and approved the submitted version.

7. Acknowledgements:

J N Shukla acknowledges Ramalingaswami grant (BT/RLF/ Re-entry/10/2015) from Department of Biotechnology, New Delhi and EMR grant (EMR/2017/001378/AS) from DST-SERB, New Delhi for their generous support. Sincere acknowledgement to the Department of Biotechnology and Department of Sport Sciences, Central University of Rajasthan for providing the infrastructural facilities to carry out this work. VV is a Ramalingaswami fellow and, CC and KKM are the Ph.D. students at CURAJ. SS and DK were M.Sc. Biotechnology students of CURAJ. JNS also acknowledges Dr. Satnam Singh (Punjab Agricultural University, Ludhiana) and Dr. Deepa Agashe (National Centre for Biological Sciences, Bangalore) for providing us the *Tribolium castaneum* culture.

8. References:

Alam, Md. Iqbal, Ojha R, Alam MA, Quasimi H, and Alam O. 2019. "Therapeutic Potential of Snake Venoms as Antimicrobial Agents." *Frontiers in Drug, Chemistry and Clinical Research* 2 (4). https://doi.org/10.15761/fdccr.1000136.

Ali, Mohamed F, Karen R Lips, Floyd C Knoop, Bernd Fritzsch, Caroline Miller, and J Michael Conlon. 2002. "Antimicrobial Peptides and Protease Inhibitors in the Skin Secretions of the Crawfish Frog, Rana Areolata." *Biochimica et Biophysica Acta (BBA)-Proteins and Proteomics* 1601 (1): 55–63.

Bouzid, Wafa, Marion Verdenaud, Christophe Klopp, Frédéric Ducancel, Céline Noirot, and Angélique Vétillard. 2014. "De Novo Sequencing and Transcriptome Analysis for Tetramorium Bicarinatum: A Comprehensive Venom Gland Transcriptome Analysis

from an Ant Species." *BMC Genomics* 15 (1). https://doi.org/10.1186/1471-2164-15-
987.

Campbell, James F, Christos G Athanassiou, David W Hagstrum, and Kun Yan Zhu. 2021.
"Annual Review of Entomology Tribolium Castaneum: A Model Insect for Fundamental
and Applied Research." https://doi.org/10.1146/annurev-ento-080921.

Chen, Dandan, Nanhai He, and Xun Xu. 2008. "Mj-DWD, a Double WAP Domain-
Containing Protein with Antiviral Relevance in Marsupenaeus Japonicus." *Fish and*
Shellfish Immunology 25 (6): 775–81. https://doi.org/10.1016/j.fsi.2008.02.017.

Coutts, M P. 1969a. "The Mechanism of Pathogenicity of Sirex Noctilio on Pinus Radiata I.
Effects of the Symbiotic Fungus Amylostereum Sp. (Thelophoraceae)." *Australian*
Journal of Biological Sciences 22 (4): 915–24. https://doi.org/10.1071/BI9690915.

Coutts, M P. 1969b. "The Mechanism of Pathogenicity of Sirex Noctilio on Pinus Radiata II.
Effects of S. Noctilio Mucus." *Australian Journal of Biological Sciences* 22 (5): 1153–
62. https://doi.org/10.1071/BI9691153.

Gantois, Inne, Richard Ducatelle, Frank Pasmans, Freddy Haesebrouck, Richard Gast, Tom J
Humphrey, and Filip Van Immerseel. 2009. "Mechanisms of Egg Contamination by
Salmonella   Enteritidis."   *FEMS   Microbiology   Reviews*   33   (4):   718–38.
https://doi.org/10.1111/j.1574-6976.2008.00161.x.

Gao, Chenglong, Lili Ren, Ming Wang, Zhengtong Wang, Ningning Fu, Huiying Wang,
Xiaochen Wang, et al. 2021. "Proteo-Transcriptomic Characterization of Sirex Nitobei
(Hymenoptera: Siricidae)   Venom."   *Toxins*   13   (8).
https://doi.org/10.3390/toxins13080562.

Hagiwara, Koichi, Tohru Kikuchi, Yoshiyuki Endo, Huqun, Kazuhiro Usui, Mitsu Takahashi,
Naoko Shibata, et al. 2003. "Mouse SWAM1 and SWAM2 Are Antibacterial Proteins
Composed of a Single Whey Acidic Protein Motif." *The Journal of Immunology* 170 (4):
1973–79. https://doi.org/10.4049/jimmunol.170.4.1973.

Hauton, C., V. Brockton, and V. J. Smith. 2006. "Cloning of a Crustin-like, Single Whey-
Acidic-Domain, Antibacterial Peptide from the Haemocytes of the European Lobster,
Homarus Gammarus, and Its Response to Infection with Bacteria." *Molecular*
Immunology 43 (9): 1490–96. https://doi.org/10.1016/j.molimm.2005.07.029.

Imjongjirak, Chanprapa, Piti Amparyup, Anchalee Tassanakajon, and Siriporn Sittipraneed. 2009. "Molecular Cloning and Characterization of Crustin from Mud Crab Scylla Paramamosain." *Molecular Biology Reports* 36 (5): 841–50. https://doi.org/10.1007/s11033-008-9253-0.

Klompen, Anna M.L., Jason Macrander, Adam M. Reitzel, and Sérgio N. Stampar. 2020. "Transcriptomic Analysis of Four Cerianthid (Cnidaria, Ceriantharia) Venoms." *Marine Drugs* 18 (8). https://doi.org/10.3390/MD18080413.

Lee, Kwang Sik, Bo Yeon Kim, Yun Hui Kim, Yong Soo Choi, and Byung Rae Jin. 2023. "Identification of Waprin and Its Microbicidal Activity: A Novel Protein Component of Honeybee (Apis Mellifera) Venom." *Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology* 266 (April). https://doi.org/10.1016/j.cbpc.2023.109561.

Li, Shuang, Xing Kun Jin, Xiao Nv Guo, Ai Qing Yu, Min Hao Wu, Shang Jian Tan, You Ting Zhu, Wei Wei Li, and Qun Wang. 2013. "A Double WAP Domain-Containing Protein Es-DWD1 from Eriocheir Sinensis Exhibits Antimicrobial and Proteinase Inhibitory Activities." *PLoS ONE* 8 (8). https://doi.org/10.1371/journal.pone.0073563.

Liu, Daixi, Yuwei Wang, Lin Wei, Huahu Ye, Huan Liu, Ling Wang, Rui Liu, Dongsheng Li, and Ren Lai. 2013. "Snake Venom-like Waprin from the Frog of Ceratophrys Calcarata Contains Antimicrobial Function." *Gene* 514 (2): 99–104. https://doi.org/10.1016/j.gene.2012.11.007.

Livak, Kenneth J., and Thomas D. Schmittgen. 2001. "Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2-ΔΔCT Method." *Methods* 25 (4): 402–8. https://doi.org/10.1006/meth.2001.1262.

Nair, Dileep G., Bryan G. Fry, Paul Alewood, Prakash P. Kumar, and R. Manjunatha Kini. 2007. "Antimicrobial Activity of Omwaprin, a New Member of the Waprin Family of Snake Venom Proteins." *Biochemical Journal* 402 (1): 93–104. https://doi.org/10.1042/BJ20060318.

Nanni, Adalena V, Natalie Martinez, Rita Graze, Alison Morse, Jeremy R B Newman, Vaibhav Jain, Srna Vlaho, et al. 2023. "Sex-Biased Expression Is Associated With Chromatin State in Drosophila Melanogaster and Drosophila Simulans." *Molecular Biology and Evolution* 40 (5): msad078. https://doi.org/10.1093/molbev/msad078.

Pierre, L. St., S. T. Earl, I. Filippovich, N. Sorokina, P. P. Masci, J. De Jersey, and M. F. Lavin. 2008. "Common Evolution of Waprin and Kunitz-like Toxin Families in Australian Venomous Snakes." *Cellular and Molecular Life Sciences* 65 (24): 4039–54. https://doi.org/10.1007/s00018-008-8573-5.

Reeks, T. A., B. G. Fry, and P. F. Alewood. 2015. "Privileged Frameworks from Snake Venom." *Cellular and Molecular Life Sciences* 72 (10): 1939–58. https://doi.org/10.1007/s00018-015-1844-z.

Romero-Gutiérrez, Maria Teresa, Carlos Eduardo Santibáñez-López, Juana María Jiménez-Vargas, Cesar Vicente Ferreira Batista, Ernesto Ortiz, and Lourival Domingos Possani. 2018. "Transcriptomic and Proteomic Analyses Reveal the Diversity of Venom Components from the Vaejovid Scorpion Serradigitus Gertschi." *Toxins* 10 (9). https://doi.org/10.3390/toxins10090359.

Romero-Gutierrez, Teresa, Esteban Peguero-Sanchez, Miguel A. Cevallos, Cesar V.F. Batista, Ernesto Ortiz, and Lourival D. Possani. 2017. "A Deeper Examination of Thorellius Atrox Scorpion Venom Components with Omic Techonologies." *Toxins* 9 (12). https://doi.org/10.3390/toxins9120399.

JN Shukla, and Subba Reddy Palli. 2012a. "Doublesex Target Genes in the Red Flour Beetle, Tribolium Castaneum." *Scientific Reports* 2. https://doi.org/10.1038/srep00948.

JN Shukla, and Subba Reddy Palli. 2012b. "Sex Determination in Beetles: Production of All Male Progeny by Parental RNAi Knockdown of Transformer." *Scientific Reports* 2. https://doi.org/10.1038/srep00602.

JN Shukla, and Subba Reddy Palli. 2013. "Tribolium Castaneum Transformer-2 Regulates Sex Determination and Development in Both Males and Females." *Insect Biochemistry and Molecular Biology* 43 (12): 1125–32. https://doi.org/10.1016/j.ibmb.2013.08.010.

JN Shukla, and Subba Reddy Palli. 2014. "Production of All Female Progeny: Evidence for the Presence of the Male Sex Determination Factor on the y Chromosome." *Journal of Experimental Biology* 217 (10): 1653–55. https://doi.org/10.1242/jeb.100438.

Smith, Valerie J., Jorge M.O. Fernandes, Graham D. Kemp, and Chris Hauton. 2008. "Crustins: Enigmatic WAP Domain-Containing Antibacterial Proteins from

Crustaceans." *Developmental and Comparative Immunology.*
https://doi.org/10.1016/j.dci.2007.12.002.

Suntravat, Montamas, Néstor L. Uzcategui, Chairat Atphaisit, Thomas J. Helmke, Sara E.
Lucena, Elda E. Sánchez, and Alexis Rodríguez Acosta. 2016. "Gene Expression
Profiling of the Venomgland from the Venezuelan Mapanare (Bothrops Colombiensis)
Using Expressed Sequence Tags (ESTs)." *BMC Molecular Biology* 17 (1).
https://doi.org/10.1186/s12867-016-0059-7.

Thankappan, Bency, and Jayaraman Angayarkanni. 2019. "Biological Characterization of
Omw1 and Omw2: Antimicrobial Peptides Derived from Omwaprin." *3 Biotech* 9 (8).
https://doi.org/10.1007/s13205-019-1801-x.

Torres, Allan M., Hui Y. Wong, Malan Desai, Shabbir Moochhala, Philip W. Kuchel, and R.
Manjunatha Kini. 2003. "Identification of a Novel Family of Proteins in Snake Venoms.
Purification and Structural Characterization of Nawaprin from Naja Nigricollis Snake
Venom." *Journal of Biological Chemistry* 278 (41): 40097–104.
https://doi.org/10.1074/jbc.M305322200.

Toutges, Michelle J., Kris Hartzer, Jeffrey Lord, and Brenda Oppert. 2010. "Evaluation of
Reference Genes for Quantitative Polymerase Chain Reaction across Life Cycle Stages
and Tissue Types of Tribolium Castaneum." *Journal of Agricultural and Food
Chemistry* 58 (16): 8948–51. https://doi.org/10.1021/jf101603j.

Zhang, Xiaoming, Raymond P. Boot-Handford, Julie Huxley-Jones, Lorna N. Forse, A. Paul
Mould, David L. Robertson, Li Li, Matthews Athiyal, and Michael P. Sarras. 2007. "The
Collagens of Hydra Provide Insight into the Evolution of Metazoan Extracellular
Matrices." *Journal of Biological Chemistry* 282 (9): 6792–6802.
https://doi.org/10.1074/jbc.M607528200.

Figure Legends:

Figure 1: Temporal and spatial expression profiles of *waprin* genes (Tc_Wap^F ,Tc_Wap^2 and
Tc_Wap^3) of *T. castaneum*.

A) Developmental profiling of *waprin* genes of *T. castaneum*. Semiquantitative RT-PCR was
performed using stage specific cDNAs (6th Instar Larvae, Q Stage, Larvae, female pupae, male
pupae, female adult and male adults) as templates. Tc_Wap^F shows female exclusive expression
while the other two *waprin* genes (Tc_Wap^2 and Tc_Wap^3) express at all the stages. B) Tissue

specific expression profile of Tc_Wap^F; Tc_Wap^F expresses in all the tissues of adult female. C) Expression profile of *waprin* genes in developing embryos (from 2 hours to 60 hours after egg laying). RPS-18 gene was used as an endogenous control. D) Relative expression of Tc_Wap^F in 6th day old adult male and female. qRT-PCR analysis was performed using the sex specific cDNA's as templates. Error bars represent the standard deviation (p value=0.0035).

Figure 2A: qRT-PCR analysis to check the expression of Tc_Wap^F in the knockdown females and control females. The templates used were cDNA of control (*ds_GFP* injected) and knockdown (Tc_Wap^F injected) females. Error bars represent s.d. **p value≤ 0.0035.

Figure 2B: Graphical representation of number of individuals injected and number of individuals survived after dsRNA injection versus the number of eggs laid by the survived females. A slight increase in egg laying was observed by Tc_Wap^F knockdown females as comparison to control females. (C1-C4: Represents control (ds_GFP injected) females; W1-W4: Represents knockdown (Tc_Wap^F) females.)

Figure 2C: Picture of eggs laid by control and Tc_Wap^F knockdown females.

Figure 3A: Graphs depicting the knockdown efficiency of *Tcdsx*. dsRNA injections targeting common region of *Tcdsx* were performed separately in female and male adults (0 day). *p value≤ 0.0311; **p value ≤ 0.0003

Figure 3B: Relative expression of Tc_Wap^F in response to *Tcdsx* knockdown, in female and male individuals. Five biological replicates were used in each experiment. Error bars represent standard deviation.

Figure 3C: Graphical representation of the effect of Tc_Tra knockdown on the expression of Tc_Wap^F. The knockdown efficiency of Tc_Tra and the expression of Tc_Wap^F was analysed both at 3[rd] day and 5[th] day after ds*Tctra* injections.

Figure 4: Phylogenetic analysis of WAP proteins across seven different orders of class 'insecta'. The circular phylogenetic tree was constructed with the MEGA11 program using the neighbour-joining method.

Figure 5: Domain architecture of various WAP domain containing proteins across different insect species. Different domains are colour coded and mentioned in the right side of the picture.

Figure 6: Schematics showing the mode of action of Tc_Wap^F. Tc_Wap^F is maternally transferred to next generation. But when Tc_Wap^F is silenced through dsRNA mediated knockdown, the maternal transcripts get degraded and therefore the embryos failed to hatch.

A)

B)

C)

Expression of *Tc_Wap^F*

D)

A)

Tc_Wap^F Knockdown

B)

Graphical representation of survival and fecundity between Control and Wap Knockdown adults

■ No. of injected female adults ■ No.of survived adults — No. of eggs laid(Fecundity)

C)

Eggs laid by ds_GFP injected females

| 0 hours egg stage | 15 hours egg stage | 30 hours egg stage | 60 hours egg stage |

Eggs laid by ds_ Tc_Wap^F knockdown females

| 0 hours egg stage | 15 hours egg stage | 30 hours egg stage | 60 hours egg stage |

A)

Knockdown efficiency of
Tcdsx

B)

Effect of *Tcdsx* knockdown on
expression of *Tc_Wap^F*

C) Effect of *Tctra* knockdown on
expression of *Tc_Wap^F*

Parental RNAi of
Tc_Wap^F

Degradation of maternally
supplied *Tc_Wap*^F

259 b.p.   51b.p.   317 b.p.

Degraded Tc_Wap^F *transcripts*

Transfer of *Tc_Wap*^F to
the embryos from mother.

Eggs fail to hatch

Development of
eggs into larvae.