Membrane contact sites regulate vacuolar fission via sphingolipid metabolism

Kazuki Hanaoka[1,†], Kensuke Nishikawa[1,†], Atsuko Ikeda[1,†], Philipp Schlarmann[1], Sayumi Yamashita[2], Aya Nakaji[2], Sotaro Fujii[1] and Kouichi Funato[1,2*]

[1]Graduate School of Integrated Sciences for Life, Hiroshima University, Kagamiyama 1-4-4, Higashi-Hiroshima 739-8528, Japan.

[2]School of Applied Biological Science, Hiroshima University, Kagamiyama 1-4-4, Higashi-Hiroshima 739-8528, Japan.

[†]These authors contributed equally to this work

* Corresponding author; Kouichi Funato

Email: kfunato@hiroshima-u.ac.jp

Abstract

Membrane contact sites (MCSs) are junctures that perform important roles including coordinating lipid metabolism. Previous studies have indicated that vacuolar fission/fusion processes are coupled with modifications in the membrane lipid composition. However, it has been still unclear whether MCS-mediated lipid metabolism controls the vacuolar morphology. Here we report that deletion of tricalbins (Tcb1, Tcb2, Tcb3), tethering proteins at endoplasmic reticulum (ER)-plasma membrane (PM) and ER-Golgi contact sites, alters fusion/fission dynamics and causes vacuolar fragmentation in the yeast *Saccharomyces cerevisiae*. In addition, we show that the sphingolipid precursor phytosphingosine accumulates in tricalbin-deleted cells, triggering the vacuolar division. Detachment of the nucleus vacuole junction (NVJ), an important contact site between the vacuole and the perinuclear ER, restored vacuolar morphology in both cells subjected to high exogenous phytosphingosine and Tcb3-deleted cells, supporting that phytosphingosine transport across the NVJ induces vacuole division. Thus, our results suggest that vacuolar morphology is maintained by MCSs through the metabolism of sphingolipids.

Introduction

Cellular organelles are formed by membranes with unique lipid compositions, morphology, and functions. The vacuole of budding yeast is an organelle that shares many similarities with mammalian lysosomes and plant vacuoles. Vacuoles possess degradative and storage capacities, and are essential for maintaining cellular homeostasis, including maintaining pH or ion homeostasis, cellular detoxification, and responses to osmotic shock and nutrient environments. Vacuoles are liable to change their morphology (size, volume, and number) through cycles of fusion and fission, to adapt to the intracellular and extracellular environments or upon vacuole inheritance to daughter cells. The steady-state morphology of the vacuole is maintained by a balance of constitutive vacuolar fusion and fission processes [1] [2].

The homotypic vacuolar membrane fusion process occurs in four stages: priming, tethering, docking and fusion [3] [4] [5] [6]. Each stage is defined as follows. "Priming" In the priming reaction, inactive fusion factors that are still assembled from previous fusion events are recycled. The yeast N-ethylmaleimide-sensitive factor (NSF), Sec18p, is activated by ATP and releases the NSF attachment protein (SNAP), Sec17p, from the protease complex, thereby activating the SNAP receptor (SNARE) molecule [3] [4] [7]. "Tethering" The HOPS/Class C Vps complex and the Rab-GTPase Ypt7p mediate the reversible contact of opposing vacuoles [4] [8]. "Docking" The coupling of three Q-SNAREs (Vam3p, Vam7p, and Vti1p) with an opposing R-SNARE (Nyv1p) to form a trans-SNARE complex between the two corresponding vacuoles. "Fusion" The final stage in which the lipid bilayer fuses and the contents of the lumen are mixed [5] [7]. For stable vacuolar fusion, several factors are required such as vacuolar acidification by V-ATPase proton pump activity [1] [9] [10] [11], ions [12] and ion transporters [13].

Furthermore, vacuolar fission is required for proper vacuolar inheritance during mitosis and acute response to osmotic shock in yeast, however the molecular mechanism of fission is not fully understood, and limited knowledge is available. Vacuolar fission is performed in two steps that require proteins and lipids [14]. The first step is the contraction and invagination of the vacuolar membrane through the involvement of Vps1p, a dynamin-like GTPase [15] and V-ATPase, which drives the proton gradient [1]. Next, vacuolar fission is promoted by Vac14p, Vac7p, and Fab1p [16], which are required for the generation of PtdIns[3,5]P2, and Atg18p, an effector of Fab1p and a sensor of PtdIns[3,5]P2 levels [17] [18]. Recently, it has also been shown that vacuolar fission is associated with nutrient status and responds to ER stress via TORC1 targets [19] [20]. Moreover, it has been proposed that vacuolar fusion and fission dynamics are regulated by the Yck3p-Env7p kinase cascade, which maintains an equilibrium between fusion and fission activities [21].

Organelles construct a dynamic network by associating with each other through membrane-contact sites (MCSs), which play an important role in lipid transport and metabolism

[22] [23]. Lipids (phosphatidic acid (PA), diacylglycerol, ergosterol, phosphatidylinositol (PtdIns)
phosphate (PtdInsP)) and lipid raft domains have been shown to be important as fusion and fission
effectors and discussed in a variety of recent reviews [2] [17] [24] [25] [26] [27]. However, it remains
unclear if MCS-mediated lipid metabolism controls the morphology of the vacuole. Vacuoles have
contact sites with other organelles. The MCS where the nuclear ER meets the vacuole is called the
nucleus-vacuole junction (NVJ), and the ER-localized Nvj1p and vacuole-membrane-localized
Vac8p act as tethers to form the NVJ [28]. Vacuoles also form MCS with mitochondria called
vacuole and mitochondria patches (vCLAMP). vCLAMP is mediated by the soluble (cytosol-
localized) HOPS complex subunit Vps39p, the vacuole-localized small Rab-GTPase Ypt7p, and by
an unknown mitochondrial factor [29] [30]. The translocase of the mitochondrial outer membrane
(TOM) subunit Tom40p has been identified as a direct binding partner of Vps39p, suggesting a
mechanism for the formation of vCLAMP by the Ypt7p-Vps39p-Tom40p tether [31]. Although it has
been suggested that MCSs with vacuoles are involved in the transport of lipids such as sterols and
precursors of sphingolipids that are involved in vacuolar morphology [32] [33] [34] [35], the roles of
those MCSs-mediated lipid metabolisms in the regulation of vacuole morphology are not
understood.

In this study, we found that tether proteins of MCS formation are involved in the regulation
of vacuolar morphology. We show that the deletion of tricalbins, tethers of the MCS between the
ER and the plasma membrane (PM) [36] and between the ER and the Golgi apparatus [37], affects
vacuolar division. Phytosphingosine, a precursor of ceramide, accumulated in the tricalbin-deleted
strain and was revealed as the underlying cause that triggers vacuolar fission. Moreover, deletion
of key tethering proteins at the NVJ recovered vacuolar morphology of cells subjected to high
exogenous phytosphingosine and tricalbin-deleted cells, indicating that the NVJ is required for
phytosphingosine-induced vacuolar fission. In summary, we propose that vacuolar morphology is
regulated by MCSs through regulation of sphingolipid metabolic pathways.

Results

Deletion of tricalbins causes vacuole fragmentation

Tricalbins (Tcb1p, Tcb2p, and Tcb3p) are ER membrane tethering proteins that connect the cortical ER (cER) with the PM [38], and contribute to PI4P turnover [36], sterol flux and transport [39] and maintain other lipid homeostasis [40]. It is also suggested that tricalbins create ER membrane curvature to maintain PM integrity [41] [42]. Tricalbins also localize to the ER-Golgi contacts and are responsible for the non-vesicular transport of ceramide from ER to Golgi apparatus [37]. However, their role in other organelle morphology and function remains unknown. Interestingly, the *NVJ1* gene, which encodes a major component of the NVJ, showed a negative synthetic interaction with the absence of all three tricalbins (*tcb1Δ2Δ3Δ*) [41]. Moreover, HOPS subunit Vam6p and Rab-GTPase Ypt7p, which are both involved in vacuolar fusion, also showed negative synthetic interaction with *tcb1Δ2Δ3Δ* [41]. Based on the findings, we assumed that tricalbins may be involved in the regulation of vacuolar morphology or function linked with the NVJ. To investigate the role of tricalbins in vacuole morphology, we analyzed the number of vacuoles per cell by a fluorescent probe FM4-64 that selectively stains yeast vacuolar membranes. We observed that compared to wild type cells, the *tcb1Δ2Δ3Δ* strain showed a phenotype characterized by a decreased percentage of cells with one vacuole and an increased percentage of cells with two or more vacuoles (Fig 1A). We refer to this phenotype as vacuolar fragmentation. In addition, analysis with single- and double deletion strains revealed that single deletion of *TCB1* or *TCB3* already exhibited strong vacuole fragmentation (Fig 1B). These results indicate that tricalbins are important to maintain vacuole morphology.

We next examined whether deletion of tricalbins affects vacuolar acidification. As homotypic vacuole-vacuole membrane fusion requires the vacuolar H^+-ATPase (V-ATPase) [11] [43] [44], tricalbins could be indirectly involved in maintaining vacuole morphology through V-ATPase. Therefore, we addressed whether tricalbin mutant strains could grow in alkaline media, because V-ATPase–deficient mutants can grow in acidic media but fail to grow in alkaline media [43]. As a negative control, we used *vma3Δ* mutant cells in which one of the V0 subunits of V-ATPase had been deleted. The *vma3Δ* mutant cells grew in acidic medium (pH 5.0), but failed to grow in alkaline medium (pH 7.5) (Fig S1A). In contrast, all tricalbin mutant strains grew like wild type in both conditions, suggesting that the V-ATPase in these mutants remains functionally normal. Thus, these results suggest that vacuole fragmentation caused by tricalbin deletion is not due to a loss of function of V-ATPase.

Furthermore, we investigated the effects of tricalbin deletion on delivery of a vacuolar protease, carboxypeptidase S (Cps1p) that is sorted into the vacuole lumen upon endosome–vacuole fusion. Deletion of HOPS complex components that mediate endosome–vacuole fusion

results in vacuole fragmentation [45]. To test the possibility that vacuole fragmentation in tricalbin

mutant cells is caused by a defect of endosome–vacuole fusion, we analyzed the localization of

GFP-Cps1p in tricalbin mutant cells. In both of WT and *tcb3Δ* cells, GFP-Cps1p was observed in

the vacuole lumen (Fig S1B), suggesting that the tricalbin mutant exhibits a normal delivery of

vacuolar proteins via endosomes to the vacuole. In addition, the fact that tricalbin deletion does not

affect the maturation of CPY, a vacuolar hydrolase carboxypeptidase Y [37], indicates that the

vacuolar degradation ability as well as protein delivery to the vacuole is normal. Taken together,

tricalbins are required for maintaining vacuole morphology, but not due to an indirect action through

vacuolar acidification or endosome-vacuole fusion.

Tricalbins negatively regulate vacuole fission in a parallel pathway with TORC1

The number and size of vacuoles within a cell are regulated by coordinated cycles of

vacuolar fission and fusion. To address whether vacuole fragmentation in the tricalbin mutant is

caused by facilitation of vacuole fission or an impaired homotypic vacuole fusion, we examined the

effect of hypotonic stress on vacuole fragmentation in *tcb1Δ2Δ3Δ* mutant cells, because low-

osmotic stimuli such as water promote homotypic vacuole fusion to maintain cytoplasmic osmolarity

[46]. We observed that addition of water to *tcb1Δ2Δ3Δ* cells partially restored vacuole

fragmentation, suggesting that vacuolar fusion machinery is functional in *tcb1Δ2Δ3Δ* cells (Fig 1C).

This result suggests that loss of tricalbins alters fusion-fission dynamics by primarily affecting the

fission machinery rather than fusion.

As the target of rapamycin complex 1 (TORC1) is a positive regulator of vacuole fission in

response to hyperosmotic shock and ER stress [19] [20], we next examined if rapamycin, which is

an inhibitor of TORC1, affects the vacuole fragmentation in *tcb1Δ2Δ3Δ* cells. As shown in Figure

1C, we observed that the vacuole fragmentation in *tcb1Δ2Δ3Δ* cells was suppressed by rapamycin,

suggesting that TORC1 may be required for tricalbin deletion-induced vacuolar fragmentation. We

attempted to characterize further the relationship between tricalbin and TORC1, and showed that

tcb1Δ2Δ3Δ cells had no significant effect on phosphorylation levels of Sch9p, a major downstream

effector of TORC1 (Fig 1D). These results suggest that deletion of tricalbins does not activate

TORC1. Therefore, we conclude that tricalbins and TORC1 act in parallel and opposite ways to

regulate vacuole fission (Fig 1E).

The TM domain of Tcb3 contributes to mediating protein interactions between the tricalbin

family to maintain vacuolar morphology

Tcb3p possesses an N-terminal TM domain, a central SMP domain, and multiple C-terminal C2

domains (Fig 2A). To identify which domain of Tcb3p is essential for regulating vacuole fission, we

replaced the C-terminal cytoplasmatic sequence of Tcb3p with a GFP binding protein (GBP). This

way, we created three strains expressing either full-length Tcb3-GBP; Tcb3(Full)-GBP, Tcb3-GBP
lacking C2 domains; Tcb3(TM-SMP)-GBP or Tcb3-GBP lacking both SMP and C2 domains;
Tcb3(TM)-GBP. Vacuole morphology in cells expressing Tcb3(Full)-GBP (Fig 2B (iii)) was almost
similar to that in WT cells (Fig 2B (i)), indicating that the addition of GBP has no effect on vacuole
morphology. Remarkably, vacuoles remained non-fragmented even in cells expressing Tcb3(TM)-
GBP, which lacked most of the C-terminal region (Fig 2B (v)), as well as Tcb3(TM-SMP)-GBP (Fig
2B (iv)). In an attempt to address the question of why the TM domain of Tcb3p is sufficient to
suppress vacuolar division, we found that cells expressing Tcb3(TM)-GBP and lacking Tcb1p and
Tcb2p (Fig 2B (vi)) showed vacuolar fragmentation to the same extent as *tcb3Δ* mutant cells (Fig
2B (ii)). These results suggest that the TM domain of Tcb3p requires Tcb1p and Tcb2p to suppress
vacuole fission. According to the structural simulation by AlphaFold2, each TM domain of Tcb1p,
Tcb2p, and Tcb3p formed alpha-helical structure, which is known to be typical for membrane
proteins (Fig 2C). It is likely that Tcb1p and Tcb2p directly interact with Tcb3p by their TM domains.
Interestingly, the complex of these three TM domains was constructed with Tcb3p between Tcb1p
and Tcb2p in most cases (Fig 2D). Together these data suggest that the TM domain of Tcb3p
contributes sufficiently to the maintenance of vacuolar morphology by mediating the tricalbin
complex formation (Fig 2E).

Accumulated phytosphingosine in tricalbin-deleted cells causes vacuole fragmentation.
182   To understand how the deletion of tricalbins leads to vacuole fragmentation, we examined
the involvement of lipids. Previous reports showed that levels of ceramides and acylceramides
increased in *tcb1Δ2Δ3Δ* cells [37], and levels of long-chain bases (LCBs) increased in Δtether cells
lacking six ER-PM tethering proteins [47]. Here we measured lipids in *tcb1Δ2Δ3Δ* cells by in vivo
labeling with [^{3}H] dihydrosphingosine (DHS), which is a precursor of phytosphingosine (PHS), and
observed significant increases in ceramide, phosphatidylethanolamine (PE), PHS,
phosphatidylinositol (PI), complex sphingolipids such as inositolphosphorylceramide (IPC) and
mannosyl-inositolphosphorylceramide (MIPC) and LCB-1P (DHS-1P/PHS-1P) levels (Fig 3A).
190   We first tested whether increased levels of PHS cause vacuolar fragmentation. Loss of
ceramide synthases could cause an increase in PHS levels. Our analysis showed that vacuoles
are fragmented in *lag1Δlac1Δ* cells, which lack both enzymes for LCBs (DHS and PHS) conversion
into ceramides (Fig 3B). This suggests that ceramide precursors, LCBs or LCB-1P, can induce
vacuolar fragmentation. Therefore, we tested if exogenously added PHS induces vacuolar
fragmentation in WT cells. As shown in Fig 3C, exogenous addition of PHS induced vacuolar
fragmentation. PHS are converted into ceramide by the ceramide synthase Lag1p, which is the
main enzyme synthesizing phytoceramide [48]. Thus, we next examined whether PHS-induced
vacuolar fragmentation occurs in *lag1Δ* cells. Our analysis showed that vacuolar fragmentation in

lag1Δ cells treated with PHS was comparable to that for WT cells (Fig 3C). These results confirm

that the increases in ceramide and subsequent product IPC/MIPC are not the cause of vacuolar

fragmentation, but rather its precursors induce vacuolar fragmentation.

Because our lipid analysis showed a strong increase in LCB-1P in *tcb1Δ2Δ3Δ* cells,

phosphorylation of PHS may be involved in PHS-induced vacuolar fragmentation. Although

exogenously added LCBs move slowly from the PM to the ER if they are not phosphorylated,

efficient utilization of exogenously added LCBs in sphingolipid synthesis requires a series of

phosphorylation and dephosphorylation steps for LCBs [49]. The major yeast LCB phosphate

phosphatase Lcb3p dephosphorylates LCB-1P (DHS-1P/PHS-1P) to yield DHS/PHS [50], while

Lcb4p and Lcb5p catalyze the reverse reaction producing DHS-1P/PHS-1P [51]. Lcb3p, Lcb4p,

and Lcb5p are localized to the ER, PM or Golgi [49] [52] [53] [54]. We examined the ability of

exogenous PHS to fragment vacuoles in cells lacking these processing factors. Our results showed

that PHS-induced vacuolar fragmentation was completely blocked in the *lcb3Δ* cells in which PHS-

1P is not dephosphorylated (Fig 3D), suggesting that PHS-induced vacuolar fragmentation requires

the reaction of dephosphorylation of PHS-1P by Lcb3p. On the other hand, we observed that

vacuoles were still fragmented in *lcb4Δ lcb5Δ* and *lcb3Δ lcb4Δ lcb5Δ* cells (Fig 3D). These results

suggest that elevated levels of non-phosphorylated PHS induces vacuolar fragmentation (Fig 3E).

Additionally, we tested whether overexpression of Rsb1p, which has been reported as a

translocase that exports LCBs from the inner to the outer leaflet of the PM [55], rescues vacuole

fragmentation in *tcb1Δ2Δ3Δ* cells. As shown in Fig 3F, we observed that Rsb1p overexpression

results in decreased vacuole fragmentation. Collectively, these results support the model in which

vacuole fragmentation in tricalbin-deleted cells is caused by increased levels of PHS.

The nucleus-vacuole junction is required for PHS- or *tcb3Δ*-induced vacuole fragmentation

To characterize further the relationship between PHS and vacuole morphology, we asked

whether PHS delivered from the ER to the vacuole induces vacuole fragmentation. We

hypothesized that if PHS are transported to the vacuole by either the vesicular transport pathway

through the Golgi apparatus or the non-vesicular transport pathway via inter-organellar MCS

triggers vacuole fission, then blockage of the transport could rescue the PHS-induced vacuole

fragmentation. Accordingly, we used two types of mutant strains, one blocking vesicular transport

and the other blocking the NVJ. In a temperature sensitive *sec18-20* mutant, vesicular transport is

abolished at the restrictive temperature of 30 ºC or higher, but even at the permissive temperature

of 25 ºC, vesicular transport is partially impaired [56]. As shown in Fig 4A (left), under the condition

without exogenous PHS, some vacuoles in *sec18-20* mutant cells became fragmented when shifted

to the non-permissive temperature of 30 ºC. We also observed that vacuoles in *sec18-20* mutant

at 30 ºC underwent fragmentation after addition of PHS to the same extent as at the permissive

temperature of 25 ºC (Fig 4A, right), suggesting that vesicle-mediated transport is not required for

PHS-induced vacuolar fragmentation. To test the role of NVJ-mediated transport for PHS-induced

vacuolar fragmentation we employed a quadruple Δ*NVJ* mutant (*nvj1*Δ *nvj2*Δ *nvj3*Δ *mdm1*Δ) that

was used in a previous study in which complete loss of the NVJ was observed [57], but thus has

a different background to other strains of this study. When PHS was added to the Δ*NVJ* mutant,

we observed a significant suppression of vacuole fragmentation compared to WT (Fig 4B). Finally,

to investigate the requirement for NVJ in tricalbin deletion-induced vacuolar fragmentation, we

constructed the *tcb3*Δ *nvj1*Δ and *tcb3*Δ *nvj1*Δ *nvj2*Δ *nvj3*Δ *mdm1*Δ mutants. *TCB3* single disruption

sufficiently induced vacuolar fragmentation (Fig 1B), whereas as expected, the fragmentation was

partially suppressed by loss of only *NVJ1* and completely suppressed by loss of all NVJ factors

(*NVJ1*, *NVJ2*, *NVJ3*, and *MDM1*) (Fig 4C). Taken together, we conclude from these findings that

accumulated PHS in tricalbin deleted cells triggers vacuole fission via non-vesicular transport of

PHS at the NVJ.

Discussion

In the present study, we found that the accumulation of PHS triggers the fission of vacuoles.

Our results suggest that MCSs are involved in this process in two steps. First, the intracellular

amount of PHS is modulated by tricalbin-tethered MCSs between the ER and PM or Golgi (Fig 4D

blue arrow). Second, the accumulated PHS in the tricalbin-deleted cells induces vacuole fission

via most likely the NVJ (Fig 4D red arrow). Thus, we propose that MCSs regulate vacuole

morphology via sphingolipid metabolism. Fundamental questions that arise from our data concern

the accumulation of PHS in the tricalbin deletion strain and the mechanism of PHS-induced vacuole

fission. Possible mechanisms for elevated PHS levels in tricalbin-deleted cells are the following.

MCS deficiency between ER and PM has been shown to reduce the activity of Sac1p, a PtdIns4P

phosphatase [36]. Sac1p disruption strain decreases the levels of complex sphingolipids such as

IPC and MIPC, while it increases the levels of their precursors, ceramide, LCB, and LCB-1P [58].

This is probably due to the loss of function of Sac1p, which reduces the level of PtdIns used for

IPC synthesis, thereby impairing IPC synthesis, and resulting in the accumulation of precursors

such as the substrate ceramide. This model is also consistent with the results of accumulated

PtdIns4P and reduced IPC synthesis in Δtether cells [47]. As tricalbins are required for ceramide

nonvesicular transport from the ER to the Golgi [37], it is also possible that impaired ceramide

nonvesicular transport due to tricalbin deficiency causes ceramide and its precursor, LCB, to

accumulate in the ER. Another possibility is that MCS facilitate PHS diffusion between the ER and

the PM, which might be coordinated with the LCB export from the PM by Rsb1p. The loss of

tricalbins partially disrupts the ER-PM tether, possibly resulting low efficiency of PHS ejection by

Rsb1p. This is supported by the result that overexpression of Rsb1p suppressed vacuolar

fragmentation in tricalbin-deleted cells (Fig 3F).

Because both PHS- and tricalbin deletion-induced vacuolar fragmentations were partially

suppressed by the lack of NVJ (Fig 4B, 4C), it is suggested that transport of PHS into vacuoles via

the NVJ is involved in triggering vacuolar fragmentation. Recently, it has been reported that

sphingoid bases are transferred between ER and vacuole via the NVJ, and that Mdm1p, a key

tethering protein for the formation of the NVJ, may play an additional role in LCB transfer [35].

Perhaps, Mdm1p may be responsible for pulling out and passing LCBs. However, we cannot rule

out the possibility that the repression of vacuolar fragmentation in the absence of NVJ is not due to

inhibition of PHS transfer, but rather to changes in the lipid composition of the vacuolar membrane

caused by the lack of supply of other substances capable of triggering vacuolar fragmentation other

than PHS, like sterols and lipids such as PtdIns[3,5]P2 and its precursors. Further analysis in this

regard is warranted.

How accumulated PHS triggers vacuolar fragmentation remains undetermined.

Sphingosine, as a bioactive lipid, has been reported to exert effects on enzyme activity in humans

and yeast [59] [60] [61], and it is possible that PHS may induce vacuolar fragmentation through

established signaling pathways. Yabuki et al [61] reported that LCB accumulation activates a

signaling pathway that includes major yeast regulatory kinases such as Pkh1/2p, Pkc1p and

TORC1, which may be a candidate to promote vacuolar fragmentation. Fab1p, a target of TORC1,

is responsible for the production of PtdIns[3,5]P2, which is a well-established inducer of vacuolar

fragmentation. Fab1p exhibits co-localization with the TORC1-activating EGO complex, and its

activity is controlled by Ivy1p [62] and TORC1 [63]. PtdIns[3,5]P2 was shown to regulate vacuole

fission employing Vps1p and Atg18p as executioners [64] while also promoting TORC1 activity in

a positive feedback loop [65]. In this context, we found that PHS-induced vacuolar fragmentation

can be suppressed by the loss of Fab1p and its regulatory binding partner Vac14p (Fig S2A). This

observation suggests that PHS-induced vacuolar fragmentation requires PI metabolism. On the

other hand, it was shown that membrane division is mediated by certain proteins that contain

amphipathic helices (AHs) and interact with lipid cofactors such as PtdIns[4,5]P2, PA, and

cardiolipin [66]. Thus, PHS may possess a similar regulatory function as a lipid cofactor for the

activity of fission-inducing proteins.

If PHS-induced vacuole fragmentation is not due to signaling, another possible model is

that PHS accumulation in the vacuolar membrane causes physical changes in the membrane

structure that result in membrane fragmentation. The accumulation of sphingosine in the of GARP

mutant *vps53Δ*, in which retrograde transport from the endosome to the Golgi is blocked, caused

vacuolar fragmentation [67], and thus also supports this model. Sphingosine stabilizes (rigidifies)
the gel domains in the membrane, leading to a structural defect between the phase separation of
"more rigid" and "less rigid" domains [68]. This structural defect may result in high membrane
permeability. Sphingosine also forms small and unstable channels (compared to the channels
formed by ceramide) in the membrane [69]. Sphingosine channels are not large enough to release
proteins, but are believed to induce a permeability transition. Other studies have suggested that
sphingosine induces non-lamellar structures by interacting with negatively charged lipids such as
PA [70]. Finally, PHS alone or in concert with negatively charged lipids such as PtdIns[3,5]P2 may
be actively involved in the fission process by inducing structural changes in the vacuolar
membrane.

Besides PHS-induced vacuolar fission, it is generally well-known that vacuolar division can
be triggered as an acute response to osmotic shock. We made an interesting observation under
hyperosmotic conditions (0.2 M NaCl), wherein the loss of NVJ led to complete suppression of
vacuolar division (Fig S2B). Although the involvement of PHS transport in this process remains
unclear, our finding suggests a significant role for NVJ in vacuolar fission as a hyperosmotic
response. Collectively, we propose that NVJ plays a general regulatory role in vacuolar
morphology.

324
325
326

Materials and Methods

Yeast strains

All strains of *S. cerevisiae* used for this work are listed in Supplementary Table 1.

Plasmids

All plasmids used for this work are listed in Supplementary Table 2.

Culture conditions

Yeast cells were grown either in rich YPD medium (2% glucose, 1% yeast extract, 2% peptone) or

in synthetic minimal SD medium (2% glucose, 0.15% yeast nitrogen base, 0.5% ammonium sulfate,

bases as nutritional requirements) and supplemented with the appropriate amino acids.

FM4-64 stain and Fluorescence Microscopy

Yeast cells were cultured in YPD medium at 25°C for 15 hours to achieve OD_{600} = 0.5. The cells

were collected and suspended in the same medium to achieve OD_{600}=20. 20 mM FM4-64 (N-(3-

Triethylammoniumpropyl)-4-(6-(4-(diethylamino) phenyl) hexatrienyl) pyridinium) dissolved in

DMSO was added (a final concentration of 20 µM) while shielded from light. The cells were

incubated with FM4-64 for 15 minutes at 25°C and then the cells were washed twice with YPD

medium. Cells were suspended to achieve OD_{600}=10 in the same YPD medium or YPD containing

reagents (such as rapamycin or PHS), and incubated at 25°C for 2 hours under light shielding to

label the vacuoles. After incubation, cells were collected and observed with a fluorescence

microscope.

Western Blotting

To analyze Sch9 phosphorylation, protein extracts from cells expressing SCH9-5HA were treated

with 2-nitro-5-thiocyanobenzoic acid overnight, resolved by SDS-PAGE, immunoblotted, and

visualized using rat anti-HA monoclonal antibody (3F10; Roche) and anti-rat IgG antibody (A9037;

Sigma-Aldrich) produced in goat. Bands were quantified using ImageJ to determine the relative

amounts of phosphorylated Sch9.

Lipid Labelling with [³H]DHS

In vivo labelling of lipids with [³H]DHS was carried out as described [71]. Radiolabeled lipids were

extracted with chloroform-methanol-water (10:10:3, vol/vol/vol), and analyzed by thin-layer

chromatography (TLC) using a solvent system (Chloroform-methanol-4.2N ammonium hydroxide

(9:7:2, vol/vol/vol)). Radiolabeled lipids were visualized and quantified on an FLA-7000 system.

Protein complex modelling

The modeled Tcb1p, Tcb2p, and Tcb3p structures were obtained from AlphaFold2 Protein

Structure Database (https://alphafold.ebi.ac.uk/). Based on the modeled structures, the amino

acids of TM regions were predicted as 79-171 for Tcb1p, 79-162 for Tcb2p, and 189-268 for Tcb3p.

The protein complex structures of the TM regions in Tcb1p, Tcb2p, and Tcb3p were modelled by

AlphaFold2   program   [72]   worked   on   the   AlphaFold   Colab   web   space

(https://colab.research.google.com/github/deepmind/alphafold/blob/main/notebooks/AlphaFold.ipy

nb#scrollTo=pc5-mbsX9PZC). The figures were drawn using PyMOL software provided by

Schrödinger, Inc. (https://pymol.org/2/).

Data and Materials Availability Statement

The data and materials that support the findings of this study are included in the article and its

Supplemental Information, or are available from the corresponding authors on request.

Acknowledgments

We thank Takashi Toda and Masashi Yukawa for GBP-plasmid, Mike Henne for the Δ*NVJ* strain, and Howard Riezman for GFP-CPS1 plasmid. This research was supported by JSPS KAKENHI 21K20572 and 22K14863 to A.I., 19H02922 and 21K19088 to K.F.

Author Contributions

K.N., A.I. and K.F. designed research; K.H., K. N., A.I., S.Y., A.N. and S.F. performed research; A.I. analyzed data; and A.I., P.S. and K.F. wrote the paper.

Competing Interest Statement

The authors declare no competing interest.

References

1. Baars, T. L., Petri, S., Peters, C., & Mayer, A. (2007). Role of the V-ATPase in regulation of the vacuolar fission-fusion equilibrium. *Molecular biology of the cell*, *18*(10), 3873–3882. doi:10.1091/mbc.e07-03-0205. https://doi.org/10.1091/mbc.e07-03-0205

2. Li, S. C., & Kane, P. M. (2009). The yeast lysosome-like vacuole: endpoint and crossroads. *Biochimica et biophysica acta*, *1793*(4), 650–663. doi:10.1016/j.bbamcr.2008.08.003. https://doi.org/10.1016/j.bbamcr.2008.08.003

3. Wickner, W., & Haas, A. (2000). Yeast homotypic vacuole fusion: a window on organelle trafficking mechanisms. *Annual review of biochemistry*, *69*, 247–275. doi:10.1146/annurev.biochem.69.1.247. https://doi.org/10.1146/annurev.biochem.69.1.247

4. Ostrowicz, C. W., Meiringer, C. T., & Ungermann, C. (2008). Yeast vacuole fusion: a model system for eukaryotic endomembrane dynamics. *Autophagy*, *4*(1), 5–19. doi:10.4161/auto.5054. https://doi.org/10.4161/auto.5054

5. Wickner W. (2010). Membrane fusion: five lipids, four SNAREs, three chaperones, two nucleotides, and a Rab, all dancing in a ring on yeast vacuoles. *Annual review of cell and*

developmental biology, *26*, 115–136. doi:10.1146/annurev-cellbio-100109-104131.

https://doi.org/10.1146/annurev-cellbio-100109-104131

6. Qiu Q. S. (2012). V-ATPase, ScNhx1p and yeast vacuole fusion. *Journal of genetics and*

genomics = Yi chuan xue bao, *39*(4), 167–171. doi:10.1016/j.jgg.2012.02.001.

https://doi.org/10.1016/j.jgg.2012.02.001

7. Müller, O., Bayer, M. J., Peters, C., Andersen, J. S., Mann, M., & Mayer, A. (2002). The Vtc

proteins in vacuole fusion: coupling NSF activity to V(0) trans-complex formation. *The*

EMBO journal, *21*(3), 259–269. doi:10.1093/emboj/21.3.259.

https://doi.org/10.1093/emboj/21.3.259

8. Ho, R., & Stroupe, C. (2016). The HOPS/Class C Vps Complex Tethers High-Curvature

Membranes via a Direct Protein-Membrane Interaction. *Traffic (Copenhagen,*

Denmark), *17*(10), 1078–1090. doi:10.1111/tra.12421.

https://doi.org/10.1111/tra.12421

9. Ungermann, C., Wickner, W., & Xu, Z. (1999). Vacuole acidification is required for trans-

SNARE pairing, LMA1 release, and homotypic fusion. *Proceedings of the National*

Academy of Sciences of the United States of America, *96*(20), 11194–11199.

doi:10.1073/pnas.96.20.11194. https://doi.org/10.1073/pnas.96.20.11194

10. Peters, C., Bayer, M. J., Bühler, S., Andersen, J. S., Mann, M., & Mayer, A. (2001). Trans-

complex formation by proteolipid channels in the terminal phase of membrane

fusion. *Nature*, *409*(6820), 581–588. doi:10.1038/35054500.

https://doi.org/10.1038/35054500

11. Desfougères, Y., Vavassori, S., Rompf, M., Gerasimaite, R., & Mayer, A. (2016). Organelle

acidification negatively regulates vacuole membrane fusion in vivo. *Scientific reports*, *6*,

29045. doi:10.1038/srep29045. https://doi.org/10.1038/srep29045

12. Starai, V. J., Thorngren, N., Fratti, R. A., & Wickner, W. (2005). Ion regulation of homotypic

vacuole fusion in Saccharomyces cerevisiae. *The Journal of biological chemistry*, *280*(17),

16754–16762. doi:10.1074/jbc.M500421200. https://doi.org/10.1074/jbc.M500421200

13. Qiu, Q. S., & Fratti, R. A. (2010). The Na+/H+ exchanger Nhx1p regulates the initiation of

Saccharomyces cerevisiae vacuole fusion. *Journal of cell science*, *123*(Pt 19), 3266–3275.

doi:10.1242/jcs.067637. https://doi.org/10.1242/jcs.067637

14. Zieger, M., & Mayer, A. (2012). Yeast vacuoles fragment in an asymmetrical two-phase
process with distinct protein requirements. *Molecular biology of the cell*, *23*(17), 3438–
3449. doi:10.1091/mbc.E12-05-0347. https://doi.org/10.1091/mbc.E12-05-0347

15. Peters, C., Baars, T. L., Bühler, S., & Mayer, A. (2004). Mutual control of membrane fission
and fusion proteins. *Cell*, *119*(5), 667–678. doi:10.1016/j.cell.2004.11.023.
https://doi.org/10.1016/j.cell.2004.11.023

16. Weisman L. S. (2006). Organelles on the move: insights from yeast vacuole
inheritance. *Nature reviews. Molecular cell biology*, *7*(4), 243–252. doi:10.1038/nrm1892.
https://doi.org/10.1038/nrm1892

17. Efe, J. A., Botelho, R. J., & Emr, S. D. (2005). The Fab1 phosphatidylinositol kinase pathway
in the regulation of vacuole morphology. *Current opinion in cell biology*, *17*(4), 402–408.
doi:10.1016/j.ceb.2005.06.002. https://doi.org/10.1016/j.ceb.2005.06.002

18. Efe, J. A., Botelho, R. J., & Emr, S. D. (2007). Atg18 regulates organelle morphology and
Fab1 kinase activity independent of its membrane recruitment by phosphatidylinositol
3,5-bisphosphate. *Molecular biology of the cell*, *18*(11), 4232–4244.
doi:10.1091/mbc.e07-04-0301. https://doi.org/10.1091/mbc.e07-04-0301

19. Michaillat, L., Baars, T. L., & Mayer, A. (2012). Cell-free reconstitution of vacuole
membrane fragmentation reveals regulation of vacuole size and number by
TORC1. *Molecular biology of the cell*, *23*(5), 881–895. doi:10.1091/mbc.E11-08-0703.
https://doi.org/10.1091/mbc.E11-08-0703

20. Stauffer, B., & Powers, T. (2015). Target of rapamycin signaling mediates vacuolar fission
caused by endoplasmic reticulum stress in Saccharomyces cerevisiae. *Molecular biology
of the cell*, *26*(25), 4618–4630. doi:10.1091/mbc.E15-06-0344.
https://doi.org/10.1091/mbc.E15-06-0344

21. Manandhar, S. P., Siddiqah, I. M., Cocca, S. M., & Gharakhanian, E. (2020). A kinase
cascade on the yeast lysosomal vacuole regulates its membrane dynamics: conserved
kinase Env7 is phosphorylated by casein kinase Yck3. *The Journal of biological
chemistry*, *295*(34), 12262–12278. doi:10.1074/jbc.RA119.012346.
https://doi.org/10.1074/jbc.RA119.012346

22. Wu, H., Carvalho, P., & Voeltz, G. K. (2018). Here, there, and everywhere: The importance
of ER membrane contact sites. *Science (New York, N.Y.)*, *361*(6401), eaan5835.
doi:10.1126/science.aan5835. https://doi.org/10.1126/science.aan5835

23. Prinz, W. A., Toulmay, A., & Balla, T. (2020). The functional universe of membrane contact
sites. *Nature reviews. Molecular cell biology*, *21*(1), 7–24. doi:10.1038/s41580-019-0180-
9. https://doi.org/10.1038/s41580-019-0180-9

24. Wickner, W., & Rizo, J. (2017). A cascade of multiple proteins and lipids catalyzes
membrane fusion. *Molecular biology of the cell*, *28*(6), 707–711. doi:10.1091/mbc.E16-
07-0517. https://doi.org/10.1091/mbc.E16-07-0517

25. Starr, M. L., & Fratti, R. A. (2019). The Participation of Regulatory Lipids in Vacuole
Homotypic Fusion. *Trends in biochemical sciences*, *44*(6), 546–554.
doi:10.1016/j.tibs.2018.12.003. https://doi.org/10.1016/j.tibs.2018.12.003

26. Ungermann, C., & Kümmel, D. (2019). Structure of membrane tethers and their role in
fusion. *Traffic (Copenhagen, Denmark)*, *20*(7), 479–490. doi:10.1111/tra.12655.
https://doi.org/10.1111/tra.12655

27. Hurst, L. R., & Fratti, R. A. (2020). Lipid Rafts, Sphingolipids, and Ergosterol in Yeast
Vacuole Fusion and Maturation. *Frontiers in cell and developmental biology*, *8*, 539.
doi:10.3389/fcell.2020.00539. https://doi.org/10.3389/fcell.2020.00539

28. Pan, X., Roberts, P., Chen, Y., Kvam, E., Shulga, N., Huang, K., Lemmon, S., & Goldfarb, D.
S. (2000). Nucleus-vacuole junctions in Saccharomyces cerevisiae are formed through the
direct interaction of Vac8p with Nvj1p. *Molecular biology of the cell*, *11*(7), 2445–2457.
doi:10.1091/mbc.11.7.2445. https://doi.org/10.1091/mbc.11.7.2445

29. Elbaz-Alon, Y., Rosenfeld-Gur, E., Shinder, V., Futerman, A. H., Geiger, T., & Schuldiner, M.
(2014). A dynamic interface between vacuoles and mitochondria in yeast. *Developmental
cell*, *30*(1), 95–102. doi:10.1016/j.devcel.2014.06.007.
https://doi.org/10.1016/j.devcel.2014.06.007

30. Hönscher, C., Mari, M., Auffarth, K., Bohnert, M., Griffith, J., Geerts, W., van der Laan, M.,
Cabrera, M., Reggiori, F., & Ungermann, C. (2014). Cellular metabolism regulates contact
sites between vacuoles and mitochondria. *Developmental cell*, *30*(1), 86–94.
doi:10.1016/j.devcel.2014.06.006. https://doi.org/10.1016/j.devcel.2014.06.006

31. González Montoro, A., Auffarth, K., Hönscher, C., Bohnert, M., Becker, T., Warscheid, B.,
Reggiori, F., van der Laan, M., Fröhlich, F., & Ungermann, C. (2018). Vps39 Interacts with
Tom40 to Establish One of Two Functionally Distinct Vacuole-Mitochondria Contact
Sites. *Developmental cell*, *45*(5), 621–636.e7. doi:10.1016/j.devcel.2018.05.011.
https://doi.org/10.1016/j.devcel.2018.05.011

32. Murley, A., Sarsam, R. D., Toulmay, A., Yamada, J., Prinz, W. A., & Nunnari, J. (2015). Ltc1
is an ER-localized sterol transporter and a component of ER-mitochondria and ER-vacuole
contacts. *The Journal of cell biology*, *209*(4), 539–548. doi:10.1083/jcb.201502033.
https://doi.org/10.1083/jcb.201502033

33. Murley, A., Yamada, J., Niles, B. J., Toulmay, A., Prinz, W. A., Powers, T., & Nunnari, J.
(2017). Sterol transporters at membrane contact sites regulate TORC1 and TORC2
signaling. *The Journal of cell biology*, *216*(9), 2679–2689.

34. Reinisch, K. M., & Prinz, W. A. (2021). Mechanisms of nonvesicular lipid transport. *The
Journal of cell biology*, *220*(3), e202012058. doi:10.1083/jcb.201610032.
https://doi.org/10.1083/jcb.201610032

35. Girik, V., Feng, S., Hariri, H., Henne, W. M., & Riezman, H. (2022). Vacuole-Specific Lipid
Release for Tracking Intracellular Lipid Metabolism and Transport in Saccharomyces
cerevisiae. *ACS chemical biology*, 17(6), 1485–1494. doi:10.1021/acschembio.2c00120.
https://doi.org/10.1021/acschembio.2c00120

36. Manford, A. G., Stefan, C. J., Yuan, H. L., Macgurn, J. A., & Emr, S. D. (2012). ER-to-plasma
membrane tethering proteins regulate cell signaling and ER morphology. *Developmental
cell*, *23*(6), 1129–1140. doi:10.1016/j.devcel.2012.11.004.
https://doi.org/10.1016/j.devcel.2012.11.004

37. Ikeda, A., Schlarmann, P., Kurokawa, K., Nakano, A., Riezman, H., & Funato, K. (2020).
Tricalbins Are Required for Non-vesicular Ceramide Transport at ER-Golgi Contacts and
Modulate Lipid Droplet Biogenesis. *iScience*, *23*(10), 101603.
doi:10.1016/j.isci.2020.101603. https://doi.org/10.1016/j.isci.2020.101603

38. Toulmay, A., & Prinz, W. A. (2012). A conserved membrane-binding domain targets
proteins to organelle contact sites. *Journal of cell science*, *125*(Pt 1), 49–58.
doi:10.1242/jcs.085118. https://doi.org/10.1242/jcs.085118

39. Quon, E., Sere, Y. Y., Chauhan, N., Johansen, J., Sullivan, D. P., Dittman, J. S., Rice, W. J.,
Chan, R. B., Di Paolo, G., Beh, C. T., & Menon, A. K. (2018). Endoplasmic reticulum-plasma
membrane contact sites integrate sterol and phospholipid regulation. *PLoS biology*, *16*(5),
e2003864. doi:10.1371/journal.pbio.2003864.
https://doi.org/10.1371/journal.pbio.2003864

40. Jorgensen, J. R., Tei, R., Baskin, J. M., Michel, A. H., Kornmann, B., & Emr, S. D. (2020).
ESCRT-III and ER-PM contacts maintain lipid homeostasis. *Molecular biology of the cell*,
31(12), 1302–1313. doi:10.1091/mbc.E20-01-0061. https://doi.org/10.1091/mbc.E20-
01-0061

41. Hoffmann, P. C., Bharat, T., Wozny, M. R., Boulanger, J., Miller, E. A., & Kukulski, W. (2019).
Tricalbins Contribute to Cellular Lipid Flux and Form Curved ER-PM Contacts that Are
Bridged by Rod-Shaped Structures. *Developmental cell*, *51*(4), 488–502.e8.
doi:10.1016/j.devcel.2019.09.019. https://doi.org/10.1016/j.devcel.2019.09.019

42. Collado, J., Kalemanov, M., Campelo, F., Bourgoint, C., Thomas, F., Loewith, R., Martínez-
Sánchez, A., Baumeister, W., Stefan, C. J., & Fernández-Busnadiego, R. (2019). Tricalbin-
Mediated Contact Sites Control ER Curvature to Maintain Plasma Membrane Integrity.
Developmental cell, 51(4), 476–487.e7. doi:10.1016/j.devcel.2019.10.018.
https://doi.org/10.1016/j.devcel.2019.10.018

43. Nelson, H., & Nelson, N. (1990). Disruption of genes encoding subunits of yeast vacuolar
H(+)-ATPase causes conditional lethality. *Proceedings of the National Academy of*
Sciences of the United States of America, *87*(9), 3503–3507. doi:10.1073/pnas.87.9.3503.
https://doi.org/10.1073/pnas.87.9.3503

44. Coonrod, E. M., Graham, L. A., Carpp, L. N., Carr, T. M., Stirrat, L., Bowers, K., Bryant, N. J.,
& Stevens, T. H. (2013). Homotypic vacuole fusion in yeast requires organelle acidification
and not the V-ATPase membrane domain. *Developmental cell*, *27*(4), 462–468.
doi:10.1016/j.devcel.2013.10.014. https://doi.org/10.1016/j.devcel.2013.10.014

45. Seeley, E. S., Kato, M., Margolis, N., Wickner, W., & Eitzen, G. (2002). Genomic analysis of
homotypic vacuole fusion. *Molecular biology of the cell*, *13*(3), 782–794.
doi:10.1091/mbc.01-10-0512. https://doi.org/10.1091/mbc.01-10-0512

46. Dove, S. K., Dong, K., Kobayashi, T., Williams, F. K., & Michell, R. H. (2009).
Phosphatidylinositol 3,5-bisphosphate and Fab1p/PIKfyve underPPIn endo-lysosome

function. *The Biochemical journal*, *419*(1), 1–13. doi:10.1042/BJ20081950.
https://doi.org/10.1042/BJ20081950

47. Omnus, D. J., Manford, A. G., Bader, J. M., Emr, S. D., & Stefan, C. J. (2016).
Phosphoinositide kinase signaling controls ER-PM cross-talk. *Molecular biology of the*
cell, *27*(7), 1170–1180. doi:10.1091/mbc.E16-01-0002.
https://doi.org/10.1091/mbc.E16-01-0002

48. Megyeri, M., Prasad, R., Volpert, G., Sliwa-Gonzalez, A., Haribowo, A. G., Aguilera-Romero,
A., Riezman, H., Barral, Y., Futerman, A. H., & Schuldiner, M. (2019). Yeast ceramide
synthases, Lag1 and Lac1, have distinct substrate specificity. *Journal of cell*
science, *132*(12), jcs228411. doi:10.1242/jcs.228411. https://doi.org/10.1242/jcs.228411

49. Funato, K., Lombardi, R., Vallee, B., & Riezman, H. (2003). Lcb4p is a key regulator of
ceramide synthesis from exogenous long chain sphingoid base in Saccharomyces
cerevisiae. *The Journal of biological chemistry*, *278*(9), 7325–7334.
doi:10.1074/jbc.M209925200. https://doi.org/10.1074/jbc.M209925200

50. Qie, L., Nagiec, M. M., Baltisberger, J. A., Lester, R. L., & Dickson, R. C. (1997). Identification
of a Saccharomyces gene, LCB3, necessary for incorporation of exogenous long chain
bases into sphingolipids. *The Journal of biological chemistry*, *272*(26), 16110–16117.
doi:10.1074/jbc.272.26.16110. https://doi.org/10.1074/jbc.272.26.16110

51. Nagiec, M. M., Skrzypek, M., Nagiec, E. E., Lester, R. L., & Dickson, R. C. (1998). The LCB4
(YOR171c) and LCB5 (YLR260w) genes of Saccharomyces encode sphingoid long chain
base kinases. *The Journal of biological chemistry*, *273*(31), 19437–19442.
doi:10.1074/jbc.273.31.19437. https://doi.org/10.1074/jbc.273.31.19437

52. Mao, C., Saba, J. D., & Obeid, L. M. (1999). The dihydrosphingosine-1-phosphate
phosphatases of Saccharomyces cerevisiae are important regulators of cell proliferation
and heat stress responses. *The Biochemical journal*, 342 Pt 3(Pt 3), 667–675. PMID:
10477278; PMCID: PMC1220508.

53. Hait, N. C., Fujita, K., Lester, R. L., & Dickson, R. C. (2002). Lcb4p sphingoid base kinase
localizes to the Golgi and late endosomes. *FEBS letters*, 532(1-2), 97–102.
doi:10.1016/s0014-5793(02)03636-0. https://doi.org/10.1016/s0014-5793(02)03636-0

54. Iwaki, S., Sano, T., Takagi, T., Osumi, M., Kihara, A., & Igarashi, Y. (2007). Intracellular
trafficking pathway of yeast long-chain base kinase Lcb4, from its synthesis to its

degradation. *The Journal of biological chemistry*, 282(39), 28485–28492.
doi:10.1074/jbc.M701607200. https://doi.org/10.1074/jbc.M701607200

55. Kihara, A., & Igarashi, Y. (2002). Identification and characterization of a Saccharomyces
cerevisiae gene, RSB1, involved in sphingoid long-chain base release. *The Journal of*
biological chemistry, *277*(33), 30048–30054. doi:10.1074/jbc.M203385200.
https://doi.org/10.1074/jbc.M203385200

56. Funato, K., & Riezman, H. (2001). Vesicular and nonvesicular transport of ceramide from
ER to the Golgi apparatus in yeast. *The Journal of cell biology*, 155(6), 949–959.
doi:10.1083/jcb.200105033. https://doi.org/10.1083/jcb.200105033

57. Henne, W. M., Zhu, L., Balogi, Z., Stefan, C., Pleiss, J. A., & Emr, S. D. (2015). Mdm1/Snx13
is a novel ER-endolysosomal interorganelle tethering protein. *The Journal of cell biology*,
210(4), 541–551. doi:10.1083/jcb.201503088. https://doi.org/10.1083/jcb.201503088

58. Brice, S. E., Alford, C. W., & Cowart, L. A. (2009). Modulation of sphingolipid metabolism
by the phosphatidylinositol-4-phosphate phosphatase Sac1p through regulation of
phosphatidylinositol in Saccharomyces cerevisiae. *The Journal of biological*
chemistry, *284*(12), 7588–7596. doi:10.1074/jbc.M808325200.
https://doi.org/10.1074/jbc.M808325200

59. Hannun, Y. A., Loomis, C. R., Merrill, A. H., Jr, & Bell, R. M. (1986). Sphingosine inhibition
of protein kinase C activity and of phorbol dibutyrate binding in vitro and in human
platelets. *The Journal of biological chemistry*, *261*(27), 12604–12609. PMID: 3462188.

60. Chang, H. C., Tsai, L. H., Chuang, L. Y., & Hung, W. C. (2001). Role of AKT kinase in
sphingosine-induced apoptosis in human hepatoma cells. *Journal of cellular*
physiology, *188*(2), 188–193. doi:10.1002/jcp.1108. https://doi.org/10.1002/jcp.1108

61. Yabuki, Y., Ikeda, A., Araki, M., Kajiwara, K., Mizuta, K., & Funato, K. (2019).
Sphingolipid/Pkh1/2-TORC1/Sch9 Signaling Regulates Ribosome Biogenesis in
Tunicamycin-Induced Stress Response in Yeast. *Genetics*, *212*(1), 175–186.
doi:10.1534/genetics.118.301874. https://doi.org/10.1534/genetics.118.301874

62. Malia, P. C., Numrich, J., Nishimura, T., González Montoro, A., Stefan, C. J., & Ungermann,
C. (2018). Control of vacuole membrane homeostasis by a resident PI-3,5-kinase
inhibitor. *Proceedings of the National Academy of Sciences of the United States of*

America, *115*(18),   4684–4689.   doi:10.1073/pnas.1722517115.

https://doi.org/10.1073/pnas.1722517115

63. Chen, Z., Malia, P. C., Hatakeyama, R., Nicastro, R., Hu, Z., Péli-Gulli, M. P., Gao, J.,

Nishimura, T., Eskes, E., Stefan, C. J., Winderickx, J., Dengjel, J., De Virgilio, C., &

Ungermann, C. (2021). TORC1 Determines Fab1 Lipid Kinase Function at Signaling

Endosomes   and   Vacuoles. *Current   biology   :   CB*, *31*(2),   297–309.e8.

doi:10.1016/j.cub.2020.10.026. https://doi.org/10.1016/j.cub.2020.10.026

64. Gopaldass, N., Fauvet, B., Lashuel, H., Roux, A., & Mayer, A. (2017). Membrane scission

driven   by   the   PROPPIN   Atg18. *The   EMBO   journal*, *36*(22),   3274–3291.

doi:10.15252/embj.201796859. https://doi.org/10.15252/embj.201796859

65. Jin, N., Mao, K., Jin, Y., Tevzadze, G., Kauffman, E. J., Park, S., Bridges, D., Loewith, R.,

Saltiel, A. R., Klionsky, D. J., & Weisman, L. S. (2014). Roles for PI(3,5)P2 in nutrient sensing

through TORC1. *Molecular biology of the cell*, *25*(7), 1171–1185. doi:10.1091/mbc.E14-

01-0021. https://doi.org/10.1091/mbc.E14-01-0021

66. Zhukovsky, M. A., Filograna, A., Luini, A., Corda, D., & Valente, C. (2019). Protein

Amphipathic Helix Insertion: A Mechanism to Induce Membrane Fission. *Frontiers in cell*

and   developmental   biology, *7*,   291.   doi:10.3389/fcell.2019.00291.

https://doi.org/10.3389/fcell.2019.00291

67. Fröhlich, F., Petit, C., Kory, N., Christiano, R., Hannibal-Bach, H. K., Graham, M., Liu, X.,

Ejsing, C. S., Farese, R. V., & Walther, T. C. (2015). The GARP complex is required for

cellular   sphingolipid   homeostasis. *eLife*, *4*,   e08712.   doi:10.7554/eLife.08712.

https://doi.org/10.7554/eLife.08712

68. Contreras, F. X., Sot, J., Alonso, A., & Goñi, F. M. (2006). Sphingosine increases the

permeability of model and cell membranes. *Biophysical journal*, *90*(11), 4085–4092.

doi:10.1529/biophysj.105.076471. https://doi.org/10.1529/biophysj.105.076471

69. Siskind, L. J., Fluss, S., Bui, M., & Colombini, M. (2005). Sphingosine forms channels in

membranes that differ greatly from those formed by ceramide. *Journal of bioenergetics*

and   biomembranes, *37*(4),   227–236.   doi:10.1007/s10863-005-6632-2.

https://doi.org/10.1007/s10863-005-6632-2

70. Jiménez-Rojo, N., Sot, J., Viguera, A. R., Collado, M. I., Torrecillas, A., Gómez-Fernández, J.

C., Goñi, F. M., & Alonso, A. (2014). Membrane permeabilization induced by sphingosine:

effect of negatively charged lipids. *Biophysical journal*, *106*(12), 2577–2584.
doi:10.1016/j.bpj.2014.04.038. https://doi.org/10.1016/j.bpj.2014.04.038

71. Ikeda, A., Hanaoka, K., & Funato, K. (2021). Protocol for measuring sphingolipid
metabolism in budding yeast. STAR protocols, 2(2), 100412.
doi:10.1016/j.xpro.2021.100412. https://doi.org/10.1016/j.xpro.2021.100412

72. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.,
Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T.,
Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M.,
Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A. W., Kavukcuoglu, K., Kohli,
P., Hassabis, D., & Hassabis, D. (2021). Highly accurate protein structure prediction with
AlphaFold. *Nature*, *596*(7873), 583–589. doi:10.1038/s41586-021-03819-2.
https://doi.org/10.1038/s41586-021-03819-2

Figures and legends

Figure 1. Deletion of tricalbin proteins causes vacuole fragmentation

(A, B) Cells (FKY2577 and FKY2927 in A; FKY2577, FKY2909, FKY3819, FKY2924, FKY3023,

FKY3820 and FKY3008 in B) were grown overnight at 25 ºC in YPD. Then vacuoles were stained

with FM4-64 and imaged by fluorescence microscopy. The number of vacuoles per cell was
counted and categorized into one of three groups. The data represent mean ± SE of three
independent experiments, each based on more than 100 cells. $*p < 0.05$, $**p < 0.01$ and $***p <$
0.001 by Student's t-test compared with WT.

(C) Cells (FKY2577 and FKY2927) were grown overnight at 25 ºC in YPD. Cells were then
incubated in sterile distilled water (SDW) for more than 45 minutes or YPD with 200 nM of
rapamycin for 2 hours. Vacuoles were stained with FM4-64 and imaged by fluorescence
microscopy. The number of vacuoles per cell was counted and categorized into one of three
groups. The data represent mean ± SE of seven (SDW) or three (Rap) independent experiments,
each based on more than 100 cells. $**p < 0.01$ by Student's t-test compared with none treated
tcb1Δ2Δ3Δ cells.

(A-C) Significant differences analysis between the pairwise combination of groups was performed
using two-way ANOVA.

(D) Cells (FKY2577 and FKY2927) transformed with pRS416-SCH9-5HA were cultured in YPD,
treated with 200 nM rapamycin (control) or untreated. The extracts from cells expressing Sch9-
5HA were reacted with 2-nitro-5-thiocyanobenzoic acid and analyzed by immunoblotting using anti-
HA. Phosphorylated Sch9 relative to the total Sch9 was calculated and shown in comparison to
untreated WT cells. The data represent mean ± SE of three independent experiments. $*p < 0.05$ by
Student's t-test.

(E) Illustration shows that tricalbin proteins negatively regulate the vacuole fission in a TORC1-
independent manner.

(A)

(B)

(C)

(D)

(E)

Figure 2. Effects of domain deletion and artificial tether on vacuole morphology

(A) Diagram of domain organization of Tcb3 protein. TM, transmembrane domain; SMP,

synaptotagmin-like mitochondrial lipid-binding protein; C2, calcium-dependent lipid-binding

domain; GBP, GFP-binding protein.

(B) Cells (FKY2577 (i), FKY2924 (ii), FKY3903 (iii), FKY3904 (iv), FKY3905 (v) and FKY4754 (vi))

were grown overnight at 25 ºC in YPD. Then vacuoles were stained with FM4-64 and imaged by

fluorescence microscopy. The number of vacuoles per cell was counted and categorized into one

of three groups. The data represent mean ± SE of three independent experiments, each based on

more than 100 cells. $*p < 0.05$ and $**p < 0.01$ by Student's t-test compared with Tcb1 Tcb2 Tcb3

(i). Significant differences analysis between the pairwise combination of groups was performed

using two-way ANOVA.

(C) The modeled structures of the Tcb1p, Tcb2p and Tcb3p proteins. The ribbons and arrows

indicate alpha-helices and beta-sheets, respectively.

(D) TM domain complex in Tcb1p (red), Tcb2p (green), and Tcb3p (blue). The rank "X" indicates

the order in which the complexes are most likely to form.

(E) Illustrations show that TM domain of Tcb3 contributes to mediating protein interactions between

the tricalbin family to maintain vacuolar morphology.

Figure 3. Accumulated PHS in *tcb1Δ2Δ3Δ* causes vacuole fragmentation.

(A) Cells (FKY2577 and FKY2927) were grown at 25˚C, and labeled with [³H]DHS for 3 hours. Labelled lipids were applied to TLC plates using solvent system (Chloroform-methanol-4.2N ammonium hydroxide (9:7:2, v/v/v)). Incorporation of [³H]DHS into each lipid was quantified and

the percentage of the total radioactivity (%) in WT cells was determined. Data represent mean ±

SE of four independent experiments. ** $P < 0.01$ by Student's t-test.

(B-D, F) Cells (FKY5687 and FKY5688 in B; FKY3340 and YKC121-59 in C; FKY36, FKY37,

FKY33 and FKY38 in D; FKY2927 in F) were grown overnight at 25 ºC in YPD. PHS was added at

160 µM (C) or 80 µM (D) for 2 hours. Vacuoles were stained with FM4-64 and imaged by

fluorescence microscopy. The number of vacuoles per cell was counted and categorized into one

of three groups. The data represent mean ± SE of three independent experiments, each based on

more than 100 cells. *$p < 0.05$, **$p < 0.01$ and ***$p<0.001$ by Student's t-test compared with WT

(B, D) or empty cells (F). Significant differences analysis between the pairwise combination of

groups was performed using two-way ANOVA.

(E) Illustration showing intracellular utilization pathway of exogenous PHS.

Figure 4. NVJ is required for PHS-induced vacuole fragmentation

(A-C) Cells (FKY2929 in A; FKY3868 and FKY5560 in B; FKY6187, FKY6189, FKY6190, FKY6188 and FKY6409 in C) were grown overnight at 25 ºC in YPD. PHS was added at 40 μM for 2 hours at 30 ºC (A) and 25 ºC (A, B). Vacuoles were stained with FM4-64 and imaged by fluorescence microscopy. The number of vacuoles per cell was counted and categorized into one of three

groups. The data represent mean ± SE of three independent experiments, each based on more

than 100 cells. *$p < 0.05$, **$p < 0.01$ and ***$p<0.001$ by Student's t-test. Significant differences

analysis between the pairwise combination of groups was performed using two-way ANOVA.

(D) Membrane contact sites regulate vacuole morphology via sphingolipid metabolism. See the

main text for details.

Supplementary information

Supplementary Figure 1

(A) Cells (FKY2577, FKY2909, FKY3819, FKY2924, FKY2927, FKY3340 and YKC112-01) were

adjusted to OD_{600} = 1.0 and fivefold serial dilutions were then spotted on YPD plates of indicated

pH, then incubated at 25 ºC.

(B) GFP-Cps1p fusion protein was transformed into WT (FKY2577) and *tcb3Δ* (FKY2924) strains

and analyzed its subcellular localization with respect to FM4-64-stained vacuoles using by

fluorescent microscopy.

Supplementary Figure 2

(A, B) Cells (FKY3340, YKC145-21 and YKC149-61 in A; FKY6187 and FKY6140 in B) were grown

overnight at 25 ºC in YPD. PHS was added at 80 µM for 2 hours, or cells were incubated with 0.2

M of NaCl for 2 hours. Vacuoles were stained with FM4-64 and imaged by fluorescence

microscopy. The number of vacuoles per cell was counted and categorized into one of three

groups. The data represent mean ± SE of three independent experiments, each based on more

than 100 cells. *p < 0.05, **p < 0.01 and ***p<0.001 by Student's t-test. Significant differences

analysis between the pairwise combination of groups was performed using two-way ANOVA.

Supplementary Table 1

Yeast strains used in this study, Related to All Figures.

Strain	Number	MAT	Genotype	Source
WT (RH)	FKY2577	a	*ura3 his3 leu2 trp1 lys2 bar1-1*	Ikeda A et al. iScience. 2020;23(10):101603.
tcb1Δ2Δ3Δ	FKY2927	a	RH except for *tcb1Δ::TRP1 tcb2Δ::HIS3 tcb3Δ::LEU2*	Ikeda A et al. iScience. 2020;23(10):101603.
tcb1Δ	FKY2909	a	RH except for *tcb1Δ::TRP1*	Ikeda A et al. iScience. 2020;23(10):101603.
tcb2Δ	FKY3819	alpha	RH except for *tcb2Δ::HIS3*	Ikeda A et al. iScience. 2020;23(10):101603.
tcb3Δ	FKY2924	a	RH except for *tcb3Δ::LEU2*	Ikeda A et al. iScience. 2020;23(10):101603.
tcb1Δ2Δ	FKY3023	a	RH except for *tcb1Δ::TRP1 tcb2Δ::HIS3*	Ikeda A et al. iScience. 2020;23(10):101603.
tcb1Δ3Δ	FKY3820	alpha	RH except for tcb1Δ::TRP1 tcb3Δ::LEU2	Ikeda A et al. iScience. 2020;23(10):101603.
tcb2Δ3Δ	FKY3008	a	RH except for *tcb2Δ::HIS3 tcb3Δ::LEU2*	Ikeda A et al. iScience. 2020;23(10):101603.
TCB3 (Full)-GBP	FKY3903	a	RH except for *TCB3-GBPΔ::KanMX*	This study
TCB3 (TM-SMP)-GBP	FKY3904	a	RH except for *TCB3* (1F-489R)-*GBP::KanMX*	This study
TCB3 (TM)-GBP	FKY3905	a	RH except for *TCB3* (1F-259R)-*GBP::KanMX*	This study
tcb1Δ2Δ TCB3 (TM)-GBP	FKY4754	a	RH except for *tcb1Δ::TRP1 tcb2Δ::HIS3 TCB3* (1F-259R)::*KanMX*	This study
WT	FKY36 / RH4423	alpha	RH except for *HIS3+ TRP1+ LYS2+ pep4-3*	Funato K et al. J Biol Chem. 2003;278(9):7325-7334.
lcb3Δ	FKY37 / RH4427	alpha	RH except for *lcb3Δ::KanMX HIS3+ LEU2+ TRP1+ LYS2+ pep4-3*	Funato K et al. J Biol Chem. 2003;278(9):7325-7334.
lcb4Δ5Δ	FKY33 / RH4402	alpha	RH except for *lcb4Δ::URA3 lcb5Δ::LEU2 HIS3+ TRP1+ LYS2+ pep4-3*	Funato K et al. J Biol Chem. 2003;278(9):7325-7334.
lcb3Δ4Δ5Δ	FKY38 / RH4464	alpha	RH except for *lcb3Δ::kanMX lcb4Δ::URA3 lcb5Δ::LEU2 HIS3+ TRP1+ LYS2+ pep4-3*	Funato K et al. J Biol Chem. 2003;278(9):7325-7334.
sec18-20	FKY2929	alpha	RH except for *sec18-20 LYS2+*	Ikeda A et al. iScience. 2020;23(10):101603.
WT (RH)	FKY6187	alpha	*ura3 his3 leu2 trp1 lys2 bar1-1*	This study
nvj1Δ	FKY6189	a	RH except for *nvj1Δ::HIS3*	This study
tcb3Δ	FKY6190	alpha	RH except for *tcb3Δ::LEU2*	This study
tcb3Δnvj1Δ	FKY6188	a	RH except for tcb3Δ::*LEU2* nvj1Δ::HIS3	This study
tcb3Δnvj1Δ2Δ3Δmdm1Δ	FKY6409	a	RH except for *nvj1Δ:: URA3 nvj2Δ::KanMX nvj3Δ::HIS3 mdm1Δ::KanMX tcb3Δ::LEU2*	This study
nvj1Δ2Δ3Δ	FKY6140	a	RH except for nvj1Δ:: URA3 nvj2Δ::KanMX nvj3Δ::HIS3	This study
WT (W303)	FKY5687 / RL1	a	*his3-11,15 leu2-3,112 trp1-1 ura3-1 ade2-1 can1-100 GAL+ bar-*	Lucena R et al. Curr Biol. 2018;28(2):196-210.
lag1Δlac1Δ	FKY5688 / RL40	a	W303 except for *lag1Δ::KanMX6 lac1Δ::HIS*	Lucena R et al. Curr Biol. 2018;28(2):196-210.
WT (BY4742)	FKY3340	alpha	*his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0*	Open biosystem, inc
lag1Δ	YKC121-59	alpha	BY4742 except for *lag1Δ::KanMx*	Open biosystem, inc
vma3Δ	YKC112-01	alpha	BY4742 except for *vma3Δ::KanMx*	Open biosystem, inc
fab1Δ	YKC145-21	alpha	BY4742 except for *fab1Δ::KanMx*	Open biosystem, inc
vac14Δ	YKC149-61	alpha	BY4742 except for *vac14Δ::KanMx*	Open biosystem, inc
WT (SEY6210)	FKY3868	alpha	*ura3-52 leu2-3,112 his3-Δ100 trp1-Δ901 lys2-801 suc2-Δ9*	Yabuki Y et al. Genetics. 2019;212(1):175-186.
ΔNVJ	FKY5560	alpha	SEY6210 except for *nvj1Δ::TRP1 nvj2Δ::HIS3 nvj3Δ::NatMX mdm1Δ::KanMX*	Henne WM et al. J Cell Biol. 2015;210(4):541-551.

767 **Supplementary Table 2**

768 Plasmids used in this study, Related to All Figures.

Plasmid	Number	Relevant information	Source
pRS426-TDH3	FKP51	2µ, yeast *TDH3* promoter, empty, *URA3*	Mumberg D et al., Gene. 1995;156(1):119-122
pRS426-TDH3/RSB1	FKP1065	2µ, yeast *TDH3* promoter, *RSB1*, *URA3*	This study
pRS316/PTDH3-EGFP-CPS1-TCMK1	FKP621	CEN, yeast *TDH3* promoter, GFP-*CPS1* *URA3*	Howard Riezman