In-vivo whole-cortex estimation of excitation-inhibition ratio indexes cortical maturation and cognitive ability in youth

Shaoshi Zhang[1,2,3,4,*], Bart Larsen[6,7,8*], Valerie J. Sydnor[6,7,8*], Tianchu Zeng[1,2,3], Lijun An[1,2,3], Xiaoxuan Yan[1,2,3,4], Ru Kong[1,2,3], Xiaolu Kong[1,2,3,5], Ruben C. Gur[7,8,9], Raquel E. Gur[7,8,9], Tyler M. Moore[7,8], Daniel H. Wolf[7], Avram J Holmes [10,11], Yapei Xie[1,2,4], Juan Helen Zhou[1,2,4], Marielle V Fortier[12,13], Ai Peng Tan[13,14], Peter Gluckman[15], Yap Seng Chong[13,16], Michael Meaney[13,17], Gustavo Deco[18,19], Theodore D. Satterthwaite[6,7,8+], B.T. Thomas Yeo[1,2,3,4,20+]

[1]Centre for Sleep & Cognition & Centre for Translational Magnetic Resonance Research, Yong Loo Lin School of Medicine, Singapore; [2]Department of Electrical and Computer Engineering, National University of Singapore, Singapore; [3]N.1 Institute for Health & Institute for Digital Medicine, National University of Singapore, Singapore; [4]Integrative Sciences and Engineering Programme (ISEP), National University of Singapore, Singapore, Singapore; [5]ByteDance, Singapore; [6]Penn Lifespan Informatics and Neuroimaging Center, University of Pennsylvania, Philadelphia, PA 19104, USA; [7]Department of Psychiatry, University of Pennsylvania, Philadelphia, PA 19104, USA; [8]Lifespan Brain Institute (LiBI) of Penn Medicine and CHOP, University of Pennsylvania, Philadelphia, PA 19104, USA; [9]Department of Radiology, University of Pennsylvania, Philadelphia, PA 19104, USA; [10]Department of Psychiatry, Brain Health Institute, Rutgers University, Piscataway, NJ, United States; [11]Wu Tsai Institute, Yale University, New Haven, CT, United States; [12]Department of Diagnostic and Interventional Imaging, KK Women's and Children's Hospital, Singapore; [13]Singapore Institute for Clinical Sciences (SICS), Agency for Science, Technology and Research (A*STAR), Singapore; [14]Department of Diagnostic Radiology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore; [15]UK Centre for Human Evolution, Adaptation and Disease, Liggins Institute, University of Auckland, Auckland, New Zealand; [16]Department of Obstetrics and Gynaecology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore; [17]Department of Neurology and Neurosurgery, McGill University, Montreal, QC, Canada;[18]Center for Brain and Cognition, Department of Technology and Information, Universitat Pompeu Fabra, Barcelona, Spain; [19]Institució Catalana de la Recerca i Estudis Avançats, Universitat Barcelona, Barcelona, Spain; [20]Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Charlestown, USA
* equal contributions; + equal contributions

Address correspondence to:
B.T. Thomas Yeo
CSC, TMR, ECE, N.1, WISDM
National University of Singapore
Email: thomas.yeo@nus.edu.sg

Theodore D. Satterthwaite
Penn Lifespan Informatics and Neuroimaging Center
University of Pennsylvania Perelman School of Medicine
Email: sattertt@pennmedicine.upenn.edu

Abstract

A balanced excitation-inhibition ratio (E/I ratio) is critical for healthy brain function, but challenging to measure *in-vivo* in humans. Here we present an approach to non-invasively estimate whole-cortex E/I ratio by fitting a large-scale biophysically plausible model to resting-state functional MRI (fMRI) data. We first confirm our model generates realistic brain dynamics in the Human Connectome Project. Next, we show that the estimated E/I ratio is sensitive to the GABA-agonist benzodiazepine alprazolam during fMRI. Furthermore, drug-induced E/I changes are spatially consistent with positron emission tomography measurement of benzodiazepine receptor density. We then leverage the model to investigate the hypothesis that the E/I ratio is normatively refined during development and supports improvement in cognitive ability. We find that the E/I ratio declines heterogeneously across the cerebral cortex during youth, with the greatest reduction occurring in sensorimotor systems relative to association systems. Importantly, among children with the same chronological age, a lower E/I ratio (especially in association cortex) is linked to better cognitive performance. This result is replicated across North American (8.2 to 23.0 years old) and Asian (7.2 to 7.9 years old) cohorts, suggesting that a more mature E/I ratio indexes improved cognition during normative development. Overall, our approach opens the door to studying cortex-wide E/I ratio changes across the lifespan and in neuropsychiatric disorders.

1. Introduction

Healthy brain function requires a delicate balance between neural excitation (E) and inhibition (I) (Nelson, 1991; Turrigiano & Nelson, 2004; Atallah & Scanziani, 2009; Dorrn et al., 2010; Sun et al., 2010). This balance – the E/I ratio – is refined during critical developmental periods of heightened experience-dependent plasticity when cortical circuits are strongly sculpted by an individual's environment (Takesian & Hensch, 2013). E/I imbalances during critical developmental periods are thought to contribute to the etiology of psychiatric disorders, such as schizophrenia (Anticevic & Lisman, 2017; Anticevic & Murray, 2017; Shaw et al., 2019; Dienel et al., 2023; Vinogradov et al., 2023) and depression (Duman et al., 2019; Anderson et al., 2020b; Larsen et al., 2022; Yao et al., 2022). Importantly, a transdiagnostic feature of psychiatric disorders linked to E/I imbalances is the presence of cognitive deficits (Fossati et al., 1999; Romer & Pizzagalli, 2021). Overall, disrupted E/I ratio development may contribute to adulthood psychopathology risk and confer vulnerability to cognitive deficits (Lam et al., 2022), but it remains sparsely explored. Here we capitalize on advances in large-scale circuit models to chart the normative development of cortex-wide E/I ratio and uncover links to cognition.

Human cortical development unfolds hierarchically – sensory systems mature earlier, while association systems follow a more protracted developmental course extending through adolescence (Gogtay & Thompson, 2010; Gennatas et al., 2017; Paquola et al., 2019; Dong et al., 2021; Pines et al., 2022; Baum et al., 2022; Bethlehem et al., 2022; Sydnor et al., 2023). This hierarchical, sensorimotor-to-association maturation axis (Sydnor et al., 2021) is thought to support the development of increasingly complex cognitive abilities throughout youth, with earlier development of unimodal cortices supporting sensory processes and later maturation of association cortices supporting higher-order cognitive abilities (Casey et al., 2008; Steinberg, 2010; Luna & Wright, 2016; Shulman et al., 2016; Larsen & Luna, 2018; Reh et al., 2020; Dong et al., 2021; Sydnor et al., 2021; Canetta et al., 2022; Tervo-Clemmens et al., 2022; Keller et al., 2023). The mechanism driving this hierarchical developmental pattern might correspond to the temporal progression of critical periods of plasticity beginning in unimodal sensory systems, progressing through heteromodal systems, and terminating in transmodal executive systems (Werker & Hensch, 2015; Larsen & Luna, 2018; Reh et al., 2020; Keller et al., 2023; Sydnor & Satterthwaite, 2023). A central feature is the strengthening of local inhibitory circuitry in juvenile cortex, which facilitates an increase in the experience-dependent pruning of excitatory pyramidal neuronal connections and

triggers the opening of a critical period of plasticity (Iwai et al., 2003; Hensch, 2005; Toyoizumi et al., 2013; Gogolla et al., 2014; Trachtenberg, 2015; Gu et al., 2016; Wu et al., 2022). As a result, a hallmark feature of critical period development is a reduction in the E/I ratio (Hensch, 2005; Hensch & Fagiolini, 2005). The decline of the E/I ratio can therefore be understood as a marker of maturation along cortical hierarchies. While the hierarchical progression of inhibitory development has been documented in sensory systems of animal models (Akil & Lewis, 1992; Condé et al., 1994; Stern et al., 2001; Reh et al., 2020), it is unclear if the same mechanisms exist in humans, extend to the evolutionarily expanded association cortices and impact cognitive ability.

Studying E/I ratio development *in-vivo* in humans is challenging due to limitations in investigating excitatory and inhibitory activities with non-invasive neuroimaging techniques. MR spectroscopy studies provided evidence for changes in the balance and covariance of excitatory and inhibitory neurotransmitter levels in single brain regions, such as the anterior insula, cingulate, and prefrontal cortex during adolescence (Silveri et al., 2013; Ghisleni et al., 2015; Perica et al., 2022). A recent study used a machine learning biomarker trained on pharmacological-fMRI data to provide initial evidence of E/I ratio reductions in the association cortices in adolescence (Larsen et al., 2022). However, past studies characterizing E/I ratio maturation were limited to individual brain regions or partial portions of the cortex, so cortex-wide E/I ratio developmental patterns have not been characterized. These prior studies also lacked a direct mapping to an underlying mechanistic model of excitatory and inhibitory dynamics.

To overcome the limitations of conventional *in-vivo* neuroimaging approaches, large-scale biophysically plausible models of coupled brain regions were developed to provide mechanistic insights into spontaneous brain dynamics (Honey et al., 2007; Ghosh et al., 2008). However, most large-scale circuit models assume that local synaptic properties are spatially homogeneous across brain regions (Deco et al., 2014; Trakoshis et al., 2020; Lam et al., 2022),which lacks biological plausibility. Spatial heterogeneity in excitatory and inhibitory cell types (Kim et al., 2017; Burt et al., 2018; Anderson et al., 2020a) might be a driver of large-scale brain dynamics (Cardin et al., 2009; Deco et al., 2021; Froudist-Walsh et al., 2021). Indeed, studies have shown that incorporating spatial heterogeneity across local synaptic parameters generates more realistic spontaneous brain dynamics (Demirtaş et al., 2019; Wang et al., 2019; Deco et al., 2021; Kong et al., 2021). Our previous study (Kong et al., 2021) demonstrated that parameterizing local synaptic parameters with anatomical and functional gradients led to dramatically more realistic brain dynamics in adults. However, we

utilized a large-scale circuit model (Deco et al., 2013) that did not differentiate among excitatory and inhibitory neural populations, so the E/I ratio could not be directly derived. Implementing a circuit model that can directly assess the E/I ratio is critical for establishing links with neurodevelopment and cognition.

Here we investigate the development of cortical E/I ratio over youth and its association with cognitive ability. To do so, we apply our previous approach (Kong et al., 2021) to the feedback inhibition control (FIC) model with coupled excitatory and inhibitory neuronal populations (Deco et al., 2014). Motivated by invasive rodent experiments (Xue et al., 2014), we hypothesize that the ratio of excitatory and inhibitory neural signals of the resulting parametric FIC (pFIC) model would provide a biologically plausible estimate of whole-cortex E/I ratio. We first confirm that our pFIC model yields realistic brain dynamics in new out-of-sample participants in a sample of healthy young adults ($n = 1,004$) from the Human Connectome Project (Van Essen et al., 2013). We next validate the pFIC model using a pharmacological fMRI dataset ($n = 45$) with the GABA-ergic drug alprazolam (Wolf et al., 2011) to establish that the pFIC model is sensitive to a pharmacological reduction in the E/I ratio. Then, using a developmental neuroimaging dataset ($n = 885$), the Philadelphia Neurodevelopmental Cohort (Satterthwaite et al., 2014; Calkins et al., 2015), we characterize the development of cortex-wide E/I ratio over youth and delineate links to individual differences in cognition. Notably, we find that the E/I ratio declines across the cortex during youth and that a lower E/I ratio is associated with greater cognitive ability, with the strongest relationships to cognition observed in association systems. Finally, we replicate the link between E/I ratio and cognitive performance in an independent Asian youth cohort, the GUSTO (Growing Up in Singapore with Healthy Outcomes) dataset ($n = 154$) (Soh et al., 2014). Overall, our study suggests that E/I ratio maturation is a driver of healthy neurocognitive development during youth and a potential target of interventions for neuropsychiatric disorders (Paus et al., 2008; Insel, 2014; Marín, 2016; Nelson & Gabard-Durnam, 2020; Vinogradov et al., 2023).

2. Results

2.1 Overview

We first evaluated the optimization of the spatially heterogeneous parametric feedback inhibition control (pFIC) model in the Human Connectome Project (HCP) dataset (**Figure 1A**). The biological plausibility of the estimated excitation-inhibition (E/I) ratio was then evaluated using pharmacological fMRI involving a GABAergic benzodiazepine challenge. Finally, we investigated developmental changes and cognitive effects of E/I ratio in the Philadelphia Neurodevelopmental Cohort (PNC); assocations with cognition were replicated in the Growing Up in Singapore with Healthy Outcomes (GUSTO) dataset.

2.2 Optimization of the parametric feedback inhibition control (pFIC) model

We began by optimizing the pFIC model. We randomly divided 1004 participants from the Human Connectome Project (HCP) S1200 release (Glasser et al., 2013; Van Essen et al., 2013) into 3 non-overlapping training (N = 335), validation (N = 335), and test (N = 334) sets. The Desikan–Killiany anatomical parcellation (Desikan et al., 2006) with 68 cortical regions of interest (ROIs) was used to generate group-level structural connectivity (SC), static functional connectivity (FC), and time-varying functional connectivity dynamics (FCD) from the training, validation, and test sets separately. See Section 4.2 for more details.

The FIC model (Deco et al., 2014) comprised differential equations at each cortical region describing the dynamics of excitatory and inhibitory neuronal populations (**Figure 1B** top panel). The local dynamics are driven by recurrent connections within the excitatory and inhibitory populations, as well as connections between excitatory and inhibitory populations. Greater excitatory-to-excitatory recurrent strength (w_{EE}) and smaller inhibitory-to-excitatory connection strength (w_{IE}) amplifies synaptic currents of the excitatory population. Similarly, greater excitatory-to-inhibitory connection strength (w_{EI}) and smaller inhibitory-to-inhibitory recurrent strength (w_{II}) amplifies synaptic currents of the inhibitory population. Neuronal noise in each cortical region is controlled by the noise amplitude σ. Finally, the excitatory populations of the regional local models are connected via the SC matrix, scaled by a global constant G.

(A) Workflow

(B) Feedback Inhibition Control (FIC) Model

(C) Parameterization of local synaptic parameters

Figure 1. Workflow and schematic of the parametric feedback inhibition control (pFIC) model. (A) Young adults from the Human Connectome Project (HCP) were used to evaluate the optimization of the spatially heterogeneous pFIC model. Pharmacological fMRI with the benzodiazepine alprazolam was then used to evaluate the biological plausibility of the estimated E/I ratio. Next, the pFIC model was used to investigate the development of E/I ratio across the cortex and its association with cognitive ability in the Philadelphia Neurodevelopmental Cohort (PNC). Finally, cognitive associations were replicated in a sample of seven-year-olds from the GUSTO dataset. HCP logo is used with permission from the HCP team. (B) The FIC model (Deco et al., 2014) consisted of differential equations at each cortical region governing the neural dynamics of excitatory and inhibitory neuronal populations ("E" and "I" respectively in the left panel). A red triangle indicates an excitatory

connection, while a blue circle indicates an inhibitory connection. "w_{xy}" indicates the strength of connection from neuronal population x to neuronal population y. For example, "w_{IE}" indicates the strength of connection from the inhibitory population to excitatory population. The regional models were connected by excitatory connections parameterized by a structural connectivity (SC) matrix. For a particular set of model parameters, time courses of excitatory (S_E) and inhibitory (S_I) synaptic gating variables (representing the fraction of open channels) were simulated. In this study, the excitation-inhibition ratio (E/I ratio) was defined as the ratio between the temporal average of S_E and S_I. (C) Local synaptic parameters were parameterized as a linear combination of the principal resting-state functional connectivity (FC) gradient (left; Margulies et al., 2016) and the T1w/T2w ratio (right) (Glasser & Van Essen, 2011) maps, following our previous study (Kong et al., 2021). We refer to the resulting model as the pFIC model.

Following previous studies (Deco et al., 2014; Demirtaş et al., 2019),w_{II} was set to one and w_{IE} was automatically set to maintain a uniform baseline excitatory firing rate of ~3Hz across regions. Excitatory-to-excitatory recurrent strength (w_{EE}), excitatory-to-inhibitory connection strength (w_{EI}), regional noise amplitude (σ) and the SC scaling constant (G) were estimated using our previous optimization approach (Kong et al., 2021). As illustrated in **Figure 1C**, w_{EE}, w_{EI} and σ were parameterized as a linear combination of the principal resting-state FC gradient (Margulies et al., 2016) and T1w/T2w myelin estimate (Glasser & Van Essen, 2011), resulting in 9 unknown linear coefficients and 1 unknown parameter G. We refer to the resulting model as the parametric FIC (pFIC) model.

The 10 parameters of the pFIC model were estimated using the covariance matrix adaptation evolution strategy (CMA-ES) (Hansen, 2006) by minimizing the difference between simulated and empirical fMRI data. More specifically, agreement between empirical and simulated FC matrices was defined as the Pearson's correlation (r) between the z-transformed upper triangular entries of the two matrices. Larger r indicates more similar static FC. However, Pearson's correlation does not account for scale difference, so we also computed the absolute difference (d) between the means of the empirical and simulated FC matrices (Demirtaş et al., 2019). A smaller d indicates more similar static FC. Finally, disagreement between the simulated and empirical FCD matrices was defined as the Kolmogorov–Smirnov (KS) distance between the upper triangular entries of the two matrices (Hansen et al., 2015; Kong et al., 2021). The overall cost was defined as $(1 - r) + d + KS$. A smaller cost indicates better agreement between simulated and empirical fMRI.

Figure 2. **The parametric feedback inhibition control (pFIC) model generates more realistic static functional connectivity (FC) and functional connectivity dynamics (FCD) than the spatially homogeneous FIC model.** (A) The CMA-ES algorithm (Hansen, 2006;

Kong et al., 2021) was applied to the HCP training set to generate 500 sets of model parameters. The top 10 parameter sets from the validation set were evaluated in the test set. (B) Empirical FC from the test set. (C) Simulated FC from the pFIC model using the best model parameters (from the validation set) and SC from the test set. (D) Agreement (Pearson's correlation) between empirical and simulated static FC in the HCP test set. (E) Empirical FCD from an HCP test participant. (F) Simulated FCD from the pFIC model using the best model parameters (from the validation set) and SC from the test set. (G) Total test cost of the pFIC model compared with three control conditions: (1) local synaptic parameters parameterized by only principal resting-state FC gradient, (2) local synaptic parameters parametrized by only T1w/T2w ratio map and (3) local synaptic parameters constrained to be spatially uniform. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers. * indicates that the pFIC model achieved statistically better (lower) total test cost.

2.3 The pFIC model generates realistic fMRI dynamics

We first validated the optimization of the pFIC model in the HCP dataset (**Figure 2A**). Our goal was to demonstrate that the parametrization of the local synaptic parameters with T1w/T2w and FC gradient led to more realistic brain dynamics than spatially homogeneous parameters. We applied CMA-ES to the HCP training set to generate 500 candidate model parameter sets. The 500 parameter sets were evaluated in the HCP validation set. The top 10 parameter sets from the HCP validation set were used to simulate FC and FCD using SC from the HCP test set, which were then compared with empirical FC and FCD from the HCP test set. A strong agreement between simulated and empirical FC (as well as between simulated and empirical FCD) would suggest that the pFIC model was able to generate realistic brain dynamics.

Figure 2B shows the empirical FC in the HCP test set, while **Figure 2C** shows the simulated FC using the best model parameters (from the validation set) and SC from the test set. Across the 10 best parameter sets from the validation set, correlation between empirical and simulated static FC was 0.71 ± 0.005 (mean $\pm$ std) in the test set (**Figure 2D**). As a reference, correlation between empirical FC and SC in the test set was 0.48. On the other hand, the absolute difference between the means of the empirical and simulated FC matrices was 0.11 ± 0.015 in the test set. This suggests that the pFIC model was able to generate realistic FC.

Figure 2E shows the empirical FCD of a representative HCP test participant. **Figure 2F** shows the simulated FCD using the best model parameters (from the validation set) and SC from the test set. Off-diagonal red blocks in both empirical and simulated FCD indicated recurring FC patterns that were not simply due to temporal autocorrelation. Therefore,

similarity in the amount of off-diagonal red blocks between empirical and simulated FCD suggests that the pFIC model was able to generate realistic FCD. Across the 10 best candidate sets from the validation set, the Kolmogorov-Smirnov (KS) distance between empirical and simulated FCD was 0.18 ± 0.028 (mean $\pm$ std) in the HCP test set.

Overall, the pFIC model was able to generate realistic FC and FCD, yielding an overall cost of 0.58 ± 0.018 in the HCP test set (**Figure 2G**). Parameterizing model parameters with only the principal FC gradient or only T1w/T2w ratio map led to worse (higher) cost in the HCP test set (**Figure 2G**). Most large-scale circuit model studies assume spatially homogeneous parameters. When local synaptic parameters (w_{EE}, w_{EI} and σ) were constrained to be uniform across brain regions (Deco et al., 2014; Kringelbach & Deco, 2020) and optimized by CMA-ES, the cost was poor in the test set (**Figure 2G**). These results emphasize the importance of parameterizing local synaptic parameters with spatial gradients that smoothly varied from sensory-motor to association cortex. Consistent with our previous study (Kong et al., 2021), the T1w/T2w and FC gradient were complementary in the sense that combining the two spatial maps led to more realistic fMRI dynamics.

2.4 Estimated E/I ratio is sensitive to the effect of benzodiazepine alprazolam

In the previous section, we showed that the pFIC model could be effectively optimized to generate realistic fMRI dynamics. Here, we evaluated the biological plausibility of the estimated E/I ratio in a pharmacological-fMRI dataset (Wolf et al., 2011), which included 45 participants who had completed a placebo-controlled double-blind fMRI study with benzodiazepine alprazolam. Alprazolam is a benzodiazepine that enhances GABAergic signaling at GABA$_A$ receptor sub-units, including $\alpha_{1,2,3,5}$ and γ_{1-3} (Sigel, 2002; Olsen & Sieghart, 2009). Alprazolam enhances GABAergic inhibitory signaling via positive allosteric modulation, thus reducing E/I ratio (Verster & Volkerts, 2006). Therefore, we hypothesized that the E/I ratio estimated with the pFIC model would be lower during the alprazolam condition compared with the placebo condition.

The 45 participants were equally divided into training, validation and test sets. Group-level SC, first principal FC gradient, and T1w/T2w ratio maps from the HCP dataset were used in the following analysis. For each experimental condition (placebo or alprazolam), 250 candidate parameter sets were generated from the condition's training set. The top 10 parameter sets from the validation set were evaluated in the test set. The costs of the 10

parameter sets generalized well to the test set (**Figure S1**), suggesting that there was no overfitting in the validation set.

One challenge in analyzing this dataset was that the fMRI data was acquired with a limited field of view. Therefore, 26 out of 68 Desikan-Killiany ROIs with less than 50% coverage (**Figure S2**) were not considered during the estimation of the model parameters. The estimated model parameters were then extrapolated to the entire cortex (see Section 4.11 for details) and used to simulate the excitatory (S_E) and inhibitory (S_I) time courses (Figure 1B). Motivated by rodent studies, the E/I ratio was defined as the ratio between the temporal average of S_E and S_I (Xue et al., 2014).

An E/I ratio contrast was computed by subtracting the E/I ratio estimated during the drug (alprazolam) session from the E/I ratio estimated during the placebo session. Since alprazolam is expected to reduce E/I ratio, we hypothesized the E/I ratio would be lower during the alprazolam condition, yielding a positive E/I ratio contrast. Consistent with our hypothesis, the E/I ratio contrasts of all regions were positive (**Figure 3B**). Specifically, 67 out of 68 regions exhibited E/I ratio contrasts statistically different from zero after correcting for multiple comparisons with a false discovery rate (FDR) of $q < 0.05$. These results suggest that the E/I ratio estimated by the pFIC model was sensitive to the pharmacological enhancement of inhibitory activities.

Since the distribution of benzodiazepine receptors (BZR) density is not spatially uniform (Nørgaard et al., 2021),we hypothesized that the E/I ratio contrast would also not be spatially uniform, and would align with BZR density. Supporting this, we found that the E/I ratio contrast exhibited a spatial gradient with strongest effects in sensory-motor networks and weakest effects in control and default networks (**Figure 3B**). **Figure 3C** shows the spatial distribution of benzodiazepine receptors (BZR) density estimated from *in-vivo* positron emission tomography in a separate group of participants (Nørgaard et al., 2021). Regions with greater BZR density exhibited greater reduction in E/I ratio during the drug session (r = 0.52; two-tail spin test p = 0.016; **Figure 3D**). Overall, the spatial distribution of E/I ratio contrast was biologically plausible.

To evaluate robustness, we repeated the above analyses 5 times with different random splits of the 45 participants into training, validation and test sets. The results were similar across the 5 splits (**Figures S3 and S4**); results from the most representative split were shown in **Figure 3**. Using this most representative split, we performed several additional sensitivity analyses. In the previous analyses, the acceptable excitatory firing rate was constrained to be between 2.7Hz and 3.3Hz. Relaxing the thresholds to between 2.5Hz and 3.5Hz yielded

similar results (**Figure S5**). Changing the ROI coverage threshold from 50% to 60% also yielded similar results (**Figure S6**). Finally, we repeated the analysis using a 100-region homotopic functional parcellation (Yan et al., 2023), which also yielded similar results (**Figure S7**). Pairwise comparisons between the control analyses are found in **Figure S8**.

Figure 3. E/I ratio estimate is sensitive to the effect of benzodiazepine alprazolam. (A) Seven resting-state networks (Yeo et al., 2011). (B) Left: Regional E/I ratio contrast overlaid with the boundaries (black) of seven resting-state networks. 67 out of 68 regions showed significant E/I ratio difference between placebo and drug sessions after FDR correction (q < 0.05). E/I ratio difference was greater than zero for all regions, consistent with lower E/I ratio during the alprazolam session. Right: E/I ratio differences exhibited a spatial gradient with higher differences in sensory-motor regions compared with regions in the control and default networks. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate

1.5 IQR. Black crosses represent outliers. (C) Spatial distribution of BZR density (pmol/ml) from *in-vivo* positron emission tomography in a separate group of participants (Nørgaard et al., 2021). (D) Higher regional BZR density was associated with larger E/I ratio changes during the drug session (r = 0.52, two-tail spin test p = 0.016).

2.5 The E/I ratio declines with development in youth

Having demonstrated that the E/I ratio estimates were sensitive to the enhancement of inhibitory activities induced by alprazolam, we next explored how the E/I ratio changes during development in the Philadelphia Neurodevelopmental Cohort (PNC) (Satterthwaite et al., 2014; Calkins et al., 2015). We hypothesized that the estimated E/I ratio would decline with age.

After data preprocessing and quality control, we obtained a sample of 885 participants ages 8-23 years old (**Figure 4A**). Participants were sorted according to age and evenly divided into 29 age groups, so each group comprised 30 or 31 participants. Within each age group, 15 participants were randomly selected as the validation set, while the remaining participants were assigned to the training set. For each age group, 250 candidate model parameter sets were generated from the group's training set using CMA-ES and evaluated in the group's validation set; the parameter set with the lowest validation cost was used to estimate regional E/I ratio across the cortex.

Mean cortical E/I ratio (i.e., E/I ratio averaged across the whole cortex) continuously declined throughout child and adolescent development (r = -0.51, p = 0.004; **Figure 4B**). This reduction in E/I ratio was statistically significant for all cortical regions (FDR q < 0.05; **Figure 4C**). Furthermore, the rate of E/I ratio decrease exhibited a spatial gradient with sensory-motor regions exhibiting greater rate of E/I ratio decrease (i.e., a more negative age slope) compared with association networks (**Figure 4D**).

To evaluate the robustness of these effects, the PNC analyses were repeated 5 times with different splits of the participants (within each age group) into training and validation sets. The results were similar across the 5 random splits of the data (**Figures S9 and S10**). We conducted several additional sensitivity analyses using the most representitve split (which was shown in **Figure 4**). Relaxing the firing rate thresholds to between 2.5Hz and 3.5Hz yielded similar results (**Figure S11**), as did using a 100-region homotopic parcellation (Yan et al., 2023) (**Figure S12**). Pairwise comparisons between the control analyses are found in **Figure S13**.

Figure 4. E/I ratio continuously declines throughout child and adolescent development.
(A) Age distribution of 885 PNC participants (mean = 15.66, std = 3.36, min = 8.17, max = 23). (B) Participants in older age groups exhibited lower E/I ratio (r = -0.51, p = 0.004). Participants were divided into 29 non-overlapping age groups. There are 29 dots in the scatter plot, corresponding to the 29 age groups. The shaded area depicts 95% confidence interval of the linear relationship. (C) Spatial distribution of linear regression slope between regional E/I ratio and age. The values represent the rate of E/I ratio changes during development. All slopes were negative and significant (FDR *q* < 0.05). (D) The slopes exhibited a spatial gradient with sensory-motor networks showing the fastest E/I ratio reduction and association networks showing slower E/I ratio reduction. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers.

2.6 Lower E/I ratio is associated with better cognition within the same age group

We have shown that older children exhibited a lower E/I ratio (**Figure 3B**). We next evaluated the cognitive implications of such a decline in the E/I ratio as part of normative development, hypothesizing that a lower E/I ratio would be associated with better cognition. To test this hypothesis, we compared the E/I ratio of PNC participants who were matched on age but differed in cognitive performance.

Participants in the PNC completed the Penn Computerized Neurocognitive Battery (CNB), a 12-task battery that has been previously summarized using an overall (domain-general) measure of accuracy as well as three domain-specific factor scores (Moore et al.,

2015). Participants were divided into 14 high-performance groups and 14 low-performance groups based on the overall accuracy measure. Each high-performance group was age-matched to a low-performance group (**Figure 5A**). Each low-performance or high-performance group comprised 31 or 32 participants. For each group, 15 participants were randomly assigned to the validation set, while the remaining participants were assigned to the training set. For each group, 250 candidate parameter sets were generated from the training set and the top parameter set from the validation set was used to estimate the E/I ratio; we compared the E/I ratio between the high and low performance groups.

We observed that the high-performance group exhibited lower mean cortical E/I ratio than the low-performance group ($p = 1.2$ x 10^{-3}; **Figure 5C**). We note that there was no motion difference between high-performance and low-performance groups (FDRMS: $p = 0.38$; DVARS: $p = 0.24$). To test for domain specificity, we also compared the E/I ratio for the three domain-specifc factor scores (complex reasoning, memory and social cognition), but observed no statistical difference after correcting for multiple comparisons (**Figure S14**).

Having found global differences in the E/I ratio between the high and low cognitive performance groups, we next evaluated regional effects (**Figure 5D**). We found that E/I ratio differences between low-performance and high-performance groups were larger in transmodal regions, such as the control and default networks compared with sensory-motor regions (**Figure 5E**; all FDR $q <0.05$). Notably, the effect sizes of these regional differences in the E/I ratio aligned well with the sensorimotor-association (S-A) axis of cortical organization, such that effect sizes were lowest at the sensorimotor pole and largest at the association pole (Sydnor et al., 2021) (**Figure 5F**). The Spearman's correlation between effect sizes and S-A axis ranks was $\rho = 0.87$ (two-tailed spin test $p < 0.001$; **Figure 5G**). Overall, these results suggest that a more mature E/I ratio – especially in higher-order association cortex – is linked to more mature cognition.

To evaluate the robustness of these results, we repeated these analyses 5 times with different random training-validation splits of participants within each high-performance group and each low-performance group. The results were similar across the 5 splits (**Figures S15 to S19**; the most representative split is displayed in **Figure 5**). Within the most representative split, we found that relaxing the thresholds to between 2.5Hz and 3.5Hz yielded similar results (**Figure S20**) as did use of a 100-region homotopic functional parcellation (Yan et al., 2023) (**Figure S21**). Pairwise comparisons between the control analyses are found in **Figure S22**.

Figure 5. Lower E/I ratio is associated with better cognitive performance within the same age group in the PNC cohort. (A) Boxplots of age, (B) 'overall accuracy' and (C) mean cortical E/I ratio of high-performance and low-performance (overall accuracy) groups. The mean cortical E/I ratio of the high-performance group was significantly lower than that of the low-performance group (FDR q < 0.05). (D) Spatial distribution of effect size of regional E/I ratio difference between high-performance and low-performance groups. All effect sizes were significant after FDR correction with q < 0.05. (E) Effect size of E/I ratio differences in cognition is larger in control and default networks compared with sensory-motor regions. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers. (F) ROI rankings along the sensorimotor-association (S-A) axis. Lower ranks were assigned to ROIs that were more towards the sensorimotor pole; higher ranks were assigned to ROIs that were more towards the association pole. (G) Agreement between effect size of E/I ratio difference and S-A axis rank. Spearman's correlation r = 0.87, two-tailed spin test p < 0.001.

2.7 Results replicate and generalize to a younger Asian cohort

As a final step, we evaluated whether the link between the E/I ratio and cognition generalized to a group of younger participants of different anscestry. This was motivated by recent concerns that relationships between resting-fMRI and behavior may not generalize well across ethnic groups (Li et al., 2022). To replicate and generalize our findings, we utilized the Growing Up in Singapore with Healthy Outcomes (GUSTO) dataset (Soh et al., 2014), which included 154 participants (after quality control) with a mean age of 7.5 years. As in the PNC, participants were divided into groups with high and low cognitive performance, which was summarized using a principal component analysis of 5 cognitive tests. The ages were well-matched between the high and low-performance groups (**Figures 6A and 6B**). There was no motion difference between the high and low-performance groups during the fMRI scans (FDRMS: $p = 0.14$; DVARS: $p = 0.11$).

Replicating previous results in the PNC, we found that the high-performance group exhibited a lower E/I ratio in higher-order association cortex than the low-performance group (**Figures 6C** to **6E**). Differences were largest in the default and control networks (**Figure 6F**). In contrast, differences in the E/I ratio between cognitive performance groups was largely not significant in sensory-motor networks after correcting for multiple comparisons. As in the PNC, we found the effect sizes of these cognitive differences aligned with the S-A axis ($r = 0.56$, two-tailed spin test $p = 0.01$; **Figure 6G**). These results in a younger Asian cohort emphasize the robustness and generalizability of our findings.

Figure 6. Lower E/I ratio is associated with better cognitive performance within the same age group in the GUSTO dataset. (A) Boxplots of age for high-performance and low-performance groups. (B) Overall cognition of high-performance and low-performance groups. (C) Spatial distribution of negative log-transformed p-values of regional E/I ratio difference between low-performance and high-performance groups. (D) Out of 68 regions, p-values of 29 regions were significant after FDR correction with q < 0.05 (blue regions). Regions with p-values that were not significant after FDR correction are shown in gray. (E) Spatial distribution of effect size of E/I ratio difference between low-performance group and high-performance group. (F) Effect size of E/I ratio differences in cognition is larger in control and default networks compared with sensory-motor regions. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers. (G) Agreement between effect size of E/I ratio difference and S-A axis rank. Spearman's correlation r = 0.56, two-tailed spin test p = 0.01.

3. Discussion

Leveraging advances in large-scale circuit models, we showed that the E/I ratio declines during youth and is associated with cognitive ability. Using a large adult fMRI dataset, we first established the pFIC model's ability to generate realistic fMRI signal that captures the dynamics of empirical fMRI data. We then demonstrated the biological validity of the pFIC model by showing it was sensitive to a pharmacological reduction of the E/I ratio induced by the benzodiazepine alprazolam. Applying the validated pFIC model to a large developmental neuroimaging sample from North America, we found that the E/I ratio decreased with age. We finally demonstrated that a lower E/I ratio—reflective of more mature cortex—was associated with better cognitive performance, particularly in executive systems within transmodal association cortex. Critically, these findings generalized to a younger Asian cohort. Together, our findings provide important evidence that refinements in the cortical E/I ratio persist into adolescence and suggest that prolonged E/I-linked developmental plasticity in the association cortex supports continued neurocognitive development. Overall, this study capitalizes upon our advanced computational modeling approach to highlight that insufficient refinement of the E/I ratio during development may create a vulnerability to cognitive deficits, with potentially important implications for transdiagnostic psychopathology.

The cortical E/I ratio has been challenging to investigate non-invasively in humans using available neuroimaging approaches. Post-mortem studies have been key to establishing how gradients in the expression of E/I relevant genes vary across the cortex (Burt et al., 2018; Gandal et al., 2018; Huntenburg et al., 2018; Deco et al., 2021; Goulas et al., 2021). The use of *in-vivo* neuroimaging has been limited by the lack of a direct mapping of signals derived from imaging techniques with excitatory and inhibitory neurobiology, as well as constrained spatial coverage and specificity of available E/I techniques (Silveri et al., 2013; Ghisleni et al., 2015; Larsen et al., 2022; Perica et al., 2022). To overcome these limitations, we capitalized on recent developments in biologically interpretable computational modeling of cortical circuits to gain insight into the E/I ratio using *in-vivo* fMRI data. Specifically, we fitted a large-scale circuit model with interacting excitatory and inhibitory populations to resting-state fMRI data and calculated the E/I ratio from the time courses of excitatory and inhibitory synaptic gating variables. The E/I ratio generated from the model parameters captured reductions in the E/I ratio induced by the benzodiazepine alprazolam, a positive allosteric modulator that increases the effectiveness of GABAergic signaling (Rudolph &

Knoflach, 2011). Bolstering the validity of our model, the spatial pattern of benzodiazepine-related E/I reductions described by the model was spatially correlated with the distribution of benzodiazepine-sensitive GABA receptors as established in positron emission tomography data (Nørgaard et al., 2021). Interestingly, one of the pharmacological targets of benzodiazepines are GABA$_A$ α_1 receptors (Rudolph & Knoflach, 2011). Increases in GABA signaling at GABA$_A$ α_1 receptors have been shown to trigger the onset of developmental critical periods in animal models (Fagiolini et al., 2004; Katagiri et al., 2007), indicating that the pFIC model is well equipped to study development-linked changes in inhibitory signaling in the human brain.

We found that the E/I ratio decreased across the cortex throughout child and adolescent development in the human brain. E/I significantly declined with age across all large-scale cortical systems; however, the magnitude of this decline varied along a unimodal-transmodal cortical hierarchy. Specifically, by age 22, the E/I ratio had declined the most in unimodal sensory territories, such as the visual and somatomotor systems, and the least in transmodal executive systems like the default and frontoparietal control systems. Differences in E/I development across these systems may be linked to differences in their maturational time courses. The development of cortical inhibitory circuitry has been well-established as a central mechanism that controls the timing and progression of critical periods of development (Hensch, 2005; Toyoizumi et al., 2013; Gogolla et al., 2014; Kalish et al., 2020). Initially, the development of inhibitory circuitry lags behind that of excitatory pyramidal cells, leading to an early increase in the local E/I ratio (Lodato et al., 2011). Later, experience and evoked activity stimulate the development of inhibitory circuitry (Hensch, 2005) — particularly fast-spiking parvalbumin positive interneurons and GABA$_A$ α_1 receptors — which begins to reduce the E/I ratio, facilitating experience-dependent plasticity and triggering the opening of the critical period window (Fagiolini et al., 2004; Hensch & Fagiolini, 2005; Katagiri et al., 2007; Toyoizumi et al., 2013). As the critical period progresses, excitatory synapses are pruned, which further reduces the E/I ratio (Han et al., 2017; Xu et al., 2020). Finally, as inhibitory circuitry begins to reach maturity, a new set of plasticity braking factors are triggered, including the formation of intracortical myelin and perineuronal nets, which stabilize cortical circuits and close the critical period window (McGee et al., 2005; McRae et al., 2007; Bavelier et al., 2010; Takesian & Hensch, 2013; Willis et al., 2022). As a result, an initial decrease in the E/I ratio can signify that a critical or sensitive period has been triggered and the cortex is in a relatively immature, plasticity-permissive state. As the E/I ratio reduces

further, it may signify that the cortex has reached a mature, plasticity-restrictive state, with pruned excitatory synapses, fully-developed inhibitory circuitry, and mature plasticity brakes that have closed the critical period (Takesian & Hensch, 2013). As such, the greater reduction in the E/I ratio we observe in sensory systems relative to association systems may reflect that sensory systems have reached a higher degree of maturity by the end of the adolescent years while association systems remain in a more immature, plasticity-permissive state. To test this hypothesis, future work could use multimodal neuroimaging that combines our pFIC approach with other markers of critical period closure -- such as intracortical myelination -- to evaluate biologically-relevent signatures of when windows of critical period plasticity open and close during youth.

Our findings align with a wealth of literature that demonstrates differences in the development of sensorimotor and association cortices. Most recently, studies have shown that other functional properties, such as functional connectivity, functional topography, structure-function coupling, and intrinsic dynamics, follow different developmental trajectories between sensory and associative cortical systems (Baum et al., 2020; Cui et al., 2020; Dong et al., 2021; Pines et al., 2022; Sydnor & Satterthwaite, 2023). For example, while the intrinsic fluctuation amplitude of sensory systems tends to linearly decline with age, functional systems within the association cortices follow curvilinear developmental trajectories which peak over adolescence before declining into adulthood (Sydnor et al., 2023). Importantly, other recent work has shown that the development of intracortical myelin, which functions as a brake on plasticity, also varies along the sensory-to-association cortical hierarchy (Grydeland et al., 2019; Paquola et al., 2019; Baum et al., 2022). Specifically, the period of peak growth in intracortical myelin was found to occur during childhood in sensorimotor cortex, yet not until adolescence in association cortex. When coupled with our current findings of a greater reduction in the E/I ratio of sensory systems (versus a weaker reduction in association systems), this work jointly indicates that sensorimotor systems are more mature by the onset of adolescence whereas association cortices may remain more plastic during the adolescent period. This interpretation aligns with a recent study showing that an fMRI marker of functional plasticity peaks during early adolescence in association cortex but continuously declines throughout childhood and adolescence in sensorimotor cortex (Sydnor et al., 2023).

The protracted development of the E/I ratio throughout adolescence may facilitate healthy cognitive development. We found that better cognitive ability was broadly associated with a lower E/I ratio across the cortex in groups of age-matched youth. Considering that the

E/I ratio normatively decreased with age, this effect may indicate that more mature cognitive performance is associated with a more mature cortical E/I ratio irrespective of age. As such, the E/I ratio may capture aspects of development independent of chronological age. Importantly, the magnitude of the effect was not uniformly distributed. We found that the greatest effect sizes were observed in association cortex, while the weakest effect sizes were observed in sensory cortex. This pattern is consistent with prior work showing that other functional properties of the association cortices are most strongly related to cognitive performance across development (Cui et al., 2020; Wang et al., 2021; Pines et al., 2022; Keller et al., 2023). Our findings also support theoretical predictions from a biophysically-based cortical circuit model of decision-making that arriving at a balanced E/I ratio supports optimal decision-making behavior (Lam et al., 2022). Together, our results suggest that although E/I ratio continues to develop throughout the cortex during adolescence, the development of the E/I ratio in association cortices is particularly relevant to maturing cognition. Critically, we replicated and generalized all associations between the E/I ratio and cognitive ability in an independent sample of youth from the GUSTO dataset, collected from a different continent, demonstrating the robustness of these effects across both populations and recruitment sites.

Our findings have important implications for understanding the emergence of psychopathology during adolescence. Though a prolonged period of developmental plasticity in the association cortices may be essential to healthy cognitive development, it may also represent a period of vulnerability to atypical developmental outcomes. A growing body of work has begun to implicate a disrupted E/I ratio in prefrontal cortex as a central mechanism in neuropsychiatric disorders such as depression and psychosis spectrum disorders (Paus et al., 2008; Anticevic & Murray, 2017; Selten et al., 2018; Sohal & Rubenstein, 2019; Vinogradov et al., 2023). These conditions are thought to involve an atypically high E/I ratio in prefrontal cortex (Jardri & Denève, 2013; Yang et al., 2016; Foss-Feig et al., 2017; Duman et al., 2019; Gao et al., 2022; Yao et al., 2022). Future studies could use our model to begin to understand how the atypical development of E/I in association cortex may lead to transdiagnostic cognitive dysfunction in developmental psychopathology.

Limitations

In this study, we applied our previous approach (Kong et al., 2021) to the feedback inhibition control (FIC) model. By parameterizing local circuit parameters as a linear combination of the known anatomical (T1w/T2w ratio) and functional (RSFC) gradients, the

resulting parametric FIC (pFIC) model generated more realistic static FC and FC dynamics. We found that using both gradients led to more realistic dynamics compared with using only one gradient or constraining local circuit parameters to be spatially uniform. However, it is not clear that the current parameterization is optimal. Future studies can explore more generic parameterizations, such as cortical geometric eigenmodes (Pang et al., 2023). Also, while our approach has the advantage of allowing for the characterization of cortex-wide patterns of E/I development, our developmental analyses were limited to linear modeling of E/I across a set of age bins. Future work in larger samples may allow for more detailed nonlinear modeling approaches that can begin to characterize potential nonlinear developmental trajectories of E/I across the cortex.

Conclusion

Our results underscore the utility of large-scale circuit models to provide insight into the underlying neurobiological mechanisms that drive neurocognitive development during youth. We provide important evidence that an essential aspect of healthy brain function—the cortical E/I ratio—is refined during childhood and adolescence. We also provide new evidence that this hallmark critical period mechanism is associated with improved cognitive ability. Our findings pave the way for future work to investigate how disrupted E/I balance may lead to cognitive dysfunction in psychopathology that emerges during youth and is characterized by atypical development of association cortices that undergo protracted maturation.

4. Methods

4.1 Human Connectome Project (HCP) dataset

We considered 1004 participants from the Human Connectome Project (HCP) S1200 release (Van Essen et al., 2013). All participants were scanned on a customized Siemens 3T Skyra using a multi-band sequence. Four resting-state fMRI (resting-fMRI) runs were collected for each participant in two sessions on two different days. Each resting-fMRI run was acquired with a repetition time (TR) of 0.72 s at 2 mm isotropic resolution and lasted for 14.4 min. The diffusion imaging consisted of 6 runs, each lasting ~9 min and 50 s. Diffusion weighting consisted of 3 shells of b = 1000, 2000, and 3000 s/mm^2 with an approximately equal number of weighting directions on each shell. Details of the data collection can be found elsewhere (Van Essen et al., 2013). The 1004 participants were randomly divided into training (N = 335), validation (N = 335) and test (N = 334) sets.

4.2 HCP preprocessing

Details of the HCP preprocessing can be found in the HCP S1200 manual. We utilized resting-fMRI data, which had already been projected to fsLR surface space, denoised with ICA-FIX and smoothed by 2 mm. For each run of each participant, the fMRI data were averaged within each Desikan–Killiany (Desikan et al., 2006) region of interest (ROI) to generate a 68 × 1200 matrix. Each 68 × 1200 matrix was used to compute 68 × 68 FC matrix by correlating the time courses among all pairs of time courses. The FC matrices were then averaged across runs of participants within the training (or validation or test) set, resulting in a group-averaged training (or validation or test) FC matrix.

Functional connectivity dynamics (FCD) were computed as follows. We defined a window with a length of 60s (equivalent to 83 time points or TRs) as recommended by previous studies (Leonardi & Van De Ville, 2015; Liégeois et al., 2017). The window was moved from the first frame to the 1118[th] frame of BOLD time series, resulting in 1118 sliding windows in total. For each run of each participant, FC was computed within each of 1118 sliding windows. Each sliding window FC matrix was then vectorized by only considering the upper triangular entries. The vectorized FCs were correlated with each other generating a 1118 × 1118 FCD matrix. Unlike static FC, we note that the FCD matrices could not be directly averaged across participants because there was no temporal correspondence between participants during the resting-state.

In the case of diffusion MRI, probabilistic tractography was run for each participant using the fiber orientation distribution (iFOD2) algorithm provided by MRtrix3 (Tournier et al., 2019). A structural connectivity (SC) matrix was generated for each participant, where each entry corresponded to the number of streamlines between two ROIs. To generate a group-level SC matrix, a thresholding procedure was employed to remove false positives. More specifically, if <50% of participants had a non-zero value in a particular entry in the SC matrix, then the entry is set to zero in all individual-level SC matrices. For each SC entry, the number of streamlines was averaged across participants with non-zero streamlines and then log-transformed (Park et al., 2021). The entries along the main diagonal were set to 0. Group-level SCs were computed by averaging individual-level SCs within the training, validation and test sets separately and normalizing the maximum value to 0.02.

4.3 Pharmacological (benzodiazepine alprazolam) dataset

The alprazolam dataset has been previously described in detail (Wolf et al., 2011). Briefly, 47 adults participated in a double-blind, placebo-controlled study using the benzodiazepine alprazolam. Each participant completed two identical experimental sessions approximately 1 week apart. In one session, participants were given a 1-mg dose of alprazolam, and in the other, they were given an identical appearing placebo. One milligram of alprazolam produces an increase in GABAergic inhibition that is considered to be clinically effective (Greenblatt et al., 1988). The order of administration was counterbalanced across participants. Alprazolam or placebo was administered 1 hour before the fMRI acquisition so that alprazolam levels and effects were near their peak at the time of data collection (Greenblatt et al., 1988). During both sessions, participants completed an emotion identification task lasting 10.5 min, while fMRI was acquired. Task-related fMRI results have been previously reported (Wolf et al., 2011). Two participants were excluded because of missing data in at least one session, yielding a final sample of 45 participants and 90 sessions in total (ages 20.9 to 56.4; mean = 39.9, standard deviation = 12.71).

All data were collected on a Siemens Trio 3T. Blood oxygen level–dependent (BOLD) fMRI data were acquired using the following parameters: TR = 3000 ms; TE = 32 ms; flip angle = 90°; FOV = 240 mm; matrix = 128 × 128; slice thickness/gap = 2/0 mm; 30 slices; effective voxel resolution = 1.875 × 1.875 × 2 mm^3; 210 volumes. The field of view (FOV) included temporal, inferior frontal, and visual cortices as well as subcortical structures, but excluded dorsal portions of the cerebral cortex.

The 45 participants were randomly divided into training, validation, and test sets with 15 participants each. The training-validation-test split was the same for both drug and placebo sessions.

4.4 Philadelphia Neurodevelopment Cohort (PNC) dataset

Neuroimaging data were obtained from a community-based sample of 1601 youth (ages 8.1 to 23.1; mean = 14.94; standard deviation = 3.69; male/ female = 764/837) that were part of the Philadelphia Neurodevelopmental Cohort (PNC). Data collection procedures and sample characteristics have been previously described in detail (Satterthwaite et al., 2014; Calkins et al., 2015). One run of resting-fMRI data was collected per participant. Following health exclusions and rigorous quality assurance, we retained 885 participants (ages 8.2 to 23.0 at first visit; mean = 15.66; standard deviation = 3.36).

All neuroimaging data were collected on the same Siemens Trio 3T scanner as was used for the alprazolam dataset. The neuroimaging procedures and acquisition parameters have been previously described in detail (Satterthwaite et al., 2014). Briefly, BOLD fMRI was acquired using similar acquisition parameters to the alprazolam dataset: TR = 3000 ms; TE = 32 ms; flip angle = 90°; FOV = 192 × 192 mm^2; matrix = 64 × 64; 46 slices; slice thickness/gap = 3/0 mm; effective voxel resolution = 3.0 × 3.0 × 3.0 mm^3; 210 volumes. The key difference is that the field of view in the PNC dataset covered the whole brain (unlike the alprazolam dataset).

4.5 Alprazolam and PNC functional image processing

Details about the alprazolam and PNC datasets have been previously provided in previous studies (Larsen et al., 2022). The alprazolam dataset consisted of two BOLD acquisitions per participant (drug and placebo session), which were preprocessed individually. BOLD runs were slice time–corrected and then motion-corrected. Susceptibility distortion was estimated and used to compute a corrected BOLD reference for more accurate co-registration with the anatomical reference. The BOLD reference was co-registered to the T1w reference using boundary-based registration (Greve & Fischl, 2009). Co-registration was configured with nine degrees of freedom to account for distortions remaining in the BOLD reference. Six head motion parameters (corresponding rotation and translation parameters) were estimated before any spatiotemporal filtering. The motion-correcting transformations, field distortion correcting warp, BOLD-to-T1w transformation, and T1w-to-template (MNI) warp were concatenated and applied to the BOLD time series in a single step

using antsApplyTransforms (ANTs) with Lanczos interpolation. Finally, the volumetric data was projected to fsaverage6 surface space (Wu et al., 2018).

Nuisance regression relied upon anatomical CompCor (aCompCor). aCompCor principal components were estimated after high-pass filtering the preprocessed BOLD time series (using a discrete cosine filter with 128-s cutoff). 5 CompCor components were extracted from the cerebrospinal fluid (CSF) and white matter (WM) masks. To remove task effects, all event conditions from the emotion identification task were modeled as 5.5-s boxcars convolved with a canonical hemodynamic response function. Each of the five emotions (fear, sad, angry, happy, and neutral) was modeled as a separate regressor. In sum, 22 regressors (6 head motion parameters and their respective temporal derivatives, top 5 aCompCor components, and 5 task regressors) were jointly regressed from the BOLD time series. The same preprocessing was performed for the PNC dataset, except that no task regressor was necessary. Overall, 17 regressors (6 head motion parameters and their respective temporal derivatives and top 5 aCompCor components) were jointly regressed from the BOLD time series.

FC and FCD of the alprazolam and PNC datasets were computed in the same manner as the HCP dataset. However, the TR was longer in the alprazolam and PNC datasets than the HCP dataset. Therefore, when computing the FCD matrices, the length of each sliding window was set to be 20 timepoints (or TRs), so that the temporal length of the window was maintained at 60s.

4.6 GUSTO dataset

To replicate our findings on the association between cognition and E/I ratio, we additionally utilized an Asian cohort, Growing Up in Singapore Towards Healthy Outcomes (GUSTO) dataset (Soh et al., 2014). We considered 389 7.5-year-old children with 1 run of resting-fMRI data and relevant cognitive scores. Participants with one or more missing cognitive scores were removed, yielding a final group of 154 participants (mean age = 7.43, std = 0.13, min age = 7.24, max age = 7.88).

All neuroimaging data were collected on a Siemens Prisma scanner. The structural data were obtained with T1w MPRAGE sequence using the following acquisition parameters: TR = 2000 ms; TE = 2.08 ms; FOV = 192×192 mm^2; matrix = 192×192; voxel resolution = $1.0 \times 1.0 \times 1.0$ mm^3. The resting-fMRI data were obtained with the following acquisition parameters: TR = 2620 ms; TE = 27 ms; flip angle = 90°; FOV = 192×192 mm^2; matrix =

64×64; 48 slices; voxel resolution $= 3.0 \times 3.0 \times 3.0$ mm³; 120 volumes. The field of view of the GUSTO dataset covered the whole brain.

4.7 GUSTO preprocessing

For each resting-fMRI run, the following sequence of preprocessing steps were performed. The first 4 frames of the run were removed. The run was then slice time-corrected and then motion-corrected. Frames with FD > 0.5mm and DVARS > 80 were marked as motion outliers. Next, the run was co-registered to the structural image with boundary-based registration (Greve & Fischl, 2009). Nuisance regression was performed with the inclusion of white matter and CSF signals, 6 head motion parameters and their respective temporal derivates as well as the top 5 aCompCor components (19 regressors in total were jointly applied). The motion outlier frames were then censored and interpolated. Finally, the run was bandpass-filtered (0.009Hz < f < 0.08Hz) and projected to Freesurfer fsaverage6 surface space.

FC and FCD of the GUSTO dataset were computed in the same manner as the HCP dataset. However, the TR was longer in the GUSTO than the HCP dataset. Therefore, when computing the FCD matrices, the length of each sliding window was set to be 23 timepoints (or TRs), so that the temporal length of the window was maintained at around 60s.

4.8 Feedback Inhibition Control (FIC) model

The derivation of the FIC model was thoroughly described in a previous study (Deco et al., 2014). Here we provide some intuition for the FIC model. The neuronal activities of the *j*-th cortical region follow the nonlinear differential equations shown below

$$I_j^{(E)} = W_E I_0 + w_{EE} J_{NMDA} S_j^{(E)} + G J_{NMDA} \sum_k C_{jk} S_k^{(E)} - w_{IE} S_j^{(I)} \qquad (1)$$

$$I_j^{(I)} = W_I I_0 + w_{EI} J_{NMDA} S_j^{(E)} - w_{II} S_j^{(I)} \qquad (2)$$

$$r_j^{(E)} = \phi\left(I_j^{(E)}\right) = \frac{a_E I_j^{(E)} - b_E}{1 - \exp\left(-d_E\left(a_E I_j^{(E)} - b_E\right)\right)} \qquad (3)$$

$$r_j^{(I)} = \phi\left(I_j^{(I)}\right) = \frac{a_I I_j^{(I)} - b_I}{1 - \exp\left(-d_I\left(a_I I_j^{(I)} - b_I\right)\right)} \qquad (4)$$

$$\frac{dS_j^{(E)}}{dt} = -\frac{S_j^{(E)}}{\tau_E} + \left(1 - S_j^{(E)}\right)\gamma r_j^{(E)} + \sigma v_j(t) \tag{5}$$

$$\frac{dS_j^{(I)}}{dt} = -\frac{S_j^{(I)}}{\tau_I} + r_j^{(I)} + \sigma v_j(t) \tag{6}$$

where S, r, and I represent synaptic gating variables, firing rate, and synaptic currents respectively. The superscripts E and I denote the excitatory and inhibitory neuronal populations respectively.

The input current $I_j^{(E)}$ of the excitatory population of the j-th cortical ROI is the sum of four inputs (Equation 1). The first input is the external input current $W_E I_0$, which might include subcortical delays. The second input is the intra-regional excitatory-to-excitatory current governed by the excitatory-to-inhibitory recurrent connection strength w_{EE} scaled by the synaptic coupling constant J_{NMDA}. The third input is the inter-regional input, which is controlled by the SC matrix (C_{jk} is the connectivity between regions j and k) and scaled by the global constant G. The fourth input is the intra-regional negative feedback from the inhibitory population governed by the inhibitory-to-excitatory connection strength w_{IE}.

The input current $I_j^{(I)}$ of the inhibitory population of the j-th cortical ROI is the sum of three inputs (Equation 2). The first input is the external input current $W_I I_0$. The second input is the intra-regional positive feedback from the excitatory population governed by the excitatory-to-inhibitory connection strength w_{EI} scaled by the synaptic coupling constant J_{NMDA}. The third input is the intra-regional inhibitory-to-inhibitory current governed by the inhibitory-to-inhibitory recurrent connection strength w_{II}.

The excitatory input current $I_j^{(E)}$ and inhibitory input current $I_j^{(I)}$ are transformed into firing rates via the input-output functions specified in Equations 3 and 4. Following previous studies (Deco et al., 2014), the parameters of the input-output function were set to be $a_E = 310$n/C, $a_I = 615$n/C, $b_E = 125$Hz, $b_I = 177$Hz, $d_E = 0.16$s and $d_I = 0.087$s.

Finally, the rate of change of the synaptic gating variables $S_j^{(E)}$ and $S_j^{(I)}$ are computed via equations 5 and 6. Following previous studies (Deco et al., 2014), the kinetic parameters for synaptic activities τ_E, τ_I and γ were set to 100ms, 10ms and 0.641 respectively. $v_j(t)$ corresponds to uncorrelated standard Gaussian noise with the noise amplitude being controlled by σ.

Following the original study (Deco et al., 2014), w_{II}, W_E, W_I, I_0 and J_{NMDA} were set to 1, 1, 0.7 0.382nA and 0.15nA respectively in the current study. The inhibitory-to-excitatory connection strength w_{IE} was computed analytically to ensure that the excitatory firing rate is maintained to be around 3Hz (Demirtaş et al., 2019). We note that this analytical computation assumes a noiseless system, so in practice, we imposed a constraint that the firing rate is between 2.7Hz to 3.3Hz. During the estimation of the pFIC model (next section), parameters were rejected if firing rates fall outside this range.

The excitatory-to-excitatory recurrent connection strength w_{EE}, excitatory-to-inhibitory connection strength w_{EI}, noise amplitude σ and global SC scaling constant G are unknown parameters, which will be estimated by fitting to empirical fMRI data (next section).

Given a fixed set of model parameters, equations 1 to 6 can be used to simulate the time courses of excitatory and inhibitory synaptic gating variables ($S_j^{(E)}$ and $S_j^{(I)}$) of each ROI. The regional E/I ratio was defined as the ratio between the temporal average of $S_j^{(E)}$ and $S_j^{(I)}$. The mean cortical E/I ratio was the average of regional E/I ratios across all cortical ROIs. The simulated excitatory synaptic gating variables ($S_j^{(E)}$) were also fed to the Balloon-Windkessel hemodynamic model to simulate fMRI BOLD signals (Stephan et al., 2007; Deco et al., 2014; Heinzle et al., 2016). The simulated fMRI BOLD signals were then used to generate simulated static FC and FCD.

4.9 Parametric FIC (pFIC) model

Recall that the FIC model was instantiated using the Desikan-Killiany parcellation with 68 regions of interest. Given that we wanted the excitatory-to-excitatory recurrent connection strength w_EE, excitatory-to-inhibitory connection strength w_EI and noise amplitude σ to be spatially heterogeneous (with G being a global constant), if we optimized each parameter independently, there would be a total of $68 \times 3 + 1 = 205$ parameters, which is computationally challenging.

In our previous study (Kong et al., 2021), we reduced the number of "free" parameters by parameterizing the local synaptic parameters with a linear combination of the first principal FC gradient and T1w/T2w ratio map. We note that in the previous study (Kong et al., 2021), we considered a highly simplified parametric mean field model that does not differentiate between excitatory and inhibitory neural populations. Here, we considered the

same approach for the FIC model by parameterizing w_{EE}, w_{EI}, and σ as a linear combination of T1w/T2w myelin map and first principal FC gradient.

$$w_{EE,j} = a + b \times \text{myelin}_j + c \times \text{FC gradient}_j \qquad (7)$$

$$w_{EI,j} = d + e \times \text{myelin}_j + f \times \text{FC gradient}_j \qquad (8)$$

$$\sigma_j = g + h \times \text{myelin}_j + i \times \text{FC gradient}_j \qquad (9)$$

where j denotes the ROI index. By adopting this parameterization approach, the number of "free" numbers was reduced to $3 \times 3 + 1 = 10$ parameters.

4.10 Optimization of the parametric FIC (pFIC) model in the HCP dataset

The 10 unknown parameters of the pFIC model were optimized using a previously published approach (Kong et al., 2021) by maximizing fit to empirical static FC and FCD. The agreement between empirical and simulated FC matrices was defined as the Pearson's correlation (r) between the z-transformed upper triangular entries of the two matrices. Larger r indicates more similar static FC. However, Pearson's correlation does not account for scale difference, so we also computed the absolute difference (d) between the means of the empirical and simulated FC matrices (Demirtaş et al., 2019). A smaller d indicates more similar static FC.

The dissimilarity between simulated and empirical FCD matrices was quantified by using the Kolmogorov-Smirnov (KS) distance. Here, the KS distance was defined as the maximum distance between the cumulative distribution functions (CDFs) constructed by collapsing the upper triangular entries of simulated and empirical FCD matrices (E. C. A. Hansen et al., 2015; Kong et al., 2021). Hence, a small KS distance indicated 2 similar CDFs, therefore 2 similar FCD matrices.

The overall cost was defined as $(1 - r) + d + KS$. A smaller cost indicates better agreement between simulated and empirical fMRI. Recall that the 1004 HCP participants were randomly divided into training (N = 335), validation (N = 335) and test (N = 334) sets. Following our previous study (Kong et al., 2021), we used the covariance matrix adaptation evolution strategy (CMA-ES) (Hansen, 2006) to minimize the overall cost function during training. Due to the lack of temporal correspondence between FCD matrices across runs and participants during rs-fMRI scans, directly averaging FCD matrices will cancel out the temporal dynamics. Representing FCD matrices using CDFs avoided such problem because averaging the CDFs across runs and participants could still largely preserve the distribution

of FCD matrix entries. We thus averaged the FCD CDFs across participants and runs separately within the training, validation and test sets.

In the HCP training set, the CMA-ES algorithm was iterated 100 times and repeated 5 times with different random initializations, yielding a total of 500 candidate parameter sets (**Figure 2A**). The 500 candidate parameter sets were evaluated in the validation set to obtain the top 10 candidate parameter sets. To ensure diversity among the parameter sets, the procedure to select the top 10 parameter sets was as follows. First, the parameter set with the lowest validation cost was selected. Then, the parameter set with the lowest validation cost and whose parameter (w_{EE}, w_{EI}, σ) maps exhibited less than 0.98 correlation with the current selected parameter set(s) was selected. This procedure was repeated until 10 parameter sets were selected.

The top 10 candidate parameter sets from CMA-ES were then applied to the HCP test set SC. For each parameter set, 1000 simulations were performed, yielding 1000 simulated static FC and FCD matrices. Pearson correlation and the absolute difference were then computed between each simulated FC and the empirical FC from the HCP test set and averaged. Similarly, KS statistics was computed between each simulated FCD CDF and the empirical FCD CDF from the HCP test set and averaged.

To speed up the computation, a step size of 6ms was used to integrate the ODEs in the training set. However, to ensure more accurate integration, a step size of 0.5ms was used for both validation and test sets. To ensure that this time step size was small enough, we repeated the experiment using a step size of 6ms for the training set and a step size of 0.1ms for the validation and test sets. The overall cost was highly similar across step size of 0.5ms and step size of 0.1ms. In particular, for both 0.5ms and 0.1ms step sizes, an overall cost of 0.58 ± 0.018 was achieved in the HCP test set across the top 10 parameter sets from the validation set.

Finally, we observed that the correlations between w_{EI} and T1w/T2w ratio were consistently negative, while the correlations between w_{EI} and RSFC gradient were consistently positive across the top 10 parameter sets. Since the training sets for the alprazolam and PNC datasets were substantially smaller than the HCP training set (~15 participants versus 335 participants), when optimizing the pFIC model in the alprazolam and PNC datasets, we additionally imposed the constraints that w_{EI} and T1w/T2w ratio should be negative, while the correlations between w_{EI} and FC gradient should be positive.

4.11   Pharmacological E/I ratio analysis

Recall that the 45 participants of the alprazolam dataset were randomly divided into training (N = 15), validation (N = 15) and test (N = 15) sets. Each participant had 2 runs of fMRI data - one run for the drug session and one run for the placebo session. The SC matrices used for training, validation, and testing were the same as HCP training, validation, and tests set respectively. Both T1w/T2w ratio and the first principal FC gradient maps were generated from HCP training set.

Due to the limited FOV of the alprazolam dataset, we only considered regions with more than 50% ROI coverage, resulting in 42 ROIs (in the case of the Desikan-Killiany parcellation). The remaining 26 ROIs were masked out for the group-level SC, static FC, FCD, T1w/T2w ratio and FC gradient. For each experimental condition (placebo or alprazolam), 250 candidate parameter sets were generated from the condition's training set. The top 10 parameter sets from the validation set were evaluated in the test set.

Although the parameters were only estimated based on the 42 ROIs, the estimated linear coefficients could be used to generate whole cortex estimates of w_{EE}, w_{EI} and σ (based on equations 7 to 9) given that the original T1w/T2w ratio and FC gradient (from the HCP training set) covered the whole cortex. The simulated time courses S_E and S_I were then generated using the 68-ROI SC and extrapolated model parameters. For a given set of parameters, 1000 simulations were performed to generate 1000 sets of E/I ratio. The final E/I ratio was the average across 1000 sets of E/I ratio.

E/I ratio contrast was defined as the difference between E/I ratio of the placebo sessions and E/I ratio of the drug sessions. To test that the E/I ratio contrast was significantly greater than 0, we performed permutation test to generate a null distribution of E/I ratio contrasts. More specifically, after dividing the participants into training, validation and test sets, the 'drug' and 'placebo' sessions were randomly permuted within each participant. The entire procedure (above) was repeated, generating a null value for the E/I ratio contrast. The permutation procedure was repeated 100 times, yielding a null distribution of regional E/I ratio contrasts. A 2-tail p-value was computed based on this null distribution.

The regional E/I ratio contrast was also correlated with benzodiazepine receptor (BZR) density. The statistical significance of this correspondence was computed using a spin test that accounts for spatial autocorrelation (Alexander-Bloch et al., 2018).

4.12   Association between age and E/I ratio in the PNC dataset

885 participants of the PNC dataset (Satterthwaite et al., 2014) were first sorted according to age in ascending order and divided into 29 groups of 30 or 31 participants. For each age group, 15 participants were randomly selected as the validation set, while the remaining participants were assigned to the training set.

For this analysis, the SC matrices used for training and validation were the same as HCP training and validation sets respectively. Both T1w/T2w ratio and the first principal FC gradient maps were generated from HCP training set. For each age group, 250 candidate model parameter sets were generated from the group's training set using CMA-ES and evaluated in the group's validation set. For each age group, the parameter set with the lowest validation cost was used to estimate regional E/I ratio across the cortex.

For each cortical ROI, we fitted a linear regression model between regional E/I ratio and the mean age of each age group. The slope of the linear regression model (for each brain region) was visualized on the cortical surface. All slopes were negative and all p-values survived FDR correction (q < 0.05). For robustness, the split of the participants into training and validation sets were repeated 5 times and the most representative split was shown in **Figure 4**.

4.13   Association between cognition and E/I ratio in the PNC dataset

Each PNC participant had completed a set of 12 tasks from 4 cognitive domains, including executive control, episodic memory, complex cognition, and social cognition. Three types of scores, including an accuracy score, a speed score, and an efficiency score, were obtained for each of the 12 tasks. Factor analyses of each type of scores were performed within each cognitive domain and across all cognitive domains to generate three domain-specific accuracy factor scores and one domain-general (overall) accuracy score (Moore et al., 2015).

To control for age, 885 participants were sorted according to age in an ascending order. For each 12-month interval, participants whose age were within this interval were extracted to form one age group. For each age group, participants with domain-general (overall) accuracy scores above the median were assigned to a high-performance group, the rest are assigned to a low-performance group. In sum, 441 participants were assigned to the high-performance group (mean age = 15.68), 444 participants were assigned to the low-performance group (mean age = 15.64). Both high and low-performance groups were then

divided into 14 subgroups of 31 (or 32) participants, yielding 14 pairs of age-matched high-performance and low-performance groups (see **Figure 6A and 6B**).

Participants of each subgroup were further randomly divided into a training set (N = 16) and a validation set (N = 15 or 16). Similar to the previous analyses, SCs used for training and validation were from the HCP training and validation sets respectively. T1w/T2w ratio and FC gradient maps were from the HCP training set. The E/I ratios of the 14 pairs of high-performance and low-performance groups were estimated separately and compared using a 2-tail 1-sample t-test. To test for domain specificity, we also repeated the analyses for the 3 domain-specific accuracy scores. FDR correction with q < 0.05 was used to correct for multiple comparisons.

4.14   Association between cognition and E/I ratio in the GUSTO dataset

Our analyses were based on the fMRI and behavioral data of 154 participants from the GUSTO dataset. The fMRI data were acquired when the participants were around 7.5 years old. We selected 5 behavioral scores which assessed participants' cognitive performances. All behavior tests were performed within 1.5 years of fMRI acquisition (i.e., age 6 to 8.5). The 5 test scores were as follows: (1) Cambridge Neuropsychological Test Automated Battery (CANTAB) Spatial Working Memory (SWM) test (completed at age 6): sum of total errors for 4 and 6 boxes trails. (2) Delayed Matching to Sample (DMS) test (completed at age 6): percentage of the total number of trials upon which a correct selection was made on the participant's first response. (3) Behavior Rating Inventory of Executive Function (BRIEF; completed at age 7): Cognition Regulation Index T-score. (4) Wechsler Abbreviated Scale of Intelligence (WASI) test (completed at age 7): sum of Block Design and Matrix Reasoning T-scores. (5) CANTAB SWM test (completed at age 8.5): sum of total errors for 4 and 8 boxes trails.

Principal component analysis (PCA) was performed on these 5 scores across all participants to derive the first principal component (PC1) score. Higher PC1 score indicated better cognitive performance across the 5 behavioral scores (on average). Participants were sorted according to their PC1 scores in an ascending order. The first 77 participants were assigned to the low-performance group, while the rest of 77 participants were assigned to the high-performance group. Ages were well-matched between high and low-performance groups (as shown in results).

Participants of high and low-performance groups were further randomly divided into a training set (N = 39) and a validation set (N = 38). Similar to the previous analyses, SCs

used for training and validation were from the HCP training and validation sets respectively. T1w/T2w ratio and FC gradient maps were from the HCP training set. For robustness, the analyses were repeated 5 times with different random training-validation splits of participants within high-performance group and each low-performance group. Results from the most representative split were shown in the results.

To compute the statistical significance of E/I ratio differences between low and high-performance groups, PC1 scores were randomly permuted across participants. The participants were again assigned to high or low-performance groups according to their permuted PC1 scores. Then the E/I ratio difference was re-estimated. This permutation process was repeated 100 times to construct a null distribution of E/I ratio difference. A 2-tail p-value was computed based on this null distribution.

4.15   Ethics

The HCP data collection was approved by a consortium of institutions institutional review boards (IRBs) in the United States and Europe, led by Washington University (St Louis) and the University of Minnesota (WU-Minn HCP Consortium). Data collection and study procedures for the Alprazolam dataset were approved by the University of Pennsylvania Institutional Review Board; data for the PNC were approved by Institutional Review Boards from both Penn and the Children's Hospital of Philadelphia. The GUSTO data collection was approved by the National Healthcare Group Domain Specific Review Board and the SingHealth Centralised Institutional Review Board. All participants provided informed consent before data collection. The current study was approved by the IRB of the National University of Singapore.

4.16   Data availability

The HCP data is publicly available (https://www.humanconnectome.org/). The GUSTO dataset can be obtained via a data transfer agreement (https://www.gusto.sg/). The developmental dataset is publicly available in the Database of Genotypes and Phenotypes (dbGaP accession phs000607.v3.p2). All pharmacological imaging data necessary to evaluate the conclusions in the paper are available here (GITHUB_LINK).

4.17 Code availability

Code for this study can be found here (GITHUB_LINK). Co-authors (TZ and LA) reviewed the code before merging into the GitHub repository to reduce the chance of coding errors.

Acknowledgements

This work was supported by the Singapore National Research Foundation (NRF) Fellowship (Class of 2017), the National University of Singapore Yong Loo Lin School of Medicine (NUHSRO/2020/124/TMR/LOA), the Singapore National Medical Research Council (NMRC) LCG (OFLCG19May-0035), NMRC STaR (STaR20nov-0003), and the United States National Institutes of Health (R01MH120080). Any opinions, findings and conclusions or recommendations expressed in this material are those of the authors and do not reflect the views of the Singapore NRF or the Singapore NMRC. In RIE2025, the GUSTO dataset was supported by funding from the NRF's Human Health and Potential (HHP) Domain, under the Human Potential Programme. The alprazolam sample was funded by AstraZeneca Pharmaceuticals LP. DHW was also supported by NARSAD and the Sidney R. Baer, Jr. Foundation. BL was funded by NIH K99MH127293 and T32MH014654. VJS was supported by a National Science Foundation Graduate Research Fellowship (DGE-1845298). The PNC was funeded via RC2 grants from the National Institute of Mental Health: MH089983 and MH089924. Additional support was provided by R01MH113550, R01MH120482, R01MH112847, R01EB022573, RF1MH116920, RF1MH121867, R37MH125829, T32 MH019112, the AE Foundation, and the Penn-CHOP Lifespan Brain Institute. Our computational work was partially performed on resources of the National Supercomputing Centre, Singapore (https://www.nscc.sg). Data were in part provided by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University.

References

Akil, M., & Lewis, D. A. (1992). Differential distribution of parvalbumin-immunoreactive pericellular clusters of terminal boutons in developing and adult monkey neocortex. *Experimental Neurology*, *115*(2), 239–249. https://doi.org/10.1016/0014-4886(92)90058-X

Alexander-Bloch, A. F., Shou, H., Liu, S., Satterthwaite, T. D., Glahn, D. C., Shinohara, R. T., Vandekar, S. N., & Raznahan, A. (2018). On testing for spatial correspondence between maps of human brain structure and function. *NeuroImage*, *178*, 540–551. https://doi.org/10.1016/j.neuroimage.2018.05.070

Anderson, K. M., Collins, M. A., Chin, R., Ge, T., Rosenberg, M. D., & Holmes, A. J. (2020a). Transcriptional and imaging-genetic association of cortical interneurons, brain function, and schizophrenia risk. *Nature Communications*, *11*(1), 2889. https://doi.org/10.1038/s41467-020-16710-x

Anderson, K. M., Collins, M. A., Kong, R., Fang, K., Li, J., He, T., Chekroud, A. M., Yeo, B. T. T., & Holmes, A. J. (2020b). Convergent molecular, cellular, and cortical neuroimaging signatures of major depressive disorder. *Proceedings of the National Academy of Sciences*, *117*(40), 25138–25149. https://doi.org/10.1073/pnas.2008004117

Anticevic, A., & Lisman, J. (2017). How Can Global Alteration of Excitation/Inhibition Balance Lead to the Local Dysfunctions That Underlie Schizophrenia? *Biological Psychiatry*, *81*(10), 818–820. https://doi.org/10.1016/j.biopsych.2016.12.006

Anticevic, A., & Murray, J. D. (2017). Rebalancing Altered Computations: Considering the Role of Neural Excitation and Inhibition Balance Across the Psychiatric Spectrum. *Biological Psychiatry*, *81*(10), 816–817. https://doi.org/10.1016/j.biopsych.2017.03.019

Atallah, B. V., & Scanziani, M. (2009). Instantaneous Modulation of Gamma Oscillation Frequency by Balancing Excitation with Inhibition. *Neuron*, *62*(4), 566–577. https://doi.org/10.1016/j.neuron.2009.04.027

Baum, G. L., Cui, Z., Roalf, D. R., Ciric, R., Betzel, R. F., Larsen, B., Cieslak, M., Cook, P. A., Xia, C. H., Moore, T. M., Ruparel, K., Oathes, D. J., Alexander-Bloch, A. F., Shinohara, R. T., Raznahan, A., Gur, R. E., Gur, R. C., Bassett, D. S., & Satterthwaite, T. D. (2020). Development of structure–function coupling in human brain networks during youth. *Proceedings of the National Academy of Sciences*, *117*(1), 771–778. https://doi.org/10.1073/pnas.1912034117

Baum, G. L., Flournoy, J. C., Glasser, M. F., Harms, M. P., Mair, P., Sanders, A. F. P., Barch, D. M., Buckner, R. L., Bookheimer, S., Dapretto, M., Smith, S., Thomas, K. M., Yacoub, E., Van Essen, D. C., & Somerville, L. H. (2022). Graded Variation in T1w/T2w Ratio during Adolescence: Measurement, Caveats, and Implications for Development of Cortical Myelin. *The Journal of Neuroscience*, *42*(29), 5681–5694. https://doi.org/10.1523/JNEUROSCI.2380-21.2022

Bavelier, D., Levi, D. M., Li, R. W., Dan, Y., & Hensch, T. K. (2010). Removing Brakes on Adult Brain Plasticity: From Molecular to Behavioral Interventions. *The Journal of Neuroscience*, *30*(45), 14964–14971. https://doi.org/10.1523/JNEUROSCI.4812-10.2010

Bethlehem, R. A. I., Seidlitz, J., White, S. R., Vogel, J. W., Anderson, K. M., Adamson, C., Adler, S., Alexopoulos, G. S., Anagnostou, E., Areces-Gonzalez, A., Astle, D. E., Auyeung, B., Ayub, M., Bae, J., Ball, G., Baron-Cohen, S., Beare, R., Bedford, S. A., Benegal, V., … Alexander-Bloch, A. F. (2022). Brain charts for the human lifespan. *Nature*, *604*(7906), 525–533. https://doi.org/10.1038/s41586-022-04554-y

Burt, J. B., Demirtaş, M., Eckner, W. J., Navejar, N. M., Ji, J. L., Martin, W. J., Bernacchia, A., Anticevic, A., & Murray, J. D. (2018). Hierarchy of transcriptomic specialization across human cortex captured by structural neuroimaging topography. *Nature Neuroscience*, *21*(9), 1251–1259. https://doi.org/10.1038/s41593-018-0195-0

Calkins, M. E., Merikangas, K. R., Moore, T. M., Burstein, M., Behr, M. A., Satterthwaite, T. D., Ruparel, K., Wolf, D. H., Roalf, D. R., Mentch, F. D., Qiu, H., Chiavacci, R., Connolly, J. J., Sleiman, P. M. A., Gur, R. C., Hakonarson, H., & Gur, R. E. (2015). The Philadelphia Neurodevelopmental Cohort: Constructing a deep phenotyping collaborative. *Journal of Child Psychology and Psychiatry*, *56*(12), 1356–1369. https://doi.org/10.1111/jcpp.12416

Canetta, S. E., Holt, E. S., Benoit, L. J., Teboul, E., Sahyoun, G. M., Ogden, R. T., Harris, A. Z., & Kellendonk, C. (2022). Mature parvalbumin interneuron function in prefrontal cortex requires activity during a postnatal sensitive period. *ELife*, *11*, e80324. https://doi.org/10.7554/eLife.80324

Cardin, J. A., Carlén, M., Meletis, K., Knoblich, U., Zhang, F., Deisseroth, K., Tsai, L.-H., & Moore, C. I. (2009). Driving fast-spiking cells induces gamma rhythm and controls sensory responses. *Nature*, *459*(7247), 663–667. https://doi.org/10.1038/nature08002

Casey, B. J., Jones, R. M., & Hare, T. A. (2008). The Adolescent Brain. *Annals of the New York Academy of Sciences*, *1124*(1), 111–126. https://doi.org/10.1196/annals.1440.010

Condé, F., Lund, J. S., Jacobowitz, D. M., Baimbridge, K. G., & Lewis, D. A. (1994). Local circuit neurons immunoreactive for calretinin, calbindin D-28k or parvalbumin in monkey prefronatal cortex: Distribution and morphology: CALCIUM-BINDING PROTEINS IN MONKEY PREFRONTAL CORTEX. *Journal of Comparative Neurology*, *341*(1), 95–116. https://doi.org/10.1002/cne.903410109

Cui, Z., Li, H., Xia, C. H., Larsen, B., Adebimpe, A., Baum, G. L., Cieslak, M., Gur, R. E., Gur, R. C., Moore, T. M., Oathes, D. J., Alexander-Bloch, A. F., Raznahan, A., Roalf, D. R., Shinohara, R. T., Wolf, D. H., Davatzikos, C., Bassett, D. S., Fair, D. A., … Satterthwaite, T. D. (2020). Individual Variation in Functional Topography of Association Networks in Youth. *Neuron*, *106*(2), 340-353.e8. https://doi.org/10.1016/j.neuron.2020.01.029

Deco, G., Kringelbach, M. L., Arnatkeviciute, A., Oldham, S., Sabaroedin, K., Rogasch, N. C., Aquino, K. M., & Fornito, A. (2021). Dynamical consequences of regional heterogeneity in the brain's transcriptional landscape. *Science Advances*, *7*(29), eabf4752. https://doi.org/10.1126/sciadv.abf4752

Deco, G., Ponce-Alvarez, A., Hagmann, P., Romani, G. L., Mantini, D., & Corbetta, M. (2014). How Local Excitation-Inhibition Ratio Impacts the Whole Brain Dynamics. *Journal of Neuroscience*, *34*(23), 7886–7898. https://doi.org/10.1523/JNEUROSCI.5068-13.2014

Deco, G., Ponce-Alvarez, A., Mantini, D., Romani, G. L., Hagmann, P., & Corbetta, M. (2013). Resting-State Functional Connectivity Emerges from Structurally and Dynamically Shaped Slow Linear Fluctuations. *Journal of Neuroscience*, *33*(27), 11239–11252. https://doi.org/10.1523/JNEUROSCI.1091-13.2013

Demirtaş, M., Burt, J. B., Helmer, M., Ji, J. L., Adkinson, B. D., Glasser, M. F., Van Essen, D. C., Sotiropoulos, S. N., Anticevic, A., & Murray, J. D. (2019). Hierarchical Heterogeneity across Human Cortex Shapes Large-Scale Neural Dynamics. *Neuron*, *101*(6), 1181-1194.e13. https://doi.org/10.1016/j.neuron.2019.01.017

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner, R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J. (2006). An automated labeling system for subdividing the human cerebral cortex on

MRI scans into gyral based regions of interest. *NeuroImage*, *31*(3), 968–980. https://doi.org/10.1016/j.neuroimage.2006.01.021

Dienel, S. J., Fish, K. N., & Lewis, D. A. (2023). The Nature of Prefrontal Cortical GABA Neuron Alterations in Schizophrenia: Markedly Lower Somatostatin and Parvalbumin Gene Expression Without Missing Neurons. *American Journal of Psychiatry*, appi.ajp.20220676. https://doi.org/10.1176/appi.ajp.20220676

Dong, H.-M., Margulies, D. S., Zuo, X.-N., & Holmes, A. J. (2021). Shifting gradients of macroscale cortical organization mark the transition from childhood to adolescence. *Proceedings of the National Academy of Sciences*, *118*(28), e2024448118. https://doi.org/10.1073/pnas.2024448118

Dorrn, A. L., Yuan, K., Barker, A. J., Schreiner, C. E., & Froemke, R. C. (2010). Developmental sensory experience balances cortical excitation and inhibition. *Nature*, *465*(7300), 932–936. https://doi.org/10.1038/nature09119

Duman, R. S., Sanacora, G., & Krystal, J. H. (2019). Altered Connectivity in Depression: GABA and Glutamate Neurotransmitter Deficits and Reversal by Novel Treatments. *Neuron*, *102*(1), 75–90. https://doi.org/10.1016/j.neuron.2019.03.013

Fagiolini, M., Fritschy, J.-M., Löw, K., Möhler, H., Rudolph, U., & Hensch, T. K. (2004). Specific GABA$_A$ Circuits for Visual Cortical Plasticity. *Science*, *303*(5664), 1681–1683. https://doi.org/10.1126/science.1091032

Fossati, P., Amar, G., Raoux, N., Ergis, A. M., & Allilaire, J. F. (1999). Executive functioning and verbal memory in young patients with unipolar depression and schizophrenia. *Psychiatry Research*, *89*(3), 171–187. https://doi.org/10.1016/S0165-1781(99)00110-9

Foss-Feig, J. H., Adkinson, B. D., Ji, J. L., Yang, G., Srihari, V. H., McPartland, J. C., Krystal, J. H., Murray, J. D., & Anticevic, A. (2017). Searching for Cross-Diagnostic Convergence: Neural Mechanisms Governing Excitation and Inhibition Balance in Schizophrenia and Autism Spectrum Disorders. *Biological Psychiatry*, *81*(10), 848–861. https://doi.org/10.1016/j.biopsych.2017.03.005

Froudist-Walsh, S., Bliss, D. P., Ding, X., Rapan, L., Niu, M., Knoblauch, K., Zilles, K., Kennedy, H., Palomero-Gallagher, N., & Wang, X.-J. (2021). A dopamine gradient controls access to distributed working memory in the large-scale monkey cortex. *Neuron*, *109*(21), 3500-3520.e13. https://doi.org/10.1016/j.neuron.2021.08.024

Gandal, M. J., Zhang, P., Hadjimichael, E., Walker, R. L., Chen, C., Liu, S., Won, H., Van Bakel, H., Varghese, M., Wang, Y., Shieh, A. W., Haney, J., Parhami, S., Belmont, J., Kim, M., Moran Losada, P., Khan, Z., Mleczko, J., Xia, Y., … Abyzov, A. (2018). Transcriptome-wide isoform-level dysregulation in ASD, schizophrenia, and bipolar disorder. *Science*, *362*(6420), eaat8127. https://doi.org/10.1126/science.aat8127

Gao, W.-J., Yang, S.-S., Mack, N. R., & Chamberlin, L. A. (2022). Aberrant maturation and connectivity of prefrontal cortex in schizophrenia—Contribution of NMDA receptor development and hypofunction. *Molecular Psychiatry*, *27*(1), 731–743. https://doi.org/10.1038/s41380-021-01196-w

Gennatas, E. D., Avants, B. B., Wolf, D. H., Satterthwaite, T. D., Ruparel, K., Ciric, R., Hakonarson, H., Gur, R. E., & Gur, R. C. (2017). Age-Related Effects and Sex Differences in Gray Matter Density, Volume, Mass, and Cortical Thickness from Childhood to Young Adulthood. *The Journal of Neuroscience*, *37*(20), 5065–5073. https://doi.org/10.1523/JNEUROSCI.3550-16.2017

Ghisleni, C., Bollmann, S., Poil, S.-S., Brandeis, D., Martin, E., Michels, L., O'Gorman, R. L., & Klaver, P. (2015). Subcortical Glutamate Mediates the Reduction of Short-Range Functional Connectivity with Age in a Developmental Cohort. *Journal of*

Neuroscience, *35*(22), 8433–8441. https://doi.org/10.1523/JNEUROSCI.4375-14.2015

Ghosh, A., Rho, Y., McIntosh, A. R., Kötter, R., & Jirsa, V. K. (2008). Noise during Rest Enables the Exploration of the Brain's Dynamic Repertoire. *PLoS Computational Biology*, *4*(10), e1000196. https://doi.org/10.1371/journal.pcbi.1000196

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J. R., Van Essen, D. C., & Jenkinson, M. (2013). The minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage*, *80*, 105–124. https://doi.org/10.1016/j.neuroimage.2013.04.127

Glasser, M. F., & Van Essen, D. C. (2011). Mapping Human Cortical Areas *In Vivo* Based on Myelin Content as Revealed by T1- and T2-Weighted MRI. *The Journal of Neuroscience*, *31*(32), 11597–11616. https://doi.org/10.1523/JNEUROSCI.2180-11.2011

Gogolla, N., Takesian, A. E., Feng, G., Fagiolini, M., & Hensch, T. K. (2014). Sensory Integration in Mouse Insular Cortex Reflects GABA Circuit Maturation. *Neuron*, *83*(4), 894–905. https://doi.org/10.1016/j.neuron.2014.06.033

Gogtay, N., & Thompson, P. M. (2010). Mapping gray matter development: Implications for typical development and vulnerability to psychopathology. *Brain and Cognition*, *72*(1), 6–15. https://doi.org/10.1016/j.bandc.2009.08.009

Goulas, A., Changeux, J.-P., Wagstyl, K., Amunts, K., Palomero-Gallagher, N., & Hilgetag, C. C. (2021). The natural axis of transmitter receptor distribution in the human cerebral cortex. *Proceedings of the National Academy of Sciences*, *118*(3), e2020574118. https://doi.org/10.1073/pnas.2020574118

Greenblatt, D. J., Harmatz, J. S., Dorsey, C., & Shader, R. I. (1988). Comparative single-dose kinetics and dynamics of lorazepam, alprazolam, prazepam, and placebo. *Clinical Pharmacology and Therapeutics*, *44*(3), 326–334. https://doi.org/10.1038/clpt.1988.158

Greve, D. N., & Fischl, B. (2009). Accurate and robust brain image alignment using boundary-based registration. *NeuroImage*, *48*(1), 63–72. https://doi.org/10.1016/j.neuroimage.2009.06.060

Grydeland, H., Vértes, P. E., Váša, F., Romero-Garcia, R., Whitaker, K., Alexander-Bloch, A. F., Bjørnerud, A., Patel, A. X., Sederevičius, D., Tamnes, C. K., Westlye, L. T., White, S. R., Walhovd, K. B., Fjell, A. M., & Bullmore, E. T. (2019). Waves of Maturation and Senescence in Micro-structural MRI Markers of Human Cortical Myelination over the Lifespan. *Cerebral Cortex*, *29*(3), 1369–1381. https://doi.org/10.1093/cercor/bhy330

Gu, Y., Tran, T., Murase, S., Borrell, A., Kirkwood, A., & Quinlan, E. M. (2016). Neuregulin-Dependent Regulation of Fast-Spiking Interneuron Excitability Controls the Timing of the Critical Period. *The Journal of Neuroscience*, *36*(40), 10285–10295. https://doi.org/10.1523/JNEUROSCI.4242-15.2016

Han, K.-S., Cooke, S. F., & Xu, W. (2017). Experience-Dependent Equilibration of AMPAR-Mediated Synaptic Transmission during the Critical Period. *Cell Reports*, *18*(4), 892–904. https://doi.org/10.1016/j.celrep.2016.12.084

Hansen, E. C. A., Battaglia, D., Spiegler, A., Deco, G., & Jirsa, V. K. (2015). Functional connectivity dynamics: Modeling the switching behavior of the resting state. *NeuroImage*, *105*, 525–535. https://doi.org/10.1016/j.neuroimage.2014.11.001

Hansen, N. (2006). The CMA Evolution Strategy: A Comparing Review. In J. A. Lozano, P. Larrañaga, I. Inza, & E. Bengoetxea (Eds.), *Towards a New Evolutionary Computation* (Vol. 192, pp. 75–102). Springer Berlin Heidelberg. https://doi.org/10.1007/3-540-32494-1_4

Heinzle, J., Koopmans, P. J., Den Ouden, H. E. M., Raman, S., & Stephan, K. E. (2016). A hemodynamic model for layered BOLD signals. *NeuroImage*, *125*, 556–570. https://doi.org/10.1016/j.neuroimage.2015.10.025

Hensch, T. K. (2005). Critical period plasticity in local cortical circuits. *Nature Reviews Neuroscience*, *6*(11), 877–888. https://doi.org/10.1038/nrn1787

Hensch, T. K., & Fagiolini, M. (2005). Excitatory–inhibitory balance and critical period plasticity in developing visual cortex. In *Progress in Brain Research* (Vol. 147, pp. 115–124). Elsevier. https://doi.org/10.1016/S0079-6123(04)47009-5

Honey, C. J., Kötter, R., Breakspear, M., & Sporns, O. (2007). Network structure of cerebral cortex shapes functional connectivity on multiple time scales. *Proceedings of the National Academy of Sciences*, *104*(24), 10240–10245. https://doi.org/10.1073/pnas.0701519104

Huntenburg, J. M., Bazin, P.-L., & Margulies, D. S. (2018). Large-Scale Gradients in Human Cortical Organization. *Trends in Cognitive Sciences*, *22*(1), 21–31. https://doi.org/10.1016/j.tics.2017.11.002

Insel, T. R. (2014). Mental Disorders in Childhood: Shifting the Focus From Behavioral Symptoms to Neurodevelopmental Trajectories. *JAMA*, *311*(17), 1727. https://doi.org/10.1001/jama.2014.1193

Iwai, Y., Fagiolini, M., Obata, K., & Hensch, T. K. (2003). Rapid Critical Period Induction by Tonic Inhibition in Visual Cortex. *The Journal of Neuroscience*, *23*(17), 6695–6702. https://doi.org/10.1523/JNEUROSCI.23-17-06695.2003

Jardri, R., & Denève, S. (2013). Circular inferences in schizophrenia. *Brain*, *136*(11), 3227–3241. https://doi.org/10.1093/brain/awt257

Kalish, B. T., Barkat, T. R., Diel, E. E., Zhang, E. J., Greenberg, M. E., & Hensch, T. K. (2020). Single-nucleus RNA sequencing of mouse auditory cortex reveals critical period triggers and brakes. *Proceedings of the National Academy of Sciences*, *117*(21), 11744–11752. https://doi.org/10.1073/pnas.1920433117

Katagiri, H., Fagiolini, M., & Hensch, T. K. (2007). Optimization of Somatic Inhibition at Critical Period Onset in Mouse Visual Cortex. *Neuron*, *53*(6), 805–812. https://doi.org/10.1016/j.neuron.2007.02.026

Keller, A. S., Sydnor, V. J., Pines, A., Fair, D. A., Bassett, D. S., & Satterthwaite, T. D. (2023). Hierarchical functional system development supports executive function. *Trends in Cognitive Sciences*, *27*(2), 160–174. https://doi.org/10.1016/j.tics.2022.11.005

Kim, Y., Yang, G. R., Pradhan, K., Venkataraju, K. U., Bota, M., García Del Molino, L. C., Fitzgerald, G., Ram, K., He, M., Levine, J. M., Mitra, P., Huang, Z. J., Wang, X.-J., & Osten, P. (2017). Brain-wide Maps Reveal Stereotyped Cell-Type-Based Cortical Architecture and Subcortical Sexual Dimorphism. *Cell*, *171*(2), 456-469.e22. https://doi.org/10.1016/j.cell.2017.09.020

Kong, X., Kong, R., Orban, C., Wang, P., Zhang, S., Anderson, K., Holmes, A., Murray, J. D., Deco, G., Van Den Heuvel, M., & Yeo, B. T. T. (2021). Sensory-motor cortices shape functional connectivity dynamics in the human brain. *Nature Communications*, *12*(1), 6373. https://doi.org/10.1038/s41467-021-26704-y

Kringelbach, M. L., & Deco, G. (2020). Brain States and Transitions: Insights from Computational Neuroscience. *Cell Reports*, *32*(10), 108128. https://doi.org/10.1016/j.celrep.2020.108128

Lam, N. H., Borduqui, T., Hallak, J., Roque, A., Anticevic, A., Krystal, J. H., Wang, X.-J., & Murray, J. D. (2022). Effects of Altered Excitation-Inhibition Balance on Decision Making in a Cortical Circuit Model. *The Journal of Neuroscience*, *42*(6), 1035–1053. https://doi.org/10.1523/JNEUROSCI.1371-20.2021

Larsen, B., Cui, Z., Adebimpe, A., Pines, A., Alexander-Bloch, A., Bertolero, M., Calkins, M. E., Gur, R. E., Gur, R. C., Mahadevan, A. S., Moore, T. M., Roalf, D. R., Seidlitz, J., Sydnor, V. J., Wolf, D. H., & Satterthwaite, T. D. (2022). A developmental reduction of the excitation:inhibition ratio in association cortex during adolescence. *Science Advances*, *8*(5), eabj8750. https://doi.org/10.1126/sciadv.abj8750

Larsen, B., & Luna, B. (2018). Adolescence as a neurobiological critical period for the development of higher-order cognition. *Neuroscience & Biobehavioral Reviews*, *94*, 179–195. https://doi.org/10.1016/j.neubiorev.2018.09.005

Leonardi, N., & Van De Ville, D. (2015). On spurious and real fluctuations of dynamic functional connectivity during rest. *NeuroImage*, *104*, 430–436. https://doi.org/10.1016/j.neuroimage.2014.09.007

Li, J., Bzdok, D., Chen, J., Tam, A., Ooi, L. Q. R., Holmes, A. J., Ge, T., Patil, K. R., Jabbi, M., Eickhoff, S. B., Yeo, B. T. T., & Genon, S. (2022). Cross-ethnicity/race generalization failure of behavioral prediction from resting-state functional connectivity. *Science Advances*, *8*(11), eabj1812. https://doi.org/10.1126/sciadv.abj1812

Liégeois, R., Laumann, T. O., Snyder, A. Z., Zhou, J., & Yeo, B. T. T. (2017). Interpreting temporal fluctuations in resting-state functional connectivity MRI. *NeuroImage*, *163*, 437–455. https://doi.org/10.1016/j.neuroimage.2017.09.012

Lodato, S., Rouaux, C., Quast, K. B., Jantrachotechatchawan, C., Studer, M., Hensch, T. K., & Arlotta, P. (2011). Excitatory Projection Neuron Subtypes Control the Distribution of Local Inhibitory Interneurons in the Cerebral Cortex. *Neuron*, *69*(4), 763–779. https://doi.org/10.1016/j.neuron.2011.01.015

Luna, B., & Wright, C. (2016). Adolescent brain development: Implications for the juvenile criminal justice system. In K. Heilbrun, D. DeMatteo, & N. E. S. Goldstein (Eds.), *APA handbook of psychology and juvenile justice.* (pp. 91–116). American Psychological Association. https://doi.org/10.1037/14643-005

Margulies, D. S., Ghosh, S. S., Goulas, A., Falkiewicz, M., Huntenburg, J. M., Langs, G., Bezgin, G., Eickhoff, S. B., Castellanos, F. X., Petrides, M., Jefferies, E., & Smallwood, J. (2016). Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proceedings of the National Academy of Sciences*, *113*(44), 12574–12579. https://doi.org/10.1073/pnas.1608282113

Marín, O. (2016). Developmental timing and critical windows for the treatment of psychiatric disorders. *Nature Medicine*, *22*(11), 1229–1238. https://doi.org/10.1038/nm.4225

McGee, A. W., Yang, Y., Fischer, Q. S., Daw, N. W., & Strittmatter, S. M. (2005). Experience-Driven Plasticity of Visual Cortex Limited by Myelin and Nogo Receptor. *Science*, *309*(5744), 2222–2226. https://doi.org/10.1126/science.1114362

McRae, P. A., Rocco, M. M., Kelly, G., Brumberg, J. C., & Matthews, R. T. (2007). Sensory Deprivation Alters Aggrecan and Perineuronal Net Expression in the Mouse Barrel Cortex. *The Journal of Neuroscience*, *27*(20), 5405–5413. https://doi.org/10.1523/JNEUROSCI.5425-06.2007

Moore, T. M., Reise, S. P., Gur, R. E., Hakonarson, H., & Gur, R. C. (2015). Psychometric properties of the Penn Computerized Neurocognitive Battery. *Neuropsychology*, *29*(2), 235–246. https://doi.org/10.1037/neu0000093

Nelson, C. A., & Gabard-Durnam, L. J. (2020). Early Adversity and Critical Periods: Neurodevelopmental Consequences of Violating the Expectable Environment. *Trends in Neurosciences*, *43*(3), 133–143. https://doi.org/10.1016/j.tins.2020.01.002

Nelson, S. (1991). Temporal interactions in the cat visual system. III. Pharmacological studies of cortical suppression suggest a presynaptic mechanism. *The Journal of*

Neuroscience, *11*(2), 369–380. https://doi.org/10.1523/JNEUROSCI.11-02-00369.1991

Nørgaard, M., Beliveau, V., Ganz, M., Svarer, C., Pinborg, L. H., Keller, S. H., Jensen, P. S., Greve, D. N., & Knudsen, G. M. (2021). A high-resolution in vivo atlas of the human brain's benzodiazepine binding site of GABAA receptors. *NeuroImage*, *232*, 117878. https://doi.org/10.1016/j.neuroimage.2021.117878

Olsen, R. W., & Sieghart, W. (2009). GABAA receptors: Subtypes provide diversity of function and pharmacology. *Neuropharmacology*, *56*(1), 141–148. https://doi.org/10.1016/j.neuropharm.2008.07.045

Pang, J. C., Aquino, K. M., Oldehinkel, M., Robinson, P. A., Fulcher, B. D., Breakspear, M., & Fornito, A. (2023). Geometric constraints on human brain function. *Nature*. https://doi.org/10.1038/s41586-023-06098-1

Paquola, C., Bethlehem, R. A., Seidlitz, J., Wagstyl, K., Romero-Garcia, R., Whitaker, K. J., Vos De Wael, R., Williams, G. B., NSPN Consortium, Vértes, P. E., Margulies, D. S., Bernhardt, B., & Bullmore, E. T. (2019). Shifts in myeloarchitecture characterise adolescent development of cortical gradients. *ELife*, *8*, e50482. https://doi.org/10.7554/eLife.50482

Park, B., Hong, S.-J., Valk, S. L., Paquola, C., Benkarim, O., Bethlehem, R. A. I., Di Martino, A., Milham, M. P., Gozzi, A., Yeo, B. T. T., Smallwood, J., & Bernhardt, B. C. (2021). Differences in subcortico-cortical interactions identified from connectome and microcircuit models in autism. *Nature Communications*, *12*(1), 2225. https://doi.org/10.1038/s41467-021-21732-0

Paus, T., Keshavan, M., & Giedd, J. N. (2008). Why do many psychiatric disorders emerge during adolescence? *Nature Reviews Neuroscience*, *9*(12), 947–957. https://doi.org/10.1038/nrn2513

Perica, M. I., Calabro, F. J., Larsen, B., Foran, W., Yushmanov, V. E., Hetherington, H., Tervo-Clemmens, B., Moon, C.-H., & Luna, B. (2022). Development of frontal GABA and glutamate supports excitation/inhibition balance from adolescence into adulthood. *Progress in Neurobiology*, *219*, 102370. https://doi.org/10.1016/j.pneurobio.2022.102370

Pines, A. R., Larsen, B., Cui, Z., Sydnor, V. J., Bertolero, M. A., Adebimpe, A., Alexander-Bloch, A. F., Davatzikos, C., Fair, D. A., Gur, R. C., Gur, R. E., Li, H., Milham, M. P., Moore, T. M., Murtha, K., Parkes, L., Thompson-Schill, S. L., Shanmugan, S., Shinohara, R. T., … Satterthwaite, T. D. (2022). Dissociable multi-scale patterns of development in personalized brain networks. *Nature Communications*, *13*(1), 2647. https://doi.org/10.1038/s41467-022-30244-4

Reh, R. K., Dias, B. G., Nelson, C. A., Kaufer, D., Werker, J. F., Kolb, B., Levine, J. D., & Hensch, T. K. (2020). Critical period regulation across multiple timescales. *Proceedings of the National Academy of Sciences*, *117*(38), 23242–23251. https://doi.org/10.1073/pnas.1820836117

Romer, A. L., & Pizzagalli, D. A. (2021). Is executive dysfunction a risk marker or consequence of psychopathology? A test of executive function as a prospective predictor and outcome of general psychopathology in the adolescent brain cognitive development study®. *Developmental Cognitive Neuroscience*, *51*, 100994. https://doi.org/10.1016/j.dcn.2021.100994

Rudolph, U., & Knoflach, F. (2011). Beyond classical benzodiazepines: Novel therapeutic potential of GABAA receptor subtypes. *Nature Reviews Drug Discovery*, *10*(9), 685–697. https://doi.org/10.1038/nrd3502

Satterthwaite, T. D., Elliott, M. A., Ruparel, K., Loughead, J., Prabhakaran, K., Calkins, M. E., Hopson, R., Jackson, C., Keefe, J., Riley, M., Mentch, F. D., Sleiman, P., Verma,

R., Davatzikos, C., Hakonarson, H., Gur, R. C., & Gur, R. E. (2014). Neuroimaging of the Philadelphia Neurodevelopmental Cohort. *NeuroImage*, *86*, 544–553. https://doi.org/10.1016/j.neuroimage.2013.07.064

Selten, M., Van Bokhoven, H., & Nadif Kasri, N. (2018). Inhibitory control of the excitatory/inhibitory balance in psychiatric disorders. *F1000Research*, *7*, 23. https://doi.org/10.12688/f1000research.12155.1

Shaw, A. D., Knight, L., Freeman, T. C. A., Williams, G. M., Moran, R. J., Friston, K. J., Walters, J. T. R., & Singh, K. D. (2019). Oscillatory, Computational, and Behavioral Evidence for Impaired GABAergic Inhibition in Schizophrenia. *Schizophrenia Bulletin*, sbz066. https://doi.org/10.1093/schbul/sbz066

Shulman, E. P., Smith, A. R., Silva, K., Icenogle, G., Duell, N., Chein, J., & Steinberg, L. (2016). The dual systems model: Review, reappraisal, and reaffirmation. *Developmental Cognitive Neuroscience*, *17*, 103–117. https://doi.org/10.1016/j.dcn.2015.12.010

Sigel, E. (2002). Mapping of the Benzodiazepine Recognition Site on GABA$_A$ Receptors. *Current Topics in Medicinal Chemistry*, *2*(8), 833–839. https://doi.org/10.2174/1568026023393444

Silveri, M. M., Sneider, J. T., Crowley, D. J., Covell, M. J., Acharya, D., Rosso, I. M., & Jensen, J. E. (2013). Frontal Lobe γ-Aminobutyric Acid Levels During Adolescence: Associations with Impulsivity and Response Inhibition. *Biological Psychiatry*, *74*(4), 296–304. https://doi.org/10.1016/j.biopsych.2013.01.033

Soh, S.-E., Tint, M. T., Gluckman, P. D., Godfrey, K. M., Rifkin-Graboi, A., Chan, Y. H., Stunkel, W., Holbrook, J. D., Kwek, K., Chong, Y.-S., Saw, S. M., the GUSTO Study Group, Sheppard, A., Chinnadurai, A., Ferguson-Smith, A., Goh, A. E. N., Biswas, A., Chia, A., Leutscher-Broekman, B., … Venkatesh, S. K. (2014). Cohort Profile: Growing Up in Singapore Towards healthy Outcomes (GUSTO) birth cohort study. *International Journal of Epidemiology*, *43*(5), 1401–1409. https://doi.org/10.1093/ije/dyt125

Sohal, V. S., & Rubenstein, J. L. R. (2019). Excitation-inhibition balance as a framework for investigating mechanisms in neuropsychiatric disorders. *Molecular Psychiatry*, *24*(9), 1248–1257. https://doi.org/10.1038/s41380-019-0426-0

Steinberg, L. (2010). A dual systems model of adolescent risk-taking. *Developmental Psychobiology*, n/a-n/a. https://doi.org/10.1002/dev.20445

Stephan, K. E., Weiskopf, N., Drysdale, P. M., Robinson, P. A., & Friston, K. J. (2007). Comparing hemodynamic models with DCM. *NeuroImage*, *38*(3), 387–401. https://doi.org/10.1016/j.neuroimage.2007.07.040

Stern, E. A., Maravall, M., & Svoboda, K. (2001). Rapid Development and Plasticity of Layer 2/3 Maps in Rat Barrel Cortex In Vivo. *Neuron*, *31*(2), 305–315. https://doi.org/10.1016/S0896-6273(01)00360-9

Sun, Y. J., Wu, G. K., Liu, B., Li, P., Zhou, M., Xiao, Z., Tao, H. W., & Zhang, L. I. (2010). Fine-tuning of pre-balanced excitation and inhibition during auditory cortical development. *Nature*, *465*(7300), 927–931. https://doi.org/10.1038/nature09079

Sydnor, V. J., Larsen, B., Bassett, D. S., Alexander-Bloch, A., Fair, D. A., Liston, C., Mackey, A. P., Milham, M. P., Pines, A., Roalf, D. R., Seidlitz, J., Xu, T., Raznahan, A., & Satterthwaite, T. D. (2021). Neurodevelopment of the association cortices: Patterns, mechanisms, and implications for psychopathology. *Neuron*, *109*(18), 2820–2846. https://doi.org/10.1016/j.neuron.2021.06.016

Sydnor, V. J., Larsen, B., Seidlitz, J., Adebimpe, A., Alexander-Bloch, A. F., Bassett, D. S., Bertolero, M. A., Cieslak, M., Covitz, S., Fan, Y., Gur, R. E., Gur, R. C., Mackey, A. P., Moore, T. M., Roalf, D. R., Shinohara, R. T., & Satterthwaite, T. D. (2023).

Intrinsic activity development unfolds along a sensorimotor–association cortical axis in youth. *Nature Neuroscience*, *26*(4), 638–649. https://doi.org/10.1038/s41593-023-01282-y

Sydnor, V. J., & Satterthwaite, T. D. (2023). Neuroimaging of plasticity mechanisms in the human brain: From critical periods to psychiatric conditions. *Neuropsychopharmacology*, *48*(1), 219–220. https://doi.org/10.1038/s41386-022-01415-0

Takesian, A. E., & Hensch, T. K. (2013). Balancing Plasticity/Stability Across Brain Development. In *Progress in Brain Research* (Vol. 207, pp. 3–34). Elsevier. https://doi.org/10.1016/B978-0-444-63327-9.00001-1

Tervo-Clemmens, B., Calabro, F. J., Parr, A. C., Fedor, J., Foran, W., & Luna, B. (2022). *A Canonical Trajectory of Executive Function Maturation During the Transition from Adolescence to Adulthood* [Preprint]. PsyArXiv. https://doi.org/10.31234/osf.io/73yfv

Tournier, J.-D., Smith, R., Raffelt, D., Tabbara, R., Dhollander, T., Pietsch, M., Christiaens, D., Jeurissen, B., Yeh, C.-H., & Connelly, A. (2019). MRtrix3: A fast, flexible and open software framework for medical image processing and visualisation. *NeuroImage*, *202*, 116137. https://doi.org/10.1016/j.neuroimage.2019.116137

Toyoizumi, T., Miyamoto, H., Yazaki-Sugiyama, Y., Atapour, N., Hensch, T. K., & Miller, K. D. (2013). A Theory of the Transition to Critical Period Plasticity: Inhibition Selectively Suppresses Spontaneous Activity. *Neuron*, *80*(1), 51–63. https://doi.org/10.1016/j.neuron.2013.07.022

Trachtenberg, J. T. (2015). Competition, inhibition, and critical periods of cortical plasticity. *Current Opinion in Neurobiology*, *35*, 44–48. https://doi.org/10.1016/j.conb.2015.06.006

Trakoshis, S., Martínez-Cañada, P., Rocchi, F., Canella, C., You, W., Chakrabarti, B., Ruigrok, A. N., Bullmore, E. T., Suckling, J., Markicevic, M., Zerbi, V., MRC AIMS Consortium, Baron-Cohen, S., Gozzi, A., Lai, M.-C., Panzeri, S., & Lombardo, M. V. (2020). Intrinsic excitation-inhibition imbalance affects medial prefrontal cortex differently in autistic men versus women. *ELife*, *9*, e55684. https://doi.org/10.7554/eLife.55684

Turrigiano, G. G., & Nelson, S. B. (2004). Homeostatic plasticity in the developing nervous system. *Nature Reviews Neuroscience*, *5*(2), 97–107. https://doi.org/10.1038/nrn1327

Van Essen, D. C., Smith, S. M., Barch, D. M., Behrens, T. E. J., Yacoub, E., & Ugurbil, K. (2013). The WU-Minn Human Connectome Project: An overview. *NeuroImage*, *80*, 62–79. https://doi.org/10.1016/j.neuroimage.2013.05.041

Verster, J. C., & Volkerts, E. R. (2006). Clinical Pharmacology, Clinical Efficacy, and Behavioral Toxicity of Alprazolam: A Review of the Literature. *CNS Drug Reviews*, *10*(1), 45–76. https://doi.org/10.1111/j.1527-3458.2004.tb00003.x

Vinogradov, S., Chafee, M. V., Lee, E., & Morishita, H. (2023). Psychosis spectrum illnesses as disorders of prefrontal critical period plasticity. *Neuropsychopharmacology*, *48*(1), 168–185. https://doi.org/10.1038/s41386-022-01451-w

Wang, H., Fan, L., Song, M., Liu, B., Wu, D., Jiang, R., Li, J., Li, A., Banaschewski, T., Bokde, A. L. W., Quinlan, E. B., Desrivières, S., Flor, H., Grigis, A., Garavan, H., Chaarani, B., Gowland, P., Heinz, A., Ittermann, B., … Jiang, T. (2021). Functional Connectivity Predicts Individual Development of Inhibitory Control during Adolescence. *Cerebral Cortex*, *31*(5), 2686–2700. https://doi.org/10.1093/cercor/bhaa383

Wang, P., Kong, R., Kong, X., Liégeois, R., Orban, C., Deco, G., Van Den Heuvel, M. P., & Thomas Yeo, B. T. (2019). Inversion of a large-scale circuit model reveals a cortical

hierarchy in the dynamic resting human brain. *Science Advances*, *5*(1), eaat7854. https://doi.org/10.1126/sciadv.aat7854

Werker, J. F., & Hensch, T. K. (2015). Critical Periods in Speech Perception: New Directions. *Annual Review of Psychology*, *66*(1), 173–196. https://doi.org/10.1146/annurev-psych-010814-015104

Willis, A., Pratt, J. A., & Morris, B. J. (2022). Enzymatic Degradation of Cortical Perineuronal Nets Reverses GABAergic Interneuron Maturation. *Molecular Neurobiology*, *59*(5), 2874–2893. https://doi.org/10.1007/s12035-022-02772-z

Wolf, D. H., Satterthwaite, T. D., Loughead, J., Pinkham, A., Overton, E., Elliott, M. A., Dent, G. W., Smith, M. A., Gur, R. C., & Gur, R. E. (2011). Amygdala abnormalities in first-degree relatives of individuals with schizophrenia unmasked by benzodiazepine challenge. *Psychopharmacology*, *218*(3), 503–512. https://doi.org/10.1007/s00213-011-2348-7

Wu, J., Ngo, G. H., Greve, D., Li, J., He, T., Fischl, B., Eickhoff, S. B., & Yeo, B. T. T. (2018). Accurate nonlinear mapping between MNI volumetric and FreeSurfer surface coordinate systems. *Human Brain Mapping*, *39*(9), 3793–3808. https://doi.org/10.1002/hbm.24213

Wu, Y. K., Miehl, C., & Gjorgjieva, J. (2022). Regulation of circuit organization and function through inhibitory synaptic plasticity. *Trends in Neurosciences*, *45*(12), 884–898. https://doi.org/10.1016/j.tins.2022.10.006

Xu, W., Löwel, S., & Schlüter, O. M. (2020). Silent Synapse-Based Mechanisms of Critical Period Plasticity. *Frontiers in Cellular Neuroscience*, *14*, 213. https://doi.org/10.3389/fncel.2020.00213

Xue, M., Atallah, B. V., & Scanziani, M. (2014). Equalizing excitation–inhibition ratios across visual cortical neurons. *Nature*, *511*(7511), 596–600. https://doi.org/10.1038/nature13321

Yan, X., Kong, R., Xue, A., Yang, Q., Orban, C., An, L., Holmes, A. J., Qian, X., Chen, J., Zuo, X.-N., Zhou, J. H., Fortier, M. V., Tan, A. P., Gluckman, P., Chong, Y. S., Meaney, M. J., Bzdok, D., Eickhoff, S. B., & Yeo, B. T. T. (2023). Homotopic local-global parcellation of the human cerebral cortex from resting-state functional connectivity. *NeuroImage*, *273*, 120010. https://doi.org/10.1016/j.neuroimage.2023.120010

Yang, G. J., Murray, J. D., Wang, X.-J., Glahn, D. C., Pearlson, G. D., Repovs, G., Krystal, J. H., & Anticevic, A. (2016). Functional hierarchy underlies preferential connectivity disturbances in schizophrenia. *Proceedings of the National Academy of Sciences*, *113*(2). https://doi.org/10.1073/pnas.1508436113

Yao, H. K., Guet-McCreight, A., Mazza, F., Moradi Chameh, H., Prevot, T. D., Griffiths, J. D., Tripathy, S. J., Valiante, T. A., Sibille, E., & Hay, E. (2022). Reduced inhibition in depression impairs stimulus processing in human cortical microcircuits. *Cell Reports*, *38*(2), 110232. https://doi.org/10.1016/j.celrep.2021.110232

Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman, J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, *106*(3), 1125–1165. https://doi.org/10.1152/jn.00338.2011

SUPPLEMENTARY INFORMATION

Figure S1. (A) Comparison of the total costs in the validation set and the test set of the alprazolam session. (B) Comparison of the total costs in the validation set and the test set of the placebo session. The total costs in the validation set correspond to the lowest 10 validation costs, generated from the top 10 sets of parameters from the validation set. The total costs of the test set were computed using the same 10 sets of parameters and the structural connectivity of the test set. For both alprazolam and placebo sessions, total costs of the test set were significantly lower than those of the validation set, suggesting that the parameters generalized well from the validation set to the test set. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers.

Desikan parcellation, ROI coverage > 50%

Figure S2. Desikan parcellation with 50% ROI coverage threshold. Due to limited FOV of the alprazolam dataset, only ROIs with coverage higher than a pre-specified threshold were included. ROIs included (excluded) for analysis are colored in cyan (grey). 42 out of 68 ROIs are included.

Figure S3. (A to D) As a control analysis of the alprazolam E/I ratio contrast result, the same analysis was replicated with 4 additional training-validation-test participant splits. E/I ratio contrast was defined as the E/I ratio difference between the placebo session and the drug

session. 45 participants were randomly assigned to a different training, validation, and test set. (Left) Spatial distribution of E/I ratio contrast between placebo and drug session. (Right) The E/I ratio contrast decreases along a sensory-to-association axis. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers.

	split #1	split #2	split #3	split #4	split #5
split #1	1	0.9936	0.9922	0.9948	0.9853
split #2		1	0.9800	0.9896	0.9949
split #3			1	0.9979	0.9802
split #4				1	0.9909
split #5					1

Figure S4. Pairwise correlation between the regional E/I ratio contrast across different training-validation-test participant splits (split 1-5). The spatial distribution of region E/I ratio contrast are highly similar across splits (r = 0.9899 ± 0.0062, mean ± std). Only the upper triangle of the matrix is shown. We chose the split that had the highest median correlation of regional E/I ratio contrast with the other 4 splits as the most representative split (i.e., Split #1) and showed as Figure 3 of the main text .

E/I ratio contrast, more relaxed acceptable r$_E$ range

Figure S5. As a control analysis of the alprazolam E/I ratio contrast result, the acceptable excitatory firing rate range was set to be less strict. The range was changed from 2.7 – 3.3Hz to 2.5 – 3.5 Hz. We repeated the same analysis, and the model parameters were retrained using the same optimization approach. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers.

(A) Desikan parcellation, ROI coverage > 60%

(B) E/I ratio contrast, higher ROI coverage threshold

Figure S6. (A) Desikan parcellation with 60% ROI coverage threshold, 39 out of 68 ROIs are left. (B) As a control analysis of the alprazolam E/I ratio contrast result, the ROI coverage threshold was changed to be stricter. The threshold was raised from 50% to 60%. As a result, 3 more ROIs were removed from analysis. We repeated the same analysis, and the model parameters were retrained using the same optimization approach. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers.

(A) Yan parcellation, ROI coverage > 50%

(B) E/I ratio contrast, Yan parcellation

Figure S7. (A) Yan parcellation with 50% ROI coverage threshold, 51 out of 100 ROIs are left. (B) As a control analysis of the alprazolam E/I ratio contrast result, we changed the parcellation scheme to a higher-resolution 100-ROI parcellation. Yan parcellation has ROIs that are symmetric for the left and right hemispheres. We repeated the same analysis, and the model parameters were retrained using the same optimization approach. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers.

	split #1	relaxed r_E range	higher ROI coverage threshold	Yan
split #1	1	0.9911	0.9941	0.8986
relaxed r_E range		1	0.9758	0.8422
higher ROI coverage threshold			1	0.9220
Yan				1

Figure S8. Pairwise correlation between the regional E/I ratio contrast across different control analyses based on split #1. Split #1 corresponds to the results shown in Figure 3 of the main text. The spatial distribution of region E/I ratio contrast are highly similar across different control analyses (r = 0.9373 ± 0.0606, mean ± std). Only the upper triangle of the matrix is shown.

Figure S9. (A to D) PNC developmental analysis results obtained from 4 additional training-validation participants splits. (Left) The mean cortical E/I ratio decreases with increasing age. (Middle) The spatial distribution of regional rate of E/I ratio reduction. (Right) The regional rate of E/I ratio reduction followed a hierarchical sensorimotor-association (S-A) axis. To generate a training-validation split, 885 PNC participants were sorted according to age in an ascending order and divided into 29 age groups. Within each age groups, participants were

randomly assigned to training and validation sets. The random participant splits and analyses were repeated 5 times (One shown in Figure 4, the other four shown here). All 5 splits exhibited the similar pattern of overall E/I ratio reduction and spatial distribution of the rate of reduction. We chose the split that had the highest median correlation of regional rate of E/I ratio reduction with the other 4 splits as the most representative split and showed in the main result section. The shaded area depicts 95% confidence interval of the linear relationship. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers.

	split #1	split #2	split #3	split #4	split #5
split #1	1	0.9776	0.9777	0.9253	0.9921
split #2		1	0.9167	0.9835	0.9501
split #3			1	0.8295	0.9954
split #4				1	0.8783
split #5					1

Figure S10. Pairwise correlation between the regional rate of E/I ratio reduction of 5 training-validation participant splits. Split #1 corresponds to the results shown in Figure 4 of the main text. The spatial distributions of the rate of E/I ratio reduction are highly similar across the 5 splits (r = 0.9426 ± 0.0551, mean ± std). The surface maps of different splits are shown in Figure S8. Only the upper triangle of the matrix is shown.

Figure S11. PNC developmental analysis results with relaxed excitatory firing rates thresholds. This figure is similar to Figure 4 but with the acceptable excitatory firing rates range was set to be less strict (2.5Hz to 3.5Hz). (Left) The scatter plot of overall E/I ratio reduction during development. The mean cortical E/I ratio is the average E/I ratio across all ROIs. (Middle) The spatial distribution of region E/I ratio reduction rate. (Right) Box plot of the vertex-level E/I ratio grouped by 7 resting-state networks. The boxplots comprised values obtained by "transferring" the parameter estimates from the 68 Desikan parcels to all vertices (from the underlying cortical meshes) comprising each anatomical parcel. The vertex wise parameter values were then segregated based on the seven resting-state networks. Therefore, there were 3203, 2478, 1523, 1520, 1067, 1438 and 2886 values comprising the boxplots for somatomotor, visual, dorsal attention, ventral attention, limbic, control and default networks respectively. The shaded area depicts 95% confidence interval of the linear relationship. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers. The rate of E/I ratio reduction follows the sensorimotor-association (S-A) axis. E/I ratio exhibits the fastest rate of reduction in sensory regions compared to association regions.

Figure S12. PNC developmental analysis results in Yan 100-ROI parcellation. This figure is similar to Figure 4 but utilizes the Yan 100-ROI parcellation with symmetric left and right hemisphere ROIs. (Left) The scatter plot of overall E/I ratio reduction during development. The mean cortical E/I ratio is the average E/I ratio across all ROIs. (Middle) The spatial distribution of region E/I ratio reduction rate. (Right) Box plot of the vertex-level E/I ratio grouped by 7 resting-state networks. The boxplots comprised values obtained by "transferring" the parameter estimates from the 100 Yan parcels to all vertices (from the underlying cortical meshes) comprising each anatomical parcel. The vertex wise parameter values were then segregated based on the seven resting-state networks. Therefore, there were 3203, 2478, 1523, 1520, 1067, 1438 and 2886 values comprising the boxplots for somatomotor, visual, dorsal attention, ventral attention, limbic, control and default networks respectively. The shaded area depicts 95% confidence interval of the linear relationship. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers. The rate of E/I ratio reduction follows the sensorimotor-association (S-A) axis. E/I ratio exhibits the fastest rate of reduction in sensory regions compared to association regions.

| | relaxed r_E | | |
	split #1	range	Yan
split #1	1	0.9887	0.7622
relaxed r_E range		1	0.6791
Yan			1

Figure S13. Pairwise correlation between the regional rate of E/I ratio reduction of different control analyses based on split #1. Split #1 corresponds to the results shown in Figure 4 of the main text. The spatial distributions of the rate of E/I ratio reduction are highly similar across different control analyses (r = 0.8100 ± 0.1603, mean ± std). The surface maps of different cases are shown in the supplementary Figure S10 and S11. Only the upper triangle of the matrix is shown.

Figure S14. Box plots of E/I ratio estimated from the three domain-specific accuracy scores from Penn Computerized Neurocognitive Battery (CNB). For each 1-year interval, participants whose age are within this interval are extracted to form one age group. For each age group and cognitive score, participants with cognitive scores above the median are assigned to a high-performance group, the rest are assigned to a low-performance group. Both high- and low-performance groups are further divided into subgroups. Participants within each subgroup are randomly assigned to training and validation sets. Each box plot shows the E/I ratio of high- and low-performance group associated with each cognitive score. We observed no significant difference between E/I ratio of high- and low-performance groups for any of the three cognitive scores. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR.

Figure S15. PNC cognition analysis results obtained from 4 additional training-validation participants splits (Figure. S15 to S18). To generate a training-validation split, 885 PNC participants were sorted according to age in an ascending order. For each 1-year interval, participants whose age are within this interval are extracted to form one age group. For each age group, participants with cognitive scores above the median are assigned to a high-performance group, the rest are assigned to a low-performance group. Both high- and low-performance groups are further divided into 14 subgroups. Participants within each subgroup

are randomly assigned to training and validation sets. The random participant splits and analyses were repeated 5 times (One shown in Figure 6, the other four shown in S5-1 to S5-4). (A) Boxplots of age, (B) 'Overall accuracy', and (C) Mean cortical E/I ratio of high- and low-performance groups. (D) Spatial distribution of effect size of regional E/I ratio difference between high-performance and low-performance groups. (E) On a network level, the effect sizes of E/I ratio differences follow a hierarchical structure. We chose the split that had the highest median correlation of regional Cohen's d values with the other 4 splits as the most representative split and showed in the main result section. The results of the 4 splits are consistent with our main results. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. (F) ROI rankings along the sensorimotor-association (S-A) axis. Lower ranks were assigned to ROIs that were more towards the sensorimotor pole; higher ranks were assigned to ROIs that were more towards the association pole. (G) Agreement between the effect size of E/I ratio difference and S-A axis rank.

Figure S16. The 3rd training-validation participant split for PNC cognition analysis.

Figure S17. The 4th training-validation participant split for PNC cognition analysis.

Figure S18. The 5th training-validation participant split for PNC cognition analysis.

	split #1	split #2	split #3	split #4	split #5
split #1	1	0.7760	0.8784	0.8339	0.7057
split #2		1	0.4565	0.6741	0.9754
split #3			1	0.7793	0.3456
split #4				1	0.6247
split #5					1

Figure S19. Pairwise correlation of regional effect sizes (i.e., Cohen's d) between E/I ratios of high- and low-performance groups across 5 participant splits (split 1-5). For each split, high- and low-performance groups were divided into 14 subgroups with training and validation sets. The spatial distributions of the rate of E/I ratio reduction are highly similar across the 5 splits ($r = 0.7050 \pm 0.1911$, mean $\pm$ std). Split #1 corresponds to the results shown in Figure 5 of the main text. The surface maps of different splits are shown in the main result (Figure 5D) and supplementary information (Figure. S15 to S18). Only the upper triangle of the matrix is shown.

Figure S20. PNC cognition analysis results with relaxed excitatory firing rates thresholds. This figure is similar to Figure 5 but with the acceptable excitatory firing rates range was set to be less strict (2.5Hz to 3.5Hz). (A) Boxplots of age, (B) 'Overall accuracy', and (C) Mean cortical E/I ratio of high- and low-performance groups. (D) Spatial distribution of effect size of regional E/I ratio difference between high-performance and low-performance groups. (E) Box plot of Cohen's d of vertex-level E/I ratio differences grouped by 7 resting-state networks. The boxplots comprised values obtained by "transferring" the parameter estimates

from the 68 Desikan parcels to all vertices (from the underlying cortical meshes) comprising each anatomical parcel. The vertex wise parameter values were then segregated based on the seven resting-state networks. Therefore, there were 3203, 2478, 1523, 1520, 1067, 1438 and 2886 values comprising the boxplots for somatomotor, visual, dorsal attention, ventral attention, limbic, control and default networks respectively. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers. The difference of E/I ratio between high- and low-performance groups follows a hierarchical structure. Cohen's d of E/I ratio differences in cognition is larger in association regions compared to sensory regions. (F) ROI rankings along the sensorimotor-association (S-A) axis. Lower ranks were assigned to ROIs that were more towards the sensorimotor pole; higher ranks were assigned to ROIs that were more towards the association pole. (G) Agreement between the effect size of E/I ratio difference and S-A axis rank.

Figure S21. PNC cognition analysis results in Yan 100-ROI parcellation. This figure is similar to Figure 5 but utilizes the Yan 100-ROI parcellation with symmetric left and right hemisphere ROIs. (A) Boxplots of age, (B) 'Overall accuracy', and (C) Mean cortical E/I ratio of high- and low-performance groups. (D) Spatial distribution of effect size of regional E/I ratio difference between high-performance and low-performance groups. (E) Box plot of Cohen's d of vertex-level E/I ratio differences grouped by 7 resting-state networks. The boxplots comprised values obtained by "transferring" the parameter estimates from the 100

Yan parcels to all vertices (from the underlying cortical meshes) comprising each anatomical parcel. The vertex wise parameter values were then segregated based on the seven resting-state networks. Therefore, there were 3203, 2478, 1523, 1520, 1067, 1438 and 2886 values comprising the boxplots for somatomotor, visual, dorsal attention, ventral attention, limbic, control and default networks respectively. The boxes show the inter-quartile range (IQR) and the median. Whiskers indicate 1.5 IQR. Black crosses represent outliers. The difference of E/I ratio between high- and low-performance groups follows the sensory-to-association (SA) axis. Cohen's d of E/I ratio differences in cognition is larger in association regions compared to sensory regions. (F) ROI rankings along the sensorimotor-association (S-A) axis. Lower ranks were assigned to ROIs that were more towards the sensorimotor pole; higher ranks were assigned to ROIs that were more towards the association pole. (G) Agreement between the effect size of E/I ratio difference and S-A axis rank.

	split #1	relaxed r_E range	Yan
split #1	1	0.9258	0.8002
relaxed r_E range		1	0.7725
Yan			1

Figure S22. Pairwise correlation of regional effect sizes (i.e., Cohen's d) between E/I ratios of high- and low-performance groups across different control analyses based on split #1. Split #1 corresponds to the results shown in Figure 6 of the main text. The spatial distributions of the rate of E/I ratio reduction are highly similar across different control analyses (r = 0.8328 ± 0.0817, mean ± std). Only the upper triangle of the matrix is shown.