Genomic characterization of a nematode tolerance locus in sugar beet

Katharina Sielemann[1,2,†], Boas Pucker[3,†], Elena Orsini[4], Abdelnaser Elashry[4], Lukas

Schulte[1], Prisca Viehöver[1], Andreas E. Müller[4], Axel Schechert[4], Bernd Weisshaar[1], and

Daniela Holtgräwe[1,*]

[1]Genetics and Genomics of Plants, Center for Biotechnology (CeBiTec) & Faculty of Biology,
Bielefeld University, 33615 Bielefeld, Germany

[2]Graduate School DILS, Bielefeld Institute for Bioinformatics Infrastructure (BIBI), Bielefeld
University, 33615 Bielefeld, Germany

[3]Plant Biotechnology and Bioinformatics, Institute of Plant Biology & Braunschweig Integrated
Centre of Systems Biology (BRICS), TU Braunschweig, 38106 Braunschweig, Germany

[4]Strube Research GmbH & Co. KG, 38387 Söllingen, Germany

[†]These authors contributed equally to this work and share first authorship.

*Correspondence: Daniela Holtgräwe, dholtgra@cebitec.uni-bielefeld.de

Email addresses:

ksielemann@cebitec.uni-bielefeld.de
b.pucker@tu-braunschweig.de
e.orsini@strube-research.net
a.elashry@strube-research.net
lschulte@techfak.uni-bielefeld.de
viehoeve@cebitec.uni-bielefeld.de
a.mueller@strube-research.net
schechert@yahoo.com
bernd.weisshaar@uni-bielefeld.de
dholtgra@cebitec.uni-bielefeld.de

Keywords: sugar beet, nematode tolerance, *Beta vulgaris*, genome resources, mapping-by-

sequencing, NLP7

Running title: Nematode tolerance in sugar beet

Abstract

Infection by beet cyst nematodes (BCN, *Heterodera schachtii*) causes a serious disease of sugar beet, and climatic change is expected to improve the conditions for BCN infection. Yield and yield stability under adverse conditions are among the main breeding objectives. Breeding of BCN tolerant sugar beet cultivars offering high yield in the presence of the pathogen is therefore of high relevance. To identify causal genes providing tolerance against BCN infection, we combined several experimental and bioinformatic approaches. Relevant genomic regions were detected through mapping-by-sequencing using a segregating F2 population. DNA sequencing of contrasting F2 pools and analyses of allele frequencies for variant positions identified a single genomic region which confers nematode tolerance. The genomic interval was confirmed and narrowed down by genotyping with newly developed molecular markers. To pinpoint the causal genes within the nematode tolerance locus, we generated long read-based genome sequence assemblies of the tolerant parental breeding line Strube U2Bv and the susceptible reference line 2320Bv. We analyzed continuous sequences of the locus with regard to functional gene annotation and differential gene expression upon BCN infection. A cluster of genes with similarity to the *Arabidopsis thaliana* gene encoding nodule inception protein-like protein 7 (NLP7) was identified. Gene expression analyses confirmed transcriptional activity and revealed clear differences between susceptible and tolerant genotypes. Our findings provide new insights into the genomic basis of plant-nematode interactions that can be used to design and accelerate novel management strategies against BCN.

1   Introduction

Sugar beet is one of the most important crops in the northern hemisphere and contributes about 20% to world-wide sugar production. The ancestor of cultivated sugar beet is the sea beet *B. vulgaris* subsp. *maritima*. White Silesian Beet, a beet segregating in the F2 from a cross of fodder beet and chard, provided the narrow genetic base for today's sugar beet breeding [1]. An intense focus on yield led to a strong domestication bottleneck [2].

Among the economically most important pests of sugar beet (*Beta vulgaris* subsp. *vulgaris*) is the beet cyst nematode (BCN, *Heterodera schachtii*). The harmful effect of this nematode

is based on nutrient competition and disturbances in the root system of the host plant, which

lead to severe growth depression and yield reduction up to 60% [3]. The growth conditions

for *H. schachtii* are expected to improve as a result of global warming during the main

vegetation period, leading to an increased yield risk. The use of pathogen resistant or tolerant

elite varieties contributes significantly to improved sustainability of sugar production through

yield stability. Such varieties address agricultural and social demands on both conventional

and organic sugar beet cultivation. Resistant sugar beet varieties are available on the market,

but they have the disadvantage of penalized yield. On the contrary, nematode-tolerant beet

varieties do not react as strongly with yield depressions when infested with *H. schachtii* [4]

and therefore represent an economically remunerable trait to breed for. From a genetic point

of view, nematode tolerance is a quantitative resistance that inhibits BCN development [4,5]

in terms of both quantity and quality.

Molecular genetic markers associated with BCN resistance or tolerance are an important step

towards the identification of trait-associated genes [6]. The first known nematode resistance

gene, *Hs1^{pro-1}*, encodes a protein harboring a leucine-rich domain. *Hs1^{pro-1}* was introduced

into sugar beet as part of a translocation from the crop wild relative *Patellifolia procumbens*

[7]. Since then, other trait-associated genomic regions have been identified, including a region

conferring nematode tolerance from sea beet [8]. Stevanato *et al.* [6] identified a region in the

sea beet genotype WB242 on chromosome 5 (*BvmHs^{-1}*) and published a molecular marker,

designated SNP192, linked to nematode tolerance. Using segregation analyses, the group

was able to show the monogenic inheritance of the trait. However, the region was not further

defined or described, and variation for the trait nematode tolerance was observed by breeders

although SNP192 was homozygous in the genotypes studied.

To gain gene-level information for traits of interest, genome sequences of accessions

harboring these traits and comprehensive annotations are needed. In combination with

methods like mapping-by-sequencing (MBS), this allows a detailed investigation of

agronomically important regions. With regard to sugar beet, high-quality genome sequences

are available for two different accessions. These are the inbred line EL10 [9] and the

'reference genotype' KWS2320 (referred to here as '2320Bv') with the sequence identifier

Refbeet-1.0 [10]. An improved version that, among other data, also incorporates the genetic

map   BeetMap-3   [11]   is   publicly   available   with   the   identifier   RefBeet-1.5

(https://jbrowse.cebitec.uni-bielefeld.de/RefBeet1.5/). In addition, draft genome assemblies
of *B. patula* and *B. vulgaris* subsp. *maritima* WB42 [12] that represent crop wild relatives of
sugar beet, are publicly available. No high-quality genome sequence of a BCN tolerant
breeding line was available until now, and short read assemblies are not suitable for MBS
approaches and subsequent detailed trait locus analysis.

Genome sequences and DNA sequence data in general enable sequence comparison to
transfer gene function information from one species to another. For example, several
Arabidopsis thaliana genes were found to be homologous to the *Lotus japonicus* gene
NODULE INCEPTION [13] encoding the plant-specific transcription factor (TF) LjNIN [14]. *A.*
thaliana NIN-LIKE PROTEIN (NLP) family TFs control nitrate-responsive gene transcription
[15]. Among the NLPs is AtNLP7 (At4g24020) which is well described as a major regulator of
nitrate signaling [16-20]. Evidence for direct target genes regulated by AtNLP7 has recently
been published [16]. NLPs including AtNLP7 contain the sequence-conserved PB1 domain
(NCBI domain cd06407; [21]) which mediates protein-protein interactions [22]. These include
NLP-NLP interactions as well as interactions between NLPs and other factors [20,23]. Four
core amino acid residues within the PB1 domain (K867, D909, D911, and E913) are thought
to be required for NLP-NLP interactions [23]. In the context of nitrate response and plant
growth, mutants with substitutions of these core amino acid residues require a higher level of
expression than wildtype NLP7 [23]. Another highly conserved residue in NLPs is S205 which
serves as regulatory phosphorylation site [24].

In this study, we generated high-continuity genomic resources of the tolerant genotype Strube
U2Bv and the susceptible sugar beet reference genotype 2320Bv. MBS of tolerant and
susceptible lines to BCN was used together with RNA-Seq data generated from infection
experiments. This allowed to further characterize the genomic locus responsible for nematode
tolerance (NT) in sugar beet. The data generated will benefit breeding approaches and enable
a better control of the yield-diminishing BCN disease.

2   Results

2.1   Generation and phenotyping of a segregating F2 population

The mapping population STR-NT was derived from the nematode susceptible maternal line Strube U1Bv and the tolerant paternal line Strube U2Bv. A total of 406 F2:3 lines segregated for tolerance to BCN with continuous variation. The distribution of adjusted means of quantified tolerance per single plant ranged from -0.105 to 16.835, corresponding to 0 to 280 cysts counted per plant (Figure 1A). Twenty tolerant F2:3 lines with low numbers of cysts and 16 susceptible F2:3 lines with high number of cysts were identified as extremes of the phenotypic distribution. Analysis of CV suggested no further segregation for NT on these families so that their further investigation was performed through MBS.

Adjusted means per line (Figure 1B) were used for QTL mapping together with the genotypic data of 194 KASPar markers. A major QTL with an additive effect corresponding to 11 cysts was detected on the north of chromosome 5 at 10 cM. This QTL explained 23% of the phenotypic variance. The LOD support interval spanned the region from 9 to 13 cM between the new marker BR1180 and SNP192. The tolerant allele was derived from the nematode tolerant parent Strube U2Bv. No dominance effects were detected, heterozygous lines display an intermediate phenotype between tolerant and susceptible lines.

Figure 1: Phenotyping of the population STR-NT segregating for BCN tolerance. A) Histogram of adjusted data of single plants. **B)** Boxplot of the most tolerant and most susceptible F2-lines represented as families chosen for MBS. Adjusted means of the most

tolerant families range from 4.39 to 5.5 corresponding to 19-30 cysts. Adjusted means of the
most susceptible families range from 8.14 to 10.08 corresponding to 66-101 cysts.

For the MBS approach, the nine most tolerant lines were represented in individual libraries
and the eleven 'second best' tolerant lines were combined and sequenced in the "pool low".
The opposite phenotypic extreme included 16 susceptible lines. The nine most susceptible
lines were used to create individual libraries, whereas the remaining seven lines were
combined in the "pool high" and sequenced. For the tolerant lines, a total of 244 Gbp of
Illumina reads totaling a calculated 20.4x genome coverage were obtained, whereas for all
susceptible lines, 189 Gbp of reads with an estimated genome coverage of 19.8x were
reached.

2.2   Assembly and annotation of genome sequences of the nematode-tolerant parent
Strube U2Bv and the reference genotype 2320Bv

High assembly continuity for both, the nematode tolerant parent Strube U2Bv and the
susceptible reference genotype 2320Bv, is indicated by the N50 values (Table 1). The
assembly sizes of 596 Mbp (U2BvONT) and 573 Mbp (2320BvONT) exceed the size of
RefBeet-1.5 by 29 Mbp and 6 Mbp, respectively. RepeatMasker results indicated that both
assemblies have a repeat content of more than 65%.

The initial assemblies of Strube U2Bv and 2320Bv prior to the scaffolding process comprised
206 and 129 sequences, respectively. In total, 87.0% total bases of U2BvONT, represented
by 60 initial contigs, were anchored to nine pseudochromosomes using Beet-Map3 markers,
whereas only a slightly higher percentage of bases (88.3%) were genetically anchored in
2320BvONT represented by 59 initial contigs. In summary, 1,237 markers anchored the
2320BvONT assembly, whereas 1,238 markers anchored the scaffolds in the U2BvONT
assembly. A new set of 187 markers was designed based on data for Strube U2Bv and
successfully applied to genotyping in the StrUBv F2-population. A subset of 165 markers
confirmed the co-linearity of both assemblies and with Beet-Map3 (Additional file S1A).
However, these markers did not always allow exact genetic anchoring because of missing
information regarding the orientation of contigs, therefore such ambiguous contigs were not
placed in the pseudochromosomes.

Table 1: Assembly statistics of U2BvONT and 2320BvONT v1.0. Assembly size, number of contigs, N50 values, BUSCO completeness, repeat content, number of predicted genes, and number of predicted mRNAs are shown.

	2320BvONT v1.0	U2BvONT v1.0
Assembly size [bp]	573,025,584	596,437,702
Sequences	79 (9 pseudochromosomes + 70 contigs larger than 100 kb)	155 (9 pseudochromosomes + 146 contigs larger than 100 kb)
N50 [bp]	54,419,778	54,373,962
BUSCOs (n=2326)	C:93.7% [S:92.1%, D:1.6%], F:3.0%, M:3.3%[*]	C:93.6% [S:92.0%, D:1.6%], F:3.2%, M:3.2%[*]
Repeat content	65.21%	65.85%
Number of predicted genes	27,840	28,871
Number of predicted mRNAs	36,350	36,728

[*]Abbreviations: C = Complete, S = Single-copy, D = Duplicated, F = Fragmented, M = Missing

Synteny analysis between 2320BvONT and RefBeet annotations revealed a mRNA-based synteny of 97.3% (35,363) with a depth of 1 indicating a high similarity. Duplicated mRNAs in 2320BvONT have, in comparison to RefBeet, a proportion of 0.6% (231), and 2.1% (756) of 2320BvONT mRNAs were not found to be syntenic with any RefBeet mRNAs.

Investigation of the synteny between 2320BvONT and U2BvONT showed a 1:1 relation of 98.3% (35,721) of all mRNAs. Duplicated mRNAs in 2320BvONT occur, in comparison to U2BvONT, with 0.6% (200); 1.2% (429) have no syntenic anchor in U2BvONT. Genes without a syntenic counterpart were investigated to identify potential functions not present in the respective other genome. However, BLAST and InterProScan revealed no unique hit in either of the genomes.

Several large structural variations differentiate the genomes of 2320BvONT and U2BvONT. In total, 63 Mbp, divided into 88 different regions, are inverted between both genomes. The largest inversion on chromosome 3 (chr3) with a size of approximately 21 Mbp spans almost three contigs. Additionally, more than 3,000 translocations with a combined size of approximately 33 Mbp were identified (Additional file S2A).

2.3 NT locus detection through mapping-by-sequencing

A total of 2,049,007 variant positions were identified by comparing variants identified as homozygous in both parents and as heterozygous in the F1. The average delta allele frequency (dAF) in 10 SNP windows across all chromosomes is approximately 0.103 (±0.079). A dAF > 0.5 in a 10 SNP window was only detected on chr5 (Figure 2A) and on no other chromosome or contig of the U2BvONT assembly. The dAF plots of all nine chromosomes are provided in Additional file S2B. The borders of our locus of interest were restricted by the occurrence of a dAF > 0.5 throughout five consecutive 10 SNP windows. This 50 SNP window delimitation resulted in a genomic interval ranging from position 452,859 bp – 4,557,625 bp on chr5 of the U2BvONT assembly.

This clear interval of about 4 Mbp from MBS was further restricted using marker analyses on a few extreme F2 genotypes and the established susceptible and tolerant standard lines. The genetic markers MH00/01, BR1180 and additional flanking markers including SNP192 enabled, by graphical genotyping of recombination events in the phenotyped lines, a further containment of the NT locus (Additional file S2C). The size was restricted to about 0.7 Mbp with coordinates 1,321,396 – 2,021,946 bp in U2BvONT and 1,389,131 – 2,154,734 bp in 2320BvONT. The published marker SNP192 is located further south on chr5 at position 2,700,363 bp in U2BvONT and 2,741,678 bp in 2320BvONT (Figure 2B). The sequences of the whole region are continuous in both assemblies.

Figure 2: Boundaries of the NT locus region. A) Delta allele frequency (y-axis) distribution on chr5 of the U2BvONT assembly. Higher dAF, purple color; lower dAF, light blue. The chromosomal position in Mbp is represented on the x-axis. **B)** Dot plot comparing the NT locus region in U2BvONT and 2320BvONT. Magenta and purple dots represent high BLAST identity between both genome sequences. The NT locus was further delimited by the genetic markers MH00/01 and BR1180. Marker SNP192 [6] is linked but located further to the south of the chromosome. The 4-letter codes represent gene IDs (see text).

2.4 Characterisation of the NT locus

Overall, very high synteny was detected within the marker-restricted NT locus in U2BvONT and 2320BvONT (Figure 2). However, reduced synteny was observed between the genes gfmq and rycp. Within this region, we identified a cluster of genes (Figure 3) with similarity to *AtNLP7*. Analysis of the direct target genes of the AtNLP7 TF revealed that genes annotated with 'response to nematode' are significantly overrepresented among the targets (Additional

file S1B). Several other terms possibly related to BCN infection, like 'response to stress', 'signal transduction', and 'response to other organism' were found to be overrepresented as well. Therefore, we focused on this region, which is smaller than the marker-restricted NT locus and call it 'functionally restricted NT locus'.

The number of *BvNLP7* genes differs in both assemblies. In total, four and three genes are functionally annotated as 'NLP7' in the U2BvONT and 2320BvONT genome assembly, respectively. All of them are located within the delimited NT locus (Figure 3, Table 2). A manual check of reading frames in the region of both assemblies revealed no additional *NLP7*-like sequence.

Figure 3: Overview of the NT locus (top) and illustration of the functional restriction of the NT locus (bottom). Each line represents the sequence region between the genes gfmq and rycp for the respective genotype/species (*B. vulgaris* subsp. *vulgaris*: U2BvONT (Strube

U2Bv); 2320BvONT (2320Bv); EL10; *B. vulgaris* subsp. *maritima* WB42 (Bmar), *B. patula*
BETA548 (Bptu), RefBeet 1.5 (RefBeet)). The grey lines between the sequences indicate
synteny. Some lines are highlighted in blue to show synteny between the genes with similarity
to *AtNLP7*. *Bidirectional sugar transporter SWEET7*: gfmq, egca, owpo; *Histone H4*: pdig,
aunc, udai, edia, ptof; *BvNLP7*: hany, tlqs, mdsr, rycp, nxxw; other: azip, rfzi, upsq.

Table 2: Gene/allele names for the *NLP7*-like genes. The short name is used for more
clarity in the text. T = tolerant genotype (Strube U2Bv); S = susceptible genotype (2320Bv).

Short name	Genotype	Gene Identifier
NLP7-T1	Strube U2Bv	Bv05_g11459_hany.t1
NLP7-T2	Strube U2Bv	Bv05_g11464_mdsr.t1
NLP7-T3	Strube U2Bv	Bv05_g11461_tlqs.t1
NLP7-T4	Strube U2Bv	Bv05_g11465_rycp.t1
NLP7-S1	2320Bv	Bv05_g11045_mdsr.t1
NLP7-S2	2320Bv	Bv05_g11050_nxxw.t1
NLP7-S3	2320Bv	Bv05_g11053_rycp.t1

The genes with similarity to *AtNLP7* were compared in multiple sequence alignments at
coding sequence (CDS) and amino acid (aa) sequence level (Additional file S3, Additional file
S4, Additional file S5, Additional file S1C).

The CDS of *NLP7-T4* and *NLP7-S3* show an identity of 99.5% (Additional file S1C) with only
15 single nucleotide variant positions. At the aa level, these encoded proteins differ only at
two amino acid positions, namely isoleucine to valine and asparagine to serine (I878V and
N927S in U2BvONT > 2320BvONT). Both conservative exchanges are a result of a single

nucleotide variant. Similarly, *NLP7-T2* shows CDS identity to *NLP7-S1* of 93.49% (Additional
file S1C). Due to the high identity, we formally consider these gene structures as allelic. *NLP7-*
S2 is truncated in comparison to *AtNLP7* and codes only for the C-terminal part of the protein.

All candidate genes were compared to *AtNLP7* via percent identity matrices (Additional file
S1C). The comparison was performed for CDS and aa sequences as well as for the PB1
domain. The U2BvONT *NLP7*-like candidate genes show a relatively low sequence identity
to *AtNLP7* (Additional file S1C, CDS), ranging from 43.2% (*NLP7-T2*), 44.6% (*NLP7-T1*), and
45.1% (*NLP7-T3*) to 63.5% (*NLP7-T4*). For the three 2320BvONT *BvNLP7* genes, the
sequence identities are comparable (44% (*NLP7-S1*), 52.6% (*NLP7-S2*) and 63.5% (*NLP7-*
S3)). The core aa positions of the PB1 domain are completely missing from the sequence of
NLP7-T2. K867, D909 and E913 are conserved in all other candidates and D911 is conserved
in the candidates except NLP7-S1 and NLP7-T2. The completely conserved residue S205 as
well as the PB1 domain are absent from NLP7-S2 due to its C-terminal truncation. The protein
sequences of NLP7-T4 and NLP7-S3 carry S205, and in NLP7-T2, NLP7-T3 and NLP7-S1
this position is substituted by threonine (S205T), another polar amino acid and possible
phosphorylation site. Only the protein sequence of NLP7-T1 holds a lysine (S205K).

2.5 Tolerant and susceptible genotypes show diverse expression of *BvNLP7* genes

Gene expression was investigated in an infection assay with *H. schachtii*. A principal
component analysis (PCA) was conducted to assess the sample distribution (Additional file
S2D). Next, differentially expressed genes (DEGs) were identified. A total number of 2,263
genes with a padj < 0.05 was detected to be differentially expressed between all samples of
the tolerant and all samples of the susceptible lines (Additional file S1D). Normalized counts
were generated for all U2BvONT genes in tolerant (BR12 and Strube U2Bv) and BCN
susceptible lines (Strube U1Bv and SUS3). All four *BvNLP7* genes are expressed in all
genotypes under both conditions. Clear differences are visible for the four *BvNLP7* genes. In
particular, the two genes *NLP7-T1* and *NLP7-T2* are significantly lower expressed in both
susceptible genotypes compared to both tolerant genotypes.

Figure 4: Mean normalized counts (n=3) for the *BvNLP7* genes. RNA-Seq mappings were performed against the U2BvONT assembly. Dark grey bars represent inoculated samples, whereas light grey bars show the counts for non-inoculated samples. BR12 and Strube U2Bv are tolerant genotypes, whereas Strube U1Bv and SUS3 are susceptible genotypes. A) *NLP7-T1*, B) *NLP7-T3*, C) *NLP7-T2*, D) *NLP7-T4*. * = padj < 0.05, ** = padj < 0.01, *** = padj < 0.001.

The genes *NLP7-T1* and *NLP7-T2* are upregulated in tolerant genotypes in comparison to susceptible genotypes, independent of the inoculation. In addition, all four *BvNLP7* genes are significantly higher expressed in inoculated samples of the most tolerant F2 line BR12 than

in inoculated SUS3 samples (Table 2; Figure 4). Further, both NLP-T1 and NLP-T2 are
significantly higher expressed in samples of the inoculated tolerant line U2Bv than in samples
of the inoculated susceptible line U1Bv.

Table 3: Log$_2$ fold change (log2FC) and adjusted p-value (padj) of the expression
analysis for all *BvNLP7* genes. A positive value for the log2FC indicates higher expression
in tolerant genotypes. The comparison 'all tolerant vs. all susceptible' involves all samples,
inoculated and not inoculated, stratified by the tolerant vs. susceptible genotypes.

Short name	U2BvONT gene identifier	log2FC	Padj	Comparison
NLP7-T1	Bv05_g11459_hany.t1	2.97	2.5e-10	inoculated tolerant line BR12 vs. inoculated susceptible line SUS3
NLP7-T2	Bv05_g11464_mdsr.t1	2.99	2.4e-12	inoculated tolerant line BR12 vs. inoculated susceptible line SUS3
NLP7-T3	Bv05_g11461_tlqs.t1	0.82	0.027	inoculated tolerant line BR12 vs. inoculated susceptible line SUS3
NLP7 -T4	Bv05_g11465_rycp.t1	1.03	6.4e-05	inoculated tolerant line BR12 vs. inoculated susceptible line SUS3
NLP7-T1	Bv05_g11459_hany.t1	1.93	7.8e-06	inoculated tolerant line U2Bv vs. inoculated susceptible line U1Bv

NLP7-T2	Bv05_g11464_mdsr.t1	1.29	5.3e-04	inoculated tolerant line U2Bv vs. inoculated susceptible line U1Bv
NLP7-T1	Bv05_g11459_hany.t1	2.34	6.5e-27	all tolerant vs. all susceptible
NLP7-T2	Bv05_g11464_mdsr.t1	1.708	2.2e-13	all tolerant vs. all susceptible

3 Discussion

Phenotypic evaluation of the 406 F2:3 families segregating for BCN tolerance, MBS and
generation of highly continuous, well annotated genome sequences allowed characterization
of the NT locus as well as development of tightly linked molecular markers. This trait region
is located further north on chr5 when compared to the published BCN tolerance marker
SNP192 [6].

The identification of extremes in the BNC tolerance distribution and the continuous variation
of cyst numbers per F2 line demonstrated that the STR-NT population as well as the method
for BNC tolerance scoring was suitable for the MBS analysis. Variant detection and dAF
calculation were aided by the two new long read-based sugar beet genome sequence
assemblies of the susceptible genotype 2320Bv and the BNC tolerant line Strube U2Bv.

In case of monogenic traits, the difference in allele frequency at the causal gene locus should
ideally be 100% and thus have a dAF value of 1. This can be achieved for phenotypic data of
traits scored without confounding environmental and/or technical effects. An example is the
the *RED* locus for sugar beet hypocotyl color which is classified as either "green" or "red" [25].
The NT locus shows up in MBS as a single locus in the north of chr5, but the locus reflects a
major QTL that independently does not fully explain tolerance to BCN in Strube U2Bv.
Therefore, dAF values at the NT locus are expected to be lower than 1.

Although a candidate interval on chr5 was already visible in a preliminary analysis when using
the initial RefBeet sequence [10], the data quality was substantially improved in the current
study. In fact, the new assemblies 2320BvONT and U2BvONT resolve previously collapsed
or duplicated regions and offer continuous sequences for the region of interest on chr5.

The *B. vulgaris* repeat content was previously estimated to be 63% [26]. The new assembly 2320BvONT contains a comparable proportion of repeats, whereas RefBeet contains 42.3% of repeats [10]. This indicates that 2320BvONT contains a substantial amount of sequences not present in the RefBeet assembly. The U2BvONT assembly, which displays similarly high quality and represents the first full assembly of a BCN tolerant line, will serve as an additional resource for genome-informed sugar beet breeding.

The slightly higher number of genes predicted for the U2BvONT assembly can be explained by the ~20 Mbp larger assembly size. This additional sequence possibly originates from remaining heterozygosity in Strube U2Bv or may be partly explained by a slightly higher repeat content. The *ab initio* gene prediction for both genome sequences allowed the functional comparison of the delimited NT locus.

The genomic region identified by MBS was further restricted by designing genetic markers that do not show recombination with the phenotype on extreme F2 genotypes analyzed together with susceptible and tolerant standard lines (Additional file S2C). This interval spans regions of 0.7 Mbp and 0.76 Mbp in U2BvONT and 2320BvONT, respectively. Since the genome sequence is continuous and colinear in both assemblies at the NT locus, there is no evidence for missing additional sequences.

The NT locus shows generally a high synteny between the genome assemblies U2BvONT and 2320BvONT. Within this region, we identified a cluster of genes related to *AtNLP7*. Within the whole genome sequence, *BvNLP7* genes were only found at the NT locus on chr5. In the published *ab initio* structural annotation of EL10, two additional gene models (compared to the liftoff from the U2BvONT annotation, Figure 3) were predicted. However, both show similarity to histone H4 and are not considered as candidates.

Despite the synteny within the target region, four *BvNLP7* genes were identified in U2BvONT, but only three in 2320BvONT. The core amino acid residues of the NLP7 PB1 domain, which is relevant for regulatory interactions, are conserved in almost all of the *BvNLP7* genes. In the context of nitrate response and plant growth, mutants with substitutions of these core amino acid residues require a higher level of expression than wildtype NLP7 [23]. The fact that *nlp7* mutants are associated with increased lateral root density as well as impaired nitrate

assimilation leading to decreased amino acid formation [17] might at least partially explain the trade-off between high yield and resistance against nematodes.

To get insights into expression patterns of the *BvNLP7* genes, an infection assay was performed. RNA-Seq analysis was used to identify up- and downregulated genes in response to BCN infection (21 dpi). A time course experiment might reveal additional insights into the plants' defense mechanisms upon infection. However, the performed experiment enabled an in depth, transcriptomic characterization of our *BvNLP7* genes. All four U2BvONT *BvNLP7* genes are expressed. Expression is significantly higher in inoculated tolerant BR12 samples compared to inoculated susceptible SUS3 standard samples. The genes *NLP7-T1* and *NLP7-T2* are significantly differentially expressed between all samples of tolerant lines and all samples of susceptible lines.

Genes directly regulated by the AtNLP7 TF in *A. thaliana* root cells [16] include several genes encoding transporters, expansins and a WRKY TF (WRKY23) which are involved in nematode-induced syncytia formation. On chr5 of U2BvONT and 2320BvONT, 12 genes were annotated as putative WRKY TFs. Two (Bv05_g11386_pnrz and Bv05_g11531_wmfg) are located close to the NT locus at approximately 1 Mbp up- and downstream of the *BvNLP7* cluster, respectively. Two additional WRKY23 homologs (Bv05_g14901_ygdi and Bv05_g14902_sogd) were identified on chr5. However, none of these four WRKYs are differentially expressed in the comparisons of all tolerant vs. all susceptible samples or inoculated tolerant line BR12 vs. inoculated susceptible line SUS3.

In oilseed rape, BCN resistance was enhanced by gene pyramiding [27]. Homologs of the plant-defense genes addressed by Zhong *et al.*, namely *AtNPR1/AT1G64280*, *AtSGT1b/AT4G11260* and *AtRAR1/AT5G51700*, were identified in U2BvONT. *BvNPR1* was detected on chr8 with 58.6% aa identity. Homologs of the other two genes were found on chr3 with 57.7% (*BvSGT1b*) and 63.6% (*BvRAR1*) aa identity. In another recent study, BCN infection phenotypes were characterized in transgenic *A. thaliana* [28]. A *AtSNAP2/AT3G56190* homolog was detected on U2BvONT chr2 (74.9% identity via blastp), a *AtSHM(T)4/AT4G13930* homolog on chr3 (89.2% identity) and a *AtPR1/AT2G14610* homolog on chr9 (58.6% identity). In summary, none of the sugar beet homologs of these six genes which were described to be associated with BCN infection/resistance in these two studies, are located within or next to the NT locus. In addition, these genes are not

375 differentially expressed when comparing all tolerant vs. all susceptible samples. Therefore,
376 these genes are highly unlikely to be causal for BCN tolerance/resistance.

377 In summary, the presence of one more *BvNLP7* (additional) gene in Strube U2Bv as well as
378 the cumulative expression of those genes might explain the BCN tolerance of the genotype
379 Strube U2Bv. Therefore, the BCN tolerance might be based on a 'gene dosage' effect.

4   Conclusion

In this study, a trait locus associated with BCN tolerance was identified via mapping-by-
sequencing. Two newly generated long read-based genome sequences of the sugar beet
reference genotype 2320Bv and the tolerant line Strube U2Bv guided the characterization of
the NT locus. Four *BvNLP7* genes in U2BvONT are upregulated in tolerant lines as revealed
by an infection assay. These four genes, *NLP7-T1-4*, combined, or a subset thereof, might
convey tolerance against the cyst nematode *H. schachtii* which infects sugar beet and is a
serious problem due to yield loss. These results have positive implications for knowledge-
based breeding of elite genotypes.

5   Methods

5.1   Plant material and growth conditions

The breeding material is a large population of 406 lines descending from a single F1 plant
obtained by crossing a highly BCN susceptible line (Strube U1Bv) with a line (Strube U2BV)
highly tolerant to *H. schachtii*. Leaf samples were collected from all F2 plants and frozen at -
20 °C until usage. Ten selfed F3 plants per line were grown at a 16 hours light, 8 hours dark
cycle at 20-22 °C in the greenhouse, and 4,060 plants were used for evaluating nematode
tolerance. Single seedlings were grown in folding boxes that guarantee separation of the root
system and all plants were randomized in larger boxes together with tolerant and susceptible
H. schachtii breeding material used as checks. A dispenser was used to inoculate each plant
with 350 *H. schachtii* second stage juvenile larvae propagated in the lab of Strube Research.

Five weeks after inoculation, roots of each plant were washed, and cysts were collected and
counted under a stereoscope.

For RNA-Seq, different BCN tolerant and susceptible lines were used. The susceptible lines
were Strube U1Bv and SUS3, an internal standard line of Strube Research. The tolerant lines
were Strube U2Bv and the best performing F2 genotype BR12. All lines were germinated and
cultivated in the greenhouse for 11 weeks. Each line was represented by 40 plants. Half of
the plants were inoculated with nematodes as described above, the other half was left
untreated. All materials were collected 8 weeks after inoculation. The sampling of infected
plants was performed by collecting tissue and washing off the cysts, which were then counted
under a stereomicroscope. After gentle washing, the shoot was removed and the roots were
immediately frozen in liquid nitrogen until RNA extraction.

The phenotypic data were analyzed using a mixed model approach [29]. Boxes were treated
as random incomplete blocks and the genotypes as fixed. The counted number of cysts (n)
was transformed using a square root transformation ($\sqrt{n + 3/8}$) to meet the assumption of
normally distributed residuals required for mixed models. Adjusted means were obtained for
each line for QTL mapping. Adjusted single plant data were used to explore within-family
variation using the coefficient of variation (CV). A small CV indicates phenotypic homogeneity
among individual plants of a line, suggesting no segregation for the trait under study.

5.2 QTL detection

Together with the genetic linkage map, the adjusted means of the F2:3 families of the STR-
NT population were used for QTL mapping. Composite interval mapping (CIM) was employed
for QTL detection and a LOD threshold of 3.5 corresponding to an experiment wise type I
error rate of 0.05 was determined using 1000 permutation runs. All QTL computations were
performed with the software package PLABQTL [30] using an additive and dominant model
and a scan of 1 cM interval.

5.3 DNA extraction and MBS pool generation

428 Genomic DNA for MBS was extracted from young leaf tissue using the CTAB method [31].
429 The nine most susceptible genotypes were used for individual library preparations, whereas
430 the gDNA from the remaining seven genotypes were equimolarly pooled before library

preparation. For the 'tolerant' pool, the gDNA from the nine most tolerant lines was used for individual library preparations, whereas the remaining 11 gDNAs were equimolarly pooled. DNA for short read sequencing and PCR-based marker analysis was extracted from 8 leaf disks with 1 cm diameters using a CTAB-based protocol [31]. High molecular weight DNA for long read ONT sequencing was extracted with a modified CTAB-based protocol as previously described [32].

5.4   Short read sequencing for MBS

Each single gDNA or gDNA pool was fragmented by sonication using a Bioruptor (Fa. Diagenode). After cleaning the DNAs by AMPureXP Beads (Fa. Beckmann-Coulther), 200 ng of fragmented DNA was used for library preparation with the TruSeq Nano DNA library preparation kit (Fa. Illumina). End-repaired fragments were size selected by AMPureXP Beads to an average size of around 700 bp. After A-tailing and ligation of barcoded adaptors, fragments were enriched by 8 cycles of PCR. The final libraries were quantified using PicoGreen (Fa. Quant-iT) on a FLUOstar plate reader (Fa. BMG labtech) and quality checked by HS-Chip on a 2100 Bioanalyzer (Fa. Agilent Technologies). After pooling of all libraries, sequencing was performed on two 2x250 nt runs on a HiSeq1500 in rapid mode over two lanes using onboard-cluster generation. Processing and demultiplexing of raw data were performed by bcl2fastq. Additional file S2E summarizes the data submitted to ENA.

5.5   Short read mapping and variant calling from MBS data

The short read WGS data from the phenotypic pools, the parental lines of the mapping population, the F1 plant, and additional standard lines were used to identify small sequence variations within the population and against the susceptible reference genotype 2320Bv. BWA MEM v0.7.13 [33] was applied with the –m option to flag small alignments as secondary to align short reads to the U2BvONT reference sequence. Picard tools v2.5.0 (https://broadinstitute.github.io/picard/) and samtools v1.15.1 [34] were used to mark PCR duplicates, sort, and index BAM files. Mappings were filtered with samtools to remove spurious hits, low quality alignments, and reads that are not properly mapped in pairs (-q 30 -b -F 0x900 -f 0x2). GATK v3.8 [35] [36] was applied for the detection of small sequence variants as previously described [37]. Sequence variants were filtered to obtain a reduced set with high confidence. The following criteria were applied to select high confidence variants:

461 (1) variants are homozygous in the parent reads, (2) variants are contrasting between the
462 parents, and (3) variants are heterozygous in the F1 reads. Python scripts developed and
463 applied for filtering are available in the corresponding GitHub repository
464 (https://github.com/bpucker/beetresmabs).

5.6   Calculation of delta allele frequencies and interval detection

466 The calculation and analysis of dAFs is generally based on a previously described workflow
467 [38]. The methods are described in detail in Additional file S2F. Figure 3, which represents
468 the comparison of the NT locus, was mostly generated with gggenomes [39].

5.7   RNA Extraction, Library Preparation, and Sequencing

Plants of the genotypes Strube U2Bv, BR12, Strube U1Bv and SUS3 were grown in the
greenhouse and either infected or not infected with *H. schachtii* as described in 5.1. In total,
24 samples (Additional file S2G) including three biological replicates for each condition of the
infection experiment were ground separately under liquid nitrogen. Total RNA was extracted
from approx. 100 mg tissue using an RNA Isolation Kit (Sigma-Aldrich Spectrum™ Plant Total
RNA) according to suppliers' instructions. Quality control, determination of RIN numbers, and
estimation of the concentrations of RNA samples was done on a Bioanalyzer 2100 (Agilent)
using RNA Nano 6000 Chips. To construct sequencing libraries according to the Illumina
TruSeq RNA Sample Preparation v2 Guide, 500 ng total RNA per subsample were used.
Further steps, like enrichment of poly-A containing mRNA, cDNA synthesis, adapter ligation,
PCR enrichment, library quantification, and equimolar pooling, were performed according to
Theine *et al.*, 2021 [40]. Single end sequencing of 100 nt was performed on an Illumina
NextSeq500.

5.8   Genetic markers and linkage mapping

Sets of small variants detected by GATK v3.8 [35,36], on the basis of short read mappings of
Strube U1Bv, Strube U2Bv, and F1 reads to RefBeet-1.2 [10], were used to design KASPar
markers. Only homozygous single nucleotide variants contrasting between the parents and
with clear heterozygosity in the F1 were taken forward as clean variants. All variants
overlapping with existing markers were excluded. Up to 1000 variants per contig were
selected based on the quality of the variant call. A total of 50 bp upstream and downstream,

respectively, were checked for other variants based on a very lenient and unfiltered variant calling. Only marker candidates without any additional variants in these flanking sequences were taken forward. Further, only marker candidates with less than 65% GC content in 100 bp of flanking sequence were considered. Finally, marker candidates were preferentially selected on unplaced contigs of RefBeet-1.2 with an upper limit of four candidates per contig. The markers included within the final list were selected based on their position on each chromosome aiming to form a well-distributed marker subset.

The linkage map was constructed using KASPar markers and the package R/qtl [41]. The final linkage map comprises a set of 194 SNP markers including SNP192 that coalesced into nine linkage groups. Each group corresponded to one of the nine chromosomes in the haploid sugar beet genome. The average distance between loci was 3 cM except for two markers at the end of chr1 in poor linkage due to distortion. The average number of markers per chromosome was 21. For chr7 and chr9 the number of markers was below average and equal to 15 and 11, respectively.

5.9 ONT sequencing and ONT assembly

ONT long-read sequencing was performed on a GridION. The initial assembly was generated with Canu and further processed. The exact methodology is described in Additional file S2F, and Additional file S2E summarizes the data submitted to ENA.

5.10 Gene prediction and functional annotation

After repeat masking, hint-based gene prediction was performed mainly with BRAKER2. All predicted genes were subsequently functionally annotated. The methodology for structural and functional annotation with BRAKER2 is described in detail in Additional file S2F.

5.11 Differential gene expression analysis

The generated RNA-Seq reads were mapped to the U2BvONT genome sequence using STAR v2.7.6a (Linux_x86_64_static/STARlong --runThreadN 8 --genomeDir /dir/ --limitBAMsortRAM 32000000000 --outBAMsortingThreadN 4 --outSAMtype BAM SortedByCoordinate --outFileNamePrefix /01 --readFilesCommand gunzip -c --readFilesIn 01.fastq.gz) [42]. FeatureCounts v2.0.0 [43] was used to quantify annotated genes in the U2BvONT GFF file (-T 8 -t gene -a annotation.gff -o readcounts_allbams.txt *.bam).

Downstream analysis was performed using the R package DESeq2 v1.26.0 [44]. A variance
stabilizing transformation was conducted. A principal component analysis (PCA) for all
samples of the infection assay (Additional file S2D) was generated using prcomp (stats-
package v3.6.3 [45]. and ggplot2 v3.3.5 [46]. Significantly differentially expressed genes
between i) all tolerant vs. all susceptible samples and ii) between inoculated SUS3 and
inoculated BR12 samples, were extracted based on an adjusted p-value < 0.05.

5.12 Rearrangement and synteny analyses

Synteny analyses between 2320BvONT and U2BvONT as well as 2320BvONT and RefBeet
were performed using JCVI MCscan v1.2.4 [47]. Unanchored mRNAs were compared for
unique functions using BLAST v2.13.0 [48] and InterProScan v5.52 [49]. Structural
rearrangements were identified with SyRI v1.4 [50].

5.13 Comparison of *BvNLP7* genes

A list of direct targets of the AtNLP7 TF has been published recently [16]. To assess a possible
role of NPL7 gene(s) in BCN tolerance, these targets were functionally investigated by
overrepresentation analysis of GO terms using PANTHER v17.0 [51]. Additionally, the
sequences were directly compared via MAFFT v7.487 [52] alignments and manual
inspection.

6   Acknowledgments

We thank the CeBiTec Bioinformatic Resource Facility team as well as the de.NBI team for
great technical support. The authors also wish to thank the members of Strube Research for
their technical support in phenotyping and genotyping large populations. We thank Prof.
Piepho from the University of Hohenheim for the constructive discussion on the liner mixed
model. We thank Britta Schulz from KWS-SAAT SE for providing seeds of the reference
genotype KWS2320. Thanks to Julia Zimmer and Marvin Hildebrandt for variant validation.

7   Author Contributions

EO, AM, AS, DH and BW are responsible for conceptualization. KS, BP, EO, EA, AS, DH and
BW developed the methodology. Validation, formal analysis and investigation were performed
by KS, BP, EO, LS, PV, EA and DH. KS, BP, EO, LS, PV, EA and DH conducted data curation.
KS, BP, EO, and DH wrote the original draft. KS, BP, EO, EA, AM, AS, PV, BW and DH
reviewed and edited the manuscript. Visualization of the data was performed by KS and BP.
Supervision and project administration was carried out by EO, AS, DH and BW. EO, AM, AS,
and BW were responsible for funding acquisition.

8   Data Availability Statement

A summary of the availability of the newly generated sequencing data, assemblies and
annotations is provided in Additional file S2E covering ENA projects PRJEB56338,
PRJEB37059, PRJEB36905, PRJEB58360 as well as DOIs for the genome sequence
annotation files (.gff). Additional RNA-Seq datasets produced for hint generation and gene
prediction are available via ENA projects PRJEB58621 and PRJEB62793, these and further
already published hint data are listed in Additional file S2H.

9   Conflict of Interest

563 The authors declare that the research was conducted in the absence of any commercial or
564 financial relationships that could be construed as a potential conflict of interest.

10   Funding

This project was funded by the Federal Ministry of Education and Research of Germany
(BMBF) in the frame of KMU-innovativ-16 under the grant numbers 031B0081A and
031B0081B (BeetRes-MaBS). This work was supported by the BMBF-funded de.NBI Cloud
within the German Network for Bioinformatics Infrastructure (grant numbers 031A532B,
031A533A, 031A533B, 031A534A, 031A535A, 031A537A, 031A537B, 031A537C,
031A537D, 031A538A). KS is funded by Bielefeld University through the Graduate School

DILS (Digital Infrastructure for the Life Sciences). We acknowledge support for the publication costs by the Deutsche Forschungsgemeinschaft and the Open Access Publication Fund of Bielefeld University.

11   Supplementary Material

Additional file S1: S1A: Genomic positions of the new set of 187 markers designed based on data for Strube U2Bv. In the first column, the name of the marker is shown, followed by the chromosome and genomic position for each assembly (2320BvONT v1.0 and U2BvONT v1.0). **S1B:** Overrepresentation analysis (GO terms) of the AtNLP7 targets identified by Alvarez et al., 2020. **S1C:** Percent identity matrices for the BvNLP7 genes. **S1D:** List of all significantly differentially expressed genes between all samples of the tolerant and all samples of the susceptible lines.

Additional file S2: S2A: Structural rearrangements between 2320BvONT and U2BvONT as identified with SyRI. **S2B:** Delta allele frequency plots for 10 SNP windows of all nine U2BvONT pseudochromosomes. **S2C:** Marker information/primer sequences used for delimitation of the NT locus and graphical genotyping with flanking and co-segregating markers. **S2D:** Principal component analysis for all samples of the RNA-Seq infection experiment. **S2E:** Availability and composition of datasets generated. **S2F:** Detailed methods for i) calculation of the delta allele frequencies and interval detection, ii) ONT sequencing, iii) ONT assembly, and iv) gene prediction and functional annotation. **S2G:** Overview of the RNA-Seq samples in the infection experiment. **S2H:** List of all RNA-Seq datasets incorporated as hints for the gene prediction including two newly submitted datasets. **S2I:** Dot plot heatmap of the closed gap region in the initial Strube U2Bv assembly.

Additional file S3: Multiple sequence alignment of coding sequences of all *BvNLP7* genes.

Additional file S4: Multiple sequence alignment of *A. thaliana* NLP7 and U2BvONT and 2320BvONT rycp aa sequences.

Additional file S5: Multiple sequence alignment of aa sequences of the *BvNLP7* genes (without rycp).

12 References

1. Fischer, H.E. Origin of the "Weisse Schlesische Ruebe" (white Silesian beet) and resynthesis of sugar beet. *Euphytica* **41**, 75-80 https://dx.doi.org/10.1007/BF00022414 (1989).

2. Biancardi, E., McGrath, J.M., Panella, L.W., Lewellen, R.T. & Stevanato, P. Sugar Beet in Root and Tuber Crops Vol. 7 (ed Bradshaw, J.E.) 173-219 Ch. Chapter 6 http://dx.doi.org/10.1007/978-0-387-92765-7_6 (Springer, 2010).

3. Richardson, K.L. Registration of sugar beet mapping populations CN239, CN240, and CN241 segregating for resistance to Heterodera schachtii from sea beet. *Journal of Plant Registrations* **16**, 459-464 https://dx.doi.org/10.1002/plr2.20152 (2022).

4. Reuther, M., Lang, C. & Grundler, F.M.W. Nematode-tolerant sugar beet varieties – resistant or susceptible to the Beet Cyst Nematode Heterodera schachtii? *Sugar Industry* **142**, 277-284 https://dx.doi.org/10.36961/si18397 (2017).

5. Holtmann, B., Kleine, M. & Grundler, F.M.W. Ultrastructure and anatomy of nematode-induced syncytia in roots of susceptible and resistant sugar beet. *Protoplasma* **211**, 39-50 https://dx.doi.org/10.1007/BF01279898 (2000).

6. Stevanato, P., Trebbi, D., Panella, L., Richardson, K., Broccanello, C., Pakish, L., Fenwick, A.L. & Saccomani, M. Identification and Validation of a SNP Marker Linked to the Gene HsBvm-1 for Nematode Resistance in Sugar Beet. *Plant Molecular Biology Reporter* **33**, 474-479 https://dx.doi.org/10.1007/s11105-014-0763-8 (2015).

7. Cai, D., Kleine, M., Kifle, S., Harloff, H.J., Sandal, N.N., Marcker, K.A., Klein Lankhorst, R.M., Salentijn, E.M.J., Lange, W., Stiekema, W.J., Wyss, U., Grundler, F.M.W. & Jung, C. Positional cloning of a gene for nematode resistance in sugar beet. *Science* **275**, 832-834 https://dx.doi.org/10.1126/science.275.5301.832 (1997).

8. Biancardi, E., Lewellen, R.T., Frese, L., Ford-Lloyd, B., de Biaggi, M., Hautekeete, N., van Dijk, H., Touzet, P., Bartsch, D., Panella, L.W., Stevanato, P., Pavli, O., Skaracis, G. & McGrath, J.M. Beta maritima: The Origin of Beets (Second Edition) http://dx.doi.org/10.1007/978-3-030-28748-1 2020).

9. McGrath, J.M., Funk, A., Galewski, P., Ou, S., Townsend, B., Davenport, K., Daligault, H., Johnson, S., Lee, J., Hastie, A., Darracq, A., Willems, G., Barnes, S., Liachko, I., Sullivan, S., Koren, S., Phillippy, A., Wang, J., Liu, T., Pulman, J., Childs, K., Shu, S., Yocum, A., Fermin, D., Mutasa-Gottgens, E., Stevanato, P., Taguchi, K., Naegele, R. & Dorn, K.M. A contiguous de novo genome assembly of sugar beet EL10 (Beta vulgaris L.). *DNA Research* **30**, https://dx.doi.org/10.1093/dnares/dsac033 (2023).

10. Dohm, J.C., Minoche, A.E., Holtgrawe, D., Capella-Gutierrez, S., Zakrzewski, F., Tafer, H., Rupp, O., Sorensen, T.R., Stracke, R., Reinhardt, R., Goesmann, A., Kraft, T., Schulz, B., Stadler, P.F., Schmidt, T., Gabaldon, T., Lehrach, H., Weisshaar, B. & Himmelbauer, H. The genome of the recently domesticated crop plant sugar beet (Beta vulgaris). *Nature* **505**, 546-9 https://dx.doi.org/10.1038/nature12817 (2014).

11. Holtgräwe, D., Sörensen, T.R., Viehöver, P., Schneider, J., Schulz, B., Borchardt, D., Kraft, T., Himmelbauer, H. & Weisshaar, B. Reliable in silico identification of sequence polymorphisms and their application for extending the genetic map of sugar beet (Beta vulgaris). *PLoS ONE* **9**, e110113 https://dx.doi.org/10.1371/journal.pone.0110113 (2014).

12. Rodriguez Del Rio, A., Minoche, A.E., Zwickl, N.F., Friedrich, A., Liedtke, S., Schmidt, T., Himmelbauer, H. & Dohm, J.C. Genomes of the wild beets Beta patula and Beta vulgaris ssp. maritima. *The Plant Journal* **99**, 1242-1253 https://dx.doi.org/10.1111/tpj.14413 (2019).

13. Schauser, L., Roussis, A., Stiller, J. & Stougaard, J. A plant regulator controlling development of symbiotic root nodules. *Nature* **402**, 191-5. https://dx.doi.org/10.1038/46058 (1999).

14. Schauser, L., Wieloch, W. & Stougaard, J. Evolution of NIN-like proteins in Arabidopsis, rice, and Lotus japonicus. *Journal of Molecular Evolution* **60**, 229-37 https://dx.doi.org/10.1007/s00239-004-0144-2 (2005).

15. Konishi, M. & Yanagisawa, S. Arabidopsis NIN-like transcription factors have a central role in nitrate signalling. *Nature Communications* **4**, 1617 https://dx.doi.org/10.1038/ncomms2621 (2013).

16. Alvarez, J.M., Schinke, A.L., Brooks, M.D., Pasquino, A., Leonelli, L., Varala, K., Safi, A., Krouk, G., Krapp, A. & Coruzzi, G.M. Transient genome-wide interactions of the master transcription factor NLP7 initiate a rapid nitrogen-response cascade. *Nature Communications* **11**, 1157 https://dx.doi.org/10.1038/s41467-020-14979-6 (2020).

17. Castaings, L., Camargo, A., Pocholle, D., Gaudon, V., Texier, Y., Boutet-Mercey, S., Taconnat, L., Renou, J.P., Daniel-Vedele, F., Fernandez, E., Meyer, C. & Krapp, A. The nodule inception-like protein 7 modulates nitrate sensing and metabolism in Arabidopsis. *The Plant Journal* **57**, 426-35 https://dx.doi.org/10.1111/j.1365-313X.2008.03695.x (2009).

18. Marchive, C., Roudier, F., Castaings, L., Brehaut, V., Blondet, E., Colot, V., Meyer, C. & Krapp, A. Nuclear retention of the transcription factor NLP7 orchestrates the early response to nitrate in plants. *Nature Communications* **4**, 1713 https://dx.doi.org/10.1038/ncomms2650 (2013).

19. Zhao, L., Zhang, W., Yang, Y., Li, Z., Li, N., Qi, S., Crawford, N.M. & Wang, Y. The Arabidopsis NLP7 gene regulates nitrate signaling via NRT1.1-dependent pathway in the presence of ammonium. *Scientific Reports* **8**, 1487 https://dx.doi.org/10.1038/s41598-018-20038-4 (2018).

20. Guan, P., Ripoll, J.J., Wang, R., Vuong, L., Bailey-Steinitz, L.J., Ye, D. & Crawford, N.M. Interacting TCP and NLP transcription factors control plant responses to nitrate availability. *Proceedings of the National Academy of Sciences of the United States of America* **114**, 2419-2424 https://dx.doi.org/10.1073/pnas.1615676114 (2017).

21. Lu, S., Wang, J., Chitsaz, F., Derbyshire, M.K., Geer, R.C., Gonzales, N.R., Gwadz, M.,
Hurwitz, D.I., Marchler, G.H., Song, J.S., Thanki, N., Yamashita, R.A., Yang, M., Zhang,
D., Zheng, C., Lanczycki, C.J. & Marchler-Bauer, A. CDD/SPARCLE: the conserved
domain database in 2020. *Nucleic Acids Research* **48**, D265-D268
https://dx.doi.org/10.1093/nar/gkz991 (2020).

22. Sumimoto, H., Kamakura, S. & Ito, T. Structure and function of the PB1 domain, a
protein interaction module conserved in animals, fungi, amoebas, and plants.
SCIENCE'S STKE Science Signaling **2007**, re6
https://dx.doi.org/10.1126/stke.4012007re6 (2007).

23. Konishi, M. & Yanagisawa, S. The role of protein-protein interactions mediated by the
PB1 domain of NLP transcription factors in nitrate-inducible gene expression. *BMC Plant*
Biology **19**, 90 https://dx.doi.org/10.1186/s12870-019-1692-3 (2019).

24. Liu, K.H., Niu, Y., Konishi, M., Wu, Y., Du, H., Sun Chung, H., Li, L., Boudsocq, M.,
McCormack, M., Maekawa, S., Ishida, T., Zhang, C., Shokat, K., Yanagisawa, S. &
Sheen, J. Discovery of nitrate-CPK-NLP signalling in central nutrient-growth networks.
Nature **545**, 311-316 https://dx.doi.org/10.1038/nature22077 (2017).

25. Ries, D., Holtgräwe, D., Viehöver, P. & Weisshaar, B. Rapid gene identification in sugar
beet using deep sequencing of DNA from phenotypic pools selected from breeding
panels. *BMC Genomics* **17**, 236 https://dx.doi.org/10.1186/s12864-016-2566-9 (2016).

26. Flavell, R.B., Bennett, M.D., Smith, J.B. & Smith, D.B. Genome size and the proportion
of repeated nucleotide sequence DNA in plants. *Biochemical Genetics* **12**, 257-69
https://dx.doi.org/10.1007/BF00485947 (1974).

27. Zhong, X., Zhou, Q., Cui, N., Cai, D. & Tang, G. BvcZR3 and BvHs1(pro-1) Genes
Pyramiding Enhanced Beet Cyst Nematode (Heterodera schachtii Schm.) Resistance
in Oilseed Rape (Brassica napus L.). *International Journal of Molecular Sciences* **20**,
https://dx.doi.org/10.3390/ijms20071740 (2019).

28. Zhao, J., Duan, Y., Kong, L., Huang, W., Peng, D. & Liu, S. Opposite Beet Cyst
Nematode Infection Phenotypes of Transgenic Arabidopsis Between Overexpressing
GmSNAP18 and AtSNAP2 and Between Overexpressing GmSHMT08 and AtSHMT4.
Phytopathology **112**, 2383-2390 https://dx.doi.org/10.1094/PHYTO-01-22-0011-R
(2022).

29. Piepho, H.P., Buchse, A. & Emrich, K. A Hitchhiker's Guide to Mixed Models for
Randomized Experiments. *Journal of Agronomy and Crop Science* **189**, 310-322
https://dx.doi.org/10.1046/j.1439-037X.2003.00049.x (2003).

30. Utz, H.F. & Melchinger, A.E. PLABQTL: a program for composite interval mapping of
QTL. *Journal of Agricultural Genomics* **2**, 1-6 (1996).

31. Rosso, M.G., Li, Y., Strizhov, N., Reiss, B., Dekker, K. & Weisshaar, B. An *Arabidopsis*
thaliana T-DNA mutagenised population (GABI-Kat) for flanking sequence tag based
reverse genetics. *Plant Molecular Biology* **53**, 247-259
https://dx.doi.org/10.1023/B:PLAN.0000009297.37235.4a (2003).

32. Siadjeu, C., Pucker, B., Viehöver, P., Albach, D.C. & Weisshaar, B. High Contiguity De
Novo Genome Sequence Assembly of Trifoliate Yam (Dioscorea dumetorum) Using
Long Read Sequencing. *Genes* **11**, E274 https://dx.doi.org/10.3390/genes11030274
(2020).

33. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv **https://arxiv.org/abs/1303.3997**, posted 2013-05-26
https://dx.doi.org/10.48550/arXiv.1303.3997 (2013).

34. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G. & Durbin, R. The Sequence Alignment/Map format and SAMtools.
Bioinformatics **25**, 2078-2079 https://dx.doi.org/10.1093/bioinformatics/btp352 (2009).

35. McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A.,
Garimella, K., Altshuler, D., Gabriel, S., Daly, M. & DePristo, M.A. The Genome Analysis
Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.
Genome Research **20**, 1297-1303 https://dx.doi.org/10.1101/gr.107524.110 (2010).

36. Van der Auwera, G.A., Carneiro, M.O., Hartl, C., Poplin, R., Del Angel, G., Levy-
Moonshine, A., Jordan, T., Shakir, K., Roazen, D., Thibault, J., Banks, E., Garimella,
K.V., Altshuler, D., Gabriel, S. & DePristo, M.A. From FastQ data to high confidence
variant calls: the Genome Analysis Toolkit best practices pipeline. *Current Protocols in
Bioinformatics* **11**, 1110 https://dx.doi.org/10.1002/0471250953.bi1110s43 (2013).

37. Baasner, J.S., Howard, D. & Pucker, B. Influence of neighboring small sequence
variants on functional impact prediction. *bioRxiv* posted 2019-06-13
https://dx.doi.org/10.1101/596718 (2019).

38. Schilbert, H.M., Pucker, B., Ries, D., Viehover, P., Micic, Z., Dreyer, F., Beckmann, K.,
Wittkop, B., Weisshaar, B. & Holtgrawe, D. Mapping-by-Sequencing Reveals Genomic
Regions Associated with Seed Quality Parameters in Brassica napus. *Genes* **13**,
https://dx.doi.org/10.3390/genes13071131 (2022).

39. Hackl, T. & Ankenbrand, M.J. gggenomes: A Grammar of Graphics for Comparative
Genomics. R package version 0.9.5.9000 edn (2022).

40. Theine, J., Holtgrawe, D., Herzog, K., Schwander, F., Kicherer, A., Hausmann, L.,
Viehover, P., Topfer, R. & Weisshaar, B. Transcriptomic analysis of temporal shifts in
berry development between two grapevine cultivars of the Pinot family reveals potential
genes controlling ripening time. *BMC Plant Biology* **21**, 327
https://dx.doi.org/10.1186/s12870-021-03110-6 (2021).

41. Arends, D., Prins, P., Jansen, R.C. & Broman, K.W. R/qtl: high-throughput multiple QTL
mapping. *Bioinformatics* **26**, 2990-2 https://dx.doi.org/10.1093/bioinformatics/btq565
(2010).

42. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M. & Gingeras, T.R. STAR: ultrafast universal RNA-seq aligner.
Bioinformatics **29**, 15-21 https://dx.doi.org/10.1093/bioinformatics/bts635 (2013).

43. Liao, Y., Smyth, G.K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930
https://dx.doi.org/10.1093/bioinformatics/btt656 (2014).

44. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology* **15**, 550
https://dx.doi.org/10.1186/s13059-014-0550-8 (2014).

45. R Core Team. R: A language and environment for statistical computing. (R Foundation
for Statistical Computing, Vienna, Austria, 2018).

46.   Wickham, H. ggplot2: Elegant Graphics for Data Analysis (Springer, 2016).

47.   Tang, H., Bowers, J.E., Wang, X., Ming, R., Alam, M. & Paterson, A.H. Synteny and
772   collinearity in plant genomes. *Science* **320**, 486-8
773   https://dx.doi.org/10.1126/science.1153917 (2008).

774   48.   Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. Basic local alignment
775   search tool. *Journal of Molecular Biology* **215**, 403-410
776   https://dx.doi.org/10.1016/S0022-2836(05)80360-2 (1990).

777   49.   Quevillon, E., Silventoinen, V., Pillai, S., Harte, N., Mulder, N., Apweiler, R. & Lopez, R.
778   InterProScan: protein domains identifier. *Nucleic Acids Research* **33**, W116-20
779   https://dx.doi.org/10.1093/nar/gki442 (2005).

780   50.   Goel, M., Sun, H., Jiao, W.B. & Schneeberger, K. SyRI: finding genomic rearrangements
781   and local sequence differences from whole-genome assemblies. *Genome Biol* **20**, 277
782   https://dx.doi.org/10.1186/s13059-019-1911-0 (2019).

783   51.   Mi, H., Muruganujan, A., Ebert, D., Huang, X. & Thomas, P.D. PANTHER version 14:
784   more genomes, a new PANTHER GO-slim and improvements in enrichment analysis
785   tools. *Nucleic Acids Research* **47**, D419-D426 https://dx.doi.org/10.1093/nar/gky1038
786   (2019).

787   52.   Katoh, K. & Standley, D.M. MAFFT multiple sequence alignment software version 7:
788   improvements in performance and usability. *Molecular Biology and Evolution* **30**, 772-
789   780 https://dx.doi.org/10.1093/molbev/mst010 (2013).