TITLE: PEO: Plant Expression Omnibus - a comparative transcriptomic database for 103

Archaeplastida

AUTHORS: Eugene Koh[1], William Goh[1], Irene Julca[1], Erielle Villanueva[1], Marek Mutwil[1*]

[1]School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive,

Singapore, 637551, Singapore

*Corresponding author:

Marek Mutwil

School of Biological Sciences,

Nanyang Technological University,60 Nanyang Drive,

637551, Singapore,

Singapore

Email: mutwil@ntu.edu.sg

Abstract

The Plant Expression Omnibus (PEO) is a web application that provides biologists with

access to gene expression insights across over 100 plant species, ~60,000 manually

annotated RNA-seq samples, and more than four million genes. The tool allows users to

explore the expression patterns of genes across different organs, identify organ-specific

genes, and discover top co-expressed genes for any gene of interest. PEO also provides

functional annotations for each gene, allowing for the identification of genetic modules

and pathways. PEO is designed to facilitate comparative kingdom-wide gene expression

analysis and provide a valuable resource for plant biology research. We provide two case

studies to demonstrate the utility of PEO in identifying candidate genes in pollen coat

biosynthesis and investigating the biosynthetic pathway components of capsaicin in

Capsicum annuum. The database is freely available at https://expression.plant.tools/.

Introduction

The earliest members of the plant kingdom evolved circa one billion years ago from photosynthetic algae. The archetypical characteristics we associate with plants, such as seeds, roots, leaves, flowers, and wood, were evolutionary innovations that allowed plants to colonize diverse habitats successfully. Today, plants form an indispensable component of life on Earth as the primary food producer in the terrestrial ecosystem. Plants being mostly sessile, have also had to evolve means of adapting quickly to their local environments. These adaptations came in the form of complex regulatory and metabolic pathways, which we are now only beginning to unravel.

Studying plant biology and genetics will allow us to improve desirable traits and engineer crops that provide higher yield and/or are more tolerant to various environmental stresses (Niazian, 2019; Nowicka et al., 2018). The use of plants as a source of various medicinal components is also exemplified in the extraction of the analgesic salicins from willow bark or the antimalarial artemisinin from *Artemisia annua* (Desborough and Keeling, 2017; Krishna et al., 2008). The elucidation of biosynthetic pathways of these medicinal compounds could then be harnessed for large-scale production and modification in recombinant systems (Cravens et al., 2019; Paddon et al., 2013; Sabzehzari et al., 2020). Currently, the study of plant gene function requires the characterization of individual genes and their interactors and regulators, which are typically limited to a handful of model plants. Consequently, bioinformatic approaches that integrate genomic, transcriptomic, metabolomic, and phenotypic data have been developed to accelerate the process of gene function prediction (Hansen et al., 2018; Radivojac et al., 2013; Rhee and Mutwil, 2014).

Studying gene co-expression can effectively address this, given the "guilt-by-association" principle, where genes with similar expression are likely to have similar functions (Rhee and Mutwil, 2014). Co-expression analysis allows one to transfer functional annotation to unannotated genes, predict undiscovered members of metabolic pathways, and identify regulator-target relationships (Rao and Dixon, 2019). Plant organs have signature gene expression patterns and similar profiles between organs are correlated to the similarity in function of the organs (Ma et al., 2005). Moreover, many genes exhibiting organ-specific expression are found to be conserved across land

plants (Julca et al., 2021). Consequently, a gene's expression profile across organs can

provide clues to a gene's function. For example, two genes that show similar gene

expression profiles across different organs and tissues are likely to be part of the same

protein complex (Persson et al., 2005) or biosynthetic pathway (Delli-Ponti et al., 2021)

or to be targeted to the same subcellular compartment (Ryngajllo et al., 2011).

Over the last decade, sequencing cost has been falling faster than Moore's law

(Wetterstrand, 2013), giving rise to immense volumes of plant genomic and transcriptomic

data. Many online databases with diverse functionalities and plant species have emerged

to provide analytical and visualization tools for the scientific community (Lim et al., 2022;

Liu et al., 2023; Robinson et al., 2018; Yu et al., 2022). However, no existing database

has been designed to cover well-annotated gene expression data. We believe that the

full value of RNA-seq repositories is only realized when insights are accessible to

biologists working on the genes, including non-model species. We created a user-friendly

tool to help non-programmer biologists extract insights about their gene of interest from

big data (Tan et al., 2020). A web application to access gene expression knowledge on

any gene would be needed to provide an effortless and instantaneous experience for

biologists, which requires large volumes of processed data. While the data is publicly

available, the interfaces have not been specifically designed for bulk data acquisition for

whole kingdoms. We addressed this by designing a parallel-processing pipeline to

automatically discover, download and process RNA-seq data for the whole plant kingdom

in less than a month (Goh and Mutwil, 2021), making the provision and maintenance of a

kingdom-wide resource feasible.

Here, we developed the Plant Gene Expression Omnibus (PEO,

https://expression.plant.tools), a web application to provide an interface for biologists to

access gene expression insights across >100 plant species. Specifically, the tool

addresses how a gene is expressed across different organs, whether it displays organ-

specific expression patterns and identifies top organ-specific genes for each organ. The

tool also automatically provides a list of the highest correlated genes based on the

Pearson Correlation Coefficient for any gene of interest, negating the need for separate

analysis. In addition, Pfam and Mapman annotations have also been linked to each gene,

allowing for functional searches to identify genetic modules or pathways. The software

design considers scalability, with the ability to include more species, other expression

conditions, and other gene annotations. The project lays the architectural foundations for

a kingdom-wide gene expression analysis resource.

Methods

Data source

Supplemental Table S1 details the sources of raw files used and contributors. These were

then reorganized into the upload format described in Supplemental Table S2, using

custom Python scripts. A total of 103 species were included. The complete list is available

in Supplemental Dataset S1. Gene expression matrices were obtained from a previous

study from public databases via our pipeline using recommended thresholds of $\log_{10}$-

normalized number of processed reads and percentage of reads pseudo-aligned to the

reference CDS (Goh and Mutwil, 2021), which removed samples with low number and

percentage of mapping reads. Gene expression was quantified as transcript per million

(TPM) via pseudoalignment to reference coding sequences (CDS) of each species, using

kallisto (Bray et al., 2016). Organ annotations were done by manually mapping samples

to Plant Ontology (PO) terms, which are precise and consistent references to plant

anatomical entities across species (Walls et al., 2019). The expression data and

annotations are available on 10.6084/m9.figshare.24257125.

Genes were functionally annotated with Mapman annotations via Mercator

(Schwacke et al., 2019) for four representative species and were assigned to the

remaining species via sequence similarity score from diamond (Buchfink et al., 2015).

Mapman was chosen over Gene Ontology (GO) as it is specific to plant pathways and

processes (Klie and Nikoloski, 2012). Genes were also annotated with protein families

(Pfam) (Mistry et al., 2021). Gene identifiers were standardized to the versions in TPM

matrices, and alternative isoforms were excluded. Co-expression analysis was performed

using Pearson Correlation Coefficient (PCC).

Database schema and software architecture

The schema defines how data is modeled and stored in our document-oriented

database and can be described with an entity relational diagram (ERD) (Supplemental

Figure S1). The chosen database technology is MongoDB, a document-oriented
database allowing flexible data modeling to optimize our specific read patterns. This is
more suitable than relational databases that enforce data schema normalization, which
would incur expensive join operations on every read. The database is hosted serverless
on MongoDB Atlas, on AWS.
129

Figure 1: RNA sequencing contents of the Plant Expression Omnibus (PEO) database. A) The heatmap shows the relative abundance of RNA-seq data for the different species (columns) and organs, tissues, and cell types (rows). The numbers in paratheses in rows indicate the number of species that have data for a given plant structure. The cell opacity indicates the relative abundance of RNA-seq samples across

organs for each species. B) The treemap showing the relative number of RNA-seq
samples for the different species. The areas of the rectangles correspond to the number
of samples, which is given in parentheses.

Results and discussion

Querying the Plant Expression Omnibus (PEO)

PEO is an interlinked database that includes 103 species, 4.6 million genes, 5,760
Mapman terms, 7,566 Pfam terms and 59,856 quality-controlled samples, of which
41,592 are annotated with 78 PO terms. To our knowledge, no species-wide gene
expression database exists for many of these 103 species. The distribution of data
illustrating species composition and organ specificity are presented as a treemap and
heatmap, respectively (Figure 1). Not surprisingly, leaves were the most sampled plant
organs (97 species contain leaf samples), followed by roots (81), flowers (70), stems (67)
and seeds (49)(Figure 1A). The less represented samples represent structures not found
in all flowering plants (strobilus, tuber, protonema, spore) or samples that are difficult to
collect (trichome, zygote, egg cell). *Arabidopsis thaliana* had the highest number of RNA-
seq samples (Ara.th, 72,732), followed by maize (35,309), rice (25,090), wheat (14,137)
and other crops (Figure 1B), with the next model organism *Brachypodium distachyon* on
the 17[th] place with 3,004 samples. The annotation depth of Mapman, Pfam and Plant
Ontology annotations are also provided (Supplemental Figure S2). The data are sorted
by several modes, such as species type, organ specificity, as well as Mapman and Pfam
annotation. Such an arrangement allows the user to query for gene expression in specific
organs of interest or look for orthologous genes in non-model species. Sequence-based
BLAST and annotation-based Pfam searches can be used to identify homologous and
orthologous genes of interest across the database. Together with Mapman annotation
and organ specificity predictions, putative functions can be ascribed to gene orthologues.
Finally, PEO also provides a list of top co-expressed genes for any gene of interest for all
species present in the database, allowing for the discovery of specific metabolic or

developmental　　　pathways　　　that　　　are　　　spatiotemporally　　　regulated.

Figure 2: Frontend of Gene Expression Omnibus. A) Genes of interest are searchable

via GeneID, protein sequence, Pfam domains, and species as query entries. Browsing of

various databases is available in the above tabs. B) Example search by organ type

(Pollen). C) The output displays a list of genes known to be present in a specific organ of

interest, searchable by species.

The home page of PEO consists of several search modes. Queries can be based
on specific GeneID or protein sequences. It is also possible to browse the database via
the 'Species', 'Organs', 'Pfam', or 'Mapman' annotations (Figure 2). For example, we can
select the 'Organs' tab, which provides a list of Plant Ontology (PO) terms. By selecting
'Pollen' as our organ of interest, we arrive at a page displaying a set of transcripts sorted
by SPM. This specificity measurement ranges between zero and one, where one
indicates the gene is exclusively expressed in the tissue (Julca et al., 2021). The page
also provides information on the Median and Mean Transcripts per Million (TPM) values
used to calculate SPM value and the number of samples the calculations are based on.
Mapman annotations are also listed for individual transcripts to provide information on
putative gene functions. This database can also be searched by species to obtain the
pollen-specific genes in other species present in the PEO.

As another example of a query strategy, we also show an example search using
the protein sequence of the Arabidopsis gene ABORTED MICROSPORES
(*AT2G16910*), a transcription factor known to be associated with the regulation of pollen
formation (Xu et al., 2014). The search output provides a list of genes sorted by %identity,
with further information on E-values, bit scores, number of mismatches, and gap
openings. The gene list includes matched genes from the other species present in PEO,
providing a means to discover gene orthologues from less common organisms
(Supplemental Figure S3).

Clicking on a gene of interest in the list (AMS, *AT2G16910*), brings us to a gene
page containing relevant information on this gene (Figure 3). The top of the page provides
information about the GeneID, Species name, and TaxonID. The next section displays
the putative organ specificity of the gene of interest based on SPM calculations. Details
on the values that the SPM calculation is based on is present in the grey subtext below
the SPM value. The top organs are reported with mean TPM, standard deviation, and
median TPM, showing the magnitude of expression. Since genes with low TPM values
can appear to be specifically expressed due to noisy expression, the mean TPM values
can be used to flag such genes (Proost and Mutwil, 2018). The sample size is also
reported. Together, these support user interpretation of whether the gene displays organ-
specific expression. The data can be visualized in a bar chart or box/scatterplot format.

Figure 3: Example gene page of the gene of interest (*AT2G16910*). A) The gene page provides organ specificity of the gene predicted by the SPM score. B) It also displays relative transcript expression in various plant organs/tissues as a box plot or bar chart. Moving the mouse pointer over the points in the box plot shows the corresponding experiment ID. C) Mapman and Pfam annotations for the gene of interest are displayed

to provide information on gene function. D) A list of top co-expressed genes sorted by
PCC value.

The boxplots depict quartiles and whiskers (hover to reveal values), indicating the skew
of data. PEO allows for automatic rescaling to exclude the outliers for the y-axis (Figure
3). To see which experiments the data stems from, the user can move the mouse pointer
over individual data points, which will reveal the run IDs. Manual zooming and panning is
also available by the control bar at the right of each chart. The charts are available to
download in .png or .svg image formats, with the raw data for the annotated samples also
available for download in tsv format. Associated Pfam and Mapman annotations are also
provided, with the relevant Mapman bin codes and Pfam names and identifiers available
for reference. Lastly, a list of top-coexpressed genes sorted by Pearson Correlation
Coefficient (PCC) is provided for the user.

Use case 1: Identification of genes involved in pollen coat biosynthesis
Here, we provide a case study on how PEO can be used to identify candidate genes in
pollen coat biosynthesis by using PEO and correlation with other evidence available in
the literature. As a starting point, we will use *ABORTED MICROSPORES* (*AMS*,
AT2G16910), a transcription factor known to regulate pollen coat biosynthesis (Xu et al.,
2014). The pollen coat is a multilayered structure that provides protection for the nascent
male gametophytes from external stresses (Scott et al., 2004). The outer layer, called
exine, is composed primarily of a polymer called sporopollenin, which is a composite of
fatty acids, phenylpropanoids, phenolics, and carotenoids (Ahlers et al., 1999). It was
found that the *ams* null mutant failed to produce functional pollen, and further
investigations led to its identification as a master regulator of various aspects of pollen
biosynthesis (Xu et al., 2014).
Using PEO, we entered its GeneID (AT2G16910) in the search bar on the home
page. We first looked at the gene page of AMS, which showed its highly specific
expression in the anthers (Figure 4). Next, we obtained the list of top co-expressed genes
against *AMS* and used it to compare against gene clusters identified by FlowerNet.
FlowerNet is a flower-specific database based on microarray data focusing on stamen-,

pollen-, or flower-specific expression (Pearce et al., 2015). By generating a correlation
network, FlowerNet generates gene clusters of transcriptionally coregulated genes, and
based on analysis of tissue specificity and expression at different pollen maturation
stages, we were able to identify several clusters associated with pollen development.
Venn diagram analysis showed that AMS co-expressed genes overlapped significantly
(55%, p = 1.4E-26) with Cluster 37, representing sporopollenin biosynthesis, but not
Clusters 21 and 116, representing pollen exine formation and pollen tube growth,
respectively (Figure 4A). This suggests that *AMS* plays a key role in regulating
sporopollenin biosynthesis in pollen development.

A

AMS Co-expression   FN – Cluster 21

39   0   31

11   0

FN – Cluster 37

AMS Co-expression   FN – Cluster 116

50   0   10

B

Gene ID	Gene name	Function	Expression
AT1G01280	CYP703A2	cytochrome P450	Anther
AT1G02050	LAP6	Chalcone and stilbene synthase family protein	Anther
AT1G62940	ACOS5	acyl-CoA synthetase	Anther
AT1G69500	CYP704B1	cytochrome P450	Anther
AT3G11980	FAR2	fatty acid reductases	Anther
AT3G13220	ABCG26	ATP-binding cassette transporter G26	Anther
AT3G23770	N.A	O-Glycosyl hydrolases family 17 protein	Anther
AT3G42960	ATA1	homolog of TASSELSEED2	Anther
AT4G14080	MEE48	Involved in the formation of the pollen wall	Anther
AT4G34850	N.A	Chalcone and stilbene synthase family protein	Anther
AT4G35420	DRL1	Dihydroflavonol 4-reductase-like1)	Anther

Figure 4: Example use case of gene *AMS* (*AT2G16910*), a transcription factor
known to regulate pollen coat biosynthesis. A) Gene expression profile by organs
shows its highly specific expression in pollen. Comparison of top co-expressed genes
with co-expression clusters extracted from FlowerNet shows strong overlap with Cluster
37 (Sporopollenin biosynthesis) but not Cluster 21 (Pollen exine formation) and Cluster
116 (Pollen tube formation/growth). B) Table shows gene names, functions, and specific

expression of genes found in the overlap between AMS co-expression and FlowerNet
Cluster 37.

A Discovery of functional orthologues using MapMan and Pfam annotations

B Output: *Solanum lycopersicum*　　　　　Output: *Oryza sativa*

Figure 5: Discovery of functional orthologues using MapMan and Pfam
annotations. A) Mapman and Pfam annotations for the gene of interest (*AT2G16910*)
provide a list of genes annotated with the bins and domains in other species. B) Example
output for functional orthologue discovery using MapMan annotation. The gene of
Solanum lycopersicum and *Oryza sativa* show specific expression in reproductive tissues
similar to *AT2G16910* in *Arabidopsis*.

Further examination of the overlapped genes (11 genes) revealed a list of genes
that were annotated as being involved in pollen development (*AT1G01280*, *AT1G69500*,

AT4G35420) or whose mutants were essential for male fertility (*AT1G62940*,
AT3G11980, *AT3G13220*, *AT4G35420*), and were all found to be predominantly
expressed in anthers (Figure 4B, Supplemental Dataset S2). This strategy of using
complementary databases can be used to narrow down large gene lists and identify key
genes involved in a pathway of interest. Finally, using the associated Mapman and Pfam
annotations available on the gene page, we can obtain gene orthologues from other
species and use that as a starting point to investigate genetic pathways in other plants
(Figure 5A). As an example, we examined the genes *SOLYC02G079810.3.1* and
LOC_OS07G36460.1 from *Solanum lycopersicum* and *Oryza sativa*, respectively (Figure
5B). Observations of the respective gene loci of *Solanum lycopersicum* and *Oryza sativa*
show transcript enrichment in reproductive tissues similar to that in Arabidopsis.
278
Use case 2: Identification of biosynthetic pathway components of capsaicin
Here, we will provide an example of using PEO to investigate capsaicin biosynthesis in
the crop plant *Capsicum annuum*. The genus Capsicum comprises a large class of crop
plants commonly known as peppers (i.e. chili peppers), and its distinctive
pungency/spiciness is due to the alkaloid capsaicin (Suzuki et al., 1980). We looked at
the literature and found that *Pun1* (*CA02G19260*) is a necessary enzymatic component
of capsaicin biosynthesis. The *Pun1* gene encodes an acyltransferase, and the deletion
of the *Pun1* gene in bell peppers is the reason for its distinctive non-pungency compared
to pungent varieties like chili peppers (Stewart Jr et al., 2005). We searched for the *Pun1*
(*CA02G19260*) gene in our database and observed that it was expressed mainly in fruit
and pericarp tissues, consistent with the site of capsaicin accumulation in the plant
(Figure 6A). Next, we obtained the list of top co-expressed genes for *Pun1* and compared
it against a list of putative capsaicin biosynthetic pathway genes previously obtained by
isolation from *Capsicum annuum* fruit tissue at specific developmental stages (Mazourek
et al., 2009; Zhang et al., 2016). This comparison yielded an overlap of 13.7% (7 genes),
which we examined individually for their function and organ expression (Figure 6B). We
observed that all the genes were predominantly found in the pericarp tissue, consistent
with the site of capsaicin accumulation (Figure 6D). We then obtained a putative map of
the capsaicin biosynthetic pathway (Zhang et al., 2016) and attempted to map our gene

list to it. Surprisingly, we observed that most of the genes mapped to the pyruvate-derived

arm of the pathway, suggesting that *Pun1* is primarily co-regulated with this arm of

capsaicin biosynthesis (Figure 6C).

Figure 6: Example use case of *Pun1* (*CA02G19260*) from *Capsicum annum*, an

important enzymatic component of capsaicin biosynthesis. A) Gene expression

profile by organs shows its highly specific expression in fruit and pericarp, which is also

the site of capsaicin accumulation. B) Comparison of top co-expressed genes with known

capsaicin biosynthetic pathway components yields an overlap of 13.7% (7 genes). C)

Mapping of genes to proposed capsaicin biosynthetic pathway based on putative gene

function (red boxes). D) Specific expression of the seven genes obtained from PEO.

Finally, we also provide several other options for general gene and pathway discovery
using either the organ, Pfam or Mapman specific annotation modes. For instance, by
selecting the 'Organs' tab, we are directed to a page listing all specific organ / tissue types
sorted by Plant Ontology (PO) number. Selecting the PO number (PO:0009005)
corresponding to root directs us to a page containing genes specific to the root, sorted by
SPM values (Supplemental Figure S4). Here, we can select the species of interest, in this
case, *Solanum lycopersicum*, and ask PEO to provide a heatmap of the top 50 genes. As
observed, the heatmap displays a set of genes showing strong enrichment in root tissue
(Supplemental Figure S4).

Next, we show an example whereby Pfam IDs are used as a starting point by selecting
the Pfam tab. Here we chose Pfam ID (PF00182) which corresponds to Chitinase class
I, which is important for defense against fungal pathogens. We are directed to a page
which displays a list of genes sorted by species that are associated with this Pfam ID
(Supplemental Figure S5). Selecting *Physcomitrella patens* as our species of interest, we
can ask PEO to provide a heat map of gene expression in various tissues. We observe
that different genes are associated with different tissue types, providing hypothesis
generation capabilities for the investigation of these genes in specific tissues.

Lastly, we also show how Mapman bins can be used as a starting point by selecting the
Mapman tab. We select the Mapman bincode (12.1.1.1 Chromatin
organisation.chromatin structure. DNA wrapping.histone *(H2A)) as an example. PEO
directs us to a page where we are provided with a list of genes associated with this
bincode sorted by species. In this case, we generate a heatmap of the species Oryza
sativa, where we observe that the majority of the genes are associated with meristematic
or reproductive tissue (Supplemental Figure 6). Again, this simple analysis can provide
useful insights into gene function and any tissue specific properties it may possess as a
starting point for experimental investigations.

Conclusion

We present here a fully annotated Gene Expression Omnibus of 103 plant species, which
provides information on tissue specificity and gene function information of the relevant
gene of interest. PEO also automatically provides a list of top-coexpressed genes for
better elucidation of gene functions and associated pathways. We provided two case
studies of how PEO can be used to study genes and pathways of interest and to narrow
down large putative gene lists based on information on tissue specificity, annotated Pfam
and Mapman functions, and comparison using complementary datasets obtained from
other analyses. These analyses are amenable to any species present in PEO and allow
researchers to build a genetic model of a pathway of interest quickly. As we expand the
number of species available in PEO, we will be able to more effectively empower
researchers working on non-model organisms and more quickly progress the pace of
translational research.

Acknowledgments

We would like to thank Riccardo Delli Ponti for the initial help in collecting the gene
expression data. M.M. is supported by the Singaporean Food Agency grant
SFS_RND_SUFP_001_05 and E.K. is supported by Ministry of Education Tier 3 Grant '
From Tough Pollen to Soft Matter'.

Conflict of interests

The authors declare no conflicts of interest.

References

Ahlers, F., Lambert, J., Wiermann, R., 1999. Structural Elements of Sporopollenin from the
Pollen of Torreya californica Torr. (Gymnospermae): Using the 1H-NMR Technique. Zeitschrift
für Naturforschung C 54, 492–495. https://doi.org/10.1515/znc-1999-7-806
Bray, N.L., Pimentel, H., Melsted, P., Pachter, L., 2016. Near-optimal probabilistic RNA-seq
quantification. Nat. Biotechnol. 34, 525–527. https://doi.org/10.1038/nbt.3519
Buchfink, B., Xie, C., Huson, D.H., 2015. Fast and sensitive protein alignment using DIAMOND.
Nat. Methods 12, 59–60. https://doi.org/10.1038/nmeth.3176
Cravens, A., Payne, J., Smolke, C.D., 2019. Synthetic biology strategies for microbial
biosynthesis of plant natural products. Nat. Commun. 10, 2142.

Delli-Ponti, R., Shivhare, D., Mutwil, M., 2021. Using Gene Expression to Study Specialized
Metabolism—A Practical Guide. Front. Plant Sci. 11, 2074.
https://doi.org/10.3389/fpls.2020.625035
Desborough, M.J., Keeling, D.M., 2017. The aspirin story–from willow to wonder drug. Br. J.
Haematol. 177, 674–683.
Goh, W., Mutwil, M., 2021. LSTrAP-Kingdom: an automated pipeline to generate annotated
gene expression atlases for kingdoms of life. Bioinforma. Oxf. Engl. btab168.
https://doi.org/10.1093/bioinformatics/btab168
Hansen, B.O., Meyer, E.H., Ferrari, C., Vaid, N., Movahedi, S., Vandepoele, K., Nikoloski, Z.,
Mutwil, M., 2018. Ensemble gene function prediction database reveals genes important for
complex I formation in Arabidopsis thaliana. New Phytol. 217, 1521–1534.
https://doi.org/10.1111/nph.14921
Julca, I., Ferrari, C., Flores-Tornero, M., Proost, S., Lindner, A.-C., Hackenberg, D.,
Steinbachová, L., Michaelidis, C., Gomes Pereira, S., Misra, C.S., Kawashima, T., Borg, M.,
Berger, F., Goldberg, J., Johnson, M., Honys, D., Twell, D., Sprunck, S., Dresselhaus, T.,
Becker, J.D., Mutwil, M., 2021. Comparative transcriptomic analysis reveals conserved
programmes underpinning organogenesis and reproduction in land plants. Nat. Plants 7, 1143–
1159. https://doi.org/10.1038/s41477-021-00958-2
Klie, S., Nikoloski, Z., 2012. The choice between MapMan and Gene ontology for automated
gene function prediction in plant science. Front. Genet. 3, 1–14.
https://doi.org/10.3389/fgene.2012.00115
Krishna, S., Bustamante, L., Haynes, R.K., Staines, H.M., 2008. Artemisinins: their growing
importance in medicine. Trends Pharmacol. Sci. 29, 520–527.
Lim, P.K., Zheng, X., Goh, J.C., Mutwil, M., 2022. Exploiting plant transcriptomic databases:
Resources, tools, and approaches. Plant Commun. 100323.
Liu, J., Zhang, Y., Zheng, Y., Zhu, Y., Shi, Y., Guan, Z., Lang, K., Shen, D., Huang, W., Dou, D.,
2023. PlantExp: a platform for exploration of gene expression and alternative splicing based on
public plant RNA-seq samples. Nucleic Acids Res. 51, D1483–D1491.
Ma, L., Sun, N., Liu, X., Jiao, Y., Zhao, H., Deng, X.W., 2005. Organ-specific expression of
Arabidopsis genome during development. Plant Physiol. 138, 80–91.
Mazourek, M., Pujar, A., Borovsky, Y., Paran, I., Mueller, L., Jahn, M.M., 2009. A Dynamic
Interface for Capsaicinoid Systems Biology. Plant Physiol. 150, 1806–1821.
https://doi.org/10.1104/pp.109.136549
Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G.A., Sonnhammer, E.L.L.,
Tosatto, S.C.E., Paladin, L., Raj, S., Richardson, L.J., Finn, R.D., Bateman, A., 2021. Pfam: The
protein families database in 2021. Nucleic Acids Res. 49, D412–D419.
https://doi.org/10.1093/nar/gkaa913
Niazian, M., 2019. Application of genetics and biotechnology for improving medicinal plants.
Planta 249, 953–973.
Nowicka, B., Ciura, J., Szymańska, R., Kruk, J., 2018. Improving photosynthesis, plant
productivity and abiotic stress tolerance–current trends and future perspectives. J. Plant
Physiol. 231, 415–433.

Paddon, C.J., Westfall, P.J., Pitera, D.J., Benjamin, K., Fisher, K., McPhee, D., Leavell, M., Tai,
A., Main, A., Eng, D., 2013. High-level semi-synthetic production of the potent antimalarial
artemisinin. Nature 496, 528–532.
Pearce, S., Ferguson, A., King, J., Wilson, Z.A., 2015. FlowerNet: A Gene Expression
Correlation Network for Anther and Pollen Development. Plant Physiol. 167, 1717–1730.
https://doi.org/10.1104/pp.114.253807
Proost, S., Mutwil, M., 2018. CoNekT: An open-source framework for comparative genomic and
transcriptomic network analyses. Nucleic Acids Res. 46, W133–W140.
https://doi.org/10.1093/nar/gky336
Radivojac, P., Clark, W.T., Oron, T.R., Schnoes, A.M., Wittkop, T., Sokolov, A., Graim, K., Funk,
C., Verspoor, K., Ben-Hur, A., Pandey, G., Yunes, J.M., Talwalkar, A.S., Repo, S., Souza, M.L.,
Piovesan, D., Casadio, R., Wang, Z., Cheng, J., Fang, H., Gough, J., Koskinen, P., Törönen, P.,
Nokso-Koivisto, J., Holm, L., Cozzetto, D., Buchan, D.W.A., Bryson, K., Jones, D.T., Limaye, B.,
Inamdar, H., Datta, A., Manjari, S.K., Joshi, R., Chitale, M., Kihara, D., Lisewski, A.M., Erdin, S.,
Venner, E., Lichtarge, O., Rentzsch, R., Yang, H., Romero, A.E., Bhat, P., Paccanaro, A.,
Hamp, T., Kaßner, R., Seemayer, S., Vicedo, E., Schaefer, C., Achten, D., Auer, F., Boehm, A.,
Braun, T., Hecht, M., Heron, M., Hönigschmid, P., Hopf, T.A., Kaufmann, S., Kiening, M.,
Krompass, D., Landerer, C., Mahlich, Y., Roos, M., Björne, J., Salakoski, T., Wong, A., Shatkay,
H., Gatzmann, F., Sommer, I., Wass, M.N., Sternberg, M.J.E., Škunca, N., Supek, F., Bošnjak,
M., Panov, P., Džeroski, S., Šmuc, T., Kourmpetis, Y.A.I., Van Dijk, A.D.J., Ter Braak, C.J.F.,
Zhou, Y., Gong, Q., Dong, X., Tian, W., Falda, M., Fontana, P., Lavezzo, E., Di Camillo, B.,
Toppo, S., Lan, L., Djuric, N., Guo, Y., Vucetic, S., Bairoch, A., Linial, M., Babbitt, P.C., Brenner,
S.E., Orengo, C., Rost, B., Mooney, S.D., Friedberg, I., 2013. A large-scale evaluation of
computational protein function prediction. Nat. Methods 10, 221–227.
https://doi.org/10.1038/nmeth.2340
Rao, X., Dixon, R.A., 2019. Co-expression networks for plant biology: why and how. Acta
Biochim. Biophys. Sin. 51, 981–988. https://doi.org/10.1093/abbs/gmz080
Rhee, S.Y., Mutwil, M., 2014. Towards revealing the functions of all genes in plants. Trends
Plant Sci. 19, 212–221. https://doi.org/10.1016/j.tplants.2013.10.006
Robinson, A.J., Tamiru, M., Salby, R., Bolitho, C., Williams, A., Huggard, S., Fisch, E.,
Unsworth, K., Whelan, J., Lewsey, M.G., 2018. AgriSeqDB: an online RNA-Seq database for
functional studies of agriculturally relevant plant species. BMC Plant Biol. 18, 1–8.
Ryngajllo, M., Childs, L., Lohse, M., Giorgi, F.M., Lude, A., Selbig, J., Usadel, B., 2011. SLocX:
predicting subcellular localization of Arabidopsis proteins leveraging gene expression data.
Front. Plant Sci. 43.
Sabzehzari, M., Zeinali, M., Naghavi, M.R., 2020. Alternative sources and metabolic
engineering of Taxol: Advances and future perspectives. Biotechnol. Adv. 43, 107569.
Schwacke, R., Ponce-Soto, G.Y., Krause, K., Bolger, A.M., Arsova, B., Hallab, A., Gruden, K.,
Stitt, M., Bolger, M.E., Usadel, B., 2019. MapMan4: A Refined Protein Classification and
Annotation Framework Applicable to Multi-Omics Data Analysis. Mol. Plant 12, 879–892.
https://doi.org/10.1016/j.molp.2019.01.003
Scott, R.J., Spielman, M., Dickinson, H.G., 2004. Stamen Structure and Function. Plant Cell 16,
S46–S60. https://doi.org/10.1105/tpc.017012

Stewart Jr, C., Kang, B.-C., Liu, K., Mazourek, M., Moore, S.L., Yoo, E.Y., Kim, B.-D., Paran, I.,
Jahn, M.M., 2005. The Pun1 gene for pungency in pepper encodes a putative acyltransferase.
Plant J. 42, 675–688. https://doi.org/10.1111/j.1365-313X.2005.02410.x
Suzuki, T., Fujiwake, H., Iwai, K., 1980. Intracellular localization of capsaicin and its analogues,
capsaicinoid, in Capsicum fruit 1. Microscopic investigation of the structure of the placenta of
Capsicum annuum var. annuum cv. Karayatsubusa1. Plant Cell Physiol. 21, 839–853.
https://doi.org/10.1093/oxfordjournals.pcp.a076058
Tan, Q.W., Goh, W., Mutwil, M., 2020. LSTrAP-Cloud: A User-Friendly Cloud Computing
Pipeline to Infer Coexpression Networks. Genes 11, 428.
https://doi.org/10.3390/genes11040428
Walls, R.L., Cooper, L., Elser, J., Gandolfo, M.A., Mungall, C.J., Smith, B., Stevenson, D.W.,
Jaiswal, P., 2019. The Plant Ontology Facilitates Comparisons of Plant Development Stages
Across Species. Front. Plant Sci. 10. https://doi.org/10.3389/fpls.2019.00631
Wetterstrand, K.A., 2013. DNA sequencing costs: data from the NHGRI Genome Sequencing
Program (GSP). Natl. Hum. Genome Res. Inst.
Xu, J., Ding, Z., Vizcay-Barrena, G., Shi, J., Liang, W., Yuan, Z., Werck-Reichhart, D.,
Schreiber, L., Wilson, Z.A., Zhang, D., 2014. ABORTED MICROSPORES Acts as a Master
Regulator of Pollen Wall Formation in Arabidopsis. Plant Cell 26, 1544–1556.
https://doi.org/10.1105/tpc.114.122986
Yu, Y., Zhang, H., Long, Y., Shu, Y., Zhai, J., 2022. Plant public RNA-seq database: a
comprehensive online database for expression analysis of~ 45 000 plant public RNA-seq
libraries. Plant Biotechnol. J. 20, 806.
Zhang, Z.-X., Zhao, S.-N., Liu, G.-F., Huang, Z.-M., Cao, Z.-M., Cheng, S.-H., Lin, S.-S., 2016.
Discovery of putative capsaicin biosynthetic genes by RNA-Seq and digital gene expression
analysis of pepper. Sci. Rep. 6, 34121. https://doi.org/10.1038/srep34121
482
483