C9orf72 polyPR directly binds to various nuclear transport components

Hamidreza Jafarinia[1], Erik Van der Giessen[1], Patrick R. Onck[1*]

[1]Zernike Institute for Advanced Materials, University of Groningen, Nijenborgh 4, 9747AG Groningen, The Netherlands.

*Corresponding author: p.r.onck@rug.nl

Abstract

The disruption of nucleocytoplasmic transport (NCT) is an important mechanism in neurodegenerative diseases. In the case of C9orf72-ALS, trafficking of macromolecules through the nuclear pore complex (NPC) might get frustrated by the binding of C9orf72-translated arginine-containing dipeptide repeat proteins (R-DPRs) to the Kapβ family of nuclear transport receptors. Beside Kapβs, several other types of transport components have been linked to NCT impairments in R-DPRs expressed cells, but the molecular origin of these observations has not been clarified. Here, we adopt a coarse-grained molecular dynamics model at amino-acid resolution to study the direct interaction between polyPR, the most toxic DPR, and various nuclear transport components to elucidate the binding mechanisms and provide a complete picture of potential polyPR-mediated NCT defects. We found polyPR to directly bind to several isoforms of the Impα family, CAS (the specific exporter of Impα) and RanGAP. We observe no binding between polyPR and Ran. Longer polyPRs at lower salt concentrations also make contact with RanGEF and NTF2. Analyzing the polyPR contact sites on the transport components reveals that polyPR potentially interferes with RanGTP/RanGDP binding, with cargo-NLS binding to Impα, with cargo-NLS release from Impα, and with Impα export from the nucleus. The abundance of polyPR binding sites on multiple transport components combined with the inherent polyPR length dependence makes direct polyPR interference of NCT a potential mechanistic pathway of C9orf72 toxicity.

Introduction

The C9orf72 G4C2 hexanucleotide repeat expansion is the most common genetic mutation in amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD) [1, 2]. This expansion can be translated into five types of dipeptide repeat proteins (DPRs): polyPR, polyGR, polyGA, polyGP, and polyPA [3]. The positively-charged arginine-containing DPRs (R-DPRs) show the highest levels of toxicity in different cell and animal models [4-10] with polyPR known to be the most toxic DPR [5, 9, 11]. R-DPRs have been linked to a wide variety of cellular defects [12-16], but a growing body of evidence suggests that neurodegenerative diseases, including C9orf72 ALS/FTD (C9-ALS/FTD), may be caused by disruption of nucleocytoplasmic transport (NCT) [17-20]. At the same time, many transport components that play a prominent role in NCT have been identified to function as modifiers of G4C2/DPR toxicity [5, 10, 21].

The regulated trafficking of proteins and RNA between the nucleus and cytoplasm occurs through nuclear pore complexes (NPCs) embedded in the nuclear membrane [22, 23]. The NPC is lined with intrinsically disordered phenylalanine-glycine-rich Nups (FG-Nups) that collectively function as a selective permeability barrier. Small molecules rapidly diffuse through the NPC, but the passage of larger cargoes across the barrier needs to be facilitated by their binding to nuclear transport receptors (NTRs) [24-26]. The β-karyopherin (Kapβ) family is the largest class of NTRs and includes both import and export receptors [27]. Another essential regulator of NCT is the GTPase Ran, a small protein bound to guanosine triphosphate (GTP) in the nucleus and to guanosine diphosphate (GDP) in the cytoplasm [24]. The directionality of NCT is mediated by the RanGTP-RanGDP gradient over the nuclear envelope, which is preserved by the cytoplasmic GTPase-activating protein RanGAP and the nuclear guanine nucleotide exchange factor RanGEF [24, 28].

In the import cycle, importins bind their cargoes directly through a nuclear localization signal (NLS) encoded on a cargo. Importin β1 (Impβ1) can also recruit importin α (Impα) that functions as a cargo-adaptor protein. Impα binds to Impβ1 through its N-terminal importin β-binding (IBB) domain [29, 30]. The importin-NLS-cargo complex then shuttles to the nucleus. The binding of RanGTP to importin in the nucleus disassembles the importin-NLS-cargo complex and the RanGTP-importin complex is recycled to the cytoplasm. When cargo is bound to Impβ1 via Impα, RanGTP dissociates Impβ1 from the Impα-NLS-cargo. This triggers a competition between the flexible IBB domain of Impα and NLS-cargo for binding to Impα,

thus facilitating the dissociation of the NLS-cargo. Nucleoporins such as Nup50/Nup2 also

catalyze this process by binding to Impα and accelerating the dissociation rate of the cargo-

NLS [22]. The specific receptor CAS bound to RanGTP is required to export Impα to the

cytoplasm. It has been proposed that CAS first displaces Nup50/Nup2 from Impα after which

the RanGTP-CAS-Impα returns to the cytoplasm. The hydrolysis of RanGTP to RanGDP in

the cytoplasm by RanGAP disassembles the RanGTP-importin and RanGTP-CAS-Impα

complexes [22]. RanGDP is transported back to the nucleus by nuclear transport factor 2

(NTF2) where the RanGDP-NTF2 complex dissociates when RanGEF regenerates RanGTP

[22]. In the export cycle, RanGTP promotes the loading of cargoes with nuclear export signal

(NES) to the exportin in the nucleus. The resulting RanGTP-exportin-NES-cargo complex

moves to the cytoplasm. Once there the complex is disassembled by RanGAP which hydrolyses

RanGTP to RanGDP [24].

In a recent study we have analyzed the binding of polyPR to the Kapβ family of importins and

exportins [13, 16, 31] by using coarse-grained (CG) molecular dynamics simulations [32].

Depending on its length, polyPR can interact with several cargo-, IBB-, RanGTP-, and FG-

Nup-binding sites on the Kapβs [32]. Beside Kapβs, there is evidence for direct binding of

Impα isomers with R-DPRs [16]. Some regulators of the Ran cycle are also affected in R-DPR-

mediated toxicity, where R-DPRs have been shown to cause mislocalization and abnormal

accumulation of RanGAP [33], and mislocalization of Ran and RanGEF in cell culture models

[5, 33, 34]. RanGAP and RanGEF also appear to be modifiers of R-DPR toxicity in genetic

studies [5, 10, 11]. It is not clear, however, whether these effects arise from a direct interaction

of R-DPRs with NCT components. The aim of the current work is to extend the findings of [32]

by investigating the interaction between polyPR and various NCT components by means of CG

molecular dynamics computations.

Results and discussion

Coarse-grained models of nucleocytoplasmic transport components

We use the residue-scale CG molecular dynamics approach developed and applied earlier to

study DPR phase separation [35] and the direct binding of polyPR to numerous members of the

Kapβ family [32]. In the present work we investigate the interaction of polyPR with unbound

human Impα isomers (Impα1, Impα3, Impα5, Impα7), Ran, CAS (the specific exporter of

Impα), RanGEF, and NTF2. We also include KAP60 (homolog of Impα), Cse1 (homolog of
CAS), and RanGAP from yeast, since contrary to the human homologs, the crystal structures
of the KAP60-Cse1 complex and the RanGAP-RanGppNHp complex (fission yeast RanGAP
bound to the non-hydrolyzable form of human RanGTP) are available in the protein data bank.
This enables us to investigate a possible polyPR interference with Impα export and the RanGAP
function in the model system of yeast. Moreover, yeast has been employed previously to study
NCT defects caused by DPRs [5, 36]. More details about the selected transport components can
be found in table S2 of the SI.

In our one-bead-per-amino-acid (1BPA) CG models of transport components, each residue is
represented by a single bead at the position of the alpha-carbon atom. The overall tertiary
structure of the NCT components is preserved through a network of stiff harmonic bonds, and
the distribution of charged and aromatic residues are included in the model. The 1BPA force
field correlates with experimental findings for polyPR-Kapβs interactions [32]. The CG models
of the NCT components studied are shown in figure 1. The Impα isomers contain a flexible N-
terminal IBB domain, followed by a helical core that is constructed from 10 Armadillo (ARM)
repeats each consisting of three alpha helices. The NLS binding sites are located on the concave
surface of the helical core [37]. The IBB domain has an autoinhibitory role and when it is not
bound to Impβ1, it binds to the ARM concave core and competes with NLS binding [22]. To
simplify the study of possible binding between polyPR and NLS binding sites of Impα, the CG
models of Impα isomers are built without their N-terminal IBB domains and referred to as
ImpαΔN. This approach enables easier investigation of polyPR's interaction with NLS binding
sites. Similar to the importins and exportins, CAS and Cse1 are constructed from HEAT repeats,
with each repeat consisting of two antiparallel α-helices, named A and B, connected by linkers
of different lengths [38]. RanGEF is mainly constructed from β-strands and has an overall
appearance of a seven-bladed propeller with each blade consisting of a four-stranded
antiparallel β-sheet [39]. The RanGAP model of fission yeast used in our study is constructed
from eleven leucine-rich repeats (LRRs) forming a symmetric crescent followed by a highly
negatively-charged C-terminal region. Each LRR motif consists of a β strand-α helix hairpin
unit [40]. NTF2 is a homodimer with each monomer consisting of a β-sheet and three α-helices
[41]. The CG model of Ran is based on the nucleotide-free state of the molecule. More details

about the CG models and forcefield are provided in the Methods section and section 1 of the
SI.

PolyPR binds to several transport components through electrostatic interactions

In investigating the direct interaction with the transport components shown in figure 1 we
include the potential effect of polyPR length, as various studies have found that the repeat length
of DPRs strongly correlates with toxicity [4, 42-44]. Simulations of polyPR with varying
numbers of repeat units, i.e., PR7, PR20, and PR50, are performed at two salt concentrations:
200 mM, similar to previous in vitro experiments performed for Kapβ importins and Impα
isomers interacting with R-DPRs [16], and a lower ion concentration of 100 mM to study the
effect of salt concentration.

To quantify the interaction between polyPR and the transport components, we calculate the
time-averaged number of contacts C_t using a cutoff of 1 nm. The number of contacts is
normalized by the sequence length of the transport components (N_{TC}) and by the polyPR length
(N_{PR}). The values of C_t are plotted against the charge parameter

$$NCPR - fM/R_g, \tag{1}$$

where the net charge per residue $NCPR$ is the total charge of the transport component (in units
of elementary charge e) divided by its sequence length, M (in units of e.nm) is the time-averaged
total dipole moment, and R_g (in units of nm) is the time-averaged radius of gyration of the
transport components in isolation. The dimensionless parameter f is a free parameter which is
calculated to be 0.0036 for the best linear fit in figures 2a for PR20 and PR50. The quality of
fit (R^2) is plotted as a function of f in figure S1. The linear correlation observed in figure 2a
confirms an electrostatically driven interaction between polyPR and the transport components.
Moreover, it highlights the importance of the spatial distribution of charge over the transport
components, as characterized here through the dipole moment. CAS and Cse1, the specific
exporters of Impα in human and yeast, respectively, are constructed from HEAT repeats and
have a super-helical conformation similar to the Kapβs studied before [32]. We therefore
present the results for these two cases jointly with the results for the Kapβ set (taken from [32])
in figure 2b. For this set, the best fit is obtained for $f = 0$ (see figure S1), which indicates the

dominant role of $NCPR$ for the number of contacts between polyPR and the Kapβs, CAS and

Cse1.

As can be seen in figure 2a, RanGAP features a much larger number of contacts with polyPR

than the other transport components, which can be related to the higher negative net charge and

the higher dipole moment of this molecule (see the dipole moments M and $NCPR$ of the

transport components in figures S2 and S3 of the SI). In contrast, polyPR makes a negligible

number of contacts with Ran since it has a relatively low dipole moment and no net charge. For

NTF2 the value of $NCPR$ is − 0.047 e, comparable to several members of the ImpαΔN family,

but the lower dipole moment of NTF2 results in a lower number of contacts with polyPR. The

binding of PR20 to the ImpαΔN isomers is consistent with the Impα1 and Impα3 binding to R-

DPRs found in experiment [16], despite the fact that the N-terminal IBB domains of Impα1,3

are excluded from our CG models. At 200 mM salt concentration, polyPR does not bind to

NTF2 and Ran. PolyPR contact with RanGEF is also very low at this salt concentration.

Reducing the salt concentration to 100 mM, increases the number of contacts, clearly indicating

that electrostatic force is the main driver for binding. At this lower salt concentration, PR7,

PR20, and PR50 make contact with RanGEF, but contact with NTF2 is only observed for longer

polyPR chains. In figure 2, the number of contacts per unit length of polyPR is often seen to be

lower for longer polyPRs especially for the lower salt concentrations, see e.g. the results in

figure 2a for the ImpαΔN family and RanGAP at 100 mM salt concentration. This is due to the

fact that most of the residues make contact with the target protein for shorter polyPRs, while

for longer polyPRs only some parts of the chain are in contact with the transport components

and other regions make less or no contact.

PolyPR interacts with important binding sites of transport components

In order to gain a better understanding of how polyPR interacts with each transport component,

we examined the polyPR contact probability of each residue in the sequence of the transport

component. The contact probability of each residue is defined as the probability of having at

least one polyPR residue within its 1 nm proximity. Figure 3a reveals our results of the

interaction between PR7 and PR50 with Impα1ΔN, KAP60ΔN, Cse1, RanGAP, RanGEF and

NTF2 (for the other transport components, see figure S4). These findings indicate that a longer

polyPR makes contact with a larger number of sites, and also exhibits a higher contact

probability with individual residues compared to a shorter polyPR. We also observe that some

regions at the C-terminal ends of the ImpαΔN family and RanGAP are permanently bound to

polyPR. We also compare the polyPR binding sites with known binding sites of transport

components (according to the PiSITE webserver), highlighting them at the bottom of each

subfigure. Figure 3b displays the number of contact residues shared between polyPR and the

native binding partners of each transport component. As expected, the general trend is that the

number of shared binding sites increases with increasing polyPR length. PolyPR interacts with

the ImpαΔN family at several known cargo-NLS, Nup50/Nup2, and Cse1 binding sites. Longer

polyPRs exhibit a significantly stronger interaction with cargo-NLS binding sites of ImpαΔN

compared to shorter polyPRs, which only interact with a limited number of sites. However, we

observe that both short and long polyPRs bind to Nup50/Nup2, Cse1, and RanGTP binding

sites particularly those located near the C-terminal end of ImpαΔN. In the case of Cse1, we

observe polyPR binding to Impα and RanGTP binding sites. PolyPR interacts with the known

RanGTP binding sites of RanGAP. We also show that polyPR is able to bind to the highly

negatively-charged region in the C-terminal domain of RanGAP that follows the LRR domain.

It has been suggested that this negatively-charged region is in close proximity to a positively-

charged region in Ran (in the complex formed by Ran and RanGAP) and plays a role in RanGTP

hydrolysis [45, 46]. Unfortunately, there is no crystal structure for this region in the PDB

structure and thus the binding sites are not known. In the case of RanGEF, a longer polyPR

interacts with a high percentage of the known Ran binding sites. For NTF2 we observe polyPR

interaction with RanGDP binding sites. PolyPR makes negligible contacts with nucleotide-free

state of Ran. For CAS the binding sites are not known. Therefore, these two cases are excluded

from figure 3b.

The findings presented in figure 3b and S4, along with previous research on the interaction

between polyPR and Kapβs (importins and exportins) [32], lead to the following mechanistic

understanding for the direct effect of polyPR on NCT as illustrated in figure 4. In this figure

the native binding interactions that are affected by polyPR are indicated by red dashed arrows

and those that are unaffected by grey arrows. In the import cycle, polyPR may impede the

loading of cargo to Kapβ and Impα families (as shown in inset A) by binding to the cargo-NLS

sites. PolyPR also binds to the RanGTP binding sites on the Kapβs and the Nup2/Nup50 binding

sites on Impα, which could result in defects in the dissociation of cargo/cargo-Impα from Kapβ

and cargo from Impα (as shown in insets B and C). PolyPR also interacts with certain CAS/Cse1

binding sites that recognize Impα and RanGTP, possibly interfering with the formation of the

RanGTP-CAS/Cse1-Impα complex (as shown in inset D). The relatively high number of contacts between polyPR and RanGAP, as well as polyPR binding to RanGTP binding sites on RanGAP, suggest a possible defect in the dissociation of the RanGTP-importin and RanGTP-CAS/Cse1-Impα complexes in the import and Ran cycle, and of the RanGTP-exportin-cargo complex in the export cycle (as shown in inset E). Longer polyPRs may also interfere with the Ran cycle by hindering the loading of RanGDP to NTF2 and the exchange of GDP to GTP in Ran by RanGEF (as shown in insets G and F). In the export cycle, as examined before [32], polyPR binding to RanGTP and FG-Nup binding sites may affect cargo-loading onto exportins (as shown in inset H) and the transport of exportins through the NPC (as shown in inset I). It should be noted that the mechanisms proposed in figure 4 do not hold equal weight, and the relative contributions based on the number of contacts (presented in figure 2), and the number of contacts with important binding sites (presented in figure 3b and S4) should be considered when comparing the relative significance of the suggested mechanisms.

Conclusion

In this study we used coarse-grained molecular dynamics simulations to show that polyPR binds to several nuclear transport components. Similar to the interaction of polyPR to the Kapβ family, the interaction of polyPR to other transport components is driven by electrostatic interactions and depends sensitively on polyPR length. Reducing the salt concentration or increasing the polyPR length increases the number of contacts with different transport components. The effect of polyPR length suggests a molecular basis for the more toxic nature of longer polyPRs in animal and cell models [4, 42-44].

We observed polyPR binding to several members of the Impα family, CAS, Cse1, and RanGAP yet no binding to Ran. PolyPR strongly binds to a highly negatively charged region in the C-terminal domain of fission yeast RanGAP. The human RanGAP also contains a similar highly negatively-charged region. Our simulations therefore suggest that a direct interaction may contribute to the observed polyPR-mediated accumulation and mislocalization of RanGAP in HeLa cells [33]. PolyPR also binds to RanGEF and NTF2 at lower salt concentrations or when the polyPR length is large enough. We also showed that including the dipole moment results in a better fit to the simulation results for the number of contacts, thus suggesting the importance of the spatial separation of charge on the transport component for polyPR binding to its targets inside the cell.

We showed that polyPR interacts with important binding sites of different transport components

in a polyPR-length-dependent manner, with polyPR interaction with RanGTP/RanGDP binding

sites being a common feature between the transport components. This suggests a strong polyPR

interference with the Ran gradient across the nuclear envelop. For the ImpαΔN family, we

observe polyPR interaction with cargo-NLS and Nup2/Nup50 binding sites. In the case of

KAP60ΔN (yeast homolog of Impα), we observe polyPR interaction with Cse1 binding sites.

For Cse1 (yeast CAS), we also observed polyPR interacting with Impα binding sites. These

findings suggest polyPR interference with the cargo-NLS association and disassociation with

Impα, and the export of Impα.

In conclusion, we showed a pronounced direct binding interaction between polyPR and a

surprisingly large number of transport components. By integrating our findings with previously

reported data, this work proposes a molecular model that explains how the binding of polyPR

might interfere with distinct stages of the transport cycle. The intrinsic length dependence of

polyPR binding to important binding sites of many transport components promotes this

mechanism to a potential target for therapeutic interventions. Overall, our results offer a basis

for future research that aims to explore the impact of C9orf72 R-DPRs on NCT disruption and

the subsequent downstream consequences.

Figure 1: Coarse-grained models of transport components used in this study

One-bead-per-amino acid (1BPA) representation of the various transport components modeled in the current study. These are several members of the Impα family (excluding the N-terminal IBB domain), specific exporters of Impα (CAS and Cse1), RanGAP, RanGEF, NTF2, and Ran. Details regarding the CG models and protein sequences are listed in tables S2 and S3.

Figure 2: The transport component's net charge per residue and dipole moment, together with polyPR length, affect polyPR interaction with various nuclear transport components

(a) and (b) show the normalized time-averaged number of contacts C_t for the interaction between polyPR with 7, 20, and 50 repeat units with different types of transport components. The results are shown for monovalent salt concentrations of $C_{\text{salt}} = 200$ mM (left panels) and $C_{\text{salt}} = 100$ mM (right panels). Subfigure (a) shows the results for the transport components shown in figure 1, excluding the specific exporters of Impα: CAS and Cse1. A linear correlation is observed between the normalized C_t and $NCPR - fM/R_g$ with f calculated to be 0.0036 for the best fit. The net charge per residue $NCPR$ is in units of elementary charge e, the dipole moment M is in units of e.nm, and the radius of gyration R_g is in units of nm. Subfigure (b) shows the results for the Kapβ data set (data points taken from [32]) together with CAS and Cse1. For this case, a linear correlation between C_t and $NCPR$ is observed. The dashed lines show linear fits for PR20 and PR50. The error bars show half the standard deviation.

Figure 3: PolyPR interacts with several known binding sites of nuclear transport components in a length-dependent manner

(a) The contact probability for each residue in the sequence of transport components interacting

with polyPR. The plot displays the contact probability for six transport components:

Impα1ΔN, KAP60ΔN, Cse1, RanGAP, RanGEF and NTF2 at a salt concentration of 100 mM.

Results for Impα3ΔN, Impα5ΔN, Impα7ΔN, CAS, and Ran are shown in figure S4. Each

figure shows two curves for PR7 and PR50. The bottom part of each figure shows the binding

sites for NLS-cargo, Impα, CAS/Cse1, RanGTP, and Nup50/Nup2 using different colors.

These binding sites are obtained from the crystal structures of the bound states of transport

components in the Protein Data Bank using PiSITE (see table S2 of the SI for more details).

For each transport component the following binding sites are marked. For the Impα family:

NLS-cargo (vertical black lines), CAS/Cse1 (vertical purple lines) and Nup50/Nup2 (vertical

orange lines) binding sites. For CAS/Cse1: Impα (vertical black lines) and RanGTP (vertical

green lines) binding sites. For RanGAP: RanGTP (vertical green lines) binding sites. For

RanGEF: Ran (vertical green lines) binding sites. For NTF2: RanGDP (vertical green lines)

binding sites. The Ran binding sites marked for RanGEF are taken from the RanGEF-Ran

complex (an intermediate step in the RanGEF function).

(b) The number of shared contact sites between polyPR and the binding partners of the transport

components, referred to as N_{shared}, are plotted for PR7, PR20, and PR50. In each bar plot, the

numbers inside the parentheses on the horizontal axis shows the number of known binding

sites obtained from PiSITE. If there is no known binding site, a (-) mark is used instead. The

results for PR7, PR20, and PR50 are reported from left to right for each set of bar plots. The

bars with darker colors represent longer polyPR chains.

Figure 4: Suggested molecular mechanism of polyPR interference with the native function of transport components in the nucleocytoplasmic transport cycle.

(a) Proposed mechanistic pathways of polyPR interference with the import cycle (left panel), Ran cycle (middle panel), and export cycle (right panel). Steps in the NCT cycle are represented with grey arrows, and a red dashed arrow indicates where polyPR may interfere with the transport cycle. The letters A-H are used to illustrate how polyPR may disrupt the function of the transport components. Each letter corresponds to a mechanistic mechanism shown at the bottom of the figure in grey circles. It should be noted that the proposed mechanisms are not equally significant. The relative significance of the suggested molecular mechanisms can be obtained by considering their relative contributions based on the number of contacts and the number of contacts with important binding sites as presented in figures 2 and 3, respectively.

Methods

Coarse-grained model

We adopt a one-bead-per-amino acid (1BPA) force field to study polyPR interaction with nucleocytoplasmic transport components. This 1BPA approach has been initially developed to simulate disordered FG-Nups [25, 47-52], and extended later to study the phase separation of DPRs [35], and the interaction of polyPR with Kapβs [32]. The force field potentials and parameters in this study are identical to those employed in [32]. The bonded potentials for polyPR are residue and sequence specific. For non-bonded polyPR-polyPR interactions, the force field accounts for hydrophobic/hydrophilic and electrostatic interactions. The crystal structure of the transport components is maintained using a stiff harmonic potential $\phi_{\text{network}} = K(r-b)^2$, where K is 8000 kJ/mol/nm^2 and b is the distance between the amino acid beads in the crystal structure. A bond is made between the beads if b is less than 1.4 nm. The unresolved regions in the crystal structure from the Protein Data Bank, and the regions with a lower prediction score (< 70 pLDDT) from Alpha Fold [53, 54] are included in the CG model as disordered regions. The CG model of fission yeast RanGAP includes two alpha helices in the C-terminal domain, as predicted by AlphaFold. The polyPR interactions with transport components are classified into three categories: (1) electrostatic interactions, (2) cation-pi interactions, and (3) excluded volume interactions. Our force field also accounts for the screening effect of ions. More details about the CG models and forcefield are provided in table S2 and section 1 of the SI.

Simulation and analysis

Langevin dynamics simulations are performed at 300 K at monovalent salt concentrations of 100 mM and 200 mM in NVT ensembles with a time-step of 0.02 ps and a Langevin friction coefficient of 0.02 ps^{-1} using GROMACS version 2018. Simulations are performed for at least 2.5 μs in cubic periodic boxes, and the last 2 μs are used for analyzing the interaction between polyPR and the transport components. The binding sites are obtained from the crystal structures of the bound states of transport components in the Protein Data Bank using PiSITE [55]. This web-based database provides interaction sites of a protein from multiple PDBs including similar proteins. The RanGTP binding data (vertical green lines) in figure 3 and S4 contains binding residues for both RanGTP and RanGppNHp, the non-hydrolysable form of RanGTP. The time-averaged number of contacts between the polyPR and transport components in figure 2 is

obtained by summing the number of contacts per time frame (i.e. the number of polyPR/transport components residue pairs that are within 1 nm) over all frames and dividing by the total number of frames. The contact probability for each transport component residue is the probability of having at least one polyPR residue within 1 nm proximity of the transport component residue. The contact probability is calculated for each transport component residue by dividing the number of frames for which this contact criterion is satisfied, by the total number of frames. Residue i is considered to be a contact site if the contact probability for this residue is larger than 0.10. N_{shared} is the number of transport component residues that make contact with polyPR (obtained in our simulations) and at the same time are known for recognition of native binding partners (according to PiSITE). The time-averaged total dipole moment of the CG models of transport components is calculated using gmx dipole in GROMACS.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

1. Renton, A.E., et al., A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron*, 2011. **72**(2): p. 257-268.

2. DeJesus-Hernandez, M., et al., Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron*, 2011. **72**(2): p. 245-256.

3. Mori, K., et al., The C9orf72 GGGGCC Repeat Is Translated into Aggregating Dipeptide-Repeat Proteins in FTLD/ALS. *Science (New York, N.Y.)*, 2013. **339**(6125): p. 1335-1338.

4. Mizielinska, S., et al., C9orf72 repeat expansions cause neurodegeneration in Drosophila through arginine-rich proteins. *Science*, 2014. **345**(6201): p. 1192-1194.

5. Jovičič, A., et al., Modifiers of C9orf72 dipeptide repeat toxicity connect nucleocytoplasmic transport defects to FTD/ALS. *Nat. Neurosci.*, 2015. **18**(9): p. 1226-1229.

6. Kwon, I., et al., Poly-dipeptides encoded by the C9orf72 repeats bind nucleoli, impede RNA biogenesis, and kill cells. *Science*, 2014. **345**(6201): p. 1139-1145.

7. Zhang, Y.J., et al., Heterochromatin anomalies and double-stranded RNA accumulation underlie C9orf72 poly(PR) toxicity. *Science*, 2019. **363**(6428).

8. Zhang, Y.J., et al., Poly(GR) impairs protein translation and stress granule dynamics in C9orf72-associated frontotemporal dementia and amyotrophic lateral sclerosis. *Nat. Med.*, 2018. **24**(8): p. 1136-1142.

9. Wen, X., et al., Antisense proline-arginine RAN dipeptides linked to C9ORF72-ALS/FTD
form toxic nuclear aggregates that initiate invitro and invivo neuronal death. *Neuron*, 2014.
84(6): p. 1213-1225.

10. Boeynaems, S., et al., Drosophila screen connects nuclear transport genes to DPR pathology in
c9ALS/FTD. *Sci. Rep.*, 2016. **6**(1): p. 20877-20877.

11. Lee, K.H., et al., C9orf72 Dipeptide Repeats Impair the Assembly, Dynamics, and Function of
Membrane-Less Organelles. *Cell*, 2016. **167**(3): p. 774-788.e17.

12. Shi, K.Y., et al., Toxic PRn poly-dipeptides encoded by the C9orf72 repeat expansion block
nuclear import and export. *Proceedings of the National Academy of Sciences of the United*
States of America, 2017. **114**(7): p. 201620293-E1117.

13. Hayes, L.R., L. Duan, K. Bowen, P. Kalab, and J.D. Rothstein, C9orf72 arginine-rich
dipeptide repeat proteins disrupt karyopherin-mediated nuclear import. *Elife*, 2020. **9**.

14. Balendra, R. and A.M. Isaacs, C9orf72-mediated ALS and FTD: multiple pathways to disease.
Nature Reviews Neurology, 2018. **14**(9): p. 544-558.

15. Freibaum, B.D., et al., GGGGCC repeat expansion in C9orf72 compromises
nucleocytoplasmic transport. *Nature*, 2015. **525**(7567): p. 129-133.

16. Hutten, S., et al., Nuclear Import Receptors Directly Bind to Arginine-Rich Dipeptide Repeat
Proteins and Suppress Their Pathological Interactions. *Cell Rep.*, 2020. **33**(12): p. 108538.

17. Kim, H.J. and J.P. Taylor, Lost in Transportation: Nucleocytoplasmic Transport Defects in
ALS and Other Neurodegenerative Diseases. *Neuron*, 2017. **96**(2): p. 285-297.

18. Mihevc, S.P., et al., Nuclear trafficking in amyotrophic lateral sclerosis and frontotemporal
lobar degeneration. *Brain : a journal of neurology*, 2016. **140**(1): p. 13-26.

19. Boeynaems, S., E. Bogaert, P. Van Damme, and L. Van Den Bosch, Inside out: the role of
nucleocytoplasmic transport in ALS and FTLD. *Acta Neuropathol.*, 2016. **132**(2): p. 159-173.

20. Jovičić, A., J.W. Paul, and A.D. Gitler, Nuclear transport dysfunction: a common theme in
amyotrophic lateral sclerosis and frontotemporal dementia. *Journal of neurochemistry*, 2016.
138(NA): p. 134-144.

21. Kramer, N.J., et al., CRISPR-Cas9 screens in human cells and primary neurons identify
modifiers of C9ORF72 dipeptide-repeat-protein toxicity. *Nat. Genet.*, 2018. **50**(4): p. 603-612.

22. Stewart, M., Molecular mechanism of the nuclear protein import cycle. *Nat. Rev. Mol. Cell*
Biol., 2007. **8**(3): p. 195-208.

23. Strambio-De-Castillia, C., M. Niepel, and M.P. Rout, The nuclear pore complex: bridging
nuclear transport and gene regulation. *Nat. Rev. Mol. Cell Biol.*, 2010. **11**(7): p. 490-501.

24. Kalita, J., L.E. Kapinos, and R.Y.H. Lim, On the asymmetric partitioning of
nucleocytoplasmic transport - recent insights and open questions. *J. Cell Sci.*, 2021. **134**(7).

25. Popken, P., A. Ghavami, P.R. Onck, B. Poolman, and L.M. Veenhoff, Size-dependent leak of
soluble and membrane proteins through the yeast nuclear pore complex. *Mol. Biol. Cell*, 2015.
26(7): p. 1386-1394.

26.   Timney, B.L., et al., Simple rules for passive diffusion through the nuclear pore complex. *J.*
494   *Cell Biol.*, 2016. **215**(1).

495   27.   O'Reilly, A.J., J.B. Dacks, and M.C. Field, Evolution of the karyopherin-β family of
496   nucleocytoplasmic transport factors; ancient origins and continued specialization. *PLoS One*,
497   2011. **6**(4): p. e19308.

498   28.   Görlich, D., M.J. Seewald, and K. Ribbeck, Characterization of Ran-driven cargo transport
499   and the RanGTPase system by kinetic measurements and computer simulation. *EMBO J.*,
500   2003. **22**(5): p. 1088-100.

501   29.   Pumroy, R.A. and G. Cingolani, Diversification of importin-α isoforms in cellular trafficking
502   and. *Biochemical Journal*, 2015. **466**: p. 13-28.

503   30.   Lott, K. and G. Cingolani, The importin β binding domain as a master regulator of
504   nucleocytoplasmic transport. *Biochim Biophys Acta*, 2011. **1813**(9): p. 1578-92.

505   31.   Nanaura, H., et al., C9orf72-derived arginine-rich poly-dipeptides impede phase modifiers.
506   *Nat Commun*, 2021. **12**(1): p. 5301.

507   32.   Jafarinia, H., E. Van der Giessen, and P.R. Onck, Molecular basis of C9orf72 poly-PR
508   interference with the β-karyopherin family of nuclear transport receptors. *Sci. Rep.*, 2022.
509   **12**(1): p. 21324.

510   33.   Ryan, S., S. Rollinson, E. Hobbs, and S. Pickering-Brown, C9orf72 dipeptides disrupt the
511   nucleocytoplasmic transport machinery and cause TDP-43 mislocalisation to the cytoplasm.
512   *Sci. Rep.*, 2022. **12**(1): p. 4799.

513   34.   Zhang, K., et al., Stress Granule Assembly Disrupts Nucleocytoplasmic Transport. *Cell*, 2018.
514   **173**(4): p. 958-971.e17.

515   35.   Jafarinia, H., E. van der Giessen, and P.R. Onck, Phase Separation of Toxic Dipeptide Repeat
516   Proteins Related to C9orf72 ALS/FTD. *Biophysical Journal*, 2020.

517   36.   Semmelink, M.F.W., et al., Nuclear transport under stress phenocopies transport defects in
518   models of C9Orf72 ALS. *bioRxiv*, 2022: p. 2022.04.13.488135.

519   37.   Pumroy, R.A., S. Ke, D.J. Hart, U. Zachariae, and G. Cingolani, Molecular determinants for
520   nuclear import of influenza A PB2 by importin α isoforms 3 and 7. *Structure*, 2015. **23**(2): p.
521   374-384.

522   38.   Cook, A., et al., The structure of the nuclear export receptor Cse1 in its cytosolic state reveals
523   a closed conformation incompatible with cargo binding. *Mol. Cell*, 2005. **18**(3): p. 355-367.

524   39.   Renault, L., et al., The 1.7 Å crystal structure of the regulator of chromosome condensation
525   (RCC1) reveals a seven-bladed propeller. *Nature*, 1998. **392**(6671): p. 97-101.

526   40.   Hillig, R.C., et al., The crystal structure of rna1p: a new fold for a GTPase-activating protein.
527   *Mol. Cell*, 1999. **3**(6): p. 781-791.

528   41.   Bullock, T.L., D.W. Clarkson, H.M. Kent, and M. Stewart, The 1.6 Å resolution crystal
529   structure of nuclear transport factor 2 (NTF2). *Journal of molecular biology*, 1996. **260**(3): p.
530   422-431.

531   42.   Bennion Callister, J., S. Ryan, J. Sim, S. Rollinson, and S.M. Pickering-Brown, Modelling
532   C9orf72 dipeptide repeat proteins of a physiologically relevant size. *Hum. Mol. Genet.*, 2016.
533   **25**(23): p. 5069-5082.

534   43.   Swaminathan, A., et al., Expression of C9orf72-related dipeptides impairs motor function in a
535   vertebrate model. *Hum. Mol. Genet.*, 2018. **27**(10): p. 1754-1762.

536   44.   White, M.R., et al., C9orf72 Poly(PR) Dipeptide Repeats Disturb Biomolecular Phase
537   Separation and Disrupt Nucleolar Function. *Mol. Cell*, 2019. **74**(4): p. 713-728.e6.

538   45.   Haberland, J.r., J.r. Becker, and V. Gerke, The acidic C-terminal domain of rna1p is required
539   for the binding of Ran· GTP and for RanGAP activity. *J. Biol. Chem.*, 1997. **272**(39): p.
540   24717-24726.

541   46.   Seewald, M.J., C. Körner, A. Wittinghofer, and I.R. Vetter, RanGAP mediates GTP
542   hydrolysis without an arginine finger. *Nature*, 2002. **415**(6872): p. 662-666.

543   47.   Ghavami, A., L.M. Veenhoff, E. Van Der Giessen, and P.R. Onck, Probing the disordered
544   domain of the nuclear pore complex through coarse-grained molecular dynamics simulations.
545   *Biophysical Journal*, 2014. **107**(6): p. 1393-1402.

546   48.   Ananth, A.N., et al., Spatial structure of disordered proteins dictates conductance and
547   selectivity in nuclear pore complex mimics. *eLife*, 2018. **7**: p. 1-24.

548   49.   Fragasso, A., et al., A designer FG-Nup that reconstitutes the selective transport barrier of the
549   nuclear pore complex. *Nat Commun*, 2021. **12**(1): p. 2010.

550   50.   Ghavami, A., E. van der Giessen, and P.R. Onck, Coarse-Grained Potentials for Local
551   Interactions in Unfolded Proteins. *J. Chem. Theory Comput.*, 2013. **9**(1): p. 432-440.

552   51.   Ghavami, A., E. van der Giessen, and P.R. Onck, Energetics of Transport through the Nuclear
553   Pore Complex. *PLoS One*, 2016. **11**(2): p. e0148876-e0148876.

554   52.   Dekker, M., E. van der Giessen, and P.R. Onck, Liquid-liquid phase separation of intrinsically
555   disordered FG-Nups is driven by highly-dynamic hydrophobic FG-motifs. *bioRxiv*, 2022: p.
556   2022.09.20.508740.

557   53.   Jumper, J., et al., Highly accurate protein structure prediction with AlphaFold. *Nature*, 2021.
558   **596**(7873): p. 583-589.

559   54.   Varadi, M., et al., AlphaFold Protein Structure Database: massively expanding the structural
560   coverage of protein-sequence space with high-accuracy models. *Nucleic acids research*, 2022.
561   **50**(D1): p. D439-D444.

562   55.   Higurashi, M., T. Ishida, and K. Kinoshita, PiSite: a database of protein interaction sites using
563   multiple binding states in the PDB. *Nucleic Acids Res.*, 2009. **37**(Database issue): p. D360-4.

Author contributions

566 H.J., E.Vd.G., and P.R.O. designed research; H.J. performed and analyzed research; and H.J.,

567 E.Vd.G, and P.R.O. wrote the paper.