Reorganization of the Flagellum Scaffolding Induces a Sperm Standstill During Fertilization

Martina Jabloñski[1]; Guillermina M. Luque[1]; Matías D. Gómez-Elías[1]; Claudia Sanchez-Cardenas[2]; Xinran Xu[3]; Jose Luis de la Vega-Beltran[2]; Gabriel Corkidi[4]; Alejandro Linares[2,6]; Victor X. Abonza Amaro[2,6]; Aquetzalli Arenas-Hernandez[7]; María Del Pilar Ramos-Godinez[8]; Alejandro López-Saavedra[8,9]; Dario Krapf[5]; Diego Krapf[3]; Alberto Darszon[2]; Adan Guerrero[2,6*†]; Mariano G. Buffone[1*†].

[1]Instituto de Biología y Medicina Experimental, Consejo Nacional de Investigaciones Científicas y Técnicas (IBYME-CONICET), Buenos Aires, Argentina.
[2]Departamento de Genética del Desarrollo y Fisiología Molecular, Instituto de Biotecnología, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, Mexico.
[3]Department of Electrical and Computer Engineering and School of Biomedical Engineering, Colorado State University, Fort Collins, Colorado, USA.
[4]Laboratorio de Imágenes y Visión por Computadora, Instituto de Biotecnología, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, México.
[5]Instituto de Biología Molecular y Celular de Rosario (IBR), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Universidad Nacional de Rosario (UNR), Rosario, Argentina.
[6]Laboratorio Nacional de Microscopía Avanzada, Instituto de Biotecnología, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, Mexico.
[7]Departamento de Microscopía Electrónica, Instituto Nacional de Cancerología, Mexico City, Mexico.
[8]Unidad de Aplicaciones Avanzadas en Microscopía, Instituto Nacional de Cancerología, Unidad de Investigación Biomédica en Cáncer, UNAM. Mexico City, Mexico
[9]Tecnologico de Monterrey, Escuela de Medicina y Ciencias de la Salud. Mexico City, Mexico.

†, These authors contributed equally to this work.

Mariano G. Buffone Ph.D.
Instituto de Biología y Medicina Experimental, CONICET. Vuelta de Obligado 2490 (1428), Buenos Aires, Argentina. Tel: (+5411)4783-2869.
E-mail: mgbuffone@ibyme.conicet.gov.ar.

Adan Guerrero Ph.D.
Laboratorio Nacional de Microscopía Avanzada, Instituto de Biotecnología, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, México.
Email: adan.guerrero@ibt.unam.mx.

Abstract

Mammalian sperm delve into the female reproductive tract to fertilize the female gamete. The available information about how sperm regulate their motility during the final journey to the fertilization site is extremely limited. In this work, we investigated the structural and functional changes in the sperm flagellum after AE and during the interaction with the eggs. The evidence demonstrates that the double helix actin network surrounding the mitochondrial sheath of the midpiece undergoes structural changes prior to the motility cessation. This structural modification is accompanied by a decrease in diameter of the midpiece and is driven by intracellular calcium changes that occur concomitant with a reorganization of the actin helicoidal cortex. Midpiece contraction occurs in a subset of cells that undergo AE, live-cell imaging during in vitro fertilization showed that the midpiece contraction is required for motility cessation after fusion is initiated. These findings provide the first evidence of the F-actin network's role in regulating sperm motility, adapting its function to meet specific cellular requirements during fertilization, and highlighting the broader significance of understanding sperm motility.

Significant statement

In this work, we demonstrate that the helical structure of polymerized actin in the flagellum undergoes a rearrangement at the time of sperm-egg fusion. This process is driven by intracellular calcium and promotes a decrease in the sperm midpiece diameter as well as the arrest in motility, which is observed after the fusion process is initiated.

Introduction

Sperm motility is required for arrival at the site of fertilization and to penetrate the different layers surrounding the egg. The temporal regulation of sperm motility involves the concerted action of multiple cell structures to enable fertilization. The initial motility of ejaculated sperm is characterized by a linear progressive movement as the cells traverse a long distance within the female reproductive tract. However, at one point, before reaching the oocyte, this initial progressive motion must be changed to a vigorous non-progressive mode of motility called hyperactivation (*1*). During that migration, most of mouse sperm undergo acrosomal exocytosis (AE), which takes place in the upper segments of the oviduct, before the sperm directly interact with the egg or its surrounding layers (*2–4*). This

exocytic event is critical because proteins involved in this process are rearranged in
preparation for fusion(*5, 6*). A second dramatic change in motility is subsequently required
for efficient sperm-egg fusion. During this event, which has not been studied in detail so
far, sperm completely stop moving (*7–9*). The cease in sperm motility is considered as an
indicative marker of an effective fusion between gametes, given that it is necessary to
complete the attachment and fusion between sperm and eggs. Fusion itself is a complex
event mediated by several proteins identified using loss of function strategies (*10, 11*).
Nevertheless, the available information about how sperm regulate their motility during the
final journey to the fertilization site and during the interaction with the female gamete is
extremely limited. In this final journey, most sperm cells migrate to the ovulated eggs after
AE, and very little is known about the motility state of those sperm and about the molecular
mechanism in charge of the motility arrest.

The regulation of sperm motility is achieved   by balancing force transduction and
the mechanical properties of the flagellum. Both effects are governed by the flagellum
cytoskeleton, which consists of two major components: a microtubule-based axoneme,
located at the flagellum axis, and actin filaments around it (*12*). The midpiece and the
principal piece also contain the outer dense fibers and the fibrous sheath that have
traditionally been referred to as part of the sperm cytoskeleton. Recently it was found that
the three-dimensional organization of polymerized actin in the flagellum midpiece of
murine sperm forms a unique double helix arrangement accompanying mitochondria (*13*).
This spatial distribution does not extend into the principal piece, where actin is uniformly
distributed between the axoneme and the plasma membrane. Across all investigated cells,
this double helix consisted of exactly 87 gyres with a pitch of 244 ± 1 nm, yielding a total
midpiece length in mice of 21.2 ± 0.3 µm (*13*). Such accurate control of sizes is not typical
in live cells given that reactions are prone to stochastic effects and entropy leads to large
cell-to-cell fluctuations. Thus, the precise control of the actin helix comes at a substantial
regulatory cost for sperm cells. Nevertheless, the function of this specialized structure of
actin filaments is largely unknown.

In this work, we find definite evidence for the role of the helical actin structure in
the final stage of motility regulation, when sperm dramatically shift from being
hyperactivated to a practically immotile stage. We have investigated the structural changes
in the actin cytoskeleton of the sperm flagellum after AE and during the interaction with the
eggs. By using a combination of single-cell imaging and super-resolution microscopy

methods, our results demonstrate that the helical structure of polymerized actin undergoes a

radical structural change at the time of sperm-egg fusion. Further, we uncover that this

process is triggered by a substantial calcium ($[Ca^{2+}]_i$) influx. This actin-dependent signaling

pathway promotes a decrease of the sperm midpiece diameter and motility arrest, both of

which are found to be required to complete the fusion process during fertilization.

Results

AE promotes a cessation of motility in a subset of sperm cells.

After AE, sperm need to accomplish migration to the ampulla, reach unfertilized

eggs, penetrate the cumulus matrix and the zona pellucida and finally, undergo fusion with

the oocyte (*14*). All these processes share the need to modulate sperm motility. Little is

known about the regulation of motility after the occurrence of AE or prior to the fusion

with the oocyte.

To simultaneously monitor the acrosomal status and sperm motility in live cells,

transgenic mice whose sperm express enhanced green fluorescent protein (EGFP) in the

acrosome and red fluorescent protein (DsRed2) in the mitochondria were used (Figure 1A,

Supplementary movie S1). EGFP-DsRed2 sperm were immobilized at the head on laminin-

coated coverslips, while still allowing free flagellar movement (Figure 1B, Supplementary

movie S2). While the presence of the acrosome is monitored using the EGFP fluorescence

signal, motility was assessed through the movement of the midpiece (DsRed2

fluorescence), which was beating in and out of the imaging plane (Figure 1B).

The majority of acrosome-intact sperm were able to move. Interestingly, Figures 1B

and 1C show the coexistence of two populations of sperm that underwent AE (cells lacking

EGFP signal) induced upon addition of progesterone, a physiological trigger of AE. Some

of the acrosome reacted sperm moved normally (34.1±3.7 %, n = 235), whereas the

majority of them remained immotile (65.9 ± 6.2 %, n = 2350) (Figures 1B and 1C, upper

and lower panel, respectively, Supplementary movie S2).

The flagellar beat cycle encompasses a self-regulatory mechanism that receives

feedback from molecular and mechanical signals (*15*). The complete abortion of flagellar

beat cycle observed in Figures 1B and 1C might be indicative of a stimulus provided by or

occurring concomitant with AE. To comprehend the connection between AE and motility,

we hypothesized the existence of a mechanical change in the flagella as a result of the AE.

For this reason, AE was studied in the presence of FM4-64 fluorescent dye (*16*) to visualize

structural changes at the plasma membrane, which can occur at macroscopic scales observed as an alteration in the shape of the flagellum, or at mesoscopic scales, monitored by local changes in FM4-64 fluorescence occurring at the vicinity or within the plasma membrane. EGFP-DsRed2 sperm were stimulated with progesterone and their 'motility' behavior was recorded for 5 min in the presence of FM4-64. The loss of EGFP fluorescence in the acrosome of transgenic mice correlated with a noticeable increase of FM4-64 fluorescence in the head, as shown in Figure 1D (see panels 0´ and 3´, Supplementary movie S3). In addition, we noticed that sperm that underwent AE, later showed an increase in the FM4-64 fluorescence intensity in the flagellum as well (see panel 14´of Figure 1D, Supplementary movie S3).

A more detailed analysis of individual cells demonstrated changes in FM4-64 fluorescence according to their motility (Supplementary movie S4). Figure 1E shows a motile acrosome-intact sperm, (a control case), where the fluorescence levels of EGFP and FM4-64 remained constant during the entire experiment (Pattern I). Interestingly, sperm that lost their acrosome (no EGFP and high FM4-64 fluorescence in the sperm head) and continued moving, presented low levels of FM4-64 fluorescence in the midpiece (Figure 1F, Pattern II). On the other hand, acrosome-reacted sperm that remained immotile showed a significant rise in the FM4-64 midpiece fluorescence (Figure 1G, Pattern III). Noteworthy, both subsets of cells (with and without motility) displayed the expected increase in FM4-64 fluorescence in the sperm head.

To deepen our understanding on how sperm lose motility during AE, we designed a sperm tracking system that can also monitor changes in fluorescence in moving cells (Figure S1B and S1C). Three parameters were then assessed in real time, namely, the beat frequency of the flagellum (by tracking it on bright field images, or the DsRed2 channel, Figure S1A), the status of the acrosome (EGFP signal), and changes occurring at or within the membrane in the midpiece (FM4-64 signal). Figure S1B shows that sperm that did not undergo AE remained motile with a stable beat frequency and low FM4-64 fluorescence over the recording time. In stark contrast, after AE, cells gradually diminished the beating frequency until a complete arrest is observed, while a gradual increase in FM4-64 fluorescence in the midpiece is observed (Figure S1C).

To evaluate the relationship between between FM4-64 and viability, a vital dye Sytox Blue was used in imaging flow cytometry experiments. Fluorescence intensities in non-capacitated sperm stimulated with ionomycin (Figure S1D) were determined and two

populations with Sytox Blue signals were clearly distinguished (Sytox+ and Sytox-),
enabling the discernment between live and dead sperm. Interestingly, the upper right panels
(Sytox Blue+ / FM4-64 high) consistently show a positive correlation between FM4-64 and
Sytox Blue. Nonetheless, the lower panels (Sytox Blue-) show no correlation with FM4-64
fluorescence, indicating that this population can exhibit either low or high FM4-64
fluorescence. Single-cell examples are shown, where the four categories are represented:
dead sperm with low FM4-64 fluorescence (Sytox Blue + / FM4-64 low), dead sperm with
high FM4-64 fluorescence (Sytox Blue + / FM4-64 high), live sperm with low FM4-64
fluorescence (Sytox Blue - / FM4-64 low), and live sperm with high FM4-64 fluorescence
(Sytox Blue - / FM4-64 high). Therefore, while the FM4-64 signal alone is not a definitive
marker for either AE or cell death, it is crucial to use additional viability assessments, such
as Sytox Blue, to accurately differentiate between live and dead sperm in studies of
acrosome exocytosis and sperm motility. Cell viability was always considered, as any
imaged sperm was chosen based on motility, indicated by a beating flagellum. The
determination of whether selected sperm die during or after AE remains to be elucidated.
The results presented in Figure 1 and Supplementary Figure S1 show examples of motile
sperm that experience an increase in FM4-64 fluorescence.

Altogether, these experiments demonstrate that AE promotes a cease of motility in a
subset of sperm cells, which coincides with an increase in FM4-64 fluorescence in the
midpiece.

AE is followed by a decrease in the midpiece diameter.

Live-cell super-resolution microscopy was used to understand the structural changes
occurring in the midpiece. Figure 2A shows a wide-field image of a sperm midpiece stained
with FM4-64 (left) and analyzed using super-resolution radial fluctuations (SRRF, right)
(*17*). AE dynamics was monitored by FM4-64 fluorescence in the head (Figure 2B, upper
right insets). A decrease in the midpiece diameter over time was observed following AE
either being spontaneous, induced by a physiological agonist (progesterone), or induced by
a non-physiological surge of Ca^{2+} (ionomycin) (Figures 2B and 2C and Supplementary
movie S5). In the negative control (no AE), the diameter remained unchanged. This
phenomenon was also observed using mean shift super-resolution (MSSR) (*18*). To further
support this observation, other plasma membrane probes of different molecular structure,
such as Memglow 700 (Figure S2A), Bodipy-GM1 (Figure S2B), and FM1-43 (Figure

S2C), were used. In all three cases, a decrease in the midpiece diameter was observed after

AE.

The FM4-64 fluorescence intensity in the head and in the midpiece (Figures 2D and

2E) were also assessed. As expected, the increase in FM4-64 fluorescence in the head

occurs after AE. Furthermore, in agreement to what was observed in Figure 1, fluorescence

intensity in the midpiece also increased over time in sperm that underwent AE (Figures 2D

and 2E).

To confirm these results, two alternative methods to visualize the change in sperm

midpiece diameter were used. In neither of them, a membrane dye was used. First, indirect

immunofluorescence to detect a membrane protein (GLUT3) was performed. As shown in

Figure 2G, a decrease in the midpiece diameter was also observed. Second, scanning

electron microscopy (SEM) was used to evaluate the midpiece in acrosome-intact or

reacted sperm (Figure 2F). The overall diameter of the midpiece in acrosome-intact sperm

was larger than in acrosome-reacted sperm, with measurements of 0.731 ± 0.008 μm and

0.694 ± 0.007 μm, respectively . Overall, we have confirmed using three different

approaches that AE is followed by a midpiece diameter.

The contraction of the midpiece initiates in the proximal part of the flagellum.

To investigate whether the contraction of the midpiece is triggered at a random

location or in a particular region of the flagellum, kymographs were used. A kymograph

allows visualization of dynamical aspects of a given phenomenon in a single figure, where

the temporal dimension is expressed as an axis of the image through a spatial-temporal

transformation of the dataset. Two types of kymographs were used (Figure 3A). First,

super-resolution microscopy kymographs were built to monitor dynamic changes in the

diameter of the flagellum along the midpiece. Second, to investigate the number of

contraction sites, fluorescence kymographs were made from diffraction-limited images to

observe changes in FM4-64 fluorescence over time and across the entire midpiece.

Interestingly, Figure S2D shows a significant negative correlation between the midpiece

diameter and the FM4-64 fluorescence, unveiling a tool to obtain an approximate value of

the midpiece diameter without the need of super-resolution microscopy.

Three experimental subsets that underwent AE were assessed: spontaneous, induced

by progesterone, and induced by ionomycin. To scrutinize randomness of focus-driven

contraction, the midpiece was segmented in three regions (Figure 3A): proximal (near the

neck), central, and distal (near the annulus). In these groups, it was then evaluated whether

cells presented one or more foci of contraction.

243   Figure 3B shows a cumulative study, which indicates that contraction is

preferentially initiated at the proximal part of the midpiece, regardless of being

progesterone-induced or spontaneous AE. When ionomycin was used as an agonist of AE,

the contraction began randomly in any segment of the midpiece. Figures 3C and 3D show

two examples of flagellar dynamics for progesterone-induced sperm, which either

presented one or two foci of contraction. In both cases, the diameter decreased along the

whole midpiece, seen as a transition from dark to bright red colors in Figures 3C and 3D.

Noteworthy, focal points of contraction were consistently observed: i.e., a case with a

single focus (Figure 3C), or a case with two foci of contraction (Figure 3D). Figures S2E

and S2F show ionomycin-induced sperm that present one or more foci. It is observed in the

normalized diameter kymograph that the flagellum presents a single focus between the

measurements of 7.5 µm and 10 µm (Figure S2E, left panel). This is reflected in the right

panel in the light blue and orange dots, which are the actual diameter measurements that

decreased first and most (Figure S2E). A representative behavior within this experimental

group is represented in Figure S2F, where the entire midpiece contracted (all its sections

simultaneously). This is observed in the color scale of the normalized diameter kymograph

and coincides with the real measurements of the diameter in the right panel, which

decreased their value. Lastly, Figures S2G and S2H show examples of spontaneous AE

with one or two foci of contraction, respectively. In particular, in Figure S2H, a first focus

of contraction between 0 and 5 µm that coincided with the diameter measurements of the

right panel that decreased their value first and to a greater extent (black, pink and green

dots). Although the entire midpiece contracted, a second focus of contraction appeared

around the measurements of 10 to 15 µm and coincidentally, these are the points with the

strongest decrease in the diameter (orange, purple and red points).

The reduction of the midpiece diameter occurs concomitantly with a $[Ca^{2+}]_i$ increase

in the flagellum.

270   It is widely accepted that an increase in $[Ca^{2+}]_i$ precedes AE (*16, 19*). It has been

shown that AE initiation by progesterone occurs after an increase in $[Ca^{2+}]_i$ in the sperm

head that is later propagated towards the midpiece (*19*). To comprehend the connection

between AE and the concomitant contraction of the midpiece, we hypothesized the

existence of a signal, i.e., Ca^{2+}, which propagates from the head to the flagellum and
modulate the architecture of the sperm flagellum.

To investigate if the increase in the concentration of $[Ca^{2+}]_i$ in the midpiece is
correlated with the midpiece contraction, sperm were incubated with FM4-64 and Fluo4, a
Ca^{2+} fluorescent sensor (Supplementary movie S6). Figure 4A shows a live-cell imaging
experiment of sperm cells that did not undergo AE (control case), nor a significant rise in
$[Ca^{2+}]_i$. Noteworthy, upon induction with progesterone (Figure 4C, Supplementary movie
S6), there was an increase in the midpiece $[Ca^{2+}]_i$ followed by an increase in FM4-64
fluorescence, which is indicative of a contraction of the flagellum (see Figure S2D). These
single cell experiments illustrate a reduction of the midpiece diameter that occurs
concomitantly with an increase in $[Ca^{2+}]_i$ within the flagellum. A large population of sperm
was analyzed using low magnification (10X). Figures 4B and 4D show a collection of
single cell kymographs, where each row represents the fluorescence of Fluo4 (green) and
FM4-64 (red) of the midpiece over time. Unstimulated cells are displayed on Figure 4B and
sperm that were exposed to progesterone are shown in Figure 4D. Most of the cells display
a transient $[Ca^{2+}]_i$ increase followed by an increase in FM4-64 fluorescence. This pattern
was also observed in ionomycin-induced cells (Figures S3A and S3B and Supplementary
movie S6) with the exception that, as expected, in this case the rise in $[Ca^{2+}]_I$ was faster and
sustained. These findings indicate that an $[Ca^{2+}]_i$ transient increase, happening in the
midpiece, precedes the contraction of the flagellum.

The relationship between the observed $[Ca^{2+}]_i$ rise and the flagellar contraction was
then assessed using live-cell super-resolution imaging. Figure 4E shows there is a $[Ca^{2+}]_i$
increase in the midpiece that coincides in space and time with a contraction of the
midpiece. To provide insight in the spatio-temporal relationship of the $[Ca^{2+}]_i$ rise and the
midpiece contraction, these results were visualized using a 3D kymograph encompassing
the dynamics happening along the entire midpiece (Figure 4F). Overall, a rise in $[Ca^{2+}]_i$
occurred along the whole midpiece, which coincided with a generalized contraction of the
midpiece (Figure 4F). Remarkably, a transient focal increase of $[Ca^{2+}]_i$ was observed at the
base of the flagellum, which correlates in space and time with a focal reduction of the
diameter (Figure 4F). We then sought to identify a molecular/structural link between both
processes.

The distance between t   he actin cytoskeleton and   the plasma membrane is decreased during the contraction of the midpiece.

The midpiece is shaped by two major structural elements: the axoneme, which consists of tubulin and its accessory components, including the outer dense fibers (ODFs); and a recently described network of filamentous actin (F-actin) arranged in a helicoidal conformation along the midpiece (*13*). We investigated whether the actin network plays a role in regulating the midpiece contraction associated with AE. We considered three possible scenarios: (i) an unknown mechanical force, mediated by a rigid matrix such as F-actin, brings the plasma membrane closer to the center of the flagellum; (ii) a dynamic relationship exists between the plasma membrane and the actin cytoskeleton of the midpiece, facilitating contraction; and (iii) the midpiece contraction is driven by an uncharacterized signaling mechanism that acts independently of the actin cytoskeleton.

The role of the actin cytoskeleton in the midpiece contraction was investigated through live-cell super-resolution using SiR-actin, a fluorescent probe that binds to F-actin (*20–22*). We used FM4-64 to visualize the plasma membrane (Figure 5) and transgenic sperm expressing DsRed2 to observe the mitochondrial network (Figure S4). Two stimulated sperm, one that experienced AE (Figure 5B) and another that did not (Figure 5A) are shown. In both cases, a network of F-actin was observed beneath the plasma membrane of the midpiece (cortical actin), enveloping the mitochondrial network (Figure S4A). As expected, in the absence of AE, the midpiece diameter (observed through FM4-64) remained unchanged (see also Figure 2, and Figure S4). In this case, both the diameter of the F-actin network and its proximity to either the plasma membrane or the mitochondrial network remained static (Figure 5A), suggesting a structural role for the F-actin network within the midpiece (scenario - (i)), such as supporting the organization of the mitochondrial network (Figure S4). Remarkably, acrosome-reacted sperm (Figure 5B) experienced an abrupt structural reorganization of the midpiece, characterized by a decrease in the distance between the plasma membrane and both the F-actin and mitochondrial networks (Figures S4 and S5). This observation suggests that the remodeling of the midpiece structure is driven by a mechanical (or molecular) interaction occurring at the boundaries of the plasma membrane (supporting scenario - (ii)).

To assess the potential interaction between the F-actin network and the plasma membrane (or its associated components), we examined their colocalization through live-cell super-resolution imaging. Figure 5C presents two Manders' colocalization coefficients

(*23*). M1 represents the proportion of F-actin (in pixels) that colocalizes with the plasma membrane, while M2 measures the proportion of plasma membrane pixels colocalizing with the F-actin network. A Manders´ value of 1 indicates full colocalization, while a value of 0 implies no colocalization.

Figure 5C reveals that acrosome-intact sperm displayed low Manders' coefficients (M1= 0.32 ± 0.03; M2= 0.28 ± 0.04). In contrast, sperm undergoing midpiece contraction exhibited a significant increase in colocalization between the F-actin network and the plasma membrane, as evidenced by the rise in M1 and M2 coefficients (M1= 0.65 ± 0.03; M2 = 0.69 ± 0.03). This increased colocalization occurred simultaneously with midpiece contraction.

Overall, these results confirm that (i) the contraction of the midpiece is linked to the remodeling of the plasma membrane and potentially the F-actin cytoskeleton, and that (ii) both structures interact with each other, directly or indirectly, at the nanoscales. We then seek to understand whether the driving force for the contraction of the midpiece emanates from the F-actin network. The following scenarios were envisaged: (a) the plasma membrane moves towards the actin cytoskeleton, (b) the actin cytoskeleton moves towards the plasma membrane, or (c) both structures come closer to the center of the flagellum. To investigate the occurrence of any of these scenarios, the positions of fluorescence peaks of FM4-64 and SiR-actin in super-resolution kymographs were tracked, as proxy of either membrane or actin cytoskeleton localization, respectively (Figures S5A and S5B). Figure S5C shows representative histograms of SiR-actin and FM4-64 over time. Both FM4-64 and SiR-actin fluorescence peaks came closer to each other. This effect can be also seen in Figure S5F, where acrosome-intact cells display a distance of 0.180 ± 0.007 μm between the plasma membrane and the actin cytoskeleton. In acrosome-reacted cells, the distance was as small as 0.074 ± 0.006 μm after midpiece contraction.

In Figure S5D, a section of the midpiece of an ionomycin-induced AE sperm is shown. In this case, the plasma membrane and the actin cytoskeleton approach move closer toward the center of the cell. (set to 0). The slope calculated from the linear fit denotes the velocity of this change. In this example, the plasma membrane is contracting at a rate of 14 μm/min, whereas the actin cytoskeleton is contracting at a rate of 3 μm/min (Figure S5E). In both cases, the signal presents a negative slope in sperm that underwent AE, consistent with a decrease in midpiece diameter. Since the results of the analysis of SiR-actin slopes were not conclusive, we studied the actin cytoskeleton structure in more detail.

The actin cytoskeleton in the midpiece is remodeled during the AE.

In the midpiece, polymerized actin forms a double helix that accompanies mitochondria (*13*). To investigate if the contraction of the midpiece is associated with structural modifications of the actin double-helix, 3D-STORM was used. Cells were exposed to progesterone, fixed, and co-stained with PNA and phalloidin to visualize the acrosomal status and the actin structure, respectively.

Figures 5D and 5F show representative images of both dyes for acrosome-intact and acrosome-reacted sperm (left panels). Different parameters were calculated: 1) Helical pitch (*l*), which is the distance between turns of the helix; 2) Frequency ($f_0 = 1/l$), obtained from the Fourier transform of the image and represented the number of turns that the helix makes per unit length; and 3) Radial distribution (*R*), to infer the distance between the center of the midpiece and the maximum of fluorescence (*13*). The frequency increased substantially in the acrosome-reacted sperm compared to intact sperm (6.29 vs 4.04 μm^{-1}, Figures 5E and 5G, red arrows in left panels). In addition, the helical pitch diminished its magnitude in acrosome-reacted sperm (159 nm vs 248 nm) (Figures 5E and 5G, left panels). The radial distribution of F-actin in cells that underwent AE indicated a smaller radius compared to acrosome-intact sperm (252 vs 300 nm, Figures 5E and 5G, right panel). Altogether, these results confirm that the actin double-helix undergoes structural changes during the contraction of the midpiece

To further investigate the structural rearrangements of the actin cytoskeleton during AE, a fluorescent molecule number and brightness analysis was performed (*24–26*). The number measure indicates the abundance or concentration of SiR-Actin molecules bound to F-actin fibers, while the brightness analysis helps reveal dynamic changes in molecular aggregation processes. This analysis is based on the idea that higher-order fluorescent complexes, which are mobile within the sample, will cause an increase in signal variability over time. An increase in brightness suggests the formation or movement of supramolecular structures, such as bundles of actin fibers bearing SiR-Actin. AE induction with progesterone and ionomycin resulted in a significant increase in actin number and brightness compared to the control (non-reacted) group (Figure S6A). Spontaneously reacted sperm also exhibited this behavior, albeit in a more moderate fashion (Figure S6A).

The observed increase in SiR-Actin number after AE induction suggests major actin monomer recruitment to the polymerizing fibers (Figure S6A and S6B), which produces an

apparent increase in local fluorophore concentration (Figure S6B). This effect positively correlated with a similar increase in SiR-actin fluorescence (Figure S6A), which is known to indicate actin polymerization due to its high affinity for F-actin.

Actin cytoskeleton polymerization in the midpiece results in a signal variation of increased magnitude (an increase in SiR-Actin brightness) as it undergoes structural rearrangements (Figures S6A and S6B), which are compatible with other observations reported in this work. Both assembly and transport of actin complexes into higher oligomeric states (F-actin) across the cellular milieu led to an apparent increase in the registered SiR-Actin brightness (Figure S6B). Additionally, actin filament displacement along the mitochondrial sheath during AE provides a SiR-Actin brightness measure with high dispersion, indicating a redistribution of this cellular structure. These results suggest that actin filaments are locally redistributed and remodeled during AE (Figure S6B).

~~Occurrence of~~ **Midpiece contraction in sperm located within the perivitelline space.**

In mice, only sperm undergoing AE prior to binding to the zona pellucida can penetrate and fertilize (*27*). We hypothesized that midpiece changes and motility cessation occur only after acrosome-reacted sperm penetrate the zona pellucida. Live imaging was performed after *in vitro* fertilization (IVF) using transgenic EGFP-DsRed2 sperm loaded with FM4-64. Eggs were inseminated with a high sperm count (200,000 cells) to increase the number of cells observed.

Figure 6A and Supplementary movie S7 show a sperm swimming within the perivitelline space. The FM4-64 fluorescence of this sperm midpiece is low (Figure 6B) which coincides with the fact that the sperm is moving (see also Figure 1F-G). Another example of this observation is shown in Figure 6C-D, an acrosome-reacted moving sperm within the perivitelline space had low FM4-64 fluorescence in the midpiece (Figure 6C). After 20 minutes, these sperm stopped moving and exhibited increased FM4-64 fluorescence, indicating midpiece contraction (Figure 6D). These results suggest that midpiece contraction and motility cessation occur after acrosome-reacted sperm penetrate the zona pellucida.

Midpiece contraction takes place following sperm-egg fusion.

Motile acrosome-reacted sperm, initially lacking midpiece contraction, can penetrate the zona pellucida. However, they eventually stop moving and exhibit an increase

in FM4-64 fluorescence. This suggests that the cessation of motility plays a crucial role in

fertilization events occurring after zona pellucida binding and penetration, and that a

similar change in midpiece architecture is necessary for successful sperm-egg fusion. To

explore this hypothesis, we designed a live imaging experiment using zona-free eggs.

Denuded oocytes were loaded with Hoechst 33342, a nuclear dye, to visualize the exact

moment of sperm-egg fusion (*28–30*). The experiment involved FM4-64, transgenic EGFP-

DsRed2 sperm, and simultaneous signal collection from five separate channels (DIC,

Hoechst, EGFP, DsRed2, and FM4-64). To enhance observation likelihood, volumetric

data was collected by imaging at different z-planes (Figure 7A).

Figure 7C presents a representative volumetric time series of sperm-egg fusion, with

only the optimal focal plane shown. The acrosome-reacted sperm (EGFP negative) initially

did not display Hoechst fluorescence (Supplementary movie S8   ), indicating a lack of

fusion. However, 14 minutes later, the dye entered the sperm and stained the nucleus,

indicating fusion initiation. At this stage, sperm remain motile while bound to the egg

plasma membrane. The midpiece diameter remained unchanged during initial sperm-egg

fusion indicated by the low FM4-64 fluorescence. Following gamete fusion initiation,

FM4-64 fluorescence increased (38 to 44 min), causing sperm motility to cease. In some

instances, the midpiece folded and extended again (Figure S7B). These findings

demonstrate that midpiece contraction occurs following sperm-egg fusion.

A decrease in $[Ca^{2+}]_i$ in the midpiece following fusion precedes the midpiece

contraction.

In previous experiments, an increase in $[Ca^{2+}]_i$ was observed throughout the

midpiece, coinciding with midpiece contraction (Figure 4 and Figure S3). To examine the

mechanism behind midpiece contraction after fusion, wild-type sperm loaded with Fluo4

were exposed to denuded oocytes loaded with Hoechst. We hypothesized that dynamic

changes in $[Ca^{2+}]_i$ during gamete fusion drive the midpiece contraction during sperm

immobilization.

Figure 8A presents representative images from a time course live imaging

experiment designed to observe sperm-egg fusion. The experiment involved simultaneous

signal collection from four separate channels over time (DIC, Hoechst, Fluo4, and FM4-

64). To enhance observation likelihood, images were taken at different z-planes, though

only the optimal focal plane is shown.

Unexpectedly, sperm binding to the egg plasma membrane displayed high levels of

$[Ca^{2+}]_i$ in the head and midpiece (Figure 8A, time 1 min, and Supplementary movie S9   ).

A few minutes after fusion began (indicated by an increase in Hoechst signal in the sperm),

a decrease in $[Ca^{2+}]_i$ in the head and midpiece was observed (21 min to 51 min). This

decrease in $[Ca^{2+}+]_i$ was followed by midpiece contraction, as evidenced by the increase in

FM4-64 fluorescence (56 min). In several cells, the midpiece folded back before

contracting. The gametes that underwent this change in the flagellum typically remained

folded, but occasionally the tail unfolded again. Figures 8B-F quantify the key events

described above during sperm-egg fusion.

In summary, 75.88   ± 12.20 % of the sperm exhibited midpiece contraction

upon fusion. In 61.83   ± 12.04 % of cases, the midpiece folded on itself, and afterward,

only 31.60   ± 7.47   % unfolded and stretched out again. All sperm bound to the plasma

membrane before fusion presented high $[Ca^{2+}]i$ in the head, and the majority displayed a

decrease in head $[Ca^{2+}]_i$ before midpiece contraction (75.13 ± 3.05 %). Concerning the

midpiece, three patterns were observed during fusion: in the majority of cases (69.67   ±

11.46 %), the midpiece experienced a decrease in $[Ca^{2+}]_i$. Additionally, 10.33 ± 10.33 % of

sperm showed a transient increase in $[Ca^{2+}]_i$, while the remaining 20.00 ± 9.41 % displayed

no changes.

Collectively, these results indicate that a decrease in $[Ca^{2+}]_i$ in the midpiece after

fusion precedes midpiece contraction and the cessation of sperm motility that precedes

sperm-egg fusion.

Discussion

In this article, we demonstrate the existence of a fundamental structural change that

occurs in the sperm flagellum at the time of fusion with the egg. Using a plethora of

advanced microscopy methods and single cell imaging, we provide insight about cellular

and molecular events that occur in acrosome-reacted sperm, which, undoubtedly, are the

ones capable of fertilizing an oocyte (*2*, *3*). At this precise moment of the fertilization

process, sperm need to stop moving to complete the fusion between the two gametes. The

cease of movement is caused by two concomitant processes that take place in the flagellar

midpiece region: a change in the F-actin helical structure and a decrease in the midpiece

diameter. To the best of our knowledge, this is the first time that a structural modification

of the sperm flagellum related to a specific cellular necessity, i.e., gamete fusion, is described.

To arrive at the site of fertilization within the female reproductive tract, sperm motility is required. It is also fundamental to penetrate the different layers surrounding the egg. In this journey, sperm sense the environment and adapt their movement to the different physiological scenarios. In addition, recent evidence also demonstrated that after being released from their site of storage in the oviductal isthmus, mouse sperm undergo AE (*2, 3*). This process occurs in the upper segments of the oviduct before any interaction with the eggs or their surrounding layers. This observation opened a new scenario about the motility of sperm in their last transit to the site of fertilization. Little is known about sperm in that region and how sperm move after AE. In this regard, most of the information about mammalian sperm comes from studies conducted *in vitro*, using a mixture of acrosome-intact and acrosome-reacted sperm. Even if their motility is analyzed in a subjective manner or using cell tracking systems, virtually all the experiments are conducted without discerning the acrosomal status of the cells. Another possible source of artifacts in this analysis is related with the fact that most of the experiments are also performed using epididymal sperm (not ejaculated) in aqueous solutions that support sperm capacitation but do not represent the natural viscous environment present in the female tract.

By performing in vitro experiments, we detected a strong association between the decrease in the midpiece diameter and the cease of sperm motility. This is observed in a subset of sperm after the occurrence of AE. This may suggest that only sperm that undergo AE in the oviduct and do not experience this midpiece contraction are capable of migrating to the ampulla and penetrate the cumulus and the zona pellucida. Those sperm that undergo this phenomenon earlier in the tract may not be suitable to continue their journey suggesting that this also may select the gametes during their transit to the ampulla. Previous observations tracking sperm within the female tract have shown the existence of acrosome-reacted sperm within the tract that remain non-motile (*2–4*).

Acrosome-reacted sperm that bind and penetrate the zona pellucida are ready to fuse with the egg. During sperm-egg fusion, several authors have reported in different species that sperm stop moving (*7–9*). However, the mechanism behind this behavior is not established. One possible explanation is that fusion promotes ion transport changes in sperm, which significantly alter flagellar movement. In this sense, it was previously demonstrated that certain ions such as Ca^{2+} may diffuse from the oocyte to the sperm (*31*).

However, a clear technical limitation in our experimental approach is that the probes that
monitor ion dynamics may be exchanged between both gametes. If the concentration of a
given probe does not remain stable, it is impossible to determine the accurate change that
occurs during fusion. Future experiments may take advantage of transgenic models that
incorporate a particular sensor to study this process in vivo, such as the one used by Cohen
and collaborators (*32*).

Regardless of the precise nature of ion exchange between sperm and eggs, the
diameter of the sperm flagellum in the midpiece is reduced. This midpiece diameter
reduction is strongly associated with cessation of sperm motility and is apparently needed
to complete the fusion process. Our observation clearly supports this notion. Importantly,
the sperm flagellum folds back during fusion coincident with the decrease in the midpiece
diameter. Interestingly, this was previously observed in mammals as well as in sea urchin
sperm (*7*, *33*, *34*).

Like other cylindrical biological structures, the sperm flagellum relies on the
cytoskeleton for its structural organization and specialized mechanical properties. In
addition to the change in midpiece diameter, a significant rearrangement of the F-actin
cytoskeleton also occurs. In the midpiece, the polymerized actin is organized in a double
helix accompanying the mitochondria (*13*). As in many other organisms, the actin
cytoskeleton possesses important structural functions, and dynamic changes of F-actin
allow the cells to conduct important cellular tasks such as exocytosis. However, less is
known about the structural changes undergone by the actin cytoskeletons in cilia and
flagella. Remarkably, it has been reported that actin forms helix-like structures in the
flagellum of the parasite *Giardia intestinalis* (*35*). These findings open the question of
whether, to some extent, flagellar helical structures are conserved among diverse species.
Our observations demonstrate that the F-actin double helix undergoes a change in the
helical pitch as well as in the radial distance to the axoneme. This is concomitant with the
decrease in midpiece diameter. Our single-cell experiments using super-resolution
microscopy also revealed that the plasma membrane approached the F-actin network during
this change. It is well established that various proteins can function as linkers between the
plasma membrane and the actin cytoskeleton (*36*), but their roles in this specific process
remain to be studied. Regardless of how these structures are connected, it is evident that
both are associated. However, our experimental data cannot determine whether the plasma

membrane is causing the change of the actin network or if the actin network influences the

plasma membrane.

Another observation emerging from our study is that a change in $[Ca^{2+}]_i$ occurs prior

to the midpiece contraction. It is well known that Ca^{2+} is important for AE, and a specific

transient rise in $[Ca^{2+}]_i$ originating in the head can trigger exocytosis (*19*). Previous

observations have demonstrated that the sperm head and tail are not isolated compartments,

and that ions and other molecules can move between them (*37*, *38*). In our single-cell

experiments, we observed that the rise in FM4-64 fluorescence in the midpiece occurs after

the increase in $[Ca^{2+}]_i$ in that region, suggesting a potential association between these

events. This could involve diffusion or active transport processes; further investigation is

required to determine the precise mechanism to demonstrate if the structural changes are

triggered by Ca2+. In addition, $[Ca^{2+}]_i$ increase and/or modification in the midpiece

architecture may result in functional changes in the mitochondria such as the status of the

mitochondrial membrane potential and the ATP production. This possibility needs to be

further explored.

The same phenomenon was studied in sperm bound to the egg plasma membrane to

evaluate if the rise in $[Ca^{2+}]_i$ also occurs at the time of fusion. Remarkably, we noticed that

most of the bound sperm that ended up fusing with the eggs displayed high levels of $[Ca^{2+}]_i$

in both the head and the flagellum. The midpiece contraction and the immobilization

occurred when the levels of $[Ca^{2+}]_i$ went down in the midpiece suggesting a possible

connection between both events. However, our experimental approach limits the

interpretation of this result. One possible explanation is that as soon as sperm bind to the

plasma membrane of the oocyte, there is a rapid increase in $[Ca^{2+}]_i$ in the sperm. On the

other hand, a massive transport of Ca^{2+} from the egg to the sperm could also occur. These

hypotheses, however, are hindered by the technical limitations mentioned above. As the

oocyte is not loaded with Fluo4, we cannot rule out that the apparent $[Ca^{2+}]_i$ decrease seen

in these experiments is due to dye diffusion into the oocyte. Either way, it is clear that the

transport of this ion into or out of the sperm is key to cease motility at this fundamental step

of fertilization (*16*). Additionally, these results demonstrate that only sperm with elevated

$[Ca^{2+}]_i$ are capable of binding to the eggs. All these possible scenarios need to be

determined in future experiments. Thus, regardless of the mechanism, a clear change in

$[Ca^{2+}]_i$ dynamics is observed at the time of midpiece contraction. Future studies will aid to

indicate if certain Ca^{2+}-dependent proteins that can modify the actin cytoskeleton are

responsible for this change.

Why sperm stop moving: We propose three possible hypotheses. The cessation of

sperm motility can be attributed to the simultaneous or not occurrence of various events. 1)

a rapid increase in $[Ca^{2+}]_i$ levels may trigger the activation of Ca^{2+} pumps within the

flagellum. This process consumes local ATP levels, disrupting glycolysis in the process. 2)

Reorganization of the actin cytoskeleton: alterations in the actin cytoskeleton can lead to

changes in the mechanical properties of the flagellum, impacting its ability to move

effectively. 3) Midpiece contraction: Contraction in the midpiece region can potentially

interfere with mitochondrial function, thereby impeding the energy production necessary

for sustained motility. In addition, we speculate that the folding of the flagellum during

fusion further facilitates sperm immobilization because it makes it more difficult for the

flagellum to beat. Such process can enhance stability and increase the probability of fusion

success. Mechanistically, the folding may occur as a consequence of the deformation-

induced stress that develops during the decrease of midpiece diameter.

In cilia, apart from the findings presented in this paper, nothing is known about the

regulation of motility by actin. Actin has been found to participate in ciliogenesis, but its

involvement in active motility regulation has not been reported. This highlights a

potentially unique role for the actin cytoskeleton in regulating sperm function during

fertilization. Our discovery of actin's dynamic reorganization in sperm suggests it could

have a more active role in regulating motility and other functions. Given the conservation

of cilia and flagella across various organisms, our discovery could prompt further research

on the role of the actin network in these structures.

In conclusion, we demonstrate that sperm undergo a structural reorganization of the

actin cytoskeleton during key events of fertilization, as summarized in the working model

shown in Figure 9. Our findings introduce a new aspect of study in reproductive biology.

Previous research has mainly focused on identifying proteins essential for sperm-egg

fusion. Our results reveal a previously unexplored biological mechanism in mammalian

fertilization, opening new avenues for contraceptive method development.

Materials and methods

Detailed methods and protocols are provided in the SI appendix. For sample details,

optical equipment, imaging conditions and probes used, see Supplementary Table S1.

Acknowledgments

We gratefully acknowledge the financial support provided by the Williams and

Rene Baron Foundations, the Male Contraceptive Initiative (MCI), and the Chan-

Zuckerberg Initiative (CZI) for this work. Our sincere thanks go to Dr. Pablo Visconti for

his valuable insights throughout the course of this project. We also thank Yoloxochitl

Sánchez-Guevara for her technical support. This paper is dedicated to the memory of OAP.

Funding

This work was supported by: Chan Zuckerberg Initiative (2021-240504 to MB and

AG, GBI-0000000093 to AG, 2022-252509 to AG); Agencia Nacional de Promoción

Científica y Tecnológica (PICT, 2017-3047, 2018-1988 and 2020-00988); Dirección

General de Asuntos del Personal Académico/Universidad Nacional Autónoma de México

(DGAPA/UNAM grant: IN105222 to GK, IN211821 to AG and IN200919 to AD);

National Institute of Health (R01HD380882 to AD, R01HD106968 to DK and MGB).

Author contributions

Investigation: MJ, GML, MDGE, CSC, JLDLVB, AL, AAH, MPRG, ALS. Data

curation: MJ and VXAA. Writing: MJ, MGB and AG. Methodology: XX, Diego K and

GK. Conceptualization: Dario K, Diego K, AD, MGB and AG. Supervision: MGB and AG.

Competing interests

The authors declare no competing interests.

Data, Materials, and Software Availability

Data acquired and analyzed during this study are included in this manuscript or will

be available at Zenodo. Github will be used for depositing the code use in analysis.

References

1.   R. P. Demott, S. S. Suarez, Hyperactivated sperm progress in the mouse oviduct.
Biol Reprod **46**, 779–785 (1992).

2.   T. Hino, Y. Muro, M. Tamura-Nakano, M. Okabe, H. Tateno, R. Yanagimachi, The
Behavior and Acrosomal Status of Mouse Spermatozoa In Vitro, and Within the
Oviduct During Fertilization after Natural Mating. *Biol Reprod* **95**, 50–50 (2016).

3.   F. A. La Spina, L. C. Puga, A. Romarowski, A. M. Vitale, T. L. Falzone, D. Krapf, N.
Hirohashi, M. G. Buffone, Mouse sperm begin to undergo AE in the upper isthmus
of the oviduct. *Dev Biol* **411**, 172–182 (2016).

4.   Y. Muro, H. Hasuwa, A. Isotani, H. Miyata, K. Yamagata, M. Ikawa, R. Yanagimachi,
M. Okabe, Behavior of Mouse Spermatozoa in the Female Reproductive Tract from
Soon after Mating to the Beginning of Fertilization. *Biol Reprod* **94**, 80–80 (2016).

5.   N. Inoue, M. Ikawa, A. Isotani, M. Okabe, The immunoglobulin superfamily protein
Izumo is required for sperm to fuse with eggs. *Nature* **434**, 234–238 (2005).

6.   T. Noda, Y. Lu, Y. Fujihara, S. Oura, T. Koyano, S. Kobayashi, M. M. Matzuk, M.
Ikawa, Sperm proteins SOF1, TMEM95, and SPACA6 are required for sperm-
oocyte fusion in mice. *Proc Natl Acad Sci U S A* **117** (2020).

7.   P. Gaddum-Rosse, R. J. Blandau, L. B. Langley, D. E. Battaglia, In vitro fertilization
in the rat: observations on living eggs. *Fertil Steril* **42**, 285–292 (1984).

8.   P. Gaddum-Rosse, R. J. Blandau, L. B. Langley, K. Sato, Sperm tail entry into the
mouse egg in vitro. *Gamete Res* **6**, 215–223 (1982).

9.   B. Ravaux, N. Garroum, E. Perez, H. Willaime, C. Gourier, A specific flagellum
beating mode for inducing fusion in mammalian fertilization and kinetics of sperm
internalization. *Sci Rep* **6** (2016).

10. V. E. Deneke, A. Pauli, The Fertilization Enigma: How Sperm and Egg Fuse. *Annu.
Rev. Cell Dev. Biol. 2021* **37**, 391–414 (2021).

11. K. K. Siu, V. H. B. Serrão, A. Ziyyat, J. E. Lee, The cell biology of fertilization:
Gamete attachment and fusion. *J Cell Biol* **220** (2021).

12. H. Otani, O. Tanaka, K. Kasai, T. Yoshioka, Development of mitochondrial helical
sheath in the middle piece of the mouse spermatid tail: regular dispositions and
synchronized changes. *Anat Rec* **222**, 26–33 (1988).

13. M. G. Gervasi, X. Xu, B. Carbajal-Gonzalez, M. G. Buffone, P. E. Visconti, D. Krapf,
The actin cytoskeleton of the mouse sperm flagellum is organized in a helical
structure. *J Cell Sci* **131**, jcs.215897 (2018).

14. R. Yanagimachi, *Mammalian Fertilization* (Raven Press, New York, 1994).

15. K. Inaba, K. Shiba, Microscopic analysis of sperm movement: links to mechanisms
and protein components. *Microscopy* **67**, 144–155 (2018).

16. C. Sanchez-Cardenas, M. R. Servin-Vences, O. Jose, C. L. Trevino, A. Hernandez-
Cruz, A. Darszon, Acrosome Reaction and Ca2+ Imaging in Single Human
Spermatozoa: New Regulatory Roles of [Ca2+]i. *Biol Reprod* **91**, 67–67 (2014).

17. N. Gustafsson, S. Culley, G. Ashdown, D. M. Owen, P. M. Pereira, R. Henriques,
Fast live-cell conventional fluorophore nanoscopy with ImageJ through super-
resolution radial fluctuations. *Nat Commun* **7**, 12471 (2016).

18. E. Torres-García, R. Pinto-Cámara, A. Linares, D. Martínez, V. Abonza, E. Brito-
Alarcón, C. Calcines-Cruz, G. Valdés-Galindo, D. Torres, M. Jabloñski, H. H.
Torres-Martínez, J. L. Martínez, H. O. Hernández, J. P. Ocelotl-Oviedo, Y. Garcés,
M. Barchi, R. D'Antuono, A. Bošković, J. G. Dubrovsky, A. Darszon, M. G. Buffone,
R. R. Morales, J. M. Rendon-Mancha, C. D. Wood, A. Hernández-García, D. Krapf,

Á. H. Crevenna, A. Guerrero, Extending resolution within a single imaging frame.
Nature Communications 2022 13:1 **13**, 1–22 (2022).

19.   A. Romarowski, C. Sanchez-Cardenas, H. V. Ramirez-Gomez, L. D. C. Puga
Molina, C. L. Trevino, A. Hernandez Cruz, A. I. Darszon, M. G. Buffone, A.
Hernandez-Cruz, A. I. Darszon, M. G. Buffone, A. Hernandez Cruz, A. I. Darszon,
M. G. Buffone, A Specific Transitory Increase in Intracellular Calcium Induced by
Progesterone Promotes AE in Mouse Sperm. *Biol Reprod* **94**, 1–12 (2016).

20.   G. Lukinavičius, L. Reymond, E. D'Este, A. Masharina, F. Göttfert, H. Ta, A. Güther,
M. Fournier, S. Rizzo, H. Waldmann, C. Blaukopf, C. Sommer, D. W. Gerlich, H. D.
Arndt, S. W. Hell, K. Johnsson, Fluorogenic probes for live-cell imaging of the
cytoskeleton. *Nature Methods 2014 11:7* **11**, 731–733 (2014).

21.   V. Magliocca, S. Petrini, T. Franchin, R. Borghi, A. Niceforo, Z. Abbaszadeh, E.
Bertini, C. Compagnucci, Identifying the dynamics of actin and tubulin
polymerization in iPSCs and in iPSC-derived neurons. *Oncotarget* **8**, 111096–
111109 (2017).

22.   S. Yamazaki, M. Harata, T. Idehara, K. Konagaya, G. Yokoyama, H. Hoshina, Y.
Ogawa, Actin polymerization is activated by terahertz irradiation. *Sci Rep* **8**, 9990
(2018).

23.   K. W. Dunn, M. M. Kamocka, J. H. McDonald, A practical guide to evaluating
colocalization in biological microscopy. *Am J Physiol Cell Physiol* **300** (2011).

24.   M. A. Digman, R. Dalal, A. F. Horwitz, E. Gratton, Mapping the Number of
Molecules and Brightness in the Laser Scanning Microscope. *Biophys J* **94**, 2320
(2008).

25.   B. Mandracchia, X. Hua, C. Guo, J. Son, T. Urner, S. Jia, Fast and accurate
sCMOS noise correction for fluorescence microscopy. *Nat Commun* **11** (2020).

26.   S. Liu, M. J. Mlodzianoski, Z. Hu, Y. Ren, K. McElmurry, D. M. Suter, F. Huang,
sCMOS noise-correction algorithm for microscopy images. *Nature Methods 2017*
14:8 **14**, 760–761 (2017).

27.   M. Jin, E. Fujiwara, Y. Kakiuchi, M. Okabe, Y. Satouh, S. A. Baba, K. Chiba, N.
Hirohashi, Most fertilizing mouse spermatozoa begin their acrosome reaction before
contact with the zona pellucida during in vitro fertilization. *Proc Natl Acad Sci U S A*
108, 4892–4896 (2011).

28.   J. C. Conover, R. B. L. Gwatkin, Pre-loading of mouse oocytes with DNA-specific
fluorochrome (Hoechst 33342) permits rapid detection of sperm-oocyte fusion. *J*
Reprod Fertil **82**, 681–690 (1988).

29.   R. E. Hinkley, B. D. Wright, J. W. Lynn, Rapid visual detection of sperm-egg fusion
using the DNA-specific fluorochrome Hoechst 33342. *Dev Biol* **118**, 148–154
(1986).

30.   J. Stewart-Savage, B. D. Bavister, A cell surface block to polyspermy occurs in
golden hamster eggs. *Dev Biol* **128**, 150–157 (1988).

31.   K. T. Jones, C. Soeller, M. B. Cannell, The passage of Ca2+ and fluorescent
markers between the sperm and egg after fusion in the mouse. *Development* **125**,
4627–4635 (1998).

32.   R. Cohen, C. Mukai, J. L. Nelson, S. S. Zenilman, D. M. Sosnicki, A. J. Travis, A
genetically targeted sensor reveals spatial and temporal dynamics of acrosomal

calcium and sperm acrosome exocytosis. *Journal of Biological Chemistry* **298**, 101868–101869 (2022).

33. A. Darszon, A. Guerrero, B. E. Galindo, T. Nishigaki, C. D. Wood, Sperm-activating peptides in the regulation of ion fluxes, signal transduction and motility. *International Journal of Developmental Biology* **52**, 595–606 (2004).

34. A. Guerrero, T. Nishigaki, J. Carneiro, Yoshiro Tatsu, C. D. Wood, A. Darszon, Tuning sperm chemotaxis by calcium burst timing. *Dev Biol* **344**, 52–65 (2010).

35. A. R. Paredeza, Z. J. Assafa, D. Sept, L. Timofejeva, S. C. Dawsond, C. J. R. Wang, W. Z. Cande, An actin cytoskeleton with evolutionarily conserved functions in the absence of canonical actin-binding proteins. *Proc Natl Acad Sci U S A* **108**, 6151–6156 (2011).

36. D. V. Köster, S. Mayor, Cortical actin and the plasma membrane: inextricably intertwined. *Curr Opin Cell Biol* **38**, 81–89 (2016).

37. M. G. Buffone, T. W. Ijiri, W. Cao, T. Merdiushev, H. K. Aghajanian, G. L. Gerton, Heads or Tails? Structural events and molecular mechanisms that promote mammalian sperm AE and motility. *Molecular Reproduction* **79**, 4–18 (2013).

38. J. L. De La Vega-Beltran, C. Sánchez-Cárdenas, D. Krapf, E. O. Hernandez-González, E. Wertheimer, C. L. Treviño, P. E. Visconti, A. Darszon, Mouse sperm membrane potential hyperpolarization is necessary and sufficient to prepare sperm for the acrosome reaction. *Journal of Biological Chemistry* **287**, 44384–44393 (2012).

39. H. Hasuwa, Y. Muro, M. Ikawa, N. Kato, Y. Tsujimoto, M. Okabe, Transgenic Mouse Sperm that Have Green Acrosome and Red Mitochondria Allow Visualization of Sperm and Their Acrosome Reaction in Vivo. *Exp. Anim* **59**, 105–107 (2010).

40. G. Corkidi, P. Hernandez-Herrera, F. Montoya, H. Gadêlha, A. Darszon, Long-term segmentation-free assessment of head-flagellum movement and intracellular calcium in swimming human sperm. *J Cell Sci* **134** (2021).

41. S. Culley, K. L. Tosheva, P. Matos Pereira, R. Henriques, SRRF: Universal live-cell super-resolution microscopy. *Int J Biochem Cell Biol* **101**, 74–79 (2018).

42. G. Campagnola, K. Nepal, B. W. Schroder, O. B. Peersen, D. Krapf, Superdiffusive motion of membrane-targeting C2 domains. *Sci Rep* **5** (2015).

43. A. V. Weigel, B. Simon, M. M. Tamkun, D. Krapf, Ergodic and nonergodic processes coexist in the plasma membrane as observed by single-molecule tracking. *Proceedings of the National Academy of Sciences* **108**, 6438–6443 (2011).

44. I. Izeddin, M. El Beheiry, J. Andilla, D. Ciepielewski, X. Darzacq, M. Dahan, PSF shaping using adaptive optics for three-dimensional single-molecule super-resolution imaging and tracking. *Opt Express* **20**, 4957 (2012).

45. B. Huang, W. Wang, M. Bates, X. Zhuang, Three-dimensional super-resolution imaging by stochastic optical reconstruction microscopy. *Science* **319**, 810–3 (2008).

46. M. Ovesný, P. Křížek, J. Borkovec, Z. Švindrych, G. M. Hagen, ThunderSTORM: a comprehensive ImageJ plug-in for PALM and STORM data analysis and super-resolution imaging. *Bioinformatics* **30**, 2389–2390 (2014).

47. G. M. Luque, X. Xu, A. Romarowski, M. G. Gervasi, G. Orta, J. L. De la Vega-Beltrán, C. Stival, N. Gilio, T. Dalotto-Moreno, D. Krapf, P. E. Visconti, D. Krapf, A. Darszon, M. G. Buffone, Cdc42 localized in the CatSper signaling complex

regulates cAMP-dependent pathways in mouse sperm. *The FASEB Journal* **35**,
e21723 (2021).
48.   C. Stival, C. Ritagliati, X. Xu, M. G. Gervasi, G. M. Luque, C. B. Graf, J. L. De La
Vega-Beltrn, N. Torres, A. Darszon, D. Krapf, M. G. Buffone, P. E. Visconti, D.
Krapf, Disruption of protein kinase A localization induces AE in capacitated mouse
sperm. *Journal of Biological Chemistry* **293**, 9435–9447 (2018).
49.   G. L. Nicolson, R. Yanagimachi, H. Yanagimachi, Ultrastructural localization of
lectin-binding sites on the zonae pellucidae and plasma membranes of mammalian
eggs. *J Cell Biol* **66**, 263–274 (1975).

810
811
812

Figures legends

Figure 1. Sperm motility loss and FM4-64 fluorescence dynamics in acrosome-reacted

transgenic EGFP-DsRed2 sperm. A) Representative time series of transgenic EGFP-

DsRed2 sperm attached to concanavalin A-coated coverslips, with AE induced by 100 μM

progesterone. White squares indicate cells with spontaneous AE (prior to induction), while

pink squares highlight cells with progesterone-induced AE. A schematic representation of

AE in this transgenic model is shown on the right side of the panel. Scale bar = 20 μm. B)

Representative time series of transgenic EGFP-DsRed2 sperm that have already

experienced AE, attached to laminin-coated coverslips. The upper panel displays a cell with

motility after AE, and the lower panel shows an immotile cell. DsRed2 is presented in

orange, and EGFP in green. Scale bar = 10 μm. C) Quantification of motile and immotile

acrosome-reacted sperm (EGFP-). A total of 235 cells were counted across at least three

independent experiments. D) Representative time series of transgenic EGFP-DsRed2 sperm

stained with 10 μM FM4-64 and attached to concanavalin A-coated coverslips, with AE

induced by 100 μM progesterone. White squares indicate cells exhibiting patterns I, II, or

III after progesterone induction. Scale bar = 20 μm. A schematic representation of AE in

this transgenic model stained with FM4-64 is shown on the right side of the panel. E-G)

Representative images of capacitated transgenic EGFP-DsRed2 sperm stained with 10 μM

FM4-64. Panel E) displays an acrosome-intact, motile sperm (Pattern I), F) shows an

acrosome-reacted sperm with motility and low FM4-64 midpiece fluorescence (Pattern II),

and G) presents an acrosome-reacted sperm with no motility and high FM4-64 midpiece

fluorescence (Pattern III). In all three cases, cells were induced with 100 μM progesterone.

Scale bar = 10 μm. Enlarged images of each pattern are shown on the right panel. Scale bar = 2 μm. Representative images from at least five independent experiments are displayed.

Figure 2. Midpiece contraction coincides with the onset of AE. A) Left panel displays a wide-field fluorescence image of capacitated CD1 sperm membrane stained with 0.5 μM FM4-64, while the right panel shows its super-resolution SRRF reconstruction. Scale bar = 1 μm. B) Representative time series of sperm midpiece with no AE (no reac), spontaneous exocytosis (spont), progesterone (prog, 100 μM) and ionomycin-induced (iono, 10 μM) exocytosis, respectively. Following acquisition, images were analyzed using SRRF. Insets in the sperm head show wide-field images of AE. The midpiece diameter value is displayed in the bottom left corner for each time point. Scale bar = 1 μm. C) Quantification of midpiece diameter changes for each experimental group across time. Data are presented as a percentage of the initial diameter value before induction for each cell. D-E) Quantification of FM4-64 fluorescence in the sperm head and midpiece, respectively, for each experimental group across time. Data are presented as times of increases compared to initial fluorescence before AE induction. *p<0.05; #p<0.01; $p<0.001 and &p<0.0001 compared to the non-reacted group. A nonparametric Kruskal-Wallis test was performed in combination with Dunn's multiple comparisons test. Representative images of at least 5 independent experiments are shown, with 36 cells analyzed. F) Comparison of the midpiece architecture in acrosome-intact (AI, upper panel) and acrosome-reacted (AR, lower panel) sperm using scanning electron microscopy (SEM). Representatives images are shown, middle panels show quantification of these images whereas the left panel shows the quantification of all replicates. Data is presented as mean ± SEM, Kruskal-Wallis test was employed, p = 0.013, (AI n=85, AR n=72). Scale bar = 2 μm. G) Capacitated transgenic EGFP-DsRed2 sperm were induced by 100 µM progesterone. Cells were fixed and immunostained against □-GLUT3 in order to see the plasma membrane in the midpiece. Representative images of at least 2 independent experiments are shown. Left panel shows quantification of midpiece diameter in acrosome-intact and acrosome-reacted EGFP-DsRed2 sperm. Data is presented as mean ± sem. A nonparametric Mann Whitney test was performed, ****p<0.001 (AI n=84, AR n=47). Scale bar = 5 μm.

Figure 3. Contraction initiation preferentially occurs near the head-midpiece junction.
A) Schematic diagram illustrating the generation of super-resolution kymographs from SRRF-processed images. Crosslines are drawn every 2.5 μm through the sperm midpiece, and the Image-J Kymograph builder plug-in is used to create kymographs. The x-axis represents time, and the y-axis shows diameter changes. For wide-field images, a line along the midpiece is drawn to create fluorescence kymographs, with the y-axis representing midpiece length. Three sections of the midpiece are defined: proximal [0-7 μm], central [7-14 μm], and distal [14-21 μm]. B) Relative frequency graph displaying the distribution of the initiation sites for midpiece contractions in sperm with no AE (no reac), spontaneous exocytosis (spont), progesterone-induced (prog, 100 μM) exocytosis, and ionomycin-induced (iono, 10 μM) exocytosis, respectively. The x-axis indicates the midpiece section where the contraction begins: proximal (P), central (C), or distal (D). A chi^2 test was performed using the R language environment. C-D) Representative contraction kymographs and diameter measurements for progesterone-induced (100 μM) AE with one or two contraction initiation sites, respectively. In contraction kymographs, yellow lines demarcate midpiece sections, and colored spots indicate where super-resolution kymographs were created. Both kymograph and diameter measurement graphs display a dotted vertical line marking the induction point. For C, horizontal scale bar = 5 min and vertical scale bar = 1 μm and for D, horizontal scale bar = 3 min and vertical scale bar = 1 μm. Data from at least 5 independent experiments are shown, with 36 cells analyzed.

Figure 4. Midpiece contraction is driven by $[Ca^{2+}]_i$ changes. A) The representative time series demonstrates $[Ca^{2+}]_i$ and AE dynamics. Capacitated F1 sperm, loaded with Fluo4 AM, were immobilized on concanavalin A-coated coverslips and incubated in a recording medium (rec) containing 10 μM FM4-64. Rec was added as indicated by arrowheads. Scale bar = 10 μm. Beneath each frame in the Fluo4 (green) and FM4-64 (red) images, a color code displays the normalized intensity of the fluorescence signal (scale bar on the right of the panel). B) Kymograph-like analysis of the midpiece of 20 sperm following the addition of recording medium. Each row depicting the $[Ca^{2+}]_i$ (upper) and membrane (lower) dynamics of a single cell over time. A white dotted line indicates the moment of addition. The images presented are representative of at least five independent experiments. C) The representative time series demonstrates $[Ca^{2+}]_i$ and AE dynamics. Progesterone (prog, 100 μM) was added as indicated by arrowheads. Scale bar = 10 μm. Beneath each frame in the

Fluo4 (green) and FM4-64 (red) images, a color code displays the normalized intensity of

the fluorescence signal (scale bar on the right of the panel). D) Kymograph-like analysis of

the midpiece of 20 sperm following the addition of prog. Each row depicting the $[Ca^{2+}]_i$

(upper) and membrane (lower) dynamics of a single cell over time. A white dotted line

indicates the moment of addition. The images presented are representative of at least five

independent experiments. Consistently, a $[Ca^{2+}]_i$ transient precedes contraction, which is

proportional to the increase in FM4-64 fluorescence, as shown in Figure S2D. E)

Representative time series of $[Ca^{2+}]_i$ and midpiece contraction dynamics. Capacitated CD1

sperm were loaded with Fluo4 AM, immobilized on concanavalin A-coated coverslips, and

incubated in a recording medium containing 0.5 μM FM4-64. AE was induced with 100

μM progesterone (prog, arrowhead). Fluo4 images are widefield images, while FM4-64

images are SRRF-processed (super- resolution). Scale bar = 1 μm. F) 3D kymographs of

$[Ca^{2+}]_i$ (left) and contraction (right) dynamics. Data are normalized to the mean of the

frames before the induction of AE. Representative images from at least 5 independent

experiments are shown, with 36 cells analyzed.

Figure 5. The flagellar membrane approaches the actin cytoskeleton in the midpiece

of the sperm flagellum during midpiece contraction and AE. A-B) Representative time

series of plasma membrane and actin cytoskeleton colocalization in the midpiece in the

absence of AE (A) and during progesterone-induced AE (B, prog, 100 μM; FM4-64 shown

in red, SiR-actin shown in cyan). Capacitated CD1 sperm were loaded with 100 nM SiR-

actin, immobilized on concanavalin A-coated coverslips, and incubated in a recording

medium containing 0.5 μM FM4-64. Scale bar = 1 μm. Representative images from at least

5 independent experiments are shown, with 36 cells analyzed. C) Manders' colocalization

coefficients for acrosome-intact and acrosome-reacted cells in the midpiece. M1 was

assigned to FM4-64, and M2 to SiR-actin. Data are presented as mean ± SEM.

****$p<0.0001$ represents statistical significance. Paired t test was performed.   D-G)

Representative images of sperm midpiece stained with the acrosome marker PNA (left

panel, upper right insets, epifluorescence) and phalloidin (actin filaments, STORM) for

acrosome-intact (D-E) and acrosome-reacted (F-G) cells. The left panel displays a

longitudinal section of the midpiece (Scale bar = 10 μm), while the right panel illustrates

the radial distribution (Scale bar = 0.2 μm). E and G) Schematics of the analyzed actin

double helix parameters in the midpiece: helical pitch (*l*, distance between turns of the

helix, left panel), helical pitch frequency (f_0, number of turns the helix makes per 1 μm),
and radial distribution (R, radius of the double helix, right panel). Representative images
from at least 3 independent experiments are shown. Four acrosome-intact cells and seven
acrosome-reacted cells were analyzed.

Figure 6. Occurrence of midpiece contraction in sperm located within the perivitelline
space.

Representative images of IVF experiments using EGFP-DsRed2 sperm. Oocyte-sperm
complexes were stained with 10 μg/ml Hoechst and 10 μM FM4-64. A) Representative
time series of DIC images showing a sperm moving within the perivitelline space. Scale bar
in right panel = 20 μm, scale bar in left panel = 10 μm. B) DIC, Hoechst, EGFP, DsRed2*,
and FM4-64 images are shown for the case depicted in Figure 6A, note that, as the sperm is
moving, it is located in a different position in the perivitelline space. The area depicted in
the upper panel is shown in higher magnification in the lower panel. Scale bar in upper
panel = 20 μm, scale bar in lower panel = 10 μm. C-D) DIC, Hoechst, EGFP, DsRed2*,
and FM4-64 images are shown for a   (C) sperm that have passed through the ZP,
displaying AE with an initially non-contracted midpiece. After 20 minutes, as shown in
(D), the midpiece becomes contracted. The area depicted in the upper panel is shown in
higher magnification in the lower panel. Scale bar in upper panel = 20 μm, scale bar in
lower panel = 10 μm.   Representative images   from at least 6 independent experiments
are shown. A total of 23 oocytes and 69 sperm were analyzed.

Figure 7. Contraction of the midpiece occurs after sperm-egg fusion. A) Schematic
representation of the acquisition settings for the sperm-oocyte fusion assay. Images were
taken every 7 μm along the z-axis. B) DIC image of a sperm-oocyte complex, with the area
depicted in higher magnification in panel C. Scale bar = 20 μm. C) Representative time
series of sperm-oocyte fusion assay experiments using EGFP-DsRed2 sperm. Oocytes were
stained with 1 μg/ml Hoechst and 10 μM FM4-64. DIC, Hoechst, EGFP, DsRed2*, and
FM4-64 images are shown over time. Scale bar = 10 μm. Note that midpiece contraction
occurs after sperm-egg fusion and is proportional to the increase in FM4-64 fluorescence,
as shown in Figure S2D, highlighting its potential importance in the fertilization process.
Representative images from at least 4 independent experiments are displayed.

Figure 8. Midpiece contraction occurs in sperm-egg fusion after a decrease in $[Ca^{2+}]_i$.
A) Representative time series of sperm-oocyte fusion assay experiments using wild-type sperm loaded with 1 μM Fluo-4. Oocytes were stained with 1 μg/ml Hoechst and 10 μM FM4-64. DIC, Hoechst, Fluo-4, and FM4-64 images are shown over time. Scale bar = 10 μm. The color code below each frame in the Hoechst (shown in blue), Fluo-4 (shown in green), and FM4-64 (shown in red) images indicates the normalized intensity of the fluorescence signal (scale bar on the right of the panel). B-F) Quantification of sperm showing midpiece contraction (B, indicated by increased FM4-64 fluorescence), midpiece folding (C), midpiece unfolding (D), Fluo-4 fluorescence dynamics in the head (E), and different patterns in the midpiece (F) during fusion. Data are presented as the mean ± SEM of the percentage of sperm counted for each experiment. Representative images and data from at least 3 independent experiments are shown. A total of 74 oocytes and 136 sperm were analyzed. Note that midpiece contraction occurs in sperm-egg fusion after a decrease in Fluo-4 fluorescence.

Figure 9: Proposed model of the structural reorganization of the sperm actin cytoskeleton during key events of fertilization. The double helix actin network surrounding the mitochondrial sheath of the midpiece undergoes structural changes prior to the motility cessation. This structural modification is accompanied by a decrease in diameter of the midpiece and is driven by intracellular calcium changes that occur concomitant with a reorganization of the actin helicoidal cortex. Although midpiece contraction may occur in a subset of cells that undergo AE (A and B), the midpiece contraction occurs prior to motility cessation observed after sperm-egg fusion (C and D).

Figure 1

A prog

B DsRed2 EGFP

C

D prog — AE — Pattern I Pattern II Pattern III — FM4-64 EGFP

E Pattern I. Acrosome-intact and motile — FM4-64 EGFP

F Pattern II. Acrosome-reacted and motile — FM4-64 EGFP

G Pattern III. Acrosome-reacted and immotile — FM4-64 EGFP

Figure 2

A Wide-field | Super-resolution

B
prog

no reac
1′ 0.77 µm | 4′ 0.81 µm | 13′ 0.79 µm | 20′ 0.77 µm

spont
1′ 0.81 µm | 5′ 0.75 µm | 10′ 0.68 µm | 15′ 0.61 µm

prog
1′ 0.81 µm | 4′ 0.77 µm | 6′ 0.63 µm | 10′ 0.58 µm

iono
1′ 0.96 µm | 6′ 0.84 µm | 10′ 0.70 µm | 14′ 0.63 µm

C Normalized diameter (%) vs Time after induction (min)
no reac / spont / prog / iono

D Head — Normalized FM4-64 fluorescence vs Time after induction (min)

E Midpiece — Normalized FM4-64 fluorescence vs Time after induction (min)

F
Acrosome-intact
Acrosome-reacted
Diameter (µm): 1.05 / 0.85 / 0.65 / 0.45
0.731 ± 0.008 µm
0.694 ± 0.007 µm
Diameter (µm) — Acrosome-intact / Acrosome-reacted

G
Acrosome-intact — 0.84 µm
Acrosome-reacted — 0.73 µm
EGFP | DsRed2 | α-GLUT3 | Merge
0.856 ± 0.005 µm
0.726 ± 0.007 µm
****
Diameter (µm) — Acrosome-intact / Acrosome-reacted

Figure 3

A

Time → | Time ↗

Diameter

Proximal

Central

Distal

Time ↗ | Time →

Proximal [0 - 7) μm

Central [7 - 14) μm

Distal [14 - 21] μm

Length across midpiece

B

● Single focus ● Multiple foci

Relative frequency (%)

prog p=0.044

spont p=0.021

iono p=0.113

Section of midpiece

C

Length across midpiece (μm)

Time (min)

single focus

1.0 — 0.65
Normalized diameter

● 0 μm ● 2.5 μm ● 5 μm

Normalized diameter

Time (min)

D

Length across midpiece (μm)

Time (min)

two foci

1.0 — 0.5
Normalized diameter

● 0 μm ● 2.5 μm ● 5 μm
● 7.5 μm ● 10 μm ● 12 μm

Normalized diameter

Time (min)

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A

B Midpiece contraction

C Midpiece folding

D Midpiece unfolding

E Head [Ca^{2+}]$_i$

F Midpiece [Ca^{2+}]$_i$

Figure 9

Midpiece contraction

A Acrosome-intact   *Motile*

F-actin
Mitochondria
Plasma membrane

No
Motile

B Acrosome-reacted   *Motile*   *Immotile*

Yes
Immotile

Zona pellucida

C Binding to oocyte´s plasma membrane   (i)   *Motile*   No

Start of fusion   (ii)   *Motile*   No

[Ca²⁺]ᵢ decreases in the midpiece*   (iii)   *Motile*   -

D Midpiece contraction and folding   (iv)   *Immotile*   Yes

Midpiece unfolding   (v)   *Immotile*   Yes

▭ Oocyte´s plasma membrane   ⋈ F-actin   ▭ Sperm´s plasma membrane