Reorganization of the Flagellum Scaffolding Induces a Sperm Standstill Required for Fertilization

Martina Jabloñski[1]; Guillermina M. Luque[1]; Matías D. Gómez-Elías[1]; Claudia Sanchez-Cardenas[2]; Xinran Xu[3]; Jose Luis de la Vega-Beltran[2]; Gabriel Corkidi[4]; Alejandro Linares[2,6]; Victor X. Abonza Amaro[2,6]; Dario Krapf[5]; Diego Krapf[3]; Alberto Darszon[2]; Adan Guerrero[2,6*†,]; Mariano G. Buffone[1*†,].

[1]Instituto de Biología y Medicina Experimental, Consejo Nacional de Investigaciones Científicas y Técnicas (IBYME-CONICET), Buenos Aires, Argentina;
[2]Departamento de Genética del Desarrollo y Fisiología Molecular, Instituto de Biotecnología, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, Mexico.
[3]Department of Electrical and Computer Engineering and School of Biomedical Engineering, Colorado State University, Fort Collins, Colorado, USA;
[4]Laboratorio de Imágenes y Visión por Computadora, Instituto de Biotecnología, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, México.
[5]Instituto de Biología Molecular y Celular de Rosario (IBR), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Universidad Nacional de Rosario (UNR), Rosario, Argentina;
[6]Laboratorio Nacional de Microscopía Avanzada, Instituto de Biotecnología, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, Mexico.

†, These authors contributed equally to this work.

Mariano G. Buffone Ph.D.
Instituto de Biología y Medicina Experimental, CONICET. Vuelta de Obligado 2490 (1428), Buenos Aires, Argentina. Tel: (+5411)4783-2869.
E-mail: mgbuffone@ibyme.conicet.gov.ar.

Adan Guerrero Ph.D.
Laboratorio Nacional de Microscopía Avanzada, Instituto de Biotecnología, Universidad Nacional Autónoma de México, Cuernavaca, Morelos, México.
Email: adan.guerrero@ibt.unam.mx.

Abstract

Mammalian sperm delve into the female reproductive tract to fertilize the female gamete. Within the midpiece of the sperm flagellum lies a cortical network of actin that is arranged as a double helix sheltering the mitochondrial sheath. This work demonstrates that the actin network of the midpiece undergoes structural changes that result in motility cessation. This structural modification is accompanied by a decrease in diameter of the midpiece and is driven by intracellular calcium changes that occur concomitant with a reorganization of the actin helicoidal cortex. Although midpiece contraction may occur in a subset of cells that undergo acrosomal exocytosis, live-cell imaging during in vitro fertilization showed that the midpiece contraction is required for motility cessation during the fusion process. These findings provide the first evidence of the F-actin network's role in regulating sperm motility, adapting its function to meet specific cellular requirements during fertilization.

Introduction

Sperm motility is required for arrival at the site of fertilization and to penetrate the different layers surrounding the egg. The initial linear movement of ejaculated sperm is changed to a vigorous non-progressive motility called hyperactivation which is required for sperm for further migration towards the fertilization site (*3*). During that migration, mouse sperm undergo acrosomal exocytosis (AE) in the upper segments of the oviduct, before any interaction with the egg or its surrounding layers (*4–6*). This exocytic event is critical for sperm-egg fusion, because proteins involved in this process are rearranged in preparation for fusion (*7, 8*).

The sperm flagellum that is used for swimming within the female reproductive tract, is divided into three segments: the midpiece, the principal piece and the end piece. To perform this unique task, sperm also possess several characteristic peri-axonemal structures that provide specific functional features. The midpiece and the principal piece are separated by the annulus and, while the axoneme is wrapped by the outer dense fibers (ODFs) all along the flagellum, in the midpiece these structures are surrounded with mitochondria (*1*). The actin cytoskeleton is another major component of the sperm flagellum, whose polymerized, tridimensional organization in the midpiece has been shown to display a double helix arrangement that accompanies mitochondria (*2*). However, this spatial

distribution is lost in the principal piece, since actin is radially distributed between the
axoneme and the plasma membrane. The function of this novel specialized structure of
actin filaments is unknown.

The detailed aspects about the motility of sperm in their final journey to the
fertilization site and during the interaction with the female gamete are limited. For example,
in this final journey, most of sperm migrate to the ovulated eggs after acrosomal exocytosis,
and very little information is available one motility in those sperm. In addition, during
sperm-egg fusion, it is well documented that sperm stop moving (*9–11*) which is often
considered as indicative of an effective fusion between gametes. The cease in sperm
motility is necessary to complete the attachment and fusion between sperm and eggs, which
is a complex event mediated by several proteins identified using loss of function strategies
(*12*, *13*). However, the molecular mechanism behind the motility arrest in the male gamete
is not established.

In this work, we have investigated the structural changes in the actin cytoskeleton of
the sperm flagellum after acrosomal exocytosis and during the interaction with the eggs. By
using a combination of single-cell imaging and super-resolution microscopy methods, our
results demonstrate that the helical structure of polymerized actin undergoes a structural
change at the time of sperm-egg fusion. This process is driven by intracellular calcium
($[Ca^{2+}]_i$) and promotes a decrease in the sperm midpiece diameter as well as the arrest in
motility, which is required to complete the fusion process during fertilization.

Results

AE promotes a cease of motility in a subset of sperm cells.

After AE, sperm need to accomplish migration to the ampulla, locate and reach
unfertilized eggs, penetrate the cumulus matrix and the ZP, and finally, undergo fusion with
the oocyte (*14*). All these processes share the need to modulate sperm motility. Little is
known about the regulation of motility after the occurrence of AE or prior to the fusion
with the oocyte.

To monitor simultaneously the acrosomal status and sperm motility in live cells,
transgenic mice whose sperm express green fluorescent protein (EGFP) in their acrosome
and red fluorescent protein (DsRed2) in their mitochondria were used (Figure 1A,
Supplementary movie S1). Transgenic EGFP-DsRed2 sperm were immobilized through

their head on laminin-coated coverslips allowing flagellar movement (Figure 1B, Supplementary movie S2). While the presence of the acrosome is monitored using the EGFP fluorescence signal, motility was assessed through the movement of the midpiece (DsRed2 fluorescence), which was beating within and outside the imaging plane (Figure 1B).

The majority of acrosome-intact sperm were able to move. Interestingly, Figure 1B and 1C shows the coexistence of two populations of sperm that underwent AE induced upon addition of progesterone (cells lacking EGFP signal; progesterone is a physiological trigger of AE). Some of the AE sperms moved normally (34.1 ± 3.7 %, n = 80/235 sperm), whereas the majority of them remained immotile (65.9 ± 6.2 %, n = 155/2350 sperm) (Figure 1B and 1C, upper and lower panel, respectively, Supplementary movie S2).

The flagellar beat cycle encompasses a self-regulatory mechanism that receives feedback from molecular and mechanical signals (*15*). The complete abortion of flagellar beat cycle observed in Figure 1B and 1C might be an indicative of a stimulus provided by or occurring concomitant with the AE process. To comprehend the connection between AE and motility, we hypothesized the existence of a mechanical change in the flagella as a result of the AE. For this reason, AE was studied in the presence of FM4-64 fluorescent dye (*16*) to visualize structural changes at the plasma membrane, which can occur at macroscopic scales observed as an alteration in the shape of the flagellum, or at mesoscopic scales, monitored by local changes in FM4-64 fluorescence occurring at the vicinity or within the plasma membrane. EGFP-DsRed2 sperm were stimulated with progesterone and their 'motility' behavior was recorded for 5 min in the presence of FM4-64. The loss of EGFP fluorescence in the acrosome of transgenic mice correlated with a noticeable increase of FM4-64 fluorescence in the head, as shown in Figure 1D (see panels 0´ and 3´, Supplementary movie S3). In addition, we noticed that sperm that underwent AE, later showed an increase in the FM4-64 fluorescence intensity in the flagellum as well (see panel 14´of Figure 1D, Supplementary movie S3).

A more detailed analysis of individual cells demonstrated changes in FM4-64 fluorescence according to their motility (Supplementary movie S4). Figure 1E shows a motile acrosome-intact sperm, (a control case), where the fluorescence levels of EGFP and FM4-64 remained constant during the entire experiment (Pattern I). Interestingly, sperm that lost their acrosome (no EGFP and high FM4-64 fluorescence in the sperm head) and continued moving, presented low levels of FM4-64 fluorescence in the midpiece (Figure

1F, Pattern II). On the other hand, acrosome-reacted sperm that remained immotile showed a significant rise in the FM4-64 midpiece fluorescence (Figure 1G, Pattern III). Noteworthy, both subsets of cells (with and without motility) displayed the expected increase in FM4-64 fluorescence in the sperm head.

To deepen our understanding on how sperm lose motility during the AE, we designed a sperm tracking system that can also monitor changes in fluorescence in moving cells (Figure S1B and S1C). Three parameters were then assessed in real time, namely, the beat frequency of the flagellum (by tracking it on bright field images, or the DsRed2* channel, Figure S1A), the status of the acrosome (EGFP signal), and changes occurring at or within the membrane in the midpiece (FM4-64 signal). Figure S1B shows that sperm that did not undergo AE remained motile with a stable beat frequency and low FM4-64 fluorescence over the recording time. In stark contrast, after AE, cells gradually diminished the beating frequency and increased FM4-64 fluorescence in the midpiece (Figure S1C).

Altogether, these experiments demonstrate that AE promotes a cease of motility in a subset of sperm cells, which coincides with an increase in FM4-64 fluorescence in the midpiece.

AE is followed by a decrease in the midpiece diameter.

Live-cell super-resolution microscopy imaging was used to understand the structural changes occurring in the midpiece. Figure 2A shows a wide-field image of a sperm midpiece stained with FM4-64 (left) and analyzed using the Super-Resolution Radial Fluctuations (SRRF) approach (right) (*17*). AE dynamics was monitored by FM4-64 fluorescence in the head (Figure 2B, up right insets). A decrease in the midpiece diameter over time was observed following spontaneous AE, AE induced by a physiological agonist (progesterone), and induced by a non-physiological surge of Ca^{2+} (ionomycin) (Figure 2B, 2C and Supplementary movie S5). In the negative control, (no AE) the diameter remained unchanged. This phenomenon was also observed using the Means Shift Super-Resolution (MSSR) approach (*18*). To further support this observation, other plasma membrane probes of different molecular structure such as Memglow 700 (Figure S2A), Bodipy-GM1 (Figure S2B) and FM1-43 (Figure S2C) were used. In all three cases, a decrease in the midpiece diameter is observed after AE.

The FM4-64 fluorescence intensity in the head and in the midpiece (Figure 2D and 2E) were also assessed. As expected, the increase in FM4-64 fluorescence in the head

occurs after AE. Furthermore, in agreement to what was observed in Figure 1, fluorescence

intensity in the midpiece also increased over time in sperm that underwent AE (Figure 2D

and 2E).

The contraction of the midpiece initiates in the proximal part of the flagellum.

To investigate whether the contraction of the midpiece is triggered at a random

location or in a particular region of the flagellum, kymographs were used. A kymograph

allows visualization of dynamical aspects of a given phenomenon in a single image, where

the temporal dimension is expressed as an axis of the image, through a spatial-temporal

transformation of the dataset. Two types of kymographs were used (Figure 3A). First,

super-resolution microscopy kymographs were built to monitor dynamic changes in the

diameter of the flagellum along the midpiece. Second, to investigate the number of

contraction sites, fluorescence kymographs were made from diffraction limited images to

observe changes in FM4-64 fluorescence over time and across the entire midpiece.

Interestingly, Figure S2D shows a significant negative correlation between the midpiece

diameter and the FM4-64 fluorescence, unveiling a tool to obtain an approximate value of

the midpiece diameter without the need of super-resolution microscopy.

Three experimental subsets that underwent AE were assessed: spontaneous, induced

by progesterone, and induced by ionomycin. To scrutinize either randomness of focus-

driven contraction, the midpiece was segmented in three regions (Figure 3A): proximal

(near the neck), central, and distal (near the annulus). In these groups, it was then evaluated

whether cells presented one or more foci of contraction.

Figure 3B shows a cumulative study, which indicates that contraction is

preferentially initiated at the proximal part of the midpiece, regardless of being

progesterone-induced or spontaneous AE. When ionomycin was used as an agonist of AE,

the contraction began randomly in any segment of the midpiece. Figure 3C and 3D shows

two examples of flagellar dynamics for progesterone-induced sperm, which either

presented one or two foci of contraction. In both cases the diameter decreased all along the

midpiece, seen as a transition from dark to bright red colors in Figure 3C and 3D.

Noteworthy, focal points of contraction were consistently observed: i.e., a case with single

focus (Figure 3C), or a case with two foci of contraction (Figure 3D). Figure S2E and S2F

shows ionomycin-induced sperm that presents one or more foci. It can be observed in the

normalized diameter kymograph that it presents a single focus between the measurements

of 7.5 µm and 10 µm (Figure S2E, left panel). This is reflected in the right panel in the light

blue and orange dots, which are the actual diameter measurements that decreased first and

most (Figure S2E). A representative behavior within this experimental group is represented

in Figure S2F, where it is observed that the entire midpiece increased the contraction (all its

sections simultaneously). This is observed in the color scale of the normalized diameter

kymograph and coincides with the real measurements of the diameter of the right panel,

which, all together decreased their value (from 0 µm to 15 µm). Lastly, Figure S2G and

S2H shows examples of spontaneous AE with one or two foci of contraction, respectively.

In particular, in Figure S2H, it can be noted that there is a first focus of contraction from 0

µm to 5 µm approximately that coincided with the actual diameter measurements of the

right panel that decreased their value first and to a greater extent (black, pinks and green

dots). Although the entire midpiece contracted, a second focus of contraction appeared

around the measurements of 10 µm to 15 µm and coincidentally, these are the points that

decreased the diameter value the most (orange, purple and red points).

The reduction of the midpiece diameter occurs concomitantly with a $[Ca^{2+}]_i$ increase

in the flagellum.

It is widely accepted that an increase in $[Ca^{2+}]_i$ precedes AE (*16*, *19*). It has been

shown that AE initiation occurs after an increase in $[Ca^{2+}]_i$ in the sperm head that is later

propagated towards the midpiece (*19*). To comprehend the connection between AE and the

concomitant contraction of the midpiece, we hypothesized the existence of a signal, i.e.,

Ca^{2+}, which propagates from the head to the flagellum and has an undescribed potential to

modulate the architecture of the sperm flagellum.

To investigate if the increase in the concentration of $[Ca^{2+}]_i$ in the midpiece is

correlated with the midpiece contraction, sperm were incubated with FM4-64 and Fluo4, a

Ca^{2+} fluorescent sensor (Supplementary movie S6). Figure 4A shows a live-cell imaging

experiment of sperm cells that did not undergo AE (control case), nor a significant rise in

$[Ca^{2+}]_i$. Noteworthy, upon induction with progesterone (Figure 4C, Supplementary movie

S6), there was an increase in the midpiece $[Ca^{2+}]_i$ followed by an increase in FM4-64

fluorescence, which is indicative of a contraction of the flagellum (see Figure S2D).

These single cell experiments illustrate a reduction of the midpiece diameter which

occurs concomitantly with an increase in $[Ca^{2+}]_i$ within the flagellum. A large population of

sperm was then analyzed using low magnification (10X). Figure 4B and 4D shows a

collection of single cell kymographs, where each row represents the fluorescence dynamics of Fluo4 (green) and FM4-64 (red) of the midpiece over time. Unstimulated cells are displayed on Figure 4B, and sperm that were exposed to progesterone are shown in Figure 4D. Most of the cells display a transient $[Ca^{2+}]_i$ increase followed by an increase in FM4-64 fluorescence. This pattern was also observed in ionomycin-induced cells (Figure S3A, S3B and Supplementary movie S6) with the exception that, as expected, the rise in $[Ca^{2+}]_i$ was faster and sustained. These findings indicate that an $[Ca^{2+}]_i$ transient increase happening in the midpiece precedes a contraction of the flagellum.

The relationship between the observed $[Ca^{2+}]_i$ rise and the flagella contraction was then assessed using live-cell super-resolution imaging. Figure 4E shows that there is a $[Ca^{2+}]_i$ increase in the midpiece that coincides in space and time with a contraction of the midpiece. To provide insight in the spatio-temporal relationship of the $[Ca^{2+}]_i$ rise and the midpiece contraction, these results were visualized using a 3D kymograph encompassing the dynamics happening along the entire midpiece (Figure 4F). Overall, a rise in $[Ca^{2+}]_i$ occurred along the whole midpiece, which coincided with a generalized contraction of the midpiece (Figure 4F). Remarkably, a transient focal increase of $[Ca^{2+}]_i$ was observed to happen at the base of the flagellum, which correlates in space and time with a focal reduction of the diameter of the flagellum (Figure 4F). We then sought to identify a molecular/structural link between both processes.

The actin cytoskeleton approaches the plasma membrane during the contraction of the midpiece.

The midpiece is shaped by two major structural elements: the axoneme, which consists of tubulin and its accessory components, including the outer dense fibers (ODFs); and a recently described network of filamentous actin (F-actin) arranged in a helicoidal conformation along the midpiece (*2*). We investigated whether the actin network plays a role in regulating the midpiece contraction associated with AE. We considered three possible scenarios: (i) an unknown mechanical force, mediated by a rigid matrix such as F-actin, brings the plasma membrane closer to the center of the flagellum; (ii) a dynamic relationship exists between the plasma membrane and the actin cytoskeleton of the midpiece, facilitating contraction; (iii) midpiece contraction is driven by an uncharacterized signaling mechanism that acts independently of the F-actin cytoskeleton.

The role of the actin cytoskeleton in midpiece contraction was investigated through
live-cell super-resolution imaging using SiR-actin, a fluorescent probe that binds to F-actin
(*20–22*). We used FM4-64 to visualize the plasma membrane (Figure 5) and transgenic
sperm expressing DsRed2 to observe the mitochondrial network (Figure S4). Figure 5A-B
shows two stimulated sperm, one that experienced AE (Figure 4B) and another that did not
(Figure 4A). In both cases, a network of F-actin was observed beneath the plasma
membrane of the midpiece (cortical actin), enveloping the mitochondrial network (Figure
S4A). As expected, in the absence of AE, the midpiece diameter (observed through FM4-
64) remained unchanged (see also Figure 2, and Figure S4). Interestingly, in this case
(Figure 5A), both the diameter of the F-actin network and its proximity to either the plasma
membrane or the mitochondrial network remained static, suggesting a structural role for the
F-actin network within the midpiece (scenario - (i)), such as supporting the organization of
the mitochondrial network (Figure S4). Remarkably, acrosome-reacted sperm (Figure 5B)
experienced an abrupt structural reorganization of the midpiece, characterized by a decrease
in the distance between the plasma membrane and both the F-actin and mitochondrial
networks (Figures S4 and S5). This observation suggests that the remodeling of the
midpiece structure might be driven by a mechanical (or molecular) interaction occurring at
the boundaries of the plasma membrane (supporting scenario - (ii)).

To assess the potential interaction between the F-actin network and the plasma
membrane (or its associated components), we examined their colocalization through live-
cell super-resolution imaging. Figure 5C presents two Manders' colocalization coefficients
(*23*). M1 represents the proportion of F-actin (in pixels) that colocalizes with the plasma
membrane, while M2 measures the proportion of plasma membrane pixels colocalizing
with the F-actin network. A Manders´ value of 1 indicates full colocalization, while a value
of 0 implies no colocalization.

Figure 5C reveals that acrosome-intact sperm displayed low Manders' coefficients
(M1= 0.32 ± 0.03; M2= 0.28 ± 0.04). In contrast, sperm undergoing midpiece contraction
exhibited a significant increase in colocalization between the F-actin network and the
plasma membrane, as evidenced by the rise in M1 and M2 coefficients (M1= 0.65 ± 0.03;
M2 = 0.69 ± 0.03). This increased colocalization occurred simultaneously with midpiece
contraction.

In fluorescence microscopy, colocalization is indicative of mutual presence of two
populations of molecules within an observation volume shaped by the point spread

function, i.e. within a disc with a radius of 200-300 nm. Super-resolution microscopy
decreases the uncertainty for colocalization studies; however, the exact gain of resolution
varies according to the optical properties of the imaging instrument, sample optical
properties and preparation protocols, and the photophysical properties of the fluorescent
probes used.

Overall, these results confirm that (i) the contraction of the midpiece is linked to the
remodeling of the plasma membrane and potentially the F-actin cytoskeleton, and that (ii)
both structures interact with each other, directly or indirectly, at the nanoscales. We then
seek to understand whether the driving force for the contraction of the midpiece emanates
from the F-actin network. The following scenarios were envisaged: (a) the plasma
membrane moves towards the actin cytoskeleton, (b) the actin cytoskeleton moves towards
the plasma membrane, or (c) both structures come closer to the center of the flagellum. To
investigate the occurrence of any of these scenarios, the positions of fluorescence peaks of
FM4-64 and SiR-actin in super-resolution kymographs were tracked, as proxy of either
membrane or actin cytoskeleton localization, respectively (Figures S5A and S5B). Figure
S5C shows representative histograms of SiR-actin and FM4-64 over time. Both FM4-64
and SiR-actin fluorescence peaks came closer to each other. This effect can be also seen in
Figure S5F, where acrosome-intact cells display a distance of 0.180 ± 0.007 μm between
the plasma membrane and the actin cytoskeleton. In acrosome-reacted cells, the distance
was as small as 0.074 ± 0.006 μm after midpiece contraction.

In Figure S5D, a section of the midpiece of an ionomycin-induced AE sperm is
shown. In this case, the plasma membrane and the actin cytoskeleton approach each other
to the center of the cell (normalized to 0). The slope calculated from the linear adjustment
denotes the velocity of this change. In this example, the plasma membrane is contracting at
a rate of 14 μm/min, whereas the actin cytoskeleton is contracting at a rate of 3 μm/min
(Figure S5E). In both cases, the signal presents a negative slope in sperm that underwent
AE, which is consistent with a decrease in midpiece diameter. Since the results of the
analysis of SiR-actin slopes were not conclusive, we decided to study the actin cytoskeleton
structure in more detail.

The actin cytoskeleton in the midpiece is remodeled during the AE.

In the midpiece, polymerized actin forms a double helix that accompanies
mitochondria ((2). To investigate if the contraction of the midpiece is associated with

structural modifications of the actin double-helix, 3D-STORM was used. Cells were exposed to progesterone, fixed, and co-stained with PNA and phalloidin to visualize the acrosomal status and the actin structure, respectively.

Figure 5D and 5F show representative images of both dyes for acrosome-intact and acrosome-reacted sperm (left panels). Different parameters were calculated: 1) Helical pitch (l), which is the distance between turns of the helix; 2) Frequency ($f_0 = 1/l$), representing the number of turns that the helix makes in one micrometer; and 3) Radial distribution (R), which is the distance between the center of the midpiece and the maximum of fluorescence (*2*). The frequency, obtained using a Fourier transform of the image, increases substantially in the acrosome-reacted sperm compared to intact sperm (6.29 vs 4.04 μm^{-1}, Figure 5E and 5G, red arrows in left panels). In addition, the helical pitch diminished its magnitude in acrosome-reacted sperm (159 nm vs 248 nm from the center of the flagellum) (Figure 5E and 5G, left panels). The radial distribution of F-actin in cells that underwent AE was smaller compared to acrosome-intact sperm (252 vs 300 nm, Figure 5E and 5G, right panel). Altogether, these results confirm that the actin double-helix undergoes structural changes following the contraction of the midpiece

To further investigate the structural rearrangements of the actin cytoskeleton during AE, a number and brightness analysis was performed (*24–26*). The number measure indicates the abundance or concentration of SiR-Actin molecules bound to F-actin fibers, while the brightness analysis helps reveal dynamic changes in molecular aggregation processes. This is based on the idea that higher-order fluorescent complexes, which are mobile within the sample, will cause an increase in signal variability over time. An increase in brightness suggests the formation or movement of supramolecular structures, such as bundles of actin fibers bearing SiR-Actin.

AE induction with progesterone and ionomycin resulted in a significant increase in actin number and brightness compared to the control (non-reacted) group (Figure S6A). Spontaneously reacted sperm also exhibited this behavior moderately (Figure S6A).

The observed increase in SiR-Actin number after AE induction suggests major actin monomer recruitment to the polymerizing fibers (Figure S6A and S6B), which produces an apparent increase in local fluorophore concentration (Figure S6B). This effect positively correlated with a similar increase in SiR-actin fluorescence (Figure S6A), which is known to indicate actin polymerization due to the probe's high affinity for F-actin.

Actin cytoskeleton polymerization in the midpiece results in a signal variation of
increased magnitude (an increase in SiR-Actin brightness) as it undergoes structural
rearrangements (Figure S6A and S6B), which are compatible with other observations
reported in this work. Both assembly and transport of actin complexes into higher
oligomeric states (F-actin) across the cellular milieu led to an apparent increase in the
registered SiR-Actin brightness (Figure S6B). Additionally, actin filament displacement
along the mitochondrial sheath during AE provides a SiR-Actin brightness measure with
high dispersion, indicating a redistribution of this cellular structure. These results suggest
that actin filaments are locally redistributed and remodeled during AE (Figure S6B).

Most sperm in the perivitelline space exhibit midpiece contraction

In mice, only sperm undergoing AE prior to binding to the ZP can penetrate and
fertilize. We hypothesized that midpiece changes and motility cessation occur only after
acrosome-reacted sperm penetrate the ZP. Live imaging was performed after *in vitro*
fertilization (IVF) using transgenic EGFP-DsRed2 sperm loaded with FM4-64. Eggs were
inseminated with a high sperm count (200,000 cells) to increase the number of cells
observed. Figure 6A shows an intact acrosome sperm with low FM4-64 fluorescence in the
midpiece. Most sperm that passed through the ZP were acrosome-reacted and displayed
high FM4-64 fluorescence (Figure 6B). Interestingly, some acrosome-reacted moving
sperm within the perivitelline space had low FM4-64 fluorescence in the midpiece (Figure
6C). After 20 minutes, these sperm stopped moving and exhibited increased FM4-64
fluorescence, indicating midpiece contraction (Figure 6D). Quantification revealed that
82.6% (57/69 sperm) of sperm within the perivitelline space displayed high FM4-64
fluorescence in the midpiece (Figure 6E). These results suggest that midpiece contraction
and motility cessation occur only after acrosome-reacted sperm penetrate the ZP.

Midpiece contraction takes place following sperm-egg fusion.

Motile acrosome-reacted sperm, initially lacking midpiece contraction, can
penetrate the ZP. However, they eventually stop moving and exhibit an increase in FM4-64
fluorescence. This suggests that the cessation of motility plays a crucial role in fertilization
events occurring after ZP binding and penetration, and that a similar change in midpiece
architecture is necessary for successful sperm-egg fusion.

To explore this hypothesis, we designed a live imaging experiment using zona-free eggs. Denuded oocytes were loaded with Hoechst 33342, a nuclear dye, to visualize the exact moment of sperm-egg fusion (27–29). The experiment involved FM4-64, transgenic EGFP-DsRed2 sperm, and simultaneous signal collection from five separate channels (DIC, Hoechst, EGFP, DsRed2*, and FM4-64). To enhance observation likelihood, volumetric data was collected by imaging at different z-planes (Figure 7A).

Figure 7C presents a representative volumetric time series of sperm-egg fusion, with only the optimal focal plane shown. The acrosome-reacted sperm (EGFP negative) initially did not display (Supplementary movie S7) Hoechst fluorescence, indicating a lack of fusion. However, 14 minutes later, the dye entered the sperm and stained the nucleus, indicating fusion initiation. At this stage, the sperm remains motile while bound to the egg plasma membrane. The midpiece diameter remained unchanged during initial sperm-egg fusion due to low FM4-64 fluorescence. Following gamete fusion initiation, the midpiece folded, and FM4-64 fluorescence increased (frames 38 to 44 min), causing sperm motility to cease. In some instances, the midpiece extended again after folding (Figure S7B). These findings demonstrate that midpiece contraction occurs subsequent to sperm-egg fusion.

A decrease in $[Ca^{2+}]_i$ in the midpiece following fusion precedes the midpiece contraction.

In previous experiments, an increase in $[Ca^{2+}]_i$ was observed throughout the midpiece, coinciding with midpiece contraction (Figure 4 and Figure S3). To further examine the mechanism behind midpiece contraction after fusion, wild-type sperm loaded with Fluo4 were exposed to denuded oocytes loaded with Hoechst. We hypothesized that dynamic changes in $[Ca^{2+}]_i$ during gamete fusion drive the midpiece contraction necessary for sperm immobilization.

Figure 8A presents representative images from a time course live imaging experiment designed to observe sperm-egg fusion. The experiment involved simultaneous signal collection from four separate channels over time (DIC, Hoechst, Fluo4, and FM4-64). To enhance observation likelihood, images were taken at different z-planes, though only the optimal focal plane is shown.

Unexpectedly, sperm binding to the egg plasma membrane displayed high levels of $[Ca^{2+}]_i$ in the head and midpiece (Figure 8A, time 1 min) (Supplementary movie S8). A few minutes after fusion began (indicated by an increase in Hoechst signal in the sperm), a

decrease in $[Ca^{2+}]_i$ in the head and midpiece was observed (21 min to 51 min). This

decrease in $[Ca^{2+}]_i$ was followed by midpiece contraction, as evidenced by the increase in

FM4-64 fluorescence (time 56 min). In several cells, the midpiece folded back before

contracting. The gametes that underwent this change in the flagellum typically remained

folded, but occasionally the tail unfolded again. Figures 8B-F quantify the key events

described above during sperm-egg fusion.

In summary, $75.8 \pm 12.2\%$ of the sperm exhibited midpiece contraction upon fusion.

In $61.8 \pm 12.0\%$ of cases, the midpiece folded on itself, and afterward, only $31.6 \pm 7.5\%$

unfolded and stretched out again. All sperm bound to the plasma membrane before fusion

presented high $[Ca^{2+}]_i$ in the head, and the majority displayed a decrease in head $[Ca^{2+}]_i$

before midpiece contraction ($75.1 \pm 3.1\%$). Concerning the midpiece, three patterns were

observed during fusion: in the majority of cases ($69.7 \pm 11.5\%$), the midpiece experienced a

decrease in $[Ca^{2+}]_i$. Additionally, $10.3 \pm 10.3\%$ of sperm showed a transient increase in

$[Ca^{2+}]_i$, while the remaining $20.0 \pm 9.4\%$ displayed no changes.

Collectively, these results indicate that a decrease in $[Ca^{2+}]_i$ in the midpiece after

fusion precedes midpiece contraction and the cessation of sperm motility necessary for

successful sperm-egg fusion.

Discussion

In this paper, we demonstrate the existence of a fundamental structural change that

occurs in the sperm flagellum at the time of fusion with the egg. Using a plethora of

advanced microscopy methods and single cell imaging, we provide insight about cellular

and molecular events that occur in acrosome-reacted sperm, which, undoubtedly, are the

ones capable of fertilizing an oocyte (*4, 5*). At this precise moment of the fertilization

process, sperm need to stop moving to complete the fusion between the two gametes. The

cease of movement is caused by two concomitant processes that take place in the flagellar

midpiece region: a change in the F-actin helical structure and a decrease in the midpiece

diameter. To the best of our knowledge, this is the first time that a structural modification

of the sperm flagellum related to a specific cellular necessity, i.e., gamete fusion, is

described.

To arrive at the site of fertilization within the female reproductive tract, sperm

motility is required. It is also fundamental to penetrate the different layers surrounding the

egg. In this journey, sperm sense the environment and adapt their movement to the different

physiological scenarios. In addition, recent evidence also demonstrated that after being

released from their site of storage, the reservoir, mouse sperm undergo AE (*4, 5*). This

process occurs in the upper segments of the oviduct before any interaction with the eggs or

their surrounding layers. This observation opened a new scenario about the motility of

sperm in their last transit to the site of fertilization. Little is known about sperm in that

region and how sperm move after AE. In this regard, most of the information about

mammalian sperm comes from studies conducted *in vitro*, using a mixture of acrosome-

intact and acrosome-reacted sperm. Even if their motility is analyzed in a subjective manner

or using cell tracking systems, virtually all the experiments are conducted without

discerning the acrosomal status of the cells. Another possible source of artifacts in this

analysis is related with the fact that most of the experiments are also performed using

epididymal sperm (not ejaculated) in aqueous solutions that support sperm capacitation but

do not represent the natural viscous environment present in the female tract.

By performing in vitro experiments, we detected a strong association between the

decrease in the midpiece diameter and the cease of sperm motility. This is observed in a

subset of sperm after the occurrence of AE. This may suggest that only sperm that undergo

AE in the oviduct and do not experience this midpiece contraction are capable of migrating

to the ampulla and penetrate the cumulus and the ZP. Those sperm that undergo this

phenomenon earlier in the tract may not be suitable to continue their journey suggesting

that this also may select the gametes during their transit to the ampulla. Previous

observations tracking sperm within the female tract have shown the existence of acrosome-

reacted sperm within the tract that remain non-motile (*4–6*).

Acrosome-reacted sperm that bind and penetrate the ZP are ready to fuse with the

egg. During sperm-egg fusion, several authors have reported in different species that sperm

stop moving (*9–11*). However, the mechanism behind this behavior is not established. One

possible explanation is that fusion promotes ion transport changes in sperm, which

significantly alter flagellar movement. In this sense, it was previously demonstrated that

certain ions such as Ca^{2+} may diffuse from the oocyte to the sperm (*30*). However, a clear

technical limitation in our experimental approach is that the probes that monitor ion

dynamics may be exchanged between both gametes. If the concentration of a given probe

does not remain stable, it is impossible to determine the accurate change that occurs during

fusion. Future experiments may take advantage of transgenic models that incorporate a

particular sensor to study this process in vivo, such as the one used by Cohen and
collaborators (*31*).

Regardless of the precise nature of ion exchange between sperm and eggs, the
diameter of the sperm flagellum in the midpiece is reduced. This midpiece diameter
reduction that completely stops sperm motility is apparently needed for sperm to complete
the fusion process. Our observation clearly supports this notion. Importantly, the sperm
flagellum folds back during fusion coincident with the decrease in the midpiece diameter.
Interestingly, this was previously observed in mammals as well as in sea urchin sperm (*9,*
32, 33).

Like other cylindrical biological structures, the sperm flagellum relies on the
cytoskeleton for its structural organization and specialized mechanical properties. In
addition to the change in midpiece diameter, a significant rearrangement of the F-actin
cytoskeleton also occurs. In the midpiece, the polymerized actin is organized in a double
helix accompanying the mitochondria (*2*). As in many other organisms, the actin
cytoskeleton possesses important structural functions, and dynamic changes of F-actin
allow the cells to conduct important cellular tasks such as exocytosis. However, less is
known about the structural changes that may undergo the actin cytoskeleton within cilia
and flagella. Remarkably, it has been reported that actin forms helix-like structures in the
flagellum of the parasite *Giardia intestinalis* (*34*). These findings open the question of
whether, to some extent, flagellar helical structures are conserved among diverse species.
Our observations demonstrate that the F-actin double helix undergoes a clear change in the
helical pitch as well as in the radial distance related to the axoneme. This is concomitant
with the decrease in midpiece diameter. Our single-cell experiments using super-resolution
microscopy also revealed that the plasma membrane approached the F-actin network during
this change. It is well established that various proteins can function as linkers between the
plasma membrane and the actin cytoskeleton (*35*), but their roles in this specific process
remain to be studied. Regardless of how these structures are connected, it is evident that
both occur in association. However, our experimental data cannot determine whether the
plasma membrane is causing this effect on the actin network or if the actin network
influences the plasma membrane instead.

Another observation emerging from our study is that a change in $[Ca^{2+}]_i$ occurs prior
to the midpiece contraction. It is well known that Ca^{2+} is important for AE, and a specific
transient rise in $[Ca^{2+}]_i$ originating in the head can trigger exocytosis (*19*). Previous

observations have demonstrated that the sperm head and tail are not separate compartments, and that ions and other molecules can move between them (*36, 37*). In our single-cell experiments, we observed that the rise in FM4-64 fluorescence in the midpiece occurs after the increase in $[Ca^{2+}]_i$ in that region, suggesting a potential association between these events. This could involve diffusion or active transport processes; further investigation is required to determine the precise mechanism. The same phenomenon was studied in sperm bound to the egg plasma membrane to evaluate if the rise in $[Ca^{2+}]_i$ also occurs at the time of fusion. Remarkably, we noticed that most of the bound sperm that ended up fusing with the eggs displayed high levels of $[Ca^{2+}]_i$ in both the head and the flagellum. The midpiece contraction and the immobilization occurred when the levels of $[Ca^{2+}]_i$ went down in the midpiece suggesting a possible connection between both events. However, our experimental approach limits the interpretation of this result. One possible explanation is that as soon as sperm binds to the plasma membrane of the oocyte, there is a rapid increase in $[Ca^{2+}]_i$ in the sperm. On the other hand, a massive transport of Ca^{2+} from the egg to the sperm could also occur. These hypotheses, however, are hindered by the technical limitations mentioned above. As the oocyte is not stained with Fluo4, we cannot rule out that the apparent $[Ca^{2+}]_i$ decrease seen in these experiments is due to dye diffusion into the oocyte. Either way, it is clear that the transport of this ion into or out of the sperm is key to cease motility at this fundamental step of fertilization. Additionally, these results may also demonstrate that only sperm with elevated $[Ca^{2+}]_i$ are capable of binding to the eggs. All these possible scenarios need to be determined in future experiments. Thus, regardless of the mechanism, a clear change in $[Ca^{2+}]_i$ dynamics is observed at the time of midpiece contraction. Future studies will aid to indicate if certain Ca^{2+}-dependent proteins that can modify the actin cytoskeleton are responsible for this change.

In conclusion, we demonstrate that sperm undergo a structural reorganization of the actin cytoskeleton during key events of fertilization, as summarized in the working model shown in Figure 9. Our findings introduce a new aspect of study in reproductive biology.

In cilia, apart from the findings presented in this paper, nothing is known about the regulation of motility by actin. Actin has been found to participate in ciliogenesis, but its involvement in active motility regulation has not been reported. This highlights a potentially unique role for the actin cytoskeleton in regulating sperm function during

fertilization. Our discovery of actin's dynamic reorganization in sperm suggests it could
have a more active role in regulating motility and other functions.

Previous research has mainly focused on identifying proteins essential for sperm-
egg fusion. Our results reveal a previously unexplored biological mechanism in mammalian
fertilization, opening new avenues for contraceptive method development

Given the conservation of cilia and flagella across various organisms, our discovery
could prompt further research on the role of the actin network in these structures.
Investigating actin's role in motility regulation and its adaptation to specific cellular
conditions has broad implications for our understanding of cellular processes and potential
therapeutic targets.
578

579   **Materials and methods**
580

581   Detailed methods and protocols are provided in the SI appendix. For sample details,
582   optical equipment, imaging conditions and probes used, see Supplementary Table S1.
583

584   **Acknowledgments**
585   We gratefully acknowledge the financial support provided by the Williams and Rene Baron
586   Foundations, the Male Contraceptive Initiative (MCI), and the Chan-Zuckerberg Initiative
587   (CZI) for this work. Our sincere thanks go to Dr. Pablo Visconti for his valuable insights
588   throughout the course of this project.
589

Funding: This work was supported by: Chan Zuckerberg Initiative (2021-240504 to MB
and AG, GBI-0000000093 to AG, 2022-252509 to AG); Agencia Nacional de Promoción
Científica y Tecnológica (PICT, 2017-3047, 2018-1988 and 2020-00988); Dirección
General de Asuntos del Personal Académico/Universidad Nacional Autónoma de México
(DGAPA/UNAM grant: IN105222 to GK, IN211821 to AG and IN200919 to AD);
National Institute of Health (R01HD380882 to AD, R01HD106968 to DK and MGB).
596

Author contributions
Investigation: MJ, GML, MDGE, CSC, JLDLVB, AL. Data curation: MJ and VXAA.
Writing: MJ, MGB and AG. Methodology: XX, Diego K and GK. Conceptualization:
Dario K, Diego K, AD, MGB and AG. Supervision: MGB and AG.

Competing interests: The authors declare no competing interests.

Data, Materials, and Software Availability: Data acquired and analyzed during this study

are included in this manuscript or available at Zenodo. Github was used for depositing the

code use in analysis.

References

1.   H. Otani, O. Tanaka, K. Kasai, T. Yoshioka, Development of mitochondrial helical
sheath in the middle piece of the mouse spermatid tail: regular dispositions and
synchronized changes. *Anat Rec.* **222**, 26–33 (1988).
2.   M. G. Gervasi, X. Xu, B. Carbajal-Gonzalez, M. G. Buffone, P. E. Visconti, D. Krapf,
J Cell Sci, in press, doi:10.1242/jcs.215897.
3.   R. P. Demott, S. S. Suarez, Hyperactivated sperm progress in the mouse oviduct.
Biol Reprod. **46**, 779–785 (1992).
4.   T. Hino, Y. Muro, M. Tamura-Nakano, M. Okabe, H. Tateno, R. Yanagimachi, The
Behavior and Acrosomal Status of Mouse Spermatozoa In Vitro, and Within the
Oviduct During Fertilization after Natural Mating. *Biol Reprod.* **95**, 50–50 (2016).
5.   F. A. La Spina, L. C. Puga, A. Romarowski, A. M. Vitale, T. L. Falzone, D. Krapf, N.
Hirohashi, M. G. Buffone, Mouse sperm begin to undergo acrosomal exocytosis in
the upper isthmus of the oviduct. *Dev Biol.* **411**, 172–182 (2016).
6.   Y. Muro, H. Hasuwa, A. Isotani, H. Miyata, K. Yamagata, M. Ikawa, R. Yanagimachi,
M. Okabe, Behavior of Mouse Spermatozoa in the Female Reproductive Tract from
Soon after Mating to the Beginning of Fertilization. *Biol Reprod.* **94**, 80–80 (2016).
7.   N. Inoue, M. Ikawa, A. Isotani, M. Okabe, The immunoglobulin superfamily protein
Izumo is required for sperm to fuse with eggs. *Nature.* **434**, 234–238 (2005).
8.   T. Noda, Y. Lu, Y. Fujihara, S. Oura, T. Koyano, S. Kobayashi, M. M. Matzuk, M.
Ikawa, Sperm proteins SOF1, TMEM95, and SPACA6 are required for sperm-
oocyte fusion in mice. *Proc Natl Acad Sci U S A.* **117** (2020),
doi:10.1073/PNAS.1922650117/-/DCSUPPLEMENTAL.
9.   P. Gaddum-Rosse, R. J. Blandau, L. B. Langley, D. E. Battaglia, In vitro fertilization
in the rat: observations on living eggs. *Fertil Steril.* **42**, 285–292 (1984).
10.   P. Gaddum-Rosse, R. J. Blandau, L. B. Langley, K. Sato, Sperm tail entry into the
mouse egg in vitro. *Gamete Res.* **6**, 215–223 (1982).
11.   B. Ravaux, N. Garroum, E. Perez, H. Willaime, C. Gourier, A specific flagellum
beating mode for inducing fusion in mammalian fertilization and kinetics of sperm
internalization. *Sci Rep.* **6** (2016), doi:10.1038/SREP31886.
12.   V. E. Deneke, A. Pauli, The Fertilization Enigma: How Sperm and Egg Fuse. *Annu.
Rev. Cell Dev. Biol. 2021.* **37**, 391–414 (2021).
13.   K. K. Siu, V. H. B. Serrão, A. Ziyyat, J. E. Lee, The cell biology of fertilization:
Gamete   attachment   and   fusion.   *J   Cell   Biol.*   **220**   (2021),
doi:10.1083/JCB.202102146.
14.   R. Yanagimachi, *Mammalian fertilization* (Raven Press, New York, 1994).
15.   K. Inaba, K. Shiba, Microscopic analysis of sperm movement: links to mechanisms
and protein components. *Microscopy.* **67**, 144–155 (2018).

16. C. Sánchez-Cárdenas, M. R. Servín-Vences, O. José, C. L. Treviño, A. Hernández-
Cruz, A. Darszon, Acrosome reaction and Ca2+ imaging in single human
spermatozoa: New regulatory roles of [Ca2+]i. *Biol Reprod.* **91** (2014),
doi:10.1095/biolreprod.114.119768.

17. N. Gustafsson, S. Culley, G. Ashdown, D. M. Owen, P. M. Pereira, R. Henriques,
Fast live-cell conventional fluorophore nanoscopy with ImageJ through super-
resolution radial fluctuations. *Nat Commun.* **7**, 12471 (2016).

18. E. Torres-García, R. Pinto-Cámara, A. Linares, D. Martínez, V. Abonza, E. Brito-
Alarcón, C. Calcines-Cruz, G. Valdés-Galindo, D. Torres, M. Jabloñski, H. H.
Torres-Martínez, J. L. Martínez, H. O. Hernández, J. P. Ocelotl-Oviedo, Y. Garcés,
M. Barchi, R. D'Antuono, A. Bošković, J. G. Dubrovsky, A. Darszon, M. G. Buffone,
R. R. Morales, J. M. Rendon-Mancha, C. D. Wood, A. Hernández-García, D. Krapf,
Á. H. Crevenna, A. Guerrero, Extending resolution within a single imaging frame.
Nature Communications 2022 13:1. **13**, 1–22 (2022).

19. A. Romarowski, C. Sanchez-Cardenas, H. V. Ramirez-Gomez, L. D. C. Puga
Molina, C. L. Trevino, A. Hernandez Cruz, A. I. Darszon, M. G. Buffone, A.
Hernandez-Cruz, A. I. Darszon, M. G. Buffone, A. Hernandez Cruz, A. I. Darszon,
M. G. Buffone, A Specific Transitory Increase in Intracellular Calcium Induced by
Progesterone Promotes Acrosomal Exocytosis in Mouse Sperm. *Biol Reprod.* **94**,
1–12 (2016).

20. G. Lukinavičius, L. Reymond, E. D'Este, A. Masharina, F. Göttfert, H. Ta, A. Güther,
M. Fournier, S. Rizzo, H. Waldmann, C. Blaukopf, C. Sommer, D. W. Gerlich, H. D.
Arndt, S. W. Hell, K. Johnsson, Fluorogenic probes for live-cell imaging of the
cytoskeleton. *Nature Methods 2014 11:7.* **11**, 731–733 (2014).

21. V. Magliocca, S. Petrini, T. Franchin, R. Borghi, A. Niceforo, Z. Abbaszadeh, E.
Bertini, C. Compagnucci, Identifying the dynamics of actin and tubulin
polymerization in iPSCs and in iPSC-derived neurons. *Oncotarget.* **8**, 111096–
111109 (2017).

22. S. Yamazaki, M. Harata, T. Idehara, K. Konagaya, G. Yokoyama, H. Hoshina, Y.
Ogawa, Actin polymerization is activated by terahertz irradiation. *Sci Rep.* **8**, 9990
(2018).

23. K. W. Dunn, M. M. Kamocka, J. H. McDonald, A practical guide to evaluating
colocalization in biological microscopy. *Am J Physiol Cell Physiol.* **300** (2011),
doi:10.1152/AJPCELL.00462.2010.

24. M. A. Digman, R. Dalal, A. F. Horwitz, E. Gratton, Mapping the Number of
Molecules and Brightness in the Laser Scanning Microscope. *Biophys J.* **94**, 2320
(2008).

25. B. Mandracchia, X. Hua, C. Guo, J. Son, T. Urner, S. Jia, Fast and accurate
sCMOS noise correction for fluorescence microscopy. *Nat Commun.* **11** (2020),
doi:10.1038/S41467-019-13841-8.

26. S. Liu, M. J. Mlodzianoski, Z. Hu, Y. Ren, K. McElmurry, D. M. Suter, F. Huang,
sCMOS noise-correction algorithm for microscopy images (2017),
doi:10.1038/nmeth.4379.

27. J. C. Conover, R. B. L. Gwatkin, Pre-loading of mouse oocytes with DNA-specific
fluorochrome (Hoechst 33342) permits rapid detection of sperm-oocyte fusion. *J
Reprod Fertil.* **82**, 681–690 (1988).

28.   R. E. Hinkley, B. D. Wright, J. W. Lynn, Rapid visual detection of sperm-egg fusion
using the DNA-specific fluorochrome Hoechst 33342. *Dev Biol*. **118**, 148–154
(1986).

29.   J. Stewart-Savage, B. D. Bavister, A cell surface block to polyspermy occurs in
golden hamster eggs. *Dev Biol*. **128**, 150–157 (1988).

30.   K. T. Jones, C. Soeller, M. B. Cannell, The passage of Ca2+ and fluorescent
markers between the sperm and egg after fusion in the mouse. *Development*. **125**,
4627–4635 (1998).

31.   R. Cohen, C. Mukai, J. L. Nelson, S. S. Zenilman, D. M. Sosnicki, A. J. Travis, A
genetically targeted sensor reveals spatial and temporal dynamics of acrosomal
calcium and sperm acrosome exocytosis. *Journal of Biological Chemistry*. **298**,
101868–101869 (2022).

32.   A. Darszon, A. Guerrero, B. E. Galindo, T. Nishigaki, C. D. Wood, Sperm-activating
peptides in the regulation of ion fluxes, signal transduction and motility. *International*
Journal of Developmental Biology. **52**, 595–606 (2004).

33.   A. Guerrero, T. Nishigaki, J. Carneiro, Yoshiro Tatsu, C. D. Wood, A. Darszon,
Tuning sperm chemotaxis by calcium burst timing. *Dev Biol*. **344**, 52–65 (2010).

34.   A. R. Paredeza, Z. J. Assafa, D. Sept, L. Timofejeva, S. C. Dawsond, C. J. R.
Wang, W. Z. Cande, An actin cytoskeleton with evolutionarily conserved functions in
the absence of canonical actin-binding proteins. *Proc Natl Acad Sci U S A*. **108**,
6151–6156 (2011).

35.   D. V. Köster, S. Mayor, Cortical actin and the plasma membrane: inextricably
intertwined. *Curr Opin Cell Biol*. **38**, 81–89 (2016).

36.   M. G. Buffone, T. W. Ijiri, W. Cao, T. Merdiushev, H. K. Aghajanian, G. L. Gerton,
Heads or tails? Structural events and molecular mechanisms that promote
mammalian sperm acrosomal exocytosis and motility. *Mol Reprod Dev*. **79**, 4–18
(2012).

37.   J. L. De La Vega-Beltran, C. Sánchez-Cárdenas, D. Krapf, E. O. Hernandez-
González, E. Wertheimer, C. L. Treviño, P. E. Visconti, A. Darszon, Mouse sperm
membrane potential hyperpolarization is necessary and sufficient to prepare sperm
for the acrosome reaction. *Journal of Biological Chemistry*. **287**, 44384–44393
(2012).

38.   H. Hasuwa, Y. Muro, M. Ikawa, N. Kato, Y. Tsujimoto, M. Okabe, Transgenic Mouse
Sperm that Have Green Acrosome and Red Mitochondria Allow Visualization of
Sperm and Their Acrosome Reaction in Vivo. *Exp. Anim*. **59**, 105–107 (2010).

39.   G. Corkidi, P. Hernandez-Herrera, F. Montoya, H. Gadêlha, A. Darszon, Long-term
segmentation-free assessment of head-flagellum movement and intracellular
calcium in swimming human sperm. *J Cell Sci*. **134** (2021),
doi:10.1242/JCS.250654.

40.   S. Culley, K. L. Tosheva, P. Matos Pereira, R. Henriques, SRRF: Universal live-cell
super-resolution microscopy. *Int J Biochem Cell Biol*. **101**, 74–79 (2018).

41.   G. Campagnola, K. Nepal, B. W. Schroder, O. B. Peersen, D. Krapf, Superdiffusive
motion of membrane-targeting C2 domains. *Sci Rep*. **5**, 17721 (2015).

42.   A. V. Weigel, B. Simon, M. M. Tamkun, D. Krapf, Ergodic and nonergodic processes
coexist in the plasma membrane as observed by single-molecule tracking.
Proceedings of the National Academy of Sciences. **108**, 6438–6443 (2011).

43. I. Izeddin, M. El Beheiry, J. Andilla, D. Ciepielewski, X. Darzacq, M. Dahan, PSF
shaping using adaptive optics for three-dimensional single-molecule super-
resolution imaging and tracking. *Opt Express.* **20**, 4957 (2012).
44. M. Ovesný, P. Křížek, J. Borkovec, Z. Švindrych, G. M. Hagen, ThunderSTORM: a
comprehensive ImageJ plug-in for PALM and STORM data analysis and super-
resolution imaging. *Bioinformatics.* **30**, 2389–2390 (2014).
45. G. M. Luque, X. Xu, A. Romarowski, M. G. Gervasi, G. Orta, J. L. De la Vega-
Beltrán, C. Stival, N. Gilio, T. Dalotto-Moreno, D. Krapf, P. E. Visconti, D. Krapf, A.
Darszon, M. G. Buffone, Cdc42 localized in the CatSper signaling complex
regulates cAMP-dependent pathways in mouse sperm. *The FASEB Journal.* **35**,
e21723 (2021).
46. C. Stival, C. Ritagliati, X. Xu, M. G. Gervasi, G. M. Luque, C. Baro Graf, J. L. Vega-
Beltran, N. I. Torres, A. Darszon, D. Krapf, M. G. Buffone, P. Visconti, D. Krapf, *J
Biol Chem*, in press, doi:10.1074/jbc.RA118.002286.
47. G. L. Nicolson, R. Yanagimachi, H. Yanagimachi, Ultrastructural localization of
lectin-binding sites on the zonae pellucidae and plasma membranes of mammalian
eggs. *J Cell Biol.* **66**, 263–274 (1975).
48. M. Jin, E. Fujiwara, Y. Kakiuchi, M. Okabe, Y. Satouh, S. A. Baba, K. Chiba, N.
Hirohashi, Most fertilizing mouse spermatozoa begin their acrosome reaction before
contact with the zona pellucida during in vitro fertilization. *Proc Natl Acad Sci U S A.*
108, 4892–4896 (2011).

759
760

Figures and tables

Figure 1. Sperm motility loss and FM4-64 fluorescence dynamics in acrosome-reacted
transgenic EGFP-DsRed2 sperm. A) Representative time series of transgenic EGFP-
DsRed2 sperm attached to concanavalin A-coated coverslips, with AE induced by 100 μM
progesterone. White squares indicate cells with spontaneous AE (prior to induction), while
pink squares highlight cells with progesterone-induced AE. A schematic representation of
AE in this transgenic model is shown on the right side of the panel. B) Representative time
series of transgenic EGFP-DsRed2 sperm that have already experienced AE, attached to
laminin-coated coverslips. The upper panel displays a cell with motility after AE, and the
lower panel shows an immotile cell. DsRed2 is presented in orange, and EGFP in green. C)
Quantification of motile and immotile acrosome-reacted sperm (EGFP-). A total of 235
cells were counted across at least three independent experiments. D) Representative time
series of transgenic EGFP-DsRed2 sperm stained with 10 μM FM4-64 and attached to
concanavalin A-coated coverslips, with AE induced by 100 μM progesterone. White
squares indicate cells exhibiting patterns I, II, or III after progesterone induction. A
schematic representation of AE in this transgenic model stained with FM4-64 is shown on

the right side of the panel. E-G) Representative images of capacitated transgenic EGFP-

DsRed2 sperm stained with 10 μM FM4-64. Panel E) displays an acrosome-intact, motile

sperm (Pattern I), F) shows an acrosome-reacted sperm with motility and low FM4-64

midpiece fluorescence (Pattern II), and G) presents an acrosome-reacted sperm with no

motility and high FM4-64 midpiece fluorescence (Pattern III). In all three cases, cells were

induced with 100 μM progesterone. Enlarged images of each pattern are shown on the right

panel. Representative images from at least five independent experiments are displayed.

Figure 2. Midpiece contraction coincides with the onset of AE. A) Left panel displays a

wide-field fluorescence image of capacitated CD1 sperm membrane stained with 0.5 μM

FM4-64, while the right panel shows its super-resolution SRRF reconstruction. B)

Representative time series of sperm midpiece with no AE (no reac), spontaneous exocytosis

(spont), progesterone (prog, 100 μM) and ionomycin-induced (iono, 10 μM) exocytosis,

respectively. Following acquisition, images were analyzed using SRRF. Insets in the sperm

head show wide-field images of AE. The midpiece diameter value is displayed in the

bottom left corner for each time point. C) Quantification of midpiece diameter changes for

each experimental group across time. Data is presented as a percentage of the initial

diameter value before induction for each cell. D-E) Quantification of FM4-64 fluorescence

in the sperm head and midpiece, respectively, for each experimental group across time.

Data is presented as times of increases compared to initial fluorescence before AE

induction. $*p<0.05$; $\#p<0.01$; $\$p<0.001$ and $\&p<0.0001$ compared to the non-reacted group.

A nonparametric Kruskal-Wallis test was performed in combination with Dunn's multiple

comparisons test. Representative images of at least 5 independent experiments are shown,

with 36 cells analyzed.

Figure 3. Contraction initiation preferentially occurs near the head-midpiece junction.

A) Schematic diagram illustrating the generation of super-resolution kymographs from

SRRF-processed images. Crosslines are drawn every 2.5 μm through the sperm midpiece,

and the Image-J Kymograph builder plug-in is used to create kymographs. The x-axis

represents time, and the y-axis shows diameter changes. For wide-field images, a line along

the midpiece is drawn to create fluorescence kymographs, with the y-axis representing

midpiece length. Three sections of the midpiece are defined: proximal (0-7 μm), central (7-

14 μm), and distal (14-21 μm). B) Relative frequency graph displaying the distribution of

the initiation sites for midpiece contractions in sperm with no AE (no reac), spontaneous

exocytosis (spont), progesterone-induced (prog, 100 μM) exocytosis, and ionomycin-

induced (iono, 10 μM) exocytosis, respectively. The x-axis indicates the midpiece section

where the contraction begins: proximal (P), central (C), or distal (D). A chi^2 test was

performed using the R language environment. C-D) Representative contraction kymographs

and diameter measurements for progesterone-induced (100 μM) AE with one or two

contraction initiation sites, respectively. In contraction kymographs, yellow lines demarcate

midpiece sections, and colored spots indicate where super-resolution kymographs were

created. Both kymograph and diameter measurement graphs display a dotted vertical line

marking the induction point. Data from at least 5 independent experiments are shown, with

36 cells analyzed.

Figure 4. Midpiece contraction is driven by $[Ca^{2+}]_i$ changes. A) The representative time

series demonstrates $[Ca^{2+}]_i$ and AE dynamics. Capacitated F1 sperm, loaded with Fluo4

AM, were immobilized on concanavalin A-coated coverslips and incubated in a recording

medium (rec) containing 10 μM FM4-64. Rec was added as indicated by arrowheads.

Beneath each frame in the Fluo4 (green) and FM4-64 (red) images, a color code displays

the normalized intensity of the fluorescence signal (scale bar on the right of the panel). B)

Kymograph-like analysis of the midpiece of 20 sperm following the addition of recording

medium. Each row depicting the $[Ca^{2+}]_i$ (upper) and membrane (lower) dynamics of a

single cell over time. A white dotted line indicates the moment of addition. The images

presented are representative of at least five independent experiments. C) The representative

time series demonstrates $[Ca^{2+}]_i$ and AE dynamics. Progesterone (prog, 100 μM) was added

as indicated by arrowheads. Beneath each frame in the Fluo4 (green) and FM4-64 (red)

images, a color code displays the normalized intensity of the fluorescence signal (scale bar

on the right of the panel). D) Kymograph-like analysis of the midpiece of 20 sperm

following the addition of prog. Each row depicting the $[Ca^{2+}]_i$ (upper) and membrane

(lower) dynamics of a single cell over time. A white dotted line indicates the moment of

addition. The images presented are representative of at least five independent experiments.

Consistently, a $[Ca^{2+}]_i$ transient precedes contraction, which is proportional to the increase

in FM4-64 fluorescence, as shown in Figure S2D. E) Representative time series of $[Ca^{2+}]_i$

and midpiece contraction dynamics. Capacitated CD1 sperm were loaded with Fluo4 AM,

immobilized on concanavalin A-coated coverslips, and incubated in a recording medium

containing 0.5 μM FM4-64. AE was induced with 100 μM progesterone (prog, arrowhead).

FM4-64 images are SRRF-processed, while Fluo4 images are widefield images. F) 3D

kymographs of $[Ca^{2+}]_i$ (left) and contraction (right) dynamics. Data is normalized to the

mean of the frames before the induction of AE. Representative images from at least 5

independent experiments are shown, with 36 cells analyzed.

Figure 5. The flagellar membrane approaches the actin cytoskeleton in the midpiece

of the sperm flagellum during midpiece contraction and AE. A-B) Representative time

series of plasma membrane and actin cytoskeleton colocalization in the midpiece in the

absence of AE (A) and during progesterone-induced AE (B, prog, 100 μM; FM4-64 shown

in red, SiR-actin shown in cyan). Capacitated CD1 sperm were loaded with 100 nM SiR-

actin, immobilized on concanavalin A-coated coverslips, and incubated in a recording

medium containing 0.5 μM FM4-64. C) Manders' colocalization coefficients for acrosome-

intact and acrosome-reacted cells in the midpiece. M1 was assigned to FM4-64, and M2 to

SiR-actin. Data is presented as mean ± SEM. Representative images from at least 5

independent experiments are shown, with 36 cells analyzed. D-G) Representative images of

sperm midpiece stained with the acrosome marker PNA (left panel, upper right insets,

epifluorescence) and phalloidin (actin filaments, STORM) for acrosome-intact (D-E) and

acrosome-reacted (F-G) cells. The left panel displays a longitudinal section of the midpiece,

while the right panel illustrates the radial distribution. E and G) Schematics of the analyzed

actin double helix parameters in the midpiece: helical pitch (*l*, distance between turns of the

helix, left panel), helical pitch frequency (f_0, number of turns the helix makes per 1 μm),

and radial distribution (R, radius of the double helix, right panel). Representative images

from at least 3 independent experiments are shown. Four acrosome-intact cells and seven

acrosome-reacted cells were analyzed.

Figure 6. Most sperm localized in the perivitelline space display midpiece contraction.

Representative images of IVF experiments using EGFP-DsRed2 sperm. Oocyte-sperm

complexes were stained with 10 μg/ml Hoechst and 10 μM FM4-64. DIC, Hoechst, EGFP,

DsRed2*, and FM4-64 images are shown for: (A) sperm attached to the ZP with an intact

acrosome and non-contracted midpiece; (B) sperm that have passed through the ZP,

exhibiting AE and contracted midpiece; (C) sperm that have passed through the ZP,

displaying AE with an initially non-contracted midpiece. After 20 minutes, as shown in

(D), the midpiece becomes contracted. The area depicted in the upper panel is shown in

higher magnification in the lower panel. E) Quantification of FM4-64 midpiece patterns in

sperm within the ZP. Representative images and data from at least 6 independent

experiments are shown. A total of 23 oocytes and 69 sperm were analyzed.

Figure 7. Contraction of the midpiece occurs after sperm-egg fusion. A) Schematic

representation of the acquisition settings for the sperm-oocyte fusion assay. Images were

taken every 7 μm along the z-axis. B) DIC image of a sperm-oocyte complex, with the area

depicted in higher magnification in panel C. C) Representative time series of sperm-oocyte

fusion assay experiments using EGFP-DsRed2 sperm. Oocytes were stained with 1 μg/ml

Hoechst and 10 μM FM4-64. DIC, Hoechst, EGFP, DsRed2*, and FM4-64 images are

shown over time. Note that midpiece contraction occurs after sperm-egg fusion and is

proportional to the increase in FM4-64 fluorescence, as shown in Figure S2D, highlighting

its potential importance in the fertilization process. Representative images from at least 4

independent experiments are displayed.

Figure 8. Midpiece contraction occurs in sperm-egg fusion after a decrease in $[Ca^{2+}]_i$.

A) Representative time series of sperm-oocyte fusion assay experiments using wild-type

sperm loaded with 1 μM Fluo-4. Oocytes were stained with 1 μg/ml Hoechst and 10 μM

FM4-64. DIC, Hoechst, Fluo-4, and FM4-64 images are shown over time. The color code

below each frame in the Hoechst (shown in blue), Fluo-4 (shown in green), and FM4-64

(shown in red) images indicates the normalized intensity of the fluorescence signal (scale

bar on the right of the panel). B-F) Quantification of sperm showing midpiece contraction

(B, indicated by increased FM4-64 fluorescence), midpiece folding (C), midpiece unfolding

(D), Fluo-4 fluorescence dynamics in the head (E), and different patterns in the midpiece

(F) during fusion. Data are presented as the mean ± SEM of the percentage of sperm

counted for each experiment. Representative images and data from at least 3 independent

experiments are shown. A total of 74 oocytes and 136 sperm were analyzed. Note that

midpiece contraction occurs in sperm-egg fusion after a decrease in Fluo-4 fluorescence,

suggesting a relationship between $[Ca^{2+}]_i$ dynamics and midpiece contraction. These

findings provide insights into the molecular mechanisms involved in sperm-egg fusion.

Figure 9: Proposed model of the structural reorganization of the sperm actin

cytoskeleton during key events of fertilization.

Figure 1

Figure 2

Figure 3

A

Time
Time

Proximal
Diameter
Central
Distal

Time
Time

Length across midpiece

Proximal
[0 - 7) µm

Central
[7 - 14) µm

Distal
[14 - 21] µm

B

● Single focus ● Multiple foci

Relative frequency (%)

prog
p=0.044

spont
p=0.021

iono
p=0.113

Section of midpiece

P C D   P C D   P C D

prog: 9, 6, 5, 4, 3
spont: 4, 7, 2, 1, 2, 1
iono: 10, 4, 8, 1, 5

C

Time (min)

Length across midpiece (µm)

single focus

5 min

1 µm

Normalized diameter
1.0   0.65

Normalized diameter

● 0 µm ● 2.5 µm ● 5 µm

Time (min)

D

Time (min)

Length across midpiece (µm)

two foci

3 min

1 µm

Normalized diameter
1.0   0.5

Normalized diameter

● 0 µm ● 2.5 µm ● 5 µm
● 7.5 µm ● 10 µm ● 12 µm

Time (min)

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A

DIC   1′   4′   11′   21′   36′   51′   56′   61′

Hoechst

Fluo4

FM4-64   10 µm

B   Midpiece contraction

Sperm (%)

Contraction   No contraction

C   Midpiece folding

Sperm (%)

Folding   No folding

D   Midpiece unfolding

Sperm (%)

Unfolding   No unfolding

E   Head $[Ca^{2+}]_i$

Sperm (%)

Decrease   No changes

F   Midpiece $[Ca^{2+}]_i$

Sperm (%)

Transitory increase   Decrease   No changes

Figure 9