Deciphering the genetic interactions between Pou4f3, Gfi1, and Rbm24 in maintaining cochlear hair cell survival

Guangqin Wang[1,2], Yunpeng Gu[1,2], Zhiyong Liu[1,2,3*]

1. Institute of Neuroscience, State Key Laboratory of Neuroscience, CAS Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences, Shanghai, 200031, China

2. University of Chinese Academy of Sciences, Beijing, 100049, China

3. Shanghai Center for Brain Science and Brain-Inspired Intelligence-Technology, Shanghai, 201210, China

* Corresponding author:

Dr. Zhiyong Liu; E-mail: Zhiyongliu@ion.ac.cn

Tel: 0086-21-54921793 (Office)

Key words: Pou4f3, Gfi1, Rbm24, hair cell, cochlea

ABSTRACT

Mammals harbor a limited number of sound-receptor hair cells (HCs) that cannot be regenerated after damage. Thus, investigating the underlying molecular mechanisms that maintain HC survival is crucial for preventing hearing impairment. Intriguingly, $Pou4f3^{-/-}$ or $Gfi1^{-/-}$ HCs form initially but then rapidly degenerate, whereas $Rbm24^{-/-}$ HCs degenerate considerably later. However, the transcriptional cascades involving Pou4f3, Gfi1, and Rbm24 remains undescribed. Here, we demonstrate that $Rbm24$ expression is completely repressed in $Pou4f3^{-/-}$ HCs but unaltered in $Gfi1^{-/-}$ HCs, and further that the expression of both POU4F3 and GFI1 is intact in $Rbm24^{-/-}$ HCs. Moreover, by using *in vivo* mouse transgenic reporter assays, we identify three $Rbm24$ enhancers to which POU4F3 binds. Lastly, through *in vivo* genetic testing of whether Rbm24 restoration alleviates the degeneration of $Pou4f3^{-/-}$ HCs, we show that ectopic Rbm24 alone cannot prevent $Pou4f3^{-/-}$ HCs from degenerating. Collectively, our findings provide new molecular and genetic insights into how HC survival is regulated.

INTRODUCTION

Housed in the mammalian cochlea, the auditory epithelium—also known as the organ of Corti—contains the mechanosensory receptors hair cells (HCs) that detect sound information [1, 2]. Two types of HCs exist, the inner HCs (IHCs) and outer HCs (OHCs), and these appear in, respectively, a single row and three rows [3]. Both OHCs and IHCs are derived from cochlear progenitors expressing high levels of ATOH1, a master transcription factor (TF) in HC development [4-6], and no HCs form in the $Atoh1^{-/-}$ cochlea, highlighting the essential role of ATOH1 in specifying the general HC fate in the undifferentiated progenitors [7, 8]. Whereas OHCs uniquely express the motor protein PRESTIN (encoded by $Slc26a5$) and function as sound amplifiers [9, 10], IHCs specifically express FGF8, vGLUT3 (encoded by $Slc17a8$), and OTOFERLIN [11-15] and act as the primary sensory cells that form synapses with distinct subtypes of type I spiral (auditory) neurons [16-19]. Moreover, INSM1 is necessary for stabilizing the OHC fate, and ~50% of $Insm1^{-/-}$ OHCs tend to transdifferentiate into IHC-like cells [20, 21]; conversely, IKZF2 is not required in early OHC development but is essential for consolidating the OHC fate, and $Ikzf2^{cello/cello}$ mutant OHCs are dysfunctional and misexpress IHC genes [22]. INSM1 is an epistatic but indirect positive regulator of $Ikzf2$, and restoration of IKZF2 expression can partially mitigate the phenotypes of $Insm1^{-/-}$ OHCs [21]. In contrast to INSM1 and IKZF2, TBX2 is required in IHC fate specification, differentiation, and maintenance, and IHCs convert into OHCs in the absence of TBX2 [23-26].

The survival of cochlear HCs must be maintained after their production. POU4F3, a POU-domain family TF encoded by the hearing-loss gene $DFNA15$, is essential for HCs to

survive [27-31]. However, unlike ATOH1, POU4F3 is not involved in the fate determination

of HCs [32]. As a target gene regulated by ATOH1 [28, 33], *Pou4f3* is expressed in HCs

between embryonic day 14.5 (E14.5) and E16 in a basal-to-apical gradient, and POU4F3 then

activates the expression of *Gfi1*, which encodes a zinc-finger TF [34, 35]; Gfi1 is

undetectable in *Pou4f3*-deficient HCs [35], and HCs, particularly OHCs, degenerate in

Pou4f3 and *Gfi1* mutants by birth [35]. Besides POU4F3 and Gfi1, the RNA-binding protein

RBM24 is also specifically expressed in cochlear HCs [36]. We recently showed that RBM24

is indispensable for maintaining OHC survival, but the OHC death in *Rbm24* mutants occurs

considerably later than in *Pou4f3* and *Gfi1* mutants; OHCs develop normally until birth but

most OHCs undergo cell death by postnatal day 19 (P19) in *Rbm24*$^{-/-}$ mice [37]. How *Rbm24*

expression is regulated during cochlear HC development is unknown.

The similar but delayed phenotype of OHC death in *Rbm24* mutants relative to that in

Pou4f3- and *Gfi1*-deficient mice prompted us to dissect the potential genetic interactions

among them by using an *in vivo* genetic approach. Our results showed that the onset of

RBM24 expression was completely repressed in *Pou4f3*$^{-/-}$ HCs, but that, unexpectedly,

RBM24 expression was normal in *Gfi1*$^{-/-}$ HCs. Moreover, the expression of neither POU4F3

nor GFI1 was altered in *Rbm24*$^{-/-}$ HCs. Thus, POU4F3, but not GFI1, is required in *Rbm24*

expression. Furthermore, we identified three *Rbm24* enhancers that were sufficient to drive

specific EGFP reporter expression in HCs, and these enhancers are likely bound by POU4F3.

Lastly, we found that restoration of RBM24 expression alone cannot alleviate the

degeneration of *Pou4f3*$^{-/-}$ HCs, which indicates that the expression of additional

POU4F3-targeted genes must be restored to enable *Pou4f3*$^{-/-}$ HCs to survive. Our study

provides new insights into the genetic interactions among *Pou4f*, *Gfi1*, and *Rbm24*, which

hold potential applications in HC protection.

RESULTS

RBM24 expression is completely repressed in *Pou4f3$^{-/-}$* cochlear HCs

HC degeneration occurs in both *Pou4f3$^{-/-}$* and *Rbm24$^{-/-}$* mice, with the phenotype

appearing earlier and being more severe in *Pou4f3$^{-/-}$* mice [35, 37]. This led us to speculate

that genetic interaction exists between *Pou4f3* and *Rbm24*. We reasoned that if POU4F3 is an

upstream positive regulator of *Rbm24*, RBM24 expression would be downregulated in the

absence of POU4F3. To rapidly test this possibility, we exploited our previously established

CRISPR-stop approach [38]. CRISPR-stop allows early stop codons to be introduced without

inducing DNA damage through Cas9, which can cause deleterious effects [39]. More

importantly, the CRISPR-stop approach can generate Founder 0 (F0) mice carrying

homozygous or mosaic homozygous gene mutations, and the F0 mice are thus immediately

ready for phenotypic analysis, considerably faster than in the case with traditional

gene-targeting methods [37, 38]. Co-injecting one sgRNA (sgRNA-1) against *Pou4f3* and

base-editor components into one-cell-stage mouse zygotes yielded F0 mice, whose tail DNA

was subject to Sanger sequencing (Figure 1-figure supplement 1A). Relative to wild-type

(WT) mice (Figure 1-figure supplement 1B), the F0 mice with homozygous premature

emergence of the stop codon TAG were defined as *Pou4f3$^{-/-}$* mice and were immediately

ready for phenotypic analysis (Figure 1-figure supplement 1C).

Triple staining of POU4F3, RBM24, and INSM1 revealed that OHCs in WT mice at

E16.5 were POU4F3+/RBM24+/INSM1+, whereas IHCs expressed POU4F3 and RBM24

but not INSM1 (Figure 1-figure supplement 1D-D'''). Conversely, POU4F3 expression was absent in *Pou4f3*$^{-/-}$ mice (Figure 1-figure supplement 1E-E'''), confirming that POU4F3 translation was blocked, and unlike in WT mice, RBM24 expression was undetectable in the *Pou4f3*$^{-/-}$ mice at E16.5 (Figure 1-figure supplement 1E-E'''). Notably, INSM1 expression appeared normal in *Pou4f3*$^{-/-}$ mice (arrows in Figure 1-figure supplement 1E-E'''). Moreover, the presence of INSM1+ OHCs in the *Pou4f3*$^{-/-}$ mice eliminated the possibility that the absence of RBM24 was due to a secondary effect of OHC death or delayed differentiation, and also agreed with previous reports that initial HC differentiation can occur without POU4F3 [32, 35]. Thus, the rapid *Pou4f3* loss-of-function analysis by using CRISPR-stop supported our hypothesis that POU4F3 is required for turning on RBM24 expression.

To further validate the observation described above, we constructed germ-line-stable *Pou4f3* mutants in which the entire *Pou4f3* genomic region between sgRNA-2 and sgRNA-3 was deleted (Figure 1A). The three obtained genotypes, *Pou4f3*$^{+/+}$, *Pou4f3*$^{+/-}$, and *Pou4f3*$^{-/-}$, were readily distinguished using tail-DNA PCR (Figure 1B). Relative to WT mice (Figure 1C), *Pou4f3*$^{-/-}$ mice again completely lacked RBM24 expression at E16.5 (Figure 1D). Similar to INSM1, BCL11B is an OHC marker [20], and triple staining of RBM24, POU4F3, and BCL11B revealed that as in WT HCs (Figure 1E-E'''), BCL11B was normally expressed in OHCs lacking both POU4F3 and RBM24 at E16.5 (Figure 1F-F'''). This result again confirmed that nascent OHCs were normally produced in the absence of POU4F3. Moreover, in contrast to WT mice at P1 (Figure 1-figure supplement 2A-A''), the germline *Pou4f3*$^{-/-}$ mutants showed HC degeneration at P0 (Figure 1-figure supplement 2B-B'') and P1 (Figure 1-figure supplement 2C-C''). Normalizing the numbers of the remaining HCs in the *Pou4f3*$^{-/-}$

mice at P0 or P1 against those in WT mice at P1 revealed that in P0 *Pou4f3*$^{-/-}$ mice, 35.94% ±

4.52%, 38.16% ± 2.79%, and 63.8% ± 5.28% of the HCs survived in basal, middle, and

apical cochlear turns, respectively, whereas in P1 *Pou4f3*$^{-/-}$ mice, 5.54% ± 3.11%, 15.79% ±

3.65%, and 65.64% ± 7.95% of the HCs survived. HC degeneration in the middle and basal

turns, but not the apical turn, was significantly more severe at P1 than at P0 (Figure 1-figure

supplement 2D). Lastly, unlike in WT cochleae in which all MYO7A+ HCs expressed

RBM24 (Figure 1-figure supplement 2E-E''), in *Pou4f3*$^{-/-}$ cochleae, the remaining HCs

lacked RBM24 expression (arrows in Figure 1-figure supplement 2F-F''). Collectively, our

results obtained using two distinct *Pou4f3*-deficient models indicated that POU4F3 is

indispensable for turning on RBM24 expression. However, whether GFI1 is also essential for

RBM24 expression remained unknown.

Construction of *Gfi1-3×HA-P2A-Cre/+* knockin mouse strain

We sought to determine whether POU4F3 regulates *Rbm24* expression through GFI1 or

independently of GFI1. Because a suitable commercial GFI1 antibody for immunostaining

was unavailable, we constructed a knockin mouse strain, *Gfi1*$^{3×HA-P2A-Cre/+}$ (*Gfi1*$^{HA-Cre/+}$ in

brief), by using our routine CRISPR/Cas9 approach (Figure 2-figure supplement 1A-C); here,

the GFI1 C-terminus was tagged with three Hemagglutinin (HA) fragments and Cre

expression was under the control of endogenous *cis*-regulatory elements (CREs) of *Gfi1*. The

obtained WT (*Gfi1*$^{+/+}$), heterozygous (*Gfi1*$^{HA-Cre/+}$), and homozygous (*Gfi1*$^{HA-Cre/HA-Cre}$) mice

were readily distinguished using tail-DNA PCR (Figure 2-figure supplement 1D). Southern

blotting revealed that in addition to being inserted in the *Gfi1* locus, the targeting vector

(Figure 2-figure supplement 1B) was randomly inserted in an unknown genomic region;

however, the random insertion likely occurred in a silent genomic region as indicated by the

analysis discussed below.

Dual staining of HA (GFI1) and the pan-HC marker MYO7A revealed that, similar to

$Gfi1^{+/+}$ mice (Figure 2-figure supplement 1E-E''), $Gfi1^{HA-Cre/HA-Cre}$ mice showed normal HC

development at P1 (Figure 2-figure supplement 1F-F''). Moreover, HA (GFI1) was highly

expressed in all MYO7A+ IHCs and OHCs in $Gfi1^{HA-Cre/HA-Cre}$ mice (arrows in Figure

2-figure supplement 1F-F''). This finding confirmed that Gfi1 endogenous expression is

unaffected in $Gfi1^{HA-Cre/HA-Cre}$ mice, which represents an advantage over the previous $Gfi1^{Cre/+}$

strain in which one copy of $Gfi1$ is lost and early onset of hearing loss occurs [40, 41].

We   also   performed   fate-mapping   analysis   in $Gfi1^{HA-Cre/+}$;

$Rosa26$-$loxp$-$stop$-$loxp$-$tdTomato$ (Ai9)/+ ($Gfi1^{HA-Cre/+}$; Ai9/+) mice at P2. The $Gfi1^{HA-Cre/+}$;

Ai9/+ model allowed us to both visualize the temporal GFI1 protein-expression pattern and

permanently mark cells expressing $Gfi1$ by using tdTomato. Neither tdTomato nor HA was

detected in the MYO7A+ HCs of control Ai9/+ mice (Figure 2-figure supplement 1G-G'''

and I-I''), whereas the majority of MYO7A+ HCs expressed tdTomato and HA in the

$Gfi1^{HA-Cre/+}$; Ai9/+ mice (Figure 2-figure supplement 1H-H''' and J-J''). HA (GFI1) was

generally expressed more highly in IHCs than OHCs (Figure 2-figure supplement 1J-J''), and

the HA (GFI1) expression level exhibited manifest heterogeneity among the OHCs. We

identified the OHCs expressing the highest (#1), intermediate (#2), and lowest (#3) levels of

HA (GFI1) (arrows in Figure 2-figure supplement 1J-J''), and we observed that, notably, such

heterogenous GFI1 expression in neonatal HCs was not detected in $Gfi1^{HA-Cre/HA-Cre}$ mice

(Figure 2-figure supplement 1F-F''), likely due to the higher level of HA-tagged GFI1 in

$Gfi1^{HA-Cre/HA-Cre}$ than $Gfi1^{HA-Cre/+}$ mice. Besides the tdTomato+ HCs, we detected tdTomato+

cells in nonsensory regions (arrows in Figure 2-figure supplement 1H), consistent with a

previous report [40]. Collectively, these results indicated that the $Gfi1^{HA-Cre/+}$ strain was

suitable for visualizing GFI1 protein by using an anti-HA antibody, although random

insertion of the targeting vector also exists here. Alternatively, $Gfi1^{HA-Cre/+}$ can at least be

treated as a pseudo-transgenic mouse strain that can be used to reliably visualize GFI1 and

trace cells expressing $Gfi1$.

GFI1 expression is prevented in $Pou4f3^{-/-}$ cochlear HCs

Because HA faithfully represented GFI1 expression in the $Gfi1^{HA-Cre/+}$ strain, we

expected to observe repression of HA expression in $Pou4f3^{-/-}$ HCs, as reported previously [34,

35]. We confirmed this by using our CRISPR-stop approach [38]. The experimental pipeline

was mostly identical to that used for producing the $Pou4f3$ mutants (Figure 1-figure

supplement 1), except that the zygotes here were derived from male $Gfi1^{HA-Cre/+}$ mice (Figure

2-figure supplement 2A). We obtained mosaic (or chimeric) $Pou4f3^{-/-}$ mice, as confirmed

through Sanger sequencing of tail-DNA PCR samples (Figure 2-figure supplement 2B and C),

partly because the injection time was the late stage of the one-cell-stage zygotes. In control

$Gfi1^{HA-Cre/+}$ mice, a pure "C" base was present (blue arrow in Figure 2-figure supplement 2B),

but a mixture of "C" and "T" double peaks existed in $Gfi1^{HA-Cre/+}$; $Pou4f3^{-/-}$ (mosaic) mice

(red arrow in Figure 2-figure supplement 2C).

In control $Gfi1^{HA-Cre/+}$ mice at E16.5, all HCs expressed POU4F3, RBM24, and HA

(GFI1), although the HA (GFI1) levels again appeared heterogenous among the OHCs

(Figure 2-figure supplement 2D-D'''). By contrast, in the $Gfi1^{HA-Cre/+}$; $Pou4f3^{-/-}$ (mosaic)

mice, we detected HCs that had either lost or maintained POU4F3 expression (Figure

2-figure supplement 2E-E'''). Notably, POU4F3+ HCs expressed HA (GFI1) and RBM24

(blue arrows in Figure 2-figure supplement 2E-E'''), whereas both HA (GFI1) and RBM24

were absent in HCs that had lost POU4F3 expression (orange arrows in Figure 2-figure

supplement 2E-E'''). This again confirmed that HA is a reliable readout for GFI1 expression

and that the expression is sensitive to the loss of POU4F3. Moreover, the results supported

the view that POU4F3 regulates *Rbm24* in a cell-autonomous manner.

GFI1 is dispensable for RBM24 expression

Next, we produced *Gfi1$^{-/-}$* mutants by using the same CRISPR-stop approach [38].

One-cell-stage zygotes derived from male *Gfi1$^{HA-Cre/+}$* mice were injected with base-editor

components and four different sgRNAs located in distinct exons of *Gfi1* (Figure 2A). The

reason for the combined use of four sgRNAs is detailed in the discussion section. With

sgRNA-6, for example, Sanger sequencing of tail DNA revealed that relative to control

Gfi1$^{HA-Cre/+}$ mice (black arrow in Figure 2B), F0 mice showed premature emergence of the

TAG stop codon, which resulted in homozygous Gfi1 inactivation (red arrow in Figure 2C).

All HCs expressed HA (GFI1) and RBM24 in control *Gfi1$^{HA-Cre/+}$* mice (Figure 2D-D''), and,

notably, RBM24 expression was maintained in *Gfi1$^{-/-}$* HCs, which showed no HA (GFI1)

expression at E16.5 (Figure 2E-E''). This finding suggested that RBM24 expression does not

depend on GFI1.

The advantage of this *Gfi1$^{-/-}$* model is that it allows rapid and direct confirmation of the

absence of GFI1 expression in F0 mice. However, complete loss of GFI1 expression in all

HCs cannot be guaranteed here at single-cell resolution through HA staining, because only

one allele of *Gfi1* was *Gfi1^HA-Cre^*. Thus, to eliminate the possibility that only the *HA*-tagged

Gfi1 allele was mutated, we generated germ-line-stable *Gfi1* mutants in which the majority of

the *Gfi1* DNA fragment between sgRNA-4 and sgRNA-9 was deleted (Figure 2-figure

supplement 3A). The WT (*Gfi1^+/+^*), *Gfi1^+/-^*, and *Gfi1^-/-^* mice were readily identified using

tail-DNA PCR (Figure 2-figure supplement 3B). In agreement with previous findings [42],

dual staining of RBM24 and POU4F3 revealed that relative to WT mice (Figure 2-figure

supplement 3C-C"), *Gfi1^-/-^* mice (Figure 2-figure supplement 3D-D") showed severe

degeneration of HCs at P1, particularly OHCs; this validated the successful generation of the

*Gfi1^-/-^* mice. Notably, the surviving POU4F3+ HCs maintained the expression of RBM24

(orange arrows in Figure 2-figure supplement 3D-D"). By contrast, we observed no

significant difference between WT and *Gfi1^-/-^* mice at E16.5: both WT and *Gfi1^-/-^* HCs

expressed POU4F3 and RBM24, and both WT and *Gfi1^-/-^* OHCs expressed BCL11B (Figure

2-figure supplement 3E-F'''). This suggests that the degeneration of *Gfi1^-/-^* HCs does not

begin by E16.5. The presence of POU4F3 in *Gfi1^-/-^* HCs agreed with the previously

mentioned notion that *Pou4f3* is epistatic to *Gfi1*. Collectively, the results of analyses of both

Gfi1 mutant models support the conclusion that GFI1 is dispensable for RBM24 expression.

POU4F3 and GFI1 expressions are normal in *Rbm24^-/-^* cochlear HCs

Considering that POU4F3 is upstream and a positive regulator of *Rbm24*, we predicted

that POU4F3 expression would be unaffected in *Rbm24^-/-^* HCs. To test this, we established an

*Rbm24^-/-^* mouse strain by injecting one *Rbm24* sgRNA (sgRNA-10) into one-cell-stage

zygotes derived from male *Gfi1^HA-Cre/+^* mice (Figure 3A). We selected *Gfi1^HA-Cre/+^* zygotes in

order to concurrently assess GFI1 and POU4F3 expression patterns in *Rbm24^-/-^* HCs. Relative

to control *Gfi1*^{HA-Cre/+}*; Rbm24*^{+/+} mice (Figure 3B), *Gfi1*^{HA-Cre/+}*; Rbm24*^{-/-} (mosaic) mice

(Figure 3C) showed premature emergence of the TAG stop codon. Notably, Sanger

sequencing of tail DNA revealed that the *Rbm24* mutation was mosaic, because mixed "T"

and "C" peaks existed (red arrow in Figure 3C).

The mosaic inactivation of RBM24 was further confirmed through triple staining of

RBM24, HA, and POU4F3 (Figure 3D-E'''). Relative to the expression in control *Gfi1*^{HA-Cre/+}*;*

Rbm24^{+/+} mice (Figure 3D-D'''), RBM24 expression disappeared in a fraction of cochlear

HCs (orange arrows in Figure 3E-E''') but remained normal in other HCs (blue arrows in

Figure 3E-E''') of the *Gfi1*^{HA-Cre/+}*; Rbm24*^{-/-} (mosaic) mice at E17. Notably, regardless of

whether RBM24 expression was inactivated or not in the cochlear HCs, the expression

patterns of both HA (GFI1) and POU4F3 remained intact. This suggests that *Pou4f3* is

epistatic to *Rbm24* and that inactivation of the *Rbm24* gene does not affect POU4F3

expression. Moreover, our results suggested that *Gfi1* and *Rbm24* do not interact genetically:

Gfi1 inactivation did not affect *Rbm24* expression (Figure 2 and Figure 2-figure supplement

3), and vice versa (Figure 3E-E''').

GATA3 is downregulated and IKZF2 is upregulated in *Rbm24*-deficient HCs

We next determined whether the expression patterns of two additional genes, *Gata3* and

Ikzf2, which are involved in OHC development and survival [21-23, 43, 44], were altered in

the absence of RBM24. For analyzing GATA3, mosaic *Rbm24* mutant mice were produced

using the CRISPR-stop approach (Figure 3-figure supplement 1A). Dual immunostaining of

GATA3 and RBM24 revealed that GATA3 was evenly expressed in control OHCs and IHCs

at P1 (Figure 3-figure supplement 1B-B''). By contrast, the GATA3 level in OHCs lacking

RBM24 expression (white arrows in Figure 3-figure supplement 1C-C") was manifestly

weaker than that in neighboring OHCs that retained RBM24 expression (yellow arrows in

Figure 3-figure supplement 1C-C"). This result suggests that GATA3 is downregulated in

Rbm24-deficient OHCs. OHCs degenerate in *Gata3*$^{+/-}$ mice [43], and thus the downregulation

of GATA3 might contribute to the cell death of *Rbm24*-deficient OHCs.

For analyzing IKZF2, mosaic *Rbm24* mutant mice were produced using zygotes obtained

by breeding male *Ikzf2*$^{V5/+}$ mice with WT female mice (Figure 3-figure supplement 1D).

Notably, the C-terminus of IKZF2 here was fused with three V5 tags to allow IKZF2 protein

to be visualized using a V5 antibody [21, 23]. In control *Ikzf2*$^{V5/+}$; *Rbm24*$^{+/+}$ mice, V5

(IKZF2) was evenly distributed in OHCs at P5 (Figure 3-figure supplement 1E-E"). By

contrast, in the mosaic *Ikzf2*$^{V5/+}$; *Rbm24*$^{-/-}$ mice, the V5 signal in the OHCs that had lost

RBM24 expression (yellow arrows in Figure 3-figure supplement 1F-F") was unfailingly

higher than the nearby OHCs expressing RBM24 (white arrows in Figure 3-figure

supplement 1F-F"). It supports that RBM24 directly or indirectly represses the IKZF2

expression.

Three *Rbm24* enhancers can drive specific EGFP expression in cochlear HCs

After showing that POU4F3, but not GFI1, is indispensable in mediating *Rbm24*

expression, we determined the mechanism by which POU4F3 controls *Rbm24* expression,

particularly the role of the CREs of *Rbm24*. According to our previous high-throughput

ATAC-seq (transposase-accessible chromatin sequencing) analysis of neonatal cochlear HCs

[8], four CREs of *Rbm24* were identified: one proximal promoter (arrow in Figure 4A), and

three distal potential enhancers that were defined as Eh1, Eh2, and Eh3 (dotted boxes in

Figure 4A). Eh1 and Eh2 were located upstream and Eh3 was downstream of the *Rbm24*

coding region. Moreover, we reanalyzed the results of POU4F3 Cut&Run assays from one

previous study [33], and we found that POU4F3 binds to Eh1, Eh2, and Eh3 but not the

Rbm24 promoter (Figure 4A).

Whether Eh1, Eh2, and Eh3 are bona-fide *Rbm24* enhancers has remained unknown. We

reasoned that if Eh1, Eh2, and Eh3 were *Rbm24* enhancers, one of these, together with the

mini-promoter of mouse heat shock protein 68 gene (*Hsp68*) [8, 45-47], would be sufficient

to drive specific reporter expression in cochlear HCs, and to test this, we established three

transgenic mouse strains: Eh1-EGFP+ (Figure 4B-C''), Eh2-EGFP+ (Figure 4D-E''), and

Eh3-EGFP+ (Figure 4F-G''); in these strains, EGFP expression would be driven by the

mini-promoter of *Hsp68* and Eh1 or Eh2 or Eh3, respectively. The mini-promoter of *Hsp68*

alone is reported to be incapable of driving EGFP expression [48], and in contrast to this, we

detected strong EGFP expression through whole-mount analysis in all three transgenic lines

(Figure 4B, D and F). Moreover, dual labeling for EGFP and MYO7A in cryosectioned

cochleae showed that EGFP was specifically expressed in IHCs and OHCs in all three strains

at P1 (Figure 4C-C'', E-E'' and G-G''). Collectively, our transgenic assay results suggest that

Eh1, Eh2, and Eh3 are *Rbm24* enhancers, and further that POU4F3 regulates *Rbm24*

expression primarily by binding to the *Rbm24* enhancers.

Forced RBM24 expression fails to alleviate the degeneration of *Pou4f3*$^{-/-}$ HCs

Our results thus far clearly showed that POU4F3 can regulate *Rbm24* expression, but

whether ectopic RBM24 expression can alleviate the degeneration of *Pou4f3*$^{-/-}$ HCs remained

unknown. Thus, we first established a new conditional mouse model,

Rosa26$^{loxp-stop-loxp-Rbm24*3×HA/+}$ (*Rosa26$^{Rbm24/+}$* in short) (Figure 5A and B), and to turn on

ectopic RBM24 expression in cochlear HCs, we further crossed the *Rosa26$^{Rbm24/+}$* strain with

the strain *Atoh1$^{Cre/+}$*, which targets a majority of cochlear HCs and a fraction of supporting

cells (SCs) [49]. Notably, in the *Rosa26$^{Rbm24/+}$* strain, RBM24 was tagged with three HA

fragments, which allowed endogenous and ectopic RBM24 to be distinguished. In control

Pou4f3$^{+/+}$ mice (Figure 5C-C") and germ-line-stable *Pou4f3$^{-/-}$* mice (Figure 5D-D"),

MYO7A+ HCs did not express HA (RBM24), whereas almost all MYO7A+ HCs expressed

HA (RBM24) in *Pou4f3$^{+/+}$; Atoh1$^{Cre/+}$; Rosa26$^{Rbm24/+}$* mice at P1 (white arrows in Figure

5E-E"). As expected, besides HCs, we detected a few SCs expressing HA (RBM24) but not

MYO7A (orange arrows in Figure 5E-E"). The cochlear HCs expressing HA (RBM24)

appeared normal, suggesting that the cells can tolerate the additional RBM24 expression by

P1. Collectively, these results supported the conclusions that the *Rosa26$^{Rbm24/+}$* model was

successfully generated and that ectopic RBM24 was expressed in cochlear HCs in *Pou4f3$^{+/+}$;*

Atoh1$^{Cre/+}$; Rosa26$^{Rbm24/+}$ mice.

In contrast to our expectation, ectopic RBM24 expression failed to mitigate the

degeneration of *Pou4f3$^{-/-}$* HCs. Severe degeneration of HCs (both IHCs and OHCs) occurred

in *Pou4f3$^{-/-}$; Atoh1$^{Cre/+}$; Rosa26$^{Rbm24/+}$* mice (Figure 5F-F") and was indistinguishable from

that in *Pou4f3$^{-/-}$* mice (Figure 5D-D"), except that the surviving HCs expressed RBM24 in

the *Pou4f3$^{-/-}$; Atoh1$^{Cre/+}$; Rosa26$^{Rbm24/+}$* mice (arrows in Figure 5F-F"). This result suggests

that restoration of RBM24 expression alone is not sufficient to prevent *Pou4f3$^{-/-}$* cochlear

HCs from undergoing degeneration.

DISCUSSION

Molecular mechanisms underlying cochlear HC survival

Mammalian sound-receptor HCs are vulnerable to various genetic mutations, environmental ototoxic factors, and aging. HC degeneration is one of the primary reasons for human sensorineural hearing impairment [50], and several genes whose mutations lead to HC degeneration starting at different ages have been identified previously. For example, HC development is severely defective in $Atoh1^{-/-}$ mutants [7, 51], and $Atoh1^{-/-}$ cochlear sensory cells undergo apoptosis. However, whether the absence of HCs in $Atoh1^{-/-}$ mice is because of HCs not being produced or dying immediately after initial emergence remains unclear, partly due to the difficulty of unambiguously defining nascent HCs by using molecular markers.

Unlike in $Atoh1^{-/-}$ mutants, in both $Pou4f3^{-/-}$ and $Gfi1^{-/-}$ mice [32, 34, 35], initial cochlear HC development is normal but becomes defective at perinatal ages, consistent with the notion that POU4F3 and GFI1 are dispensable for HC fate specification but necessary for subsequent differentiation and survival. Intriguingly, caspase-3 is active during HC death in $Pou4f3^{-/-}$ mutants, and the antiapoptotic factor z-VAD-fmk exerts a protective effect on $Pou4f3^{-/-}$ HCs between E14.5 and E16.5 [52]. Moreover, we noted here milder overall HC degeneration in $Gfi1^{-/-}$ mice than in $Pou4f3^{-/-}$ mice at P1, because only OHC degeneration occurred in $Gfi1^{-/-}$ mice but both IHCs and OHCs were degenerated in $Pou4f3^{-/-}$ mice (Figure 1-figure supplement 2 and Figure 2-figure supplement 3).

$Gfi1$ is a recognized target of POU4F3 [35], but whether forced GFI1 expression can alleviate the degeneration of $Pou4f3^{-/-}$ HCs is unknown. Similarly, $Pou4f3$ is one of the target genes regulated by ATOH1 [33], but whether forced expression of POU4F3 or GFI1, or both,

can mitigate the developmental defects of *Atoh1*[-/-] HCs remains undetermined. Future studies

must address these questions to enable comprehensive understanding of the mechanisms

underlying the maintenance of cochlear HC survival.

Roles of RBM24 and its regulation during cochlear HC development

Shortly after cochlear HCs emerge, RBM24 expression begins and then is maintained

permanently, as revealed by transcriptomic analyses and antibody staining [23, 36, 44, 53, 54].

RBM24 is not necessary in the early phase of cochlear HC development, because *Rbm24*[-/-]

HC development is normal by P1, but RBM24 is required for OHC survival after birth [37];

Rbm24[-/-] OHCs, but not IHCs, are degenerated by P19 [37]. RBM24 is also involved in

mRNA stability and pre-mRNA alternative splicing of several genes, including *Cdh23* and

Pcdh15, which are crucial for the development of HC stereocilia [55-57]. Our study also

revealed that GATA3 expression is decreased and IKZF2 expression is increased in OHCs in

the absence of RBM24, which could be a direct or indirect effect of RBM24 loss.

Nonetheless, single-cell transcriptomic analysis has not yet been performed on *Rbm24*[-/-] IHCs

or OHCs, and future investigation is necessary to clarify the detailed molecular mechanism

underlying *Rbm24*[-/-] HC degeneration.

What are the *trans*-acting factors involved in regulating *Rbm24* expression? First, this

study has provided strong genetic evidence indicating that POU4F3 is necessary for turning

on *Rbm24* expression; in the absence of POU4F3, *Rbm24* expression was not triggered.

Moreover, the normal expression of POU4F3 in *Rbm24*[-/-] HCs confirmed that *Pou4f3* is

epistatic to *Rbm24* and that RBM24 is dispensable for *Pou4f3* expression. Besides POU4F3,

ATOH1 appears to regulate *Rbm24* expression, as per two lines of evidence: one, RBM24

expression is lost in *Atoh1*$^{-/-}$ cochlear HCs [58]; and two, *Rbm24* is one of the ATOH1

binding targets revealed by the ATOH1 Cut&Run assay [8, 33]. Notably, ATOH1 binds to

both the *Rbm24* promoter and the three *Rbm24* enhancers (Eh1, Eh2, and Eh3) [8, 33],

whereas POU4F3 only binds to the *Rbm24* enhancers (Figure 4). Thus, POU4F3 and ATOH1

likely cooperate to regulate *Rbm24* expression, and either *Pou4f3* or *Atoh1* mutation leads to

repression of *Rbm24* expression. Future conditional *Pou4f3* loss-of-function studies are

necessary to determine whether *Rbm24* expression in adult HCs requires POU4F3.

Gfi1 and _Rbm24_ are expressed independently of each other

We initially hypothesized that POU4F3 regulates *Rbm24* expression through GFI1.

However, this hypothesis was not supported by our observation that RBM24 expression is

normal in *Gfi1*$^{-/-}$ HCs (Figure 2 and Figure 2-figure supplement 3). Moreover, in the absence

of RBM24, GFI1 expression was not altered in cochlear HCs. Thus, *Rbm24* expression and

Gfi1 expression appear to be independent of each other. This might be due to the functional

difference between POU4F3 and GFI1 during cochlear HC development. Although both

POU4F3 and GFI1 are necessary for promoting the expression of genes involved in HC

differentiation [29, 31, 35, 42], GFI1, but not POU4F3, also represses the preceding

expression of neural genes in nascent HCs [42].

Another observation here relevant to *Gfi1* isoforms is noteworthy. We showed previously

that one efficient sgRNA is adequate for inducing homozygous gene inactivation by using

CRISPR-stop [37]. Here, we used four *Gfi1* sgRNAs distributed across distinct exons (Figure

2A). Interestingly, we successfully established one *Gfi1* mutant by using sgRNA-5 alone, and

this, in principle, should effectively pre-stop *GFI1* translation in exon 1, which codes for the

SNAG repressor domain; the obtained *Gfi1* mutant presented the HC degeneration phenotype,

although HA (GFI1) remained detectable in HCs. This agrees with the notion that the *Gfi1*

mutant lacking the SNAG domain is equivalent to the *Gfi1*-null model [59]. Thus, *Gfi1* is

likely expressed as multiple unknown isoforms, many of which might be recognized by the

HA-tag antibody because HA is tagged to the last exon (exon 6). However, when we used the

four *Gfi1* sgRNAs together, we were able to both reproduce the HC degeneration phenotype

and completely block HA (GFI1) expression.

Why does ectopic RBM24 fail to alleviate the degeneration of *Pou4f3*$^{-/-}$ HCs?

Upon confirming the epistatic genetic interaction between *Pou4f3* and *Rbm24*, we

predicted that ectopic RBM24 expression should be able to, at least partly, alleviate the

degeneration of *Pou4f3*$^{-/-}$ HCs, similar to what we recently reported in our study on INSM1

and IKZF2 in cochlear OHCs [21]. However, RBM24 failed to rescue the HC degeneration in

Pou4f3$^{-/-}$ mice. Two potential explanations for this finding are the following: One, although

Rbm24 expression is modulated by POU4F3, RBM24 is not directly involved in the pathway

regulating HC survival; accordingly, cell death occurs later in *Rbm24*$^{-/-}$ HCs than in *Pou4f3*$^{-/-}$

HCs. Two, *Rbm24* is a key POU4F3 downstream target, but forced expression of RBM24

alone cannot compensate for the loss of POU4F3 or other known POU4F3 targets such as

orphan thyroid nuclear receptor *Nr2f2* and *Caprin-1* [60, 61]; CAPRIN-1 is recruited to stress

granules in cochlear HCs exposed to ototoxic trauma [61].

Potential application of the three *Rbm24* enhancers in cochlear HC gene therapy

Gene therapy is a promising strategy for restoring hearing capacity in humans with

inherited gene mutations causing hearing impairment [50], and a few such therapy examples

have been reported to date, including *Otoferlin* and *vGlut3* gene-replacement therapies

[62-65]. Currently, therapeutic cDNAs are primarily delivered into HCs by using an

adeno-associated virus (AAV) vector. Although several AAVs have been reported [66, 67],

the vectors transfect cochlear cells non-selectively and HC-specific AAVs are not yet

available. A solution to this problem is the following: instead of the CAG/CMV ubiquitous

promoter widely used in current AAVs [68, 69], using any one of the three *Rbm24* enhancers

together with the *Hsp68* mini-promoter (in brief *Rbm24-Hsp68*) should generate an AAV that

would allow HC-specific transfection. Moreover, *Rbm24* is permanently expressed in

cochlear HCs, and if *Rbm24-Hsp68* AAV acts as expected, it would represent a powerful tool

for future HC-specific gene therapy at all postnatal ages, with potential applications in

treating clinical human deafness.

METHODS AND MATERIALS

Mice

The *Atoh1$^{Cre/+}$* model was kindly provided by Dr. Lin Gan (Augusta University, USA).

The *Rosa26-loxp-stop-loxp-tdTomato* (Ai9)*/+* strain (Jax#: 007905) was from The Jackson

Laboratory. The *Ikzf2$^{V5/+}$* mouse strain is described in detail in our previous reports [21, 23].

All mice were bred and raised in an SPF-level animal room and all animal procedures were

performed according to the guidelines (NA-032-2022) of the IACUC of the Institute of

Neuroscience (ION), Center for Excellence in Brain Science and Intelligence Technology,

Chinese Academy of Sciences.

One-step generation of homozygous *Pou4f3, Gfi1,* or *Rbm24* mutants by using

CRISPR-stop

The detailed protocol for using CRISPR-stop to generate homozygous gene mutants is described in our previous reports [37, 38]. Briefly, efficient pre-tested sgRNAs (Table S1) and hA3ABE3 were co-injected into one-cell-stage mouse zygotes that were then transplanted into pseudopregnant female mice, which gave birth to F0 mice. The F0 mice carrying the expected homozygous mutation (pre-emergence of protein translation stop codon) were identified using Sanger sequencing of tail-DNA PCR samples and were immediately ready for analysis.

Generating *Pou4f3-* or *Gfi1*-null mutants harboring large DNA fragment deletions by using CRISPR/Cas9

To construct germ-line-stable null mutants of either *Pou4f3* or *Gfi1*, *Cas9* mRNA, two efficient pre-tested sgRNAs located at the proximal and distal ends of the targeted gene, and a single-stranded DNA donor (120 bp) (Table S3) were co-injected into one-cell-stage WT zygotes. Notably, the left and right halves (60 bp each) of the single-stranded DNA donor were homologous to the 5′ and 3′ ends of the targeted gene, respectively. The post-injected zygotes were transplanted into pseudopregnant females, which gave birth to F0 mice; the F0 mice were subject to tail-DNA PCR screening with the primers listed in Table S2, and the mice harboring the designed large DNA deletion between the two sgRNAs were identified and further bred with WT mice to establish the germ-line-stable mutants (F1 or afterward).

Construction of *Gfi1*$^{3 \, \Diamond HA\text{-}P2A\text{-}Cre}$ knockin mouse strain

An sgRNA against *Gfi1* (5′-ATGGACTCAAATGAGTACCC-3′), *Cas9* mRNA, and the targeting vector (Figure 2-figure supplement 1B) were co-injected into one-cell-stage WT

mouse zygotes. The targeting vector comprised three portions: the 5′ homologous arm (800

bp), the 3′ homologous arm (800 bp), and the region between the 5′ and 3′ arms that

contained three HA fragments followed by 2A-Cre. The F0 mice with the potential gene

targeting (Figure 2-figure supplement 1C) were screened using tail-DNA PCR and then

crossed with WT mice to produce F1 mice; these F1 mice were confirmed using tail-DNA

PCR again and further screened using Southern blotting. The detailed Southern blotting

protocol is described in our previous report [70]. Tail-DNA PCR was used for routine

genotyping, and the knockin ($Gfi1^{HA-Cre/+}$) and WT alleles were distinguished using the

primers F4, R5, and R6 (Table S2).

Construction of Eh1-EGFP+, Eh2-EGFP+, and Eh3-EGFP+ transgenic reporter lines

The three transgenic reporter mouse lines used here, Eh1-EGFP+, Eh2-EGFP+, and

Eh3-EGFP+, were produced using the same procedures. The core DNA sequences of each

enhancer (Eh1, Eh2, or Eh3 in Figure 4A), the mRNA encoding PiggyBac transposase, and

the PiggyBac vector (Figure 4B, D and F) were co-injected into one-cell-stage WT mouse

zygotes. The PiggyBac vector contained the Eh1/Eh2/Eh3 core DNA sequence (Table S3),

the mini-promoter of mouse *Hsp68*, and the EGFP coding sequence, and the vector was

randomly integrated into the mouse genome by the transposase. The F0 mice harboring the

PiggyBac vector were screened using tail-DNA PCR and further bred with WT mice to

produce germ-line-stable F1 transgenic reporter strains. The samples analyzed in Figure

4B-G'' were from F1 or later generations. The primers used for genotyping transgenic

reporter strains were F8, F9, F10, and the common primer R11 (primer sequences are listed in

Table S2).

Generation of *Rosa26*$^{CAG\text{-}lsl\text{-}Rbm24*3\times HA/+}$ mouse model

The *Rosa26*$^{CAG\text{-}lsl\text{-}Rbm24*3\times HA/+}$ (*Rosa26*$^{Rbm24/+}$) knockin mouse strain was constructed through homologous recombination mediated by CRISPR/Cas9. A pre-tested *Rosa26* sgRNA (5′-*ACTCCAGTCTTTCTAGAAGA*-3′), the targeting vector (Figure 5A), and *Cas9* mRNA were co-injected into one-cell-stage WT mouse zygotes, and similarly as in other cases, F0 mice with potentially correct gene targeting were screened using tail-DNA PCR and further bred with WT mice to establish germ-line-stable F1 mice. The WT and knockin (*Rosa26*$^{Rbm24/+}$) alleles were distinguished using the primers F12, R13, and R14 (Table S2).

Sample processing and immunofluorescence assay

Inner ears were dissected out and fixed in 4% paraformaldehyde (PFA) in PBS (E607016-0500, Sangon Biotech) at 4℃ overnight. For obtaining cochlear cryosections, inner ears were dehydrated in 30% sucrose (V900116, Sigma) at 4℃ before embedding into optimal cutting temperature (OCT) compound (4583, SAKURA), and then slices were cut at 14 μm thickness. The detailed protocol for immunofluorescence staining is described in our previous report [71]. The following primary antibodies were used: anti-RBM24 (rabbit, 1:500, 18178-1-AP, Proteintech), anti-POU4F3 (mouse, 1:500, sc-81980, Santa Cruz), anti-POU4F3 (rabbit, 1:500, NBP1-88349, Novus Biologicals), anti-HA (rat, 1:500, 11867423001, Roche), anti-MYO7A (rabbit, 1:500, 25-6790, Proteus Biosciences), anti-MYO7A (mouse, 1:500, MYO7A 138-1, Developmental Studies Hybridoma Bank), anti-CTIP2 (BCL11B) (rat, 1:500, ab18465, Abcam), anti-INSM1 (guinea pig, 1:6000, a kind gift from Dr. Carmen Birchmeier from Max Delbrueck Center for Molecular Medicine, Germany), anti-GFP (chicken, 1:500, ab13970, Abcam), anti-GATA3 (goat, 1:500, AF2605, R&D Systems), and anti-V5 (mouse,

1:500, MCA1360, Bio-Rad). Various corresponding Alexa Fluor-conjugated secondary

antibodies were used for detecting the primary antibodies, and Hoechst 33342 (1:1000,

H3570, Thermo Fisher Scientific) was used for nuclear DNA staining.

After staining, the whole-mount or cryosection samples were mounted with Prolong

Gold antifade medium (P36930, Thermo Fisher Scientific) at room temperature for 12 h.

Samples were scanned using a Nikon C2 or Nikon NiE-A1 Plus confocal microscope, and

ImageJ software was used to process the confocal images.

Cell quantification and statistical analysis

Before immunofluorescence staining, each cochlear sample was grossly separated into

three portions of distinct lengths, and each portion of the same cochlea was initially scanned

using a confocal microscope at low magnification (10× lens). After calculating the total

length of each cochlea, the cochlear sample was precisely divided into basal, middle, and

apical turns of equal length. Subsequently, for cell counting in experiments (Figure 1-figure

supplement 2D), an ~200 μm stretch of the sensory epithelium in each turn was scanned

using a confocal microscope at high magnification (60× lens) and the numbers of HCs were

determined. In $Pou4f3^{-/-}$ cochleae, the percentage of surviving HCs was calculated by

normalizing the number of remaining HCs against their counterparts in WT mice. All cell

numbers are presented as means ± SD. For statistical analyses, we used GraphPad Prism 6.0

software and performed Student's *t* tests with Bonferroni correction.

REFERENCES

1.   Wu, D.K. and M.W. Kelley, *Molecular mechanisms of inner ear development.* Cold Spring Harb
Perspect Biol, 2012. **4**(8): p. a008409.

2.   Qiu, X. and U. Muller, *Sensing sound: Cellular specializations and molecular force sensors.*
Neuron, 2022. **110**(22): p. 3667-3687.

3.   Groves, A.K., K.D. Zhang, and D.M. Fekete, *The genetics of hair cell development and*
regeneration. Annu Rev Neurosci, 2013. **36**: p. 361-81.

4.   Li, S., et al., *Fate-mapping analysis of cochlear cells expressing Atoh1 mRNA via a new*
Atoh1(3*HA-P2A-Cre) knockin mouse strain. Dev Dyn, 2022: p. DOI: 10.1002/dvdy.453.

5.   Driver, E.C., et al., *The Atoh1-lineage gives rise to hair cells and supporting cells within the*
mammalian cochlea. Dev Biol, 2013. **376**(1): p. 86-98.

6.   Matei, V., et al., *Smaller inner ear sensory epithelia in Neurog 1 null mice are related to earlier*
hair cell cycle exit. Dev Dyn, 2005. **234**(3): p. 633-50.

7.   Bermingham, N.A., et al., *Math1: an essential gene for the generation of inner ear hair cells.*
Science, 1999. **284**(5421): p. 1837-41.

8.   Luo, Z., et al., *Three distinct Atoh1 enhancers cooperate for sound receptor hair cell*
development. Proc Natl Acad Sci U S A, 2022. **119**(32): p. e2119850119.

9.   Zheng, J., et al., *Prestin is the motor protein of cochlear outer hair cells.* Nature, 2000.
405(6783): p. 149-55.

10.   Liberman, M.C., et al., *Prestin is required for electromotility of the outer hair cell and for the*
cochlear amplifier. Nature, 2002. **419**(6904): p. 300-4.

11.   Pan, Y., et al., *Fgf8(P2A-3xGFP/+): A New Genetic Mouse Model for Specifically Labeling and*
Sorting Cochlear Inner Hair Cells. Neurosci Bull, 2023. **39**(12): p. 1762-1774.

12.   Roux, I., et al., *Otoferlin, defective in a human deafness form, is essential for exocytosis at the*
auditory ribbon synapse. Cell, 2006. **127**(2): p. 277-89.

13.   Jacques, B.E., et al., *Fgf8 induces pillar cell fate and regulates cellular patterning in the*
mammalian cochlea. Development, 2007. **134**(16): p. 3021-9.

14.   Ruel, J., et al., *Impairment of SLC17A8 encoding vesicular glutamate transporter-3, VGLUT3,*
underlies nonsyndromic deafness DFNA25 and inner hair cell dysfunction in null mice. Am J Hum
Genet, 2008. **83**(2): p. 278-92.

15.   Seal, R.P., et al., *Sensorineural deafness and seizures in mice lacking vesicular glutamate*
transporter 3. Neuron, 2008. **57**(2): p. 263-75.

16.   Petitpre, C., et al., *Neuronal heterogeneity and stereotyped connectivity in the auditory afferent*
system. Nat Commun, 2018. **9**(1): p. 3691.

17.   Shrestha, B.R., et al., *Sensory Neuron Diversity in the Inner Ear Is Shaped by Activity.* Cell, 2018.
174(5): p. 1229-1246 e17.

18. Sun, S., et al., *Hair Cell Mechanotransduction Regulates Spontaneous Activity and Spiral*
Ganglion Subtype Specification in the Auditory System. Cell, 2018. **174**(5): p. 1247-1263 e15.

19. Zhang, D., et al., *Lypd1-DTR/+: A New Mouse Model for Specifically Damaging the Type Ic*
Spiral Ganglion Neurons of the Cochlea. Neurosci Bull, 2023.

20. Wiwatpanit, T., et al., *Trans-differentiation of outer hair cells into inner hair cells in the absence*
of INSM1. Nature, 2018. **563**(7733): p. 691-695.

21. Li, S., et al., *Epistatic genetic interactions between Insm1 and Ikzf2 during cochlear outer hair*
cell development. Cell Rep, 2023. **42**(5): p. 112504.

22. Chessum, L., et al., *Helios is a key transcriptional regulator of outer hair cell maturation.* Nature,
2018. **563**(7733): p. 696-700.

23. Bi, Z., et al., *Development and transdifferentiation into inner hair cells require Tbx2.* Natl Sci
Rev, 2022. **9**(12): p. nwac156.

24. Garcia-Anoveros, J., et al., *Tbx2 is a master regulator of inner versus outer hair cell*
differentiation. Nature, 2022. **605**(7909): p. 298-303.

25. Kaiser, M., et al., *TBX2 specifies and maintains inner hair and supporting cell fate in the Organ*
of Corti. Nat Commun, 2022. **13**(1): p. 7628.

26. Li, X., et al., *In situ regeneration of inner hair cells in the damaged cochlea by temporally*
regulated co-expression of Atoh1 and Tbx2. Development, 2023. **150**(24).

27. Vahava, O., et al., *Mutation in transcription factor POU4F3 associated with inherited*
progressive hearing loss in humans. Science, 1998. **279**(5358): p. 1950-4.

28. Masuda, M., et al., *Regulation of POU4F3 gene expression in hair cells by 5' DNA in mice.*
Neuroscience, 2011. **197**: p. 48-64.

29. Xiang, M., et al., *Essential role of POU-domain factor Brn-3c in auditory and vestibular hair*
cell development. Proc Natl Acad Sci U S A, 1997. **94**(17): p. 9445-50.

30. Zhu, G.J., et al., *Aldh inhibitor restores auditory function in a mouse model of human deafness.*
PLoS Genet, 2020. **16**(9): p. e1009040.

31. Erkman, L., et al., *Role of transcription factors Brn-3.1 and Brn-3.2 in auditory and visual*
system development. Nature, 1996. **381**(6583): p. 603-6.

32. Xiang, M., et al., *Requirement for Brn-3c in maturation and survival, but not in fate*
determination of inner ear hair cells. Development, 1998. **125**(20): p. 3935-46.

33. Yu, H.V., et al., *POU4F3 pioneer activity enables ATOH1 to drive diverse mechanoreceptor*
differentiation through a feed-forward epigenetic mechanism. Proc Natl Acad Sci U S A, 2021.
118(29).

34. Wallis, D., et al., *The zinc finger transcription factor Gfi1, implicated in lymphomagenesis, is required for inner ear hair cell differentiation and survival.* Development, 2003. **130**(1): p. 221-32.

35. Hertzano, R., et al., *Transcription profiling of inner ears from Pou4f3(ddl/ddl) identifies Gfi1 as a target of the Pou4f3 deafness gene.* Hum Mol Genet, 2004. **13**(18): p. 2143-53.

36. Grifone, R., et al., *Expression patterns of Rbm24 in lens, nasal epithelium, and inner ear during mouse embryonic development.* Dev Dyn, 2018. **247**(10): p. 1160-1169.

37. Wang, G., et al., *Mosaic CRISPR-stop enables rapid phenotyping of nonsense mutations in essential genes.* Development, 2021.

38. Zhang, H., et al., *Simultaneous zygotic inactivation of multiple genes in mouse through CRISPR/Cas9-mediated base editing.* Development, 2018. **145**(20).

39. Kuscu, C., et al., *CRISPR-STOP: gene silencing through base-editing-induced nonsense mutations.* Nat Methods, 2017. **14**(7): p. 710-712.

40. Matern, M., et al., *Gfi1(Cre) mice have early onset progressive hearing loss and induce recombination in numerous inner ear non-hair cells.* Sci Rep, 2017. **7**: p. 42079.

41. Yang, H., et al., *Gfi1-Cre knock-in mouse line: A tool for inner ear hair cell-specific gene deletion.* Genesis, 2010. **48**(6): p. 400-6.

42. Matern, M.S., et al., *GFI1 functions to repress neuronal gene expression in the developing inner ear hair cells.* Development, 2020. **147**(17).

43. Bardhan, T., et al., *Gata3 is required for the functional maturation of inner hair cells and their innervation in the mouse cochlea.* J Physiol, 2019. **597**(13): p. 3389-3406.

44. Sun, S., et al., *Dual expression of Atoh1 and Ikzf2 promotes transformation of adult cochlear supporting cells into outer hair cells.* Elife, 2021. **10**.

45. Kothary, R., et al., *Inducible expression of an hsp68-lacZ hybrid gene in transgenic mice.* Development, 1989. **105**(4): p. 707-14.

46. Xu, J., et al., *Chromatin remodelers and lineage-specific factors interact to target enhancers to establish proneurosensory fate within otic ectoderm.* Proc Natl Acad Sci U S A, 2021. **118**(12).

47. Shu, M., et al., *Single-cell chromatin accessibility identifies enhancer networks driving gene expression during spinal cord development in mouse.* Dev Cell, 2022. **57**(24): p. 2761-2775 e6.

48. Sun, Y., et al., *In vivo CRISPR-Cas9-mediated DNA chop identifies a cochlear outer hair cell-specific enhancer.* FASEB J, 2022. **36**(4): p. e22233.

49. Yang, H., et al., *Generation and characterization of Atoh1-Cre knock-in mouse line.* Genesis, 2010. **48**(6): p. 407-13.

50. Petit, C., C. Bonnet, and S. Safieddine, *Deafness: from genetic architecture to gene therapy.* Nat
Rev Genet, 2023.

51. Fritzsch, B., et al., *Atoh1 null mice show directed afferent fiber growth to undifferentiated ear*
sensory epithelia followed by incomplete fiber retention. Dev Dyn, 2005. **233**(2): p. 570-83.

52. Atar, O. and K.B. Avraham, *Anti-apoptotic factor z-Val-Ala-Asp-fluoromethylketone promotes the*
survival of cochlear hair cells in a mouse model for human deafness. Neuroscience, 2010. **168**(3): p.
851-7.

53. Liu, H., et al., *Molecular and cytological profiling of biological aging of mouse cochlear inner*
and outer hair cells. Cell Rep, 2022. **39**(2): p. 110665.

54. Ranum, P.T., et al., *Insights into the Biology of Hearing and Deafness Revealed by Single-Cell*
RNA Sequencing. Cell Rep, 2019. **26**(11): p. 3160-3171 e3.

55. Wang, Y., et al., *RBM24 is required for mouse hair cell development through regulating*
pre-mRNA alternative splicing and mRNA stability. J Cell Physiol, 2023.

56. Zheng, L., et al., *Rbm24 regulates inner-ear-specific alternative splicing and is essential for*
maintaining auditory and motor coordination. RNA Biol, 2021. **18**(4): p. 468-480.

57. Liu, L., et al., *Template-independent genome editing in the Pcdh15(av-3j) mouse, a model of*
human DFNB23 nonsyndromic deafness. Cell Rep, 2022. **40**(2): p. 111061.

58. Cai, T., et al., *Characterization of the transcriptome of nascent hair cells and identification of*
direct targets of the Atoh1 transcription factor. J Neurosci, 2015. **35**(14): p. 5870-83.

59. Fiolka, K., et al., *Gfi1 and Gfi1b act equivalently in haematopoiesis, but have distinct,*
non-overlapping functions in inner ear development. EMBO Rep, 2006. **7**(3): p. 326-33.

60. Tornari, C., et al., *Regulation of the orphan nuclear receptor Nr2f2 by the DFNA15 deafness*
gene Pou4f3. PLoS One, 2014. **9**(11): p. e112247.

61. Towers, E.R., et al., *Caprin-1 is a target of the deafness gene Pou4f3 and is recruited to stress*
granules in cochlear hair cells in response to ototoxic damage. J Cell Sci, 2011. **124**(Pt 7): p.
1145-55.

62. Akil, O., et al., *Dual AAV-mediated gene therapy restores hearing in a DFNB9 mouse model.*
Proc Natl Acad Sci U S A, 2019. **116**(10): p. 4496-4501.

63. Tang, H., et al., *Hearing of Otof-deficient mice restored by trans-splicing of N- and C-terminal*
otoferlin. Hum Genet, 2023. **142**(2): p. 289-304.

64. Al-Moyed, H., et al., *A dual-AAV approach restores fast exocytosis and partially rescues auditory*
function in deaf otoferlin knock-out mice. EMBO Mol Med, 2019. **11**(1).

65. Akil, O., et al., *Restoration of hearing in the VGLUT3 knockout mouse using virally mediated*

gene therapy. Neuron, 2012. **75**(2): p. 283-93.

66. Tan, F., et al., *AAV-ie enables safe and efficient gene transfer to inner ear cells.* Nat Commun, 2019. **10**(1): p. 3733.

67. Wu, J., et al., *Single and Dual Vector Gene Therapy with AAV9-PHP.B Rescues Hearing in Tmc1 Mutant Mice.* Mol Ther, 2021. **29**(3): p. 973-988.

68. Isgrig, K., et al., *AAV2.7m8 is a powerful viral vector for inner ear gene therapy.* Nat Commun, 2019. **10**(1): p. 427.

69. Landegger, L.D., et al., *A synthetic AAV vector enables safe and efficient gene transfer to the mammalian inner ear.* Nat Biotechnol, 2017. **35**(3): p. 280-284.

70. Li, C., et al., *Characterizing a novel vGlut3-P2A-iCreER knockin mouse strain in cochlea.* Hear Res, 2018. **364**: p. 12-24.

71. Liu, Z., et al., *Dynamic expression pattern of Sonic hedgehog in developing cochlear spiral ganglion neurons.* Dev Dyn, 2010. **239**(6): p. 1674-83.

Figure Legends

Figure 1. RBM24 expression is repressed in cochlear HCs in the absence of POU4F3. (A) Simple illustration of how *Pou4f3*$^{-/-}$ strain was generated. **(B)** One example gel image of tail-DNA PCR used to distinguish WT (*Pou4f3*$^{+/+}$), heterozygous (*Pou4f3*$^{+/-}$), and homozygous (*Pou4f3*$^{-/-}$) mice. **(C-F''')** Triple staining of RBM24, POU4F3, and BCL11B in WT (C and E-E''') and *Pou4f3*$^{-/-}$ (D and F-F''') mice (n=3 for each) at E16.5; only the RBM24 (green) channel (together with nuclear staining) is shown in (C and D) at low magnification. (E-E''') and (F-F'''): High-magnification images of middle turn in WT and *Pou4f3*$^{-/-}$ mice, respectively. In the absence of POU4F3, RBM24 expression completely disappeared, although nascent BCL11B+ OHCs were present. OHC: outer hair cell; IHC: inner hair cell. Scale bars: 100 μm (D), 20 μm (F''').

Figure 2. GFI1 is dispensable for RBM24 expression. (A) Simple cartoon depicting how

Gfi1 was inactivated in one-cell-stage zygotes derived from male *Gfi1$^{HA-Cre/+}$* mice by using

the CRISPR-stop approach. **(B-C)** Using sgRNA-6 as an example, Sanger sequencing

chromatograms of control *Gfi1$^{HA-Cre/+}$* (B) and F0 *Gfi1$^{-/-}$* (C) samples are presented. The base

"C" (black arrow in B) was converted to "T" (red arrow in C), resulting in pre-emergence of

the translation stop codon TAG. The red "T" appears as a single peak, indicating that the stop

codon pre-emerged in both alleles. **(D-E")** Dual staining of HA (GFI1) and RBM24 in

cochleae from control *Gfi1$^{HA-Cre/+}$* (D-D", n=3) and F0 *Gfi1$^{-/-}$* (E-E") mice (n=4) at E16.5.

RBM24 expression (E') was not altered in the absence of HA (GFI1) expression (E). OHC:

outer hair cell; IHC: inner hair cell. Scale bar: 20 μm (E").

Figure 3. POU4F3 and GFI1 expressions are normal in the absence of *Rbm24*. (A)

Simple cartoon showing how CRISPR-stop was used to induce mosaic inactivation of

RBM24 in one-cell-stage mouse zygotes harboring *Gfi1$^{HA-Cre/+}$*. Notably, the injection was in

the late one-cell stage of the zygotes, which resulted in high rates of final mosaic inactivation.

(B-C) Sanger sequencing chromatograms of control *Gfi1$^{HA-Cre/+}$; Rbm24$^{+/+}$* (B) and

Gfi1$^{HA-Cre/+}$; Rbm24$^{-/-}$ (mosaic) (C) samples. The base "C" (black arrow in B) was converted

to "T" (red arrow in C) in a fraction of cochlear cells. Mosaic inactivation of *Rbm24* is

evidenced by the double peaks of "T" and "C" (red arrow in C). **(D-E''')** Triple staining of

RBM24, HA, and Pou4f3 in cochlear samples (basal turns) of control *Gfi1$^{HA-Cre/+}$; Rbm24$^{+/+}$*

(D-D''', n=3) and *Gfi1$^{HA-Cre/+}$; Rbm24$^{-/-}$* (mosaic) (E-E''', n=3) mice at E17. Consistent with

the Sanger sequencing results (C), RBM24 expression was found to be inactivated in cochlear

HCs in a mosaic pattern (E). Orange arrows in (E-E'''): one OHC that lost RBM24

expression; blue arrows in (E-E'''): one OHC that maintained RBM24 expression. Regardless

of whether RBM24 expression was inactivated or not, the expression of HA (GFI1) and

POU4F3 was unaltered. OHC: outer hair cell; IHC: inner hair cell. Scale bar: 20 μm (E''').

Figure 4. Three *Rbm24* enhancers are sufficient to drive specific EGFP expression in

cochlear HCs. (A) Visualization of ATAC-seq (top row, orange) of P1 cochlear HCs (from

our previous study, GSE181311) and POU4F3 Cut&Run assay (bottom row, blue) of E17.5

cochlear HCs (from another study, GSE150391) by using Integrative Genomics Viewer.

Black arrow: *Rbm24* promoter; dotted boxes: three *Rbm24* enhancers (Eh1, Eh2, and Eh3).

(B-G'') Dual labeling for EGFP and MYO7A in Eh1-EGFP+ (B-C'', n=7), Eh2-EGFP+

(D-E'', n=6), and Eh3-EGFP+ (F-G'', n=4) mice at P1. Only the EGFP channel (plus nuclear

staining) is shown for whole-mount samples (B, D, F). In cryosection cochlear samples,

EGFP labeling overlapped with HC marker MYO7A. OHCs: outer hair cells; IHC: inner hair

cell; HC: hair cell. Scale bars: 100 μm (F), 20 μm (G'').

Figure 5. Forced expression of RBM24 fails to rescue OHC degeneration in

***Pou4f3$^{-/-}$*mutants. (A)** Generation of *Rosa26$^{loxp-stop-loxp-Rbm24*3×HA/+}$* (*Rosa26$^{Rbm24/+}$*) mouse

strain. The C-terminus of ectopic Rbm24 was fused with three HA tags. **(B)** One example gel

image of tail-DNA PCR used to distinguish WT (+/+) and knockin (*Rosa26$^{Rbm24/+}$*) mice.

(C-F'') Dual staining of HA (RBM24) and HC marker MYO7A in mice featuring four

distinct genotypes at P1: *Pou4f3$^{+/+}$* (C-C'', n=3), *Pou4f3$^{-/-}$* (D-D'', n=3), *Pou4f3$^{+/+}$;*

Atoh1$^{Cre/+}$; Rosa26$^{Rbm24/+}$ (E-E'', n=3), and *Pou4f3$^{-/-}$; Atoh1$^{Cre/+}$; Rosa26$^{Rbm24/+}$* (F-F'', n=3).

Ectopic HA (RBM24) was not detected in *Pou4f3$^{+/+}$* (C) and *Pou4f3$^{-/-}$* (D) mice, but was

detected in both *Pou4f3$^{+/+}$; Atoh1$^{Cre/+}$; Rosa26$^{Rbm24/+}$* (E) and *Pou4f3$^{-/-}$; Atoh1$^{Cre/+}$;*

Rosa26$^{Rbm24/+}$ (F) mice. White arrows in (E-E''): one MYO7A+ IHC that expressed ectopic

HA (RBM24). HCs appeared normal by P1, despite the induction of ectopic RBM24

expression. Orange arrows in (E-E''): one SC that expressed HA (RBM24) but not MYO7A.

White arrows in (F-F''): one remaining MYO7A+ HC that expressed ectopic RBM24.

Overall, many of the *Pou4f3$^{-/-}$* HCs degenerated even though ectopic RBM24 expression was

induced. OHC: outer hair cell; IHC: inner hair cell; HC: hair cell. Scale bar: 20 μm (F'').

Figure supplement Legends

Figure 1-figure supplement 1. RBM24 expression is repressed in cochlear HCs at E16.5

when POU4F3 is inactivated. (A) Illustration of how *Pou4f3$^{-/-}$* mice (Founder 0, F0) were

rapidly generated using CRISPR-stop. **(B-C)** Sanger sequencing chromatograms of WT (B)

and F0 *Pou4f3$^{-/-}$* (C) samples. The base "C" (black arrow in B) was completely converted to

"T" (red arrow in C), resulting in pre-emergence of a protein translation stop codon. The

Pou4f3$^{-/-}$ mutants (F0 mice) were immediately ready for phenotypic analysis. **(D-E''')** Triple

staining of pan-HC marker POU4F3, RBM24, and OHC marker INSM1 in whole-mount

cochlear samples (basal turns) of WT (D-D''', n=3) and *Pou4f3$^{-/-}$* (E-E''', n=9) mice at E16.5.

Relative to WT HCs, *Pou4f3$^{-/-}$* HCs showed no expression of POU4F3 or RBM24, but

expressed INSM1 normally. Orange arrows in (E-E'''): one INSM1+ OHC that lost POU4F3

and RBM24 expression. OHC: outer hair cell; IHC: inner hair cell; HC: hair cell. Scale bar:

20 μm (E''').

Figure 1-figure supplement 2. Severe degeneration of *Pou4f3$^{-/-}$* HCs occurs by perinatal

ages. (A-C") Dual staining of POU4F3 and MYO7A in middle-turn regions of cochleae from

WT mice (A-A") at P1 and *Pou4f3$^{-/-}$* mice at P0 (B-B") and P1 (C-C"). The *Pou4f3$^{-/-}$* mice

were from a germ-line-stable strain produced through CRISPR/Cas9-mediated large DNA

fragment deletion (Figure 1A). Arrows in (B-C"): surviving HCs in the absence of POU4F3

expression. **(D)** Percentages of remaining HCs in basal, middle, and apical turns in *Pou4f3$^{-/-}$*

mice at P0 (n=4) or P1 (n=3), relative to WT mice at P1 (n=3). Data are presented as Means ±

SD. Student's *t* test; ** $p<0.01$; **** $p<0.0001$; n.s., no significant difference. HC

degeneration in middle and basal turns was more severe at P1 than at P0, but this difference

was not significant in the apical turn (D). **(E-F")** Dual staining of RBM24 and MYO7A in

WT (E-E", n=3) and *Pou4f3$^{-/-}$* (F-F", n=3) mice at P1. White arrow in (F-F"): one remaining

Pou4f3$^{-/-}$ HC that lost RBM24 expression. OHC: outer hair cell; IHC: inner hair cell; HC:

hair cell. Scale bars: 20 μm (C" and F").

Figure 2-figure supplement 1. Construction of *Gfi1$^{3 \times HA-P2A-Cre/+}$* (*Gfi1$^{HA-Cre/+}$*) mouse

model in which HA expression faithfully represents GFI1 expression. (A-C) Simple

cartoon illustrating construction of *Gfi1$^{HA-Cre/+}$* strain. In the WT *Gfi1* allele (A), a DNA

fragment containing 3×HA-P2A-Cre (B) was inserted immediately before the TGA stop

codon (red in A) to generate the post-targeted *Gfi1$^{HA-Cre/+}$* allele (C). **(D)** One example gel

image of tail-DNA PCR used to distinguish $Gfi1^{+/+}$, $Gfi1^{HA-Cre/+}$, and $Gfi1^{HA-Cre/HA-Cre}$ mice.

(E-F") Dual staining of HA (GFI1) and HC marker MYO7A in control $Gfi1^{+/+}$ (E-E", n=3)

and homozygous $Gfi1^{HA-Cre/HA-Cre}$ mice (F-F", n=3) at P1. Arrows in (F-F"): one OHC

expressing HA (GFI1) and MYO7A. **(G-J")** Fate-mapping analysis of cells expressing $Gfi1$

by using $Gfi1^{HA-Cre/+}$; $Rosa26^{loxp-stop-loxp-tdTomato}$ (Ai9)/+ ($Gfi1^{HA-Cre/+}$; Ai9/+) strain; confocal

low-magnification and high-magnification images are presented in (G-H"') and (I-J"),

respectively. In Ai9/+ control mice at P2 (n=5), no tdTomato+ cells were detected (G-G"' and

I-I"). By contrast, in $Gfi1^{HA-Cre/+}$; Ai9/+ mice (n=5), tdTomato was detected in nonsensory

cells (white arrows in H) as well as in MYO7A+ HCs, which expressed heterogeneous levels

of HA (GFI1) and are indicated by different arrows in (J-J"): white arrows (#1), one OHC

with highest HA (GFI1) expression; orange arrows (#3), one OHC with lowest (or

undetectable) HA (GFI1) expression; and blue arrows (#2), one OHC with intermediate HA

(GFI1) expression. Scale bars: 100 μm in (H"'), 20 μm (F" and J").

Figure 2-figure supplement 2. GFI1 expression is lost in $Pou4f3^{-/-}$ HCs. (A) One-step

generation of $Pou4f3$ mutants by using CRISPR-stop with zygotes derived from $Gfi1^{HA-Cre/+}$

mice. **(B-C)** Sanger sequencing chromatograms of tail-DNA samples from control

$Gfi1^{HA-Cre/+}$ (B) and $Gfi1^{HA-Cre/+}$; $Pou4f3^{-/-}$ (mosaic) (C) mice. The base "C" (blue arrow in B)

was converted to "T" (red arrow in C), but the conversion was not complete—a double peak

existed and indicated mosaic inactivation of $Pou4f3$ expression. **(D-E"')** Triple staining of

POU4F3, RBM24, and HA (GFI1) in basal region of cochleae from $Gfi1^{HA-Cre/+}$ (D-D"', n=3)

and $Gfi1^{HA-Cre/+}$; $Pou4f3^{-/-}$ (mosaic, E-E"', n=3) mice at E16.5. Blue arrows in (E-E"'): one

OHC that expressed POU4F3, RBM24, and HA (GFI1); orange arrows: one OHC that lost

POU4F3, RBM24, and HA (GFI1) expression. This confirmed that not only does HA

faithfully represent GFI1 expression, but also that HA expression depends on POU4F3

expression. Scale bar: 20 μm (E''').

Figure 2-figure supplement 3. RBM24 expression is normal in the absence of *Gfi1*. (A)

The DNA fragment between sgRNA-4 and sgRNA-9 within the *Gfi1* locus was deleted using

the CRISPR/Cas9 method. **(B)** One example gel image of tail-DNA PCR used to distinguish

$Gfi1^{+/+}$, $Gfi1^{+/-}$, and $Gfi1^{-/-}$ mice. **(C-D'')** Dual staining of POU4F3 and RBM24 in the middle

turn of cochleae from WT (C-C'', n=3) and $Gfi1^{-/-}$ (D-D'', n=3) mice at P1. Severe

degeneration of HCs, particularly OHCs, occurred in $Gfi1^{-/-}$ cochleae. Orange arrows in

(D-D''): one surviving $Gfi1^{-/-}$ OHC that expressed POU4F3 and RBM24. **(E-F''')** Triple

staining of POU4F3, RBM24, and OHC marker BCL11B in cochleae from WT (E-E''', n=3)

and $Gfi1^{-/-}$ (F-F''', n=3) mice at E16.5. POU4F3, RBM24, and BCL11B expression patterns

did not differ markedly between WT and $Gfi1^{-/-}$ cochleae at E16.5, suggesting that $Gfi1^{-/-}$ HC

degeneration had not yet started. Scale bars: 20 μm (D'' and F''').

Figure 3-figure supplement 1. GATA3 is downregulated but IKZF2 is upregulated in

***Rbm24*-deficient HCs. (A)** Simple illustration of how mosaic *Rbm24* mutant mice were

generated. **(B-C'')** Dual staining of RBM24 and GATA3 in cochlear tissues from control

$Rbm24^{+/+}$ (B-B'') and $Rbm24^{-/-}$ (C-C'') mice at P1. **(D)** Cartoon depicting how mosaic *Rbm24*

mutant mice were produced using zygotes derived by crossing female wild-type (WT) mice

with male *Ikzf2$^{V5/+}$* mice. **(E-F")** Dual staining of RBM24 and V5 (IKZF2) in cochleae from

control *Ikzf2$^{V5/+}$; Rbm24$^{+/+}$* (E-E") and *Ikzf2$^{V5/+}$; Rbm24$^{-/-}$* (F-F") mice at P5. Scale bars: 20

μm (C" and F").

Figure 1

Figure 2

A

Gfi1^HA-Cre

sgRNA-6   sgRNA-8

ATG

E1 E2 E3 E4 E5 E6 3xHA-P2A-Cre TGA

sgRNA-5   sgRNA-7

Gfi1^HA-Cre/+ male × WT → one-cell stage zygotes → F0 pups Tail PCR Gfi1^-/-

hA3ABE3 mRNA
sgRNA: 5+6+7+8

B

sgRNA-6

G G G A C A G C C A G G G A G C T G C G G G G

PAM

Gfi1^HA-Cre/+

160   170

C

G G G A T A G C C A G G G A G C T G C G G G G

Gfi1^-/-

160   170

D HA (Gfi1) **D'** Rbm24 **D''** Merge + Hoechst

Gfi1^HA-Cre/+   E16.5

OHC
IHC

E **E'** **E''**   E16.5

Gfi1^-/-

OHC
IHC

Figure 3

Figure 4

Figure 5

Figure 1- figure supplement 1

Figure 1-figure supplement 2

Figure 2-figure supplement 1

A Gfi1 allele

B Targeting vector

C Gfi1 HA-Cre

D Tail PCR

Gfi1 +/+   Gfi1 HA-Cre/+   Gfi1 HA-Cre/HA-Cre

— 735 bp
— 429 bp

Gfi1 +/+   Gfi1 HA-Cre/HA-Cre

E / F   HA (Gfi1)   OHC / IHC

E' / F' Myo7a

E'' / F'' Merge + Hoechst   P1

Ai9/+   Gfi1 HA-Cre/+;Ai9/+

tdTomato   HA (Gfi1)   Myo7a   Merge+ Hoechst

G / G' / G'' / G'''   Ai9/+   P2

H / H' / H'' / H'''   Gfi1 HA-Cre/+;Ai9/+   P2

Middle turn

Ai9/+   Gfi1 HA-Cre/+;Ai9/+

I / J   tdTomato   #1 #2 #3   OHC / IHC

I' / J'   HA (Gfi1)

I'' / J'' Merge + Myo7a + Hoechst   P2

Figure 2-figure supplement 2

A

Gfi1 alelle

Gfi1 null

B
Tail PCR
Gfi1 +/+ Gfi1 +/- Gfi1 -/-
429bp
364bp

Pou4f3 | Rbm24 | Merge + Hoechst

WT

Gfi1 -/-

OHC
IHC
P1

Pou4f3 | Rbm24 | Bcl11b | Merge + Hoechst

WT

Gfi1 -/-

OHC
IHC
E16.5

Figure 3-figure supplement 1