Early Deficits in Dentate Circuit and Behavioral Pattern Separation after Concussive Brain Injury

Lucas Corrubia[1,2*], Andrew Huang[2*], Susan Nguyen[2*], Michael W. Shiflett[3], Mathew V. Jones[4], Laura A. Ewell[5] and Vijayalakshmi Santhakumar[1,2]

[1]Department of Pharmacology, Physiology and Neuroscience, Rutgers New Jersey Medical School, Newark, New Jersey 07103, [2]Department of Molecular, Cell and Systems Biology, University of California Riverside, Riverside, California 92521. [3]Department of Psychology, Rutgers University, Newark, NJ 07102.[4]Department of Neuroscience, University of Wisconsin, Madison, WI, 53705. [5]Department of Anatomy and Neurobiology, University of California Irvine, Irvine, California 92697.

Short Title Dentate Pattern Separation after FPI

*Authors Contributed Equally

Word count for Abstract: 235

Full Text Character count: 5690 words   Number of pages: 36   Number of figures: 5

Number of Supplementary Figures: 1   Color figures:5   Number of Tables: 0

Number of Supplementary Tables: 0

Author Contributions: L.C. A.H L.A.E and S.N. Investigation; Formal analysis; Data curation; Methodology; Figures; A.H. Software; L.C. A.H., S.N. M.W.S., M.V.J., L.A.E and V.S. interpreted results of experiments; A. H., L.C. and S.N. prepared figures; A.H., L.C., S.N., and V.S. Conceptualization V.S. Funding acquisition; Project administration; Resources; Supervision; L.C. A.H., S.N. M.W.S., M.V.J., L.A.E and V.S. Writing - review & editing.

Declaration of Interest: None

Correspondence:

Vijayalakshmi Santhakumar, PhD
Department of Molecular, Cell and Systems Biology
University of California, Riverside
3401 Watkins Drive (Rm 1308 Spieth Hall) Riverside, CA 92521
E-mail: vijayas@ucr.edu

Abstract:

Traumatic brain injury leads to cellular and circuit changes in the dentate gyrus, a gateway to

hippocampal information processing. Cell intrinsic sparse granule cell firing and strong feedback

inhibition in the dentate are proposed as critical to its ability to generate unique representation of

similar inputs by a process known as pattern separation. Here we evaluate the impact of brain

injury on cellular decorrelation of temporally patterned inputs in slices and behavioral

discrimination of spatial locations *in vivo* one week after concussive lateral fluid percussion

injury (FPI) in mice. Despite posttraumatic increases in perforant path evoked excitatory drive to

granule cells and enhanced ΔFosB labeling, indicating sustained increase in excitability, the

reliability of granule cell spiking was not compromised after FPI. Although granule cells

continued to effectively decorrelate output spike trains recorded in response to similar temporally

patterned input sets after FPI, their ability to decorrelate highly similar input patterns was

reduced. In parallel, encoding of similar spatial locations in a novel object location task that

involves the dentate inhibitory circuits was impaired one week after FPI. Injury induced changes

in pattern separation were accompanied by loss of somatostatin expressing inhibitory neurons in

the hilus. Together, these data suggest that the early posttraumatic changes in the dentate circuit

undermine dentate circuit decorrelation of temporal input patterns as well as behavioral

discrimination of similar spatial locations, both of which could contribute to deficits in episodic

memory.

Keywords: granule cell, dentate gyrus, electrophysiology, traumatic brain injury, memory,

behavior, fluid percussion injury, temporal pattern separation

Introduction

Traumatic brain injury (TBI) is a growing silent epidemic (Vaishnavi et al., 2009; Rusnak, 2013)

with an annual global incidence of over 60 million patients (McAllister, 1992) who are at risk for

a variety of neuropsychological sequelae (Rao and Lyketsos, 2000). While neurological

complications of TBI increase with injury severity (Annegers and Coan, 2000), even mild to

moderate concussive brain injuries can lead to early and long term deficits in memory function

(Kumar et al., 2013; Arciniega et al., 2021; Alosco et al., 2023). The dentate gyrus, a critical

node in memory processing, receives extensive inputs from the entorhinal cortex and develops

discrete sparse representations for downstream processing in hippocampal circuits (Leutgeb et

al., 2007; Neunuebel and Knierim, 2014). The dentate gyrus is also the focus of neuronal loss

and circuit reorganization early after human and experimental concussive brain injury

(Lowenstein et al., 1992; Santhakumar et al., 2000; Meier et al., 2016; Leh et al., 2017;

Neuberger et al., 2017a; Folweiler et al., 2020), suggesting that dentate dependent memory

functions are likely compromised after brain injury.

In particular, the dentate gyrus has been shown to be essential for distinguishing between similar

memory episodes by encoding partially overlapping memories into discrete neuronal

representations (Gilbert et al., 2001; McHugh et al., 2007; Yassa and Stark, 2011; Cayco-Gajic

and Silver, 2019). At the circuit level, this is proposed to be achieved by pattern separation, a

process of minimizing interference by segregating similar input patterns into discrete neuronal

activity patterns. Whereas behaviorally relevant inputs and the resulting neuronal input patterns

occur in space and time (Hainmueller and Bartos, 2020), their neuronal representations have

been evaluated mostly in the spatial domain as changes in firing or activity rates in spatially

discrete neuronal groups or ensembles (Leutgeb et al., 2007; Larimer and Strowbridge, 2010;

GoodSmith et al., 2017). However, whether brain injury impairs the ability of the dentate circuit

to faithfully represent inputs and disambiguate similar neuronal input patterns remains unknown.

Most behavioral studies have focused on discrimination of spatial and contextual patterns to

evaluate mechanisms of dentate pattern separation and assess the impact of disease (McHugh et

al., 2007; Morris et al., 2012; Bekinschtein et al., 2013; Kahn et al., 2019). Studies in the fluid

percussion injury (FPI) model of concussive brain injury have revealed early deficits in a

sequential spatial navigation in a dentate-dependent radial arm maze task, indicating that

concussive brain injury compromises the ability to process and recall subtle spatial differences

(Correll et al., 2021). Similarly, brain injury is also known to compromise working memory

function (Smith et al., 2015; Korgaonkar et al., 2020a). Still, whether episodic processing of

spatial novelty by behavioral discrimination of spatial location is altered after brain injury

remains to be determined.

In an approach complementary to behavioral studies, computational analyses of mechanisms for

disambiguating overlapping memories have adopted spatially and temporally patterned inputs to

assess "pattern separation". In this approach, model networks are activated by inputs with

various degrees of similarity in spike timing and/or activated cell ensembles and the ability of the

network to generate distinct spatio-temporal firing patterns is used as a measure of pattern

separation (Rolls and Kesner, 2006; Myers and Scharfman, 2011; Madar et al., 2019b; Braganza

et al., 2020). These studies have shown that dentate granule cells (GCs) decorrelate input

patterns both in terms of the active ensembles recruited as well as the pattern of firing in the

active neurons (Myers and Scharfman, 2009, 2011; Yim et al., 2015). However, few

experimental studies have evaluated the ability of the dentate circuit to decorrelate spatially and

temporally patterned inputs (Larimer and Strowbridge, 2010; Madar et al., 2019a). Recently,

Madar et al. (2019a) adopted an experimental paradigm in hippocampal slices to demonstrate

that GCs could undertake input decorrelation at the level of individual neurons. These studies

identified a role for inhibition in supporting temporal pattern separation in GCs and revealed

cell-type specific and disease related changes in input decorrelation by dentate neurons (Madar et

al., 2019a; Madar et al., 2021). These findings are consistent with predictions of simulations

(Braganza et al., 2020) and with behavioral deficits in discrimination of novel spatial location in

mice during suppression of dentate somatostatin interneuron activity in vivo (Morales et al.,

2021). Since experimental concussive brain injury has been shown to result in early loss of

dentate inhibitory neuron subtypes (Lowenstein et al., 1992; Toth et al., 1997; Santhakumar et

al., 2001; Folweiler et al., 2020), we examined whether the resulting changes in inhibition

(Folweiler et al., 2020; Gupta et al., 2020; Gupta et al., 2022) could alter dentate dependent

pattern separation in the temporal and spatial domains one week after brain injury. Because the

relation between neuronal spike train timing and behavioral spatial information is currently

unclear, we will denote GC pattern separation of temporally patterned inputs in slice as

"decorrelation" and behavioral spatial pattern separation in vivo as "spatial discrimination".

MATERIALS AND METHODS

All experiments were performed in accordance with IACUC protocols approved by the University of California, Riverside, CA in keeping with the ARRIVE guidelines.

Fluid Percussion Injury: Randomly selected 8 to 10-week-old littermate male and female C57BL6/J mice were subjected to lateral fluid percussion injury (FPI) or sham injury. Briefly, mice were placed in a stereotaxic frame under isoflurane anesthesia and administered the local anesthetic 0.25% bupivacaine (subcutaneous). A 2.7 mm hole was trephined on the left side of the skull 2 mm caudal to bregma and 1 mm lateral from the sagittal suture. A Luer-Lok syringe hub with a 3 mm inner diameter was placed over the exposed dura and bonded to the skull with cyanoacrylate adhesive. The animals were returned to their home cage following recovery. One day later, animals were anesthetized with isoflurane and attached to a fluid percussion injury device (Virginia Commonwealth University, Richmond, VA, Model 01-B) directly at the metal nozzle. A pendulum was dropped to deliver a brief (20 ms) 1.5–1.7 atm impact on the intact dura. For sham injury, the animals underwent surgical implantation of the hub and were anesthetized and attached to the FPI device, but the pendulum was not dropped (Smith et al., 2012; Gupta et al., 2022).

Slice Preparation: Naive C57BL6/J mice and mice one week after FPI or sham injury were anesthetized with isoflurane and decapitated for slice preparation using established protocols (Korgaonkar et al., 2020b; Afrasiabi et al., 2021). Horizontal brain slices (300 μm) were prepared in ice-cold sucrose artificial CSF (sucrose-aCSF) containing the following (in mM): 85 NaCl, 75 sucrose, 24 $NaHCO_3$, 25 glucose, 4 $MgCl_2$, 2.5 KCl, 1.25 NaH_2PO_4, and 0.5 $CaCl_2$ using a Leica VT1200S Vibratome (Wetzlar, Germany). Slices were incubated at 34°C for 30 min in an interface holding chamber containing an equal volume of sucrose-aCSF and recording

aCSF, and subsequently were held at room temperature. The recording aCSF contained the

following (in mM): 126 NaCl, 2.5 KCl, 2 CaCl$_2$, 2 MgCl$_2$, 1.25 NaH$_2$PO$_4$, 26 NaHCO$_3$, and 10

D-glucose. All solutions were saturated with 95% O$_2$ and 5% CO$_2$ and maintained at a pH of 7.4

for 1– 6 h. Only sections on the side of injury were used in experiments.

Physiological Recordings

Voltage clamp recordings of perforant path evoked excitatory postsynaptic currents (eEPSCs)

were obtained from dentate GCs using recording electrodes (3 - 6 MΩ) containing a Cs-

Methanesulfonate Internal (in mM): 140 Cs-Methane sulfonate, 5 NaCl, 10 HEPES, 0.2 EGTA,

4 Mg-ATP, 0.2 Na-GTP, 5 Qx-314 with 0.2% biocytin. GCs were held at -70 mV and EPSCs

were recorded in response to 0.1 to 2.0 mA stimuli, applied in 100 pA steps, through a concentric

stimulus electrode positioned at the perforant path. Peak amplitude of eEPSC was measured from

the average of 3 to 5 responses at each step using ClampFit 10.

Current clamp recordings to assess GC pattern separation were obtained using filamented

borosilicate glass electrodes (3 - 6 MΩ) containing (in mM): 126 K-gluconate, 4 KCl, 10

HEPES, 4 Mg-ATP, 0.3 Na-GTP, 10 PO-creatinine with 0.2% biocytin. To assess cell health,

GCs were held at -70 mV and responses to 1500 ms current steps from -200 pA to 200 pA in 40

pA steps were recorded to examine intrinsic properties. Cells with resting membrane potential

positive to -55 mV were excluded from analysis.

Pattern Separation Paradigm and Analysis

To assess GC pattern separation, input trains of varying correlations were adopted from Madar et

al. (2019a). Each input correlation set included 5 temporally distinct trains at 10 Hz with an

overall mean Pearson's correlation coefficient (R_{in}) of 0.25, 0.50, 0.75, or 0.95, designated

henceforth as R25, R50, R75 and R95. The individual input trains (#1-#5) within an input

correlation set were each 2 seconds long and were delivered at 2 second intervals. The 5 distinct

input trains were delivered in sequence and the set was repeated 10 times for a total of 50

recorded trains (Madar et al., 2019a; Madar et al., 2021).

GCs were held at -70 mV in whole cell configuration and stimuli were delivered through a

concentric stimulating electrode (10 µm tip diameter) placed in the outer molecular layer to

activate perforant path fibers. Stimulus trains were delivered with intensity ranging from 0.1-2

mA so as to elicit firing in response to approximately 50% of stimuli. To assess the role of

inhibition in temporal pattern separation, GCs were recorded and stimulated with input

correlations of R75 in aCSF and then perfused with a sub-saturating concentration of gabazine

(SR95531, 100nM), a $GABA_A$ receptor antagonist.

For quantification of GC pattern separation function, Pearson's R correlation was calculated on

output spike train rasters as previously defined in Madar et al. (2019a) using custom MATLAB

code modified from: https://github.com/antoinemadar/PatSepSpikeTrains. Unless indicated

otherwise, data were binned at 10 ms. For validation of pattern separation paradigm in naïve

mice, pairwise correlation of each GC spike output train (R_{out}) was plotted against the

predetermined correlation in the corresponding pairwise input spike trains. Because the output

spike trains generated by each input pattern are compared in a pairwise manner, the five patterns

per input correlation yielded a total of 10 R_{out} values. Individual correlation values were used to

compare data from sham and FPI mice.

To assess the reliability of GC input-output coupling between experimental groups, we evaluated

spike train reliability (R_w) as a measure of the cell's ability to faithfully respond to multiple

presentations of the same input train (effectively $R_{in}=1$). R_w was calculated as the average

correlation of output of a given GC in response to 10 presentations of the same input pattern in

the R95 input set and then averaged across the 5 patterns.

Novel Object Location Task and Analysis

To assess behavioral pattern separation, we implemented a novel object location task to test

spatial discrimination (Bekinschtein et al., 2013; Morales et al., 2021) in mice one week post FPI

or sham injury. The task consisted of a circular arena 50 cm in diameter and 47 cm high with

laboratory tape in varying patterns in four distinct locations on the white opaque arena walls

serving as spatial visual cues. The experimenter room was separated from the testing room with

the test arena by a closed door. The testing room was dimly lit with overhead lights using

dimmer switch on lowest setting. Mice underwent four days of habituation followed by a sample

and choice phase on day five. The testing arena and objects were cleaned with 70% ethanol in

between each animal throughout the entire experiment. During habituation, mice were brought to

the testing room from the vivarium, handled by the experimenter, and exposed to the testing

arena for 10 min each day where they were allowed to explore freely. On test day, mice were

first exposed to a sample phase for 10 min, with three identical objects placed 120 degrees and

approximately 16 cm apart and allowed to freely explore the objects. Mice were then returned to

their home cage for 30 min before the choice phase. During the choice phase, one object was

placed in its original (familiar) location, one object was removed, and the last object was shifted

by 60° to a novel (unfamiliar) location. The mice were placed into the arena and allowed to

explore the objects for five minutes. Videos were manually coded and analyzed by an

experimenter blinded to animal injury status. Object exploration was counted as the number of

times the animal's nose was within 0.5 cm of the object, with each continuous second counted as

an additional exploration. Discrimination ratio was calculated as [Exploration of Object in

Unfamiliar Location – Exploration of Object in Familiar Location] / [Total Exploration]. Total

exploration was also compared between groups.

Immunohistochemistry:

Immunohistochemistry for ΔFosB and somatostatin (SST) was used to evaluate persistent

neuronal activity (You et al., 2017) in interneuron loss. Immediately following the novel object

location task, mice were perfused transcardially with ice cold PBS followed by 4% PFA. Brains

were excised and incubated in 4% PFA overnight and transferred to 30% sucrose for 2-3 days or

until tissue sank. Brains were then flash frozen in OCT using liquid nitrogen and stored at -80°C

prior to cryosectioning. 20 µm serial sections were collected and mounted on SuperFrost slides

and stored at -80°C prior to staining. Slides were stained for ΔFosB or SST following standard

procedures (Neuberger et al., 2017b; Korgaonkar et al., 2020b). Briefly, slides were allowed to

thaw to room temperature (RT) for 10 min, washed with 1xPBS 3x5min to wash off residual

OCT. Slides were blocked in 10% normal goat serum in PBS + 0.3% Triton-X for one hour at

RT, then incubated in ΔFosB or SST antibody (1:500, Cell Signaling D3S8R; SST: 1:100,

Millipore MA5-16987) overnight at 4°C. Slides were then washed 3x10 min with PBS and then

incubated in goat anti-rabbit AF594 secondary antibody (1:1000, Invitrogen A11012) for ΔFosB

or goat anti-rat AF647 antibody (1:1000, Invitrogen A-21247) for SST for 1.5 hours at RT,

washed 3x10 min with PBS, and coverslipped using VectaShield with DAPI mounting media.

Slides were imaged using Zeiss Axioscope at 10x and 20x. Analysis was performed using

automated algorithms in QuPath 0.4.2 (https://qupath.github.io) by drawing a ROI around the

granule cell layer (GCL) using DAPI labeling and detection for ΔFosB positive cells and in the

hilus to detect SST neurons. A range of 3 to 6 level matched sections were averaged in each

mouse for analysis. Confocal images were obtained on Zeiss 880 Upright Airyscan for

illustration.

Statistical Analysis: Statistical analyses were conducted in GraphPad Prism 9. The Shapiro-

Wilk test was used to test for normality, and appropriate parametric or non-parametric statistical

analyses were conducted. F-test to compare variances was used to assess whether both sample

groups undergoing statistical comparisons have equal standard deviations. All statistical

comparisons were conducted at an alpha level of $\alpha = 0.05$. Two-way repeated measures ANOVA

(TW-RM ANOVA), paired and unpaired Student's or Welch's t-test and Mann-Whitney U test

were used where appropriate.

RESULTS

Dentate gyrus excitability is increased one week after concussive brain injury in mice.

Previous studies have identified an early increase in dentate network excitability after concussive brain injury in rat (Santhakumar et al., 2001; Neuberger et al., 2017b; Korgaonkar et al., 2020b) and mice (Smith et al., 2012). To determine if perforant path mediated drive of GCs is enhanced after FPI in mice, we performed whole-cell voltage clamp recordings one week after FPI or sham injury. As reported following FPI in rat (Korgaonkar et al., 2020b), perforant path-evoked excitatory post-synaptic current (eEPSC) amplitudes in response to increasing current injection, was significantly enhanced in mice one week after FPI compared to sham injured controls (Figure 1A-C, n = 6 cells from sham and 4 cells from FPI, $F(1, 156)=29.79$, $p<0.0001$ for effect of treatment by TW-RM ANOVA). To assess whether FPI results in sustained increase in excitability one week after injury, we examined dentate sections for expression of ΔFosB, a uniquely stable activity-dependent immediate early gene product shown to be persistently enhanced in epilepsy (You et al., 2017). Immunohistochemical staining of sections from a cohort of sham and FPI mice sacrificed one week after FPI identified an increase in the number of ΔFosB expressing putative GCs in the granule cell layer compared to that in sham mice (Fig 1 D-E, number of ΔFosB + GCs: sham: 24.11±5.98, n = 8 mice FPI: 52.51±15.25, n = 7 mice, p=0.02 by Mann-Whitney U test), which is consistent with sustained enhancement of dentate excitability. There was an increase in variance in the number of ΔFosB expressing GCs after brain injury ($F_{6,7}$ = 5.6, p=0.04) indicative of individual differences in dentate hyperexcitability. Together, these data establish that the mouse model of moderate FPI used in our study results in both an increase in excitatory drive to GCs and an overall sustained increase in dentate excitability one week after injury.

Inhibition is critical for granule cell decorrelation of temporal spike train patterns.

The characteristic low GC excitability and robust inhibition are critical for maintaining sparse GC firing and pattern separation. To determine if temporal input decorrelation by GCs is impacted by FPI, we implemented an ex-vivo temporal pattern separation paradigm (Madar et al., 2019a; Madar et al., 2021) to assess the ability of GCs, within the local dentate microcircuit, to disambiguate temporally patterned inputs. GC firing was recorded in response to stimulation of perforant path fibers in the outer molecular layer, using sets of stimulus patterns with known degrees of similarity, to quantify similarity between output spikes. For initial validation, four sets of Poisson input stimulus patterns with 10 Hz average frequency and input correlations of R25, R50, R75, or R95 were used to elicit GC responses (Fig 2A). Similarity indices of the GC "output" spike trains (R_{out}) were computed as the pairwise Pearson's correlation coefficient on output spike train data binned at 10, 20 and 100 ms to assess the effect of bin-width on R_{out} (Fig 2B). Consistent with Madar et al. (2019a), GC output similarity was consistently lower than the corresponding pairwise input similarity (Fig 2B, R_{in} vs. R_{out}, n =4 cells / 3 mice). Since the reduction in output similarity was most robust when spike data was binned over 10 ms (Fig. 2B), a bin width of 10 ms was adopted in subsequent experiments. We then examined the potential contribution of the local inhibitory circuit to GC temporal pattern separation by partially decreasing GABAergic inhibition. To compare similarity indices before and after partial inhibitory block within the same cell, we restricted the inputs to a set of 5 stimuli with R75, reflecting a 75% similarity of across the inputs. GC responses were recorded first in aCSF and then in a non-saturating concentration of gabazine (100 nM). Under both recording conditions, GC output correlation remained consistently lower than the input correlation of 0.75 suggesting contributions from cell intrinsic and/or afferent synaptic characteristics in mediating GC temporal pattern separation. However, pairwise

255 comparison of R_{out} within the same cell and pattern in aCSF versus gabazine revealed a

256 significant increase in GC output correlation (Fig.2C-E, Mean pairwise R_{out}, aCSF: 0.20±0.02,

257 Gabazine: 0.30±0.01, n = 4 cells /3 mice; p < 0.0001 by Paired t-test, Cohen's D =1.7). These

258 results validate the role for inhibition in the ability of GCs to decorrelate temporally patterned

259 inputs as reported previously (Madar et al., 2019a).

Reliability of dentate granule cell spiking is not altered early after FPI.

After validating the experimental brain injury model in mouse (Fig. 1) and establishing the

temporal pattern separation paradigm in slice recordings (Fig. 2), we examined GCs in mice one

week after sham or FPI for their ability to decorrelate input spike train sets with 25% and 95%

similarity. To ensure that there were no systematic differences in GC recordings between sham

and FPI, we confirmed that the access resistance was not different between recordings from the

two experimental groups (in MΩ, sham: 13.98±1.38, FPI: 15.57±1.97, p=0.54 by Student's t-

test). Additionally, examination of the GC intrinsic parameters failed to reveal differences in

resting membrane potential (in mV, sham: -72.18±1.14, , FPI: -75.86±2.06, p=0.16 by Student's

t-test), input resistance (in MΩ, sham: 102.3±8.61, FPI: 121.1±13.34, p=0.28 by Student's t-test),

firing frequency (in Hz, in response to +160 pA, sham: 4.6±1.98, FPI: 5.6±1.44, p=0.85 by

unpaired Student's t-test) and action potential threshold (in mV, sham: -31.49±2.13, FPI: -

32.15±1.78, p=0.81 by unpaired Student's t-test) between groups (based on 10-11 cells from 3

sham mice and 12-14 cells from 4 FPI mice). Given the neuronal loss and circuit alterations

after FPI (Bonislawski et al., 2007; Neuberger et al., 2017b; Folweiler et al., 2020; Gupta et al.,

2020), we examined whether the fidelity of GC response to a given input train is fundamentally

altered by injury. Under optimal conditions, if a local dentate network receives identical inputs, a

given GC within the receiving network should maintain indistinguishable output responses. To

determine if injury alters the ability of the GC network to reliably respond to a given input train,

we calculated spike train reliability (R_w) as the average R_{out} of a given GC in response to

identical input pattern using the R95 input set (Fig 3A, B). Note that because stimulus intensity

was set to achieve a 50% spike success in each cell, the enhanced eEPSC (Fig. 1) should not

impact evaluation of spike train reliability. Indeed, the average firing elicited by the input spike

trains was not different between GCs from sham and FPI (Fig. 3C, average firing frequency in

Hz, sham: 3.55 ± 0.38, FPI: 2.99 ± 0.19, p=0.17 by unpaired Student's t-test). Interestingly, the

variance of the firing frequency trended towards a reduction after FPI ($F_{10,13}$= 3.10, p = 0.06, F-

test for variance) suggesting a potential decrease in a source of input decorrelation after FPI.

Despite the dentate circuit changes and enhanced ΔFosB labeling in GCs after FPI, spike train

reliability was not different between the GCs from sham and FPI mice (Fig. 3C, average spike

train reliability: sham: 0.47 ± 0.05, FPI: 0.51 ± 0.03, p=0.44 by unpaired Student's t-test; variance

($F_{10,13}$= 1.92, p = 0.27 by F-test for variance).

Granule cell decorrelation of temporal patterned inputs is reduced after FPI.

To compare temporal pattern separation between GCs from sham and FPI mice, GCs were

stimulated using input stimulus train sets with 25% and 95% similarity (Fig. 4A-B). Plots of

averaged pairwise R_{out} values against R_{in}, for each input set showed that the GCs from both sham

and FPI mice were still able to decorrelate inputs ($R_{out} < R_{in}$) (Fig. 4C). However, they suggested

a trend towards higher R_{out} (decrease in decorrelation) in GCs from FPI mice in response to

inputs with higher similarity (n = 11 cells / 3 mice) and FPI (n = 14 cells / 4 mice). To better

resolve this, we directly compared the pairwise R_{out} between groups for both low and high

similarity input sets. Indeed, R_{out} values in response to inputs with low similarity (R25) were not

different between FPI and sham (Fig 4D and Supplementary Fig 1A, p=0.41 by unpaired

Student's t-test) indicating that GCs from FPI mice are still effective at decorrelating dissimilar

patterns. However, in response to highly similar input trains (R95), the R_{out} values in GCs from

FPI mice were significantly higher than in sham controls (Fig 4D and Supplementary Fig 1B,

p=0.04 by unpaired Welch's t-test, Cohen's D = 0.26) indicating a reduction in the ability of the

dentate circuit to decorrelate highly similar inputs after injury. The posttraumatic decrease in

input decorrelation was accompanied by a reduction in the variance ($F_{109,139}$= 1.90, p = 0.0004,

F-test for variance). These data show that although GCs are continuing to disambiguate similar

spike train inputs early after injury, their critical ability to do so when inputs are highly similar is

compromised after injury.

Behavioral spatial pattern separation is degraded early after FPI.

To determine whether the reduction in the ability of the dentate circuit to decorrelate input

patterns after injury was accompanied by deficits in behavioral spatial discrimination, we

adopted a novel object location task, which relies on dentate function and tests for episodic rather

than sequential memory (Morales et al., 2021). Mice one week after FPI or sham surgery were

assessed in the novel object location task following appropriate habituation (Fig. 5A). The mice

were exposed to three identical objects in the sample phase followed, 30 minutes later, by a

choice phase in which one object was removed and one of the remaining objects was moved to a

novel location (Fig. 5A-B). Compared to sham mice, which showed significantly greater

interaction with the object in the novel location (p=0.0002 by one sample t-test), FPI mice failed

to show preference for the object in the novel location (p=0.67 by one sample t-test).

Consistently, the discrimination ratio, defined as the difference between number of interactions

with the object in the novel location and the familiar location divided by the total interaction with

both objects, was lower in FPI mice (Fig. 5C, discrimination ratio: sham: 0.28±0.04, FPI:

0.04±0.08, n = 8 mice/group, p=0.02 by unpaired Student's t-test). However, the total

exploration assessed as the number of contacts with objects in the 5-minute exploration period

was not different between groups (Fig. 5D, number of contacts: sham: 39.25±5.58, FPI:

41.13±6.6, n = 8 mice each, p=0.83 by unpaired Student's t-test) indicating no difference in

exploration between sham and FPI mice.

Finally, because, previous studies have identified a role for dentate somatostatin interneurons in

performance of the novel object location task (Morales et al., 2021) and somatostatin neurons are

vulnerable to brain injury (Lowenstein et al., 1992; Santhakumar et al., 2000; Frankowski et al.,

2019), we examined whether there was a loss of somatostatin neurons in the FPI mice that

performed the behavioral task. Staining for somatostatin revealed a significant reduction labeled

neurons in the dentate hilus one week after FPI (Fig 5E-F, cells/section, sham: 12.09±0.63, FPI:

9.39±1.04, based on within animal averages of level matched sections from 8 sham and 6 FPI

mice, p=0.04 by unpaired Student's t-test). These data suggest that injury-induced loss of dentate

somatostatin interneurons could contribute to the observed impairment in location discrimination

in mice early after brain injury.

Discussion:

Changes in episodic memory are a major neurocognitive consequence of traumatic brain injury with early and persistent deficits observed in patients and in animal models (Hamm et al., 1996; McHugh et al., 2006; Tsirka et al., 2010; Kumar et al., 2013; Smith et al., 2015; Korgaonkar et al., 2020a; Alosco et al., 2023). Although the dentate gyrus is known to play an essential role in encoding and disambiguating spatial memories, the underlying circuit mechanisms are not fully understood (Rolls and Kesner, 2006; Morris et al., 2012; Kesner, 2013; Cayco-Gajic and Silver, 2019). Here we implemented parallel circuit and behavioral assays of pattern separation in mice one week after concussive brain injury. Interestingly, we find that the reliability of GC firing is not altered after brain injury. Moreover, the GC output correlation was consistently lower than input spike train correlation in cells from both control and injured mice, indicating that temporal pattern separation at the cellular level continued to occur after brain injury. However, the critical ability of dentate GCs to decorrelate highly similar input patterns was reduced after brain injury. These results are analogous to the effect of partial $GABA_A$ receptor antagonism where GC output correlation is higher than in aCSF, yet, remains lower than input correlation indicating deficits in decorrelation of inputs by GCs (Fig. 2 and Madar et al., 2019a). Despite the limited decrease in GC cellular temporal pattern decorrelation after injury, behavioral discrimination of spatial location in an episodic memory test was severely compromised. It is possible that the posttraumatic increase in the number of GCs showing sustained increase in excitability, identified by ΔFosB labeling, degrades sparse coding in the dentate and contributes to decline in spatial location discrimination. Furthermore, we demonstrate a decrease in dentate hilar somatostatin neurons in brain injured mice that showed deficits in spatial discrimination. Since somatostatin neurons have been shown to contribute to the in the novel object location task

(Morales et al., 2021) used in our study, posttraumatic somatostatin neuron loss could contribute

to deficits in behavioral spatial discrimination. Together these findings identify early changes in

dentate inhibition and increase in GC activity that could compromise the ability of the dentate to

effectively decorrelate spike trains and engage distinct GC ensembles, likely contributing to

deficits in encoding episodic memories.

Posttraumatic changes in dentate circuit impact temporal decorrelation of input patterns.

There is considerable evidence for the dentate gyrus playing a critical role in distinguishing

context and spatial locations (Leutgeb et al., 2007; McHugh et al., 2007; Kesner, 2013; Senzai

and Buzsaki, 2017). Because events occur in space and time, patterned neuronal firing can be

essential for encoding spatial and temporal information (Kobayashi and Poo, 2004; Buzsaki,

2010; Tort et al., 2011; Eichenbaum, 2013). Experimental and computational studies examining

dentate gyrus processing of temporally patterned inputs have identified that individual GCs can

effectively orthogonalize temporal patterns in input spike trains. The ability of GCs to

decorrelate temporal patterns likely depends on a combination of their intrinsic properties, local

circuit inhibition, probabilistic synaptic release and spike-timing history resulting from short-

term synaptic dynamics (Yim et al., 2015; Madar et al., 2019b, a). By adopting a temporal

pattern separation paradigm, we confirmed that GCs effectively decorrelate input patterns,

particularly those with high similarity (Fig. 2). Although inhibition contributed to input

decorrelation (Fig. 2), GCs still exhibited some output decorrelation in gabazine (100 nM)

indicating that perforant path synaptic dynamics and GC intrinsic physiology contribute to

temporal pattern separation in GCs.

In mice one week after concussive brain injury, we demonstrated an increase in perforant path

evoked excitatory drive (Fig 1). These results and the lack of changes in GC intrinsic physiology

and loss of hilar somatostatin neurons early after FPI are consistent with earlier studies in rats

(Lowenstein et al., 1992; Santhakumar et al., 2000; Korgaonkar et al., 2020b). Moreover,

labeling for ΔFosB, which indicates sustained increases in excitability (You et al., 2017), is

increased one week after FPI. Therefore, it may seem surprising that the reliability and average

firing frequency of GC responses to temporally patterned inputs remains unchanged after brain

injury (Fig 3). Indeed, spike train reliability, a measure of the output correlation in response to

identical inputs, is consistently lower than unity due to contributions from network noise and

probabilistic transmitter release. A possible reason for maintenance of GC reliability despite the

posttraumatic increase in afferent drive (Fig. 1) could be because, in the absence of post-injury

changes in GC intrinsic physiology, setting the input stimulus intensity to elicit response in 50%

of trials limited the effect of the increased excitatory input. Moreover, GCs continue to

effectively decorrelate input spike patterns as demonstrated by the consistent observation that

output correlation remains lower than the input correlation even after brain injury (Fig 4C). Our

finding that GCs continue to decorrelate input patterns early after brain injury is also consistent

with recent findings in computational and experimental models of temporal lobe epilepsy (Yim

et al., 2015; Madar et al., 2021). However, our results reveal a significant decrease in temporal

pattern separation in response to input trains with high degrees of similarity, which suggests that

the network level changes including a partial loss of hilar somatostatin neurons do impact

decorrelation of input information. Although injury did not decrease GC spike reliability the

variance in firing frequency in response to identical input trains trended to decrease after injury.

It is interesting to speculated that the mechanisms contributing to reduced variance in GC firing

after injury could also reduce GC decorrelation of highly similar inputs.

Distinguishing between cellular decorrelation of spike timing and behavioral discrimination of

temporally ordered memories.

It must be noted that the temporal pattern separation paradigm used in this study focused on

timing of spike trains and does not directly correspond to the encoding of temporal sequences of

events. Indeed, it is unclear whether the dentate gyrus is involved in distinguishing the temporal

order of sequential events occurring over a span of minutes to days (Gilbert et al., 2001).

However, the dentate gyrus, and particularly the ongoing generation of adult-born neurons in the

dentate, has been shown to play a role in stratification of memories over the time course of

weeks (Clelland et al., 2009; Aimone et al., 2011; Nakashiba et al., 2012; Rangel et al., 2014;

Miller and Sahay, 2019). Interestingly, adult neurogenesis is altered after brain injury with early

increase at one week followed by a progressive decline at one month (Yu et al., 2008; Kernie and

Parent, 2010; Shapiro, 2016; Neuberger et al., 2017b; Ngwenya and Danzer, 2018; Clark et al.,

2020). Whether injury induced changes in neurogenesis alter stratification or timestamping of

contemporary memories and/or compromise spatial and contextual pattern separation at later

time points after injury remains to be examined. In this study we focused on the early time point,

one week after injury, when the injury-induced neurons are believed to be immature and unlikely

to contribute to local dentate feedback inhibition (Overstreet-Wadiche and Westbrook, 2006;

Temprana et al., 2015; Miller and Sahay, 2019). Additionally, we recorded GCs in the middle

and outer third of the granule cell layer so as to avoid immature adult-born neurons, typically

located in the inner third of the granule cell layer (Kerloch et al., 2018; Save et al., 2018).

Crucially, all GCs included in analysis had input resistance less than 300 MΩ consistent with

mature GC phenotype. Thus, our results identify that, at the level of individual putative mature

dentate GCs, spike reliability and ability to decorrelate spike train patterns persist despite the

early circuit changes after brain injury. Still, considering that temporal correlations between

dentate neurons at the sub-second time scale are important for spatial memory discrimination

(van Dijk and Fenton, 2018), our slice physiology data suggest that the dentate ability to encode

input timing is largely retained early after brain injury. Because the dentate gyrus is particularly

important for disambiguating highly similar patterns, the posttraumatic reduction in GC ability to

decorrelate highly similar inputs could contribute to deficits in pattern separation after brain

injury.

Mechanisms underlying altered behavioral pattern separation after brain injury.

In contrast to GC temporal pattern separation, we find that the ability to encode and discriminate

subtle differences in spatial location is profoundly compromised one week after brain injury (Fig.

5). Our results complement a previous study which showed that mice one week after mild to

moderate concussive brain injury are impaired in a sequential location discrimination task using

the radial arms maze (Correll et al., 2021). Unlike the radial arm maze, the novel object location

adopted in our study examines episodic memory and does not rely on repetition of discrete trial

procedures (Bekinschtein et al., 2013; Morales et al., 2021). At the medium difficulty level (60°

relocation) adopted in our study, the task has been shown to rely on dentate processing and is

compromised by selective suppression of dentate somatostatin neurons. Consistently, we find

that the behavioral deficit in novel object location one week after brain injury is accompanied by

loss of hilar somatostatin neurons suggesting that interneuron loss may drive deficits in spatial

pattern separation. Whereas loss of immature and highly excitable adult born neurons born prior

to injury could contribute to deficits in encoding novel object location, studies using the cortical

impact injury have reported no loss of GCs born prior to injury (Kang et al., 2022). Rather, these

neurons support enhanced feedback inhibition 8-10 weeks after injury. Moreover, it should be

noted that neurogenesis is increased rather than decreased one week after injury (Shapiro, 2016;

Neuberger et al., 2017b; Clark et al., 2020; Correll et al., 2021) and adult born neurons support

local circuit feedback inhibition of mature GCs 4-6 weeks into development (Temprana et al.,

2015; Miller and Sahay, 2019). Taken together, it is unlikely that injury induced changes in

dentate neurogenesis underlie deficits in spatial pattern separation observed one week after

injury. Instead, post-injury reduction of feedback inhibition, due to loss of somatostatin neurons,

and increases in basal activity of GCs, revealed by the enhanced ΔFosB staining, likely impair

the ability of the dentate circuit to effectively decorrelate entorhinal inputs into discrete active

GC ensembles and firing patterns.

Reconciling the modest loss in cellular temporal decorrelation with profound impairment in

behavioral spatial discrimination.

The apparent divergence in effect of brain injury on decorrelation of spike trains at the cellular

level and discrimination of spatial location at a behavioral level indicate that disambiguation of

information likely involves activation of distinct neuronal ensembles by input patterns. This is

consistent with findings in computational models (Myers and Scharfman, 2009, 2011).

Experimentally, due to the time for genetically encoded reporters to express (Guenthner et al.,

2013; Ramirez et al., 2013), it may be challenging to identify neurons activated by two distinct

and recent spatial inputs. However, there is some evidence for sparse but shared activation of

neurons by different spatial experiences (Tashiro et al., 2007). The basal increase in neuronal

excitability, identified by increase in ΔFosB labeled GCs after brain injury, could corrupt this

sparse and selective activity necessary for discrimination of spatial memories. A potential caveat

is that, whereas the circuit experiments focused on the outer molecular layer, which receives the

lateral entorhinal cortical fibers carrying contextual information, the spatial information

examined in the behavioral spatial discrimination study is thought to be carried by inputs from

the medial entorhinal cortex. Future studies examining decorrelation of medial perforant path

inputs and behavioral analysis of context encoding could resolve whether the differences

identified here are pathway specific. Analysis of spatiotemporal activity patterns of neuronal

ensembles during behaviors or imaging circuit activity in vitro in response to spatiotemporal

input patterns would help resolve how the dentate disambiguates input patterns.

In summary, our study identifies that early posttraumatic changes, including persistent increase

in excitability and loss of somatostatin interneurons, reduce temporal pattern separation by

dentate GCs and degrade the ability to discriminate between similar spatial locations. These

changes likely contribute to working memory deficits after brain injury.

Acknowledgements

We thank Dr. Krista Marrero for thoughtful comments and discussion.

Funding

The project was supported by National Institutes of Health (NIH) NINDS R01 NS069861,

R01NS097750 to V.S., NIH/NINDS F31NS110220 to L.C, F31NS120620 to S.N, and American

Epilepsy Society (AES) #957615 to A.H.

Reference List

Afrasiabi M, Gupta A, Xu H, Swietek B, Santhakumar V (2021) Differential Activity-Dependent

Increase in Synaptic Inhibition and Parvalbumin Interneuron Recruitment in Dentate Granule

Cells and Semilunar Granule Cells. Journal of Neuroscience.

Aimone JB, Deng W, Gage FH (2011) Resolving new memories: a critical look at the dentate

gyrus, adult neurogenesis, and pattern separation. Neuron 70:589-596.

Alosco ML et al. (2023) Neuropsychological test performance of former American football

players. Alzheimers Res Ther 15:1.

Annegers JF, Coan SP (2000) The risks of epilepsy after traumatic brain injury. Seizure 9:453-

457.

Arciniega H, Shires J, Furlong S, Kilgore-Gomez A, Cerreta A, Murray NG, Berryhill ME

(2021) Impaired visual working memory and reduced connectivity in undergraduates with a

history of mild traumatic brain injury. Sci Rep 11:2789.

Bekinschtein P, Kent BA, Oomen CA, Clemenson GD, Gage FH, Saksida LM, Bussey TJ (2013)

BDNF in the dentate gyrus is required for consolidation of "pattern-separated" memories. Cell

reports 5:759-768.

Bonislawski DP, Schwarzbach EP, Cohen AS (2007) Brain injury impairs dentate gyrus

inhibitory efficacy. Neurobiol Dis 25:163-169.

Braganza O, Mueller-Komorowska D, Kelly T, Beck H (2020) Quantitative properties of a

feedback circuit predict frequency-dependent pattern separation. eLife 9.

Buzsaki G (2010) Neural syntax: cell assemblies, synapsembles, and readers. Neuron 68:362-

385.

Cayco-Gajic NA, Silver RA (2019) Re-evaluating Circuit Mechanisms Underlying Pattern

Separation. Neuron 101:584-602.

Clark LR, Yun S, Acquah NK, Kumar PL, Metheny HE, Paixao RCC, Cohen AS, Eisch AJ

(2020) Mild Traumatic Brain Injury Induces Transient, Sequential Increases in Proliferation,

Neuroblasts/Immature Neurons, and Cell Survival: A Time Course Study in the Male Mouse

Dentate Gyrus. Front Neurosci 14:612749.

Clelland CD, Choi M, Romberg C, Clemenson GD, Jr., Fragniere A, Tyers P, Jessberger S,

Saksida LM, Barker RA, Gage FH, Bussey TJ (2009) A functional role for adult hippocampal

neurogenesis in spatial pattern separation. Science 325:210-213.

Correll EA, Ramser BJ, Knott MV, McCullumsmith RE, McGuire JL, Ngwenya LB (2021)

Deficits in pattern separation and dentate gyrus proliferation after rodent lateral fluid percussion

injury. IBRO Neurosci Rep 10:31-41.

Eichenbaum H (2013) Memory on time. Trends Cogn Sci 17:81-88.

Folweiler KA, Xiong G, Best KM, Metheny HE, Nah G, Cohen AS (2020) Traumatic Brain

Injury Diminishes Feedforward Activation of Parvalbumin-Expressing Interneurons in the

Dentate Gyrus. eNeuro 7.

Frankowski JC, Kim YJ, Hunt RF (2019) Selective vulnerability of hippocampal interneurons to

graded traumatic brain injury. Neurobiol Dis 129:208-216.

Gilbert PE, Kesner RP, Lee I (2001) Dissociating hippocampal subregions: double dissociation

between dentate gyrus and CA1. Hippocampus 11:626-636.

GoodSmith D, Chen X, Wang C, Kim SH, Song H, Burgalossi A, Christian KM, Knierim JJ

(2017) Spatial Representations of Granule Cells and Mossy Cells of the Dentate Gyrus. Neuron

93:677-690 e675.

Guenthner CJ, Miyamichi K, Yang HH, Heller HC, Luo L (2013) Permanent genetic access to

transiently active neurons via TRAP: targeted recombination in active populations. Neuron

78:773-784.

Gupta A, Dovek L, Proddutur A, Elgammal FS, Santhakumar V (2022) Long-Term Effects of

Moderate Concussive Brain Injury During Adolescence on Synaptic and Tonic GABA Currents

in Dentate Granule Cells and Semilunar Granule Cells. Front Neurosci 16:800733.

Gupta A, Proddutur A, Chang YJ, Raturi V, Guevarra J, Shah Y, Elgammal FS, Santhakumar V

(2020) Dendritic morphology and inhibitory regulation distinguish dentate semilunar granule

cells from granule cells through distinct stages of postnatal development. Brain structure &

function 225:2841-2855.

Hainmueller T, Bartos M (2020) Dentate gyrus circuits for encoding, retrieval and discrimination

of episodic memories. Nat Rev Neurosci 21:153-168.

Hamm RJ, Temple MD, Pike BR, O'Dell DM, Buck DL, Lyeth BG (1996) Working memory

deficits following traumatic brain injury in the rat. J Neurotrauma 13:317-323.

Kahn JB, Port RG, Yue C, Takano H, Coulter DA (2019) Circuit-based interventions in the

dentate gyrus rescue epilepsy-associated cognitive dysfunction. Brain 142:2705-2721.

Kang YJ, Lee SH, Boychuk JA, Butler CR, Juras JA, Cloyd RA, Smith BN (2022) Adult Born

Dentate Granule Cell Mediated Upregulation of Feedback Inhibition in a Mouse Model of

Traumatic Brain Injury. J Neurosci 42:7077-7093.

Kerloch T, Clavreul S, Goron A, Abrous DN, Pacary E (2018) Dentate Granule Neurons

Generated During Perinatal Life Display Distinct Morphological Features Compared With Later-

Born Neurons in the Mouse Hippocampus. Cereb Cortex.

563 Kernie SG, Parent JM (2010) Forebrain neurogenesis after focal Ischemic and traumatic brain

564 injury. Neurobiol Dis 37:267-274.

565 Kesner RP (2013) An analysis of the dentate gyrus function. Behav Brain Res 254:1-7.

566 Kobayashi K, Poo MM (2004) Spike train timing-dependent associative modification of

567 hippocampal CA3 recurrent synapses by mossy fibers. Neuron 41:445-454.

568 Korgaonkar AA, Nguyen S, Li Y, Sekhar D, Subramanian D, Guevarra J, Pang KCH,

569 Santhakumar V (2020a) Distinct cellular mediators drive the Janus faces of toll-like receptor 4

570 regulation of network excitability which impacts working memory performance after brain

571 injury. Brain Behav Immun.

572 Korgaonkar AA, Li Y, Sekhar D, Subramanian D, Guevarra J, Swietek B, Pallottie A, Singh S,

573 Kella K, Elkabes S, Santhakumar V (2020b) Toll-like Receptor 4 Signaling in Neurons Enhances

574 Calcium-Permeable alpha-Amino-3-Hydroxy-5-Methyl-4-Isoxazolepropionic Acid Receptor

575 Currents and Drives Post-Traumatic Epileptogenesis. Ann Neurol 87:497-515.

576 Kumar S, Rao SL, Chandramouli BA, Pillai S (2013) Reduced contribution of executive

577 functions in impaired working memory performance in mild traumatic brain injury patients. Clin

578 Neurol Neurosurg 115:1326-1332.

579 Larimer P, Strowbridge BW (2010) Representing information in cell assemblies: persistent

580 activity mediated by semilunar granule cells. Nature neuroscience 13:213-222.

581 Leh SE, Schroeder C, Chen JK, Mallar Chakravarty M, Park MT, Cheung B, Huntgeburth SC,

582 Gosselin N, Hock C, Ptito A, Petrides M (2017) Microstructural Integrity of Hippocampal

583 Subregions Is Impaired after Mild Traumatic Brain Injury. J Neurotrauma 34:1402-1411.

584 Leutgeb JK, Leutgeb S, Moser MB, Moser EI (2007) Pattern separation in the dentate gyrus and

585 CA3 of the hippocampus. Science 315:961-966.

Lowenstein DH, Thomas MJ, Smith DH, McIntosh TK (1992) Selective vulnerability of dentate

hilar neurons following traumatic brain injury: a potential mechanistic link between head trauma

and disorders of the hippocampus. J Neurosci 12:4846-4853.

Madar AD, Ewell LA, Jones MV (2019a) Pattern separation of spiketrains in hippocampal

neurons. Sci Rep 9:5282.

Madar AD, Ewell LA, Jones MV (2019b) Temporal pattern separation in hippocampal neurons

through multiplexed neural codes. PLoS computational biology 15:e1006932.

Madar AD, Pfammatter JA, Bordenave J, Plumley EI, Ravi S, Cowie M, Wallace EP, Hermann

BP, Maganti RK, Jones MV (2021) Deficits in Behavioral and Neuronal Pattern Separation in

Temporal Lobe Epilepsy. J Neurosci 41:9669-9686.

McAllister TW (1992) Neuropsychiatric sequelae of head injuries. Psychiatr Clin North Am

15:395-413.

McHugh T, Laforce R, Jr., Gallagher P, Quinn S, Diggle P, Buchanan L (2006) Natural history

of the long-term cognitive, affective, and physical sequelae of mild traumatic brain injury. Brain

Cogn 60:209-211.

McHugh TJ, Jones MW, Quinn JJ, Balthasar N, Coppari R, Elmquist JK, Lowell BB, Fanselow

MS, Wilson MA, Tonegawa S (2007) Dentate gyrus NMDA receptors mediate rapid pattern

separation in the hippocampal network. Science 317:94-99.

Meier TB, Savitz J, Singh R, Teague TK, Bellgowan PS (2016) Smaller Dentate Gyrus and CA2

and CA3 Volumes Are Associated with Kynurenine Metabolites in Collegiate Football Athletes.

J Neurotrauma 33:1349-1357.

Miller SM, Sahay A (2019) Functions of adult-born neurons in hippocampal memory

interference and indexing. Nat Neurosci 22:1565-1575.

Morales C, Morici JF, Espinosa N, Sacson A, Lara-Vasquez A, Garcia-Perez MA, Bekinschtein

P, Weisstaub NV, Fuentealba P (2021) Dentate Gyrus Somatostatin Cells are Required for

Contextual Discrimination During Episodic Memory Encoding. Cereb Cortex 31:1046-1059.

Morris AM, Churchwell JC, Kesner RP, Gilbert PE (2012) Selective lesions of the dentate gyrus

produce disruptions in place learning for adjacent spatial locations. Neurobiol Learn Mem

97:326-331.

Myers CE, Scharfman HE (2009) A role for hilar cells in pattern separation in the dentate gyrus:

a computational approach. Hippocampus 19:321-337.

Myers CE, Scharfman HE (2011) Pattern separation in the dentate gyrus: a role for the CA3

backprojection. Hippocampus 21:1190-1215.

Nakashiba T, Cushman JD, Pelkey KA, Renaudineau S, Buhl DL, McHugh TJ, Rodriguez

Barrera V, Chittajallu R, Iwamoto KS, McBain CJ, Fanselow MS, Tonegawa S (2012) Young

dentate granule cells mediate pattern separation, whereas old granule cells facilitate pattern

completion. Cell 149:188-201.

Neuberger EJ, Gupta A, Subramanian D, Korgaonkar AA, Santhakumar V (2017a) Converging

early responses to brain injury pave the road to epileptogenesis. J Neurosci Res.

Neuberger EJ, Swietek B, Corrubia L, Prasanna A, Santhakumar V (2017b) Enhanced Dentate

Neurogenesis after Brain Injury Undermines Long-Term Neurogenic Potential and Promotes

Seizure Susceptibility. Stem Cell Reports 9:972-984.

Neunuebel JP, Knierim JJ (2014) CA3 retrieves coherent representations from degraded input:

direct evidence for CA3 pattern completion and dentate gyrus pattern separation. Neuron 81:416-

427.

Ngwenya LB, Danzer SC (2018) Impact of Traumatic Brain Injury on Neurogenesis. Front

Neurosci 12:1014.

Overstreet-Wadiche LS, Westbrook GL (2006) Functional maturation of adult-generated granule

cells. Hippocampus 16:208-215.

Ramirez S, Tonegawa S, Liu X (2013) Identification and optogenetic manipulation of memory

engrams in the hippocampus. Front Behav Neurosci 7:226.

Rangel LM, Alexander AS, Aimone JB, Wiles J, Gage FH, Chiba AA, Quinn LK (2014)

Temporally selective contextual encoding in the dentate gyrus of the hippocampus. Nat Commun

5:3181.

Rao V, Lyketsos C (2000) Neuropsychiatric sequelae of traumatic brain injury. Psychosomatics

41:95-103.

Rolls ET, Kesner RP (2006) A computational theory of hippocampal function, and empirical

tests of the theory. ProgNeurobiol 79:1-48.

Rusnak M (2013) Traumatic brain injury: Giving voice to a silent epidemic. Nat Rev Neurol

9:186-187.

Santhakumar V, Ratzliff AD, Jeng J, Toth K, Soltesz I (2001) Long-term hyperexcitability in the

hippocampus after experimental head trauma. AnnNeurol 50:708-717.

Santhakumar V, Bender R, Frotscher M, Ross ST, Hollrigel GS, Toth Z, Soltesz I (2000)

Granule cell hyperexcitability in the early post-traumatic rat dentate gyrus: the 'irritable mossy

cell' hypothesis. J Physiol 524 Pt 1:117-134.

Save L, Baude A, Cossart R (2018) Temporal Embryonic Origin Critically Determines Cellular

Physiology in the Dentate Gyrus. Cereb Cortex.

Senzai Y, Buzsaki G (2017) Physiological Properties and Behavioral Correlates of Hippocampal

Granule Cells and Mossy Cells. Neuron 93:691-704 e695.

Shapiro LA (2016) Altered hippocampal neurogenesis during the first 7 days after a fluid

percussion traumatic brain injury. Cell Transplant.

Smith CJ, Xiong G, Elkind JA, Putnam B, Cohen AS (2015) Brain Injury Impairs Working

Memory and Prefrontal Circuit Function. Frontiers in neurology 6:240.

Smith CJ, Johnson BN, Elkind JA, See JM, Xiong G, Cohen AS (2012) Investigations on

alterations of hippocampal circuit function following mild traumatic brain injury. Journal of

visualized experiments : JoVE:e4411.

Tashiro A, Makino H, Gage FH (2007) Experience-specific functional modification of the

dentate gyrus through adult neurogenesis: a critical period during an immature stage. J Neurosci

27:3252-3259.

Temprana SG, Mongiat LA, Yang SM, Trinchero MF, Alvarez DD, Kropff E, Giacomini D,

Beltramone N, Lanuza GM, Schinder AF (2015) Delayed coupling to feedback inhibition during

a critical period for the integration of adult-born granule cells. Neuron 85:116-130.

Tort AB, Komorowski R, Kopell N, Eichenbaum H (2011) A mechanism for the formation of

hippocampal neuronal firing patterns that represent what happens where. Learning & memory

18:718-727.

Toth Z, Hollrigel GS, Gorcs T, Soltesz I (1997) Instantaneous perturbation of dentate

interneuronal networks by a pressure wave-transient delivered to the neocortex. J Neurosci

17:8106-8117.

Tsirka V, Simos P, Vakis A, Vourkas M, Arzoglou V, Syrmos N, Stavropoulos S, Micheloyannis

S (2010) Material-specific difficulties in episodic memory tasks in mild traumatic brain injury.

Int J Neurosci 120:184-191.

Vaishnavi S, Rao V, Fann JR (2009) Neuropsychiatric problems after traumatic brain injury:

unraveling the silent epidemic. Psychosomatics 50:198-205.

van Dijk MT, Fenton AA (2018) On How the Dentate Gyrus Contributes to Memory

Discrimination. Neuron 98:832-845 e835.

Yassa MA, Stark CE (2011) Pattern separation in the hippocampus. Trends Neurosci 34:515-

525.

Yim MY, Hanuschkin A, Wolfart J (2015) Intrinsic rescaling of granule cells restores pattern

separation ability of a dentate gyrus network model during epileptic hyperexcitability.

Hippocampus 25:297-308.

You JC, Muralidharan K, Park JW, Petrof I, Pyfer MS, Corbett BF, LaFrancois JJ, Zheng Y,

Zhang X, Mohila CA, Yoshor D, Rissman RA, Nestler EJ, Scharfman HE, Chin J (2017)

Epigenetic suppression of hippocampal calbindin-D28k by DeltaFosB drives seizure-related

cognitive deficits. Nat Med 23:1377-1383.

Yu TS, Zhang G, Liebl DJ, Kernie SG (2008) Traumatic brain injury-induced hippocampal

neurogenesis requires activation of early nestin-expressing progenitors. JNeurosci 28:12901-

12912.

Figure Legends

Figure 1. Enhanced dentate excitability one week after brain injury. (A) Schematic of slice

recording. (B) Representative eEPSC traces from dentate granule cells in sham (black) and FPI

(red). (C) Plot of eEPSC peak amplitude in granule cells recorded in responses to increasing

stimulus intensities (sham: 509.4, n =11; FPI: 759.4, n = 4, p<0.0001 by TW-RM ANOVA). D.

Representative images of staining for ΔFosB in sections (20 µm) from mice one week after sham

and FPI. Contour lines illustrate ROI drawn around granule cell layer (GCL) using DAPI to

visualize cell layer and circles around the cells detected by the automated algorithm. Scale

bar=100µm. (E) Histogram showing average number of ΔFosB-positive cells in GCL of sham

and FPI mice.

Figure 2. Robust temporal pattern separation by dentate granule cells. (A) Representative

input spike trains from the R75 input set are illustrated with each pattern (input trains #1-5) in a

distinct color. The GC firing evoked by three (of 10) repetitions of each input set is illustrated

with output trace color matched to corresponding input spike train. (B) R_{out} vs R_{in} plots

calculated with bin widths for correlation set to 10, 20, and 100 ms. Responses were elicited by

input sets with similarity indices ranging from R25 to R95. (C-D) Example traces of GC firing

responses elicited during 10 repetitions of the same representative (R75, pattern #4) input spike

train recorded in aCSF (C) and in 100 nM gabazine (D). (E) Summary of pairwise R

comparisons in aCSF and 100 nM gabazine (n = 4 cells). * Indicates p < 0.05 by Paired T-test.

Figure 3. Assessment of GC firing fidelity after FPI. (A-B) Example of a single input spike

train at R95 and the corresponding membrane voltage traces recorded in GC during 10

repetitions of the same stimulus in sham (A) and FPI (B) mice. (C-D) Summary plot of GC firing

frequency (C) and reliability index (D) of GC firing during R95 inputs in sham (n = 11 cells

from 3 mice) and FPI (n = 14 cells from 3 mice).

Figure 4. GC temporal pattern separation is maintained after brain injury. (A-B)

Representative input spike trains from the R95 input set are illustrated with each pattern (input

trains #1-5) in a distinct color. GC firing evoked by three repetitions of each input set in sham

(A) and FPI (B). Each membrane voltage trace is color matched to corresponding input spike

train. (C) Summary plot of R_{out} vs R_{in} recorded during presentation of stimulus sets with R25 and

R95 similarity indices in GCs from sham (black) and FPI (red) mice. (D-E) Summary plots

compare pairwise R_{out} in GCs from sham and FPI mice during presentation of input sets with

R25 (D, 90 recordings from 9 cells in 3 sham mice and 110 pairwise correlates from 11 cells in 3

FPI mice) and R95 (E, 110 pairwise correlates from 11 cells in 3 sham mice n = 140 pairwise

correlates from 14 cells in 3 FPI mice) * Indicates $p < 0.05$ by unpaired Welch's t-test.

Figure 5. Spatial recognition memory is impaired following FPI (A) Timeline of behavioral

task and tissue processing for histological staining. (B) Schematic representation of the novel

object location task performed one week post injury showing 4-day habituation in open chamber

and test day timeline and location of objects in sample and choice phases. (C-D) Summary plot

of discrimination ratio (C) and total duration of exploration (D) in sham (n = 8) and FPI (n=8)

mice. (E) Representative images of dentate sections immunostained for somatostatin illustrates

reduction in hilar somatostatin neurons one week after FPI, scale bar=100µm. (F) Summary

histogram of somatostatin cell counts in sections from the dentate hilus in sham and FPI mice.

Supplementary Figures

Supplementary Figure 1. Decrease in GC decorrelation of highly similar temporal input

patterns after brain injury. (A-B) Summary plots compare Pairwise R_{out}/R_{in} in GCs from sham

and FPI mice during presentation of input sets with R=0.25 (A, 90 pairwise correlates from 9

cells in 3 sham mice and 110 pairwise correlates from 11 cells in 3 FPI mice) and R=0.95 (B,

110 pairwise correlates from 11 cells in 3 sham mice n = 140 pairwise correlates from 14 cells in

3 FPI mice) * Indicates $p < 0.05$ by unpaired Student's t-test.

A SHAM

B FPI

C

D

A SHAM

B FPI

C
- Control
- FPI

D
R25

ns

R95

*