1 **Methods to Utilize Pulse Wave Velocity to Measure Alterations in Cerebral and**
2 **Cardiovascular Parameters**
3
4 **Authors and Affiliations:**
5

6 Andrea G. Marshall[1],*, Kit Neikirk[1],*, Bryanna Shao[1], Amber Crabtree[1], Zer Vue[1], Heather K.
7 Beasley[1], Edgar Garza-Lopez[2], Estevão Scudese[3], Celestine N. Wanjalla[4], Annet Kirabo[4], Claude
8 F Albritton[4,5], Sydney Jamison[4,5], Mert Demirci[4], Sandra A. Murray[6], Anthonya T. Cooper[6],
9 George E Taffet[7], Antentor O. Hinton, Jr[1,#], Anilkumar K. Reddy[7,#]
10
[1] Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN,
37232, USA.
[2] Department of Internal Medicine, University of Iowa, Iowa City, IA, 52242, USA.
[3] Laboratory of Biosciences of Human Motricity (LABIMH) of the Federal University of State of
Rio de Janeiro (UNIRIO), Rio de Janeiro, Brazil; Sport Sciences and Exercise Laboratory
(LaCEE), Catholic University of Petrópolis (UCP), Brazil
[4] Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, 37232, USA
[5] School of Graduate Studies, Meharry Medical College, Nashville, TN
[6] Department of Cell Biology, School of Medicine, University of Pittsburgh, Pittsburgh, PA, 15260, USA
[7] Department of Medicine, Baylor College of Medicine, One Baylor Plaza, Houston, TX 77030,
USA.
22
23
24 **Running Title:** CEREBRAL PWV METHOD
25
26
27 **Corresponding Authors**
28
Anilkumar K. Reddy, Ph.D.
Department of Medicine
Baylor College of Medicine
One Baylor Plaza, MS: BCM285
Houston, TX 77030, USA.
Ph: 713-798-6831
Email: areddy@bcm.edu
36
Antentor O. Hinton, Jr, Ph.D.
Department of Molecular Physiology and Biophysics
Vanderbilt School of Medicine Basic Sciences
The Vanderbilt Diabetes Research and Training Center
750 Robinson Research Building
2200 Pierce Ave
Nashville, TN 37232-0615
319-383-3095 (cell)
antentor.o.hinton.jr@Vanderbilt.Edu
46

Abstract:
Alzheimer's Disease (AD) is a global health issue, affecting over 6 million in the United States,
with that number expected to increase as the aging population grows. As a neurodegenerative
disorder that affects memory and cognitive functions, it is well established that AD is associated
with cardiovascular risk factors beyond only cerebral decline. However, the study of
cerebrovascular techniques for AD is still evolving. Here, we provide reproducible methods to
measure impedance-based pulse wave velocity (PWV), a marker of arterial stiffness, in the
systemic vascular (aortic PWV) and in the cerebral vascular (cerebral PWV) systems. Using
aortic impedance and this relatively novel technique of cerebral impedance to comprehensively
describe the systemic vascular and the cerebral vascular systems, we examined the sex-
dependent differences in 5x transgenic mice (5XFAD) with AD under normal and high-fat diet,
and in wild-type mice under a normal diet. Additionally, we validated our method for measuring
cerebrovascular impedance in a model of induced stress in 5XFAD. Together, our results show
that sex and diet differences in wildtype and 5XFAD mice account for very minimal differences
in cerebral impedance. Interestingly, 5XFAD, and not wildtype, male mice on a chow diet show
higher cerebral impedance, suggesting pathological differences. Opposingly, when we subjected
5XFAD mice to stress, we found that females showed elevated cerebral impedance. Using this
validated method of measuring impedance-based aortic and cerebral PWV, future research may
explore the effects of modifying factors including age, chronic diet, and acute stress, which may
mediate cardiovascular risk in AD.

New and Noteworthy: Here, we presented a new technique which is an application of the
concept of aortic impedance to determining cerebral impedance. While aortic PWV is typically
utilized to study aortic stiffness, we also developed a technique of cerebral PWV to study
cerebral vascular stiffness. This method may be useful in improving the rigor of studies that seek
to have a dual focus on cardiovascular and cerebral function.

Keywords: Cerebral Blood Flow, Aortic Impedance, Cerebral Impedance, Doppler Flow System

Introduction:

Alzheimer's disease (AD) is a progressive neurodegenerative disorder that affects millions
of people worldwide, a rate that is expected to grow with the aging population (1). AD causes
deficits in quality of life and daily living marked by a decline of memory and cognitive functions
(2). A hallmark feature of AD is the accumulation of aggregated amyloid beta (Aβ) protein in the
brain, which ultimately leads to synaptic dysfunction, neuronal loss, and cognitive impairment (3).
In addition to genetic factors, lifestyle and environmental factors such as stress and diet have been
implicated in the development and progression of AD (4). Beyond dementia, AD also impacts
cardiovascular function, as the amyloid beta plaques also accumulate in the heart (1). A key risk
factor of AD is hypertension, and it has been hypothesized that neuroinflammatory effects link
these pathologies (5). However, the full extent of the mechanisms which link AD and
cardiovascular disease remains poorly understood. Here, we sought to expand the understanding
of how AD alters cardiovascular and cerebral blood flow in conditions of chronic diet changes or
acute stress.

AD is well understood to cause neuropsychologic decline but is also understood to affect
cardiovascular dynamics. AD importantly is shown to potentially be associated with hypertension
(HTN) through the formation of neuritic plaques in cerebral vasculature and potentially limiting
cerebral blood flow (6). In human populations, even with correcting for risk factors such as age,
HTN reduction is correlated with increased retention of cognitive ability, indicating blood pressure
treatment as a potential mechanism to treat AD (1). This blood pressure linkage to AD is especially
pronounced when it comes to the systolic pressure (7). However, a past review on the matter shows
the issue to be far more complicated, as many previous studies are not using standardized HTN
measurements and there may be age-dependent and ethnicity-dependent influences of HTN on AD
development (8). Together, this underscores the need to better understand hemodynamics in AD.

Hemodynamics that includes pulse wave velocity (PWV) is important to study
aortic/arterial stiffness and cardiovascular function (9). Even when adjusting for factors including
age, race, and disease states, National Health and Nutrition Examination Survey studies indicate PWV
remains a strong predictor of overall mortality in human populations (10). Past studies looking at
echocardiographic parameters in humans with AD showed that diastolic function was more heavily
impacted in AD pathology, marked by increased arterial stiffness, increased atrial conduction
times, reduced blood flow, and altered mitral valve velocities (11). Carotid-femoral pulse wave
velocity increased aortic stiffness has also been linked as an independent predictor of cognitive
impairment for both dementia and AD (12). Another longitudinal study across over 1700
participants found that arterial stiffness and alterations in pulse pressure are antecedents for
cognitive decline (13). PVW can further be used to distinguish vascular dementia and AD, with
the former often expressing increased vascular stiffness relative to AD (14). Past studies have also
highlighted the importance of looking at both PWV as well as cerebral blood flow, through MRI,
with hypertensive symptoms altering heart and brain structure (15). Yet there remains controversy,
as some studies have found no correlation between carotid-femoral PWV and AD (16).

Our primary way to study AD cardiovascular-related parameters is through 5x transgenic
(5xFAD) mice, which have 5 AD-linked mutations. The 5xFAD mouse model expresses human
amyloid precursor protein (APP) and presenilin 1 (PS1) mutations, leading to the rapid
accumulation of Aβ and the development of AD-like pathology (17). This model also bears
numerous similarities to human models including Aβ-butyrylcholinesterase association and
hallmarks of progressive loss of cognitive function concomitant with reduced synaptic markers
(18). Due to apoptotic neuron loss, 5xFAD mice display memory deficits by 4 months of age (19).
This mouse model has been widely used to study the mechanisms underlying AD and to test
potential therapeutic interventions. Past studies using 3x transgenic mice models highlight that
with induced hypertension, mice had faster development of AD marked by upticks in Aβ, amyloid
plaque load, and phosphorylated tau (20). This validates that transgenic mice models retain the
relationship between cerebrovascular and cardiovascular dysfunction that has been observed in
AD.

To advance a new way of studying cerebrovascular dynamics in both wildtype and 5XFAD
mice, we developed a technique to measure cerebral pulse wave velocity. The relationship between
PWV and AD pathology in 5xFAD models remains unclear, especially as it comes to cerebral
pulse wave velocity (cPWV), a hemodynamic parameter that has been increasingly recognized as
an important predictor of cerebrovascular disease. While aortic stiffness can serve as a predictor
for strokes, it is limited in detection of minor changes which precede cerebrovascular changes such

as microbleeds (21). Similar to its aortic counterpart, cPWV measures the speed at which arterial
pressure or velocity waves propagate through the cerebral arteries. Measuring cerebral impedance
is a novel measure of the comprehensive characterization of cerebral blood vessels, which is highly
relevant as the brain is a highly vascularized organ that requires a constant supply of oxygen and
nutrients, with disruptions in blood flow being linked to cognitive impairment. In AD, this remains
highly relevant as it may relate to Aβ clearance (8). A recent clinical trial protocol has proposed
utilizing carotid-cerebral PWV as a mechanism to better understand acute ischemic strokes (22).
Here, we use pulsed Doppler ultrasound to simultaneously measure the arrival of the velocity wave
at two different arterial sites, one at the aortic arch and another at ophthalmic artery in the distal
internal carotid artery and measure the physical distance between the two sites to estimate cerebral
PWV (cPWV). Also, using carotid flow velocity and aortic blood pressure we measured cerebral
impedance and impedance-based PWV ($cPWV_{Zc}$) which is crucial for improving our
understanding of the pathophysiology of AD (Table 1).

To validate our methods, we studied sex-dependent, diet-dependent, and stress-induced
differences in WT and 5XFAD mice. While it is clear that females are more likely to get AD in
humans, there remains a gap in the literature in understanding sex-dependent differences in
cerebral hemodynamics in 5xFAD mouse models (18). Using a combination of aortic and cerebral
PWV, our findings provide important insights into the relationship between PWV and AD
pathology, as well as potential novel mechanisms of protection. By understanding how these
factors interact to influence arterial stiffness and cognitive function, we may be able to identify
new therapeutic targets for the treatment and prevention of AD.

Methods Before You Begin:

Animals. All animal protocols were approved by the Institutional Animal Care and Use Committee
of Baylor College of Medicine in accordance with the National Institutes of Health Guide for the
Care and Use of Laboratory Animals. We used 3 groups of 5xFAD mice at 4-5 months of age. The
diets of the 3 groups of mice consisted of standard commercial chow (2920X Harlan Teklad,
Indianapolis, IN, USA), high-fat diet (60% kCal from fat), and Equi diet () with free access to food
and water (**Figure S1**). Separately, we also used 8-9 month C57BL6J mice with chow diet only.
All the mouse groups are shown below.

Animals	Strain	Male	Female	Age	Diet
Group1	5xFAD	n=10	n=9	4-5mo	Normal Chow
Group2	5xFAD	n=10	n=9	4-5mo	HFD
Group3	5xFAD	n=5	n=5	4-5mo	Equi Diet (Pre & Post Stress)
Group4	C57BL6/J	n=5	n=5	8-9mo	Normal Chow

Mice were initially anesthetized with 2.5% isoflurane in the induction chamber and then
transferred to a heated (37±1°C) electrocardiography (ECG) board (MouseMonitor S, Indus
Instruments, Webster, TX) with the paws taped to the ECG electrodes and isoflurane maintained
at 1.5% via nose cone.

Doppler flow velocity measurements. We used a 20 MHz Doppler probe to measure doppler aortic
aortic arch velocity, ophthalmic artery velocity (representative of cerebral blood flow), and
abdominal aortic velocity signals to determine cerebral PWV (cPWV) and aortic PWV (aPWV)
(figure 1a). We also measured aortic outflow velocity and carotid flow velocity along with aortic
blood pressure, and mitral blood flow velocity (figure 1b) to determine cardiac and cerebral
hemodynamics. All signals were acquired and stored using Doppler Flow Velocity System (DFVS;
Indus Instruments, Webster, TX). We measured separation distance between aortic arch and OA
sites and between aortic arch and abdominal aortic sites to determine cPWV and aPWV,
respectively. We measured peak and mean aortic velocities, stroke distance (Sd), aortic ejection
time (ET), peak and mean aortic accelerations from aortic outflow velocity, and early peak and
atrial peak velocities, E/A ratio, E deceleration time, isovolumic contraction (IVCT) & relaxation
(IVRT) times from mitral inflow signals, and myocardial performance index (also known as Tei
index = (IVCT + IVRT)/ET). From the carotid flow velocity signal, we calculated peak, minimum,
and mean velocities, pulsatility index, and resistivity index are calculated.

Blood pressure measurements. Blood pressure measurements were made as previously described
(23–25), Briefly, a 1-French (0.33mm diameter) blood pressure catheter (SPR-1,000: Millar
Instruments, Inc., Houston, TX) was introduced via the isolated right carotid artery and advanced
into the ascending aorta to measure aortic pressure. About 2-3 second segments of blood pressure
signals were acquired (simultaneously with either aortic flow velocity or carotid flow velocity and
ECG signals) with the DFVS system. Systolic (SBP), diastolic (DBP), mean (MBP), pulse
pressures (PP), end-systolic pressure (ESP) and rate-pressure product (RPP) were calculated from
the recorded aortic blood pressure signals.

Determination of aortic and carotid impedance. The method to determine aortic impedance was
described elsewhere (22, 24-26). Aortic impedance is determined using aortic pressure-velocity
relationship (figure 2a). The foot of aortic pressure waveform was aligned with the foot of the
aortic velocity waveform to avoid potential errors in phase relation between pressure and velocity
signals. The signals are converted to frequency domain using fast Fourier transform and impedance
($|Z|$ $|P|/|V|$) parameters (peripheral vascular resistance [aZ_0], characteristic impedance [aZ_C], and
impedance at first harmonic [aZ_1]) are calculated. Aortic pulse wave velocity was calculated as
aZ_C/ρ (ρ-density of blood). The foot of the blood pressure waveform was aligned with the foot of
carotid flow velocity waveform and cerebral impedance was calculated in the same way as aortic
impedance (figure 2b) and the cerebral impedance parameters (cerebral vascular resistance [cZ_0],
cerebral characteristic impedance [cZ_C], and cerebral impedance at first harmonic [cZ_1]) were
calculated. Cerebral pulse wave velocity was calculated as cZ_C/ρ (ρ-density of blood).

Calculation of parameters to determine VVC. Elastance was determined as previously discussed
(25, 26). Arterial elastance (Ea) was calculated as ESP/SV (stroke volume, SV = Sd * aortic cross-
sectional area), end-systolic elastance (Ees) was calculated as ESP/ESV, ventricular-vascular
coupling (VVC) was calculated as Ea/Ees, and stroke work (SW) was calculated as ESP*SV (27).

Statistical analyses. All the data are presented as mean ± standard error of the mean (SEM). Dots
represent each sample, as sample size varies throughout. Statistical analyses were performed via
analysis using an unpaired T-test to compare conditions in each sex through Prism (GraphPad
Software; La Jolla, USA).

Step-By-Step to Measure Aortic and Carotid Velocity and Blood Pressure to Determine
Aortic and Cerebral Impedance
220
221
Non-invasive:
NOTE: Only allows for PWV measurement.
Before you begin: This protocol needs a Doppler system (see **Figure S2** – which shows the
workflow of signals, but it does not require the pressure system for the noninvasive measurement
of PWV).
1. On the day of the study, weigh the animal and anesthetize it in the induction chamber
using 3.0% isoflurane (mixed with 1L/min 100% O2).
2. Transfer the animal to the heated ECG board, and place in supine position with 1.5%
isoflurane supplied via a nose cone.
3. Apply artificial tears lubricant gel to the eyes to prevent dryness.
4. Apply ECG cream to the four paws and tape them to the ECG electrodes.
NOTE: Ensure excess gas is scavenged for the safety of the operator.
5. Remove the hair from a small area near sternal border and a small area near mid
abdominal area.
NOTE: The temporal canthus site near the eye used for the internal carotid artery branch of the
ophthalmic artery does not need hair removal.
6. Using a 20 MHz pulsed Doppler probe, place the tip on the chest at the aortic arch site
and aim toward mid-line to find an aortic arch signal. Place the probe in a tightly held
holder after the signal is found.
7. Holding the second probe by hand, place its tip at the temporal canthus of the mouse eye
and aim the probe toward the internal carotid artery to measure ophthalmic artery (OA)
flow velocity. Measure the separation distance between the two sites (where Doppler tips
are placed).
8. Treat the Arch & OA signals as I & Q (**Figure S1**) and combine to produce signals
(**Figure 1A**); save a 2-3 second segment of these signals. Determine the pulse transit time
offline and calculate cPWV (see upper right-hand panel **Figure 1A**).
9. Move the second probe to the abdominal location to find the abdominal aortic signal.
Measure the separation distance between the two sites.
10. Treat the Arch & Abd signals as I & Q and combine them to produce signals. Again, save
a 2-3 second segment of these signals, determine pulse transit time offline, and calculate
aPWV (see lower right panel in **Figure 1A**).
11. Once the measurements are made, wake the mouse up and return to the cage.
254
Invasive:
Before you begin: This specific protocol needs a Doppler system (see **Figure S2**- which shows
the workflow of signals), that allows for the measurement and acquisition of Doppler velocity
signals and ECG and blood pressure signals, simultaneously. A Millar pressure catheter and
amplifier system are needed to measure blood pressure signals. The Doppler velocity signal and
blood pressure are acquired simultaneously along with ECG for the temporal alignment of
signals.
NOTE: Avoid the usage of analgesics as this may suppress blood pressure.

1. On the day of the study, weigh the animal and anesthetize in the induction chamber using
3.0% isoflurane (mixed with 1L/min 100% O2).
2. Transfer to heated ECG board and place in supine position with 1.5% isoflurane supplied
via nose cone.
3. Apply artificial tears lubricant gel to eyes to prevent dryness.
4. Apply ECG cream to the four paws and tape them to the ECG electrodes.
NOTE: Ensure excess gas is scavenged for the safety of the operator.
5. Shave the hair from the neck area and apply hair removal cream to ensure all body hair is
removed.
6. Perform a pinch test to make sure the animal is unresponsive. If responsive, adjust
isoflurane level to 2.0% and then return to 1.5% and redo the pinch test.
7. Use a rectal temperature probe to monitor body temperature and adjust the heated board
to maintain body temperature at 37.0±0.5 °C.
NOTE: The heated board temperature is usually higher than the body temperature, so this should
be closely monitored.
8. Make a 60-70mm cut in the skin of the neck, to the right of the midline.
9. Expose the right carotid artery and cannulate with a pre-calibrated saline-soaked 1F
Millar catheter and secure with a suture.
10. Advance the catheter advanced into the aorta.
NOTE: Make sure the open skin is covered with wet gauze.
283
Waveforms: (See **Figure 1B** for devices/probe placements for respective measurements)
11. Using the Millar catheter, measure aortic blood pressure signals continuously.
12. Aim a 20 MHz pulsed Doppler probe tip at right suprasternal notch, caudally towards the
heart with a low angle for the measurements of aortic blood flow velocity.
13. For the measurement of carotid blood flow velocity, place the Doppler probe tip left of
the midline in the neck and aim caudally towards the heart at as low an angle as possible.
14. For the measurement of mitral flow velocity, reposition the Doppler probe tip under the
xiphoid and aim rostrally toward the heart.
15. Record Doppler signals. Doppler signals are processed in real time and displayed as
Doppler spectrograms on the screen along with the blood pressure and ECG waveforms.
16. Record the following sets of signals/waveforms: **a**. Aortic blood pressure & flow velocity
and ECG; **b**. Aortic blood pressure & carotid flow velocity and ECG; and **c**. Mitral flow
velocity signals with ECG.
NOTE: For each of the signal sets, collect 2-3 second segments of data for offline analysis.
17. Extract aortic and carotid flow velocity waveforms along with respective blood pressure
waveforms.
18. Calculate aortic impedance and cerebral impedance, as shown in **Figures 2A & 2B**,
respectively. Using a similar method to how aortic impedance is traditionally measured,
cerebral impedance is derived from pressure waveforms of carotid velocity (**Table 1**).
Use aortic pressure and carotid signal to equalize these values.
NOTE: Discrete Fourier transform can be computed using a simple custom code, MATLAB, or
other programs.
306

Representative Results

While we measured general parameters, such as body weight and aortic cross-sectional
area, the aim of this study was to validate our technique across experimental conditions (**Figure
S1**). To begin with, we looked at wildtype mice with normal chow diet at an age of 8 months old,
representing a standard model. Sex-dependent differences in LV afterload were evaluated using
aortic impedance, which showed non-statistically significant differences of impedance parameters
Zp, Z1, Zc, or PWVz (**Figure 3A-D**). When performing the cerebral counterpart to this
measurement, while many measurements remained non-significant, females exhibited a lower
cerebral Zp (**Figure 3E-H**).

To recapitulate these findings in a transgenic model, we looked at sex-dependent
differences in aortic and cerebral impedance in 5XFAD mice chow-fed cohort that was 5-months
old. LV afterload was evaluated using aortic impedance, which showed non-statistically significant
differences in impedance parameters Zp, Z1, Zc, or PWVz (**Figure 4A-D**), but slightly more intra-
cohort variability. When examining cerebral PWV, we found that both Zc and PWVz were both
significantly lower in female mice (**Figure 4E-H**). These findings indicate that fine differences at
baseline conditions may be detected, as well as showing that the 5XFAD mice differed in sex-
dependent differences, as compared to WT mice. From there, we sought to validate the
applicability of this aortic and cerebral measurement method with changes in diets.

A cohort of 8-week-old 5XFAD mice was subject to an HFD for 12 weeks and compared
to the aforementioned cohort of 5XFAD mice on a chow diet. Generally, as compared to the chow
cohort, the HFD diet cohort did not exhibit a significant difference in aortic or cerebral PWV
parameters (**Figure S3**). However, when looking at sex-dependent differences within the 5XFAD
cohort fed an HFD, we observed that cerebral parameters no longer showed a significant
difference, but aortic Zc and PWVz were significantly lower in female mice (**Figure 5A-H**). These
results that our aortic and cerebral methods have the sensitivity to detect minute sex-dependent
differences that arise as a result of changes in diet.

Finally, we sought to apply this method to mice undergoing an acute chronic stress
condition. Prior to subjecting 5XFAD mice to stress, non-invasive parameters were collected, as
described above. Looking at Tei index, a measure of myocardial performance and cerebral
separation distance, another measurement allowed for by non-invasive techniques, we saw no
significant differences (**Figure S4A-B**). Additionally, measuring pulse transit time (PTT), which
represents the interval for the aortic or cerebral pulse pressure wave to travel from the aortic or
cerebral valve to a peripheral site (27), and PWV showed no significant differences for aortic or
cerebral measurements (**Figure S4C-D**).

Given that, this non-invasive method validated sex-dependent PWV was minimally
different before stress, we subjected a cohort of 5XFAD mice to 1 hour of acute stress. When sexes
were grouped together, we saw that every measurement of aortic impedance, Zp, Z1, Zc, or PWVz,
was significantly higher in stressed mice (**Figure 5A-D**). Additionally, cerebral Zp and Z1 were
also found to be significantly higher in the cohort subjected to stress, as compared to a non-acutely
stressed chow-fed 5XFAD cohort (**Figure 5E-H**). While we investigated sex-dependent
differences in the stressed cohort and between cohorts, generally these were non-significant
(**Figure S5**). Together, these results validate that our method of aortic and cerebral methods can
effectively use Doppler measurements to observe differential chronic diet-dependent and acute
stress-induced changes in hemodynamics in 5xFAD and WT mice cohorts.

Discussion:

<u>Aortic and Cerebral impedance in 5XFAD</u>

When evaluating the aortic peripheral vascular resistance (Zp,), strength of wave reflections from the periphery (Z1), characteristic impedance (Zc,), and PWV we did not note a significant sex-dependent difference in both control and 5XFAD mice compared to their control counterpart (**Figure 3-4**). Notably, in both control and 5XFAD mice we did see that only females displayed increased cerebral impedance. In the case of the WT, females show a slightly lower cerebral impedance Zp, while the 5XFAD model differentially shows decreased Zc and PWV. This can imply that neurovascular coupling or conduction of electrical signals within the brain may be uniquely protective in females with AD. Interestingly, one past study suggested that cerebral blood flow is increased in AD risk states, suggesting a complex relationship between cPWV and AD that must be further investigated (28).

While 5XFAD mice serve as a strong model for human AD that has been used in the past, it lacks neurofibrillary tangles which human models would otherwise have (17). Additionally, while AD in humans has clear sex-dependent differences with increased CVD risk (33, 34), a lesser effect was observed here. This may be due to mouse estrous cycles which we did not consider here (35); thus, it is unclear whether our findings can be extrapolated to humans. Another limitation of our study is that we did not measure Aβ deposition levels, which may be correlated with PWV (30), or other molecular markers of AD pathology. Therefore, it is possible that our findings do not fully capture the complex relationship between hemodynamic parameters and AD pathology. Finally, our study only examined one-time points (12 weeks), and it is possible that longer development of mice may have different effects on hemodynamic parameters and AD pathology, as the latter is linked to aging.

Beyond this, future studies may look at the compounding effects of other experimental changes, such as induced hypertension, as the effectors that modulate sex-dependent differences in aortic and cerebral PWV of 5XFAD mice. Furthermore, there is an unmet need of understanding how aortic and cerebral PWV changes across the aging process for 5XFAD. Aging is especially important to study in the context of 5XFAD as they present a truncated development of AD pathology compared to human counterparts, marked by lower levels of Aβ with isomerized D7 and logarithmic plaque deposition beginning at 2 months of age, indicating that aging can uniquely affect 5XFAD mice (29, 30). Furthermore, past studies have shown that while changes in pulse pulsatility, are more common in younger individuals, stiffness is more likely to arise in older AD patients (31, 32). While 5xFAD females had less severe cardiovascular function impairments than is observed in human AD (33, 34), it is possible that compounding factors such as age and estrous cycles are implicated in our findings (35), highlighting the importance of future studies considering stratifications across sex of PWV while considering these factors. Investigating these effectors may also pave the way for future therapies, as PWV-dependence on estrogen receptors suggests a future avenue of estrogen-driven therapies to mitigate sex-dependent differences in AD.

<u>Aortic and Cerebral impedance in HFD</u>

While a diet-dependent difference was not observed in 5XFAD, we did not that sex-
dependent differences did not follow the same pattern in the chow and HFD cohorts. Notably,
characteristic impedance (Zc; average of 2–10 harmonics of impedance modulus), and pulse wave
velocity (PWV; aortic stiffness index) were both significantly lower in 5xFAD HFD females mice
compared to their male counterparts (**Figure 5**). Both of these measures of aortic impedance are
methods of considering the change of pressure in relation to changes in velocity to consider
hydraulic external load experienced by the left ventricle (36). Reduced parameters of aortic
impedance generally signify a reduced aortic wall stiffness (36). Similarly, PWV typically changes
concomitantly with pulse pressure, and high PWV, a marker of arterial stiffness, is known to occur
across aging and with AD (13). Even in healthy adults, PWV generally increases with aging (37),
so it is possible that this sex-dependent difference may lessen in an older mice cohort. Furthermore,
PWV commonly decreases alongside blood pressure in the case of medications such as anti-
hypertensives (38). Notably, PWV, which is measured at end-diastolic pressure, can be
independent of blood pressure, which is measured at peak systolic pressure, due to less noticeable
effects of stiffness in end-diastolic (24). It is still, however, possible that the HFD-induced increase
in PWV in male is simply indicative of the higher BP in the 5xFAD HFD cohort. However, given
that increased PWV is commonly observed in AD (13), it may also point toward certain
cardiovascular differences in the 5xFAD model.

Contrastingly, when looking at cerebral impedance measurements, a sex-dependent
difference is no longer observed in the HFD cohort. While variability in the male cohort is high,
this is unlikely to account for this difference. Thus, it is possible that at baseline, while 5XFAD
females are more resistant to elevated cerebral impedance, chronic diet change ameliorates this
advantage. This finding highlights the potential of understanding the confluence of sex and gender.
Notably, a prior study showed that while N-acetylneuraminic acid is responsible for HFD-induced
increased cognitive impairment in 5xFAD, RNA-sequencing showed diet and AD have different
roles in the microglia (39). This suggests that while HFD and AD may affect each other in the
brain, they may remain independent effectors in the heart. Other studies have found that HFD
modulates plasma metabolites more so than other factors in 5xFAD mice and amplifies the role of
sex (40), showing the necessity of exploring sex-dependent differences on diet in the future. Past
studies have suggested that female 5xFAD mice, while more susceptible to metabolic dysfunction,
HFD may serve a protective mechanism (41).

One other potential avenue in which AD pathology has an interplay with HFD-induced
changes are through mitochondria. Previous studies have shown that HFD had depressed
respiration concomitant with reduced bioenergetics in the liver (42). In Alzheimer's Disease,
mitochondria are known to decline in axons (43), in 5xFAD mice mitochondria undergo fission in
an age-dependent manner which is a driving force between loss of cognitive function and
pathology progression observed in 5xFAD mice (44). This loss of function may be partially
attributed to oxidative stress with accumulates across aging in 5xFAD mice (45). The role of
mitochondria in AD pathology is further underscored by drugs that target voltage-dependent anion
channel-1 to protect against pathology by stopping mitochondria dysfunction (46). While these
effects have been observed mainly to occur in neurons, future studies may explore if mitochondrial
function is also affected in the heart. Notably, in HFD cases alone, mitochondrial oxidative
phosphorylation is actually increased which protects against contractile function changes in heart

failure (47). This is despite increased oxidative stress, which impairs mitochondrial function,
induced by HFD in the brain in mice (48). One consideration is that few studies have yet looked
at oxidative stress in cardiac tissue, which may have differential regulation than neuronal oxidative
stress (49). Nonetheless, these contrasting roles between HFD and 5XFAD conditions may
partially explain some of the altered PWV that we observed.

Future studies may look at the compounding effects of other experimental changes, such
as high-salt diet changes in addition to HFD, as well as measuring if there is differential plaque
formation under experimental conditions of 5xFAD mice. Beyond this, the specific composition
of the diet may strongly affect 5xFAD's response. For example, while in the past a HFD causes
increased neuroinflammation and plaque accumulation (50), a ketogenic diet (high fat/low-
carbohydrate) improved cognition through inverse effects of reducing neuroinflammation (51),
which may reduce cardiovascular deficits. This may be due to the glycemic content of a diet more
heavily influencing the effects on neuroinflammation (52). Thus, this suggests that depending the
composition of high-fat relative to salt and carbs can alter positive or negative effects of it; thus,
future studies may examine hemodynamic alterations with a high-fat, salt, and sugar diet model
(53), which causes increased neuroinflammation in WT mice, to better understand the influence
of diet akin to that of the western-populations on AD pathology. Beyond this, since we had middle-
aged 5xFAD mice (i.e., 4-5 months), it is possible that younger or older mice would respond
differently to an HFD due to altered non-linear Aβ accumulation that occurs in 5xFAD mice (29).

<u>Aortic and Cerebral Impedance in Acute Stress Conditions</u>

Finally, we examined acute stress conditions. Despite baseline 5XFAD conditions before
stress exposure, stress was significant different than the 5XFAD mice they were compared to.
These results generally showed that all measurements of aortic impedance had a significant
increase following stress (**Figure 6**). Additionally, cerebral impedance measures of Zp and Z1
were increased. Given that chronic stress has arisen as a risk factor for AD (54)p, these findings
suggest a poorly explicated interplay through which stress may contribute to AD pathology
through cerebrovascular effects.

Notably, sex-dependent differences were not drastic in cerebral impedance of the HFD
cohort (**Figure S5**). This suggests that, like an HFD challenge, while females may have slightly
lower cerebral impedance at baseline in 5XFAD, these sex-dependent differences may fade under
certain chronic or acute challenges. These findings suggest that the relationship between PWV and
AD pathology in animal models may be more complex than previously thought and highlight the
importance of studying multiple hemodynamic parameters in the context of AD. Females may be
less vulnerable to AD partially due to reduced cerebral impedance Zp, however global increases
in aortic and cerebral PWV following stress induction may erase sex-dependent differences.
Interestingly, past studies have shown that stress in females only, and not males, leads to elevated
beta-amyloid (55). This highlights that PWV and cPWV are distinct from overall beta-amyloid
accumulation, and not necessarily correlative, emphasizing the importance of continuing to study
PWV and cPWV in a range of conditions.

Further studies are needed to elucidate the mechanisms underlying these findings and their
implications for the development and progression of AD. In conclusion, our study showed no
significant changes in most of the sex-dependent hemodynamic parameters examined, with the
most pertinent change but confirmed the deleterious effects of stress, which displayed slight sex-
dependent differences.

Conclusions

5XFAD mice are important new experimental models for the study of Alzheimer's Disease
(17), but such models have limited research evaluating their *in situ* global cardiovascular function.
Together our data also highlight that AD pathology should not only be considered in the context
of tau pathology and cognitive decline but aortic and cerebral impedance metrics as well. In
example, recent studies have demonstrated that Resveratrol can reduce HFD-induced accelerated
cognitive decline in 5xFAD through proteolytic mechanisms (56), but these same pathways may
not change HFD-induced cardiovascular remodeling. Thus, the impact of AD on cardiovascular
parameters, especially under altered environmental states must be considered. Beyond using
noninvasive measurements of Doppler mitral inflow and aortic flow velocity, invasive
measurements of aortic and LV pressure, and the calculated aortic impedance, we also evaluated
cerebral impedance.

In the future, cPWV may also aid in the early detection and monitoring of other
neurological disorders. While arterial PWV may serve as a mechanism to explore cerebral
perfusion, cerebral microbleeds have arisen as markers that are important to study in AD (8). Other
studies have also shown that glutamate chemical exchange saturation transfer has been observed
to concomitantly decrease with cerebral blood flow in 5XFAD mice, suggesting other novel
imaging techniques that can potentially be used alongside cPWV (57). Another promising
technique is 4D flow MRI, which has been used to show increased transcranial PWV in AD (58).
Therefore, further studies are needed to elucidate the mechanisms underlying how cPWV and other
innovative techniques may explore AD in 5XFAD and other models.

Limitations:

Study: There are several limitations to this study that must be considered when interpreting the
results. While 5xFAD mice have many similarities to AD pathology in humans, it lacks
neurofibrillary tangles and may have altered cardiac pathways which are poorly elucidated.
Different mouse models of AD also respond to HFD differently (59, 60), while it remains unclear
which most closely mimics that of humans which limits the ability to extrapolate findings to AD
(17, 61). Beyond this, while past studies have clearly shown HFD-induced accelerated cognitive
decline in 5xFAD mice (56), we did not specifically look at this or correlate brain hemodynamics
or other neuronal molecular markers of AD pathology with cardiovascular stiffness changes. While
we sought more to understand the impact of 5xFAD condition on diet-induced changes, it is
possible there are additional neglected links between hemodynamics in the opposite regulation of
AD. While we began an HFD at 5 months of age, the age mice can affect response to HFD, with
past studies showing that HFD feeding before 3 months of age can have protective effects against
cognitive decline (62). Past studies also observed high- and low-weight cohort differences in
response to HFD in 5xFAD, which we did not see, is exacerbated in female mice (62).

<u>Methods:</u> Generally, these models are highly reproducible and adapted from existing aortic PWV
measurements, so they require minimal new equipment. However, an important limitation is that
wall motion must be converted to wave form in mice.

FUNDING

This work is supported by National Institute of Health (NIH) NIDDK T-32, number DK007563
entitled Multidisciplinary Training in Molecular Endocrinology to Z.V.; NSF MCB #2011577I
to S.A.M.; The UNCF/Bristol-Myers Squibb E.E. Just Faculty Fund, Career Award at the
Scientific Interface (CASI Award) from Burroughs Welcome Fund (BWF) ID # 1021868.01,
BWF Ad-hoc Award, NIH Small Research Pilot Subaward to 5R25HL106365-12 from the
National Institutes of Health PRIDE Program, DK020593, Vanderbilt Diabetes and Research
Training Center for DRTC Alzheimer's Disease Pilot & Feasibility Program. CZI Science
Diversity Leadership grant number 2022- 253529 from the Chan Zuckerberg Initiative DAF, an
advised fund of Silicon Valley Community Foundation to A.H.J.; National Institutes of Health
grants R01HL147818 and R01HL144941 (A. Kirabo). Its contents are solely the responsibility of
the authors and do not necessarily represent the official view of the NIH. The funders had no role
in study design, data collection and analysis, decision to publish, or preparation of the
manuscript.

Disclosures

Dr. Reddy is a collaborator and consultant with Indus Instruments, Webster, TX. All other authors
have no competing interests.

Author Contributions

A–M - 3456, –N - 3456, –V - 3456, H–B - 34, –G - 34, –V - 34, –B - 34, –E - 34, –A - 34, AM –
34, ES-34, AC-34, JD -34, DS – 34, SD – 34, T–P - 23, J–G - 678, –E - 678, –D - 678, M–E - 7,8,
G–T - 5678, A–H - 5678, A–R - 12345678

1 Conceived and designed research, 2 Performed experiments and Data collection, 3 Data analysis,
4 Prepared figures, 5 Interpretation of results, 6 Drafted manuscript, 7 Edited and revised
manuscript, 8 Approved final version of manuscript.

References

Figure Legend:

Figure 1: (A) Experimental setup to noninvasively measure cerebral (at ophthalmic artery, OA)
& aortic pulse wave velocity. One probe is fixed at the aortic arch and the other probe is switched
between ophthalmic artery or abdominal aorta sites for Arch-OA combined signals or Arch-Abd
combined signals. Electrocardiography (ECG) waveform is shown for timing. **(B)** Experimental
setup to measure aortic blood flow velocity, aortic blood pressure and) ECGin mice. The probe is
repositioned at carotid artery (shown in the right panel along with pressure & ECG signals). The
probe is repositioned again to measure mitral flow velocity (waveform not shown).

Figure 2: (A) Procedure to calculate aortic impedance – conversion of time domain signals to
frequency domain spectrums and calculation of impedance modulus ($|Z(f)| = |P(f)|/|V(f)|$). Z_c is
calculated as the average of Z_2 to Z_{10} harmonics. **(B)** Procedure to calculate cerebral impedance –
conversion of time domain signals to frequency domain spectrums and calculation of impedance
modulus ($|Z(f)| = |P(f)|/|V(f)|$). Z_c is calculated as the average of Z_2 to Z_{10} harmonics.

Figure 3: Parameters of aortic and cerebral impedance sex-dependent differences in chow-fed 8-
month-old wildtype (C57BL6J) mice. **(A)** Total peripheral resistance (Z_P), **(B)** impedance at first
harmonic (Z_1), **(C)** characteristic impedance, (Z_C) **(D)** and impedance-based pulse wave velocity
(PWV) in aortic and **(E-H)** cerebral impedance. Data are presented as mean±SEM (n = 5/group).
* represents $p < 0.05$, ns indicates a statistically non-significant relationship, as determined
through an unpaired t-test.

Figure 4: Parameters of aortic and cerebral impedance sex-dependent differences in chow-fed 5-
month-old 5XFAD mice. **(A)** Total peripheral resistance (Z_P), **(B)** impedance at first harmonic
(Z_1), **(C)** characteristic impedance, (Z_C) **(D)** and impedance-based pulse wave velocity (PWV) in
aortic and **(E-H)** cerebral impedance. Data are presented as mean±SEM (n = 9-10/group). *
represents $p < 0.05$, ns indicates a statistically non-significant relationship, as determined through
an unpaired t-test.

Figure 5: Parameters of aortic and cerebral impedance sex-dependent differences in high fat
diet-fed 5-month-old 5XFAD mice. **(A)** Total peripheral resistance (Z_P), **(B)** impedance at first
harmonic (Z_1), **(C)** characteristic impedance, (Z_C) **(D)** and impedance-based pulse wave velocity
(PWV) in aortic and **(E-H)** cerebral impedance. Data are presented as mean±SEM (n = 9-
10/group). * represents $p < 0.05$, ns indicates a statistically non-significant relationship, as
determined through an unpaired t-test.

Figure 6: Parameters of aortic and cerebral in pre-stress and stressed state of 5-month-old
5XFAD mice. **(A)** Total peripheral resistance (Z_P), **(B)** impedance at first harmonic (Z_1), **(C)**
characteristic impedance, (Z_C) **(D)** and impedance-based pulse wave velocity (PWV) in aortic
and **(E-H)** cerebral impedance. Data are presented as mean±SEM (n = 10/group). *, **
represents $p < 0.05$, $p < 0.01$, respectively, and ns indicates a statistically non-significant
relationship, as determined through an unpaired t-test.

Supplementary Files:

Supplementary Figure 1: Workflow of inducing a high-fat diet or stress experimental
conditions in 5XFAD mice.

Supplementary Figure 2: Flow of Doppler, electrocardiography (ECG), and blood pressure
(BP) signals from the animal to the transceiver and amplifier to generate the Doppler inphase (I)
and quadrature (Q) audio signals, ECG, and BP signals. The Doppler Flow Velocity System
(DFVS) hardware consists of a high-speed digitizer which sends the digitized I/Q, ECG, & BP to
the DFVS software for acquisition, display, storage, and analysis.

Supplementary Figure 3: Parameters of aortic and cerebral impedance in a mixture of male and
female 5XFAD mice on chow and high-fat diets. (**A**) Total peripheral resistance (Z_P), (**B**)
impedance at first harmonic (Z_1), (**C**) characteristic impedance, (Z_C) (**D**) and impedance-based
pulse wave velocity (PWV) in aortic and (**E-H**) cerebral impedance. Data are presented as
mean±SEM (n = 19/group). ns indicates a statistically non-significant relationship, as determined
through an unpaired t-test.

Supplementary Figure 4: Parameters of aortic and cerebral impedance in 5XFAD mice
measured through noninvasive methods in males and females. (**A**) Myocardial performance
index (Tei index), a measure of the overall function of the heart, which takes into account
systolic and diastolic function, as calculated by (IVCT+IVRT)/ET. (**B**) Cerebral separation
distance. (**C**) Pulse transit time and (**D**) impedance-based pulse wave velocity (PWV) in aortic
and (**E-F**) cerebral parameters. Data are presented as mean±SEM (n = 5/group). ns indicates a
statistically non-significant relationship, as determined through an unpaired t-test.

Supplementary Figure 5: Parameters of aortic and cerebral impedance sex-dependent
differences in non-stressed and stressed 5XFAD mice. (**A**) Total peripheral resistance (Z_P), (**B**)
impedance at first harmonic (Z_1), (**C**) characteristic impedance, (Z_C) (**D**) and impedance-based
pulse wave velocity (PWV) in aortic and (**E-H**) cerebral impedance. Data are presented as
mean±SEM (n = 5/group). * represents p <0.05, ns indicates a statistically non-significant
relationship, as determined through an unpaired t-test.

Supplementary Table 1: Summary of Pulse Wave and Cerebral Impedance Measurements
Utilized.

Figure 1

A

cPWV = 20/7.5
= 2.67m/s

Doppler probe

20mm

25mm

Sample volumes

Dual
20 MHz
Pulsed
Doppler
Transceiver

Aortic arch velocity

Ophthalmic
artery velocity

7.5ms

Aortic arch velocity

Signals from
mouse limbs

ECG

10ms

Abdominal
aortic velocity

aPWV = 25/10
= 2.5m/s

99 ms

B

Pressure
catheter

Doppler
probe

Aortic flow
velocity
Doppler
sample
volume site

Mitral flow
velocity
Doppler sample
volume site

Carotid flow
velocity
Doppler sample
volume site

20 MHz
Pulsed
Doppler
Module

Pressure
Amplifier

Signals from
Mouse limbs

ECG

Waveforms to determine aortic impedance

Waveforms to determine cerebral impedance

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6